Conformational heterogeneity allows access to DNA in longer Transcription

Activator-Like Effector (TALE) arrays.

Kathryn Geiger-Schuller[1,2], Jaba Mitra[3], Taekjip Ha[1,4,5,6], & Doug Barrick[1]*

[1]T.C. Jenkins Department of Biophysics and Program in Molecular Biophysics, Johns

Hopkins University, 3400 N. Charles St, Baltimore MD 21218.

[2]Current address: Broad Institute of Harvard and Massachusetts Institute of

Technology, Cambridge, MA 02142.

[3]Materials Science and Engineering, University of Illinois Urbana-Champaign, Urbana IL

61801.

[4]Department of Physics, Center for the Physics of Living Cells and Institute for Genomic

Biology, University of Illinois at Urbana Champaign, Urbana IL 61801.

[5]Department of Biophysics and Biophysical Chemistry and Department of Biomedical

Engineering, Johns Hopkins University, Baltimore MD 21205.

[6]Howard Hughes Medical Institute, Baltimore, MD, 21205, USA

*Corresponding Author: barrick@jhu.edu, (410) 516-0409

Keywords: TALE repeat, Single-molecule biophysics, FRET, Functional instability,

Deterministic modeling

Abstract

Transcription activator-like effectors (TALEs) bind DNA through an array of

tandem 34-residue repeats. Here, we examine the kinetics of DNA binding for a set of

TALE arrays with identical repeats of varying length (and repeat number) using single

molecule microscopy. Using a new deterministic modeling approach to test kinetic

models consistent with data, we find evidence for conformational heterogeneity in both

the free- and DNA-bound TALE arrays. We connect these results with previous work

demonstrating populations of partly folded TALE states. TALEs forming less than one

superhelical turn around DNA access partly folded open states that inhibit DNA binding,

whereas TALEs forming more than one turn access partly folded open states that

facilitate DNA binding. Overall, we find that increasing repeat number results in

significantly slower interconversion between and among DNA-free and DNA-bound

states. These findings highlight the role conformational dynamics can play in facilitating

the assembly of large complexes.

Introduction

Transcription activator-like effectors (TALEs) are bacterial proteins containing a

domain of tandem DNA-binding repeats as well as a eukaryotic transcriptional activation

domain[1,2]. The repeat domain binds double stranded DNA with a register of one repeat

per base pair. Specificity is determined by the sequence identity at positions twelve and

thirteen in each TALE repeat, which are referred to as repeat variable diresidues

(RVDs)[3–5]. This specificity code has enabled design of TALE-based tools for

transcriptional control[6–10], DNA modifications[11], in-cell microscopy[12,13], and genome

editing (TALENs)[14,15].

TALE repeat domains wrap around DNA in a continuous superhelix of 11.5 TALE

repeats per turn[16,17]. Because TALEs contain on average 17.5 repeats[18], most form

over 1.5 full turns around DNA. Many proteins that form rings around DNA require

energy in the form of ATP to open or close around DNA, yet TALEs are capable of

wrapping around DNA without energy from nucleotide triphosphate hydrolysis. One

possibility is that TALEs bind DNA through an energetically accessible open

conformation. Consistent with this possibility, we previously demonstrated that TALE

arrays can populate partly folded or broken states[19].

Consensus TALEs (cTALEs) are homopolymeric arrays composed of the most

commonly observed residue at each of the 34 positions of the repeat[19]. In addition to

simplifying analysis of folding and conformational heterogeneity, the consensus

approach simplifies analysis of DNA binding, eliminating contributions from sequence

heterogeneity and providing an easy means of site-specific labeling.

Here we characterize DNA binding kinetics of cTALEs using total internal

reflection fluorescence single-molecule microscopy. We find that consensus TALE

arrays bind to DNA reversibly, with high affinity. Analysis of the dwell-times of the on-

and off-states reveals multiphasic binding and unbinding kinetics, suggesting

conformational heterogeneity in both the free and DNA bound state. Here we describe a

deterministic optimization analysis that supports such a model, and provides rate

constants for both conformational changes and binding. Comparing the dynamics

observed here to previously characterized local unfolding suggests that locally unfolded

states inhibit binding of short cTALE arrays (less than one full superhelical turn around

DNA), whereas they promote binding of long arrays (more than 1 full superhelical turn).

Results

cTALE design

Consensus TALE (cTALE) repeat sequence was design previously described[19].

To avoid self- association of cTALE arrays, we fused arrays to a conserved N-terminal

extension of the PthXo1 gene. Although the sequence of this domain differs from a

TALE repeat sequence, the structure of this domain closely mimics four TALE

repeats[17,20] and is required for full transcriptional activation[20]. In this study, all repeat

arrays contain this solubilizing N-terminal domain.

cTALE local instability promotes population of partly folded states

Figure 1A depicts types of partly folded states of a generic repeat protein. In the

fully folded state, all repeats are folded, and all interfaces are intact. In the end-frayed

states, one or more terminal repeats are unfolded and all interfaces, except the

interface(s) between the unfolded and adjacent folded repeat(s), are intact. In the

internally unfolded states, a central repeat is unfolded and all interfaces, except the

interfaces involving the unfolded repeats, are intact. In the interfacially ruptured state, all

repeats are folded but one interface is disrupted due to local structural distortion.

Figure 1B shows calculated free energy differences between various partly

folded states and the fully folded repeat array for two different RVDs (NS and HD) in an

otherwise identical consensus sequence background, using the intrinsic and interfacial

engeries we determined previously[19]. The distribution of partly folded states is

calculated for different 20-repeat arrays containing two types of TALE arrays (with the

NS RVD in red and with the HD RVD in blue) as well as consensus ankyrin arrays

(cAnk in black). For cTALE arrays, end frayed states are within a few RT, internally

unfolded states are highest in energy, and interfacially ruptured states fall energetically

between end frayed and internally unfolded states.

Changing the RVD sequence affects the distribution of these partly folded states:

HD repeat containing arrays are more likely to internally unfold or interfacially rupture

than NS repeat containing arrays. Both types of cTALEs are more likely to populate

many of these partly folded states than cAnk is to populate even the lowest energy

partly folded state, the end frayed state. Thus, compared to ankyrin repeats, cTALEs

are locally unstable, meaning they are likely to form partly folded states. As these

states disrupt the superhelix, they may facilitate DNA binding.

Single-molecule studies of cTALE binding to DNA

To ask if cTALE local instability is relevant for DNA binding kinetics, DNA binding

trajectories were measured using single molecule total internal reflection fluorescence

(smTIRF). Figure 2A shows a schematic of the smTIRF experiments performed to

measure DNA binding. For site-specific cTALE labeling, R30 is mutated to cysteine in a

single repeat. Position 30 is frequently a cysteine in naturally occurring TALEs (in earlier

folding studies, arginine was chosen in the consensus sequence to avoid disulfide

formation)[19]. This cysteine was Cy3 (FRET donor)-labelled using maleimide chemistry,

and was attached to biotinylated slides via the C-terminal His_6 tag and α-Penta•His

antibodies. At salt concentrations below 300 mM NaCl, cTALEs aggregate. Because

DNA binding is weak at high salt concentrations, measuring binding kinetics in bulk at

high salt is not possible. However, tethering cTALEs to the quartz slide at high salt

prevents self-association, even at the low salt concentrations required to study DNA

binding kinetics. A histogram of $NcTALE_8$ (8 NS-type repeats and the N-terminal

domain) labeled via a cysteine in the first repeat shows a single peak at zero FRET

efficiency, as expected for donor-only constructs (Figure 2B).

To test for DNA binding to tethered cTALE constructs, we added Cy5 (FRET

acceptor)-labeled 15 bp-long DNA ($Cy5.A_{15}/T_{15}$) to tethered $NcTALE_8$. This results in a

new peak at a FRET efficiency of 0.45, indicating that DNA binds directly to cTALE

arrays. As DNA concentration in solution is increased, the peak at 0.45 FRET efficiency

increases in population (Figure 2C-D), suggesting a measurable equilibrium between

free and bound DNA rather than saturation or irreversible binding. In support of this,

single molecule time trajectories show interconversion between bound and unbound

states, providing access to rates of binding and dissociation.

As expected for reversible complex formation, the peak at 0.45 FRET efficiency

can be competed away by adding mixtures of labeled and unlabeled DNA to pre-formed

cTALE-labelled DNA complexes (schematic shown in Figure 2E; pre-formed complex

shown in Figure 2F, competition data shown in Figure 2G-H). Challenging pre-formed

complexes with a mixture of 5 nM unlabeled DNA and 15 nM labeled DNA results in a

slight decrease in the population of the peak at 0.45 FRET (compare Figures 2F and

2G). Challenging with a mixture of 50 nM unlabeled DNA and 15 nM labeled DNA

further decreases the peak at 0.45 FRET (Figure 2H).

cTALE arrays display multiphasic DNA-binding kinetics.

In addition to the short smTIRF movies used to generate smFRET histograms

from many molecules, long movies were also collected to examine the extended

transitions of individual molecules between the low- and high-FRET (0 and 0.45) (Figure

3A-B). A transition from low to high FRET (0 to 0.45) indicates that the acceptor

fluorophore on DNA moved close enough to the donor on the protein for FRET and is

likely a binding event. A transition from high to low FRET (0.45 to 0.0) indicates the

acceptor fluorophore on DNA moved too far away from the donor on the protein for

FRET and is likely an unbinding event. Low-FRET states show low colocalization with

signal upon direct excitation of the acceptor, confirming that high-FRET states are DNA-

bound states and low-FRET states are DNA-free states (Figure S1). These long single

molecule traces show both long- and short-lived low- and high-FRET states, indicating

that kinetics are multi-phasic (Figure 3A-B). Binding events (transitions from low to high

FRET) become more frequent as bulk DNA concentration increases (compare

representative traces at 1 nM dsDNA to 15 nM dsDNA; Figure 3A and Figure 3B).

Cumulative distributions generated from dwell times in the low FRET state at a given

DNA concentration are best-fit by a double-exponential decay, indicating a minimum of

two kinetic phases associated with binding events (Figure 3C). Cumulative distributions

generated from dwell times in the high FRET state are best-fit by a double-exponential

decay, indicating that there are a minimum of two kinetic phases for unbinding as well

(Figure 3D).

The rate constant for the fast phase in DNA binding shows a linear increase with

DNA concentration (Figure 3E), indicating that this step involves an associative binding

mechanism. The slope of these rate constants as a function of DNA concentration gives

a bimolecular rate constant of 5.9×10^8 $nM^{-1}s^{-1}$, close to the diffusion limit. The rate

constant for the slower phase (0.59 s^{-1}) is independent of DNA concentration indicating

a unimolecular isomerization mechanism (Figure 3E).

In contrast, neither of the two fitted rate constants for transitions from high to low

FRET (0.45 to 0.0; unbinding events) depends on DNA concentration, suggesting that

unbinding involves two (or more) unimolecular processes (Figure 3F). The rate

constants of these two phases are 1.2 s^{-1} and 0.13 s^{-1} respectively.

To rule out kinetic contributions of TALEs threading axially onto the ends of short

DNAs, binding kinetics were measured with capped double-helical DNA sites. Capped

DNA was generated by forming 5'digoxygenin-A_5-Cy5-A_{15} duplexed with 5'-digoxygenin-

T_{26} and adding a three-fold molar excess of anti-Digoxygenin. Low and high FRET dwell

time cumulative distributions generated from capped DNA-binding kinetics are bi-

phasic, similar to distributions from uncapped DNA (Figure S2). The DNA concentration-

independent rate constant for binding is roughly the same for capped DNA as for

uncapped DNA (compare $FRET_{L \to H}$ red and blue triangles in Figure S2), as are the

dissociation rate constants (compare $FRET_{H \to L}$ red and blue triangles in as well as

$FRET_{H \to L}$ red and blue circles in Figure S2). The rate constant for bimolecular binding of

capped DNA decreases compared to that for uncapped DNA (compare $FRET_{L \to H}$ red

and blue circles in Figure S2), which is consistent with the expected decrease in the

rate of diffusion of the larger capped DNA. To assess the effect of molecular weight

increase on diffusion of capped versus uncapped DNA, Sednterp[21], a program

commonly used to estimate sedimentation and diffusion properties of biomolecules, was

used to estimate maximum diffusion coefficients. Including the two antibodies bound on

the ends of capped DNA (320 kDa total) gives an estimated diffusion coefficient of

4.7×10^{-7} cm^{2}s^{-1}, which is much lower than the estimated diffusion coefficient of the

uncapped DNA (1.5×10^{-6} cm^{2}s^{-1}). This ~3.6-fold decrease in the diffusion constant for

capped DNA is similar to the 6.7-fold decrease in the bimolecular rate constant for

binding of capped DNA (Figure S2).

Longer cTALEs have slower DNA binding and unbinding kinetics

To examine how increasing the length of the cTALE array influences DNA

binding, we generated Cy3-labelled constructs with 16 and 12 cTALE repeats, and

measured binding to a longer Cy5-labelled DNA (A_{23}/T_{23}). Because we did not observe

FRET for these longer constructs, a fluorescence colocalization microscopy protocol

was used instead (Figure S3). In this protocol, Cy3 was first imaged for ten camera

frames (1017.5 msec total) to identify positions of single TALE molecules. Then a long

time series of fluorescence images of Cy5 signal were collected through directly exciting

Cy5 on the DNA, and time trajectories of Cy5 signal were generated from the initially

identified single TALE positions.

Increasing the number of cTALE repeats from 8 to 12 and 16 dramatically affects

DNA binding kinetics. Long movies collected over a range of DNA concentrations show

short- and long-lived Cy5 signal on and off states, indicating a level of kinetic

heterogeneity similar to NcTALE$_8$ (Figure S3). Single molecule traces were analyzed

using a thresholding filter (see Materials and Methods and Figure S3) to identify states

and dwell times. Cumulative distributions were generated from dwell times at low Cy5

signal (unbound states, with lifetimes representing binding kinetics), and at high Cy5

signal (bound states, with dwell times representing unbinding kinetics). As with the eight

repeat constructs, unbound cumulative distributions for these longer TALE arrays are

best-fit by double exponential decays, particularly at high DNA concentrations (compare

the cumulative distribution at low DNA concentration, Figure S4A, to cumulative

distribution at 5 nM DNA, Figure S4B). Bound cumulative distributions for longer TALE

arrays are best-fit by double exponential decays (Figure S4C-D). All apparent rate

constants are much smaller for NcTALE$_{16}$ and NcTALE$_{12}$ (green/black circles and

triangles, Figure 4A-B) compared to NcTALE$_8$, indicating that binding and unbinding is

impeded by increasing the length of the binding surface between cTALEs and their

cognate DNA (Figure 4C). To address whether differences in binding kinetics are

related to experimental differences between colocalization and FRET assays,

alternating laser experiments were performed by switching between FRET and

colocalization detection (every 5 frames) within single molecule trajectories (Figure S1).

Changes in FRET and colocalization signals occurred simultaneously according to

single molecule time traces, showing that differences in binding and unbinding kinetics

of short and longer cTALEs are not due to differences in colocalization and FRET

assays (Figure S1).

A deterministic approach to modeling cTALE-DNA binding kinetics.

To determine how the kinetic changes above are partitioned into underlying

kinetic steps in binding, we fitted various kinetic models to the cumulative distributions

for binding and unbinding. In addition to providing information about the mechanism of

binding, this approach allows us to estimate the underlying microscopic rate constants

and compare them for different constructs. This approach is generally applicable to

studies of complex single molecule kinetics. Numerical integration was used to calculate

the relative population of cTALE states as a function of time (Figures 5A-C and 5G-H),

given a binding mechanism, an associated set of rate laws, and a set of initial

conditions. Cumulative distributions of unbound dwell times represent the distribution of

times single molecules spent in the unbound state before transitioning into the bound

state, allowing us to split the kinetic scheme when fitting to single-molecule dwell times.

Among the various models tested, the model that is most consistent with the data

has two unbound DNA-free states and two DNA-bound states. This is consistent with

alternating laser experiments showing that DNA is only colocalized when cTALEs are in

the high FRET state (Figure S1). This four-state model includes a TALE isomerization

step in the absence of DNA from a DNA-binding incompetent conformation (which we

refer to as TALE) to DNA-binding competent conformation (which we refer to as TALE*).

The DNA-binding competent TALE* conformer binds and unbinds DNA (called TALE*

when DNA free and TALE*~DNA when DNA-bound). Before unbinding, a fraction of

TALE*~DNA isomerizes to a longer-lived DNA-bound state called TALE‡~DNA.

Based on this mechanism, the rate laws for binding are given in equations 1a -

1d.

$$\frac{d[TALE]}{dt} = -k_1[TALE] + k_{-1}[TALE^*] \tag{1a}$$

$$\frac{d[TALE^*]}{dt} = k_1[TALE] - k_{-1}[TALE^*] - k_2[TALE^*][DNA] \tag{1b}$$

$$\frac{d[TALE^* \sim DNA]}{dt} = k_2[TALE^*][DNA] \tag{1c}$$

252
$$K_{eq,DNA-free} = \frac{k_1}{k_{-1}}$$
(1d)

Since the single-molecule dwell-time histograms of the unbound states are insensitive

to the isomerization after DNA binding, the equation describing the time evolution of the

long-lived bound state (TALE‡~DNA) is not relevant to our analysis of unbound-state

lifetimes.

To determine microscopic rate constants k_1, k_{-1}, and k_2, equations 1a-1c were

numerically integrated in Matlab, and the fraction of TALE*~DNA as a function of time

was fitted to the low-FRET cumulative distributions (NcTALE$_8$; Figure 5D-E) or to the no

colocalization cumulative distributions (NcTALE$_{16}$; Figure 5J-K). Microscopic rate

constants were adjusted to reduce sum of the squared residuals between the

concentration of TALE*~DNA (the direct product of binding) as a function of time and

single-molecule cumulative distributions. In both cases, cumulative distributions at

different bulk DNA concentrations were fitted globally. Initial fractions of TALE and

TALE*~DNA were set to zero, and the initial fraction of TALE* was set to one.

Confidence intervals (CI) were estimated by bootstrapping (Table 1; mean and 68% CI

from 2000 or 8000 bootstrap iterations).

Rate laws for dissociation are given in equations 2a - 2d

269
$$\frac{d[TALE^* \sim DNA]}{dt} = -k_{-2}[TALE^* \sim DNA] - k_3[TALE^* \sim DNA] + k_{-3}[TALE^{\ddagger} \sim DNA]$$ (2a)

270
$$\frac{d[TALE^{\ddagger} \sim DNA]}{dt} = k_3[TALE^* \sim DNA] - k_{-3}[TALE^{\ddagger} \sim DNA]$$
(2b)

271
$$\frac{d[TALE^*]}{dt} = k_{-2}[TALE^* \sim DNA]$$
(2c)

272
$$K_{eq,DNA-bound} = \frac{k_3}{k_{-3}}$$
(2d)

As with the system of equations above (1a-d), the equation describing the time

evolution of the binding-incompetent free state (TALE) is not relevant to our analysis of

bound-state lifetimes.

To determine microscopic rate constants k_{-2}, k_{-3}, and k_3, equations 2a-2c were

numerically integrated in Matlab, and the fraction of TALE* as a function of time was

fitted to the high-FRET cumulative distributions (NcTALE$_8$; Figure 5F) or to the low

colocalization cumulative distributions (NcTALE$_{16}$; Figure 5L). Microscopic rate

constants were adjusted to reduce sum of the squared residuals between the

concentration of TALE* (the direct product of dissociation) as a function of time and

single-molecule cumulative distributions. In both cases, cumulative distributions at

different bulk DNA concentrations were fitted globally. The initial fraction of

TALE*~DNA conformer was set at one; all other initial fractions were set to zero.

Confidence intervals were estimated by bootstrapping (Table 1; mean and 68% CI from

2000 iterations).

Fitted curves reproduce the experimental cumulative distributions for binding and

unbinding (Figure 5), both for the short and long cTALE arrays, with reasonably small

residuals, over a range of DNA concentrations. Generally, fitted rate constants have

confidence intervals of 10% or smaller (Table 1).

Comparison of microscopic rate constants for 8, 12, and 16 repeats show some

significant differences. The bimolecular microscopic binding rate constant, k_2, is slightly

larger for 8 repeats than for 12 and 16 repeats (1.1, 0.31, and 0.39 nM^{-1}s^{-1} for 8, 12, and

16 repeats respectively). However, microscopic unbinding rate constant, k_{-2}, is higher

for 8 repeat cTALEs (0.66 s^{-1} for NcTALE$_8$ versus 0.13 s^{-1} for NcTALE$_{12}$ and 0.299 s^{-1} for

NcTALE$_{16}$). Also, bound state isomerization (interconversion between TALE*~DNA and

TALE‡~DNA) is 5-10 times slower for 16 and 12 repeat cTALEs than 8 repeat cTALEs.

The value of $K_{eq, DNA-free}$, which is a measure of the equilibrium proportion of the

unbound TALE that is DNA-binding competent (TALE*) to that which is binding-

incompetent (TALE), is larger for cTALEs with 8 repeats ($K_{eq, DNA-free}$ = 1.32) than for

cTALEs with 12 and 16 repeats ($K_{eq, DNA-free}$ = 0.11 and $K_{eq, DNA-free}$ = 0.61, respectively).

Discussion

By measuring DNA-binding kinetics of cTALE arrays that form 0.7, 1, and 1.4

superhelical turns, we probe the functional relevance of locally unfolded TALE states.

We describe a novel method to glean mechanistic details from complex single molecule

kinetics. In our simplified cTALE system, we find conformational heterogeneity in both

DNA free and DNA-bound states. We find that association is slowed in arrays

containing one full turn of repeats or more. Because most natural and designed TALEs

contain more than a full turn of repeats, this finding has important implications for design

of high affinity TALE molecules, suggesting that placement of destabilized repeats at

specific positions may increase activity.

cTALEs containing NS RVD bind DNA with high affinity

NS is an uncommon RVD in natural TALEs. Previous reports suggest that NS is

fairly nonspecific, but may bind with higher affinity than other common RVDs (NG, NI,

NN, and HD)[5]. Our fitted rate constants can be used to calculate the apparent K_d (K_{app})

calculated from using equation as follows:

$$K_{app} = \frac{[TALE-DNA+TALE*-DNA]}{[DNA][TALE+TALE*]}$$

319
$$= \frac{K_{eq,DNA-free}K_2 + K_{eq,DNA-free}K_2 K_{eq,DNA-bound}}{1+K_{eq,DNA-free}}$$ (3)

$$= \frac{\frac{k_1}{k_{-1}} \times \frac{k_2}{k_{-2}}(1+\frac{k_3}{k_{-3}})}{1+\frac{k_1}{k_{-1}}}$$

where $K_2 = k_2/k_{-2}$.

Using fitted rate constants from Table 1 in the final equality in equation 3 gives values

for K_{app} of 2.5 nM for the 8 repeat cTALE array, 0.5 nM for the 12 repeat cTALE array,

and 1.0 nM for the 16 repeat cTALE array. Increasing the number of repeats has a

modest affect on the apparent K_d due to the increased population of binding

incompetent DNA free TALE in the 12, and 16 repeat arrays. This affinity change is

small compared to a previous report studying length dependence on affinity of designed

TALEs (dTALEs) showing the K_d of a dTALE decreased by a factor of two with the

addition of only 1.5 repeats[22].

TALEs are believed to read out sequence information from one strand[3]. Due to

the asymmetry of our DNA sequences (poly-dA base-paired with poly-dT), in principle,

the FRET efficiency contains information on the binding orientation (and thus strand

preference). However, based on the crystal structure of the DNA-bound state of TAL-

effector PthXo1[17], we estimate that the distance between the donor site of NcTALE$_8$

(repeat 1) to the 5' acceptor site on the DNA (Cy5-A$_{15}$/T$_{15}$) should be similar for both the

dA-sense or dT-sense orientations (Figure S5A). Thus, the FRET data does not

discriminate between the two modes of binding for the eight-repeat construct. However,

for the 16 repeat NS RVD cTALE arrays, the PthXo1 model suggests very different

distances (25 Å versus 73 Å for the dT-sense or dA-sense respectively, Figure S5B)

between the donor site (TALE repeat 14) and the acceptor site (5' Cy5-A$_{23}$/T$_{23}$). To

restrict the number of binding positions available to longer cTALE arrays, the 23 base

pair DNA used for NcTALE$_{16}$ measurements (as well as DNA depicted in Figure S5B)

has the same number of additional base pairs as repeats (8 additional repeats and 8

additional base pairs) compared to the 15 base pair DNA used for NcTALE$_8$

measurements (as well as DNA depicted in Figure S5A). While we limited the number of

available binding positions, it may be possible for cTALEs to slide along DNA. However,

taking into account the four repeat N-terminal capping domain, there are only three

available base pairs in the bound complex. Thus we don't expect the distance

measurements to change by more than 10Å (~3 base pairs) if sliding occurs. The

observation that there is colocalization but no measurable FRET when NcTALE$_{16}$ is

bound to DNA suggests that cTALEs containing the NS RVD prefer adenine (the dA-

sense mode) compared with thymine bases, consistent with previous reports[3].

Conformational heterogeneity in the unbound state could be the result of local unfolding

The cumulative distributions of dwell-times in Figure 3 provide clear evidence for

conformational heterogeneity in both the free and DNA-bound cTALEs. Although the

deterministic modeling supports such heterogeneity, puts it in the framework of a

molecular model, and provides a means to determine the microscopic rate and

equilibrium constants, such analysis provides little information about the structural

nature of TALE conformational heterogeneity.

Figure 6 shows a model of cTALE conformational change consistent with DNA

binding kinetics. In this model there are four TALE states. DNA-free cTALEs comprise

both incompetent and binding competent states. DNA-bound cTALES comprise

encounter and locked complexes with DNA. For 8 repeat cTALE arrays, the DNA-

binding competent state is more highly populated than the DNA-binding incompetent

state. In this reaction scheme, the DNA-binding incompetent state can be regarded as

an off-pathway conformation that inhibits DNA binding (Figure 6A).

Because the 8 repeat cTALE array does not form multiple turns of a superhelix,

unfolding to bind DNA is not required. In the model in Figure 6, the binding competent

state is the fully folded conformation, whereas the binding incompetent state includes

partly folded conformations. Consistent with this interpretation, increasing populations of

partly folded states through addition of 1M urea and through entropy enhancing

mutations decreases apparent binding rates of 8 repeat cTALEs (Figure S6). This is

also consistent with a partly folded DNA-binding incompetent state in shorter cTALE

arrays.

For 12 and 16 repeat cTALE arrays, the DNA-binding incompetent state is more

highly populated than the DNA-binding competent state. In the model in Figure 6, the

DNA-binding competent state is a high-energy conformation required for DNA binding

(Figure 6B-C). Because 12 and 16 repeat cTALEs are expected to form 1 and 1.4 turns

(excluding the N-terminal domain), we hypothesize that the binding competent state

includes some partly folded states that allow access to DNA. Not all partly folded states

open the array to access DNA; therefore, the binding incompetent state includes some

nonproductive partly folded states in addition to the fully folded state.

In arrays containing 12 or more cTALEs, the binding competent and binding

incompetent states likely include mixtures of many specific partly folded states. Because

the types of partly folded states are unknown, connecting equilibria between binding

competent and binding incompetent states to calculated partly folded equilibria (using

folding free energies similar to Figure 1) is challenging. Future work towards

understanding the structural characteristics of the binding competent state in TALE

arrays of one or more turns would inform which partly folded states to include in the

calculation, making this comparison meaningful.

Conformational heterogeneity in the bound state

Previous reports show that TALEs have multiple diffusional modes when

searching nonspecific DNA[23]. Our work suggests that cTALEs have multiple binding

modes (encounter and locked states in Figure 6) indicating cTALEs undergo a

conformational change also when bound to specific DNA sequences. Table 1 shows

that microscopic rate constants for transition into and out of longer lived locked bound

states become much slower in 12 and 16 repeat cTALES compared with 8 repeat

cTALEs (k_{-3} and k_3). These rate constants decrease much more than the microscopic

unbinding rate constant (the k_{-2} values are 0.66 s^{-1}, 0.13 s^{-1}, and 0.299 s^{-1} for NcTALE$_8$,

NcTALE$_{12}$, and NcTALE$_{16}$ respectively) suggesting a large conformational change that

depends on the number of repeats. Although the model does not provide information on

structure of this conformational change, it is possible this conformational change

involves a slinky motion to decrease helical rise. Consistent with this hypothesis, crystal

structures of TALEs in the free and DNA-bound state show 11.5 repeats per turn in both

states, but the helical rise decreases upon binding[16]. Another possibility involves

specific interaction with RVDs and bases in the major groove of DNA. Crystal structures

show little deformation of DNA structure, so bending of DNA seems unlikely. While we

can only hypothesize about the structural nature of the conformational changes,

deterministic simulations show that cTALEs bind DNA through short encounters which

occasionally become long-lived locked conformations (Figure 6). Taken together, these

findings indicate that functional stability plays a crucial role in cTALE DNA binding, and

demonstrate the importance of conformational dynamics in complex assembly.

Materials and Methods

Cloning, expression, purification, and labeling

Consensus TALE repeat constructs were cloned with C-terminal His_6 tags via an

in-house version of Golden Gate cloning[24]. TALE constructs were grown in BL21(T1R)

cells at 37°C to an OD of 0.6-0.8 and induced with 1 mM IPTG. Following cell pelleting

and lysis, proteins were purified by resuspending the insoluble material in 6M urea, 300

mM NaCl, 0.5 mM TCEP, and 10 mM $NaPO_4$ pH 7.4. Constructs were loaded onto a Ni-

NTA column. Protein was eluted using 250 mM imidazole and refolded during buffer

exchange into 300 mM NaCl, 30% glycerol, 0.5 mM TCEP, and 10 mM $NaPO_4$ pH 7.4.

Labelling of cTALE arrays followed a previously reported protocol[25]. $NcTALE_8$

and $NcTALE_{12}$ were labeled at residue R30C in the first repeat, while $NcTALE_{16}$ was

labeled at residue R30C in the fourteenth repeat. 1 mg protein was loaded onto 500 uL

NiNTA spin column. The column as washed with 10 column volumes of 300 mM NaCl,

0.5 mM TCEP, and 10 mM $NaPO_4$ pH 7.4. Tenfold molar excess Cy3 maleimide dye

was resuspended in 10 μL DMSO and added to column. The column was rocked at

room temperature for 30 minutes, then at 4°C overnight. Cy3-labeled protein was eluted

with 250 mM imidazole, 300 mM NaCl, 30% glycerol, 0.5 mM TCEP, and 10 mM $NaPO_4$

pH 7.4. Protein was stored in 300 mM NaCl, 30% glycerol, 0.5 mM TCEP, and 10 mM

$NaPO_4$ pH 7.4 at -80°C.

Oligonucleotides

Sequences used for binding studies were 5'-Cy5-A_{15}-3' and 5' T_{15}-3' duplex

(Cy5-A_{15}/T_{15}) for 8 repeat binding studies, and 5'-Cy5-A_{23}-3' and 5' T_{23}-3' duplex (Cy5-

A_{23}/T_{23}) for 12 and 16 repeat binding studies. DNA was annealed at 5 μM concentration

with 1.2-fold molar excess unlabeled strand in 10 mM Tris pH 7.0, 30 mM NaCl.

Single-molecule detection and data analysis

Biotinylated quarts slides and glass coverslips were prepared as previously

described[25]. Cy3-labeled cTALEs were immobilized on biotinylated slides taking

advantage of neutravidin interaction with biotinylated α-penta•His antibody which binds

the His$_6$ cTALE tag. Slides were pretreated with blocking buffer (5 μL yeast tRNA, 5 μL

BSA, 40 μL T50) before addition of 250 pM labeled cTALE. Cy5-labeled duplex DNA

was mixed with imaging buffer (20 mM Tris pH 8.0, 200 mM KCl, 0.5 mg mL^{-1} BSA, 1

mg mL^{-1} glucose oxidase, 0.004 mg mL^{-1} catalase, 0.8% dextrose and saturated Trolox

~1mg mL^{-1}) and molecules were imagined using total internal reflection fluorescence

microscopy. The time resolution was 50 msec for NcTALE$_8$ and 100 msec for NcTALE$_{16}$

and NcTALE$_{12}$. Collection and analysis was performed as previously described[26].

FRET histograms

A minimum of 20 short movies were collected, and the first 5 frames (50 msec

exposure time) were used to generate smFRET histograms. FRET was calculated as

$I_A/(I_A+I_D)$ where I_A and I_D are donor-leakage and background corrected fluorescence

emission of acceptor (Cy5) and donor (Cy3) fluorophores. In competition experiments,

unlabeled DNA with the same sequence as labeled DNA was mixed at indicated

concentrations with labeled DNA prior to imaging.

Dwell time analysis

Long movies were collected with 50 msec exposure time for NcTALE$_8$ and 100

msec exposure time for NcTALE$_{16}$ and NcTALE$_{16}$. At least 20 representative traces at

each DNA concentration were selected and dwell times were determined by fitting as

previously described using HaMMy[27] for FRET in NcTALE$_8$. Dwell times in NcTALE$_{12}$

and NcTALE$_{16}$ colocalization experiments are determined by using a thresholding

procedure for Cy5 excitation (Figure S3). The algorithm used to identify low and high

emission states here is slightly different than previously described thresholding

algorithms[28]. To reduce the number of incorrectly identified transitions arising from

increased background and noise at higher Cy5-labeled DNA concentrations, a

thresholding algorithm with two limits was implemented (see Figure S3). All FRET and

colocalization data are well described by models with two distinct states (0.0 FRET and

~0.45 FRET as well as low colocalization and high colocalization). Dwell times of the

same state (low versus high FRET or low versus high colocalization) for all traces at a

given DNA concentration are compiled, and cumulative distribution is generated with

spacing equal to imaging exposure time.

To determine apparent rate constants using model-independent analysis,

cumulative distributions were fitted with single and double exponential decays (Figure 3

and 4). Observed rates from exponential decay fits were plotted as a function of DNA

concentration. Apparent rate constants were calculated as slope of DNA concentration-

dependent observed rates or average of DNA concentration-independent observed

rates.

Deterministic modeling

Equations 1a-1c and 2a-2c were numerically integrated using the ODE15s and

ODE45 solver in MATLAB. Microscopic rate constants were adjusted to minimize the

sum of squared residuals between ODE-determined concentration of bound or free

TALE and single molecule cumulative distributions using lsqnonlin in MATLAB. 68%

confidence intervals were estimated by performing 2000 or 8000 bootstrap iterations in

which residuals from the best fit of the model to the data were randomly re-sampled

(with replacement) and re-fitted.

Acknowledgements

The authors thank members of the Barrick and Ha lab for their input on this work.

The authors acknowledge the support of the Center for Molecular Biophysics at Johns

Hopkins and Dr. Katherine Tripp for instrumental and technical support. Support to

KGS was provided by NIH training grant T32-GM008403. Support for this project was

provided by NIH grant 1R01-GM068462 to DB and GM112659 to TH.

References

1. Kay, S., Hahn, S., Marois, E., Hause, G. & Bonas, U. A bacterial effector acts as a

plant transcription factor and induces a cell size regulator. *Science* **318,** 648–651

(2007).

2. Römer, P. *et al.* Plant pathogen recognition mediated by promoter activation of the

pepper Bs3 resistance gene. *Science* **318,** 645–648 (2007).

3. Boch, J. *et al.* Breaking the code of DNA binding specificity of TAL-type III effectors.

Science **326,** 1509–1512 (2009).

4. Moscou, M. J. & Bogdanove, A. J. A simple cipher governs DNA recognition by TAL

effectors. *Science* **326,** 1501 (2009).

5. Miller, J. C. *et al.* Improved specificity of TALE-based genome editing using an

expanded RVD repertoire. *Nat. Methods* **12,** 465–471 (2015).

6. Cong, L., Zhou, R., Kuo, Y.-C., Cunniff, M. & Zhang, F. Comprehensive interrogation

of natural TALE DNA-binding modules and transcriptional repressor domains. *Nat.*

Commun. **3,** 968 (2012).

7. Geissler, R. *et al.* Transcriptional activators of human genes with programmable

DNA-specificity. *PloS One* **6,** e19509 (2011).

8. Li, Y., Moore, R., Guinn, M. & Bleris, L. Transcription activator-like effector hybrids for

conditional control and rewiring of chromosomal transgene expression. *Sci. Rep.* **2,**

897 (2012).

9. Mahfouz, M. M. *et al.* Targeted transcriptional repression using a chimeric TALE-

SRDX repressor protein. *Plant Mol. Biol.* **78,** 311–321 (2012).

10.   Zhang, F. *et al.* Efficient construction of sequence-specific TAL effectors for

modulating mammalian transcription. *Nat. Biotechnol.* **29,** 149–153 (2011).

11.   Maeder, M. L. *et al.* Targeted DNA demethylation and activation of endogenous

genes using programmable TALE-TET1 fusion proteins. *Nat. Biotechnol.* **31,** 1137–

1142 (2013).

12.   Miyanari, Y., Ziegler-Birling, C. & Torres-Padilla, M.-E. Live visualization of

chromatin dynamics with fluorescent TALEs. *Nat. Struct. Mol. Biol.* **20,** 1321–1324

(2013).

13.   Ma, H., Reyes-Gutierrez, P. & Pederson, T. Visualization of repetitive DNA

sequences in human chromosomes with transcription activator-like effectors. *Proc.*

Natl. Acad. Sci. U. S. A. **110,** 21048–21053 (2013).

14.   Li, T. *et al.* TAL nucleases (TALNs): hybrid proteins composed of TAL effectors

and FokI DNA-cleavage domain. *Nucleic Acids Res.* **39,** 359–372 (2011).

15.   Christian, M. *et al.* Targeting DNA Double-Strand Breaks with TAL Effector

Nucleases. *Genetics* **186,** 757–761 (2010).

16.   Deng, D. *et al.* Structural basis for sequence-specific recognition of DNA by TAL

effectors. *Science* **335,** 720–723 (2012).

17.   Mak, A. N.-S., Bradley, P., Cernadas, R. A., Bogdanove, A. J. & Stoddard, B. L.

The crystal structure of TAL effector PthXo1 bound to its DNA target. *Science* **335,**

716–719 (2012).

18.   Boch, J. & Bonas, U. Xanthomonas AvrBs3 family-type III effectors: discovery

and function. *Annu. Rev. Phytopathol.* **48,** 419–436 (2010).

19.   Geiger-Schuller, K. & Barrick, D. Broken TALEs: Transcription Activator-like

542   Effectors Populate Partly Folded States. *Biophys. J.* **111,** 2395–2403 (2016).

543   20.   Gao, H., Wu, X., Chai, J. & Han, Z. Crystal structure of a TALE protein reveals an

544   extended N-terminal DNA binding region. *Cell Res.* **22,** 1716–1720 (2012).

545   21.   edited by S. E. Harding, A. J. R. *Analytical ultracentrifugation in biochemistry and*

546   *polymer science.* (Royal Society of Chemistry, 1992).

547   22.   Rinaldi, F. C., Doyle, L. A., Stoddard, B. L. & Bogdanove, A. J. The effect of

548   increasing numbers of repeats on TAL effector DNA binding specificity. *Nucleic Acids*

549   *Res.* (2017). doi:10.1093/nar/gkx342

550   23.   Cuculis, L., Abil, Z., Zhao, H. & Schroeder, C. M. Direct observation of TALE

551   protein dynamics reveals a two-state search mechanism. *Nat. Commun.* **6,** 7277

552   (2015).

553   24.   Cermak, T. *et al.* Efficient design and assembly of custom TALEN and other TAL

554   effector-based constructs for DNA targeting. *Nucleic Acids Res.* **39,** e82 (2011).

555   25.   Rasnik, I., Myong, S., Cheng, W., Lohman, T. M. & Ha, T. DNA-binding

556   Orientation and Domain Conformation of the E.coli Rep Helicase Monomer Bound to

557   a Partial Duplex Junction: Single-molecule Studies of Fluorescently Labeled

558   Enzymes. *J. Mol. Biol.* **336,** 395–408 (2004).

559   26.   Roy, R., Hohng, S. & Ha, T. A practical guide to single-molecule FRET. *Nat.*

560   *Methods* **5,** 507–516 (2008).

561   27.   McKinney, S. A., Joo, C. & Ha, T. Analysis of single-molecule FRET trajectories

562   using hidden Markov modeling. *Biophys. J.* **91,** 1941–1951 (2006).

563   28.   Blanco, M. & Walter, N. Analysis of Complex Single Molecule FRET Time

564   Trajectories. *Methods Enzymol.* **472,** 153–178 (2010).

Figure 1. cTALEs populate partly folded states. (A) Cartoon of different partly folded

TALE conformational states. End-frayed states have one (or more) terminal repeats

unfolded. Internally unfolded states have a central repeat unfolded. Interfacially

fractured states have a disrupted interface between adjacent repeats. (B) Free energies

of partly folded states, calculated from previously published measurements and

analysis[19], relative to the fully folded state, for consensus TALE repeats with the NS

repeat-variable diresidue sequence (cTALE(NS), red), consensus TALE repeats with

the HD repeat-variable diresidue sequence (cTALE(HD), blue), and consensus ankyrin

repeats (cAnk, black).

Figure 2. cTALEs bind dsDNA reversibly. (A) Schematic of single-molecule FRET

assay, with donor-labelled cTALE attached to a surface, and acceptor-labelled DNA free

in solution. (B-D) Single molecule FRET histograms show the appearance of a peak at

a FRET efficiency of 0.45 with increasing labelled DNA, consistent with a DNA-bound

cTALE. (E) Schematic of single-molecule FRET competition assay, with donor-labelled

cTALE attached to a surface, and acceptor-labelled DNA as well as competitor

unlabeled DNA free in solution. (F-H) Single molecule FRET histograms show the

disappearance of the peak at 0.45 FRET efficiency with increasing unlabeled competitor

DNA. Conditions: 20 mM Tris pH 8.0, 200 mM KCl.

Figure 3. Single Molecule (SM) kinetics show multiple phases in binding and unbinding kinetics. (A-B) Long time trajectories showing transitions between low- and high-FRET states (efficiencies of 0 and 0.45). The top panel shows calculated FRET efficiency in blue and two-state Hidden Markov Model fit in green[27]. The bottom panels show Cy3 and Cy5 fluorescence emission in green and red respectively. At low DNA

concentration (A), the low FRET state predominates. As DNA concentration is

increased (B), more time is spent in the high FRET state, because the dwell times in the

low FRET state are shorter. At low DNA concentrations, there appears to be long- and

short-lived high-FRET states. Likewise, at near-saturating DNA concentrations, there

appear to be long and short-lived low FRET states. (C, D) Cumulative distributions of

low- and high-FRET dwell times (blue circles). Fits to single-exponentials (black) show

large nonrandom residuals (lower panels), consistent with the heterogeneity noted in (A)

and (B). Double-exponentials (red) give smaller, more uniform residuals. (E) Apparent

association rate constants as a function of DNA concentration. The apparent rate

constants for the fast phase depend on DNA concentration (blue circles), indicating a

bimolecular step binding event. The apparent rate constants for the slow phase do not

depend on DNA-concentration (blue triangles), suggesting an isomerization event. (F)

Apparent dissociation rate constants as a function of DNA concentration (phase 1

shown in blue circles, and phase 2 shown in blue triangles). Neither phase shows a

DNA concentration dependence, indicating a dissociation and/or isomerization events.

Conditions: 20 mM Tris pH 8.0, 200 mM KCl.

Figure 4. A 16-repeat TALE protein binds and unbinds DNA more slowly than an

eight repeat protein. (A) Apparent association rate constants as a function of DNA

concentration for an 8 repeat cTALE (blue), a 12 repeat cTALE (black), and a 16 repeat

cTALE (green). 8 repeat TALE kinetics are measured by FRET ($FRET_{L \to H}$) while 12 and

16 repeat TALE kinetics are measured by colocalization ($E_{L \to H}$). The apparent rate

constants for the fast phase of binding are DNA concentration dependent (blue, black,

and green circles), indicating a bimolecular binding event. The DNA concentration-

dependence is greatest (larger slope) for the 8 repeat cTALE. The apparent rate

constants for the slow phase do not depend on DNA-concentration (blue, black, and

green triangles), suggesting an isomerization event. (B) Apparent dissociation rate

constants as a function of DNA concentration (phase 1 shown in circles, and phase 2

shown in triangles). Neither phase shows a DNA concentration dependence, indicating

a dissociation and/or isomerization events. Rate constants for all phases are slower for

the 12-repeat construct (black) and 16-repeat construct (green) than for the 8-repeat

construct (blue), particularly for the bimolecular binding step. (C) Log_{10} of rate constants

for 8 (blue), 12 (black), and 16(green) repeat cTALEs. Units of the bimolecular binding

rate constant are $nM^{-1}s^{-1}$, other unimolecular rate constants have units s^{-1}. Conditions:

20 mM Tris pH 8.0, 200 mM KCl.

Figure 5. Deterministic simulations provide evidence for conformational

heterogeneity in the unbound state. The model most consistent with data is shown at

the top. Unbound TALEs can exist in DNA-binding competent (TALE*) or DNA-binding

incompetent (TALE) states. DNA-bound TALEs can exist in short-lived (TALE*~DNA) or

long-lived (TALE‡~DNA) DNA-bound states. Cumulative distributions of dwell-times

(shown as blue points) from 8 repeat single-molecule time trajectories (A-F) and 16

repeat single-molecule time trajectories (G-L) were analyzed with the model (best-fit

shown in black). (A-C and G-I) Populations of states as a function of time, generated by

numerical integration in Matlab. (D-F and J-L) Cumulative distributions in blue circles

and best fit lines are shown in black. Best-fit microscopic rate constants and 68%

confidence intervals are listed in Table 1.

A

BINDING COMPETENT   ENCOUNTER

k_2
1.1 nM^{-1}s^{-1}

k_{-2}
0.66 s^{-1}

k_1
0.17 s^{-1}

k_{-1}
0.13 s^{-1}

k_{-3}
0.22 s^{-1}

k_3
0.36 s^{-1}

BINDING INCOMPETENT   LOCKED

B

BINDING COMPETENT   ENCOUNTER

k_2
0.31 nM^{-1}s^{-1}

k_{-2}
0.13 s^{-1}

k_1
0.135 s^{-1}

k_{-1}
1.26 s^{-1}

k_{-3}
0.0435 s^{-1}

k_3
0.043 s^{-1}

BINDING INCOMPETENT   LOCKED

C

BINDING COMPETENT   ENCOUNTER

k_2
0.39 nM^{-1}s^{-1}

k_{-2}
0.30 s^{-1}

k_1
0.26 s^{-1}

k_{-1}
0.43 s^{-1}

k_{-3}
0.078 s^{-1}

k_3
0.074 s^{-1}

BINDING INCOMPETENT   LOCKED

Figure 6. TALEs with multiple superhelical turns must break to bind DNA. Single-

molecule FRET studies and deterministic modeling support a model where TALEs exist

in four states: binding incompetent, binding competent, encounter complex, and locked.

In this model, for TALEs that form less than one full superhelical turn (8 repeats, A),

partly folded states are off-pathway and slow down binding. For longer TALEs that form

one (12, B) or more (16, C) complete superhelical turns, partial unfolding is required for

binding. DNA-bound TALEs form both encounter complexes and higher-affinity locked

conformations. Dynamics of long (12 and 16-repeat; B-C) TALEs bound to DNA are

significantly slower than for the shorter (8-repeat; A) TALE.

Table 1. Kinetic parameters obtained from deterministic simulation fits.

	k_1 (sec^{-1})	k_{-1} (sec^{-1})	$K_{eq, DNA-free}$	k_2 (sec^{-1}nM^{-1})	k_{-2} (sec^{-1})	k_3 (sec^{-1})	k_{-3} (sec^{-1})	$K_{eq, DNA-bound}$ (nM^{-1})
NcTALE$_8$[a]	0.17 [0.16, 0.18]	0.13 [0.12, 0.14]	1.32 [1.26, 1.39]	1.1 [1.08, 1.12]	0.66 [0.65, 0.67]	0.36 [0.35, 0.37]	0.222 [0.218, 0.227]	1.62 [1.58, 1.66]
NcTALE$_{12}$[b]	0.135 [0.133, 0.137]	1.26 [1.09, 1.34]	0.11 [0.10, 0.12]	0.31 [0.28, 0.33]	0.130 [0.129, 0.131]	0.043 [0.042, 0.044]	0.0435 [0.0428, 0.0442]	0.99 [0.97, 1.00]
NcTALE$_{16}$[a]	0.26 [0.25, 0.27]	0.43 [0.39, 0.47]	0.61 [0.57, 0.64]	0.39 [0.38, 0.41]	0.299 [0.298, 0.300]	0.074 [0.073, 0.076]	0.078 [0.076, 0.079]	0.96 [0.95, 0.97]

68% confidence intervals shown in brackets are from 2,000[a] and 8,000[b] iterations of bootstrap analysis.

Supplemental Material

Figure S1. Alternating laser experiments show agreement between cTALE₈ FRET

and colocalization kinetics. (A-C) Three representative time trajectories. In these

trajectories, excitation alternated between red and green with 250 msec at each color.

Top panels show detection of red and green light resulting from green laser excitation.

High red (and low green) fluorescence result from FRET. Lower panels show detection

of red light resulting from red laser excitation. There is strong correlation between

periods of high FRET in the upper panels (high red fluorescence, low green

fluorescence) and periods of high red fluorescence (from DNA binding) in the lower

panel. Conditions: 200 mM KCl, 20 mM Tris pH 8.0.

Figure S2. cTALEs do not slide onto ends of short dsDNA. Apparent association

and dissociation rate constants as a function of DNA concentration for an 8 repeat

cTALE array (NcTALE$_8$) binding to uncapped (blue) and capped (red) DNA, measured

by single molecule FRET dwell time analysis. (A) Rate constants for conversion from

the low to the high FRET state (FRET$_{L \to H}$). Rate constants for the faster phase increase

with DNA concentration (circles), indicating a bimolecular event. The DNA

concentration-dependence is stronger (larger slope, k=0.59 ± 0.08 nM^{-1} sec^{-1}) with

uncapped DNA than with capped DNA (k=0.09 ± 0.05 nM^{-1} sec^{-1}), which is likely a result

of the faster diffusion of small, uncapped DNA (10 kDa) compared to large, capped

DNA (320 kDa). Rate constants for the slower phase are not DNA-concentration

dependent (triangles). (B) Rate constants for conversion from the high to the low FRET

state (FRET$_{H \to L}$). Neither phase shows a DNA concentration dependent, indicating

unimolecular steps. Conditions: 200 mM KCl, 20 mM Tris pH 8.0.

Figure S3. Colocalization trajectories show TALE-DNA binding and unbinding

events. (A-C) One representative time trajectory and colocalization analysis for a

sixteen repeat TALE, NcTALE$_{16}$, incubated with 1 nM Cy5-A$_{15}$/T$_{15}$. In the colocalization

protocol, both the green and red lasers were used in the first ten frames (not shown) to

identify single molecule locations. For all subsequent frames, Cy5-labeled DNA was

continuously excited with the red laser. Red light emission images (from Cy5) were

collected with time steps of 100 msec. Trajectories were generated from sites of

red/green colocalization in the first ten frames, and were corrected for background Cy5

emission (red trajectories above). For each trajectory, we set two thresholds for low and

high Cy5 emission (C). A molecule is designated to be in the high emission (DNA-

bound) state above the high threshold, and in the low emission state below the low

threshold. From this state-assignment procedure, dwell-times were determined in the

697   high (bound) and low (unbound) states (C). Conditions: 200 mM KCl, 20 mM Tris pH

698   8.0.

Figure S4. Bound and unbound lifetimes of 16- and 12-repeat TALE proteins are consistent with multiphasic binding and unbinding. (A, B) Cumulative distributions of low-FRET dwell times (blue circles) for a 16-repeat construct at low and high DNA concentration. Fits to single-exponentials (black) show large nonrandom residuals (lower panels) which are most pronounced at high DNA concentrations, consistent with the binding heterogeneity observed in eight repeat cTALEs (Figure 3). Double-exponentials (red) give smaller, more uniform residuals. (C, D) Cumulative distributions of high-FRET dwell times (blue circles) for 16- and 12-repeat constructs at high DNA

709   concentration. Fits to single-exponentials (black) show large nonrandom residuals

710   (lower panels), consistent with the dissociation heterogeneity observed in eight repeat

711   cTALEs. Double-exponentials (red) give smaller, more uniform residuals.

Figure S5. Distance estimates between labeling sites for NcTALE$_8$ and NcTALE$_{16}$

and the 5' ends of bound DNA. The crystal structure of PthXo1 (PDB: 3UGM) [17] in

complex with DNA (colored by repeat) as a model of donor-acceptor fluorophore

distances . (A) The distance from C30 in the first repeat (the site of labelling in

NcTALE$_8$) to the 5' base of the sense strand (light grey) is 38.3Å; the distance to the 5'

base of the antisense strand (dark grey) is 37.4Å. (B) The distance from C30 in the

fourteenth repeat (the site of labelling in NcTALE$_{16}$) to the 5' base of the sense strand

(light grey) is 72.9Å; the distance to the 5' base of the antisense strand (dark grey) is

25.1Å. Note that although there is likely to be some variation these distances since the

DNAs used here are longer (15 and 23 bases) than the TALE arrays (8 and 16 repeats,

each with a four-repeat N-capping domain), the differences in donor-acceptor distances

for the 16 repeat construct (B) are likely to be robust to registry shifts of a few bases.

Figure S6. Urea and destabilizing mutations decrease apparent binding rate of

cTALE₈. (A) Apparent association rate constants as a function of DNA concentration for

an 8 repeat cTALE in 0 M urea (blue), 1 M urea (pink), and with destabilizing point

mutations (yellow; mutations substitute leucine at position 1 in the fourth repeat with a

glycine). The DNA concentration dependence is strongest (steepest slope) in the

absence of urea and destabilizing point substitutions (circles). Rates in the slower

phase (triangles) appear unaffected by urea or mutational destabilization. (B) Apparent

dissociation rate constants as a function of DNA concentration (phase 1 shown in

circles, and phase 2 shown in triangles). Rates in the both phases appear unaffected by

urea and mutational destabilization. Conditions: 200 mM KCl, 20 mM Tris pH 8.0.