1

High throughput single cell sequencing of both *T-cell-receptor-beta* alleles

Tomonori Hosoya[*,‡], Hongyang Li[†,‡], Chia-Jui Ku[*], Qingqing Wu[*], Yuanfang Guan[†] and

James Douglas Engel[*,§]

[*]Department of Cell and Developmental Biology

[†]Department of Computational Medicine and Bioinformatics

University of Michigan

3035 BSRB

109 Zina Pitcher Place

Ann Arbor, Michigan 48109-2200

[‡]These authors contributed equally to this work.

[§]Corresponding author

James Douglas Engel

3035 BSRB, 109 Zina Pitcher Place

Ann Arbor, MI   48109

Email: engel@umich.edu

Telephone: 734-647-0803

Running title: Validated sequencing of both *Trb* alleles in single cells

23 **ABSTRACT**

24 Allelic exclusion is a vital mechanism for the generation of monospecificity to foreign

25 antigens in B- and T-lymphocytes. Here we developed a high-throughput barcoded

26 method to simultaneously analyze the VDJ recombination status of both mouse *T cell*

27 *receptor beta* alleles in hundreds of single cells using Next Generation Sequencing.

INTRODUCTION

Vertebrates have evolved both innate and adaptive immune systems to protect

individuals against infection, cancer and invasion by parasites. B and T lymphocytes

comprise the central components of adaptive immunity and every individual lymphocyte

harbors specific reactivity to a single antigen that is conferred by individually unique

antigen receptors expressed on the cell surface. The diversity of antigen receptors is

generated by DNA recombination, called VDJ rearrangement in jawed vertebrates(1). To

maintain the required monospecificity of mature lymphocytes, only one of the two

autosomal alleles is allowed to express a functionally rearranged beta- and alpha- chain T

cell receptors (TCRβ and TCRα) in T cells, or immunoglobulin heavy and light chain

receptor (IgH and IgL) in B cells: both are controlled by a historically opaque mechanism

referred to as "allelic exclusion". Allelic exclusion occurs at the genetic level for IgH,

IgL and TCRβ(2), while at the level of protein localization on the cell surface for

TCRα(3). Loss of allelic exclusion results in dual-TCR expression, which can lead to

autoimmunity(4, 5).

A high-throughput method to study the diversity of lymphocytes at the single-cell

level has been reported recently(6), yet how one might analyze their mono-specificity

remains unclear. Since the abundance of transcripts from a non-functional allele is lower

than that of the functional allele(7, 8), traditional RNA-based methods cannot detect the

mechanisms underlying allelic exclusion. One major challenge in analyzing the genes

themselves is that there is only one copy of DNA representing each allele, unlike the

existence of multiple transcribed RNA species. Therefore, extremely accurate DNA

sequencing is required to avoid erroneously mis-assigning nucleotide-level mutations,

which would alter the interpretation of TCR locus activity. Additionally, no reference

genome is available for merging multiple sequencing reads, due to the many millions of

possible sequences that can be generated from VDJ rearrangement(9). Thus, *de novo*

sequence assembly or a long-read approach is required to retrieve the original genomic

sequence of both alleles in single cells. To significantly improve the traditional Sanger

sequencing method employed by us and others(10-12), we developed a high-throughput

method that enables analysis of *Trb* allelic exclusion status by sequencing both alleles of

the genome in single cells, enabling us to determine whether each allele in those cells

underwent either no, unproductive or productive rearrangement.

MATERIALS AND METHODS

Single cell isolation. Staged thymocytes were first isolated from mice (C57BL/6J,

Jackson Laboratories) between 5 and 8 weeks old using a cell sorter (BD FACSAria III).

Lin⁻CD4⁻CD8a⁻ Thy1.2⁺cKit⁻CD25⁺CD28⁻ (DN3a stage), Lin⁻CD3⁻TCRβ⁻ CD8a⁻

Thy1.2⁺cKit⁻CD25⁻ (DN4 stage), CD4⁺CD8⁺ (DP stage) and TCRβ⁺CD4⁺CD8⁺ (late DP

stage) thymocytes (10,000 to 100,000 cells at each stage) were isolated as described

previously(10). Next, single cells were directly sorted into 20 μl of lysis buffer

[containing 1x Q5® Reaction Buffer (NEB), 4 μg Proteinase K and 0.1% Triton X100] in

one well of a 96 well PCR plate using a Synergy cell sorter (Sony iCyt SY3200). The cell

sorter setting was carefully aligned so that sorted cells were precisely deposited into the

center of each well. Sorted single cells in the lysis solution were kept on ice, and then

digested at 55ºC for 60 minutes followed by 95ºC for 15 minutes (to inactivate Proteinase

K) using a PCR thermal cycler within 6 hours after sorting.

Multiplex nested PCR. For the first round of PCR, primers were selected to amplify all

potential VDJ rearrangement at the *Trb* locus: 31 V region primers covering all 35 *Trbv*

genes(13), 2 D region primers, 2 J region primers and 2 control primers used to detect

sequences 3' of the *Actb* gene (Figs. 1a and 1b). The full list of primers used are

deposited at (https://umich.box.com/s/3f5q64u2i68dn2i7hlneucxph9oetpvb). Since this

method was designed to recover not only rearranged but also germ line configured

genomic DNA, the two J primers were designed to be 3' of J1-7 and J2-7. The following

PCR condition amplifies the entire J1 (Jβ1 primer coupled with a V or D region primer)

as well as J2 genomic DNA region (Jβ2 primer coupled with a V or D segment primer).

Since VDJ will be spliced to a C region at the RNA level, the selected Jβ primer

sequences remain in genomic DNA after recombination. D region primers were designed

5' of D1 and D2. The D segment primers amplify only a D-to-J rearranged genome, but

not a V-to-DJ rearranged genome. After V-to-D1J rearrangement, the D1 primer

sequence is removed from the genome. Similarly after V-to-D2J2 rearrangement, the D2

primer sequence is removed from the genome. The first round of PCR was performed in a

60 μl final reaction volume containing of 50 nM of each primer, 1x Q5® Reaction

Buffer, 200 μM each dNTPs, 0.4 unit Q5® High-Fidelity DNA Polymerase (NEB) and

20 μl of the lysed single cell solution. The PCR condition was 30 sec at 98 ºC followed

by 30 cycles of 5 sec at 98 ºC, 10 sec at 66-58 ºC and 2 min at 72 ºC, and then final

extension for 2 min at 72 ºC. During the first 5 cycles, the annealing temperature was

reduced 2 ºC per cycle from 66 ºC to 58 ºC, and then performed at 56ºC for the last 25

cycles.

For the second round of PCR, nested primers were selected: 32 nested V region

primers containing forward adapter sequences (AF-Vβn), 2 nested D region forward

adapter primers (AF-Dβn) and 2 nested J region primers in a reverse adapter orientation

(AR3-Jβn). The second round of PCR was performed in a 20 μl final reaction volume

containing 50 nM of each primer (Fig. 1a), 1x Q5® buffer, 200 μM of each dNTP, 0.4

units Q5® High-Fidelity DNA Polymerase (NEB) and 1 μl of the first round PCR

product. The 2nd PCR condition was same as for the first round PCR but for only 25

cycles.

Finally, in the third round of PCR, unique barcoded-AF and barcoded-AR3

combinations of primers were selected for each single cell (Fig. 1a). Barcodes were

adopted from a published report(6). The 3rd PCR condition was same as for the second

round PCR (5 + 20 cycles). Hot start PCR was performed for all PCR reactions using

either a BioRad T100™ or Applied Biosystems 2720 Thermal Cycler. All primers were

purchased from Integrated DNA Technologies, Inc. The 37 (for 1st round) or 36 (for 2nd

round) PCR primers at 200 μM final concentrations were mixed, aliquoted and frozen for

subsequent use. Each PCR mix was prepared immediately before initiating PCR

reactions. (The amplification efficiency was reduced if the PCR mix was prepared in

advance and repeatedly frozen and thawed. Amplification efficiency was also reduced

when 500 nM of each primer was used in the 2nd round PCR reaction). Successful DNA

amplification was confirmed by running 2 μl of 3rd round PCR product on agarose gels

with 0.4 μg of 2-Log DNA Ladder (NEB) (Fig. 1c): 8 samples were selected from each

96 well PCR plate. Of note, the PCR conditions employed here allow amplification of

unrearranged (germline configuration) genomic DNA (Fig. 1c left panel).

Recovery frequency

To analyze the frequency of recovered wells, a primer pair amplifying the 3' region of the

Actb gene was added to the 1st round PCR. 1 μl of the 1st round PCR product was used

in a PCR reaction with nested *Actb* primers: 5'-CATAGGCTTCACACCTTCCT-3', 5'-

CTTTGCCTCCATCTGCATAAC-3' and FAM-labeled probe

TGCTAGTCTGAAGCTGCCCTTTCC (ZEN / Iowa Black FQ) purchased from

Integrated DNA Technologies, Inc. Luna Universal Probe qPCR Master Mix (NEB,

M3004) was used in a 20 μl reaction. The PCR condition was 30 sec at 95 ºC followed

by 45 cycles of 1 sec at 95 ºC and 20 sec at 60 ºC using StepOnePlus™ Real-Time PCR

System (Thermo Fisher Scientific). A 90-95% recovery efficiency was routinely obtained

from each plate containing 96 single cell sorted wells.

PacBio high-throughput sequencing

Since the length of the PCR products described above ranged from 200 to 2700 bp (Fig.

1b), a sequencer with the ability to read long DNA fragments is preferred. To this end,

Pacific Biosciences single molecule real-time (SMRT) sequencing technology(14) was

employed. A PacBio smartbell adapter was ligated to the barcoded PCR product mixture

and then sequenced on PacBio RSII sequencer at the University of Michigan Sequencing

Facility. Thus each circular consensus sequencing (CCS) read generated from PacBio

sequencing corresponds to a single molecule of initial PCR product (Fig. 2a).

PacBio raw reads analysis

To obtain DNA sequences amplified from genomic DNA in single cells, we developed an

in-house analysis pipeline (Fig. 2a; the full version is available at

https://github.com/Hongyang449/scVDJ_seq). To calculate CCS reads of inserts, PacBio

ConsensusTools software version 2.0.0 was used with the following settings: --

minFullPasses=2, --minPredictedAccuracy=90, --minLength=10 and default for other

parameters (https://github.com/PacificBiosciences/SMRT-

Analysis/wiki/Documentation).

First, to demultiplex CCS reads, we grouped the CCS reads based on the presence of

a barcode primer set flanking each end of the CCS. We obtained approximately 35,000

CCSs per PacBio RS II SMRTcell on average, and 46% of them started with a barcode-

AF primer and ended with a barcode-AR3 primer. The remaining 54% had mutations,

deletions, or additions within the barcoded primer sequence or bore truncated barcode

primer sequences, and were excluded from the analysis.

Next, we generated sub-groups in each single cell (unique barcode set) based on the

presence of a V or D primer sequence, and sequences in each sub-group were aligned

using multiple sequence alignment software MUSCLE(15)

(http://www.drive5.com/muscle/). For each group, hierarchical clustering based on

sequence length was performed to create initial clusters, then k-means clustering was

recursively performed on the DNA sequences of each initial cluster to generate the final

multiple clusters, until every nucleotide position in each cluster achieved >51% identity.

Clusters containing only 1 CCS read were excluded. Aberrant CCSs with two PCR

products ligated together probably resulted during the smartbell adapter ligation step, and

were also excluded from the analysis.

To obtain accurate genomic DNA sequence, we calculated a consensus sequence

from multiple CCSs in each cluster. Of note, each CCS file that was generated from

single zero-mode waveguide (ZMW) corresponds to one copy of PCR product. If a

mutation happened during the first PCR cycle, 50% of the final PCR products inherit the

mutation at a specific base position in the final PCR product. Similarly, a mutation

during the second cycle is inherited by 25% of the final PCR products, 12.5% after the

third cycle and so on. After adapter and J region genomic DNA sequences (except the J

region used for rearrangement) were removed, the cluster consensus with V primer

sequences were submitted to IMGT HighV-Quest analysis(16) (http://www.imgt.org/) to

determine whether a specific VDJ sequence in a cluster was predicted to encode a

productive or unproductive TCRβ protein. For example, when DNA sequences

rearranged with J1-3, J1-4 to J1-7 regions as well as 3' of J1-7 to J1 primer sequences

were removed. The removal of the extra J region genomic DNA is essential to accurately

determine the productivity of rearrangement, especially when some bases are truncated

from J regions.

Based on sequence and functionality predicted by IMGT analysis, each cluster

consensus sequence was categorized into a tag shown in Supplementary Fig. S1a. Since

the status of an allele (GL, DJ, uVDJ and pVDJ) can be determined from possible

combinations of 1-2 cluster tags (a total of 24 different patterns are shown in

Supplementary Fig. S1b), the rearrangement status from within a single cell (the

combination of two separate allelic patterns) was calculated from 2-4 tags recovered from

each cell (for a total 253 different combinations of tags). From those, the summary tables

were generated.

In this study, the rearrangement status of both alleles was obtained from 909 single

cells (25% of 3,664 wells) for thymocytes post beta-selection and 193 DN3a stage

thymocytes (25% of 784 wells) when V-to-DJ rearrangement takes place. The frequency

depends on sequence depth and was 40% when we recovered 29,643 CCSs from 576

wells. We obtained 0-4 clusters per single cell (1.62 clusters per cell on average) in this

depth of sequencing. PacBio CCS data, the barcode list for each sample, genomic DNA

and CDR3-IMGT amino acid sequences for all clusters generated from this study are

deposited at

(https://umich.box.com/s/3f5q64u2i68dn2i7hlneucxph9oetpvb).

RESULTS

High-throughput *Trb* genomic DNA sequencing from single cells

To sequence both alleles of genomic DNA in the *Trb* gene VDJ region, we employed

a multiplex strategy (Fig. 1a; details described in Methods). Since mouse thymocytes

were directly sorted into lysis buffer in individual wells of 96-well PCR plates.

Multiplex, nested PCR reactions followed by barcoding PCR amplified the germline (GL:

unrearranged configuration) genomic DNA, D-to-J rearranged (DJ) DNA, and all

potential V-to-DJ rearranged (VDJ) DNA in the *Trb* locus(13) (Figs. 1b and 1c). The

PCR products were mixed and then sequenced in a PacBio RS II long-read sequencer

(www.pacb.com). Each circular consensus specification (CCS) read of an insert

generated by the PacBio system corresponds to a single molecule of the original PCR

product (Fig. 2a). Based on barcoding and sequence similarity, multiple CCSs were

grouped into clusters (Fig. 2a). For each cluster, the consensus sequence was retrieved,

which corresponds to an original PCR product. The consensus sequences were

subsequently submitted to IMGT HighV-Quest analysis(16) (http://www.imgt.org/) to

determine whether the VDJ sequences in any cluster predicted that a functional TCRβ

protein could be generated in that cell. From one *Trb* allele, two [(V-)D1-J1 and (V31-

)D2-J2 rearrangement] or one [(V-)D-J2] PCR products are amplified (Fig. 1b and

Supplementary Fig. S1). Theoretically, between two and four PCR products (clusters) can

be recovered from two alleles in each cell. Details for the computational processing of the

PacBio output data to calculate allele rearrangement status are described in Materials and

Methods.

To calculate the PCR error rate under our experimental conditions, we took advantage

of the unrearranged (germline configuration) genomic DNA corresponding to parts of

Trbd1 to *Trbj1-7* and *Trbd2* to *Trbj2-7* (Fig. 1b) in single bone marrow (BM) cells. From

single BM cells we obtained two PCR bands (Fig. 1c), which correspond to the lengths of

germline DNA amplified using Dβ1 forward and Jβ1 reverse nested primers (2,672 bp)

as well as Dβ2 forward and Jβ2 reverse nested primers (1,896 bp). The accuracy of the

resultant sequences was 99.768% for each individual CCS, including both PCR and

PacBio sequencing errors. To increase the accuracy, we employed an approach to obtain

the consensus sequence from more than 2 CCSs. If we obtain only 2 CCSs per cluster, an

error at a position results in 50% identity at the position (ex. An A in 1 CCS and a G in

the other CCS); such a cluster was then removed because it is not >51% identical. If we

recover 3 CCSs, an error at any position results in a 66.6% identity at the position (e.g.

An A in 2 CCSs and a G in 1 CCS). In this case A is the consensus sequence at the

position in this cluster. When a consensus was generated from 3 randomly chosen

synonymous CCSs, the accuracy increased to 99.991%. These data demonstrate that this

method using PCR with Q5® High-Fidelity DNA polymerase (NEB) followed by

calculating the consensus sequence from multiple CCSs yields >99.99% accuracy, even

after a total of 80 PCR cycles.

Trb VDJ rearrangement status in single thymocytes

To precisely determine the genomic status at the *Trb* loci following VDJ

rearrangement in single cells, we analyzed wild type DN4, DP and late DP stage

thymocytes, in which *Trb* VDJ rearrangement had already been completed (i.e. post-β-

selection). Since DJ joining takes place before the DN3 stage on both alleles, all of the

cells analyzed here would be predicted to have undergone at least one productive VDJ

recombination event on one allele. We obtained a total of 5,877 clusters and assessed the

rearrangement status of both alleles from 909 single cells (Fig. 3). 52% (472) of the

single thymocytes were in a [pVDJ (productive VDJ rearranged)/DJ] configuration and

41% (377) were in a [uVDJ (unproductively VDJ rearranged)/pVDJ] configuration, as

expected from the 60/40-rule(2). 4% (37) of the post-β-selection thymocytes were in a

pVDJ/pVDJ arrangement, which was similar to the 3% dual-TCRβ bearing cells as

analyzed using cell surface antibodies(17). Stage-specific data is shown in

Supplementary Figs. S2 and S3. The V regions that were most frequently utilized for

recombination were V13-2 (Fig. 4 and Supplemental Fig. S4), which is also the most

frequently used in inflamed lung epitope specific CD8+ T cells(8). 6,415 V-to-DJ

rearranged genomic DNA and complementarity determining region 3 (CDR3) amino acid

sequences (4,258 productive and 2,158 unproductive) recovered from single thymocytes

in this analysis are available online (see Data Access for detail). Surprisingly, we

recovered D2-to-J1 rearranged sequences, although only rarely (Supplementary Fig. S2).

At the double negative 3a (DN3a) stage, when V-to-DJ rearrangement takes place, only

9% of the cells had generated a productive (pVDJ) allele [0.5% (pVDJ/GL), 7.2% had

generated a (pVDJ/DJ) recombinant and 1.5% were in a (uVDJ/pVDJ) configuration,

Supplementary Fig. S3], all of which probably represent cells that were about to develop

to the next (DN3b) stage.

Next, to determine whether the methods developed here can capture an artificial

allelic exclusion phenomenon, we analyzed late DP cells isolated from Vb8 transgenic

mice(18) (Tg^{Vb8}), which express a productively rearranged TCRβ protein; expression of

that transgene has been shown to repress endogenous *Trb* VDJ rearrangement(19). Out of

2,429 clusters generated, 722 clusters bore the DNA sequence of the Tg^{Vb8} transgene

(Supplementary Figs. S4a) and were removed from further analysis. We then determined

the rearrangement status of both alleles from 127 single cells (Supplementary Figs. S4b

and S4c). V-to-DJ rearrangement was not observed at the endogenous locus in 80% of

the thymocytes, in agreement with published observations(19). However, 20% of the

cells had either unproductively (13% uVDJ/DJ) or productively (6% pVDJ/DJ) escaped

transgene-suppressed allelic exclusion to allow V-to-DJ rearrangement on one of the

endogenous *Trb* loci. In the Tg^{Vb8}, D2-to-J2 rearrangement was observed less often than

D1-to-J1. This supports the idea that D2-to-J2 rearrangement was more frequently

repressed when compared to D1-to-J1 region recombination in the presence of a

functional TCRβ transgene, in agreement with published analysis of 9 T cell clones(19).

DISCUSSION

In summary, the method developed here provides an extremely accurate, rapid high-

throughput approach for the analysis of *Trb* genomic DNA status at the single cell level.

This strategy allowed us to analyze the rearrangement status in thousands of single cells,

which is more than 10-fold greater than the number from Sanger sequencing methods

previously employed by us and others(10-12) and requiring far less time. This strategy

would also be directly applicable to the analysis of the *Igl* and *Igh* genes (encoding

immunoglobulins in B cells), which are also regulated by recombination and allelic

exclusion at the genetic level(2).

REFERENCES

1.   Tonegawa, S. 1983. Somatic generation of antibody diversity. *Nature* 302: 575-

581.

2.   Mostoslavsky, R., F. W. Alt, and K. Rajewsky. 2004. The lingering enigma of the

allelic exclusion mechanism. *Cell* 118: 539-544.

3.   Alam, S. M., and N. R. Gascoigne. 1998. Posttranslational regulation of TCR

Valpha allelic exclusion during T cell differentiation. *J Immunol* 160: 3883-3890.

4.   Hinz, T., E. Weidmann, and D. Kabelitz. 2001. Dual TCR-expressing T

lymphocytes in health and disease. *Int Arch Allergy Immunol* 125: 16-20.

5.   Auger, J. L., S. Haasken, E. M. Steinert, and B. A. Binstadt. 2012. Incomplete

TCR-beta allelic exclusion accelerates spontaneous autoimmune arthritis in

K/BxN TCR transgenic mice. *Eur J Immunol* 42: 2354-2362.

6.   Han, A., J. Glanville, L. Hansmann, and M. M. Davis. 2014. Linking T-cell

receptor sequence to functional phenotype at the single-cell level. *Nat Biotechnol*

32: 684-692.

7.   Weischenfeldt, J., I. Damgaard, D. Bryder, K. Theilgaard-Monch, L. A. Thoren,

F. C. Nielsen, S. E. Jacobsen, C. Nerlov, and B. T. Porse. 2008. NMD is essential

for hematopoietic stem and progenitor cells and for eliminating by-products of

programmed DNA rearrangements. *Genes Dev* 22: 1381-1396.

8.   Dash, P., J. L. McClaren, T. H. Oguin, 3rd, W. Rothwell, B. Todd, M. Y. Morris,

J. Becksfort, C. Reynolds, S. A. Brown, P. C. Doherty, and P. G. Thomas. 2011.

Paired analysis of TCRalpha and TCRbeta chains at the single-cell level in mice.

J Clin Invest 121: 288-295.

9.   Robins, H. S., P. V. Campregher, S. K. Srivastava, A. Wacher, C. J. Turtle, O.

Kahsai, S. R. Riddell, E. H. Warren, and C. S. Carlson. 2009. Comprehensive

assessment of T-cell receptor beta-chain diversity in alphabeta T cells. *Blood* 114:

4099-4107.

10.   Ku, C. J., J. M. Sekiguchi, B. Panwar, Y. Guan, S. Takahashi, K. Yoh, I. Maillard,

T. Hosoya, and J. D. Engel. 2017. GATA3 Abundance Is a Critical Determinant

of T Cell Receptor beta Allelic Exclusion. *Mol Cell Biol* 37.

11.   Aifantis, I., J. Buer, H. von Boehmer, and O. Azogui. 1997. Essential role of the

pre-T cell receptor in allelic exclusion of the T cell receptor beta locus. *Immunity*

7: 601-607.

12.   Aifantis, I., V. I. Pivniouk, F. Gartner, J. Feinberg, W. Swat, F. W. Alt, H. von

Boehmer, and R. S. Geha. 1999. Allelic exclusion of the T cell receptor beta locus

requires the SH2 domain-containing leukocyte protein (SLP)-76 adaptor protein. *J*

Exp Med 190: 1093-1102.

13.   Bosc, N., and M. P. Lefranc. 2000. The mouse (Mus musculus) T cell receptor

beta variable (TRBV), diversity (TRBD) and joining (TRBJ) genes. *Exp Clin*

Immunogenet 17: 216-228.

14.   Roberts, R. J., M. O. Carneiro, and M. C. Schatz. 2013. The advantages of SMRT

sequencing. *Genome Biol* 14: 405.

15.   Edgar, R. C. 2004. MUSCLE: multiple sequence alignment with high accuracy

and high throughput. *Nucleic Acids Res* 32: 1792-1797.

16.   Li, S., M. P. Lefranc, J. J. Miles, E. Alamyar, V. Giudicelli, P. Duroux, J. D.

Freeman, V. D. Corbin, J. P. Scheerlinck, M. A. Frohman, P. U. Cameron, M.

Plebanski, B. Loveland, S. R. Burrows, A. T. Papenfuss, and E. J. Gowans. 2013.

IMGT/HighV QUEST paradigm for T cell receptor IMGT clonotype diversity and

next generation repertoire immunoprofiling. *Nat Commun* 4: 2333.

17.   Balomenos, D., R. S. Balderas, K. P. Mulvany, J. Kaye, D. H. Kono, and A. N.

Theofilopoulos. 1995. Incomplete T cell receptor V beta allelic exclusion and

dual V beta-expressing cells. *J Immunol* 155: 3308-3312.

18.   Shinkai, Y., S. Koyasu, K. Nakayama, K. M. Murphy, D. Y. Loh, E. L. Reinherz,

and F. W. Alt. 1993. Restoration of T cell development in RAG-2-deficient mice

by functional TCR transgenes. *Science* 259: 822-825.

19.   Uematsu, Y., S. Ryser, Z. Dembic, P. Borgulya, P. Krimpenfort, A. Berns, H. von

Boehmer, and M. Steinmetz. 1988. In transgenic mice the introduced functional T

cell receptor beta gene prevents expression of endogenous beta genes. *Cell* 52:

831-841.

375 **FOOTNOTES**

376 This work was supported by National Institute of Health Grant AI094642 (to T.H. and

377 J.D.E). The research was also supported in part by the University of Michigan

378 Comprehensive Cancer Center Support Grant (P30 CA046592) for use of the Flow

379 Cytometry and the Sequencing Cores at the University of Michigan.

AUTHOR CONTRIBUTIONS

T.H. designed the study, performed experiments, wrote computer script, analyzed the

data, and wrote the manuscript. H.L. designed the study, wrote computer script and

edited the paper. C.K. designed the study, performed experiments, analyzed the data and

edited the manuscript. Q.W performed experiments and edited the paper. Y.G. designed

the study and edited the paper. J.D.E. designed the study and edited the paper.

DISCLOSURE DECLARATION

The authors declare no competing interests.

FIGURE LEGENDS

Fig. 1. High-throughput *Trb* genomic DNA sequencing from single-cells. (a) Strategy

for multiplex PCR amplification of genomic DNA sequences at the *Trb* gene VDJ loci

from single cells. Final PCR products were mixed and then sequenced by PacBio RS II

sequencer. Vβn: V region primers. Dβn: D region primers. Jβn: J region primers. AF:

adapter forward. AR: adapter reverse. BC: barcode. Details are described in Methods. (**b**)

Schematic presentation of VDJ rearrangement at the *Trb* locus with simplified illustration

of multiplex primer location. Genomic DNA recombination events are initiated by

recombining Dβ (diversity) and Jβ (joining) segments on both chromosomes.

Subsequently, on only one chromosome, one of the Vβ (variable) segments is joined to

the previously rearranged DJ recombinant. Not only the selection of each VDJ segment,

but also multiple lengths of spacer sequence between V, D and J, generates an

incalculable number of VDJ sequence possibilities. mRNA splicing joins Cβ (constant)

segments to the rearranged VDJ DNA recombinants to generate a final TCRβ protein.

VDJ rearrangement that generates a stop codon or elongated transcript results in a

predictedly unproductive TCRβ. (**c**) Successful amplification of rearranged VDJ, DJ and

germline D-to-J regions were confirmed by agarose gel electrophoresis with NEB 2-log

DNA ladder in the last (left panel) and first (right panel) lanes.

Fig. 2. A schematic outline of the PacBio raw data processing. Details are described

in Methods.

Fig. 3. *Trb* VDJ rearrangement status in single thymocytes. (**a**) Cluster tags in a total

of 5,877 clusters recovered from single wild type thymocytes at DN4, DP and late DP

(post β-selection) stages. Each cluster consensus sequence was categorized into one tag

shown on the x-axis (see Supplementary Fig. S1 for detail). (**b**) Calculated *Trb* allele

status for 909 single wild type thymocytes at DN4, DP and late DP (post β-selection)

stages. Legend: GL: germline allele status; DJ: D-to-J recombined allele; pVDJ: V-to-DJ

recombined allele encoding a productive TCRβ protein; uVDJ: V-to-DJ recombined

allele encoding a predicted unproductive TCRβ. The data in (**b**) and (**c**) represent a

summary of 4 mice for DN4, 5 mice for DP and 4 mice for late DP stages, and details are

shown in Supplementary Figs. S2 and S3.

Fig. 4. V and J region usage. Frequency of each V region (a) and J region (b) utilization

was calculated out of 3,356 VDJ clusters recovered from wild type single thymocytes at

DN4, DP and late DP stages. Stage-specific data are shown in Supplementary Fig. S4.

Fig. 5. Endogenous *Trb* rearranged status in thymocytes of Vb8 transgenic mice.

Number of tags (a) for recovered PCR fragments and rearrangement status (b) calculated

from late DP stage single cells isolated from wild type (black) and Vb8 transgenic (Tg,

white) mice. Data represent the summary from 4 mice.

Figure 1

a Multiplex PCR from single cell

1st PCR

2nd PCR

3rd PCR

c single BM cells   single thymocytes

Figure 2

Figure 3

a

b

Figure 4

a **V region usage**

b **J region usage**

Figure 5

a

cluster tag	V x J1	D1 x J1	1 x 2	D2 x J2	V x J2
tag1		D1_germ_J1			
tag2				D2_germ_J2	
tag3		D1_x_J1			
tag4			D1_x_J2		
tag5				D2_x_J2	
tag6			D2_x_J1		
tag7 (V1 to V30)	uV_x_J1				
tag8 (V1 to V30)					uV_x_J2
tag9	uV31_x_J1				
tag10					uV31_x_J2
tag11 (V1 to V30)	pV_x_J1				
tag12 (V1 to V30)					pV_x_J2
tag13	pV31_x_J1				
tag14					pV31_x_J2

b

status	pattern	V x J1	D1 x J1	1 x 2	D2 x J2	V x J2
GL	g1	__	D1_germ_J1	__	D2_germ_J2	__
DJ	d1	__	D1_x_J1	__	D2_germ_J2	__
DJ	d2	__	D1_germ_J1	__	D2_x_J2	__
DJ	d3	__	D1_x_J1	__	D2_x_J2	__
DJ	d4	__	__	D1_x_J2	__	__
DJ	d5	__	__	D2_x_J1	__	__
uVDJ	u1	__	__	__	__	uV_x_J2
uVDJ	u2	uV_x_J1	__	__	D2_germ_J2	__
uVDJ	u3	uV_x_J1	__	__	D2_x_J2	__
uVDJ	u4	uV31_x_J1	__	__	__	__
uVDJ	u5	uV31_x_J2	__	__	__	__
uVDJ	u6	__	D1_germ_J1	__	__	uV31_x_J2
uVDJ	u7	__	D1_x_J1	__	__	uV31_x_J2
uVDJ	u8	uV_x_J1	__	__	__	uV31_x_J2
pVDJ	p1	__	__	__	__	pV_x_J2
pVDJ	p2	pV_x_J1	__	__	D2_germ_J2	__
pVDJ	p3	pV_x_J1	__	__	D2_x_J2	__
pVDJ	p4	pV31_x_J1	__	__	__	__
pVDJ	p5	pV31_x_J2	__	__	__	__
pVDJ	p6	__	D1_germ_J1	__	__	pV31_x_J2
pVDJ	p7	__	D1_x_J1	__	__	pV31_x_J2
pVDJ	p8	uV_x_J1	__	__	__	pV31_x_J2
pVDJ	p9	pV_x_J1	__	__	__	uV31_x_J2
pVDJ	p10	pV_x_J1	__	__	__	pV31_x_J2

Supplementary Figure S1

(**a**) Full list of cluster tags (PCR products). PCR fragments amplified using Dβ forward (D1 or D2) and Jβ reverse (J1 or J2) nested primers include both germline configuration (germ) and rearranged DNA (x). PCR fragments amplified using Vβ forward (V) and Jβ reverse (J1 or J2) nested primers encode either productive (p, green) or unproductive (u, red) TCRβ protein. (**b**) Full list of *Trb* gene status and patterns. Status of an allele are shown with possible combination of 1-2 PCR fragments. GL: germline configuration (patter g). DJ: D-to-J rearranged (pattern d). uVDJ: V-to-DJ rearranged DNA encoding unproductive TCRβ (pattern u). pVDJ: V-to-DJ rearranged DNA encoding productive TCRβ (pattern p). Rearrangement with *Trbv31* region (V31) increases the complexity. Full list of possible combination of PCR fragment tags per cell is found in tag_status.csv file in code.zip file. Combination of 2-4 tags used to identify allele status of each cell.

a

sampleID	D1 germ	D2 germ	D1 J1	D1 J2	D2 J2	D2 J1	uV J1	uV J2	uV31 J1	uV31 J2	pV J1	pV J2	pV31 J1	pV31 J2		total
DN3a_WT_mouse2	50	108	93	44	21	0	22	10	0	1	4	5	0	0		358
DN3a_WT_mouse3	18	65	93	49	28	1	29	37	1	1	10	2	1	0		335
DN3a_WT_mouse4	21	70	87	52	23	0	26	17	2	0	5	7	0	0		310
DN3a_WT_mouse6	31	78	86	31	24	1	25	16	1	3	12	9	0	0		317
DN4_WT_mouse62	0	16	47	26	48	0	19	28	4	1	59	107	2	4		361
DN4_WT_mouse64	0	28	69	39	58	0	32	36	0	0	66	98	3	3		432
DN4_WT_mouse67	0	18	70	30	63	0	34	44	0	1	50	118	1	3		432
DN4_WT_mouse72	1	36	59	39	53	0	35	41	2	0	71	104	1	5		447
DP_WT_mouseWT	0	91	167	89	163	0	81	99	4	5	169	239	13	5		1125
DP_WT_mouse2	2	34	57	25	37	0	22	19	1	1	59	64	4	2		327
DP_WT_mouse3	0	26	35	34	35	0	22	15	1	0	54	53	4	0		279
DP_WT_mouse4	0	29	60	24	39	0	11	18	0	2	60	57	1	6		307
DP_WT_mouse6	0	46	48	34	50	0	21	24	0	0	59	75	3	0		360
lateDP_WT_mouse20	0	27	67	42	63	0	25	46	0	0	76	111	6	9		472
lateDP_WT_mouse25	1	18	58	29	52	0	41	53	1	2	53	143	1	11		463
lateDP_WT_mouse57	0	16	44	30	57	0	31	36	3	2	44	108	5	3		379
lateDP_WT_mouse77	0	18	54	38	81	0	37	48	0	4	67	139	1	6		493

count	D1 germ	D2 germ	D1 J1	D1 J2	D2 J2	D2 J1	uV J1	uV J2	uV31 J1	uV31 J2	pV J1	pV J2	pV31 J1	pV31 J2		total
DN3a	120	321	359	176	96	2	102	80	4	5	31	23	1	0		1320
DN4	1	98	245	134	222	0	120	149	6	2	246	427	7	15		1672
DP	2	226	367	206	324	0	157	175	6	8	401	488	25	13		2398
late DP	1	79	223	139	253	0	134	183	4	8	240	501	13	29		1807

%	D1 germ	D2 germ	D1 J1	D1 J2	D2 J2	D2 J1	uV J1	uV J2	uV31 J1	uV31 J2	pV J1	pV J2	pV31 J1	pV31 J2		total
DN3a	9%	24%	27%	13%	7%	0%	8%	6%	0%	0%	2%	2%	0%	0%		
DN4	0%	6%	15%	8%	13%	0%	7%	9%	0%	0%	15%	26%	0%	1%		
DP	0%	9%	15%	9%	14%	0%	7%	7%	0%	0%	17%	20%	1%	1%		
late DP	0%	4%	12%	8%	14%	0%	7%	10%	0%	0%	13%	28%	1%	2%		

b

Supplementary Figure S2

Tags for PCR fragments. Number (**a**) and frequency (**b**) of cluster tags recovered from wild type single thymocytes at DN3a, DN4, DP and late DP stages.

a

sampleID	GL GL	DJ GL	DJ DJ	uVDJ GL	pVDJ GL	uVDJ DJ	pVDJ DJ	uVDJ uVDJ	pVDJ uVDJ	pVDJ pVDJ	ID12 to 15	total
DN3a_WT_mouse2	33	10	15	0	0	6	4	2	1	0	0	71
DN3a_WT_mouse3	11	1	10	1	0	16	3	3	0	0	0	45
DN3a_WT_mouse4	8	7	12	0	0	7	3	3	0	0	0	40
DN3a_WT_mouse6	16	2	7	0	1	4	4	2	2	0	0	38
DN4_WT_mouse62	0	0	0	0	0	1	20	0	17	3	0	41
DN4_WT_mouse64	0	0	2	0	0	1	26	0	30	3	0	62
DN4_WT_mouse67	0	0	0	0	0	1	28	1	24	2	0	56
DN4_WT_mouse72	0	0	1	0	0	1	34	1	31	1	0	69
DP_WT_mouseWT	0	0	1	0	0	2	122	3	96	9	0	233
DP_WT_mouse2	1	0	1	0	0	0	29	0	15	2	1	49
DP_WT_mouse3	0	0	0	0	0	0	21	0	8	1	0	30
DP_WT_mouse4	0	0	0	0	0	1	36	0	11	0	0	48
DP_WT_mouse6	0	0	0	0	0	0	37	0	21	2	0	60
lateDP_WT_mouse20	0	0	0	0	0	0	38	0	27	1	0	66
lateDP_WT_mouse25	1	0	0	0	0	0	25	1	37	3	0	67
lateDP_WT_mouse57	0	0	0	0	0	0	18	0	27	1	0	46
lateDP_WT_mouse77	0	0	0	0	0	1	38	1	33	9	0	82

count	GL GL	DJ GL	DJ DJ	uVDJ GL	pVDJ GL	uVDJ DJ	pVDJ DJ	uVDJ uVDJ	pVDJ uVDJ	pVDJ pVDJ	ID12 to 15	total
DN3a	68	20	44	1	1	33	14	10	3	0	0	194
DN4	0	0	3	0	0	4	108	2	102	9	0	228
DP	1	0	2	0	0	3	245	3	151	14	1	420
late DP	1	0	0	0	0	1	119	2	124	14	0	261

%	GL GL	DJ GL	DJ DJ	uVDJ GL	pVDJ GL	uVDJ DJ	pVDJ DJ	uVDJ uVDJ	pVDJ uVDJ	pVDJ pVDJ	ID12 to 15	total
DN3a	35%	10%	23%	1%	1%	17%	7%	5%	2%	0%	0%	
DN4	0%	0%	1%	0%	0%	2%	47%	1%	45%	4%	0%	
DP	0%	0%	0%	0%	0%	1%	58%	1%	36%	3%	0%	
late DP	0%	0%	0%	0%	0%	0%	46%	1%	48%	5%	0%	

b

Supplementary Figure S3

VDJ rearrangement status of both *Trb* alleles. *Trb* VDJ rearrangement status for both alleles calculated from wild type single thymocytes at DN3a, DN4, DP and late DP stages. Number (**a**) and frequency (**b**) are shown.

Supplementary Figure S4

V gene segment usage in wild type single thymocytes at DN4, DP and late DP stages.
Frequency was calculated from 971 (DN4), 1,273 (DP) and 1,112 (late DP4) VDJ sequence recovered for each stage.