Title: Bidirectional Long-Term Synaptic Zinc Plasticity at Mouse Glutamatergic

Synapses

Running title: Mechanisms of Synaptic Zinc Plasticity

Authors: Nathan W. Vogler[1] and Thanos Tzounopoulos[*1]

[1]Pittsburgh Hearing Research Center and Department of Otolaryngology, University of

Pittsburgh, Pittsburgh, PA 15261

*Correspondence: thanos@pitt.edu

Keywords: zinc; auditory synapses; synaptic plasticity

Table of Contents category: Neuroscience

Length: 45 pages, plus 7 figures

Funding and Acknowledgements. This work was supported by NIH grants F31-

DC015924 (to N.W.V.) and R01-DC007905 (to T.T.). We thank Drs. Stephen Lippard and

Jacob Goldberg from Massachusetts Institute of Technology for generously providing DA-

ZP1. We thank Patrick Cody and Dr. Ross Williamson for technical help with data analysis.

We thank Dr. Elias Aizenman for helpful discussions. The authors declare no competing

financial interests.

Key points summary:

- Synaptic zinc is coreleased with glutamate to modulate neurotransmission and
auditory processing. Sensory experience causes long-term changes in synaptic
zinc signaling, termed synaptic zinc plasticity.

- At zinc-containing glutamatergic synapses in the dorsal cochlear nucleus (DCN),
we show that high-frequency stimulation reduces synaptic zinc signaling (Z-LTD),
whereas low-frequency stimulation increases synaptic zinc signaling (Z-LTP).

- Group 1 metabotropic glutamate receptor (mGluR) activation is necessary and
sufficient to induce Z-LTP and Z-LTD. Z-LTP and Z-LTD are associated with
bidirectional changes in presynaptic zinc levels.

- Sound-induced Z-LTD at DCN synapses requires Group 1 mGluR activation.

- Bidirectional synaptic zinc plasticity is a previously unknown mechanism of LTP
and LTD at zinc-containing glutamatergic synapses.

Abstract

Synaptic zinc is coreleased with glutamate to modulate neurotransmission in many excitatory synapses. In the auditory cortex, synaptic zinc modulates sound frequency tuning and enhances frequency discrimination acuity. In auditory, visual, and somatosensory circuits, sensory experience causes long-term changes in synaptic zinc levels and/or signaling, termed here *synaptic zinc plasticity*. However, the mechanisms underlying synaptic zinc plasticity and the effects of this plasticity on long-term glutamatergic plasticity remain unknown. To study these mechanisms, we used male and female mice and employed *in vitro* and *in vivo* models in zinc-rich, glutamatergic dorsal cochlear nucleus (DCN) parallel fiber (PF) synapses. High-frequency stimulation of DCN PF synapses induced long-term depression of synaptic zinc signaling (Z-LTD), as evidenced by reduced zinc-mediated inhibition of AMPA receptor (AMPAR) excitatory postsynaptic currents (EPSCs). Low-frequency stimulation induced long-term potentiation of synaptic zinc signaling (Z-LTP), as evidenced by enhanced zinc-mediated inhibition of AMPAR EPSCs. Thus, Z-LTD is a new mechanism of LTP and Z-LTD is a new mechanism of LTP. Pharmacological inhibition of Group 1 metabotropic glutamate receptors (G1 mGluRs) eliminated Z-LTD and Z-LTP. Pharmacological activation of G1 mGluRs induced Z-LTD and Z-LTP, associated with bidirectional changes in presynaptic zinc levels. Finally, exposure of mice to loud sound caused G1 mGluR-dependent Z-LTD in DCN PF synapses, consistent with our *in vitro* results. Together, we show that G1 mGluR activation is necessary and sufficient for inducing bidirectional long-term synaptic zinc plasticity.

Introduction

In many brain areas, including the neocortex, limbic structures, and the auditory brainstem, glutamatergic vesicles are loaded with zinc (Danscher & Stoltenberg, 2005; Frederickson *et al.*, 2005). This pool of mobile, synaptic zinc is coreleased with glutamate. Synaptically released zinc inhibits synaptic and extrasynaptic NMDA receptor (NMDAR) EPSCs, and modulates AMPA receptor (AMPAR) EPSCs (Vogt *et al.*, 2000; Vergnano *et al.*, 2014; Anderson *et al.*, 2015; Kalappa *et al.*, 2015; Kalappa & Tzounopoulos, 2017). Namely, synaptic zinc inhibits AMPAR EPSCs during baseline synaptic activity via postsynaptic mechanisms, but enhances steady-state AMPAR EPSCs during higher frequencies of synaptic stimulation (Kalappa *et al.*, 2015; Kalappa & Tzounopoulos, 2017). The enhancing effect of synaptic zinc on AMPAR EPSCs is short-lasting and is mediated by short-term, zinc-mediated changes in presynaptic glutamatergic neurotransmission (Perez-Rosello *et al.*, 2013; Kalappa & Tzounopoulos, 2017). Thus, synaptic zinc is a major modulator of baseline neurotransmission and short-term plasticity of glutamatergic synapses.

In awake mice, synaptic zinc enhances the responsiveness (gain) of auditory cortical principal neurons to sound, but reduces the gain of cortical interneurons (Anderson *et al.*, 2017). Furthermore, synaptic zinc sharpens the sound frequency tuning of auditory cortical principal neurons, and enhances frequency discrimination acuity (Kumar *et al.*, 2019). Sensory experience bidirectionally modulates the levels of vesicular zinc and synaptic zinc signaling in several sensory brain areas (Nakashima & Dyck, 2009; Kalappa *et al.*, 2015; Li *et al.*, 2017; McAllister & Dyck, 2017). In the somatosensory cortex, whisker plucking increases zinc levels, whereas whisker stimulation reduces zinc levels (Brown & Dyck, 2002, 2005). In the primary visual cortex, monocular deprivation increases vesicular zinc

levels (Dyck *et al.*, 2003). In the retina, optic nerve damage increases zinc levels, which in

turn inhibit optic nerve regeneration and promote cell death (Li *et al.*, 2017). In the dorsal

cochlear nucleus (DCN), an auditory brainstem nucleus, exposure to loud sound reduces

vesicular zinc levels and synaptic zinc signaling (Kalappa *et al.*, 2015). Yet, the cellular

and molecular mechanisms underlying the long-term experience-dependent plasticity of

synaptic zinc signaling, termed here *synaptic zinc plasticity*, and the relationship of

synaptic zinc plasticity to long-term glutamatergic synaptic plasticity remain unknown.

Elucidating these mechanisms is crucial for understanding how the brain adapts during

normal sensory processing, and why it fails to properly adjust in sensory disorders

associated with pathological central adaptation, such as in tinnitus (Auerbach *et al.*, 2014).

To determine the mechanisms of long-term synaptic zinc plasticity and its effects on LTP

and LTD, we developed *in vitro* and *in vivo* models. Namely, we used electrophysiology,

pharmacology, and fluorescent imaging in the DCN, which contains granule cell endings,

parallel fibers (PFs), with high levels of synaptic zinc (Frederickson *et al.*, 1988; Rubio &

Juiz, 1998; Kalappa *et al.*, 2015). We investigated these mechanisms *in vitro*, in response

to electrical synaptic activation that induces synaptic plasticity such as LTP and LTD in

brain slices, as well as *in vivo*, in response to loud sound exposure. Our results

demonstrate that bidirectional activity-dependent synaptic zinc plasticity is a previously

unknown, Group 1 mGluR-dependent mechanism of LTP and LTD at zinc-containing

glutamatergic synapses.

Materials and Methods

Animals. Male or female ICR mice (Envigo) were used in this study, aged between postnatal day 17 (P17) to P28. All animal procedures were approved by the Institutional Animal Care and Use Committee of the University of Pittsburgh, Pittsburgh, PA.

Brain slice preparation. Mice were deeply anesthetized with isoflurane (3% in O_2), then immediately decapitated and their brains were removed. Brain slices were prepared in artificial cerebrospinal fluid (ACSF, 34°C) containing the following (in mM): 130 NaCl, 3 KCl, 1.2 $CaCl_2$·$2H_2O$, 1.3 $MgCl_2$·$6H_2O$, 20 $NaHCO_3$, 3 HEPES, and 10 D-Glucose, saturated with 95% O_2/5% CO_2 (vol/vol), pH = 7.25-7.35, ~300 mOsm. Using a Vibratome (VT1200S; Leica), coronal brain slices (210 μm thickness) containing the left dorsal cochlear nucleus (DCN) were cut, then placed in a chamber containing warm (34°C) ACSF, and incubated for 60 min at 34°C, then room temperature (no longer than 3 hours) before beginning electrophysiology experiments. Incubating ACSF was the same as cutting ACSF, except it was stirred with Chelex 100 resin (Bio-Rad) for 1 hour to remove contaminating zinc, then filtered using Nalgene rapid flow filters lined with polyethersulfone (0.2 µm pore size). After filtering, high purity $CaCl_2$·$2H_2O$ and $MgCl_2$·$6H_2O$ (99.995%; Sigma Aldrich) were added. All plastic and glassware used for these experiments were washed with 5% nitric acid.

Electrophysiology. *Whole-cell recordings*. DCN slices were transferred to the recording chamber and perfused with ACSF (1-2 mL/min), maintained at ~34°C using an inline heating system (Warner Instruments). Recording ACSF was the same as incubating ACSF (see above), except it contained 2.4 mM $CaCl_2$·$2H_2O$. Whole-cell recordings from cartwheel cells were performed using glass micropipettes (3-6 MΩ; Sutter Instruments).

Cartwheel cells were identified by the presence of complex spikes in cell-attached

configuration before break-in or in response to current injections in current-clamp mode

after break-in (Zhang & Oertel, 1993; Manis *et al.*, 1994; Tzounopoulos *et al.*, 2004).

Recording pipettes were filled with a potassium-based internal solution (except for Figure

6, see below) containing the following (in mM): 113 K-gluconate, 4.5 $MgCl_2 \cdot 6H_2O$, 14 Tris-

phosphocreatine, 9 HEPES, 0.1 EGTA, 4 Na_2ATP, 0.3 Tris-GTP, and 10 sucrose (pH =

7.25, 295 mOsm). For experiments shown in Figure 6 measuring NMDAR EPSCs,

recordings were performed using a cesium-based internal solution containing the following

(in mM): 128 $Cs(CH_3O_3S)$, 10 HEPES, 4 $MgCl_2 \cdot 6H_2O$, 4 Na_2ATP, 0.3 Tris-GTP, 10 Tris-

phosphocreatine, 1 EGTA, 1 QX-314, and 3 Na-ascorbate (pH = 7.25, 300 mOsm).

Voltages were not corrected for junction potentials. Recordings were performed using

ephus (Suter *et al.*, 2010) and a MultiClamp 700B amplifier (Axon Instruments). Data were

sampled at 10 kHz and low-pass-filtered at 4 kHz. Series resistance (R_s) and input

resistance (R_m) were monitored during the recording period by delivering -5 mV voltage

steps for 50 ms. R_s was calculated by dividing the -5 mV voltage step by the peak current

generated immediately after the voltage step. R_m was calculated by dividing the -5 mV

voltage step by the difference between the baseline and steady-state hyperpolarized

current, then subtracting R_s. Data were excluded if R_s or R_m changed by more than 20%

from the baseline period. EPSCs were evoked using an Isoflex stimulator (A.M.P.I., 0.1 ms

pulses) through a glass ACSF-containing theta electrode to stimulate the zinc-rich parallel

fibers. All EPSCs were recorded in the presence of SR95531 (20 μM, $GABA_AR$ antagonist)

and strychnine (1 μM, GlyR antagonist). AMPAR EPSCs were recorded in voltage-clamp

mode at -70 mV. For paired-pulse experiments, the inter-stimulus interval was 50 ms.

NMDAR EPSCs were evoked by a 5-pulse stimulus train (20 Hz) (Anderson *et al.*, 2015),

recorded in voltage clamp mode at +40 mV, and in the presence of DNQX (20 μM,

AMPA/kainate receptor antagonist). All drugs were always bath applied.

Induction of plasticity. High-frequency stimulation (HFS) consisted of 3 trains of 100

Hz pulses for 1 sec, with 10 sec between trains. For the experiments shown in Figure 1 C,

a subset of cells (n=5) were depolarized to -10 mV during each HFS train, while the other

subset (n=6) were held at -70 mV during HFS. Because we observed no difference in the

zinc plasticity (% potentiation by ZX1) between these subsets (depolarized = 2.68 ±

5.36%, non-depolarized = 8.76 ± 8.64%, *p* = 0.58, unpaired *t* test), they were grouped

together for subsequent analysis. ZX1 (100 μM) is a fast, high-affinity extracellular zinc

chelator (Anderson *et al.*, 2015; Kalappa *et al.*, 2015; Kalappa & Tzounopoulos, 2017). For

all other experiments, cells were voltage-clamped at -70 mV during HFS. For experiments

measuring NMDAR EPSCs after HFS (Figure 6), DNQX (20 μM) was added after HFS,

then cells were voltage-clamped at +40 mV to record NMDAR EPSCs. For ifenprodil

experiments (Figure 6 D-E), ZX1 was applied prior to ifenprodil to chelate extracellular

zinc, because zinc affects NMDAR ifenprodil sensitivity (Hansen *et al.*, 2014). In these

experiments after HFS, ZX1 was applied with DNQX, after the HFS. Low-frequency

stimulation (LFS) consisted of 5 Hz pulses for 3 min. During LFS, cells were held at -80

mV in current-clamp mode. To isolate mGluR-mediated plasticity, all LFS experiments

were performed in the presence of APV (50 μM, NMDAR antagonist), and with external

ACSF containing 4 mM $CaCl_2 \cdot 2H_2O$ and 4 mM $MgCl_2 \cdot 6H_2O$ (Oliet *et al.*, 1997). The

interleaved experiments shown in Figure 4 D, examining the effect of 50 μM DHPG

application, were also performed in these conditions. For normalized EPSCs (% baseline),

EPSC amplitudes were normalized to the average EPSC amplitude during the 5 min

baseline period before HFS/LFS, DHPG, ifenprodil, or ZX1 application. To quantify ZX1

potentiation after HFS/LFS or DHPG application, EPSC amplitudes were renormalized to

the average EPSC amplitude of the new baseline period 5 min before ZX1 application.

ZX1 potentiation (shown in bar graphs) was quantified as the percent increase in the

average EPSC amplitude during the last 5 min of ZX1 application compared to the 5 min

baseline period before ZX1 application.

Vesicular zinc imaging with DA-ZP1. After preparation and incubation of DCN slices

(described above), slices were transferred to the imaging chamber and perfused with

recirculating ACSF (2-3 mL/min) maintained at ~34°C. Imaging of presynaptic vesicular

zinc levels in DCN parallel fibers was performed using DA-ZP1, a high-affinity, membrane

permeable fluorescent zinc sensor (Zastrow *et al.*, 2016). DA-ZP1 (0.5-1.0 µM) was added

to the ACSF, and allowed to incubate for at least 20 min before imaging. Images were

acquired using an upright microscope (Olympus BX5) with epifluorescence optics through

a 20x water immersion objective (Olympus). Green fluorescent signals were isolated using

a Pinkel filter set (Semrock LF488/543/625-3X-A-000) in response to excitation by an

ephus-driven blue LED (M470L2; Thorlabs), and images were acquired using a CCD

camera (Retiga 2000R, QImaging). Images consisted of 20 frames captured at 0.067 Hz

which were then averaged together and analyzed in MATLAB (Mathworks). The DCN

molecular layer, which contains the vesicular zinc-rich parallel fibers, extends ~75 µm

deep from the ependymal surface, while deeper layers lack vesicular zinc (zinc-free

region) (Ryugo & Willard, 1985; Frederickson *et al.*, 1988; Rubio & Juiz, 1998). Thus, DA-

ZP1 produces a band of fluorescence within the molecular layer near the ependymal

surface, consistent with the distribution of zinc-rich parallel fiber terminals (Frederickson *et*

al., 1988; Kalappa *et al.*, 2015; Zastrow *et al.*, 2016). The DA-ZP1 fluorescence band is

absent in ZnT3 KO mice lacking vesicular zinc, indicating that it specifically labels vesicular

zinc (Kalappa *et al.*, 2015; Zastrow *et al.*, 2016). To control for slice-to-slice variability in

the molecular layer volume, which in turn might lead to variability in DA-ZP1 brightness, we

compared DA-ZP1 fluorescence in the same region of the same slice before and after

DHPG application (Figure 5). DA-ZP1 fluorescence 15-20 min after DHPG application was

normalized to baseline fluorescence before DHPG application. To quantify DA-ZP1

fluorescence, we quantified two ROIs within each slice: one within the zinc-containing

molecular layer (zinc ROI) and the other within the zinc-free region (zinc-free ROI)

(Kalappa *et al.*, 2015; Zastrow *et al.*, 2016). Because the DCN molecular layer is curved

along the ependymal surface, to define the zinc ROI, we used a MATLAB routine to

automatically detect the abrupt increase in fluorescence intensity between the background

and the ependymal surface of the slice. Then the zinc ROI was automatically selected to

include 50 μm depth from the ependymal surface, consistent with the extent of the zinc-

containing parallel fiber terminals (Frederickson *et al.*, 1988). The length of the ROI was

450 μm. The zinc-free ROI was identical to the zinc ROI, except located 200-250 μm from

the border of the slice, within the zinc-free region (deep or fusiform cell layers) (Ryugo &

Willard, 1985; Frederickson *et al.*, 1988). Thus, all ROIs contained the same cross-

sectional area. The automatically generated ROI borders are shown with yellow lines in

Figure 5. Fluorescence intensity was averaged within each ROI, and the zinc-sensitive

fluorescence was calculated by subtracting the zinc-free ROI fluorescence from the zinc

ROI fluorescence.

Noise exposure. Noise exposure was performed based on previously published methods

(Kalappa *et al.*, 2015). Sham- or noise-exposed mice were anesthetized using 3%

isoflurane during induction and 1-1.5% during maintenance. Noise-exposed mice were

exposed for 4 hours to narrow bandpass noise at 116 dB sound pressure level (SPL),

248 centered at 16 kHz with a 1.6 kHz bandwidth. Noise was presented unilaterally (left ear)

249 through a pipette tip inserted into the left ear canal, with the other end attached to a

250 calibrated speaker (CF-1; Tucker Davis Technologies). Insertion of the pipette tip into the

251 ear canal did not produce a seal. Sham-exposed mice underwent an identical procedure

252 except without any noise exposure. For mice given intraperitoneal injections of AIDA (2

253 mg/kg), one injection was given 30 min prior to exposure, and a second injection was

254 given 2 hours later. After noise- or sham-exposure, ABRs were collected and mice

255 recovered from anesthesia, then DCN slices were prepared (within 30 min after exposure).

257 **ABRs.** Auditory Brainstem Responses (ABRs) were measured based on previously

258 published methods (Kalappa *et al.*, 2015). ABRs were recorded immediately after noise- or

259 sham-exposure. During ABR measurements, mice were anesthetized using 3% isoflurane

260 during induction and 1-1.5% during maintenance. Mice were placed in a sound attenuating

261 chamber and temperature was maintained at ~37°C using a heating pad. A subdermal

262 electrode was placed at the vertex, the ground electrode placed ventral to the right pinna,

263 and the reference electrode placed ventral to the left pinna (sham- or noise-exposed ear).

264 In noise-exposed mice, because no ABRs were detected when recording from the

265 exposed (ipsilateral) ear, we recorded ABRs from the non-exposed (contralateral) ear

266 (Figure 7 D). For ABR measurements from contralateral ears of noise-exposed mice, the

267 reference electrode was placed ventral to the right pinna (contralateral ear) and the ground

268 electrode placed ventral to the left pinna. ABRs were detected in response to 1 ms click

269 sound stimuli, presented through a pipette tip inserted into the ear canal, with the other

270 end attached to the speaker (CF-1; Tucker Davis Technologies). ABRs were recorded in

271 response to clicks presented in 10 dB steps, ranging from 0-80 dB SPL. 1 ms clicks were

272 presented at a rate of 18.56/sec using System 3 software package from Tucker Davis

Technologies, and ABRs were averaged 512 times and filtered using a 300-3,000 Hz

bandpass filter. ABR threshold was defined as the lowest stimulus intensity which

generated a reliable Wave 1 in the response waveform. Wave 1 amplitude was measured

as the peak-to-trough amplitude of the first wave in the ABR waveform (latency ~2 ms), in

response to 80 dB SPL clicks.

Drugs. All chemicals used for ACSF and internal solutions were purchased from Sigma-

Aldrich. The following drugs were purchased from HelloBio: SR95531 hydrobromide, DL-

AP5, DNQX disodium salt, ifenprodil, MPEP hydrochloride, LY367385, and (S)-3,5-

Dihydroxyphenylglycine (DHPG). Strychnine hydrochloride was purchased from Abcam.

(RS)-1-Aminoindan-1,5-dicarboxylic acid (AIDA) was purchased from Tocris. ZX1 was

purchased from STREM Chemicals.

Statistical Analysis

All data analysis was performed using Matlab (Mathworks), Excel (Microsoft), or Prism 7

(GraphPad). For statistical tests for normalized data, or within groups, we used one-

sample t tests (for normally distributed data) or Wilcoxon signed rank tests (for non-

normally distributed data). Data were considered normally distributed if they passed the

Shapiro-Wilk normality test. For comparisons between two (normally distributed) groups,

we used unpaired t tests. All t tests were two-tailed. For comparisons between three

groups, we used ordinary one-way ANOVA with Bonferroni's multiple comparisons test (for

normally distributed data), or Kruskal-Wallis test with Dunn's multiple comparisons test (for

non-normally distributed data). IC_{50} was calculated using the Hill equation by fitting the

dose-response curve with a nonlinear least squares fit. The IC_{50} of each fit was compared

using the extra sum-of-squares F test. Significance levels are defined as *p* < 0.05. Group

data are presented as mean ± SEM.

Detailed values and statistical tests for Figures. Figure 1: **(1C)** EPSC % baseline after HFS

(average of mins. 19-23): 115.1 ± 6.43%, n=11, t=2.34 df=10, *p=0.041, one-sample t test

vs. 100%. **(1D)** ZX1 potentiation (%): 'Control': 34.47 ± 5.7%, n=10, t=6.049 df=9,

*p=0.0002, one-sample t test vs. 0%. 'HFS': 6.0 ± 5.15%, n=11, t=1.165 df=10, n.s.

p=0.27, one-sample t test vs. 0%. 'Control' vs. 'HFS': t=3.719 df=19, *p=0.0015, unpaired t

test.

Figure 2: **(2A)** EPSC % baseline after HFS (average of mins. 19-23): 124.8 ± 4.76%, n=9,

t=5.198 df=8, *p=0.0008, one-sample t test vs. 100%. **(2B)** EPSC % baseline after HFS

(average of mins. 19-23): 126.2 ± 9.54%, n=6, t=2.743 df=5, *p=0.041, one-sample t test

vs. 100%. One cell was included for analysis of EPSCs following HFS, but did not remain

stable throughout subsequent ZX1 application and was excluded from analysis following

ZX1 application, quantified in C. **(2C)** ZX1 potentiation (%): 'HFS + APV': 4.28 ± 6.08%,

n=9, t=0.7039 df=8, n.s. p=0.502, one-sample t test vs. 0%. 'HFS + LY367385, MPEP,

APV': 36.07 ± 9.05%, n=5, t=3.987 df=4, *p=0.016, one-sample t test vs. 0%. One-way

ANOVA: F= 7.737, *p=0.003. 'Control' vs. 'HFS + APV': *p=0.0038; 'HFS + APV' vs. 'HFS

+ LY367385, MPEP, APV': *p= 0.0115; Bonferroni's multiple comparisons test.

Figure 3: **(3A)** EPSC % baseline after LFS (average of mins. 19-23): 95.97 ± 3.5%, n=8,

t=1.155 df=7, n.s. p=0.29, one-sample t test vs. 100%. Two cells were included for

analysis of EPSCs following LFS, but did not remain stable throughout subsequent ZX1

application and were excluded from analysis following ZX1 application, quantified in C.

(3B) EPSC % baseline after LFS (average of mins. 20-24): 73.67 ± 5.8%, n=6, t=4.528

df=5, *p=0.006, one-sample t test vs. 100%. **(3C)** ZX1 potentiation (%): 'Control': 19.65 ±

4.3%, n=5, t=4.567 df=4, *p=0.01, one-sample t test vs. 0%. 'LFS': 57.86 ± 12.4%, n=6,

t=4.681 df=5, *p=0.005, one-sample t test vs. 0%. 'LFS + LY367385, MPEP': 22.18 ±

8.3%, n=6, t=2.663 df=5, *p=0.04, one-sample t test vs. 0%. One-way ANOVA: F=5.257,

*p=0.0198. 'Control' vs. 'LFS': *p=0.0276; 'LFS' vs. 'LFS + LY367385, MPEP': *p=0.0309;

Bonferroni's multiple comparisons test.

Figure 4: **(4A)** EPSC % baseline after 50 μM DHPG (average of mins. 16-20): 49.66 ±

5.1%, n=6, t=9.945 df=5, *p=0.0002, one-sample t test vs. 100%. One cell was included for

analysis of EPSCs following 50 μM DHPG, but did not remain stable throughout

subsequent ZX1 application and was excluded from analysis following ZX1 application,

quantified in C. **(4B)** EPSC % baseline after 5 μM DHPG (average of mins. 16-20): 82.41 ±

7.4%, n=5, t=2.376 df=4, n.s. p=0.076, one-sample t test vs. 100%. **(4C)** ZX1 potentiation

(%): 'DHPG (50 μM)': 93.51 ± 10.92%, n=5, t=8.561 df=4, *p=0.001, one-sample t test vs.

0%. 'DHPG (5 μM)': 0.44 ± 7.08%, n=5, t=0.06273 df=4, n.s. p=0.95, one-sample t test vs.

0%. One-way ANOVA: F=30.22, *p<0.0001. 'Control' vs. 'DHPG (50 μM)': *p<0.0001;

'Control' vs. 'DHPG (5 μM)': *p=0.01; Bonferroni's multiple comparisons test. **(4D)** EPSC %

baseline after 50 μM DHPG (average of mins. 16-20): 83.05 ± 5.9%, n=5, t=2.893 df=4,

*p=0.044, one-sample t test vs. 100%. **(4E)** EPSC % baseline after LFS and 50 μM DHPG

(average of mins. 20-24): 81.43 ± 9.1%, n=5, t=2.051 df=4, n.s. p=0.11, one-sample t test

vs. 100%. **(4F)** ZX1 potentiation (%): 'DHPG (50 μM)': 55.83 ± 17.9%, n=5, t=3.12 df=4,

*p=0.036, one-sample t test vs. 0%. 'LFS + DHPG (50 μM)': 74.65 ± 17.6%, n=5, t=4.246

df=4, *p=0.013, one-sample t test vs. 0%. One-way ANOVA: F=0.4126, n.s. p=0.6703.

'LFS + DHPG (50 μM)' vs. 'LFS': n.s. p=0.9181; 'LFS + DHPG (50 μM)' vs. 'DHPG (50

μM)': n.s. p=0.8553; Bonferroni's multiple comparisons test.

Figure 5: **(5C)** DA-ZP1 fluorescence (% control): '+ DHPG (50 μM)': 132.3 ± 9.096%, n=9,

*p=0.0039, Wilcoxon signed rank test vs. 100%. '+ DHPG (5 μM)': 68.73 ± 11.99%, n=8,

*p=0.0078, Wilcoxon signed rank test vs. 100%.

Figure 6: **(6C)** ZX1 potentiation (%): 'Control': 37.1 ± 3.1%, n=5, t=12.06 df=4, *p=0.0003,

one-sample t test vs. 0%. 'HFS': 9.2 ± 5.2%, n=6, n.s. p=0.16, Wilcoxon signed rank test

vs. 0%. 'HFS + LY267385, MPEP': 42.6 ± 8.2%, n=5, t=5.197 df=4, *p=0.007, one-sample

t test vs. 0%. Kruskal-Wallis test: *p=0.0102. 'Control' vs. 'HFS': *p=0.0242; 'HFS' vs. 'HFS

+ LY367385, MPEP': *p=0.0428; Dunn's multiple comparisons test. **(6D)** EPSC (%

baseline): 'Control': 300nM: n=3, 90.68 ± 1.051%; 1μM: n=5, 70.89 ± 3.943%; 3μM: n=5,

54.61 ± 2.791%; 10μM: n=3, 40.72 ± 4.845%. 'HFS': 300nM: n=3, 90.24 ± 4.327%; 1μM:

n=4, 69.52 ± 2.208%; 3μM: n=4, 52.1 ± 3.214%; 10μM: n=3, 37.89 ± 1.533%. Nonlinear

fits: 'Control': Hill Slope=1.095, R^2=0.9472. 'HFS': Hill Slope=1.128, R^2=0.9765. **(6E)** IC_{50}

(μM): 'Control': 1.284 ± 0.3566. 'HFS': 1.267 ± 0.2321. Extra sum-of-squares F test: n.s.

p=0.9687.

Figure 7: **(7B)** ZX1 potentiation (%): 'N.E.': 11.7 ± 8.56%, n=5, t=1.373 df=4, n.s. p=0.24,

one-sample t test vs. 0%. 'N.E. + AIDA': 43.8 ± 8.05%, n=6, t=5.447 df=5, *p=0.003, one-

sample t test vs. 0%. 'N.E.' vs. 'N.E. + AIDA': t=2.724 df=9, *p=0.024, unpaired t test. **(7C)**

PPR: 'N.E.': 1.896 ± 0.19, n=5. 'N.E. + AIDA: 2.056 ± 0.12, n=6. 'N.E.' vs. 'N.E. + AIDA':

t=0.7446 df=9, n.s. p=0.476, unpaired t test. Normalized $1/CV^2$: 'N.E. + AIDA': 0.78 ± 0.22,

n=6; n.s. p=0.44, Wilcoxon signed rank test vs. 1. **(7E)** ABR threshold (dB SPL): 'Sham

ipsi.': 43.75 ± 3.24, n=8. 'N.E. contra.': 68.33 ± 3.07, n=6. 'N.E. + AIDA contra.': 65.71 ±

2.97, n=7. Kruskal-Wallis test: *p=0.0002. 'Sham ipsi.' vs. 'N.E. contra.': *p=0.0042; 'Sham

ipsi.' vs. 'N.E. + AIDA contra.': *p=0.0076; 'N.E. contra.' vs. 'N.E. + AIDA contra.': n.s.

p>0.9999; Dunn's multiple comparisons test. ABR Wave I (μV): 'Sham ipsi.': 2.67 ± 0.31,

n=8. 'N.E. contra.': 1.23 ± 0.13, n=6. 'N.E. + AIDA contra.': 1.25 ± 0.27, n=7. Kruskal-

Wallis test: *p=0.0024. 'Sham ipsi.' vs. 'N.E. contra.': *p= 0.0387; 'Sham ipsi.' vs. 'N.E. +

AIDA contra.': *p=0.0107; 'N.E. contra.' vs. 'N.E. + AIDA contra.': n.s. p>0.9999; Dunn's

multiple comparisons test.

Results

Bidirectional activity-dependent long-term synaptic zinc plasticity requires Group 1

mGluR activation. To investigate the mechanisms underlying synaptic zinc plasticity, we

first determined whether we could induce long-term synaptic zinc plasticity in DCN PF

synapses in mouse brain slices. In these synapses, synaptic zinc inhibits AMPAR and

NMDAR EPSCs via postsynaptic mechanisms. This has been evidenced by application of

ZX1, a fast, high-affinity extracellular zinc chelator, which potentiates AMPAR and NMDAR

EPSCs (Anderson *et al.*, 2015; Kalappa *et al.*, 2015; Kalappa & Tzounopoulos, 2017). This

ZX1 potentiation of AMPA and NMDAR EPSCs is dependent on ZnT3, the transporter that

loads zinc into synaptic vesicles (Palmiter *et al.*, 1996; Cole *et al.*, 1999). Moreover,

reductions in ZX1 potentiation reflect reductions in synaptic zinc levels and release

(Kalappa *et al.*, 2015). Therefore, in this study, we used the amount of ZX1 potentiation of

AMPAR and NMDAR EPSCs to monitor synaptic zinc signaling, and long-term synaptic

zinc plasticity, in DCN PF synapses.

Consistent with previous studies, we found that ZX1 potentiated postsynaptic PF AMPAR

EPSCs in DCN cartwheel cells (CWCs), a class of inhibitory interneurons (Figure 1 A-B)

(Kalappa *et al.*, 2015; Kalappa & Tzounopoulos, 2017). We then tested whether we can

induce long-term synaptic zinc plasticity by using patterns of synaptic activation that induce

long-term plasticity of glutamatergic synaptic strength in DCN PF synapses, such as LTP

and LTD (Fujino & Oertel, 2003; Tzounopoulos *et al.*, 2004; Tzounopoulos *et al.*,

2007). We started by examining the effect of ZX1 on AMPAR EPSCs following high-

frequency stimulation of PFs (HFS, 3 x 100 Hz for 1 sec, 10 sec inter-stimulus interval),

which induces LTP (Fujino & Oertel, 2003). After applying HFS and inducing LTP (Figure 1

C), we renormalized AMPAR EPSC amplitude to quantify the amount of ZX1 potentiation.

(Figure 1 C). After HFS, ZX1 application did not potentiate AMPAR EPSCs (Figure 1 C-D).

The loss of ZX1 potentiation indicates a loss of zinc-mediated inhibition of AMPARs,

suggesting that HFS caused a long-term reduction in synaptic zinc signaling, termed *Z-*

LTD (Figure 1 D). Furthermore, these results suggest that Z-LTD, by reducing zinc-

mediated inhibition of AMPAR EPSCs and thus enhancing baseline synaptic strength, is a

new mechanism of HFS-induced LTP. For a discussion on the impact of synaptic zinc

plasticity in the context of other long-term plasticity mechanisms, see *Discussion* section,

Implications of Z-LTP and Z-LTD for LTD and LTP.

After establishing that HFS caused Z-LTD, we then studied the underlying mechanisms.

NMDARs contribute to the induction of LTP and LTD in the DCN and most central

synapses (Malenka & Nicoll, 1993; Fujino & Oertel, 2003; Tzounopoulos *et al.*, 2004;

Tzounopoulos *et al.*, 2007). To test the role of NMDARs in the induction of Z-LTD, we

blocked NMDARs with APV (NMDAR antagonist, 50 μM; Figure 2 A). As evidenced by the

lack of ZX1 potentiation of AMPAR EPSCs after HFS, APV did not affect Z-LTD (Figure 2

A, C), indicating that NMDARs are not required for the induction of Z-LTD.

Parallel fiber synapses in the DCN also exhibit glutamatergic plasticity that involves

metabotropic glutamate receptor (mGluR) signaling (Fujino & Oertel, 2003). Furthermore,

Group 1 (G1) mGluRs are expressed in CWCs and in the DCN molecular layer, where PF

terminals reside (Wright *et al.*, 1996; Bilak & Morest, 1998). We therefore tested whether

G1 mGluR activation is necessary for Z-LTD. To test this hypothesis, we repeated the

experiment shown in Figure 2A, but we now blocked G1 mGluRs with MPEP (4 μM,

mGluR5-selective antagonist) and LY367385 (100 μM, mGluR1-selective antagonist)

(Figure 2 B). Under these conditions, ZX1 potentiation was observed after HFS, indicating

that HFS did not induce Z-LTD (Figure 2 B-C). This result demonstrates that G1 mGluR

activation is necessary for the induction of Z-LTD.

Glutamatergic plasticity is bidirectional: synapses undergo LTP or LTD in response to high-

or low-frequency stimulation, respectively (Mulkey & Malenka, 1992; Malenka & Nicoll,

1993; Fujino & Oertel, 2003). To determine whether long-term synaptic zinc plasticity is

bidirectional, we tested whether low-frequency stimulation (LFS) increases zinc signaling

(Figure 3 A). Because the induction of zinc plasticity depends on mGluR activation, we

used conditions that favor mGluR-dependent LTD, such as LFS (5 Hz, 3 min), blockade of

NMDARs with APV, and high extracellular concentrations of divalent ions (4 mM Ca^{2+} and

Mg^{2+}) (Oliet *et al.*, 1997). Compared to interleaved control experiments, LFS increased the

amount of subsequent ZX1 potentiation (Figure 3 A, C). Increased ZX1 potentiation

indicates increased zinc-mediated inhibition of AMPARs, suggesting that LFS caused a

long-term increase in synaptic zinc signaling, termed *Z-LTP* (Figure 3 C). By enhancing

zinc-mediated inhibition of AMPAR EPSCs, Z-LTP is a new mechanism of LFS-induced

LTD.

Note that control ZX1 potentiation in these conditions (Figure 3 A, C) was slightly less,

albeit not significantly different (p=0.11, unpaired t test), than previous control experiments

performed in ACSF with 2.4/1.3 mM of extracellular Ca^{2+}/Mg^{2+} (Figure 1 D). This is likely

due to reduced neuronal excitability in higher divalent concentrations (Oliet et $al.$, 1997;

Kalappa et $al.$, 2015). Together, these results show that LFS induced Z-LTP, thus

demonstrating that activity-dependent plasticity of zinc signaling is bidirectional: HFS

induces long-term depression of zinc signaling (Z-LTD), whereas LFS induces long-term

potentiation of zinc signaling (Z-LTP).

We next tested whether G1 mGluR activation is necessary for the induction of Z-LTP. In

the presence of MPEP and LY367385, LFS did not increase the amount of ZX1

potentiation (Figure 3 B-C), indicating that G1 mGluR activation is necessary for the

induction of Z-LTP. Together, these results reveal that activation of G1 mGluR signaling is

necessary for the induction of Z-LTP and Z-LTD.

Group 1 mGluR activation is sufficient to induce bidirectional long-term synaptic

zinc plasticity. Is activation of G1 mGluRs sufficient to induce Z-LTP and Z-LTD?

Because G1 mGluRs are required for both increases and decreases in synaptic zinc

signaling by different stimulation paradigms, we hypothesized that the direction of plasticity

depends on the differential activation of G1 mGluRs during HFS and LFS. To test this, we

applied high or low concentrations of DHPG (G1 mGluR agonist, 50 μM or 5 μM).

Consistent with previous studies, application of 50 μM DHPG caused a significant

depression of synaptic strength (Figure 4 A) (Huber et $al.$, 2001; Snyder et $al.$, 2001;

Wisniewski & Car, 2002). After applying 50 μM DHPG, obtaining a new stable baseline,

and then applying ZX1, we observed that the ZX1 potentiation of EPSCs was significantly

increased compared to control experiments (Figure 4 A, C). This result indicates that a

high concentration of DHPG increases synaptic zinc signaling: G1 mGluR activation is

sufficient to induce Z-LTP.

Because Z-LTP and Z-LTD induced by LFS and HFS depend on G1 mGluR activation

(Figure 2 and 3), we next tested whether application of a lower concentration of DHPG

causes Z-LTD. After applying 5 μM DHPG and obtaining a new stable baseline, ZX1 did

not potentiate EPSCs, consistent with Z-LTD induction (Figure 4 B-C). Together, these

results demonstrate that G1 mGluR activation is sufficient to cause bidirectional zinc

plasticity. Furthermore, the direction of zinc plasticity depends on the concentration of

DHPG: 50 μM DHPG causes Z-LTP, whereas 5 μM DHPG causes Z-LTD (Figure 4 C).

These results are consistent with the notion that bidirectional zinc plasticity depends on

differential activation of G1 mGluRs by either LFS/HFS or high/low concentrations of

DHPG.

Electrical synaptic stimulation with LFS/HFS or pharmacological activation of G1 mGluRs

with high/low concentrations of DHPG induce bidirectional synaptic zinc plasticity;

however, it is unknown whether these two different methods induce mechanistically similar

synaptic zinc plasticity. To explore this, we compared the amount of Z-LTP elicited by

applying sequential LFS and 50 μM DHPG to the amount of Z-LTP elicited by LFS or 50

μM DHPG alone. If electrical and pharmacological manipulations induce Z-LTP by different

mechanisms, then LFS and 50 μM DHPG application should yield an additive effect on Z-

LTP, and subsequent ZX1 potentiation should be greater than that following LFS alone or

application of 50 μM DHPG alone. To test this, we performed interleaved experiments to

determine the effect of 50 μM DHPG alone, under the conditions used for LFS-induced Z-

LTP as in Figure 3, with experiments involving stimulation with LFS and subsequent DHPG

application (Figure 4 D-E). Under these conditions, ZX1 potentiation following application

of 50 μM DHPG was similar to ZX1 potentiation following LFS (Figure 4 F). Importantly,

ZX1 potentiation after sequential LFS and 50 μM DHPG was not significantly greater than

ZX1 potentiation after LFS or DHPG alone (Figure 4 F). Together, these results show that

LFS occluded the effect of 50 μM DHPG; thus, LFS and DHPG induce Z-LTP likely via a

common mechanistic pathway.

Group 1 mGluR activation bidirectionally modulates presynaptic zinc levels. We

used activity-dependent changes in the amount of ZX1 potentiation of AMPAR EPSCs for

assessing changes in synaptic zinc signaling (Z-LTP and Z-LTD). However, ZX1

potentiation is determined by the postsynaptic zinc-mediated inhibition of AMPAR EPSCs,

as well as the amount of presynaptic zinc release (Kalappa *et al.*, 2015). Because previous

studies demonstrated sensory experience-dependent, long-term modulation of presynaptic

zinc levels (Nakashima & Dyck, 2009; Kalappa *et al.*, 2015), we hypothesized that Z-LTP

and Z-LTD are expressed, at least in part, by the modulation of presynaptic zinc levels. To

quantify potential changes in presynaptic zinc levels, we used DA-ZP1, a fluorescent

intracellular zinc sensor capable of tracking presynaptic zinc levels in PF terminals

(Kalappa *et al.*, 2015; Zastrow *et al.*, 2016). DA-ZP1 produces a band of fluorescence

within the DCN molecular layer in wild type mice. This fluorescent signal is absent in mice

lacking the vesicular ZnT3 transporter, thus demonstrating that the signal is due to ZnT3-

dependent, synaptic zinc (Kalappa *et al.*, 2015; Zastrow *et al.*, 2016). To induce Z-LTP and

Z-LTD, we applied DHPG, which is mechanistically similar to electrically-induced Z-LTP

and Z-LTD (Figure 4 F) and capable of inducing robust synaptic zinc plasticity in many

terminals in the slice. To test for changes in presynaptic zinc levels, we imaged DA-ZP1

fluorescence in the same region of the same DCN slice before and after DHPG application

(50 μM or 5 μM) (Figure 5 A; see *Materials and Methods*). Application of 50 μM DHPG

increased DA-ZP1 fluorescence, indicating increased presynaptic zinc levels in PF

terminals, which is consistent with Z-LTP (Figure 5 A, C). In contrast, application of 5 μM

DHPG reduced DA-ZP1 fluorescence, indicating reduced zinc levels, which is consistent

with Z-LTD (Figure 5 B-C).

Together, these results demonstrate that differential activation of G1 mGluRs, by

application of different concentrations of DHPG, causes bidirectional modulation of

presynaptic zinc levels. Furthermore, these results are consistent with our

electrophysiological experiments: 50 μM DHPG results in Z-LTP by increasing presynaptic

zinc levels, whereas 5 μM DHPG results in Z-LTD by reducing presynaptic zinc levels.

Although these results do not rule out potential postsynaptic mechanisms of Z-LTP and Z-

LTD, they demonstrate that Z-LTP and Z-LTD are associated with modulation of

presynaptic zinc levels.

G1 mGluR-dependent Z-LTD reduces zinc-mediated inhibition of NMDARs. Z-LTP

and Z-LTD involve modulation of presynaptic zinc signaling (Figure 5). Based on this

finding, the induction of long-term synaptic zinc plasticity should also affect postsynaptic

NMDAR EPSCs, which are inhibited by zinc via direct high-affinity NMDAR allosteric

modulation (Paoletti *et al.*, 1997; Vergnano *et al.*, 2014). To test this prediction, we

quantified the ZX1 potentiation of NMDAR EPSCs after inducing Z-LTD with HFS. To

monitor NMDAR EPSCs, we used a short train of presynaptic stimulation (5 pulses at 20

Hz) to activate extrasynaptic NMDARs, for NMDAR EPSCs recorded in somata of CWCs

are mostly mediated by extrasynaptic NMDARs activated by glutamate spillover during this

short train (Anderson *et al.*, 2015). To avoid keeping CWCs at +40 mV for too long while

recording NMDAR EPSCs, and to maintain the same induction protocol used in our

previous experiments, we initially recorded AMPAR EPSCs at -70 mV and then applied

HFS (Figure 6 A). Subsequently, we blocked AMPARs with DNQX (20 μM, AMPA/kainate

receptor antagonist) and recorded at +40 mV to obtain a stable baseline of NMDAR

EPSCs before applying ZX1 (Figure 6 A). Consistent with our results on AMPAR EPSCs,

after HFS, ZX1 no longer potentiated NMDAR EPSCs, whereas ZX1 potentiated NMDAR

EPSCs in interleaved control experiments, where HFS was not applied (Figure 6 A, C).

These results demonstrate that Z-LTD reduces zinc-mediated inhibition of NMDARs. To

determine whether this plasticity shares the same mechanism as Z-LTD evidenced by

changes in the ZX1 potentiation of AMPAR EPSCs, we tested whether G1 mGluR

signaling is required. Indeed, application of MPEP and LY367385 blocked the observed Z-

LTD, evidenced by the ZX1 potentiation of NMDAR EPSCs after HFS (Figure 6 B-C).

Together, our results suggest that G1 mGluR-dependent synaptic zinc plasticity modulates

zinc-mediated inhibition of AMPARs and NMDARs similarly, suggesting that it is

independent of the mode of action of synaptic zinc on its postsynaptic targets. This

supports our findings that zinc plasticity is expressed, at least in part, by changes in

presynaptic zinc levels.

However, the contribution of postsynaptic mechanisms in synaptic zinc plasticity cannot be

excluded. To address this possibility, we tested whether activity-dependent changes in

postsynaptic NMDAR subunit composition could modulate zinc sensitivity. NMDARs are

composed of two GluN1 subunits and two GluN2 subunits (Traynelis *et al.*, 2010). GluN2A-

containing   NMDARs   (GluN1/GluN2A   diheteromers   and   GluN2/GluN2A/GluN2B

triheteromers) have nanomolar affinity for zinc, whereas GluN1/GluN2B diheteromers have

micromolar affinity (Paoletti *et al.*, 1997; Rachline *et al.*, 2005; Tovar & Westbrook, 2012;

Hansen *et al.*, 2014). Therefore, the reduced zinc-mediated inhibition of NMDAR EPSCs

after HFS, evidenced by reduced ZX1 potentiation (Figure 6 A), could be explained by an

increase in the proportion of GluN2B subunits. We therefore tested whether HFS increases

the sensitivity of NMDAR EPSCs to ifenprodil, a GluN2B-selective antagonist (Figure 6 D-

E) (Tovar & Westbrook, 2012; Hansen *et al.*, 2014). Compared to controls, HFS did not

affect the ifenprodil sensitivity (IC_{50}) of NMDAR EPSCs (Figure 6 D-E). This indicates that

HFS-induced plasticity does not alter the proportions of GluN2B vs. GluN2A NMDAR

subunits, suggesting that Z-LTD is not due to reduced zinc sensitivity caused by a

decrease in the relative contribution of GluN2A vs. GluN2B in the NMDAR EPSC.

Therefore, these results further support that zinc plasticity is expressed by changes in

presynaptic zinc levels, rather than postsynaptic receptor modifications.

Sound-induced zinc plasticity requires Group 1 mGluRs *in vivo*. Our experiments

described here, using *in vitro* brain slice electrophysiology in the DCN point toward a

mechanism of bidirectional long-term synaptic zinc plasticity dependent on G1 mGluR

activation. We therefore hypothesized that G1 mGluR activation may also be necessary for

the reduction in synaptic zinc signaling observed in the DCN after sound exposure

(Kalappa *et al.*, 2015). To test this hypothesis, we quantified the ZX1 potentiation of PF

EPSCs in DCN slices from mice exposed to loud sound (116 dB, 4 hours). Consistent with

sound-induced LTD and previous studies (Kalappa *et al.*, 2015), we did not observe ZX1

potentiation in slices from noise-exposed (N.E.) mice (Figure 7 A-B). To test whether G1

mGluRs are necessary for the reduced zinc signaling in slices from N.E. mice, we

administered a systemic, blood brain barrier-permeable G1 mGluR antagonist (AIDA, i.p.,

2 mg/kg; twice: 30 min before and 1.5 hours after beginning the noise exposure). Indeed,

we observed ZX1 potentiation in slices from N.E. mice treated with AIDA (Figure 7 A-B),

suggesting that *in vivo* inhibition of G1 mGluR activity blocked the sound-induced Z-LTD.

Although AIDA treatment blocked Z-LTD in DCN PF synapses (Figure 7 A-B), it did not

affect assays that are sensitive to presynaptic glutamate release probability, such as

paired-pulse ratio (PPR) and coefficient of variation (CV) analysis (Figure 7 C). This

indicates that sound-induced G1 mGluR-dependent Z-LTD specifically modulates synaptic

zinc signaling, without affecting presynaptic glutamate signaling in PFs. Furthermore, AIDA

treatment did not affect sound-induced hearing loss in N.E. mice, quantified with Auditory

Brainstem Responses (ABRs) (Figure 7 D). ABRs reflect the synchronous activity, arising

from the auditory nerve (Wave I), of auditory brainstem nuclei to the inferior colliculus

(Waves II-V) in response to sound stimuli. Elevated ABR thresholds indicate increased

hearing thresholds. However, similar ABR thresholds may be accompanied by differences

in the suprathreshold response of Wave I, which could reflect differential degeneration of

the auditory nerve (Kujawa & Liberman, 2009). AIDA treatment did not affect noise-

induced changes in either ABR thresholds or Wave I amplitude (Figure 7 E), thus

indicating that the effect of AIDA on blocking Z-LTD is not due to differential noise-induced

hearing loss after AIDA treatment. Together, these results demonstrate that sound-induced

Z-LTD requires G1 mGluR activation, consistent with our *in vitro* results.

Discussion

Our results show that long-term synaptic zinc plasticity is an experience-, G1 mGluR-dependent mechanism that bidirectionally modulates synaptic zinc signaling in the DCN. Is this a general mechanism that applies to all synaptic zinc-containing brain areas? Synaptic zinc is present throughout the neocortex and other brain structures, such as the amygdala and the hippocampus (McAllister & Dyck, 2017). Moreover, synaptic zinc is modulated by sensory activity throughout the sensory cortex (McAllister & Dyck, 2017), shapes the gain of central sensory responses (Anderson *et al.*, 2017), and when upregulated by optic nerve injury, it inhibits retinal ganglion cell survival and axon regeneration (Li *et al.*, 2017). It is therefore likely, although not tested here, that the reported long-term synaptic zinc plasticity mechanism is a general mechanism that dynamically modulates sensory processing for adaptation to different sensory environments and injury.

Whereas the exact synaptic, natural, and ethologically relevant stimuli that elicit Z-LTP and Z-LTD remain unknown, here we developed *in vitro* and *in vivo* models for studying Z-LTP and Z-LTD. This is a crucial step towards further elucidation of the detailed natural stimuli eliciting long-term synaptic zinc plasticity, as well as the precise cellular and molecular mechanisms underlying the induction and expression of Z-LTP and Z-LTD. Moreover, our model will be useful for probing the unknown behavioral consequences of Z-LTP and Z-LTD.

Mechanisms of Group 1 mGluR-dependent Z-LTP and Z-LTD. Our results show that differential activation of G1 mGluRs, by either LFS/HFS or high/low concentrations of DHPG, determines the induction and direction of long-term synaptic zinc plasticity. Prolonged LFS causes Z-LTP, similarly to G1 mGluR activation with 50 μM DHPG;

whereas, brief HFS causes Z-LTD, similarly to activation with 5 μM DHPG. This suggests

that prolonged LFS activates G1 mGluR signaling differently than brief HFS.

Group 1 mGluRs, mGluR1 and mGluR5, are linked to the IP_3-Diacylglycerol (DAG)

signaling pathway, leading to intracellular rises in Ca^{2+} from intracellular stores (Abdul-

Ghani *et al.*, 1996; Conn & Pin, 1997; Kim *et al.*, 2008). In the hippocampus, LFS induces

G1 mGluR-mediated LTD via postsynaptic AMPAR endocytosis involving Ca^{2+} release

from endoplasmic reticulum (ER) stores and dendritic protein synthesis (Huber *et al.*, 2000;

Holbro *et al.*, 2009; Luscher & Huber, 2010; Pick & Ziff, 2018). Moreover, in the

hippocampus, HFS or theta-burst stimulation induces G1 mGluR-mediated LTP, also

involving ER Ca^{2+} release, resulting in postsynaptic AMPAR/NMDAR trafficking or

enhanced presynaptic glutamate release (Topolnik *et al.*, 2006; Wu *et al.*, 2008; Anwyl,

2009). It remains unknown whether G1 mGluR-dependent Z-LTP and Z-LTD are

downstream effects of the same signaling pathways that induce LTD and LTP, or occur

through separate mechanisms. Nonetheless, we propose, albeit not tested here, that

differential G1 mGluR activation, by LFS/HFS, leads to subsequent release of different

amounts or types of intracellular Ca^{2+} signals. Different Ca^{2+} signals may in turn activate

diverse signaling pathways that ultimately lead to increased and decreased synaptic zinc

signaling. An analogue that comes to mind is the mechanism via which differential

activation of NMDARs, by various levels of synaptic activity, leads to variable Ca^{2+} levels

and signaling, ultimately determining the induction of both LTP and LTD (Malenka & Bear,

2004).

Our results suggest that increases or decreases in synaptic zinc signaling, evidenced by

increased or decreased ZX1 potentiation of EPSCs, are mediated by bidirectional

modulation of vesicular zinc levels and subsequent synaptic zinc release. High or low

concentrations of DHPG, which induce Z-LTP and Z-LTD, increase or decrease

presynaptic zinc levels in PF terminals (Figure 5). Furthermore, synaptic zinc plasticity

modulates zinc-mediated inhibition of NMDARs as well as AMPARs, and this effect on

NMDARs cannot be explained by postsynaptic changes in the relative contributions of

GluN2A vs. GluN2B subunits in the NMDAR EPSCs (Figure 6). Although we cannot fully

exclude potential contributions of postsynaptic mechanisms in synaptic zinc plasticity, our

results support that synaptic zinc plasticity is mainly mediated by activity-dependent

modulation of presynaptic zinc levels and signaling, and are consistent with previous

studies demonstrating experience-dependent modulation of vesicular zinc levels in the

somatosensory cortex (Brown & Dyck, 2002, 2005), visual cortex (Dyck *et al.*, 2003), optic

nerve (Li *et al.*, 2017), and the DCN (Kalappa *et al.*, 2015).

Cartwheel cells express G1 mGluRs, particularly mGluR1, suggesting that the locus of

induction of zinc plasticity is postsynaptic (Wright *et al.*, 1996). Because Z-LTP and Z-LTD

involve modulation of presynaptic zinc levels, one suggestion is the presence of a

retrograde signal from CWCs involved in the expression of Z-LTP and Z-LTD in PFs.

Alternatively, the presence of mGluR1 on axon terminals in the DCN molecular layer may

support a presynaptic locus of induction (Bilak & Morest, 1998). Because ZnT3 determines

vesicular zinc levels (Palmiter *et al.*, 1996; Cole *et al.*, 1999), modulation of ZnT3

expression or function may underlie the expression of Z-LTP and Z-LTD. In the retina,

optic nerve injury increases ZnT3 immunostaining, supporting that increases in ZnT3

expression mediate increases in synaptic zinc levels (Li *et al.*, 2017). However, in the

barrel cortex, whisker plucking increases the vesicular zinc content and the density of zinc-

containing synapses, but does not alter either ZnT3 protein or mRNA levels (Brown &

Dyck, 2002; Liguz-Lecznar *et al.*, 2005; Nakashima & Dyck, 2010; Nakashima *et al.*,

2011). Furthermore, in barrel cortical layers IV and V, the density of excitatory synapses

remains unchanged despite the increased density of zinc-containing synapses, indicating

that some previously excitatory non-zinc-containing synapses were converted to zinc-

containing synapses (Nakashima & Dyck, 2010). Together, these studies suggest that

changes in vesicular zinc content can occur without affecting glutamatergic synapses,

likely via functional modulation of pre-existing ZnT3. This may also explain our

electrophysiological results after sound exposure, because sound exposure caused Z-LTD

without affecting presynaptic glutamate dynamics (Figure 7 C). In the context of ZnT3

modulation, it is interesting that the vesicular glutamate transporter 1 (VGlut1), which is co-

targeted to synaptic vesicles with ZnT3, increases ZnT3 zinc transport in cultured cells

(Salazar *et al.*, 2005). Because VGlut1 is highly expressed in the DCN molecular layer

(Zhou *et al.*, 2007), one hypothesis is that modulation of VGlut1 may modulate ZnT3

function in PF terminals. However, the independent modulation of presynaptic glutamate

and zinc dynamics after sound exposure (Figure 7 C) suggests a VGlut1-independent

mechanism of ZnT3 modulation. While our results reveal a role for G1 mGluRs in Z-LTP

and Z-LTD, future experiments will be necessary to determine the detailed induction and

expression mechanisms.

Implications of Z-LTP and Z-LTD for short-term plasticity. Previous studies in DCN PF

synapses revealed that synaptic zinc triggers endocannabinoid synthesis, which inhibits

presynaptic glutamate release and modulates short-term plasticity (Perez-Rosello *et al.*,

2013; Kalappa & Tzounopoulos, 2017). During high-frequency (50 Hz) trains, synaptic zinc

inhibits AMPAR EPSCs during the first few stimuli, but enhances steady-state EPSCs in

subsequent stimuli by recruiting endocannabinoid signaling and enhancing synaptic

facilitation (Kalappa & Tzounopoulos, 2017). Therefore, long-term increases in zinc

signaling, via Z-LTP, would enhance endocannabinoid activation during subsequent

stimulus trains, increase synaptic facilitation, and further enhance steady-state EPSCs.

Conversely, long-term decreases in zinc signaling, via Z-LTD, would reduce

endocannabinoid activation, decrease synaptic facilitation, and suppress steady-state

EPSCs.

Following stimulus trains, zinc-mediated endocannabinoid activation causes short-term

depression and inhibits short-term facilitation (Perez-Rosello *et al.*, 2013). Therefore, Z-

LTP and Z-LTD are expected to shift the balance between short-term facilitation and short-

term depression in DCN synapses. Z-LTP will enhance subsequent zinc-mediated short-

term depression, whereas Z-LTD will enhance short-term facilitation. Taken together, our

results highlight a powerful mechanism by which long-term bidirectional zinc plasticity may

modulate short-term glutamatergic synaptic plasticity.

Implications of Z-LTP and Z-LTD for LTD and LTP. In central synapses, including DCN

PF synapses, the direction and size of LTP or LTD are determined by the combination of

multiple simultaneous LTP and LTD mechanisms (O'Connor *et al.*, 2005; Bender *et al.*,

2006; Tzounopoulos *et al.*, 2007; Shen *et al.*, 2008; Zhao & Tzounopoulos, 2011). In DCN

PF synapses, LTP and LTD are influenced by the coactivation of pre- and postsynaptic

signaling mechanisms including NMDARs, mGluRs, muscarinic acetylcholine receptors,

and endocannabinoid signaling (Fujino & Oertel, 2003; Tzounopoulos *et al.*, 2007; Zhao &

Tzounopoulos, 2011). Therefore, bidirectional zinc plasticity likely acts together with these

other known mechanisms to shape the size and direction of synaptic plasticity.

Several of our results are consistent with this notion. As shown in Figure 2, blockade of

NMDARs did not block either HFS-induced LTP or Z-LTD. This indicates that NMDAR-

independent LTP was induced, suggesting that Z-LTD contributes to NMDAR-independent

LTP (Figure 2 A). G1 mGluR antagonists blocked HFS-induced Z-LTD (Figure 2 B).

Therefore, the induced LTP under these conditions is NMDAR-, G1 mGluR-, and Z-LTD-

independent. As shown in Figure 3 A, LFS induced Z-LTP; however, LFS did not induce

LTD. This suggests that LFS also induced an LTP that counterbalances the LTD effect of

Z-LTP. This is consistent with previous studies showing that LTP and LTD mechanisms

occur simultaneously in DCN PF synapses (Tzounopoulos *et al.*, 2007). G1 mGluR

antagonists blocked Z-LTP, but LTD was induced under these conditions (Figure 3 B),

suggesting that this LTD is NMDAR-, G1 mGluR-, and Z-LTP-independent. Taken

together, all these results are consistent with previous studies and further support the

notion that LTP and LTD are the result of coactivation of different signaling pathways of

long-term plasticity in the DCN (Fujino & Oertel, 2003; Tzounopoulos *et al.*, 2007; Zhao &

Tzounopoulos, 2011). Nevertheless, our results add Z-LTP and Z-LTD as new

mechanisms of LTD and LTP at zinc-containing glutamatergic synapses.

In DCN PFs, mGluR activation contributes to both HFS-induced LTP and LFS-induced

LTD (Fujino & Oertel, 2003). Our findings on HFS-induced LTP in CWCs are consistent

with Fujino & Oertel, 2003 (Fujino & Oertel, 2003), demonstrating mGluR- and NMDAR-

independent LTP (Figure 2 B). However, Fujino & Oertel showed LFS-induced NMDAR-

dependent LTD, whereas here we observed LFS-induced NMDAR-independent LTD

(Figure 3 B). This discrepancy could be explained by the use of different LFS induction

protocols (5 Hz for 3 min here, vs. 1 Hz for 5 min. paired with postsynaptic depolarization)

and extracellular solutions (4/4 mM Ca^{2+}/Mg^{2+} here, vs. 2.4/1.3 mM Ca^{2+}/Mg^{2+}).

In addition to DCN PF synapses, Z-LTP and Z-LTD may contribute to LTD and LTP in

other synaptic zinc-containing brain areas which express G1 mGluR-dependent LTD and

LTP, such as the hippocampus, amygdala, and striatum (Oliet *et al.*, 1997; Huber *et al.*,

2000; Gubellini *et al.*, 2003; Topolnik *et al.*, 2006; Wu *et al.*, 2008; Anwyl, 2009; Luscher &

Huber, 2010; Chen *et al.*, 2017; McAllister & Dyck, 2017). In the hippocampus, LFS

induces G1 mGluR-mediated LTD, whereas HFS induces LTP (Oliet *et al.*, 1997; Huber *et*

al., 2000; Topolnik *et al.*, 2006; Wu *et al.*, 2008; Anwyl, 2009). Therefore, LFS-induced Z-

LTP would likely further enhance the effects of G1 mGluR-LTD, by increasing zinc

inhibition of AMPARs; whereas HFS-induced Z-LTD would further enhance the effects of

LTP, by reducing zinc inhibition of AMPARs. Thus, synaptic zinc plasticity likely serves as

a positive feedback mechanism to enhance the effects of G1 mGluR-dependent LTP or

LTD on glutamatergic synaptic transmission.

Implications of Z-LTP and Z-LTD for metaplasticity. Our results reveal that the

induction of Z-LTP and Z-LTD is NMDAR-independent. However, zinc inhibits NMDARs

and thus modulates the induction of NMDAR-dependent LTP and LTD in the hippocampus

(Izumi *et al.*, 2006; Takeda *et al.*, 2009; Vergnano *et al.*, 2014). As such, long-term

synaptic zinc plasticity may contribute to 'metaplasticity', the modulation of subsequent

LTP and LTD (Abraham & Tate, 1997). Z-LTD, by reducing the inhibitory effect of zinc on

NMDARs, may promote subsequent NMDAR-dependent LTP and decrease subsequent

NMDAR-dependent LTD. Conversely, Z-LTP, by enhancing the inhibitory effect of zinc on

NMDARs, may promote subsequent NMDAR-LTD over NMDAR-LTP. Therefore, zinc

plasticity likely serves as a positive feedback mechanism for NMDAR-dependent

metaplasticity.

Synaptic zinc contributes to mossy fiber presynaptic LTP in response to HFS, via

activation of TrkB receptors (Huang *et al.*, 2008; Pan *et al.*, 2011). Therefore, if HFS

induces Z-LTD in mossy fiber synapses, it would act as a negative feedback signal for

subsequent LTP induction. Taken together, we propose that the role of Z-LTD and Z-LTP

in LTP and LTD depends on the specific mechanisms underlying LTP and LTD, but

overall, Z-LTD and Z-LTP likely act as positive feedback signals in G1 mGluR-dependent

synaptic plasticity and NMDAR-dependent metaplasticity.

Clinical and translational implications of zinc plasticity. In the context of zinc plasticity

as a positive feedback signal for NMDAR-dependent metaplasticity, it is interesting that

exposure to loud sound – known to induce tinnitus – causes Z-LTD in the DCN. Although

not tested here, it is possible that Z-LTD could potentially lead to runaway excitation due to

enhanced LTP and decreased LTD, and thus to pathological DCN hyperactivity associated

with tinnitus (Tzounopoulos, 2008). Noise-induced pathological hyperexcitability through

LTP/LTD-like mechanisms in the DCN PF synapses has been hypothesized and recently

implicated in tinnitus treatment (Tzounopoulos, 2008; Marks *et al.*, 2018), therefore

suggesting that noise-induced reductions in synaptic zinc might contribute to tinnitus.

References

Abdul-Ghani MA, Valiante TA, Carlen PL & Pennefather PS. (1996). Metabotropic
glutamate receptors coupled to IP3 production mediate inhibition of IAHP in
rat dentate granule neurons. *J Neurophysiol* **76,** 2691-2700.
809
810 Abraham WC & Tate WP. (1997). Metaplasticity: a new vista across the field of
811 synaptic plasticity. *Prog Neurobiol* **52,** 303-323.
812
813 Anderson CT, Kumar M, Xiong S & Tzounopoulos T. (2017). Cell-specific gain
814 modulation by synaptically released zinc in cortical circuits of audition. *Elife* **6**.
815

816   Anderson CT, Radford RJ, Zastrow ML, Zhang DY, Apfel UP, Lippard SJ & Tzounopoulos
817   T. (2015). Modulation of extrasynaptic NMDA receptors by synaptic and tonic
818   zinc. *Proc Natl Acad Sci U S A* **112,** E2705-2714.
819
820   Anwyl R. (2009). Metabotropic glutamate receptor-dependent long-term potentiation.
821   *Neuropharmacology* **56,** 735-740.
822
823   Auerbach BD, Rodrigues PV & Salvi RJ. (2014). Central gain control in tinnitus and
824   hyperacusis. *Front Neurol* **5,** 206.
825
826   Bender VA, Bender KJ, Brasier DJ & Feldman DE. (2006). Two coincidence detectors
827   for spike timing-dependent plasticity in somatosensory cortex. *J Neurosci* **26,**
828   4166-4177.
829
830   Bilak SR & Morest DK. (1998). Differential expression of the metabotropic glutamate
831   receptor mGluR1alpha by neurons and axons in the cochlear nucleus: in situ
832   hybridization and immunohistochemistry. *Synapse* **28,** 251-270.
833
834   Brown CE & Dyck RH. (2002). Rapid, experience-dependent changes in levels of
835   synaptic zinc in primary somatosensory cortex of the adult mouse. *J Neurosci*
836   **22,** 2617-2625.
837
838   Brown CE & Dyck RH. (2005). Modulation of synaptic zinc in barrel cortex by whisker
839   stimulation. *Neuroscience* **134,** 355-359.
840
841   Chen A, Hu WW, Jiang XL, Potegal M & Li H. (2017). Molecular mechanisms of group I
842   metabotropic glutamate receptor mediated LTP and LTD in basolateral
843   amygdala in vitro. *Psychopharmacology (Berl)* **234,** 681-694.
844
845   Cole TB, Wenzel HJ, Kafer KE, Schwartzkroin PA & Palmiter RD. (1999). Elimination of
846   zinc from synaptic vesicles in the intact mouse brain by disruption of the ZnT3
847   gene. *Proc Natl Acad Sci U S A* **96,** 1716-1721.
848
849   Conn PJ & Pin JP. (1997). Pharmacology and functions of metabotropic glutamate
850   receptors. *Annu Rev Pharmacol Toxicol* **37,** 205-237.
851
852   Danscher G & Stoltenberg M. (2005). Zinc-specific autometallographic in vivo
853   selenium methods: tracing of zinc-enriched (ZEN) terminals, ZEN pathways,
854   and pools of zinc ions in a multitude of other ZEN cells. *J Histochem Cytochem*
855   **53,** 141-153.
856
857   Dyck RH, Chaudhuri A & Cynader MS. (2003). Experience-dependent regulation of the
858   zincergic innervation of visual cortex in adult monkeys. *Cereb Cortex* **13,** 1094-
859   1109.

Frederickson CJ, Howell GA, Haigh MD & Danscher G. (1988). Zinc-containing fiber
systems in the cochlear nuclei of the rat and mouse. *Hear Res* **36,** 203-211.

Frederickson CJ, Koh JY & Bush AI. (2005). The neurobiology of zinc in health and
disease. *Nat Rev Neurosci* **6,** 449-462.

Fujino K & Oertel D. (2003). Bidirectional synaptic plasticity in the cerebellum-like
mammalian dorsal cochlear nucleus. *Proc Natl Acad Sci U S A* **100,** 265-270.

Gubellini P, Saulle E, Centonze D, Costa C, Tropepi D, Bernardi G, Conquet F &
Calabresi P. (2003). Corticostriatal LTP requires combined mGluR1 and
mGluR5 activation. *Neuropharmacology* **44,** 8-16.

Hansen KB, Ogden KK, Yuan H & Traynelis SF. (2014). Distinct functional and
pharmacological properties of Triheteromeric GluN1/GluN2A/GluN2B NMDA
receptors. *Neuron* **81,** 1084-1096.

Holbro N, Grunditz A & Oertner TG. (2009). Differential distribution of endoplasmic
reticulum controls metabotropic signaling and plasticity at hippocampal
synapses. *Proc Natl Acad Sci U S A* **106,** 15055-15060.

Huang YZ, Pan E, Xiong ZQ & McNamara JO. (2008). Zinc-mediated transactivation of
TrkB potentiates the hippocampal mossy fiber-CA3 pyramid synapse. *Neuron*
57, 546-558.

Huber KM, Kayser MS & Bear MF. (2000). Role for rapid dendritic protein synthesis in
hippocampal mGluR-dependent long-term depression. *Science* **288,** 1254-
1257.

Huber KM, Roder JC & Bear MF. (2001). Chemical induction of mGluR5- and protein
synthesis--dependent long-term depression in hippocampal area CA1. *J*
Neurophysiol **86,** 321-325.

Izumi Y, Auberson YP & Zorumski CF. (2006). Zinc modulates bidirectional
hippocampal plasticity by effects on NMDA receptors. *J Neurosci* **26,** 7181-
7188.

Kalappa BI, Anderson CT, Goldberg JM, Lippard SJ & Tzounopoulos T. (2015). AMPA
receptor inhibition by synaptically released zinc. *Proc Natl Acad Sci U S A* **112,**
15749-15754.

Kalappa BI & Tzounopoulos T. (2017). Context-Dependent Modulation of Excitatory
Synaptic Strength by Synaptically Released Zinc. *eNeuro* **4**.

Kim CH, Lee J, Lee JY & Roche KW. (2008). Metabotropic glutamate receptors:
phosphorylation and receptor signaling. *J Neurosci Res* **86,** 1-10.

Kujawa SG & Liberman MC. (2009). Adding insult to injury: cochlear nerve
degeneration after "temporary" noise-induced hearing loss. *J Neurosci* **29,**
14077-14085.

Kumar M, Xiong S, Tzounopoulos T & Anderson CT. (2019). Fine Control of Sound
Frequency Tuning and Frequency Discrimination Acuity by Synaptic Zinc
Signaling in Mouse Auditory Cortex. *J Neurosci* **39,** 854-865.

Li Y, Andereggen L, Yuki K, Omura K, Yin Y, Gilbert HY, Erdogan B, Asdourian MS,
Shrock C, de Lima S, Apfel UP, Zhuo Y, Hershfinkel M, Lippard SJ, Rosenberg PA
& Benowitz L. (2017). Mobile zinc increases rapidly in the retina after optic
nerve injury and regulates ganglion cell survival and optic nerve regeneration.
Proc Natl Acad Sci U S A **114,** E209-E218.

Liguz-Lecznar M, Nowicka D, Czupryn A & Skangiel-Kramska J. (2005). Dissociation of
synaptic zinc level and zinc transporter 3 expression during postnatal
development and after sensory deprivation in the barrel cortex of mice. *Brain*
Res Bull **66,** 106-113.

Luscher C & Huber KM. (2010). Group 1 mGluR-dependent synaptic long-term
depression: mechanisms and implications for circuitry and disease. *Neuron* **65,**
445-459.

Malenka RC & Bear MF. (2004). LTP and LTD: an embarrassment of riches. *Neuron* **44,**
5-21.

Malenka RC & Nicoll RA. (1993). NMDA-receptor-dependent synaptic plasticity:
multiple forms and mechanisms. *Trends Neurosci* **16,** 521-527.

Manis PB, Spirou GA, Wright DD, Paydar S & Ryugo DK. (1994). Physiology and
morphology of complex spiking neurons in the guinea pig dorsal cochlear
nucleus. *J Comp Neurol* **348,** 261-276.

Marks KL, Martel DT, Wu C, Basura GJ, Roberts LE, Schvartz-Leyzac KC & Shore SE.
(2018). Auditory-somatosensory bimodal stimulation desynchronizes brain
circuitry to reduce tinnitus in guinea pigs and humans. *Sci Transl Med* **10**.

McAllister BB & Dyck RH. (2017). Zinc transporter 3 (ZnT3) and vesicular zinc in
central nervous system function. *Neurosci Biobehav Rev* **80,** 329-350.

Mulkey RM & Malenka RC. (1992). Mechanisms underlying induction of homosynaptic
long-term depression in area CA1 of the hippocampus. *Neuron* **9,** 967-975.
950
951 Nakashima AS, Butt RH & Dyck RH. (2011). Alterations in protein and gene expression
952   within the barrel cortices of ZnT3 knockout mice: experience-independent and
953   dependent changes. *Neurochem Int* **59,** 860-870.
954
955 Nakashima AS & Dyck RH. (2009). Zinc and cortical plasticity. *Brain Res Rev* **59,** 347-
956   373.
957
958 Nakashima AS & Dyck RH. (2010). Dynamic, experience-dependent modulation of
959   synaptic zinc within the excitatory synapses of the mouse barrel cortex.
960   *Neuroscience* **170,** 1015-1019.
961
962 O'Connor DH, Wittenberg GM & Wang SS. (2005). Dissection of bidirectional synaptic
963   plasticity into saturable unidirectional processes. *J Neurophysiol* **94,** 1565-
964   1573.
965
966 Oliet SH, Malenka RC & Nicoll RA. (1997). Two distinct forms of long-term depression
967   coexist in CA1 hippocampal pyramidal cells. *Neuron* **18,** 969-982.
968
969 Palmiter RD, Cole TB, Quaife CJ & Findley SD. (1996). ZnT-3, a putative transporter of
970   zinc into synaptic vesicles. *Proc Natl Acad Sci U S A* **93,** 14934-14939.
971
972 Pan E, Zhang XA, Huang Z, Krezel A, Zhao M, Tinberg CE, Lippard SJ & McNamara JO.
973   (2011). Vesicular zinc promotes presynaptic and inhibits postsynaptic long-
974   term potentiation of mossy fiber-CA3 synapse. *Neuron* **71,** 1116-1126.
975
976 Paoletti P, Ascher P & Neyton J. (1997). High-affinity zinc inhibition of NMDA NR1-
977   NR2A receptors. *J Neurosci* **17,** 5711-5725.
978
979 Perez-Rosello T, Anderson CT, Schopfer FJ, Zhao Y, Gilad D, Salvatore SR, Freeman BA,
980   Hershfinkel M, Aizenman E & Tzounopoulos T. (2013). Synaptic Zn2+ inhibits
981   neurotransmitter release by promoting endocannabinoid synthesis. *J Neurosci*
982   **33,** 9259-9272.
983
984 Pick JE & Ziff EB. (2018). Regulation of AMPA receptor trafficking and exit from the
985   endoplasmic reticulum. *Mol Cell Neurosci*.
986
987 Rachline J, Perin-Dureau F, Le Goff A, Neyton J & Paoletti P. (2005). The micromolar
988   zinc-binding domain on the NMDA receptor subunit NR2B. *J Neurosci* **25,** 308-
989   317.
990

991 Rubio ME & Juiz JM. (1998). Chemical anatomy of excitatory endings in the dorsal
992   cochlear nucleus of the rat: differential synaptic distribution of aspartate
993   aminotransferase, glutamate, and vesicular zinc. *J Comp Neurol* **399,** 341-358.
994
995 Ryugo DK & Willard FH. (1985). The dorsal cochlear nucleus of the mouse: a light
996   microscopic analysis of neurons that project to the inferior colliculus. *J Comp*
997   *Neurol* **242,** 381-396.
998
999 Salazar G, Craige B, Love R, Kalman D & Faundez V. (2005). Vglut1 and ZnT3 co-
1000   targeting mechanisms regulate vesicular zinc stores in PC12 cells. *J Cell Sci* **118,**
1001   1911-1921.
1002
1003 Shen W, Flajolet M, Greengard P & Surmeier DJ. (2008). Dichotomous dopaminergic
1004   control of striatal synaptic plasticity. *Science* **321,** 848-851.
1005
1006 Snyder EM, Philpot BD, Huber KM, Dong X, Fallon JR & Bear MF. (2001).
1007   Internalization of ionotropic glutamate receptors in response to mGluR
1008   activation. *Nat Neurosci* **4,** 1079-1085.
1009
1010 Suter BA, O'Connor T, Iyer V, Petreanu LT, Hooks BM, Kiritani T, Svoboda K &
1011   Shepherd GM. (2010). Ephus: multipurpose data acquisition software for
1012   neuroscience experiments. *Front Neural Circuits* **4,** 100.
1013
1014 Takeda A, Fuke S, Ando M & Oku N. (2009). Positive modulation of long-term
1015   potentiation at hippocampal CA1 synapses by low micromolar concentrations
1016   of zinc. *Neuroscience* **158,** 585-591.
1017
1018 Topolnik L, Azzi M, Morin F, Kougioumoutzakis A & Lacaille JC. (2006). mGluR1/5
1019   subtype-specific calcium signalling and induction of long-term potentiation in
1020   rat hippocampal oriens/alveus interneurones. *J Physiol* **575,** 115-131.
1021
1022 Tovar KR & Westbrook GL. (2012). Amino-terminal ligands prolong NMDA Receptor-
1023   mediated EPSCs. *J Neurosci* **32,** 8065-8073.
1024
1025 Traynelis SF, Wollmuth LP, McBain CJ, Menniti FS, Vance KM, Ogden KK, Hansen KB,
1026   Yuan H, Myers SJ & Dingledine R. (2010). Glutamate receptor ion channels:
1027   structure, regulation, and function. *Pharmacol Rev* **62,** 405-496.
1028
1029 Tzounopoulos T. (2008). Mechanisms of synaptic plasticity in the dorsal cochlear
1030   nucleus: plasticity-induced changes that could underlie tinnitus. *Am J Audiol*
1031   **17,** S170-175.
1032
1033 Tzounopoulos T, Kim Y, Oertel D & Trussell LO. (2004). Cell-specific, spike timing-
1034   dependent plasticities in the dorsal cochlear nucleus. *Nat Neurosci* **7,** 719-725.

1035
1036   Tzounopoulos T, Rubio ME, Keen JE & Trussell LO. (2007). Coactivation of pre- and
1037   postsynaptic signaling mechanisms determines cell-specific spike-timing-
1038   dependent plasticity. *Neuron* **54,** 291-301.
1039
1040   Vergnano AM, Rebola N, Savtchenko LP, Pinheiro PS, Casado M, Kieffer BL, Rusakov
1041   DA, Mulle C & Paoletti P. (2014). Zinc dynamics and action at excitatory
1042   synapses. *Neuron* **82,** 1101-1114.
1043
1044   Vogt K, Mellor J, Tong G & Nicoll R. (2000). The actions of synaptically released zinc at
1045   hippocampal mossy fiber synapses. *Neuron* **26,** 187-196.
1046
1047   Wisniewski K & Car H. (2002). (S)-3,5-DHPG: a review. *CNS Drug Rev* **8,** 101-116.
1048
1049   Wright DD, Blackstone CD, Huganir RL & Ryugo DK. (1996). Immunocytochemical
1050   localization of the mGluR1 alpha metabotropic glutamate receptor in the dorsal
1051   cochlear nucleus. *J Comp Neurol* **364,** 729-745.
1052
1053   Wu J, Harney S, Rowan MJ & Anwyl R. (2008). Involvement of group I mGluRs in LTP
1054   induced by strong high frequency stimulation in the dentate gyrus in vitro.
1055   *Neurosci Lett* **436,** 235-238.
1056
1057   Zastrow ML, Radford RJ, Chyan W, Anderson CT, Zhang DY, Loas A, Tzounopoulos T &
1058   Lippard SJ. (2016). Reaction-Based Probes for Imaging Mobile Zinc in Live Cells
1059   and Tissues. *ACS Sens* **1,** 32-39.
1060
1061   Zhang S & Oertel D. (1993). Cartwheel and superficial stellate cells of the dorsal
1062   cochlear nucleus of mice: intracellular recordings in slices. *J Neurophysiol* **69,**
1063   1384-1397.
1064
1065   Zhao Y & Tzounopoulos T. (2011). Physiological activation of cholinergic inputs
1066   controls associative synaptic plasticity via modulation of endocannabinoid
1067   signaling. *J Neurosci* **31,** 3158-3168.
1068
1069   Zhou J, Nannapaneni N & Shore S. (2007). Vessicular glutamate transporters 1 and 2
1070   are differentially associated with auditory nerve and spinal trigeminal inputs
1071   to the cochlear nucleus. *J Comp Neurol* **500,** 777-787.
1072
1073
1074
1075

Figure Legends

Figure 1. High-frequency stimulation (HFS) induces Z-LTD in DCN parallel fiber synapses. (A) Schematic of experimental setup illustrating stimulation of zinc-rich glutamatergic DCN parallel fibers (PFs) and whole-cell recording of a postsynaptic cartwheel cell (CWC). **(B)** *Left:* Time course of AMPAR EPSC amplitude before and after ZX1 application, normalized to baseline before ZX1 application (100 μM). *Right:* Example AMPAR EPSCs before and after ZX1 application, showing ZX1 potentiation. **(C)** *Left:* Time course of AMPAR EPSC amplitude before and after HFS, and before and after subsequent ZX1 application (blue). After obtaining a stable baseline, HFS was delivered (3 x 100 Hz for 1 sec, 10 sec ISI). EPSC % baseline after HFS (mins. 19-23): n=11, *p=0.041, one-sample t test vs. 100%. Star (*) indicates significant LTP. To examine ZX1 potentiation after HFS, after obtaining a stable baseline after HFS, AMPAR EPSC amplitude was renormalized to the new baseline before ZX1 application. The renormalization is indicated by a gap and restart of timing in the x-axis. For comparison, red line shows normalized time course of AMPAR EPSC amplitude before and after ZX1 application in controls replotted from **B**. *Right:* Example AMPAR EPSCs showing no ZX1 potentiation after HFS. **(D)** Average ZX1 potentiation (% increase from baseline) during the last 5 min of ZX1 application. 'Control' (n=10) vs. 'HFS' (n=11): *p=0.002, unpaired t test. The reduction in ZX1 potentiation is termed Z-LTD. Values represent mean ± SEM. Star (*) indicates p<0.05. For detailed values and statistical tests for all figures, see Materials and Methods, Statistical Analysis section.

Figure 2. Group 1 mGluR activation is required for HFS-induced Z-LTD. (A) Time course of AMPAR EPSC amplitude before and after HFS in the presence of APV (50 μM),

and before and after subsequent ZX1 application. EPSC % baseline after HFS (mins. 19-

23): n=9, *p=0.0008, one-sample t test vs. 100%. **(B)** Same time course as in **A** but in the

presence of LY367385 (100 μM), MPEP (4 μM), and APV (50 μM). EPSC % baseline after

HFS (mins. 19-23): n=6, *p=0.041, one-sample t test vs. 100%. For **(A-B)**, star (*) indicates

significant LTP. To examine ZX1 potentiation after HFS, similar approach and

renormalization as in **1C** was performed. Red line shows the time course of AMPAR EPSC

amplitude before and after ZX1 application in control replotted from **1B**. Example traces

show AMPAR EPSCs before and after ZX1. **(C)** Average ZX1 potentiation (% increase

from baseline) during the last 5 min of ZX1 application, with control data from **1D.** 'HFS +

APV' (n=9) reduced ZX1 potentiation compared to control; this reduction was blocked by

LY367385 and MPEP (n=5). One-way ANOVA/Bonferroni, *p=0.003. Values represent

mean $\pm$ SEM. Star (*) indicates p<0.05.

1112

Figure 3. Low frequency stimulation (LFS) induces Z-LTP, which requires Group 1

mGluR activation. (A) Time course of AMPAR EPSC amplitude before and after LFS (5

Hz, 3 min), and before and after subsequent ZX1 application (cyan); and similar time

course in interleaved control experiments (without LFS, red). **(B)** Same time course as in **A**

but in the presence of LY367385 (100 μM) and MPEP (4 μM) (green). EPSC % baseline

after LFS (mins. 20-24): n=6, *p=0.006, one-sample t test vs. 100%. Star (*) indicates

significant LTD. Red line shows similar time course in controls replotted from **A**. For **(A-B)**,

to examine the ZX1 potentiation after LFS, similar approach and renormalization as in **1C**

was performed. Example traces show AMPAR EPSCs before and after ZX1. **(C)** Average

ZX1 potentiation (% increase from baseline) during the last 5 min of ZX1 application.

'Control': n=5; 'LFS': n=6; 'LFS + LY367385, MPEP': n=6. LFS increased ZX1 potentiation

compared to control; this increase was blocked by LY367385 and MPEP. One-way

ANOVA/Bonferroni, *p=0.02. The increase in ZX1 potentiation is termed Z-LTP. Values

represent mean $\pm$ SEM. Star (*) indicates p<0.05.

1127

Figure 4. Group 1 mGluR activation is sufficient to induce Z-LTP and Z-LTD. (A) Time

course of AMPAR EPSC amplitude before and after application of 50 μM DHPG, and

before and after subsequent ZX1 application. EPSC % baseline after 50 μM DHPG (mins.

16-20): n=6, *p=0.0002, one-sample t test vs. 100%. Star (*) indicates significant synaptic

depression. **(B)** Time course of AMPAR EPSC amplitude before and after application of 5

μM DHPG, and before and after subsequent ZX1 application. For **(A-B)**, to examine the

ZX1 potentiation after DHPG application, after obtaining a stable baseline after DHPG,

AMPAR EPSC amplitude was renormalized to the new baseline before ZX1 application.

The renormalization is indicated by a gap and restart of timing in the x-axis. Example

traces show AMPAR EPSCs before and after ZX1. **(C)** Average ZX1 potentiation (%

increase from baseline) during the last 5 min of ZX1 application, with control data from **1D**.

'DHPG (50 μM)': n=5; 'DHPG (5 μM)': n=5. DHPG (50 μM) increased ZX1 potentiation

compared to control, whereas DHPG (5 μM) reduced ZX1 potentiation compared to

control. One-way ANOVA/Bonferroni, *p<0.0001. Increased and decreased ZX1

potentiation correspond to Z-LTP and Z-LTD, respectively. **(D)** Similar time course as in **A**,

but in same extracellular conditions as in **3A-B**. EPSC % baseline after 50 μM DHPG

(mins. 16-20): n=5, *p=0.044, one-sample t test vs. 100%. Star (*) indicates significant

synaptic depression. **(E)** Time course of AMPAR EPSC amplitude before and after

sequential LFS (5 Hz, 3 min) and application of 50 μM DHPG, and before and after

subsequent ZX1 application, in same conditions as in **D**. For **(D-E)**, to examine the ZX1

potentiation, similar approach and renormalization as in **A-B** was performed. Example

traces show AMPAR EPSCs before and after ZX1. **(F)** Average ZX1 potentiation (%

increase from baseline) during the last 5 min of ZX1 application for the experiments in **D-E**,

with LFS data from **3C**. 'DHPG (50 μM)': n=5; 'LFS + DHPG (50 μM)': n=5. Sequential LFS

and DHPG (50 μM) did not increase ZX1 potentiation compared to LFS or DHPG (50 μM)

alone. One-way ANOVA/Bonferroni: n.s. p=0.67. Values represent mean $\pm$ SEM. Star (*)

indicates p<0.05.

1155

Figure 5. Group 1 mGluR activation bidirectionally modulates presynaptic zinc

levels. (A) *Left:* Schematic of the DCN, showing the presynaptic zinc-containing region,

and the zinc-free region. *Right:* 20x image of DA-ZP1 fluorescence, demonstrating the

zinc-containing ROI (zinc ROI) and the zinc-free ROI, before and after application of 50 μM

DHPG. **(B)** Same approach as in **A**, before and after application of 5 μM DHPG. **(C)**

Average DA-ZP1 fluorescence after application of 50 μM or 5 μM DHPG, normalized to

baseline fluorescence before DHPG application. '+ DHPG (50 μM)': n=9, *p=0.004,

Wilcoxon signed rank test vs. 100%. '+ DHPG (5 μM)': n=8, *p=0.008, Wilcoxon signed

rank test vs. 100%. Values represent mean $\pm$ SEM. Star (*) indicates p<0.05.

1165

Figure 6. Group 1 mGluR-dependent Z-LTD reduces zinc inhibition of NMDARs. (A)

Left: Time course of AMPAR EPSC amplitude before and after HFS, and NMDAR EPSC

amplitude before and after subsequent ZX1 application (blue); and similar time course in

interleaved control experiments (without HFS, red). *Right:* Example NMDAR EPSCs before

and after ZX1 application. **(B)** *Left:* Same time course as in **A** but in the presence of

LY367385 (100 μM) and MPEP (4 μM) (green). Red line shows controls replotted from **A**.

Right: Example NMDAR EPSCs before and after ZX1 application. For **(A-B)**, after

obtaining a stable baseline of AMPAR EPSCs, HFS was delivered, then DNQX (20 μM)

was applied. NMDAR EPSCs were then recorded at +40 mV normalized to the baseline

NMDAR EPSC amplitude before ZX1 application. The switch from AMPAR to NMDAR

EPSC time course, and the renormalization of EPSC amplitude are indicated by a gap and

restart of timing in the x-axis. **(C)** Average ZX1 potentiation (% increase from baseline)

during the last 5 min of ZX1 application. 'Control': n=5; 'HFS': n=6; 'HFS + LY367385,

MPEP': n=5. HFS reduced ZX1 potentiation compared to control; this reduction was

blocked by LY367385 and MPEP. Kruskal-Wallis test/Dunn: *p=0.01. **(D)** Dose-response

of NMDAR EPSCs (% baseline) for increasing concentrations of ifenprodil, in controls (red)

and after HFS (blue). 'Control': n=3-5 per concentration; 'HFS': n=3-4 per concentration.

(E) IC_{50} of ifenprodil, from dose-responses in **D**. n.s. p=0.97, comparison of fits, extra sum-

of-squares F test. Values represent mean $\pm$ SEM. Star (*) indicates p<0.05.

1185

Figure 7. Sound-induced Z-LTD requires Group 1 mGluR activation. (A) Time course

of AMPAR EPSC amplitude before and after ZX1 application in slices from N.E. mice

(gray) and N.E. AIDA-treated mice (orange). Example traces show AMPAR EPSCs before

and after ZX1. **(B)** Average ZX1 potentiation (% increase from baseline) during the last 5

min of ZX1 application. 'N.E.' (n=5) vs. 'N.E. + AIDA' (n=6): *p=0.024, unpaired t test. **(C)**

Left: Average paired-pulse ratio (PPR, pulse 2 / pulse 1) of baseline AMPAR EPSCs in

slices from N.E. mice and N.E. AIDA-treated mice. 'N.E.' (n=5) vs. 'N.E. + AIDA' (n=6): n.s.

p=0.476, unpaired t test. Example traces show AMPAR EPSCs in response to two pulses.

Right: coefficient of variation (CV) analysis ($1/CV^2$) of baseline AMPAR EPSCs (pulse 1) in

slices from N.E. mice and N.E. AIDA-treated mice, normalized to N.E. mice. 'N.E. + AIDA':

n=6; n.s. p=0.44, Wilcoxon signed rank test vs. 1. **(D)** Example Auditory Brainstem

Responses (ABRs, 10-80 dB SPL sound stimuli) from sham-exposed mice (recorded from

sham-exposed, ipsilateral ear, black), N.E. mice (gray), and N.E. AIDA-treated mice

(orange). Because no ABRs were detected in the ipsilateral ears of N.E. mice, ABRs were

measured from ears contralateral to noise exposure. **(E)** *Left:* Average ABR thresholds (dB

SPL). 'Sham ipsi.': n=8; 'N.E. contra.': n=6; 'N.E. + AIDA contra.': n=7. N.E. increased ABR

thresholds compared to sham-exposed (*p=0.0002), but AIDA and N.E. did not affect

increases in ABR thresholds compared to N.E. alone, Kruskal-Wallis test/Dunn. *Right:*

Average ABR Wave I amplitude (μV). 'Sham ipsi.': n=8; 'N.E. contra.': n=6; 'N.E. + AIDA

contra.': n=7. N.E. decreased ABR Wave I amplitude compared to sham-exposed

(*p=0.0024), but AIDA and N.E. did not affect decreases in ABR Wave I amplitude

compared to N.E. alone, Kruskal-Wallis test/Dunn. Values represent mean $\pm$ SEM. Star (*)

indicates p<0.05.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7