Tumor suppressor DAPK1 catalyzes Adhesion Assembly on Rigid but Anoikis on Soft Matrices

Ruifang Qin[1], Haguy Wolfenson[1,2,*], Mayur Saxena[3], Michael P. Sheetz[1,4,*]

[1]Department of Biological Sciences, Columbia University, New York, NY10027.

[2]Current address: Rappaport Faculty of Medicine, Technion-Israel Institute of Technology, Haifa, Israel 31096.

[3]Department of Biomedical Engineering, Columbia University, New York, NY10027.

[4]Mechanobiology Institute, National University of Singapore, Singapore 117411.

Abstract

Cancer cells will normally grow on soft surfaces, but if rigidity sensing modules are restored in cancer cells, they will undergo apoptosis on soft surfaces (*anoikis)* like most normal cells. DAPK1 is a major tumor suppressor that activates cell death, but it is unclear how DAPK1 could activate *anoikis* through rigidity sensing. Here we find that when rigidity sensing is decreased through inhibition of DAPK1 activity, cells are transformed for growth on soft matrices. Further, DAPK1 catalyzes matrix adhesion assembly and is part of adhesions on rigid surfaces. Additional factors include DAPK1 phosphorylation of tropomyosin2.1, talin1 head domain and tyrosine phosphorylation of DAPK1 by Src. On soft surfaces, DAPK1 rapidly dissociates from the adhesion complexes and activates apoptosis that requires PTPN12 activity and talin1 head. Thus, DAPK1 is important for adhesion assembly on rigid surfaces and the activation of *anoikis* on soft surfaces through its binding to rigidity-sensing modules.

Introduction

In the early studies of cancer cells in vitro, it was evident that they could grow on widely different substrates, particularly on soft agar, which typically caused apoptosis of normal cells (a process called *anoikis*). This result indicated that either the cancer cells could not sense the surface properly, or that they were activated for growth under all conditions through receptor tyrosine kinases (RTK), Ras, or other mutations. Evidence for the former possibility came from recent studies showing that metastatic cancer cells (e.g., MDA-MB-231, HT1080 and SKOV3) undergo *anoikis* on soft agar after restoration of rigidity sensing modules. This was done through normalization of the expression levels of cytoskeletal proteins involved in rigidity sensing modules. Subsequent depletion of other proteins involved in the modules restored transformed growth[1]. However, the mechanism by which the rigidity sensing modules activated *anoikis* was unclear.

DAPK1 is a calcium/calmodulin-dependent protein kinase with cell death-inducing functions[2,3]. Not only is DAPK1 activity linked to *anoikis* but also it is a tumor suppressor that is depleted in many cancers[4-7]. Further, DAPK1 is linked to autophagy and apoptosis activated by TNF-alpha[8]. Classically, the activation of DAPK1 is believed to occur by calcium-calmodulin binding to DAPK1 to enable the dephosphorylation of the auto-inhibitory S308 site[9]. There is, however, additional evidence of inhibition through phosphorylation of vicinal tyrosines (Y490/Y491 in human DAPK1) and activation in

immune cells by tyrosine phosphatase activity[10]. Still, no direct link between DAPK1

activity and rigidity sensing has been established. Some studies have linked DAPK1 to

cytoskeletal proteins and motility, including the ability of DAPK1 to phosphorylate myosin

light chain and tropomyosin2.1 (Tpm2.1)[11,12] (formerly known as Tm1[13]). Notably,

Tpm2.1 expression is sufficient to restore *anoikis* in several metastatic cancer lines[1,14].

Further, DAPK1 has a role in cell polarization and migration through an interaction with

the talin1 head domain[15]. These observations raise the possibility that there may be a

role for DAPK1 in rigidity sensing, as well as a role for rigidity sensing complexes in

activating DAPK1 on soft surfaces.

Rigidity sensors in differentiated fibroblasts are relatively complicated protein modules

that are capable of transducing matrix rigidity into biochemical signals to either activate

growth on rigid matrices or apoptosis on soft matrices. The rigidity sensors are non-

muscle sarcomeric-like units of about 2 μm in length, with anti-parallel actin filaments

anchored to the matrix adhesions and a bipolar myosin IIA filament in the middle

(reviewed in Wolfenson et al., Ann Rev Physiol In Press). To test the rigidity, these units

contract the matrix to a constant distance of about 100 nm in total, and the force that is

developed is sensed. Receptor tyrosine kinases, AXL and ROR2, regulate the

displacement and the duration of the contractions, respectively[16]. EGFR and HER2 will

activate contractions on rigid but not on soft matrices downstream of Src

phosphorylation[17]. Surprisingly, normal rigidity sensing involves force-dependent

cleavage of talin1 by calpain that catalyzes growth on rigid surfaces through the

generation of free talin1 rod domains[18]. The velocity of the contraction depends upon

Tpm2.1 binding to the actin filaments, and AXL phosphorylation of Tpm2.1 on Y214 is

needed for adhesion formation[16]. Perhaps because this is a complicated sensory module,

depletion of this module's activity follows upon the depletion of any one of many

cytoskeletal proteins, including Tpm2.1, myosin IIA, and α-actinin 4[19,20]. Loss of rigidity

sensing activity, however, enables cell growth on soft surfaces, as is the case with many

cancers. Upon restoration of rigidity sensing in cancer cells, many will die on soft

surfaces, which begs the question of how does rigidity sensing activate apoptosis.

Here we studied the activation of *anoikis* through rigidity sensors, and found that DAPK1

activated *anoikis* and inhibition of DAPK1 was sufficient to enable growth on soft surfaces.

Under control conditions, assembly of rigidity sensors and adhesion formation normally

involved DAPK1 activity and particularly required the phosphorylation of Tpm2.1. On rigid

substrates, DAPK1 associated with adhesions in a Tpm2.1-dependent process that
involved Tpm2.1 phosphorylation by DAPK1. On soft substrates, DAPK1 rapidly
dissociated from the adhesion complexes and activated apoptosis. Apoptosis was
catalyzed by the constitutively phosphorylated mutant of Tpm2.1 and talin1 cleavage, but
it was inhibited by Src and EGFR activity. DAPK1 assembly in adhesions depended on
Y490/Y491 phosphorylation by Src or EGFR and tyrosine (Y) to phenylalanine (F)
mutations catalyzed apoptosis. Inhibition of the tyrosine phosphatase, PTPN12, inhibited
anoikis because it blocked DAPK1 activation. Thus, we suggest that DAPK1 is an
important component in the assembly of rigidity sensors and adhesions through
interactions with Tpm2.1 and talin1 head, leading to growth on rigid matrices and *anoikis*
on soft matrices.

Results

Inhibition of DAPK1 contributes to cell growth on soft matrices

Because DAPK1 activity correlated with *anoikis*, autophagy, and TNF-alpha-induced
apoptosis[8, 21, 22], we wanted to test if it was responsible for apoptosis of cells on soft
substrates. As shown in Fig. 1a, mouse embryonic fibroblasts (MEFs) on soft substrates
(0.2 kPa) activated apoptosis with the characteristic changes in morphology, including
membrane blebbing and cell rounding. Apoptosis of MEFs on soft substrates was
blocked by inhibition of DAPK1 (Fig. 1a). Similar results were observed in MCF10A,
normal human epithelial cells (Fig. 1a). Further quantitative analysis of cellular apoptosis
based on Annexin V immunostaining matched cell apoptotic morphology results
(Supplementary Fig. 1a). Thus, in both fibroblasts and epithelial cells, the cells were
protected from apoptosis on soft matrices by an inhibitor of DAPK1.

The autophosphorylation of DAPK1 at S308 was found to be the major mechanism of
DAPK1 inhibition in cells and that site was exposed for dephosphorylation upon binding
of calcium/calmodulin[9]. To test if phospho-S308 decreased in *anoikis*, we followed the
levels of pS308 by immunostaining with specific antibodies during the first 18 hours on
soft surfaces. As predicted, the level of pS308 phosphorylation dropped on soft surfaces,
indicating that DAPK1 was activated (Fig. 1b). Thus, there was activation of DAPK1 on
soft surfaces with dephosphorylation of S308.

DAPK1 regulates adhesion growth and force production

Inhibition of DAPK1 activity also slowed cell spreading and particularly adhesion

formation (Fig. 2a-c). Further, the knockdown of DAPK1 by shRNA produced similar

results (Supplementary Fig. 1b). Careful examination of the spreading process indicated

that the early spreading phase was nearly normal without DAPK1 activity, but the later

spreading phase that was associated with rigidity-sensing, contractile activity (Phase 2

of spreading[23]) was inhibited in correlation with decreased adhesion size. Thus, it

seemed that DAPK1 activity was needed to form mature adhesions.

We previously showed that local contractions to sense substrate rigidity correlated with

adhesion formation and maturation[24]. To test if DAPK1 inhibition altered adhesion

formation by affecting rigidity sensing contractions, cells were spread on fibronectin-

coated 500 nm diameter PDMS pillars. When cells reached phase 2 of spreading, they

contracted pairs of pillars toward each other to a peak distance of 60 nm each for about

30 seconds and then relaxed pillars at about the same velocity (2-3 nm/s), consistent

with previous studies[19,24,25]. Inhibition of DAPK1 caused a dramatic decrease in the

velocity of pulling (0.3-0.5 nm/s) with repeated release events (Fig. 2d-f), but did not alter

the maximum pillar displacement (~60 nm) or contractile unit density (Supplementary Fig.

1c). Thus, the decrease in adhesion formation after DAPK1 inhibition correlated with a

decrease in the rate of rigidity-sensing contractions.

DAPK1 co-localizes with Tpm2.1 and targets to focal adhesion sites

Because the rigidity sensing contractions depended upon Tpm2.1[19] and DAPK1, we

then analyzed the distribution of DAPK1 and Tpm2.1 in early spreading of

MCF10A epithelial cells. Both DAPK1 and Tpm2.1 colocalized at the spreading cell

edges, indicating that they could be in a complex there (Fig. 3a). In Tpm2.1-transfected

MDA-MB-231 cells (human breast cancer cell line) which normally lacked endogenous

Tpm2.1[26] but were rigidity-dependent for growth with Tpm2.1[1], DAPK1 overlapped with

Tpm2.1 at the cell edge. In contrast, further back from the edge, where Tpm2.1 was also

present, decorating actin filaments, no overlap was observed (Fig. 3b). Thus, it seemed

that Tpm2.1 and DAPK1 concentrated at active edge regions but not at all sites of

Tpm2.1 concentration.

To dynamically track DAPK1 in cells without causing cell blebbing and apoptosis due to

its overexpression, we next used an EGFP-DAPK1 K42A construct (the inactive form of

DAPK1)[27] in MEFs. During early spreading of MEFs on fibronectin-coated glass, EGFP-
DAPK1 K42A showed significant overlap with paxillin in focal adhesion sites, although
the majority of DAPK1 was in the older portion of the adhesions (the edge closer to the
nucleus) (Fig. 3c). We further confirmed in static images that DAPK1 WT (in the few
cells that did not die) localized to focal adhesions like DAPK1 K42A (Fig. 3d;
Supplementary Fig. 2a). Similar results were observed on fibronectin-coated pillars with
either the DAPK1 construct or anti-DAPK1 antibody (Supplementary Fig. 2b, c). These
results all indicated that DAPK1 was part of adhesions on fibronectin-coated surfaces in
cells that were rigidity-dependent for growth.
174

The phosphorylation of Tpm2.1 by DAPK1 is important for adhesion formation but
increases sensitivity to *anoikis*
Since DAPK1 was found to phosphorylate Tpm2.1 on S283[11], the inhibition of DAPK1
activity may have altered cell adhesion formation through the inhibition of Tpm2.1
phosphorylation. To test this possibility, we transfected MDA-MB-231 cells with mutant
forms of Tpm2.1 which were blocked for phosphorylation, Tpm2.1 S283A, or mimicked
the phosphorylated form, Tpm2.1 S283E. Adhesion size depended upon Tpm2.1
transfection and with the non-phosphorylatable mutant, small adhesions formed that
were similar in size to non-transfected MDA-MB-231 cells. In contrast, with the Tpm2.1
S283E mutant, larger adhesions formed and cells spread to larger areas similar to
Tpm2.1 WT (Fig. 4a, b). In the Tpm2.1 S283A-transfected cells, there was increased
resistance to apoptosis on soft surfaces, whereas the Tpm2.1 S283E cells had greater
sensitivity to apoptosis on soft surfaces than Tpm2.1 WT transfected ones (Fig. 4c).
Thus, the phosphorylation of Tpm2.1 at the DAPK1 phosphorylation site was important
for adhesion formation in MDA-MB-231 cells but facilitated *anoikis* on soft substrates.
190

Talin1 head aids in the localization of DAPK1 to the adhesions and facilitates
apoptosis
Previous studies reported that DAPK1 interacted with talin1 through the head domain[15].
In a separate study, we also showed that the assembly of adhesions depended upon
talin1 cleavage by calpain and cell growth depended upon the presence of the talin1 rod
in the absence of the head[18]. To test if talin1 cleavage was needed for DAPK1 assembly
into adhesions, we transfected non-cleavable (NC) talin1, WT talin1 and the rod and
head fragments separately in talin1-/- fibroblasts. With WT talin1 and the talin1 head, we

found that DAPK1 assembled into focal adhesions; however, neither the talin1 rod nor
NC talin1 supported adhesion assembly with DAPK1 (Fig. 5a). Thus, although the rod
was needed for growth, the head fragment was needed for DAPK1 assembly in
adhesions, which supported previous findings of an interaction of DAPK1 with the talin
head[15]. In DAPK1 WT transfected talin1-/- cells with talin1 constructs or MEFs treated
with a calpain inhibitor (to prevent talin cleavage), talin1 head and talin1 cleavage
caused an increase in DAPK1 apoptotic function (Fig. 5b, c). When talin1-/- cells were
cultured on soft surfaces, they were protected against apoptosis whereas with the WT
talin1 they apoptosed on soft surfaces. Rescue of talin1-/- cells with NC-talin1 did not
restore apoptosis on soft surfaces but expression of the talin1 head domain caused
greater apoptosis on soft surfaces than did WT talin1 (Supplementary Fig. 3). Thus, the
cleavage of talin1 by calpain produced the talin1 head domain fragment that was needed
for DAPK1 to assemble in focal adhesions and to catalyze apoptosis on soft surfaces.
212

Dynamics of DAPK1 during cell spreading on dual stiffness pillars
To confirm that DAPK1 recruitment to adhesions was dependent upon local rigidity, we
prepared pillar substrates that had sub-cellular regions with a 20-fold difference in pillar
rigidity by exposing PDMS pillars to UV light in localized regions[28]. When MEF cells
transfected with EGFP-DAPK1 K42A and mCherry-paxillin were spread on those dual
stiffness pillars for 30 minutes, both DAPK1 and paxillin staining was lower on the soft
than the rigid pillars (Fig. 6a). In time-lapse studies of DAPK1 recruitment to pillars,
EGFP-DAPK1 K42A accumulated equally rapidly on soft and stiff pillars at early times,
but was then released from the soft pillars after 1-2 minutes whereas it remained
associated with the rigid pillars (Fig. 6b, c). We then followed single pillars during the
contraction and relaxation to correlate the recruitment level of DAPK1 with contractile
unit force. After the peak of contraction force, DAPK1 intensity on the soft pillars
decreased rapidly (Fig. 6d). Thus, the recruitment of DAPK1 to adhesions was rapid but
there was also a rapid release from soft pillars. To further determine if DAPK1
recruitment to or release from soft pillars was altered, we measured photobleaching
recovery of EGFP-DAPK1 at soft and rigid pillars. The recovery rates were the same on
both, indicating that binding was unaltered but release was accelerated on soft surface
(Supplementary Fig. 4a, b). Thus, it seems that soft matrices cause the rapid release of
DAPK1 from adhesion sites but recruitment continues.
232

DAPK1 apoptotic function is controlled by Src phosphorylation and PTPN12

In previous studies, tyrosine kinase/phosphatase activities were implicated in the regulation of DAPK1[10]. To test if the vicinal tyrosines in DAPK1 (Y490/Y491) were involved, we overexpressed EGFP-DAPK1 mutants Y490D/Y491D (DYD, phospho-mimic mutant), Y490F/Y491F (DYF, non-phosphorylated mutant) or Y490F/Y491F-K42A (DYF-K42A, non-phosphorylated inactive mutant) in MEFs. Most of the Y-D mutation transfected cells (>80%) had concentrations of DAPK1 in adhesion sites, whereas there was much less DAPK1 assembly in adhesion sites with the DYF (~30%) or DYF-K42A (~20%) transfected cells caused (Fig. 7a). In an apoptosis assay, the Y-F mutation enabled normal apoptosis whereas the Y-D mutation inhibited apoptosis as did the Y-F double mutation of the inactive DAPK1-K42A (Fig. 7b). To test for the possible involvement of the Src family tyrosine kinases (SFK), we added the general SFK inhibitor PP2 and found that there was less DAPK1 assembled into adhesions (Supplementary Fig. 5a, b), as well as a dramatic decrease in adhesion size as had been reported before because of the inhibition of rigidity sensing contractions[17]. We next tested for apoptosis in the presence of PP2 and found a dramatic increase in DAPK1-induced cell death (Supplementary Fig. 5c). Since the SFKs also activated EGFR in early spreading[17], we tested for the effect of EGFR inhibition on apoptosis and found enhanced cell death after EGFR inhibition (Supplementary Fig. 5c). Thus, tyrosine phosphorylation of DAPK1 on the vicinal tyrosines enabled assembly of DAPK1 into adhesions and inhibited apoptosis whereas the unphosphorylated DAPK1 couldn't assemble in adhesions but supported apoptosis.

There was reason to believe that DAPK1 assembly into the adhesions was a normal part of the cell-matrix interaction and that a phosphatase was needed to disassemble adhesions on soft surfaces and to activate DAPK1. A major tyrosine phosphatase that was implicated in cancer and motility was PTPN12[29,30]. To test for its possible involvement, we added an inhibitor of PTPN12 and found that it blocked DAPK1 apoptotic function in MEFs. Further, we observed an enhancement of apoptosis if we overexpressed PTPN12 (Fig. 7c). Thus, it seemed that tyrosine phosphorylation inhibited DAPK1 and apoptosis whereas PTPN12 activity was needed for activation of apoptosis on soft surfaces. These findings were summarized in a model diagram (Fig. 7d).

Discussion

These studies show that DAPK1 has a major role both in the formation of matrix adhesions on rigid surfaces at early times and in the apoptosis of cells on soft surfaces. Both processes involve DAPK1 assembly in adhesions through its phosphorylation of Tpm2.1 at S283, as well as the proteolysis of talin1 to produce the head fragment. DAPK1 activity is needed for normal assembly of focal adhesions and proper rigidity sensing contractions. Further, the phosphorylation of DAPK1 on the vicinal tyrosine residues (Y490/491) by Src or EGFR is critical for its inactivation at adhesions. On sub-micrometer pillars with sub-cellular areas of different stiffness, the binding of DAPK1 to both soft and rigid pillars is initially similar, but once a maximal force is reached, rapid release is only found in the soft regions. DAPK1 activation, release, and adhesion disassembly all appear to correlate with the dephosphorylation of DAPK1 by PTPN12. Thus, the rigidity dependent activation of apoptosis involves DAPK1 activity in the formation of adhesions through Tpm2.1 and rigidity-dependent dephosphorylation of DAPK1.

An earlier study found that DAPK1 had a role in cell migration and polarization through an interaction with integrin and the talin1 head domain[15]. This is consistent with its function in the formation of contractile adhesions and rigidity sensing; however, the longer-term reactions to rigid surfaces involve activation of motility responses beyond early rigidity sensing. Actin-dependent membrane extension correlates with rigidity sensing and fully spread cells have few rigidity-sensing contractions because they have few extensions[17]. The cycle of mechano-testing and response can result in migration if there is polarization with a single leading edge. In the case of transformed cancer cells, there is a major change in cell state since they lack rigidity sensing contractions and Tpm2.1 in adhesions. It seems that normal rigidity sensing involves the recruitment of tyrosine phosphorylated DAPK1 along with Tpm2.1 to the contractile units. If the surface is rigid, then endogenous DAPK1 is stabilized in the adhesions along with Tpm2.1, talin1 head, and other focal adhesion proteins. If the surface is soft, then a phosphatase, such as PTPN12, dephosphorylates DAPK1 and causes its activation as well as the disassembly of the adhesions. In studies of immune cells, the phosphatase LAR dephosphorylates DAPK1 but that is not related to cell mechanics[10]. The role of PTPN12 in activating *anoikis* is consistent with its role as a tumor suppressor[30] and it also has major effects on adhesion lifetime[31].

The role of the tyrosine kinases in inactivating DAPK1 is consistent with previous studies

that show high levels of phosphorylation of the vicinal tyrosine residues in cancer as well

as fibroblasts[10]. In these studies, DAPK1 is inactivated upon phosphorylation by Src

either directly or downstream through EGFR activation. In earlier studies, we found that

Src activated EGFR or HER2 in cell spreading to catalyze rigidity sensing[17]. Here we

find that inhibitors of either will cause increased apoptosis, indicating that DAPK1

normally is phosphorylated in the adhesions. Further, the inhibition of PTPN12 will block

DAPK1 activation. The level of pS308 decreases on soft surfaces over time and the level

of DAPK1 expression appears to increase as observed in other studies of DAPK1

activation but that may be through continued expression without assembly into

adhesions on the soft surfaces. There are few rigidity sensing contractions on soft

surfaces and few chances for assembly of DAPK1 into adhesion complexes with

tyrosine phosphorylation[17].

It is logical that the activation of apoptosis on soft surfaces would involve engagement of

the death-associated protein kinase with the rigidity sensing process at integrin

adhesions. Rigidity sensing involves the assembly of sarcomere-like contractile units at

integrin adhesion sites and the subsequent pulling of those adhesions to produce a

displacement of 100 nm. If the force developed is less than about 25 pN, then the matrix

is considered soft and the adhesions disassemble, releasing activated DAPK1. Based

upon these findings we hypothesize that Tpm2.1 and talin1 head domain form a complex

with tyrosine phosphorylated DAPK1 that dissociates on soft surfaces due to the tyrosine

phosphatase, PTPN12 (Fig. 7d).

The complex nature of the rigidity sensing module indicates that it integrates

mechanosensing with several output signals depending upon the matrix rigidity.

Although those output signals can vary dramatically for different cell types, the basic

sensory machine appears to have many common elements. For example, the receptor-

like-protein-tyrosine-phosphatase alpha (RPTPα) is needed for rigidity sensing in both

fibroblasts and neurons but in hippocampal neurons the loss of rigidity sensing results in

long straight neurites whereas in fibroblasts it results in transformed growth[32]. Since

DAPK1 is part of rigidity sensing modules in fibroblasts, those cells can activate

apoptosis on soft surfaces whereas DAPK1 is not linked to matrix rigidity when the

sensory modules are missing cytoskeletal components and do not assemble as in
cancer cells. Signaling from Src and RTKs is part of the rigidity-sensing module and that
is needed to inactivate DAPK1 as well as activate growth because of force-dependent,
tyrosine phosphorylation on rigid substrates. The competing tyrosine phosphatases such
as PTPN12 and possibly LAR are designed to keep cells from growing inappropriately
when the matrix is soft but are not operative in the absence of the rigidity sensing
modules. This raises many questions about how DAPK1 is inactivated when the rigidity
sensors are missing and how are the phosphatases recruited to the adhesions on soft
surfaces.
344
Materials and Methods
Cell culture
Mouse embryonic fibroblast cells (MEFs) were generated by J. Sap's laboratory[33]. MDA-
MB-231 and MCF10A cells were obtained from J. Groves (University of California, USA
and Mechanobiology Institute, National University of Singapore, Singapore). The
talin1−/− fibroblast cell line dj26.28 was generated and maintained as described
previously[34]. The MEFs and MDA-MB-231 cells were cultured in Dulbecco's Modified
Eagle Medium (DMEM) (Thermo Fisher Scientific) supplemented with 10% fetal bovine
serum (FBS) (Atlanta Biologicals) and 100 IU ml^{-1} Penicillin-Streptomycin (Sigma) at
37°C and 5% CO_2. MCF10A cells were cultured at 37°C in a 5% CO_2 incubator in
DMEM (Thermo Fisher Scientific) supplemented with 20 ng ml^{-1} EGF (Thermo Fisher
Scientific), 10 ng ml^{-1} bovine insulin (Sigma), 500 ng ml^{-1} hydrocortisone, 5% horse
serum albumin (Thermo Fisher Scientific), and 100 IU ml^{-1} Penicillin-Streptomycin
(Sigma). Imaging experiments were conducted using starvation medium without phenol
red or in Ringer's buffer (150 mM NaCl, 5 mM KCl, 1 mM $CaCl_2$, 1 mM $MgCl_2$, 20 mM
Hepes, and 2 g/L glucose, pH 7.4). Pharmacological inhibitors were as follows: DAPK1
inhibitor (100 nM, EMD Millipore), EGFR inhibitor Gefitinib (10 nM, Santa Cruz), PP2
(200 nM, abcam), calpain Inhibitor ALLN (100 μM, Santa Cruz), PTPN12 inhibitor (1.5
μM, EMD Millipore). Cells were trypsinized using TrypLE (Thermo Fisher Scientific) on
the following day and suspended in Ringer's buffer at 37°C for 30 minutes prior to plating
on human plasma fibronectin (10 μg/ml, Roche) coated glass-bottom dishes (MatTek) or
pillars. The 0.2 kPa PDMS gels used for testing cells' growth on soft matrices were
purchased from Soft Substrates^{TM}.
368

Transfection and plasmids

Transfections were carried out 1 day before measurements using Lipofectamine Plus Reagent (Invitrogen) according to the manufacturer's instructions. Expression vectors encoding the following fluorescent fusion proteins were used: EGFP-DAPK1 WT, EGFP-DAPK1 K42A, mCherry-DAPK1 WT, mCherry-DAPK1 K42A, YFP-Tpm2.1 WT, YFP-Tpm2.1 S283E (phospho-mimic mutant), YFP-Tpm2.1 S283A (non-phosphorylated mutant), mCherry-paxillin, GFP-talin1 WT, mRuby-talin1 head, mCherry-talin1 rod, GFP-non-cleavable talin1, EGFP-DAPK1-DYD (phospho-mimic mutant), EGFP-DAPK1-DYF (non-phosphorylated mutant), EGFP-DAPK1-DYF-K42A (non-phosphorylated inactive mutant), mCherry-PTPN12. Cells were seeded into a 6-well dish on day 0 and transfected with control shRNA (Sigma) or DAPK1 shRNA (Sigma) using Lipofectamine Plus Reagent (Invitrogen) on day 1, followed by selection in 2 µg/ml puromycin for 72 hours.

Pillar fabrication

Fabrication of pillar arrays with diameters of 500 nm was performed as previously reported[24]. Polydimethylsiloxane (PDMS, Sylgard 184, Dow Corning) was mixed thoroughly with its curing agent (10:1) for at least 5 minutes, centrifuged (2000 RPM, 3 minutes, room temperature) and degassed in a vacuum for about 10 minutes. A drop of PDMS was then placed on the top of the Poly (methyl methacrylate) (PMMA) mold and degassed again in a vacuum for a few minutes. At the same time, a glass-bottom petri dish (No.0 Coverslip, MatTek) was treated with O_2 plasma for 1 minute, in order to increase adhesion of the PDMS to the glass after curing. The mold with PDMS was then inverted and placed onto the bottom of the petri dish, and a 1 gram weight was placed on the top of mold to make the PDMS thinner. The PDMS was then cured at 70°C for 12-14 hours, in order to achieve a Young's modulus of 2 ± 0.1 MPa. The mold was then peeled off the PDMS while immersed in ethanol. Individual pillars were 500 nm in diameter with a center to center distance of 1µm.

For the dual stiffness pillar fabrication, Coumarin 343 (Sigma-Aldrich), which will be bleached under UV treatment, was premixed in PDMS for the marking of areas with different rigidity. The dyed PDMS was applied to the master molds, which were spun at 6600 RPM for 3 seconds to ensure even coating. After the PDMS was cured at 70°C for 12-14 hours, it was carefully peeled off the molds and placed (pillar-side up) onto the O_2

plasma-treated glass-bottom petri dish. Prior to UV/Ozone treatment, nickel TEM grids

(SPI supplies) were laid on the pillar tops. Pillar substrates covered by grids were then

placed in the UV/Ozone chamber (Bioforce Nanosciences UV/Ozone ProCleaner Plus)

for 2 hours.

Traction force measurements

Cells were spreading on pillar arrays coated with fibronectin (10 μg/ml, Roche). Time

lapse imaging of pillars was performed with bright-field microscopy using an Orca-flash

2.8 camera (Hamamatsu) attached to an inverted microscope (Olympus IX81)

maintained at 37° C running MicroManager software (UCSF). Images were recorded at 1

Hz using a 100x objective (1.4 NA oil immersion, Olympus). Videos were processed with

ImageJ (National Institutes of Health) using the Nano Tracking plugin to track the

position of pillars. The time-series positions of all pillars in contact with the cell were fed

into a MatLab program (MathWorks) to generate displacement maps as explained

previously[17,19].

Immunostaining and fluorescence microscopy

Cells were fixed for 15 minutes with 4% formaldehyde solution (Sigma, diluted from 37%

in PBS). Fixed cells were then permeabilized with 0.2% Triton X-100 in PBS (V/V,

Sigma) for 5 minutes, and blocked in 2% bovine serum albumin (BSA) in PBS (W/V,

Sigma) at room temperature for one hour. Primary antibody was diluted in 1% BSA and

incubated on samples overnight at 4° C. Secondary antibody was diluted in 1% BSA and

incubated on samples for 1 hour at room temperature. For immunostaining, the following

primary antibodies were used: anti-paxillin (Abcam, dilution 1:250), anti-DAPK1 (BD

Biosciences, dilution 1:50), anti-Tpm2.1 (Abcam, dilution 1:1000), anti-pSer308 DAPK1

(LSBio, dilution 1:100), anti-Annexin V (Abcam, dilution 1:100). Secondary antibodies

were Alexa Fluor 488 goat anti-mouse IgG (H+L) and Alexa Fluor 555 goat anti-rabbit

IgG (H+L) (Thermo Fisher Scientific).

Total internal reflection fluorescence (TIRF) images were taken using an Olympus IX81

fluorescence microscope with a 60x, 1.45 NA oil-immersion objective and a Cool Snap

FX cooled CCD camera (Photometrics) controlled by MicroManager software (UCSF).

Confocal microscopy was performed on a Zeiss LSM 700 laser-scanning confocal

microscope, using a 63x/1.4 NA oil-immersion objective. Quantification of images was performed with ImageJ (National Institutes of Health).

For the fluorescence recovery after photobleaching (FRAP) experiment, MEFs transfected with EGFP-DAPK1 K42A or EGFP-DAPK1 WT were allowed to spread on fibronectin-coated pillars for 20 minutes and then the focal adhesion areas were bleached using fast bleach on Zeiss LSM 700. Fluorescence intensity recovery was simultaneously recorded at 1 frame per 2 seconds. The images were then corrected for background using ImageJ (National Institutes of Health). Single exponentials were then fitted to the recovery profiles in Excel and SigmaPlot 12.0 software (Systat Software Inc.) to calculate half-times ($t_{1/2}$).

Statistical analysis

Statistical analysis and graph plotting were performed using SigmaPlot 12.0 software (Systat Software Inc.) and Matlab (Math Works). Two-tailed Student's t test was performed when two cases were compared. One-way-ANOVA was performed for multi-group comparison. P values <0.05 were considered statistically significant.

Acknowledgements

We thank all the members of Sheetz lab for their kind help. This work was funded by National Institutes of Health (NIH) grant "Analysis of 120 nm local contractions linked to rigidity sensing" (1 R01 GM100282-01), National Institutes of Health (NIH) grant "Tropomyosin and tyrosine kinases in mechanics of cancer" (5 R01 GM113022-02). H.W. was supported by a Marie Curie International Outgoing Fellowship within the Seventh European Commission Framework Programme (PIOF-GA-2012-332045), and is an incumbent of the David and Inez Myers Career Advancement Chair in Life Sciences. M.P.S is supported by NUS grants and MBI, National University of Singapore.

Author Contributions

R.Q. H.W. M.S. and M.P.S conceived the study and designed the experiments; R.Q. and H.W. performed the experiments; R.Q. and H.W. analyzed the data; R.Q. H.W. and M.P.S wrote and prepared the manuscript.

Competing interests

The authors declare no conflict of interest.

Materials & Correspondence

Correspondence and requests for materials should be addressed to M.P.S.
(ms2001@columbia.edu) or H.W. (haguyw@technion.ac.il).

References

1. Yang, B. et al. Stopping Transformed Growth with Cytoskeletal Proteins: Turning a
Devil into an Angel. *bioRxiv* doi: 10.1101/221176 (2018).

2. Bialik, S., Kimchi, A. The DAP-kinase interactome. *Apoptosis* **19,** 316-328 (2014).

3. Cohen, O., Feinstein, E., Kimchi, A. DAP-kinase is a Ca2+/calmodulin-dependent,
cytoskeletal-associated protein kinase, with cell death-inducing functions that depend on
its catalytic activity. *EMBO J* **16,** 998-1008 (1997).

4. Raval, A. et al. Downregulation of death-associated protein kinase 1 (DAPK1) in
chronic lymphocytic leukemia. *Cell* **129,** 879-890 (2007).

5. Michie, A.M., McCaig, A.M., Nakagawa, R., Vukovic, M. Death-associated protein
kinase (DAPK) and signal transduction: regulation in cancer. *FEBS J* **277,** 74-80 (2010).

6. Lévy, D. et al. Death-associated protein kinase loss of expression is a new marker for
breast cancer prognosis. *Clin Cancer Res* **10,** 3124-3130 (2004).

7. Ivanovska, J. et al. DAPK loss in colon cancer tumor buds: implications for migration
capacity of disseminating tumor cells. *Oncotarget* **6,** 36774-36788 (2015).

8. Cohen, O. et al. DAP-kinase participates in TNF-alpha- and Fas-induced apoptosis
and its function requires the death domain. *J Cell Biol* **146,** 141-148 (1999).

9. Shohat, G. et al. The pro-apoptotic function of death-associated protein kinase is

controlled by a unique inhibitory autophosphorylation-based mechanism. *J Biol Chem*

276, 47460-47467 (2001).

10. Wang, W.J. et al. The tumor suppressor DAPK is reciprocally regulated by tyrosine

kinase Src and phosphatase LAR. *Mol Cell* **27,** 701-716 (2007).

11. Houle, F., Poirier, A., Dumaresq, J., Huot, J. DAP kinase mediates the

phosphorylation of tropomyosin-1 downstream of the ERK pathway, which regulates the

formation of stress fibers in response to oxidative stress. *J Cell Sci* **120,** 3666-3677

(2007).

12. Bialik, S., Bresnick, A.R., Kimchi, A. DAP-kinase-mediated morphological changes

are localization dependent and involve myosin-II phosphorylation. *Cell Death Differ* **11,**

631-644 (2004).

13. Geeves, M.A., Hitchcock-DeGregori, S.E., Gunning, P.W. A systematic

nomenclature for mammalian tropomyosin isoforms. *J Muscle Res Cell Motil* **36,** 147-

153 (2015).

14. Bharadwaj, S., Thanawala, R., Bon, G., Falcioni, R., Prasad, G.L. Resensitization of

breast cancer cells to *anoikis* by tropomyosin-1: role of Rho kinase-dependent

cytoskeleton and adhesion. *Oncogene* **24,** 8291-8303 (2005).

15. Kuo, J.C., Wang, W.J., Yao, C.C., Wu, P.R., Chen, R.H. The tumor suppressor

DAPK inhibits cell motility by blocking the integrin-mediated polarity pathway. *J Cell Biol*

172, 619-631 (2006).

16. Yang, B. et al. Mechanosensing Controlled Directly by Tyrosine Kinases. *Nano Lett*

16, 5951-5961 (2016).

17. Saxena, M. et al. EGFR and HER2 activate rigidity sensing only on rigid matrices.

Nat Mater **16,** 775-781 (2017).

18. Saxena, M., Changede, R., Hone, J., Wolfenson, H., Sheetz, M.P. Force-Induced

Calpain Cleavage of Talin Is Critical for Growth, Adhesion Development, and Rigidity

Sensing. *Nano Lett* **17,** 7242-7251 (2017).

19. Wolfenson, H. et al. Tropomyosin controls sarcomere-like contractions for rigidity

sensing and suppressing growth on soft matrices. *Nat Cell Biol* **18,** 33-42 (2016).

20. Meacci, G. et al. α-Actinin links extracellular matrix rigidity-sensing contractile units

with periodic cell-edge retractions. *Mol Biol Cell* **27,** 3471-3479 (2016).

21. Wang, W.J., Kuo, J.C., Yao, C.C., Chen, R.H. DAP-kinase induces apoptosis by

suppressing integrin activity and disrupting matrix survival signals. *J Cell Biol* **159,** 169-

179 (2002).

22. Inbal, B., Bialik, S., Sabanay, I., Shani, G., Kimchi, A. DAP kinase and DRP-1

mediate membrane blebbing and the formation of autophagic vesicles during

programmed cell death. *J Cell Biol* **157,** 455-468 (2002).

23. Döbereiner, H.G., Dubin-Thaler, B., Giannone, G., Xenias, H.S., Sheetz, M.P.

Dynamic phase transitions in cell spreading. *Phys Rev Lett* **93,** 108105 (2004).

24. Ghassemi, S. et al. Cells test substrate rigidity by local contractions on

submicrometer pillars. *Proc Natl Acad Sci U S A* **109,** 5328-5333 (2012).

25. Iskratsch, T., Wolfenson, H., Sheetz, M.P. Appreciating force and shape-the rise of

mechanotransduction in cell biology. *Nat Rev Mol Cell Biol* **15,** 825-833 (2014).

26. Raval, G.N. et al. Loss of expression of tropomyosin-1, a novel class II tumor

suppressor that induces *anoikis*, in primary breast tumors. *Oncogene* **22,** 6194-6203

(2003).

27. Chen, C.H. et al. Bidirectional signals transduced by DAPK-ERK interaction promote

the apoptotic effect of DAPK. *EMBO J* **24,** 294-304 (2005).

28. Song, J., Tranchida, D., Vancso, G.J. Contact Mechanics of UV/Ozone-Treated

PDMS by AFM and JKR Testing: Mechanical Performance from Nano- to Micrometer

Length Scales. *Macromolecules* **41,** 6757-6762 (2008).

29. Sastry, S.K., Lyons, P.D., Schaller, M.D., Burridge, K. PTP-PEST controls motility

through regulation of Rac1. *J Cell Sci* **115,** 4305-4316 (2002).

30. Li, J. et al. Loss of PTPN12 Stimulates Progression of ErbB2-Dependent Breast

Cancer by Enhancing Cell Survival, Migration, and Epithelial-to-Mesenchymal Transition.

Mol Cell Biol **35,** 4069-4082 (2015).

31. Angers-Loustau, A. et al. Protein tyrosine phosphatase-PEST regulates focal

adhesion disassembly, migration, and cytokinesis in fibroblasts. *J Cell Biol* **144,** 1019-

1031 (1999).

32. Kostic, A., Sap, J., Sheetz, M.P. RPTPalpha is required for rigidity-dependent

inhibition of extension and differentiation of hippocampal neurons. *J Cell Sci* **120,** 3895-

3904 (2007).

33. Su, J., Muranjan, M., Sap, J. Receptor protein tyrosine phosphatase alpha activates

Src-family kinases and controls integrin-mediated responses in fibroblasts. *Curr Biol* **9,**

505-511 (1999).

34. Priddle, H. et al. Disruption of the talin gene compromises focal adhesion assembly

in undifferentiated but not differentiated embryonic stem cells. *J Cell Biol* **142,** 1121-1133

(1998).

Figure Legends

Fig. 1 Inhibition of DAPK1 contributes to cell growth on soft matrices. (**a**) DIC images of

MEFs and M10A cells with or without DAPK1 inhibitor after 2 days in culture on 0.2 kPa

gels. Scale bar: 10 µm. Apoptotic cells were identified for appearance of apoptotic

morphology including membrane blebbing and cell rounding. Live cells retained their

normal spread morphology. Graphs represent means ± SEM of at least two independent

experiments. ***P <0.001. (**b**) DAPK1-Ser308 phosphorylation was monitored by

immunofluorescence in MEFs plated on 0.2 kPa gels. Immunofluorescence shows that

DAPK1 was dephosphorylated at Ser-308 and thus activated on soft surfaces over time.

Representative pictures are shown on the left panel for each condition. Scale bar: 10 µm.

Graphs represent means ± SEM. Experiment was repeated twice.

Fig. 2 DAPK1 regulates adhesion growth and force production. (**a**) Micrographs showing

the distribution of paxillin in MEFs treated with or without DAPK1 inhibitor. MEFs treated

with or without DAPK1 inhibitor were fixed after spreading on fibronectin-coated glass

dishes for 30 minutes, 60 minutes or 16 hours, followed by anti-paxillin/AlexaFluor 488

immunostaining. (**b**) The adhesion sizes were quantified (n>400 adhesions from >10

cells in each case, mean ± SEM). ***P <0.001. (**c**) Average area ± s.e.m of cells were

also determined (n>10 cells in each case). ***P <0.001. Experiment was repeated three

times. (**d**) Displacements (red arrows) of pillars that show the contractile pair (green

arrows marked with yellow oval) at the periphery of a spreading MFF. Scale bar: 1 µm.

(**e**) Displacements versus time of contractile pillars in MEFs treated with or without

DAPK1 inhibitor. (**f**) Mean ± SEM of the T1/2 (time of contraction above half-maximal

displacement) distributions of pillar displacements by MEFs before and after treatment

with DAPK1 inhibitor. ***P < 0.001. N > 30 pillars in each case. Experiment was

repeated twice.

Fig. 3 DAPK1 co-localizes with tropomyosin2.1 and targets to focal adhesion sites. (**a**)

MCF10A cells were fixed after spreading on fibronectin-coated glass dishes for 30

minutes, followed by anti-tropomyosin2.1/AlexaFluor 488 and anti-DAPK1/AlexaFluor

555 immunostaining. Scale bar: 10 µm. (**b**) MDA-MB-231 cells were transfected with

YFP-Tpm2.1 WT. Cells were further analyzed by anti-DAPK1/AlexaFluor555

immunostaining. Scale bar: 10 µm. (**c**) Total internal reflection fluorescence (TIRF)

image of MEF co-transfected with EGFP-DAPK1 K42A and mCherry-paxillin after 30

minutes spreading on fibronectin-coated glass dish. Fluorescence intensity line profiles

on the right are from the area of the representative adhesions covered by the white

arrow in the merged images. Scale bar: 10 µm. (**d**) MEFs were co-transfected with

EGFP-talin and mCherry-DAPK1 WT. Cells were subsequently allowed to spread for 30

minutes on fibronectin-coated glass dish. Scale bar: 10 µm.

Fig. 4 The phosphorylation of Tpm2.1 by DAPK1 is important for adhesion formation but
facilitates *anoikis* on soft substrates. (**a**) Left: micrographs showing the distribution of
paxillin in MDA-MB-231 cells transfected with YFP-Tpm2.1 WT, YFP-Tpm2.1 S283E
(phospho-mimic mutant), YFP-Tpm2.1 S283A (non-phosphorylated mutant) or YFP-
Vector respectively after 30 minutes spreading on fibronectin-coated glass dishes,
followed by anti-paxillin/AlexaFluor 555 immunostaining. Scale bar: 10 μm. Right: the
adhesion sizes were quantified (n>400 adhesions from >10 cells in each case, mean ±
SEM), ***P <0.001. (**b**) Average area ± s.e.m of transfected cells in each group were
determined (n>10 cells in each case). *P <0.05, **P <0.01. (**c**) MDA-MB-231 cells were
transfected with YFP-Tpm2.1 WT, YFP-Tpm2.1 S283E, YFP-Tpm2.1 S283A or YFP-
Vector respectively for one day and replated on fibronectin-coated 0.2 kPa gels for one
day. The percentage of transfected cells that were still alive was quantified. The mean ±
SEM of at least two independent experiments is described. For each experiment,
100~150 cells were analyzed for each transfection point. **P <0.01, ***P <0.001.

Fig. 5 Talin1 head aids in the localization of DAPK1 to the adhesions and facilitates
apoptosis. (**a**) Talin1-/- cells were co-transfected with DAPK1 and talin1 constructs
(talin1 WT, talin1 head, talin1 rod, non-cleavable talin1) or vector. Cells were spread on
fibronectin (30 minutes). Scale bar: 10 μm. (**b**) Talin1-/- cells were co-transfected with
DAPK1 WT and talin1 constructs (talin1 WT, talin1 head, talin1 rod, NC talin1) or vector
for one day. The percentage of co-transfected cells that were still alive was quantified
(mean ± SEM of ≥2 experiments with 100~150 cells; *P <0.05, **P <0.01) (**c**) MEFs
transfected with EGFP-DAPK1 WT treated with DMSO or calpain inhibitor ALLN (100
μM). The percentage of transfected cells that were still alive was quantified. (mean ±
SEM of ≥2 experiments with 100~150 cells; *P <0.05).

Fig. 6 Dynamics of DAPK1 during cell spreading on dual stiffness pillars. (**a**) MEFs were
co-transfected with EGFP-DAPK1 K42A and mCherry-paxillin. Cells were spread (30
minutes) on dual stiffness pillars (9kPa (Coumarin 343) and 170kPa), Scale bar: 10 μm.
The leading edge protein relative intensity was normalized by dividing the average
intensity value within 3 micrometers of the cell edge in stiff region or soft region by the
average intensity value of the cell edge in stiff region of each cell (stiff /stiff or soft /stiff).
The relative intensity of DAPK1 or paxillin is presented in the lower panels. **P <0.01.
N=at least 5 different cells in each group. Graphs represent mean ± SEM. (**b**) Left:

confocal image showing a cell transfected with EGFP-DAPK1 K42A and mCherry-
paxillin on dual stiffness pillars. Scale bar: 10 μm. Right: one adhesion site of the left cell
corresponding to a single pillar in a stiff or soft region was tracked during cell spreading.
The tracked pillar is marked by a yellow box. The relative fluorescence of both molecules
was plotted. (**c**) Quantification of relative increase in DAPK1 and paxillin fluorescence
over time on dual stiffness pillars within 3 micrometers of outward curving cell edges.
Four cells were measured on 2 different days. Graphs represent means ± SEM. (**d**) The
time course of recruitment of DAPK1 was measured during pillar contraction and
relaxation. Displacements versus time of contractile pillars are shown in the lower panel.
The peak of the pillar contraction force is marked by a red line.

Fig. 7 DAPK1 apoptotic activity is controlled by Src phosphorylation and PTPN12. (**a**)
Micrographs showing the distribution of DAPK1 in MEFs transfected with EGFP-DAPK1-
DYD (phospho-mimic mutant), EGFP-DAPK1-DYF (non-phosphorylated mutant) or
EGFP-DAPK1-DYF-K42A (non-phosphorylated inactive mutant) respectively after 30
minutes spreading on fibronectin-coated glass dishes. Scale bar: 10 μm. (**b**) MEFs were
transfected with EGFP-DAPK1 WT, EGFP-DAPK1-DYD, EGFP-DAPK1-DYF or EGFP-
DAPK1-DYF-K42A respectively for one day. The percentage of transfected cells that
were still alive was quantified. The mean ± SEM of at least two independent experiments
is described. For each experiment, 100~150 cells were analyzed for each transfection
point. *$P < 0.05$, ***$P < 0.001$. (**c**) MEFs were transfected with EGFP-DAPK1 WT either
co-transfected with mCherry-PTPN12 or treated with PTPN12 inhibitor. The percentage
of co-transfected cells that were still alive was quantified. The mean ± SEM of at least
two independent experiments is described. For each experiment, 100~150 cells were
analyzed for each transfection point. **$P < 0.01$, ***$P < 0.001$. (**d**) Schematic of a working
model of DAPK1 shuttling depending on substrate rigidity. The presence of a rigid
substrate keeps DAPK1 at the FA through mechanical tension. In contrast, a soft
substrate induces DAPK1 disassembly from FA and activates DAPK1 apoptotic function.

Figure 1

Figure 2

a Control   DAPK1 inhibitor

Anti-paxillin

10um

b Control, DAPK1 inhibitor

FA Area (μm²) vs 30 min, 60 min, 16 h; *** significance markers

c Control, DAPK1 inhibitor

Cell Area (μm²) vs 30 min, 60 min, 16 h; *** significance markers

d 1um

e Control, DAPK1 inhibitor

Displacement (nm) vs Time (s)

f T1/2 (s) vs Control, DAPK1 inhibitor; *** significance marker

Figure 3

Figure 4

Figure 5

a

b

  c

Figure 6

Figure 7

a

DAPK1-DYD   DAPK1-DYF-K42A   DAPK1-DYF

10um

MEFs

b

Live transfected cells / total transfected cells (100%)

DAPK1 WT   DAPK1-DYD   DAPK1-DYF   DAPK1-DYF-K42A

MEFs

*   *   *   ***

c

Live transfected cells / total transfected cells (100%)

mCherry-vector   mCherry-vector +PTPN12 inhibitor   mCherry-PTPN12

**   ***   ***

EGFP-DAPK1 WT in MEFs

d

Cell Rigidity sensing
P2

EGFR HER2   Activates   DAPK   Tropomyosin   Tropomodulin
α-Actinin   Actin   FLNA
SFK   Myosin II   Adhesion complex
Integrin   AXL   ROR2   Fibronectin

Rigid surface

Talin rod   GROWTH
Calpain 2   Reinforcement
Talin head
VPNM4

Soft surface

↑PTPN12 → ↑DAPK → DEATH
Talin head   Dissasembly
PTPN12