Arginine to glutamine mutation in olfactomedin like 3 (*OLFML3*) is a candidate for severe

goniodysgenesis and glaucoma in the Border Collie dog breed.

Carys A Pugh*, Lindsay L Farrell* [1], Ailsa J Carlisle*, Stephen J Bush*†, Violeta Trejo-

Reveles*, Oswald Matika*, Arne de Kloet‡, Caitlin Walsh‡, Stephen C Bishop* [2], James GD

Prendergast*, Jeffrey J Schoenebeck*, Joe Rainger*, Kim M Summers*§

* The Roslin Institute, Easter Bush EH25 9RG, United Kingdom

† Nuffield Department of Clinical Medicine, University of Oxford, John Radcliffe Hospital,

Oxford OX3 9DU, United Kingdom

‡ Animal Genetics, 1336 Timberlane Rd, Tallahassee, FL 32312, USA

§ Mater Research Institute-UQ, Translational Research Institute, Brisbane, Qld 4012,

Australia

[1] Present address: British Columbia Centre on Substance Abuse, St Paul's Hospital,

Vancouver, BC V6Z 1Y6, Canada

[2] Deceased

Running title: *OLFML3* mutation in Border Collie glaucoma

Key words: Glaucoma, goniodysgenesis, olfactomedin like 3, Border Collie, eye development

Corresponding authors:

Dr Carys Pugh, The Roslin Institute, University of Edinburgh, Easter Bush EH25 9RG, UK

Phone: +441316519221

Email: carys.pugh@roslin.ed.ac.uk

Professor Kim M Summers, Mater Research Institute-UQ, Translational Research Institute,

37 Kent St, Woolloongabba, QLD 4102, Australia

Phone: +61734437655

Email: kim.summers@mater.uq.edu.au

Abstract

Goniodysgenesis is a developmental abnormality of the anterior chamber of the eye. It is

generally considered to be congenital in dogs (*Canis lupus familiaris*), and has been

associated with glaucoma and blindness. Goniodysgenesis and early-onset glaucoma initially

emerged in Border Collies in Australia in the late 1990s and has subsequently been found in Europe

and the USA. The objective of the present study was to determine the genetic basis of

goniodysgenesis in Border Collies. Clinical diagnosis was based on results of examinations by

veterinary ophthalmologists of affected and unaffected dogs from eleven different

countries. Genotyping using the Illumina high density canine SNP chip and whole genome

sequencing were used to identify candidate genetic regions. Expression profiles and

evolutionary conservation of candidate genes were assessed using public databases.

Analysis of pedigree information was consistent with an autosomal recessive mode of

inheritance for severe goniodysgenesis (potentially leading to glaucoma) in this breed. There

was a highly significant peak of association over chromosome 17, with a *p*-value of 2×10^{-13}.

Whole genome sequences of three dogs with glaucoma, three severely affected and three

unaffected dogs identified a missense variant in the olfactomedin like 3 (*OLFML3*) gene in all

six affected animals. This was homozygous in all nine cases with glaucoma and 12 of 14

other severely affected animals. Of 67 reportedly unaffected animals, only one (offspring of

two homozygous affected parents) was homozygous for this variant. The identification of a

candidate genetic region and putative causative mutation will aid breeders to reduce the

frequency of goniodysgenesis and the risk of glaucoma in the Border Collie population.

Introduction

Companion animals including cats, dogs and horses, suffer from a range of diseases, with both genetic and environmental aetiologies. The breed barrier created by registration requirements and breeding practices seeking to maximise compliance of animals with breed specifications, results in increased levels of matings between relatives [1] and means that purebred companion animals are particularly likely to suffer from recessive genetic conditions. Understanding the genetic factors underlying these diseases is important to improve welfare within breeds and the strong linkage disequilibrium in such inbred populations means that small numbers of pedigree animals can be used to identify genetic regions of interest, making them excellent genetic models if the same disease is present in humans.

In response to the identification of causal variants, breeding strategies can be put in place to reduce the prevalence of conditions that impact on animal welfare, especially when a genetic test can be developed. However, the success of these approaches depends on understanding the mode of inheritance and level of genetic contribution to the disease, and ideally on identifying a causative mutation that can be used to develop a test for genetic status.

Primary glaucoma is a condition in which increased ocular pressure damages the retinal ganglion, leading to blindness (reviewed in [2]). In dogs (*Canis lupus familiaris*) it can be preceded by goniodysgenesis (also known as mesodermal dysgenesis), a developmental abnormality of the eye characterised by narrowing or closure of the iridocorneal angle through which the aqueous humour drains. This is associated with alterations in the structure of the pectinate ligament which traverses the drainage angle, called pectinate

ligament dysplasia (PLD) [3] (more properly pectinate ligament abnormality [4]). In the

Leonberger dog breed, goniodysgenesis has been associated with an increased risk of

developing primary closed angle glaucoma (PCAG) [2] and in Flat-Coated Retrievers,

development of glaucoma is positively and significantly related to the severity of

goniodysgenesis [5]. Goniodysgenesis is generally considered to be congenital in dogs,

although its subsequent progression is different among breeds. The severity is thought to

increase with age in Leonbergers [2] and Flat-Coated Retrievers [6] and was shown to be

strongly associated with age in Dandie Dinmont Terriers, Basset Hounds, Flat-Coated

Retrievers, Hungarian Vizslas and Golden Retrievers [7, 8]. However, goniodysgenesis

remained stable in Samoyeds [9] and was not associated with age in a small study of Border

Collies [8].

Sudden onset glaucoma leading to blindness with bilateral loss of eyes began to be seen in

young Border Collies in Australia about 15 years ago. A similar, sudden onset glaucoma in

young animals subsequently emerged in the UK Border Collie population, often in dogs that

were related to the affected Australian dogs. A number of dogs in the USA and Europe have

now also been diagnosed with severe goniodysgenesis and glaucoma. Goniodysgenesis in

Border Collies has been added to the Schedule B list of "Conditions Under Investigation" in

the British Veterinary Association (BVA) Eye Scheme (http://www.bva.co.uk/Canine-Health-

Schemes/Eye-scheme/) and it is highly prevalent in some Border collie lineages (see

Anadune Border collie database: http://www.anadune.com/; free access but registration

required). In a recent study [8], 11 of 102 Border Collies (10.8%) were reported to have

moderate or severe PLD (associated with goniodysgenesis). Seven (6.9%) were mildly

affected. Because of the association with a small number of popular sires and the high

prevalence in some lineages, a genetic aetiology is strongly suspected in Border Collies, but

the heritability and mode of inheritance of the condition is currently unknown. It is also not

known why a proportion of dogs diagnosed with severe goniodysgenesis go on to develop

glaucoma, but some do not, and nor do those with mild goniodysgenesis, over a life span of

15 or more years.

The aim of the current study was to perform a genome-wide analysis to find genetic regions

that are associated with severe goniodysgenesis and glaucoma in the Border Collie and

identify candidate mutations that might be responsible for this condition.

Materials and Methods

Full details of the Materials and Methods are available in **Supplemental File S1**.

Ethical approval

All studies were approved by the Veterinary Ethical Review Committee of the University of

Edinburgh (approval number VERC 2012-8) and the Animal Ethics Committee of the

University of Queensland (approval number ANFRA/MRI-UQ/565/17).

Ascertainment of clinical status and pedigree

Goniodysgenesis was diagnosed by examination of the eyes with gonioscopy to assess the

status of the drainage angle and the pectinate ligament fibres that span it. Since we had

results in multiple formats from many veterinary ophthalmologists, each report was

manually assessed to assign clinical status to each subject. Only individuals who had had

gonioscopy analysis were included. Breeding records and pedigree data were obtained from

public databases and from the breeders' websites. Websites were accessed at various times

up to February 2018. Full details including exclusion criteria are given in **Supplemental File**

S1.

Genetic analysis

DNA samples were from buccal cells collected by owners and occasionally from blood. Full

details of collection and extraction procedures are given in **Supplemental File S1**.

Genotyping used the Illumina 173k CanineHD Whole-Genome Genotyping Bead Chip

(Illumina, San Diego, CA, USA). Results were filtered in PLINK v1.07 [10] to remove

individuals that had more than 10% of missing genotypes and SNPs that had rates of

genotyping < 0.95, had minor allele frequency < 0.05 in this population or deviated from the

Hardy-Weinberg equilibrium in the controls with P value of less than 0.0001. Only the

autosomes were considered.

Cases for the analysis were defined as severely affected and consisted of dogs that had

goniodysgenesis of grades four or five under an early grading scheme, dogs whose

goniodysgenesis was described as "severe", those where the veterinary ophthalmologist

report indicated PLD over more than 70% of the iridocorneal angle and those that

developed glaucoma following a diagnosis of goniodysgenesis. Controls were dogs that were

assessed as clear or clinically unaffected based on at least one gonioscopy examination.

Previous work indicated that goniodysgenesis was not progressive in Border Collies [8] so

the controls were categorised on the basis of a clear gonioscopy, regardless of age. The SNP

analysis was performed in GEMMA v0.94.1 [11] using a linear mixed model approach,

controlling for relatedness using a SNP-based relatedness matrix as a random effect. The

significance threshold was based on a Bonferroni correction of 0.05. Additionally, a more

conservative, regional approach was undertaken using sliding windows of 40 adjacent SNPs,

overlapping by 20 SNPs in REACTA v0.9.7 [12] with the first two genomic principal

components controlling for relatedness. Whole genome sequencing (30X coverage) of DNA

from three dogs with glaucoma, three severely affected and three unaffected animals was

performed using the Illumina HiSeq X platform. Library insert sizes were 450 bp and paired

reads of 150 bp were sequenced. Reads were processed by standard procedures, detailed in

Supplemental File S1 and mapped to the CanFam3.1 reference genome. The significance of

variants was assessed using the Ensembl Variant Effect Predictor [13].

Transcriptomic analysis of candidate region genes

A time course of mouse eye development from embryonic day 12 to adult was examined

using the FANTOM5 database [14-16] (http://fantom.gsc.riken.jp/zenbu). FANTOM5 data

were also used to obtain expression levels for human eye-related samples (**Supplemental**

Table S1). Transcriptomic data for the developing chicken eye were generated for another

project by RNA sequencing methodology (Rainger et al, in preparation). Details are given in

Supplemental File S1. Microarray-based expression data for a range of tissues in human and

mouse were from BioGPS (http://biogps.org) as were RNA sequencing data for chicken

tissues. Microarray expression data for immune cell types was derived from the

Immunological Genome Project (https://www.immgen.org/).

Analysis of canine OLFML3 gene

The region of the canine *OLFML3* gene containing the candidate causative mutation was

amplified by the polymerase chain reaction and the amplicon sequenced by chain

termination sequencing. Full details of primers and protocols are given in **Supplementary**

File S1. Sequences for analysis of nucleotide and amino acid conservation across species

were downloaded from Ensembl (http://ensembl.org) and alignment was performed using

Clustal Omega (http://www.ebi.ac.uk/Tools/msa/clustalo/).

Results

Familial clustering of goniodysgenesis and glaucoma

Inspection of pedigrees revealed that there were three common ancestors in both the sire

and dam lineages for all cases of glaucoma on the goniodysgenesis database (https://bc-

glaucomadatabase.synthasite.com/) (N=12) including two pairs of full siblings both affected

with glaucoma, as well as additional cases not on the database (N=4). These common

ancestors dated back to the late 1970s and early 1980s and were found in multiple ancestral

lineages for the glaucoma cases. All the glaucoma cases had been diagnosed with

goniodysgenesis and subsequently had one or both eyes removed. However, several

individuals diagnosed with severe goniodysgenesis had not gone on to develop glaucoma.

This included two popular sires who reached the age of 15 years without a diagnosis of

glaucoma. It also included a full sibling of two dogs with glaucoma. Three dogs with severe

goniodysgenesis had produced a total of six offspring with glaucoma.

Genome wide association analysis

To identify genetic regions associated with goniodysgenesis in this breed, we performed

SNP analysis on 17 severely affected cases (8 female, 9 male) and 42 unaffected dogs (26

female, 16 male) who had been passed as "clear" at veterinary ophthalmology testing. The

cases all had goniodysgenesis described as "severe" (N=4), Grade 4 or 5 by the earlier

grading scheme (N=1 and 5 respectively), or goniodysgenesis followed by glaucoma (N=7).

A strong association of goniodysgenesis with a region of chromosome 17 was found (**Figure**

1A). The lowest *p*-value was for SNP rs22561716 at 17:51,919,221 (P=2 x 10^{-13}) included in a

window of 40 SNPs that had a *p*-value of 3 x 10^{-07} according to REACTA. SNPs with the lowest

p-values were in a region of about 1 x 10^6 base pairs (**Figure 1B**). Sixteen of the severely

affected and glaucoma cases were homozygous for this haplotype and one case was

heterozygous in this region. Within the genomic region identified by the association analysis

there were a number of coding genes (**Figure 1A**), although none was known to be involved

in eye development or glaucoma in animals or humans. Several had strong expression in the

developing mouse eyeball (**Supplemental Figure F1**).

Whole genome sequencing

We then obtained the whole genome sequence of this region in three glaucoma, three

severely affected and three unaffected individuals, who were members of four nuclear

families (**Supplemental Figure F2**). All were distantly related through multiple lineages in

the extended pedigree. Relatives were chosen to highlight variants that were different

between affected and unaffected family members [17]. Within the region of the lowest *p*-

value SNP, a missense variant was found in the olfactomedin like 3 (*OLFML3*) gene, at

position 17:51,786,924. The mutation was c.590G>A with the predicted change of arginine

at position 197 to glutamine (p.R197Q). This arginine is in the olfactomedin (OLF) domain

[18, 19] (**Figure 2A**) and was found to be highly conserved throughout mammals, birds and

reptiles (**Figure 2B**). The impact predicted by the Ensembl Variant Effect Predictor [13] was

given as "moderate" indicating "a non-disruptive variant that might change protein

effectiveness", consistent with the non-lethal phenotype of homozygous animals. The SIFT

score was 0.55, indicating a mild effect. In humans, the mutation (c.587G>A) corresponds

with rs377336789 (https://www.ncbi.nlm.nih.gov/snp/) and was seen in four out of 121,412

chromosomes with a minor allele frequency of 3.3×10^{-5}. Genotypes are not provided for

the human data, but these were presumably all heterozygous, given the minor allele

frequency. SIFT score was 0.54 and PolyPhen score was 0.392, also consistent with a mild

effect. The mutation was heterozygous in four of 35 Border Collies from the Dog Biomedical

Variant Database Consortium (giving an allele frequency of 0.06 in this breed) but absent in

504 genotypes of other breeds. Notably, it was not present in 51 Leonbergers, three Basset

Hounds, eight Golden Retrievers or one Dandie Dinmont Terrier, all breeds reported to be

associated with goniodysgenesis. In addition, 350 dogs recorded as Border Collies but of

unknown clinical status were genotyped by Animal Genetics (Talahassie, Florida, USA) via

allele-specific PCR run on an ABI 3730 machine. There were 2 homozygotes and 29

heterozygotes among these 350 dogs, giving an allele frequency of 0.05, consistent with the

smaller sample from the Consortium.

OLFML3 genotypes of severely affected and unaffected Border Collies

The results for the mutation c.590G>A in glaucoma, severe and unaffected dogs are shown

in Table 1. The individual who was heterozygous for the region based on SNP chip was also

AG at *OLFML3*. This animal had been diagnosed with severe goniodysgenesis but did not

develop glaucoma over a 15-year lifespan. In addition, one unaffected dog was AA

genotype. This animal was the offspring of a severely affected (grade 5) father and a mother

diagnosed as affected (with no grade). Both parents were also AA. Under some testing

schemes animals could pass with some degree of goniodysgenesis [2, 4] which may be the

case with this dog. In general, the genotype for *OLFML3* segregated with that for

rs22561716, indicating strong linkage disequilibrium between them. However, an

unaffected mother and son were homozygous for G at *OLFML3* but heterozygous at

rs22561716 suggesting that the ancestral haplotype was G at *OLFML3* with T at rs22561716.

In addition, two unaffected individuals were homozygous for several SNPs either side of

OLFML3 but heterozygous for the *OLFML3* mutation and rs22561716 and appeared to carry

a different haplotype.

To validate these results, we genotyped 31 additional Border Collies that had had

gonioscopy investigations: two with glaucoma (both female), four severely affected (all

female) and 25 unaffected (10 male, 15 female). The genotypes of these dogs are shown in

Table 1 and are consistent with the results suggesting that severe

goniodysgenesis/glaucoma was associated with homozygosity for the A allele of *OLFML3*.

OLFML3 genotypes in mildly affected and uncategorised affected Border Collies

Our initial analysis compared severely or glaucoma affected dogs with dogs passed as

unaffected at gonioscopy examination. Because of changes in the reporting scheme in the

UK and differences among countries, there were a number of equivocal diagnoses. These

were dogs where either the level of goniodysgenesis was given as mild (Grade 1 under the

former scheme or stated as mild or marginal on the report; N = 9; five female, four male),

moderate (Grade 2 – 3 under the old scheme, or stated as moderate or between 25 and

70% blocked on the report; N = 9; eight female, one male), or the dogs were diagnosed as

affected with no description given (N = 15; nine female, six male). To examine whether

OLFML3 was involved in these cases we genotyped these 33 dogs. The results are shown in

Table 2. All three genotypes were found among these cases.

Expression of OLFML3 in the eye

In expression data generated by the FANTOM5 project, the *Olfml3* gene was strongly

expressed in mouse eyeball at embryonic and neonatal stages and declined in the adult

sample (**Figure 3A**). FANTOM5 data showed that the gene was expressed in human lens,

corneal epithelium and retina (**Figure 3B**), although the highest expression was in amniotic

and placental epithelial cells and in mesenchymal stem cells (not shown). *OLFML3* was also

expressed in the developing chick eye, with highest expression at embryonic day 6 (**Figure**

3C). In mouse microarray data (mouse probeset 1448475_at; http://biogps.org) expression

of *Olfml3* was highest in osteoblasts undergoing calcification, but it was also found in

eyecup, iris, ciliary bodies and lens (**Supplemental Figure S3A**). There was negligible

expression in immune system cells, except for macrophages, osteoclasts and particularly

microglia, which had expression comparable to lens and eyecup. Strong expression in

microglia was also seen in the Immunological Genome data set (**Supplemental Figure S3B**).

Microarray data also available through BioGPS showed strongest expression of *OLFML3* in

human adipocytes, uterus and retina, the only eye tissue represented (human probeset

218162_at; http://biogps.org) (**Supplemental Figure S3C**). RNA sequencing data compiled

on BioGPS revealed expression in many samples from the chicken, notably the whole

embryo and connective tissues and the cornea of the adult (**Supplemental Figure 3D**). Other

OLF family members were also expressed in the developing mouse eyeball with a similar

pattern to *Olfml3*, peaking in the early neonatal period and declining in the adult eye

(**Supplemental Figure S4**). In contrast, the myocilin gene (encoding a related protein which

is mutated in human open angle glaucoma) had negligible expression in embryo and early

neonate but increased expression at neonatal day 16 and in adult eyeball (**Supplemental**

Figure S4).

Discussion

In a number of dog breeds development of glaucoma has been associated with the presence

of goniodysgenesis. In the Flat-Coated Retriever, the likelihood of glaucoma was positively

and significantly related to the severity of goniodysgenesis [5] and in the Leonberger five

glaucoma-affected individuals all had a high grade of goniodysgenesis in the contralateral

eye [2]. An abnormal drainage angle is considered typical of some breeds, with reports of up

to 81% of individuals showing evidence of goniodysgenesis [9, 20]. However, the highest

prevalence of glaucoma reported in any single breed is only 5.52% [21], indicating that

severe goniodysgenesis does not necessarily proceed to glaucoma.

An increasing incidence of glaucoma in young animals has been reported in the Border

Collie breed over the last 15 years. Consistent with results in other breeds, our survey of

goniodysgenesis in the Border Collie found some individuals who lived entire lifetimes with

apparently severe goniodysgenesis without developing glaucoma. This would suggest that

although severe goniodysgenesis is important in the progression to glaucoma, it is not the

sole cause. The aetiology may involve a genetic predisposition to goniodysgenesis combined

with genetic, environmental or random factors influencing development of glaucoma.

The diagnosis of goniodysgenesis was made by veterinary ophthalmologists (such as

members of the BVA Eye Panel in the UK or the equivalent certifying body in other

countries) who performed gonioscopies to assess the status of the drainage angle and the

pectinate ligament fibres that cross it. Since the diagnoses were made by a number of

different clinicians over a period of years, there may be considerable noise in the

phenotype. An initial eye scheme developed by the BVA in association with the Kennel Club

of the United Kingdom involved assessing whether the dog had no eye abnormality (a pass),

or some abnormality graded from one to five according to the extent of PLD and the width

of the iridocorneal angle, in increasing levels of severity. Since there was a high degree of

inter-observer variability in assigning a grade, the scheme was subsequently amended such

that the dog would either pass or fail. Dogs that passed could have a drainage angle that

was abnormal to some extent ("mild" goniodysgenesis; see information available at

https://www.bva.co.uk/Canine-Health-Schemes/Eye-scheme/). The pass/fail scheme was

expected to have a higher level of inter-observer agreement. This was confirmed in a recent

study of Leonberger dogs [2] where there was a high correlation in proportion affected with

goniodysgenesis between a prospective study where examinations were performed by a

single specialist (2012-2014; 18% affected) and a retrospective study of BVA eye scheme

certificates (2009-2014; 22% affected). In a study of PLD in Welsh Springer Spaniels, there

was a good correlation between two examiners in determining unaffected eyes, although

there was considerable variability between examiners in assigning the degree of PLD [4].

This variability in categorising the extent of goniodysgenesis means that the classification of

phenotype as "affected" or "unaffected" can result in mild PLD or angle narrowing being

recorded as clinically unaffected [2, 4]. For this reason, we focussed our genome-wide

analysis on severely affected dogs including several that had gone on to develop glaucoma.

Using severely affected cases and unaffected controls, we found a strong peak of

association on chromosome 17, in the region of the *OLFML3* gene, where we detected a

missense mutation p.R197Q. Although this gene has not been implicated in goniodysgenesis

or glaucoma in humans, it is strongly expressed in tissues of the anterior segment of the

human and baboon eye including lens, iris, sclera and trabecular meshwork [18, 22] and in

the developing mouse eye [23] . The *Olfml3* gene was expressed throughout mouse eyeball

development (**Figure 3A**) and during chick eye development (**Figure 3C**). In mouse,

inactivation of the *Olfml3* gene by insertion of a *Lac* cassette resulted in viable and fertile

homozygotes [23], consistent with the relatively mild selective disadvantage of dogs with

goniodysgenesis. Expression has also been observed in human tissues of mesenchymal

origin such as bone and adipose. Olfactomedin like 3 is an extracellular matrix protein that

has been implicated in the epithelial to mesenchyme transition in cancer [24], in keeping

with its mesenchymal expression and suggesting that abnormality of this protein could be

associated with the formation or retention of abnormal sheets of mesenchyme in the

drainage angle. Olfactomedin like 3 is also proangiogenic [25] and interacts with a member

of the bone morphogenic protein (BMP) family, BMP4. Since BMPs including BMP4 are

involved in eye development (see [26] for a review), this may indicate the pathway for

olfactomedin like 3 action.

In addition, expression of *OLFML3* is also characteristic of microglia in humans [27, 28] and

mice (http://biogps.org; http://www.immgen.org), although it is not strongly expressed by

other macrophage lineages. Genetic manipulation resulting in absence of microglia is

associated with loss of *Olfml3* expression in mice (R Rojo et al, in revision). The retina

contains microglia which are important for retinal health [29, 30], and they could be one

source of *OLFML3* mRNA in the canine eye. Opening of the iridocorneal angle involves

considerable remodelling of the mesenchymal tissue within the angle, including thinning

and extending of the pectinate ligament and opening the trabecular meshwork [31].

Microglia may contribute to this process as eye development progresses by phagocytosing

apoptotic cells and debris from the remodelling. Abnormality of olfactomedin like 3 in the

microglia of the eye could interfere with this function and therefore disrupt the opening of

the iridocorneal angle and formation of the proper drainage channels.

Olfactomedin like 3 is a member of a large family characterised by the olfactomedin (OLF)

domain [19, 32], with several members expressed in eye structures. In particular, the

glucocorticoid-inducible family member myocilin (*MYOC* gene) has been implicated in

dominant open angle glaucoma in humans [33] and is strongly expressed in the trabecular

meshwork of the drainage angle [34, 35]. As with *Olfml3* mutants, there is no gross

phenotype in *Myoc* knockout mice [36]. Increased expression of *Myoc* mRNA resulted in a

reduction in *Olfml3* mRNA [37] in a mouse model, suggesting that there is an inverse

association between these two OLF proteins, consistent with the expression profiles in the

developing mouse eye (**Supplemental Figure S4**) where *Olfml3* declined in the adult while

Myoc rose. *In vitro* experiments indicated that myocilin interacts with olfactomedin 3 (also

known as optimedin, encoded by the *OLFM3* gene) which was co-expressed with myocilin in

the eye [38]. Mutation in another OLF family member, *OLFM2,* has been associated with

human open angle glaucoma in a small number of Japanese patients (rs779032127;

p.R144Q) [39] and identified as contributing to eye development [40]. These other OLF

family members may compensate for olfactomedin like 3 abnormalities, or may interact

with olfactomedin like 3 in the development of the eye. Hence genetic variation in these

family members may modify the impact of the *OLFML3* mutation in the Border Collies,

explaining the variable phenotype of homozygotes and heterozygotes carrying the

mutation.

We propose that the p.R197Q mutation in Border Collies is responsible for the presence of

severe goniodysgenesis predisposing to glaucoma in homozygous animals. We have also

seen some heterozygous animals with mild to relatively severe phenotypes (although none

with glaucoma), but most heterozygotes have been passed as clinically unaffected at

gonioscopy testing. This suggests that there may be variants at modifier loci that can

374 compensate for the *OLFML3* mutation. In particular, the presence of many OLF proteins in

375 the eye and the mild phenotype associated with relatively broadly expressed gene family

376 members suggests redundancy for at least some functions. It is also possible that

377 inadequate compensation for the putative effects of the p.R197Q mutation may be

378 responsible for the progression from severe goniodysgenesis to glaucoma.

379 There were too few glaucoma cases in our study to allow identification of loci involved in

380 progression from goniodysgenesis to glaucoma in the Border Collie and ambiguous

381 phenotyping complicates the investigation of mild goniodysgenesis. The genetic relationship

382 between the mild forms and the more severe forms is unknown, but the *OLFML3* variant

383 described here does not appear to be associated with most cases of mild goniodysgenesis

384 indicating that genetic predisposition to mild goniodysgenesis may be independent of the

385 findings of this study. Although developing a test for the *OLFML3* mutation would allow

386 breeders to select against homozygotes to decrease the prevalence of the risk genotype for

387 severe goniodysgenesis, it is important not to reduce the breeding pool too much because

388 of the risk of other recessive conditions resulting from homozygosity for alleles that are

389 identical by descent. Therefore, we need to understand why some of our cases progressed

390 to glaucoma at a young age and some did not before we can make recommendations to the

391 breeders of Border Collies worldwide.

393 **Author contributions**

394 KMS, CAP and LLF conceived the study. AJC, LLF and KMS curated and prepared the DNA

395 samples. OM, SCB and LLF performed initial genetic analysis. LLF and CAP performed the

396 SNP analysis. CAP and SJB performed the genome wide sequencing analysis. KMS and AJC

designed, performed and analysed the *OLFML3* genotyping. JJS analysed the data from the

Dog Genome Consortium. AdeK and CW analysed the Animal Genetics samples. KMS

analysed the public gene expression data. JR and VT-R created and analysed the chicken

eye data. KMS, LLF and CAP drafted the text and all authors read and edited the final

manuscript.

Acknowledgements

We are grateful to Dogs Trust (UK) for a Bateson Canine Welfare Grant for this project and

to the Pastoral Breeds Health Foundation (UK) for pump priming funding. The Roslin

Institute is supported by core funding from the Biotechnology and Biological Sciences

Research Council (UK) [Grant numbers BB/J004235/1, BB/J004316/1, BB/P013732/1]. JR is

funded by Fight for Sight (UK) [Grant number1590/1591]. VT-R was funded by a CONACYT

(Mexico) international studentship. KMS is supported by the Mater Foundation (Brisbane,

Australia). We would like to thank all the Border Collie owners and breeders who have

contributed samples and information to this investigation and the Pastoral Breeds Health

Foundation for continuing support and encouragement during the course of this project. We

are grateful to Dr Alan Wilton (now deceased) and Dr Barbara Zangerl for initiating and

supporting the project.

Conflict of Interest

AdeK and CW are employed by Animal Genetics, Talahassie, FL, USA, which is now offering a

test for genetic predisposition to severe goniodysgenesis based on these results.

Figure legends

Figure 1. Genome wide association with severe goniodysgenesis and glaucoma

A. Manhattan plot for the canine genome; chromosome 17 is shown in more detail and

genes in the region are shown below.

B. Homozygosity within the region of interest. Each row is a DNA sample and each column

shows a SNP. Red and yellow indicate homozygosity for the major and minor allele

respectively and green shows heterozygosity. The upper block shows severely affected and

glaucoma animals; the lower block shows those passed as clinically unaffected at

gonioscopy testing. The most significant SNP is shown as yellow in the cases (homozygous

for the minor allele); one clinically unaffected animal was homozygous for this allele and

one severely affected animal was heterozygous.

Figure 2. Conservation of *OLFML3* predicted amino acid sequence

A. Amino acid sequence of the canine OLFML3 protein. Functional domains are highlighted:

blue – signal peptide; pink – coiled coil domain; yellow – olfactomedin like domain [18, 19].

Within the yellow region, bold lettering shows the core OLF region. Green shows the

arginine at position 197.

B. Conservation of arginine 197 (green) in OLFML3 of vertebrates. Arginine is found at the

equivalent position in all mammal, bird and reptile species for which there is an annotated

OLFML3 gene, and in coelacanth but not in other fish. The region around this arginine is also

highly conserved.

Figure 3. Expression of *OLFML3* genes in eye regions

A. Expression of *Olfml3* in developing mouse eyeball. Results are based on CAGE and were downloaded from the FANTOM5 database (http://fantom.gsc.riken.jp/5/tet). Y axix shows RLE normalised values. Light grey – embryonic samples; dark grey – neonatal samples; black – adult sample.

B. Expression of *OLFML3* in human eye tissues and cells. Results are based on CAGE and were downloaded from the FANTOM5 database. Y axis shows RLE normalised values. Light grey – optic nerve and retina; mid grey – cornea; dark grey – lens.

C. Expression of *OLFML3* in the developing chick eye. Results are based on RNA sequencing. Y axis shows transcripts per million (TPM) values. Light grey – embryonic day 5; mid grey – embryonic day 6; dark grey – embryonic day 7.

Supplemental Material

Supplemental Table S1. FANTOM5 samples used in analysis of gene expression. Details can be found at fantom.gsc.riken.jp/5/sstar/.

Supplemental Figure S1. Expression of genes in the candidate region in the developing mouse eye. Upper panel shows the canine genes in the region. Lower panel shows expression of the genes in the developing mouse eyeball. Y axis shows RLE normalised expression levels. Data taken from FANTOM5 (see **Supplemental Table S1**). Light grey – embryonic; dark grey – neonatal; black – adult.

Supplemental Figure S2. Pedigrees of animals used for whole genome sequencing. Grey fill

indicates severe goniodysgenesis; black fill indicates glaucoma; white fill indicates

unaffected or unknown status. Double line indicates a first cousin mating; all glaucoma

individuals can be traced back to the same three distant common ancestors on both sides.

Animals with a BC ID were sequenced.

Supplemental Figure S3. Expression of olfactomedin like 3 gene in cells and tissues

A. Expression in mouse tissues and cell lines (http://biogps.org; microarray probeset

1448475_at). Y axis shows relative intensity of expression (arbitrary fluorescence

units).

B. Expression in mouse myeloid cells (http://www.immgen.org, dataset "MFs,

Monocytes, Neutrophils"; microarray probeset 10500808). Y axis shows relative

intensity of expression (arbitrary fluorescence units).

C. Expression in human tissues and cell lines (http://biogps.org; microarray probeset

218162_at). Y axis shows relative intensity of expression (arbitrary fluorescence

units).

D. Expression in chicken tissues (RNA sequencing data curated from public databases

and analysed using Kallisto, available at

http://biogps.org/dataset/BDS_00031/chicken-atlas/; Bush et al, in revision). Y axis

shows tags per million (TPM).

In A, C and D, blue – nervous tissues; dark green ☐ connective tissues; light green ☐

digestive tract and renal; yellow – reproductive; orange – immune; red – ocular.

Supplemental Figure S4. Expression of olfactomedin family genes in the developing mouse

eye (data from FANTOM5; http://fantom.gsc.riken.jp/zenbu/). Y axis shows RLE normalised

expression levels based on CAGE analysis. Light grey – embryonic; dark grey – neonatal;

black – adult.

1.   Asher L, Diesel G, Summers JF, McGreevy PD, Collins LM: **Inherited defects in pedigree dogs.**
489   **Part 1: disorders related to breed standards**. *Vet J* 2009, **182**(3):402-411.

490   2.   Fricker GV, Smith K, Gould DJ: **Survey of the incidence of pectinate ligament dysplasia and**
491   **glaucoma in the UK Leonberger population**. *Vet Ophthalmol* 2016, **19**(5):379-385.

492   3.   Plummer CE, Regnier, A., Gellatt, K.N.: **The canine glaucomas**. In: *Veterinary Ophthalmology*.
493   Edited by Gelatt KN, 5th edn. Iowa: John Wiley & Sons Inc; 2013: 1050-1145.

494   4.   Oliver JAC, Cottrell BC, Newton JR, Mellersh CS: **Gonioscopy in the dog: inter-examiner**
495   **variability and the search for a grading scheme**. *J Small Anim Pract* 2017, **58**(11):652-658.

496   5.   Wood JL, Lakhani KH, Read RA: **Pectinate ligament dysplasia and glaucoma in Flat Coated**
497   **Retrievers. II. Assessment of prevalence and heritability**. *Vet Ophthalmol* 1998, **1**(2-3):91-
498   99.

499   6.   Pearl R, Gould D, Spiess B: **Progression of pectinate ligament dysplasia over time in two**
500   **populations of Flat-Coated Retrievers**. *Vet Ophthalmol* 2015, **18**(1):6-12.

501   7.   Oliver JA, Ekiri A, Mellersh CS: **Prevalence of pectinate ligament dysplasia and associations**
502   **with age, sex and intraocular pressure in the Basset hound, Flatcoated retriever and**
503   **Dandie Dinmont terrier**. *Canine Genet Epidemiol* 2016, **3**:1.

504   8.   Oliver JA, Ekiri AB, Mellersh CS: **Pectinate ligament dysplasia in the Border Collie,**
505   **Hungarian Vizsla and Golden Retriever**. *Vet Rec* 2017, **180**(11):279.

506   9.   Ekesten B, Narfstrom K: **Correlation of morphologic features of the iridocorneal angle to**
507   **intraocular pressure in Samoyeds**. *Am J Vet Res* 1991, **52**(11):1875-1878.

508   10.   Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MA, Bender D, Maller J, Sklar P, de
509   Bakker PI, Daly MJ *et al*: **PLINK: a tool set for whole-genome association and population-**
510   **based linkage analyses**. *Am J Hum Genet* 2007, **81**(3):559-575.

511   11.   Zhou X, Stephens M: **Genome-wide efficient mixed-model analysis for association studies**.
512   *Nat Genet* 2012, **44**(7):821-824.

513   12.   Cebamanos L, Gray A, Stewart I, Tenesa A: **Regional heritability advanced complex trait**
514   **analysis for GPU and traditional parallel architectures**. *Bioinformatics* 2014, **30**(8):1177-
515   1179.

516   13.   McLaren W, Gil L, Hunt SE, Riat HS, Ritchie GR, Thormann A, Flicek P, Cunningham F: **The**
517   **Ensembl Variant Effect Predictor**. *Genome Biol* 2016, **17**(1):122.

518   14.   Forrest AR, Kawaji H, Rehli M, Baillie JK, de Hoon MJ, Haberle V, Lassmann T, Kulakovskiy IV,
519   Lizio M, Itoh M *et al*: **A promoter-level mammalian expression atlas**. *Nature* 2014,
520   **507**(7493):462-470.

521   15.   Arner E, Daub CO, Vitting-Seerup K, Andersson R, Lilje B, Drablos F, Lennartsson A,
522   Ronnerblad M, Hrydziuszko O, Vitezic M *et al*: **Gene regulation. Transcribed enhancers lead**
523   **waves of coordinated transcription in transitioning mammalian cells**. *Science* 2015,
524   **347**(6225):1010-1014.

525   16.   Summers KM, Hume DA: **Identification of the macrophage-specific promoter signature in**
526   **FANTOM5 mouse embryo developmental time course data**. *J Leukoc Biol* 2017,
527   **102**(4):1081-1092.

528   17.   Lander ES, Botstein D: **Homozygosity mapping: a way to map human recessive traits with**
529   **the DNA of inbred children**. *Science* 1987, **236**(4808):1567-1570.

530   18.   Rodriguez-Sanchez IP, Garza-Rodriguez ML, Mohamed-Noriega K, Voruganti VS, Tejero ME,
531   Delgado-Enciso I, Perez-Ibave DC, Schlabritz-Loutsevitch NE, Mohamed-Noriega J, Martinez-
532   Fierro ML *et al*: **Olfactomedin like 3 (OLFML3) gene expression in baboon and human**
533   **ocular tissues: cornea, lens, uvea, and retina**. *J Med Primatol* 2013, **42**(3):105-111.

534   19.   Zeng LC, Han ZG, Ma WJ: **Elucidation of subfamily segregation and intramolecular**
535   **coevolution of the olfactomedin like proteins by comprehensive phylogenetic analysis and**
536   **gene expression pattern assessment**. *FEBS Lett* 2005, **579**(25):5443-5453.

537   20.   Kato K, Sasaki N, Matsunaga S, Mochizuki M, Nishimura R, Ogawa H: **Possible association of**
538   **glaucoma with pectinate ligament dysplasia and narrowing of the iridocorneal angle in**
539   **Shiba Inu dogs in Japan**. *Vet Ophthalmol* 2006, **9**(2):71-75.
540   21.   Gelatt KN, MacKay EO: **Prevalence of the breed-related glaucomas in pure-bred dogs in**
541   **North America**. *Vet Ophthalmol* 2004, **7**(2):97-111.
542   22.   Tomarev SI, Nakaya N: **Olfactomedin domain-containing proteins: possible mechanisms of**
543   **action and functions in normal development and pathology**. *Mol Neurobiol* 2009,
544   **40**(2):122-138.
545   23.   Ikeya M, Kawada M, Nakazawa Y, Sakuragi M, Sasai N, Ueno M, Kiyonari H, Nakao K, Sasai Y:
546   **Gene disruption/knock-in analysis of mONT3: vector construction by employing both in**
547   **vivo and in vitro recombinations**. *Int J Dev Biol* 2005, **49**(7):807-823.
548   24.   Schmitz S, Bindea G, Albu RI, Mlecnik B, Machiels JP: **Cetuximab promotes epithelial to**
549   **mesenchymal transition and cancer associated fibroblasts in patients with head and neck**
550   **cancer**. *Oncotarget* 2015, **6**(33):34288-34299.
551   25.   Miljkovic-Licina M, Hammel P, Garrido-Urbani S, Lee BP, Meguenani M, Chaabane C,
552   Bochaton-Piallat ML, Imhof BA: **Targeting olfactomedin like 3 inhibits tumor growth by**
553   **impairing angiogenesis and pericyte coverage**. *Mol Cancer Ther* 2012, **11**(12):2588-2599.
554   26.   Pillai-Kastoori L, Wen W, Morris AC: **Keeping an eye on SOXC proteins**. *Dev Dyn* 2015,
555   **244**(3):367-376.
556   27.   Butovsky O, Jedrychowski MP, Moore CS, Cialic R, Lanser AJ, Gabriely G, Koeglsperger T,
557   Dake B, Wu PM, Doykan CE *et al*: **Identification of a unique TGF-beta-dependent molecular**
558   **and functional signature in microglia**. *Nat Neurosci* 2014, **17**(1):131-143.
559   28.   Gosselin D, Skola D, Coufal NG, Holtman IR, Schlachetzki JCM, Sajti E, Jaeger BN, O'Connor C,
560   Fitzpatrick C, Pasillas MP *et al*: **An environment-dependent transcriptional network**
561   **specifies human microglia identity**. *Science* 2017, **356**(6344).
562   29.   Hume DA, Perry VH, Gordon S: **Immunohistochemical localization of a macrophage-specific**
563   **antigen in developing mouse retina: phagocytosis of dying neurons and differentiation of**
564   **microglial cells to form a regular array in the plexiform layers**. *J Cell Biol* 1983, **97**(1):253-
565   257.
566   30.   Karlstetter M, Ebert S, Langmann T: **Microglia in the healthy and degenerating retina:**
567   **insights from novel mouse models**. *Immunobiology* 2010, **215**(9-10):685-691.
568   31.   Trejo-Reveles V, McTeir L, Summers K, Rainger J: **An analysis of anterior segment**
569   **development in the chicken eye**. *Mech Dev* 2018, **150**:42-29.
570   32.   Anholt RR: **Olfactomedin proteins: central players in development and disease**. *Frontiers in*
571   *cell and developmental biology* 2014, **2**:6.
572   33.   Stone EM, Fingert JH, Alward WL, Nguyen TD, Polansky JR, Sunden SL, Nishimura D, Clark AF,
573   Nystuen A, Nichols BE *et al*: **Identification of a gene that causes primary open angle**
574   **glaucoma**. *Science* 1997, **275**(5300):668-670.
575   34.   Tamm ER, Russell P, Epstein DL, Johnson DH, Piatigorsky J: **Modulation of myocilin/TIGR**
576   **expression in human trabecular meshwork**. *Invest Ophthalmol Vis Sci* 1999, **40**(11):2577-
577   2582.
578   35.   Tomarev SI, Wistow G, Raymond V, Dubois S, Malyukova I: **Gene expression profile of the**
579   **human trabecular meshwork: NEIBank sequence tag analysis**. *Invest Ophthalmol Vis Sci*
580   2003, **44**(6):2588-2596.
581   36.   Kim BS, Savinova OV, Reedy MV, Martin J, Lun Y, Gan L, Smith RS, Tomarev SI, John SW,
582   Johnson RL: **Targeted Disruption of the Myocilin Gene (Myoc) Suggests that Human**
583   **Glaucoma-Causing Mutations Are Gain of Function**. *Mol Cell Biol* 2001, **21**(22):7707-7713.
584   37.   Paper W, Kroeber M, Heersink S, Stephan DA, Fuchshofer R, Russell P, Tamm ER: **Elevated**
585   **amounts of myocilin in the aqueous humor of transgenic mice cause significant changes in**
586   **ocular gene expression**. *Exp Eye Res* 2008, **87**(3):257-267.

38.   Torrado M, Trivedi R, Zinovieva R, Karavanova I, Tomarev SI: **Optimedin: a novel**
olfactomedin-related protein that interacts with myocilin. *Hum Mol Genet* 2002,
11(11):1291-1301.
39.   Funayama T, Mashima Y, Ohtake Y, Ishikawa K, Fuse N, Yasuda N, Fukuchi T, Murakami A,
Hotta Y, Shimada N *et al*: **SNPs and interaction analyses of noelin 2, myocilin, and**
optineurin genes in Japanese patients with open-angle glaucoma. *Invest Ophthalmol Vis Sci*
2006, **47**(12):5368-5375.
40.   Holt R, Ugur Iseri SA, Wyatt AW, Bax DA, Gold Diaz D, Santos C, Broadgate S, Dunn R, Bruty J,
Wallis Y *et al*: **Identification and functional characterisation of genetic variants in OLFM2 in**
children with developmental eye disorders. *Hum Genet* 2017, **136**(1):119-127.

Genotype Clinical status	AA	AG	GG	TOTAL
SNP-chip dogs[1]				
Glaucoma	7	0	0	7
Severe goniodysgenesis	9	1	0	10
Unaffected	1	19	22	42
Other dogs				
Glaucoma	2	0	0	2
Severe goniodysgenesis	3	1	0	4
Unaffected	0	17	8	25
TOTALS				
Glaucoma	9	0	0	9
Severe goniodysgenesis	12	2	0	14
Unaffected	1	36	30	67

601   **Table 1.** Genotype frequencies for *OLFML3* mutation c.590G>A in severely affected and unaffected

602   dogs.

603   Dogs that were included in the genome wide analysis (SNP-chip dogs) are shown separately from

604   dogs that were only tested for *OLFML3.*

Genotype Clinical status	AA	AG	GG	TOTAL
Mild	3	3	3	9
Moderate	4	2	3	9
Affected, no grade	7	3	5	15
TOTALS	14	8	11	33

608   **Table 2.** Genotype frequencies for *OLFML3* mutation c.590G>A in mild, moderate and

609   uncharacterised affected dogs.

A

MGPYTPLLTMFLLSWWLRPLQGQQHHLVEYMERRLAALEERLAQCQDQSSRHAAELRDFR

NKMLPLLEVAEKEREALRTEADTISGRVDRLEREVDYLETQNPALPCVEVDEKVTGGPGI

KGKGRRNEKYDMLTDCGYTISQVRSMKILKRFGGPAGLWTKDPMGPTEKIYVLDGTQNDT

AFVFPRLRDFTLAMAARKASRVRVPFPWVGTGQLVYGGFLYYARRPPGGPGGGGGLENTL

QLIKFHLANRTVVDSSVFPAEGLIPPYGLTADTYIDLAADEEGLWAVYATREDDRHLCLA

KLDPQTLDTEQQWDTPCPRENAEAAFVICGTLYVVYNTRPASRARIQCSFDASGTLTPER

AALPYFPRRYGAHASLRYNPRERQLYAWDDGYQIVYKLEMRRKEEEV

B

```
Opossum_OLFML3        VRSMKILKRFGSPAGMWTKDPLGPTEKIFVLDGTQNDTAYVFPRLRDFTLSTTARKASRV
Wallaby_OLFML3        VRSMKILKRFGSSAGMWTKDPLGPSEKIFVFDGTQNDTAYVFPRLRDFTLSSAARKASRV
Tasmanian_Devil_OLFML3 VRSMKILKRFGSSAGMWTKDPLGHSEKIFVLDGTQNDTAYVFPRLRDFTLSTAARKASRV
Shrew_OLFML3          VRSMKILKRFGGPTGLWTKDPLGPTEKIFVLDGTHNDTAFIFPRLRDFTLAMAARKAARV
Kangaroo_rat_OLFML3   VRSMKILKRFGGPAGMWTKDPLGPAEKIYFDGTQNDTALVFPRLRDFTLAMAARKASRI
Hedgehog_OLFML3       VRSMKILKRFGSPPGLWTKDPLGTTEKIYVLD-PQNDTAFIFPRLRDFTLAMAARKASRV
Armadillo_OLFML3      VRSMKILKRFGGPAGLWTKDPLGPAERIYVLAGTQNDTALVFPKLRDFTLATAARKASRV
Mouse_Olfml33         VRSMKILKRFGGSVGLWTKDPLGPAEKIYVLDGTQNDTAFVFPRLRDFTLAMAARKASRI
Rat_Olfml3            VRSMKILKRFGGSAGLWTKDPLGPAEKIYVLDGTQNDTAFVFPRLRDFTLTMAARKASRI
Elephant_OLFML3       VRSMKILKRFGGPAGLWTKDPLGPTEKIYVLDGTQNDTAFVFPRLRDFSLSTATRKASRV
Guinea_pig_OLFML3     VRSMKILKRFGGPAGLWTKDPLGPTEKIYVLDGTQNDTAFVFPRLRDFTLAMAARKASRV
Hyrax_OLFML3          VRSMKILKRFGGPAGLWTKDPLGPAEKIYVLDGTQNDTAFVFPRLRDFSLSMATRKASRV
Microbat_OLFML3       VRSMKILKRFGGSAGLWTKDPLGPTEKIYVLDGTQNDTAFVFPRLRDFTLAMAARKASRV
Squirrel_OLFML3       VRSMKILKRFGGPAGLWTKDPLGPTEKIYVLDGTQNDTAFVFPRLRDFTLTMAARKASRV
Pig_OLFML3            VRSMKILKRFGGPAGLWTKDPLGPTEKIFVLDGTQNDTAFVFPRLRDFTLAMAARKASRI
Horse_OLFML3          VRSMKILKRFGGPAGLWTKDPMGPTEKIYVLDGTQNDTAFVFPRLRDFTLAMAARKASRV
Dog_OLFML3            VRSMKILKRFGGPAGLWTKDPMGPTEKIYVLDGTQNDTAFVFPRLRDFTLAMAARKASRV
Ferret_OLFML3         VRSMKILKRFGGPAGLWTKDPMGPTEKIYVLDGTQNDTAFVFPRLRDFTLAMAARKASRV
Panda_OLFML3          VRSMKILKRFGGPAGLWTKDPMGPTEKIYVLDGTQNDTAFIFPRLRDFTLAMAARKASRV
Cat_OLFML3            VRSMKILKRFGGPAGLWTKDPMGPAERIYVLDGTQNDTAFVFPRLRDFTLAMAARKASRV
Mouse_lemur_OLFML3    VRSMKILKRFGGPAGLWTKDPLGPTEKIYVLDGTQNDTAFVFPRLRDFTLAMAARKASRV
Human_OLFML3          VRSMKILKRFGGPAGLWTKDPLGQTEKIYVLDGTQNDTAFVFPRLRDFTLAMAARKASRV
Cow_OLFML3            VRSMKILKRFGGPAGLWTKDPLGPAEKIYVLDGTQNDTAFVFPRLRDFTLAMAARKASRV
Sheep_OLFML3          VRSMKILKRFGGPAGLWTKDPLGPAEKIYVLDGTQNDTAFVFPRLRDFTLTMAARKASRV
Dolphin_OLFML3        VRSMKILKRFGGPAGLWTKDPLGPAEKIYVLDGTQNDTAFVFPRLRDFTLAMAARKASRV
Coelacanth_OLFML3     IKSMKILKRIGSSTGMWARDPLGDSDNIYVLEGTGNDTVYEYSRARDFTQSPVMRARRI
Xenopus_OLFML3        VTAMKILKRFGSSAGLWTKDLAGNSDRIYVFDGAGNDTVYVPRMKEFTLSSSTRKAAKI
Chicken_OLFML3        VRAMKILKRFGSSAGLWTKDAAGSSEKIYVFDGTANDTVYIFPRMREFTLFSATRRAARI
Flycatcher_OLFML3     VRAMKILKRFGSASGLWTKDAAGSSEKIYVFDGTANDTVYVFPRMREFTLFSATRRAARI
Lizard_OLFML3         VKAMKILKRFGSTAGLWTKDPVGNSEKIYVFDGTSNDTVYVFPRMREFTLFSATRRAARI
Turtle_OLFML3         VKAMKILKKFGSSAGLWTKDPVGSSEKIYVFDGTSNDTVYVFPRMREFTLFSATRRAARI
                       : :*****::*.  *:*::*  *  ::.*:*:    ***. : : ::*:    .*:* ::
```