Highly diverged novel subunit composition of apicomplexan F-type ATP synthase identified from *Toxoplasma gondii*

Rahul Salunke[1], Tobias Mourier[2], Manidipa Banerjee[3], Arnab Pain[2] and Dhanasekaran Shanmugam[#1]

[1]Biochemical Sciences Division, CSIR-National Chemical Laboratory, Pune, Maharashtra, India.

[2]Pathogen Genomics Laboratory, BESE Division, King Abdullah University of Science and Technology (KAUST), Thuwal, Saudi Arabia.

[3]Kusuma School of Biological Sciences, Indian Institute of Technology-Delhi, Hauz Khas, New Delhi, India.

[#]Correspondence to be addressed to d.shanmugam@ncl.res.in

Short title: Novel subunits of *T. gondii* F-type ATP synthase

Key words: *Toxoplasma gondii*, Apicomplexa, Mitochondrion, F-type ATP synthase, Oxidative Phosphorylation

Abstract

The mitochondrial F-type ATP synthase, a multi-subunit nanomotor, is critical for maintaining cellular ATP levels. In *Toxoplasma gondii* and other apicomplexan parasites, many subunit components, necessary for proper assembly and functioning of this enzyme, appear to be missing. Here, we report the identification of 20 novel subunits of *T. gondii* F-type ATP synthase from mass spectrometry analysis of partially purified monomer (~600 kDa) and dimer (>1 MDa) forms of the enzyme. Despite extreme sequence diversification, key F_O subunits, a, b and d, can be identified from conserved structural features. Orthologs for these proteins are restricted to apicomplexan, chromerid and dinoflagellate species. Interestingly, their absence in ciliates indicates a major diversion, with respect to subunit composition of this enzyme, within the alveolate clade. Discovery of these highly diversified novel components of the apicomplexan F-type ATP synthase complex will facilitate the development of novel anti-parasitic agents. Structural and functional characterization of this unusual enzyme complex will advance our fundamental understanding of energy metabolism in apicomplexan species.

Introduction

The F-type ATP synthase is a ubiquitous nanomotor present on the inner membrane
of mitochondria, chloroplast and bacterial plasma membrane, and catalyzes ATP formation
from ADP and Pi [1,2]. In mitochondria, the energy required to drive the nanomotor is
harnessed from the proton motive force and associated membrane potential, which are
generated by the mitochondrial respiratory complexes [3,4]. The mitochondrial F-type ATP
synthase complex consists of two sub-complexes called the F_1 and F_O sectors. The
hydrophilic globular F_1 sector includes three each of α and β subunits, and a central stalk
containing one each of γ, δ and ε subunits. The site of ATP formation resides in the catalytic
center located in the β subunit [1,2,4,5]. Unlike the F_1 sector subunit components, which are
well conserved across various eukaryotic phyla, many F_O sector subunit components are
highly diverse [6-9] and are not readily identified based on sequence similarity. The complete
set of F_O sector subunits have been mapped only in a few species [10,11]. For example, in the
yeast *Saccharomyces cerevisiae*, the F_O sector is composed of the hydrophobic membrane
spanning oligomeric subunit c (10-12 monomers), along with subunits a, b, d, f, 8, h and
OSCP (Oligomycin sensitivity confirming protein) [10]. In addition, other accessory subunits
such as e, g, i, j, and k are also associated with the enzyme. Orthologs of all yeast F_O sector
subunits, except subunits j and k, are present in the mammalian counterpart (bovine enzyme),
which additionally has subunits AGP and MLQ [11].

Numerous biochemical and structural studies have addressed the assembly and
interaction between the various subunit components of F-type ATP synthases [1,12]. These
studies have provided in-depth insights on the functional coupling of the proton motive force
to ATP synthesis. The F_O sector subunits c and a interact to form the proton channel, and
proton translocation, which occurs along the interface of these two subunits, is coupled to the
rotary motion of the c-ring and central stalk of the enzyme [13-16]. The peripheral stalk

structure of the enzyme, comprised of F_O sector subunits b, d, h (f6 in bovine) and OSCP, has

the critical role of holding the $\alpha_3\beta_3$ catalytic domain in place during the clockwise rotation of

the central stalk. These events elicit conformational changes in the $\alpha_3\beta_3$ catalytic domain,

which then results in ATP synthesis [2,4,17,18]. F_O sector subunits e and g are known to be

involved in dimer assembly of the enzyme complex in yeast [19,20], which appears to

facilitate cristae formation by the mitochondrial inner membrane [21-24].

However, despite the importance of the various F_O sector subunits, only subunit c and

OSCP can be detected from sequence conservation in the vast majority of unicellular

eukaryotes, including in the alveolate infrakingdom, which includes the phylum *Apicomplexa*

[25-27]. It is unlikely that the missing subunits are totally absent in these species. Plausible

scenarios are that they have either diverged in sequence beyond recognition or that a

completely new set of proteins have substituted for the missing subunits in these species. This

is indeed the case in organisms such as *Euglena gracilis* [9], *Trypanosoma brucei* [7],

Tetrahymena thermophile [8] and *Chlorophyceae* algae [6]. Hence, our interest was to

characterize the complete subunit composition of the F-type ATP synthase from the

apicomplexan parasite *Toxoplasma gondii*.

T. gondii is an obligate intracellular protozoan parasite, responsible for toxoplasmosis

in humans and animals [28]. The parasite completes its life cycle into two host species; the

definitive hosts are naïve feline species which support the sexual development of the parasite,

and all warm blooded animals are intermediate hosts and support its asexual development

[29]. During asexual development, the parasites can reversibly differentiate between fast-

growing virulent tachyzoites and slow growing latent bradyzoites [30]. By virtue of their fast-

growing nature, tachyzoites are metabolically more active and glycolysis is the primary

source for carbon and energy for optimal growth of this parasite [31-33]. However, recent

findings have revealed that glucose is not an essential nutrient [34], and parasites survive in

the presence of glutamine as an alternative nutrient source [35,36]. In the absence of glucose

oxidation *via* glycolysis, oxidative phosphorylation is the only source for bulk cellular ATP.

Evidence for an active oxidative phosphorylation in *T. gondii* exists [37-39], and treatment

with atovaquone, a potent inhibitor of mitochondrial electron transport chain (mtETC),

results in inhibition of ATP synthesis and parasite growth ($EC_{50} < 10$ nM) [38].

Very little is known about the structure and function of the F-type ATP synthase from

T. gondii and other apicomplexan parasites. Comparative genomics has revealed that, while

the five canonical F_1 sector subunits - α, β, γ, δ, and ε - are readily identified in all

apicomplexan parasites based on amino acid sequence conservation, only two F_O sector

subunits - c and OSCP - can be identified by sequence [25-27]. Characterization of F-type

ATP synthase from solubilized mitochondria lysates of the human malaria parasite

Plasmodium falciparum revealed that the enzyme assembles into monomer and dimer forms

[26]. This suggests that a full complement of subunits, typical for the eukaryotic enzyme, is

present in *P. falciparum* and likely in other apicomplexan parasites as well.

Here, we have investigated the subunit composition of the enzyme from *T. gondii*. We

were able to identify and partially purify the F-type ATP synthase enzyme complex from

solubilized mitochondrial preparation of *T. gondii* using blue native PAGE separation, by

immuno-precipitation and by chromatographic enrichment. LC-MS analysis of the enzyme

preparations revealed the identity of the proteins associated with the *T. gondii* F-type ATP

synthase. We have identified 20 novel proteins (of unknown function) as being *bona fide*

subunit constituents of *T. gondii* F-type ATP synthase based on consensus from multiple and

independent experiments. Phyletic profiling revealed that orthologs for many of these

proteins are present in most apicomplexan species, except in those *Cryptosporidium sp* which

posses a degenerate mitochondrion devoid of all proteins involved in oxidative

phosphorylation [40]. Further, the phyletic profiles of these novel proteins reveal their origin

to be ancient, probably even ancestral to alveolate organisms. The identification of these

novel protein components of the apicomplexan F-type ATP synthase will facilitate further

structural, functional and inhibitor discovery studies on this important enzyme complex.

Results

Identification of F-type ATP synthase subunits from T. gondii proteome – Comprehensive *in*

silico analysis on the presence and absence of key enzymes involved in mitochondrial

metabolism in apicomplexan species has been previously carried out [25,27,41]. These

studies have highlighted the missing subunits of the F-type ATP synthase enzyme from these

organisms. We have further confirmed these findings from our efforts to identify *T. gondii*

orthologs for yeast and bovine F-type ATP synthase subunits. Orthologs for F_1 sector

subunits   α (TGME49_204400),   β (TGME49_261950),   δ (TGME49_226000),

ε (TGME49_314820),   and   γ (TGME49_231910)   and F_O sector   subunits   c

(TGME49_249720) and OSCP (TGME49_284540) were readily identified from *T. gondii*.

This suggested that all the subunits necessary for assembling the catalytic core (α3β3),

central stalk (δ, ε and γ) and the subunit c oligomer portions of the F-type ATP synthase (Fig

1A) are encoded by *T. gondii*. However, as reported previously, we were unable to identify

the orthologs for critical F_O sector subunits involved in proton translocation and formation of

the peripheral stalk structure of the enzyme in *T. gondii* through routine bioinformatic

analysis. This appears to be the case in all apicomplexan species and in other taxa within the

alveolate infrakingdom, such as the dinoflagellates, ciliates and chromerids (Fig 1B). It is

likely that the missing F_O subunits are either highly divergent or completely novel.

Interestingly, this situation is not exclusive to alveolates, since many other unicellular

eukaryotes also appear not to posses the corresponding orthologs for many of the yeast or

bovine F_O subunits. In fact, novel F-type ATP synthase associated proteins have been

previously identified from *Tetrahymena* [8], *Chlamydomonas* [6] and *Trypanosoma* [7].

Therefore, we attempted to purify the F-type ATP synthase enzyme from isolated *T. gondii*

mitochondria in native form and identify its protein composition by mass spectrometry.

Generating transgenic parasites expressing YFP-HA tagged TgATP–β and TgATP-OSCP

proteins – To facilitate the purification of the *T. gondii* F-type ATP synthase enzyme in

native form, we engineered in-frame genomic YFP-HA tags in the 3' end of the respective

genes encoding the F_1 β and F_O OSCP subunits (Fig 2A). The modified genes continued to be

expressed under the control of their endogenous promoters. The correct insertion of the tags

was confirmed by genomic PCRs (S1 Data) from clonal isolates of the respective transgenic

parasites. Expression of the desired YFP-HA tagged proteins, and their mitochondrial

localization, were confirmed by Western blotting (Fig 2B) and microscopy (Fig 2C),

respectively. We then compared the ability of the parental and transgenic strains of the

parasites to maintain total cellular ATP levels in the presence/absence of glucose. Although

T. gondii tachyzoites are known to prefer glucose as the primary nutrient source, in the

absence of glucose they are known to switch to glutaminolysis for carbon and energy supply.

In the later case, ATP is obtained by the parasites *via* oxidative phosphorylation, and this can

be inhibited using atovaquone [25,42,43]. Like the parental parasites, transgenic parasites

expressing YFP-HA tagged F_1 β and F_O OSCP subunits were capable of maintaining cellular

ATP levels *via* oxidative phosphorylation in the absence of glucose, which was inhibited by

atovaquone (Fig 2D). These results confirm that the modification of either F_1 β or F_O OSCP

proteins with YFP-HA tag had no detrimental effect on the function of the enzyme,

implicating that the structure of the tagged enzyme remained intact.

Identification of monomer and dimer forms of T. gondii F-type ATP synthase by Blue Native

Page (BNP) analysis – The fully assembled F-type ATP synthase is known to exist as dimers

on the inner mitochondrial membrane and this dimerization is known to influence the

characteristic cristae formation by the membrane [21,24]. In case of the human malaria

parasite *P. falciparum* F-type ATP synthase, dimer formation has been observed previously

[26]. To find out whether the *T. gondii* enzyme can assemble into dimers, we carried out BNP

analysis of detergent solubilized mitochondrial preparations. After resolving the samples on a

3% - 12% gradient BNP gel capable of resolving a molecular weight range between 20 kDa

to ~1200 kDa (Fig 3A lane M), Coomassie staining revealed prominent bands at the sizes

corresponding to dimer and monomer forms of the enzyme (Fig 3A lane A). This was further

confirmed by Western blotting after BNP separation using α-HA antibodies (Fig 3A lane B),

which indicated that the dimer form of the enzyme is more abundant than the monomer form.

Next, we carried out in-gel activity assays, to detect the ATPase activity associated with the

isolated enzyme. Results from these assays confirmed that the ATPase activity was intact, for

both dimer and monomer forms of the enzyme, after BNP separation (Fig 3B). We found

that, although the dimer form of the enzyme was more abundant, its ATPase activity was less

than that of the monomer form. This is supported by the fact that F_1 ATPase activity is

inhibited *in vivo* to minimize the risk of ATP hydrolysis and favor ATP synthesis [44]. Since

the in-gel activity assay is an ATP hydrolysis assay, the dimer has less of this activity. Based

on BNP mobility, we deduce the size of the dimer form of the enzyme to be between 1 – 1.2

MDa and that of the monomer form as ~600 kDa. This is in agreement with what has been

reported previously for *P. falciparum* [26], *S. cerevisiae* [19], and bovine [45] enzymes, and

suggests that a full complement of F_O subunits is present in *T. gondii* F-type ATP synthase.

Therefore, we excised the regions in the BNP corresponding to the dimer and monomer

forms of the enzyme and proceeded to identify the proteins in the gel band by LC-MS

analysis (Fig 3A).

Identification of T. gondii F-type ATP synthase associated proteins by LC-MS/MS proteomics

– A standard protocol (see methods section) was followed for in-gel LC MS/MS analysis of

the gel bands corresponding to the dimer and monomer forms of the enzyme identified by

BNP. This was done on samples prepared from both *Tgatp−β−YFP-HA* and *Tgatp-*

OSCP−YFP-HA transgenic parasites. Since the dimer form of the enzyme was more abundant

than the monomer form after BNP separation, we had better success with LC-MS/MS

analysis of the gel band corresponding to the dimer form of the enzyme. A total of 96

proteins were identified with high confidence from consensus data obtained from multiple in-

gel LC MS/MS analysis (Fig 4). Proteins corresponding to all F_1 subunits and F_O OSCP were

detected, while F_O subunit c was not detected in any of the experiments. In addition, many

proteins of unknown function were also detected. However, due to co-migration of several

non-specific proteins, such as myosin, in the area corresponding to the gel band processed for

LC MS/MS analysis, it is likely that some of these proteins of unknown function are not

bona-fide subunits of the F-type ATP synthase. A full list of the proteins identified from these

samples, along with the details of the peptides detected is given in S1 Table.

In order to specifically identify the protein subunits of F-type ATP synthase, we resorted to

partially purify the enzyme before LC MS/MS analysis by chromatographic separation (Fig

4A, B) and immunoprecipitation (IP; using the α-HA antibody). In contrast to BNP analysis,

after size exclusion chromatography we observed that the dimer form of the enzyme was less

abundant than the monomer form (Fig 4B). This is likely due to stability issues with the

dimer, which might progressively fall apart as monomers during the process of

chromatographic separation. The fractions corresponding to the dimer and monomer forms of

the enzyme were pooled and concentrated separately before processing for LC-MS/MS

analysis. A total of 64 proteins were detected (Fig 5A) from the fractions corresponding to

monomer form of the enzyme and we did not get reliable data from the dimer fractions owing

to very low protein concentration.

LC-MS/MS analysis on the enzyme enriched by IP resulted in the identification of a total of

29 proteins, of which 28 were also detected from either BNP or chromatography samples (Fig

5A). Details for the peptides detected from chromatography and IP samples, is provided in

S1 Table. It is notable that 19 proteins, including most F_1 subunits and F_O OSCP, were

detected in samples prepared by all three methods. We detected very few non-specific

proteins from the IP samples, i.e., functionally not related to F-type ATP synthase (3 out of

29). Out of the remaining 26 proteins, 5 were known F_1 subunits (α, β, δ, ϵ, γ), one was F_O

OSCP, and the remaining 20 were proteins of unknown function. It should be noted that we

were unable to identify F_O subunit c in any of our LC-MS/MS analysis, probably owing to its

highly hydrophobic nature. Based on the consensus of proteins identified by the three

different approaches, the F-type ATP synthase from *T. gondii* appears to be comprised of at

least 27 protein subunits. We have coined the termed 'ATP synthase Associated Proteins'

(ASAP) to refer to the 20 novel subunit components of the enzyme identified in this study. A

complete list of all *T. gondii* F-type ATP synthase subunits identified by mass spectrometry,

along with their annotation from *ToxoDB* and the experiments from which they were

detected, is given in the table in Fig 5B.

Identification of F_O subunits a, b and d – As shown in Fig 1, in addition to subunits c and

OSCP, the F_O sector consists of at least 6 other subunits in yeast, mammalian and plant

enzymes. Subunit a is essential for proton conductance, which it facilitates along with subunit

c [13-15]. Subunit b forms the core of the stator structure, which is essential for holding the

catalytic $\alpha_3\beta_3$ structure in place during the rotary motion of the central stalk [18]. Subunit d is

part of the stator structure. We attempted to find out which of the ASAPs correspond to these

three subunits in *T. gondii* F-type ATP synthase. Since none of the ASAPs had any sequence

identity to the yeast F_O a, b and d subunits, we resorted to conserved structure based

identification using pairwise comparison of profile hidden Markov models as implemented in

HHPred [46,47]. For this analysis we used the corresponding tool available from web based

MPI bioinformatics toolkit (*toolkit.tuebingen.mpg.de*). We analyzed each of the ASAPs using

this tool and successfully identified hits for F_O subunits a, b and d based on previously known

structures for these proteins from other species. The ASAP TGME49_310360 was identified

as F_O subunits a (Fig 6A), and we were also able to generate sequence alignments for the C-

terminal domain of this protein which showed the conservation of key amino acid residues

arginine (important for proton translocation) and glutamine (yellow highlight in Fig 6B).

Similarly, searches with TGME49_231410 and TGME49_268830 came up with hits for F_O

subunits b and d, respectively (S2 Data and S3 Data). We further validated the predictions by

HHPred with Phyre^2 [48], which is a structure-based remote homology prediction tool, and

confirmed the identification of F_O subunits a, b and d from *T. gondii* (data not shown).

Mitochondrial localization of ASAPs – Confirming the mitochondrial localization for the

ASAPs is essential for validating their subunit membership in *T. gondii* F-type ATP synthase.

We first predicted the presence of signal sequence for mitochondrial targeting using the

Mitoprot tool [49] in twelve ASAPs (Fig 7). Five ASAPs, including TGME49_231410 (F_O

subunit b like protein), were previously shown to localize in the mitochondrion [50], and we

experimentally confirmed the mitochondrial localization for another four (S4 Data).

Moreover, unpublished results from Hyper-LOPIT analysis [51] of *T. gondii* tachyzoite

proteome in an independent study (personal communication, Dr. Ross Waller, Cambridge

University, UK), confirmed that all ASAP proteins (except TGME49_310360 and

TGME49_249720, for which no data were available) are localized to the mitochondria (Fig

7).

Phylogenetic analysis reveals conserved F-type ATP synthase subunit composition in three

major alveolate taxons – From our ortholog identification protocol, and from previous studies

[27,41], it was apparent that the F-type ATP synthase subunits missing from *T. gondii* were

also missing from all other species grouped within the *Alveolata* infrakingdom. In fact, the

novel subunit components identified from *Tetrahymena* (*Alveolata*; *Ciliophora*) enzyme were

found to be unique to ciliates and not conserved in other alveolate organisms [8]. Therefore,

we were interested in finding out whether the novel ASAPs identified in this study are unique

to *T. gondii* F-type ATP synthase. Interestingly, we were able to identify orthologs for many

of the ASAPs in three major alveolate taxons – *Apicomplexa*, *Chromerida* and *Dinoflagellata*

(Fig 7). A list of all orthologs identified from selected species belonging to these taxon are

given in S2 Table. Out of the 20 ASAPs, 15 were conserved in all apicomplexan clades,

except in case of *Cryptosporidium*, where only *C. muris* contained orthologs for 10 of these

proteins. A few ASAPs were unique to the *Coccidian* clade. More importantly, all ASAPs,

except one, were conserved in *Chromerida*, and at least 9 and 12 ASAPs were also conserved

in *Symbiodinium* and *Perkinsus* respectively.

To obtain further insights on the evolutionary origin of these proteins, we generated

neighbor-joining phylogenetic trees for ortholog sequences of all F_1 and F_O subunits from

representative species of *Haemosporida*, *Piroplasmida*, *Coccidia*, *Cryptosporidiidae*,

Chromerida, Dinoflagellata and *Ciliophora* (Fig 7 and S5 Data). *Stramenopile* (*Ectocarpus*

siliculosis, *Phaeodactylum tricornumtum*, *Pythium ultimum*, *Thalassiosira pseudonana*) and

Plantae (*Chlamydomonas reinhardtii* and *Arabidopsis thaliana*) species were included as

outgroups while constructing the phylogenetic trees for the highly conserved F_1 subunits.

Even though the topology of most trees did not reflect the expected evolutionary relationship

between the included species, monophyletic grouping was observed in general at the taxon

level, and importantly, this was evident for the conserved F_1 / F_O subunits, as well as the

novel ASAPs (S5 Data). Thus, the evolutionary origin of the newly identified highly

divergent ASAPs in *Apicomplexa*, *Chromerida* and *Dinoflagellata* clades appears to be

ancient.

Essentiality and gene co-expression analysis of ASAPs from T. gondii and P. falciparum –

Due to the functional importance of the F-type ATP synthase enzyme, it is reasonable to

expect that the enzyme would be essential in *T. gondii*. This is indeed the case, and all known

subunits and ASAPs (except ASAP-19 and 20; Fig 7) of the enzyme were found to be

essential in a previous study [50]. In order to obtain further independent evidence for ASAPs

as *bona fide* subunits of F-type ATP synthase, we carried out transcriptome co-expression

correlation analysis using publicly available gene expression datasets for *T. gondii* and *P.*

falciparum [52,53]. We assumed that the ASAPs interacting with one other would show

significant pair-wise correlation profiles in their expression levels. Strikingly, we found very

high correlation of gene co-expression profiles among the F-type ATP synthase subunits, in

comparison to co-expression with other unrelated gene pairs, in both *T. gondii* and *P.*

falciparum transcriptome datasets (Fig 8). This finding further supports the fact that the novel

ASAPs are indeed co-expressed together and are likely *bona fide* subunits of the unusual F-

type ATP synthase enzyme from *T. gondii*.

Discussion

Mitochondrial oxidative phosphorylation is an important source of ATP in most

eukaryotic organisms, and is facilitated by the multimeric F-type ATP synthase enzyme

complex [1,2,5]. The enzyme consists of two functionally distinct F_1 and F_O sectors, which

act in concert to convert the electrochemical energy into mechanical energy to facilitate ATP

synthesis. F_O sector subunits a and c come together to form the proton translocation channel

across the inner mitochondrial membrane. Subunit c, which forms a membrane embedded

oligomeric ring, converts the proton gradient energy into mechanical energy *via* subunit

rotation. The F_1 sector $\alpha_3\beta_3$ catalytic core is connected to the rotary motor formed by F_O

subunit c ring *via* the rotating central stalk structure comprised of the γ, δ and ϵ F_1 subunits

[1,2,5]. The stator structure, comprised primarily of the F_O subunits b, d, h and OSCP, along

with other accessory subunits helps in holding the $\alpha_3\beta_3$ catalytic core in place while the

central stalk rotates [17]. The asymmetric structure of the γ subunit of the rotating central

stalk induces conformational changes in the $\alpha_3\beta_3$ catalytic core, which facilitates ATP

synthesis. Thus all subunit components and structural features of this unique enzyme are

critical for efficient ATP synthesis.

In addition to the core subunit composition described, other accessory proteins

participate in assembling the V shaped dimer form of the enzyme complex [19]. The paired

arrangement of the F_1 complex (spaced by 12 nm) was observed in *Paramecium* inner

mitochondrial membrane by freeze-fracture and deep-etching studies [54]. Subsequent

studies on yeast, bovine and other species confirmed the dimeric arrangement of the F-type

ATP synthase complex by BNP, atomic force microscopy and cryo-electron microscopy

[19,55,56]. In fact, the dimer form of the enzyme appears to facilitate cristae formation by the

inner mitochondrial membrane [21,22,24] and is implicated in maintenance of the

mitochondrial membrane potential [57]. However, the identity of the proteins responsible for

dimer formation has not been ascertained in most species. In case of the yeast enzyme,

subunits e, g, k were identified as dimer specific components, and studies with genetic

mutants revealed that subunits e and g are essential in dimer formation [19,20]. Orthologs for

these proteins are present in mammalian species as well.

BNP and size exclusion chromatography studies on purified yeast F-type ATP

synthase revealed that the average molecular size of the dimer and monomer form of the

enzyme is around 1 MDa and between 500 – 600 kDa, respectively [19]. Given the complex

structure and assembly of the enzyme, the expected number of protein subunits in an intact

dimer is around twenty, based on yeast and mammalian enzyme compositions [10,11].

Interestingly, while the F_1 subunits are well conserved across various taxa, the F_O subunits,

except c and OSCP, appear to be poorly conserved and in many instances are not readily

identified from sequence. This is especially true for a variety of unicellular eukaryotes,

including many free living and parasitic protists, as evident from KEGG data (*kegg.jp*).

Comparative genomics studies revealed that orthologs for yeast and mammalian F_O subunits,

except c and OSCP, are missing in the entire apicomplexan phylum [25-27]. In fact, along

with *Apicomplexa*, this appears to be the case for all other taxa belonging to the alveolata

infrakingdom. A study on the subunit composition for F-type ATP synthase from the ciliate

Tetrahymena thermophila, an alveolate species, resulted in the identification of 13 novel

proteins [8]. Surprisingly, these proteins were unique to ciliates, indicating the possible

existence of unique set of F_O components in other alveolate species as well.

The *P. falciparum* enzyme was identified in dimer form with a molecular size of > 1

MDa, confirming the presence of novel F_O subunits [26]. This study also revealed that the

enzyme is essential in *P. falciparum*. However, in case of *P. berghei*, F_1 subunit β was found

to be essential for development in mosquito, but not for survival of blood stage parasites in

mammalian host [58]. Studies with metabolic mutants in *T. gondii* have revealed that

mitochondrial oxidative phosphorylation is an important source of ATP, especially during

glucose deprivation and in the absence of a glycolytic flux [34,35]. Mitochondrial ATP

synthesis is also likely essential for formation and maintenance of tissue cyst forms of *T.*

gondii in infected hosts. Despite the importance of F-type ATP synthase in apicomplexan

parasites, not much is known about its subunit composition, structure and function in these

parasites.

In this study, we have successfully identified the subunit constituents of F-type ATP

synthase from *T. gondii*, a model apicomplexan parasite. Intact monomer and dimer forms of

the enzyme can be identified by BNP analysis from detergent solubilized parasite

mitochondria preparations. We have identified 20 novel proteins (ASAPs) as subunit

constituents of the *T. gondii* F-type ATP synthase by LC-MS/MS analysis of partially

purified enzyme using BNP, IP and chromatographic techniques. While some of these

proteins could be counterparts of the missing F_O subunits, others are most likely accessory

proteins which may be involved in dimer formation. Importantly, we were able to identify

putative F_O subunits a (TGME49_310360), b (TGME49_231410) and d (TGME49_268830),

based on conserved structural features. Although the F_O subunit a from yeast is known to

have six transmembrane domains, the *T. gondii* protein has only three. Nevertheless, the

critical arginine residue, which is important for proton translocation [14], appears to be

conserved (Fig 6). The hallmark feature of F_O subunit b from yeast and other species is the

presence of an extended helix that spans the distance between the membrane and the $\alpha_3\beta_3$

catalytic core. Although there was very poor sequence conservation, very high structural

similarity (>97% probability in HHPred assignment) can be found between yeast and *T.*

gondii F_O subunit b proteins. F_O subunit d has no transmembrane domain and is known to

interact with F_O subunit b *via* parallel/anti-parallel coiled coil helical domains [2]. The

putative *T. gondii* F_O subunit d was highly similar in secondary structure features over the

conserved regions with the bovine enzyme. Further follow up work on the 3D structure of the

enzyme is needed to verify these findings.

Interestingly, all F_1 subunits, and F_O c and OSCP subunits, along with 18 ASAPs,

were found to be essential in *T. gondii*, in a genome wide CRISPR/Cas9 mediated gene

knockout screen [50] (Fig 7). In the same study, some ASAPs were shown to be localized in

the mitochondrion. We interrogated the mitochondrial localization for all subunits of *T.*

gondii F-type ATP synthase by first considering evidence from *in silico* prediction of

mitochondrial localization signals, followed by experimental localization of selected ASAPs

(Fig 7). We also obtained confirmation for mitochondrial localization of all but two ASAPs

from Hyper-LOPIT analysis of tachyzoite stage *T. gondii* (unpublished data from an

independent study, Dr. Ross Waller, University of Cambridge, UK; personal

communication). Further, we found that the transcript co-expression for the subunits of F-

type ATP synthase was highly correlated, independently in both *T. gondii* and *P. falciparum*

transcriptome datasets (Fig 8), thus providing additional evidence for functional interaction

between these proteins across apicomplexan parasite taxa. In summary, evidence from gene

essentiality, subcellular localization and transcript co-expression together provide

independent levels of support for our identification of ASAPs as *bona fide* subunits of *T.*

gondii F-type ATP synthase.

As can be expected for such an important protein complex playing a fundamental

biological role, phylogenetic analysis revealed that most ASAPs identified from *T. gondii* are

conserved across the phylum *Apicomplexa*. This suggests that the origin of a divergent F-type

ATP synthase may predate the origin of apicomplexan parasites. In support of this

conclusion, we find that many of the ASAPs are also conserved in two other alveolate

phylums, *Chromerida* and *Dinoflagellata*. This opens up the opportunity for studying the

structure and function of this unusual enzyme from free-living alveolate species, from which

the enzyme can be isolated in native conditions at higher yields and purity. Moreover, a

deeper understanding of the structure and function of this enzyme will facilitate the discovery

of anti-parasitic compounds with pan-apicomplexan effect.

Methods

Sequence analysis and ortholog identification -

Molecular reagents and methods – Genomic DNA and total RNA from tachyzoite stage *T. gondii*, and plasmid DNA from recombinant *E. coli* (DH5α) were isolated using Qiagen kits (Germany); cDNA was prepared from total RNA using the reverse transcription kit from Thermo Fisher Scientific (USA). The manufacturer's protocol was followed while using the various kits. Polymerase chain reactions (PCR) was done using a proof reading DNA polymerase obtained from Takara Bio (Japan), and all primers used in this study (S3 Table) were obtained from Integrated DNA Technologies (USA). Ligase and various restriction endonucleases were purchased from New England BioLabs (USA) or Promega (USA).

Plasmid constructs – Various PCR products were cloned into the Topo 2.1 plasmid backbone (Thermo Fisher Scientific, USA) and sequenced before sub-cloning into other plasmids used for either modification of target genomic loci or ectopic expression of full-length cDNA for gene of interest in *T. gondii*. The plasmid construct used for tagging the 3' end of F-type ATP synthase subunits β (*Tg*atpβ) and OSCP (*Tg*atpOSCP) with yellow fluorescent protein plus hemagglutinin (YFP-HA) tag had the following features: 2606 bp (BglII and AvrII) and 1831 bp (BamHI and AvrII) PCR fragments amplified from the 3' end of *Tg*atpβ and *Tg*atpOSCP loci respectively, excluding the stop codon, were cloned into the Topo 2.1 plasmid, upstream of a YFP-HA tag coding sequence, followed by the 3' UTR of the *T. gondii* dihydrofolate reductase-thymidilate synthase (*Tg*dhfr-ts) gene [59]. A modified pBluescript plasmid backbone was used for ectopic expression of full-length cDNA of selected *T. gondii* genes encoding the novel F-type ATP synthase subunits identified from this work. The cDNAs were cloned as BamHI and XbaI/NheI fragments in-frame to a HA tag, for constitutive

expression using the *T. gondii* β-tubulin gene promoter and *Tg*dhfr-ts gene 3' UTR. For

isolating clonal lines of transgenic parasites, the respective plasmid constructs include either

the DHFR cassette (for endogenous tagging; expresses a mutant version of the *Tg*dhfr cDNA

that confers pyrimethamine resistance [60] or the CAT cassette (for ectopic expression;

expresses the chloramphenicol acetyl transferase gene, which confers resistance to

chloramphenicol [61].

T. gondii culture and genetic manipulation – Tachyzoite stage of *T. gondii* parasites (Type-I

RH (RH) and Type-I RH ΔKu80 [62] (RHΔKu80)) strains were propagated following

published protocols [63]. Briefly, Human foreskin fibroblast (HFF), which were used as host

cells for parasite infection were grown in Dulbecco's modified Eagle medium (DMEM High

glucose) supplemented with 10% heat-inactivated fetal bovine serum, 2 mM GlutaMAX, 25

mM HEPES and 50 µg/ml Gentamicin. All cell culture grade reagents were procured from

Thermo Fisher Scientific, USA. The HFF cells were maintained at 37°C in a humidified

atmosphere containing 5% CO_2. *T. gondii* tachyzoites were propagated by inoculating ~10^5

freshly isolated tachyzoites into T25 flasks containing two-weeks old confluent HFF cell

monolayer. The culture medium used for parasite growth is similar to that used for host cell

growth except that it lacks serum. Parasites are harvested 48 h after infection by scrapping the

host cell monolayer, physically disrupting the HFF cell suspension by passing it repeatedly

through a 22-gauge needle, and filtering it through a 3 µM nucleopore membrane (Whatman,

GE Healthcare ,USA (to obtain a suspension of parasites devoid of HFF cell debris .The

number of parasites present in 1 ml of suspension is estimated from cell counts obtained

using a hemocytometer.

For generating transgenic parasites expressing the *Tg*ATPβ and *Tg*ATPOSCP proteins from

the endogenous loci, the RHΔKu80 parasites were transfected [62] with the respective

tagging plasmids. Transfection was achieved by electroporating 10^7 freshly harvested

tachyzoite stage parasites resuspended in 400 µl parasite culture medium containing 50 µg of

linearized sterile plasmid DNA, using a BioRad (USA) Gene Pulser system using 10µF

capacitance, ∞ ohms resistance and 1.5 kV voltage settings. Transfected parasites were

immediately inoculated into a T25 flask containing HFF monolayer and allowed to invade

and replicated for 12-15 hours before beginning drug selection with 1 µM pyrimethamine.

For ectopic expression of cDNA with HA tags for selected genes, RH parasites were

transfected with the respective plasmid constructs and stable transformants were selected

using 20 µM chloramphenicol. Clonal lines of stable transgenic parasites were isolated using

the limiting dilution technique [63].

Measurement of intracellular ATP - For this assay, freshly isolated extracellular tachyzoite

stage parasites were incubated in culture media which either contained 5.5 mM glucose and 4

mM glutamine or 0 mM glucose and 4 mM glutamine, in order to evaluate the ability of the

parasites to maintain ATP homeostasis. The anti-parasitic drug atovaquone [43], an inhibitor

of mtETC, was used to specifically inhibit mitochondrial ATP synthesis. After 2 h of

incubation, total cellular ATP was measured using the ViaLight Plus Cell Proliferation and

Cytotoxicity Bioassay Kit (Lonza, Switzerland) as per the manufacture protocol. The

luminescence readout was quantified using the Varioskan Flash plate reader (Thermo Fisher

Scientific, USA).

Microscopy and Western blotting – For visualizing the expression and subcellular

localization of selected ASAPs fused to a reporter tag, replicating intracellular tachyzoite

stage parasites stably expressing these proteins were imaged by microscopy. Before imaging

the parasite infected HFF monolayers cultured on glass coverslips were washed with 1X

phosphate buffered saline (PBS), and fixed using 4% paraformaldehyde for 20 min, before

again washing and mounting on glass slides using the fluoroshield reagent (Sigma, USA). To

counterstain the parasite mitochondrion, the cells were treated with 250 nM Mitotracker Red

(Thermo Fisher Scientific, USA) for 20 minutes before fixing. The slides were imaged using

the 63X oil immersion objective fitted to the Axio Observer inverted fluorescent microscope

(Carl Zeiss, Germany), and images were processed using the Zen software (Carl Zeiss,

Germany). Intrinsic fluorescence from YFP and Mitotracker Red were visualized using the

excitation/emission filter combination of 493/520 and 578/599 respectively. HA tagged

proteins were visualized by immunofluorescence staining, using rabbit α-HA primary

antibodies (1:1000) followed by Alexa 488 conjugated goat anti rabbit secondary antibody

(1:1000), both purchased from Thermo Fisher Scientific, USA. After fixing, the cells were

permeabilized for 5-10 minutes using 0.25% Triton X-100 in 1X PBS. They were then treated

with 2% fetal bovine serum in 1X PBS for 30 minutes, followed by primary antibody for 1 h,

3X wash with 1X PBS containing 0.25% Triton X-100, and secondary antibody for 1 h.

Finally, the cells were washed 3X with 1X PBS containing 0.25% Triton X-100, followed by

a water wash, before mounting the cover slip on glass slides for imagining.

For Western blotting, purified mitochondria were solubilized in 1X Laemmli buffer (120 mM

Tris-HCL pH 6.8, 2% SDS, 10% glycerol and 0.01% w/v bromophenol blue) and resolved by

SDS-PAGE, followed by electro-transfer to PVDF membrane (using Tris-glycine buffer pH 8

containing 20% methanol) at 65 mA for ~1 h 20 min. The blots were kept in blocking buffer

(5% skimmed milk in 1X PBS) overnight at 4°C and then treated with primary antibody

(1:5000; rabbit α-HA monoclonal antibody, Thermo Fisher Scientific, USA) for 1 h at

ambient temperature. This was followed by 3X wash with 1X PBS containing 0.1% Tween

20, before incubating with horse radish peroxidase coupled secondary antibody (1:5000;

donkey anti-rabbit antibody, Nif 824 GE Healthcare, USA) for 1 h at ambient temperature,

and 3X wash with 1X PBS containing 0.1% Tween 20. The blot was then developed using

either the 3,3'-diaminobenzidine substrate (Sigma, USA) or the chemiluminescent ECL

Western blotting kit (GE Healthcare, USA).

Preparation of mitochondria from T. gondii – Freshly harvested tachyzoite stage transgenic

parasites ($\sim 10^9$), expressing either *Tg*ATPβ-YFP-HA or *Tg*ATPOSCP-YFP-HAHA, were

washed with PBS and resuspended in 1-2 ml of hypotonic lysis buffer (15 mM phosphate

buffer pH 7.5 and 2 mM Glucose). The cells were lysed by sonication in an iced water bath

for 30 minutes, and samples were centrifuged at 2000 g for 2 minutes at 4°C to remove

unbroken cells and large cell debris. The supernatant was recovered into a new tube and

centrifuged at 21,000 g for 15 minutes at 4°C to pellet the mitochondria, which was then re-

suspended in 500 µl of mitochondria storage buffer (320 mM sucrose, 1 mM EDTA, 10 mM

Tris pH 7.4). Total protein in the mitochondria suspension was estimated by Bradford method

[64] and the lysate was stored in -80°C until further use. When required, the mitochondria

were recovered from storage buffer by centrifuging at 21,000 g for 15 minutes at 4°C and

lysed using mitochondria solubilization buffer A (MSB-A; 50 mM NaCl, 50 mM Imidazole,

2 mM 6-Aminohexoanic acid, 1 mM EDTA, pH 7.0) containing β-dodecyl maltoside (DDM)

detergent in total protein to detergent ratio of 1:5. Solubilization was allowed to continue

overnight with rocking at 4°C, and the solubilized fraction was separated by centrifuging at

100,000 g for 20 minutes at 4°C.

Blue Native Page separation, in-gel ATPase activity assay and Western blotting – In order to

identify the monomeric and dimeric forms of the *T. gondii* F-type ATP synthase, the

solubilized mitochondrial lysate was subject to one-dimensional Blue Native PAGE (BNP)

analysis [65,66]. Samples were prepared by adding glycerol to a final concentration of 5%

and Coomassie Blue G 250 5% (w/v) dye was added such that DDM detergent to dye ratio

was 8:1. Samples were separated on a 3-12 % native PAGE (Thermo Fisher Scientific, USA)

at 150 V setting for 30 minutes in cathode buffer A, which was then switched to cathode

buffer B and separation was continued till the blue dye exited the gel. The gel was visualized

under white light to identify the various dye stained bands. 25 mM imidazole pH 7.0 was

used as the anode buffer, while the composition of cathode buffers A and B were 50 mM

Tricine, 7.5 mM imidazole pH 7.0, 0.02% Coomassie Blue G 250 dye and 50 mM Tricine,

7.5 mM imidazole pH 7.0, 0.002% Coomassie Blue G 250 dye, respectively.

For in-gel ATPase assay, after sample separation by BNP, the gel was incubated overnight in

assay buffer (35 mM Tris-HCL pH 7.8, 270 mM glycine, 14 mM MgSO4, 0.2% $Pb(NO_3)_2$,

and 8 mM ATP). The gel was then washed with water and the ATPase activity of F-type ATP

synthase was observed by the formation of a milky white precipitate, visible against a black

background, at the region corresponding to the expected monomeric and dimeric protein

bands. For Western blotting after BNP separation, proteins were electro-transferred onto a

PVDF membrane using Tris-glycine buffer pH 8.0 without methanol at 65 mAh for ~1hour

20 minutes. The membrane was then fixed in 8 % acetic acid for 15 minutes, air dried at

room temperature for 30 minutes, and washed with methanol several times to remove the

Coomassie dye stain [26]. Further steps were similar to that described above for SDS PAGE

Western blotting.

Immunoprecipitation of F-type ATP synthase complex – The supernatant from solubilized

mitochondria preparation from tachyzoite stage transgenic *T. gondii* expressing YFP-HA

tagged *Tg*ATPOSCP subunit was used for immunoprecipitating the F-type ATP synthase

using α-HA antibodies (Thermo Fisher Scientific, USA). The antibodies were first cross-

linked to Protein A/G Plus agarose beads (Santa Cruz Biotechnology, USA) using the

following protocol. ~50 µl of Protein A/G Plus agarose beads were washed twice in 500 µl

1X PBS (pH 7.4) at 4°C and mixed with 1 ml of 1X PBS containing 5-7 µg of rabbit α-HA

antibody and incubate overnight at 4°C with gentle mixing on rocker. The beads were then

separated by centrifugation (1000g for 2 minute at 4°C), equilibrate with 0.2 M

Triethanolamine (pH 8.2) for 2 minutes and then washed twice with the same buffer. The

bound antibodies were then cross-linked to the beads using 20 mM DMP (Sigma, USA) in 1

ml of 0.2 M Triethanolamine at room temperature for 45 minutes, followed by washing with

1ml of 50 mM Tris pH 7.5 twice for 15 minutes to quench the cross-linking reaction.

Unbound antibodies are removed using three quick washes with 0.2 M glycine buffer pH 2.3.

The antibody conjugated beads were then washed twice with 1X PBS and equilibrate with

MSB containing the detergent DDM at critical micelle concentration. The supernatant from

solubilized mitochondria was added to the antibody coupled beads and incubated overnight at

4°C with gentle mixing on rocker. The beads were then washed three times with 1X PBS

containing DDM at critical micelle concentration. Proteins captured by the antibodies were

eluted in 100 µl of 0.2 M glycine buffer (pH 2.3) in three rounds, and the pooled eluate was

neutralized with 1M Tris (pH 8.0) and concentrated using a 3 KDa cut-off concentrator

(Amicon Millipore, Germany). The samples were equilibrated with 0.1% RapiGest (Waters,

USA) in 50 mM ammonium bicarbonate buffer in preparation for LC-MS analysis.

Purification of native F-type ATP synthase by chromatography – We followed a protocol that

was previously reported for the purification of the F-type ATP synthase from the *Polytomella*

sp. [67]. About 13 mg (wet weight) of mitochondrial preparation from transgenic *T. gondii*

expressing YFP-HA tagged *Tg*atpβ subunit was solubilized overnight in 3 ml of

mitochondrial solubilization buffer B (MSB-B; 50 mM Tris HCl (pH 8.0), 1 mM $MgCl_2$, and

DDM at protein to detergent ration 1:5). The supernatant was loaded onto a HiTrap DEAE

sepharose column (GE Healthcare Life sciences, USA) having a bed volume of 1 ml and

equilibrated with buffer A (50 mM Tris HCl (pH 8.0) $MgCl_2$ 1mM, 0.05 % (w/v) DDM).

After completion of loading, the column was first washed with 10 column volumes of buffer

B_{20} (50 mM Tris HCl (pH 8.0), 1 mM $MgCl_2$, 20 mM NaCl, and 0.05 % (w/v) DDM) and

eluted with a 10X column volume linear gradient using buffer B_{20} and buffer B_{500} (50 mM

Tris HCl (pH 8.0), 1 mM $MgCl_2$, 500 mM NaCl, and 0.05 % (w/v) DDM) at a flow rate of

0.5 ml per minute. 500 µl fractions were collected and checked by Western blotting using α-

HA antibodies. Peak positive fractions were identified, pooled and concentrated to a final

volume of 200 µl using the Vivaspin concentrator (100 kDa cut-off; GE Healthcare Life

sciences, USA) and separated by size exclusion chromatography using the Superose 6

Increase 3.2/300 column (GE Healthcare Life sciences, USA) having a bed volume of 2.4 ml.

50 µl sample was loaded on the column previously equilibrated with buffer B_{20}, and then

eluted with 1.5 column volumes of the same buffer at 50 µl flow rate. 100 µl fractions were

collected and positive fractions, identified by Western blotting using α-HA antibodies, were

pooled and concentrated using a 3 KDa cut-off concentrator (Amicon Millipore, Germany).

Samples were then equilibrated with 0.1% RapiGest in 50 mM ammonium bicarbonate buffer

in preparation for LC-MS analysis. All chromatography steps were carried out using the

AKTApure system (GE Healthcare Life sciences, USA).

LC-MS proteomics – In order to identify the subunit components of the F-type ATP synthase

from *T. gondii*, we performed LC-MS analysis on the partially purified enzyme obtained

from BNP, immunoprecipitation and chromatographic purification. Sample preparation for

LC-MS analysis was based on a previous report [68].

Sample preparation after BNP separation: The bands corresponding to the dimer and

monomer forms of the protein were excised out from BNP gel, cut to small pieces, destained

with 50% acetonitrile in 50 mM ammonium bicarbonate, and dehydrate with 100 %

acetonitrile. The gel pieces were then treated with 10 mM DTT in 50 mM ammonium

bicarbonate for 45-60 minutes at 56ºC to reduce the proteins, followed by alkylation in dark

with 55 mM Iodoacetamide in 50 mM ammonium bicarbonate at ambient temperature for 45

minutes. Then, trypsinization of protein was carried out overnight at 37°C, and peptides were

extracted using 50% acetonitrile in 2% formic acid. The extracts were dried using a speedvac

and the peptides were reconstituted in 50 µl of 50 mM ammonium bicarbonate, acidified with

HCl and desalted using the C_{18} Zip tip columns (Millipore, Germany). The samples were then

dried in a speedvac and stored at -80°C until further use.

Sample preparation after Immunoprecipitation and size exclusion chromatography:

Samples obtained from immunoprecipitation and size exclusion chromatography were

immediately equilibrated with 0.1 % RapiGest as stated above. The mixtures were then

heated to 80°C and maintained at this temperature for 15 minutes. The denatured proteins

were reduced by 100 mM DTT at 60°C for 15 minutes, followed by alkylation with 200 mM

iodoacetamide at ambient temperature and in dark for 30 minutes. Samples were then treated

overnight with trypsin (Sigma Aldrich, USA) at 20:1 substrate to enzyme ratio at 37°C.

Trypsinization was stopped by addition of 2 µl of 1N HCL and incubating at 37°C for 20

minutes, after which they were vortexed and centrifuged. The peptide samples were desalted

using the C_{18} Zip tip (Millipore, Germany), dried and stored at -80°C until further use.

LC-MS analysis: The peptide samples were analyzed using LC-MS[E] workflow on the

nano ACQUITY (UPLC) Synapt system (Waters Corporation, USA). 4 µl of the digested

peptide sample was injected into a 5 µm Symmetry C_{18} trapping column (180 µm x 20 mm)

at a flow rate of 5 nl/min and peptides were eluted by using the following protocol: 3-40 % B

(B - 100% acetonitrile with 0.1% formic acid) for 90 minutes, 40-85% B from 90-105

minutes and 97% A (A - 0.1% formic acid in water) and 3% B from 105-120 minutes, on a

bridge ethyl hybrid C_{18} column (75 μm x 250 mm, 1.7 μm) at flow rate of 250 nl/min. The

system was coupled to the Synapt High Definition Mass Spectrometer (Waters Corporation,

USA) with a nonoLock spray as the ion source. Standard glu-fibrinopeptide B (Sigma

Aldrich, USA) as the lockmass calibrate peptide was infused into ion source having 500

nl/min flow rate and sampled every 30 second. Acquisition of LC-MS^E data was performed in

positive V mode with mass range m/z 50-1900 having scan time of 0.75 second, a constant

low energy of 4 V for MS mode, and 20-40 V of collision energy during high energy MS^E

mode scan. A capillary voltage of 3.5 kV and cone voltage 38 V were maintained during

analysis. LC-MS^E data were processed using Protein Lynx Global Server (PLGS Version

2.5.3, Waters Corporation, USA) for identifying the proteins, with reference to the *T. gondii*

proteome taken from *ToxoDB.org* release 36.

Gene co-expression analysis and phylogenetic studies – Normalized signal intensity values

from microarray hybridizations performed at 13 different time points in the tachyzoite stage

[52] were obtained directly from *EupathDB.org*. Normalized microarray intensities from time

series data for intraerythrocytic stage *P. falciparum* were retrieved from reference [53].

Pearson correlation coefficients were calculated for all gene pairs using R (*R-project.org*). P-

values were calculated using the Mann-Whitney U test [69] for co-expression between ATP

synthase genes vs. co-expression between ATP synthase genes and non- ATP synthase genes,

and for co-expression between ATP synthase genes vs. co-expression between non- ATP

synthase genes. Since minute differences in very large samples may lead to very small p-

values, the effect size was therefore tested using Cohen's d test [70], where values above 0.8

denote large effect size. Orthologs for the novel subunits of *T. gondii* ATP synthase were

identified as reported previously [41]. Sequences were aligned with mafft (v7.222) [71] and

Neighbor-Joining trees were constructed using Clustalw (2.1) [72] using 1000 bootstrap

replicates. The expected 'true' phylogeny was adopted from a previous study [41].

Acknowledgements

The project is supported by funding from Science & Engineering Research Board (SERB),

India (EMR/2016/003588), awarded jointly to DS and MB and This publication is also based

upon work supported by the King Abdullah University of Science and Technology (KAUST)

Office of Sponsored Research (OSR) under Award No. OCRF-2014-CRG3-2267 and the

faculty baseline funding (BAS/1/1020-01-01) awarded to AP. RS received fellowship from

Council of Scientific and Industrial Research, India. Authors would like to thank Dr. Mahesh

J. Kulkarni, CSIR-National Chemical Laboratory, Pune, India, for facilitating the mass

spectrometry work, and Prof. David. S. Roos, University of Pennsylvania, Philadelphia,

USA, for providing *T. gondii* strains and other biological materials as gifts.

Author contributions

DS conceived the project and designed the experiments along with RS, MB and AP. RS and

TM executed the various experiments. DS, RS, AP and TM were involved in manuscript

preparation.

Competing interests

The authors declare no competeing interests, financial or otherwise, related to this

manuscript.

References

1. Capaldi RA, Aggeler R (2002) Mechanism of the F(1)F(0)-type ATP synthase, a biological
rotary motor. Trends Biochem Sci 27: 154-160.
2. Devenish RJ, Prescott M, Rodgers AJ (2008) The structure and function of mitochondrial
F1F0-ATP synthases. Int Rev Cell Mol Biol 267: 1-58.
3. Mitchell P (1966) Chemiosmotic coupling in oxidative and photosynthetic
phosphorylation. Biol Rev Camb Philos Soc 41: 445-502.
4. Stock D, Leslie AG, Walker JE (1999) Molecular architecture of the rotary motor in ATP
synthase. Science 286: 1700-1705.
5. Boyer PD (1997) The ATP synthase--a splendid molecular machine. Annu Rev Biochem
66: 717-749.
6. Vazquez-Acevedo M, Cardol P, Cano-Estrada A, Lapaille M, Remacle C, et al. (2006) The
mitochondrial ATP synthase of chlorophycean algae contains eight subunits of unknown
origin involved in the formation of an atypical stator-stalk and in the dimerization of the
complex. J Bioenerg Biomembr 38: 271-282.
7. Zikova A, Schnaufer A, Dalley RA, Panigrahi AK, Stuart KD (2009) The F(0)F(1)-ATP
synthase complex contains novel subunits and is essential for procyclic Trypanosoma
brucei. PLoS Pathog 5: e1000436.
8. Balabaskaran Nina P, Dudkina NV, Kane LA, van Eyk JE, Boekema EJ, et al. (2010)
Highly divergent mitochondrial ATP synthase complexes in Tetrahymena thermophila.
PLoS Biol 8: e1000418.
9. Yadav KNS, Miranda-Astudillo HV, Colina-Tenorio L, Bouillenne F, Degand H, et al.
(2017) Atypical composition and structure of the mitochondrial dimeric ATP synthase
from Euglena gracilis. Biochim Biophys Acta 1858: 267-275.
10. Velours J, Arselin G (2000) The Saccharomyces cerevisiae ATP synthase. J Bioenerg
Biomembr 32: 383-390.
11. Meyer B, Wittig I, Trifilieff E, Karas M, Schagger H (2007) Identification of two proteins
associated with mammalian ATP synthase. Mol Cell Proteomics 6: 1690-1699.
12. Ruhle T, Leister D (2015) Assembly of F1F0-ATP synthases. Biochim Biophys Acta
1847: 849-860.
13. Cain BD, Simoni RD (1989) Proton translocation by the F1F0ATPase of Escherichia coli.
Mutagenic analysis of the a subunit. J Biol Chem 264: 3292-3300.
14. Valiyaveetil FI, Fillingame RH (1997) On the role of Arg-210 and Glu-219 of subunit a
in proton translocation by the Escherichia coli F0F1-ATP synthase. J Biol Chem 272:
32635-32641.
15. Rastogi VK, Girvin ME (1999) Structural changes linked to proton translocation by
subunit c of the ATP synthase. Nature 402: 263-268.
16. Diez M, Zimmermann B, Borsch M, Konig M, Schweinberger E, et al. (2004) Proton-
powered subunit rotation in single membrane-bound F0F1-ATP synthase. Nat Struct Mol
Biol 11: 135-141.
17. Weber J (2006) ATP synthase: subunit-subunit interactions in the stator stalk. Biochim
Biophys Acta 1757: 1162-1170.
18. Walker JE, Dickson VK (2006) The peripheral stalk of the mitochondrial ATP synthase.
Biochim Biophys Acta 1757: 286-296.
19. Arnold I, Pfeiffer K, Neupert W, Stuart RA, Schagger H (1998) Yeast mitochondrial
F1F0-ATP synthase exists as a dimer: identification of three dimer-specific subunits.
EMBO J 17: 7170-7178.

20. Brunner S, Everard-Gigot V, Stuart RA (2002) Su e of the yeast F1Fo-ATP synthase
forms homodimers. J Biol Chem 277: 48484-48489.
21. Paumard P, Vaillier J, Coulary B, Schaeffer J, Soubannier V, et al. (2002) The ATP
synthase is involved in generating mitochondrial cristae morphology. EMBO J 21: 221-
230.
22. Dudkina NV, Heinemeyer J, Keegstra W, Boekema EJ, Braun HP (2005) Structure of
dimeric ATP synthase from mitochondria: an angular association of monomers induces
the strong curvature of the inner membrane. FEBS Lett 579: 5769-5772.
23. Dudkina NV, Sunderhaus S, Braun HP, Boekema EJ (2006) Characterization of dimeric
ATP synthase and cristae membrane ultrastructure from Saccharomyces and Polytomella
mitochondria. FEBS Lett 580: 3427-3432.
24. Davies KM, Anselmi C, Wittig I, Faraldo-Gomez JD, Kuhlbrandt W (2012) Structure of
the yeast F1Fo-ATP synthase dimer and its role in shaping the mitochondrial cristae. Proc
Natl Acad Sci U S A 109: 13602-13607.
25. Mather MW, Henry KW, Vaidya AB (2007) Mitochondrial drug targets in apicomplexan
parasites. Curr Drug Targets 8: 49-60.
26. Balabaskaran Nina P, Morrisey JM, Ganesan SM, Ke H, Pershing AM, et al. (2011) ATP
synthase complex of Plasmodium falciparum: dimeric assembly in mitochondrial
membranes and resistance to genetic disruption. J Biol Chem 286: 41312-41322.
27. Danne JC, Gornik SG, Macrae JI, McConville MJ, Waller RF (2013) Alveolate
mitochondrial metabolic evolution: dinoflagellates force reassessment of the role of
parasitism as a driver of change in apicomplexans. Mol Biol Evol 30: 123-139.
28. Weiss LM, Dubey JP (2009) Toxoplasmosis: A history of clinical observations. Int J
Parasitol 39: 895-901.
29. Dubey JP (1998) Advances in the life cycle of Toxoplasma gondii. Int J Parasitol 28:
1019-1024.
30. Dubey JP, Lindsay DS, Speer CA (1998) Structures of Toxoplasma gondii tachyzoites,
bradyzoites, and sporozoites and biology and development of tissue cysts. Clin Microbiol
Rev 11: 267-299.
31. Polonais V, Soldati-Favre D (2010) Versatility in the acquisition of energy and carbon
sources by the Apicomplexa. Biol Cell 102: 435-445.
32. Pomel S, Luk FC, Beckers CJ (2008) Host cell egress and invasion induce marked
relocations of glycolytic enzymes in Toxoplasma gondii tachyzoites. PLoS Pathog 4:
e1000188.
33. Al-Anouti F, Tomavo S, Parmley S, Ananvoranich S (2004) The expression of lactate
dehydrogenase is important for the cell cycle of Toxoplasma gondii. J Biol Chem 279:
52300-52311.
34. Blume M, Rodriguez-Contreras D, Landfear S, Fleige T, Soldati-Favre D, et al. (2009)
Host-derived glucose and its transporter in the obligate intracellular pathogen
Toxoplasma gondii are dispensable by glutaminolysis. Proc Natl Acad Sci U S A 106:
12998-13003.
35. MacRae JI, Sheiner L, Nahid A, Tonkin C, Striepen B, et al. (2012) Mitochondrial
metabolism of glucose and glutamine is required for intracellular growth of Toxoplasma
gondii. Cell Host Microbe 12: 682-692.
36. Nitzsche R, Zagoriy V, Lucius R, Gupta N (2016) Metabolic Cooperation of Glucose and
Glutamine Is Essential for the Lytic Cycle of Obligate Intracellular Parasite Toxoplasma
gondii. J Biol Chem 291: 126-141.
37. Vercesi AE, Rodrigues CO, Uyemura SA, Zhong L, Moreno SN (1998) Respiration and
oxidative phosphorylation in the apicomplexan parasite Toxoplasma gondii. J Biol Chem
273: 31040-31047.

38. Saleh A, Friesen J, Baumeister S, Gross U, Bohne W (2007) Growth inhibition of
Toxoplasma gondii and Plasmodium falciparum by nanomolar concentrations of 1-
hydroxy-2-dodecyl-4(1H)quinolone, a high-affinity inhibitor of alternative (type II)
NADH dehydrogenases. Antimicrob Agents Chemother 51: 1217-1222.
39. Lin SS, Gross U, Bohne W (2009) Type II NADH dehydrogenase inhibitor 1-hydroxy-2-
dodecyl-4(1H)quinolone leads to collapse of mitochondrial inner-membrane potential and
ATP depletion in Toxoplasma gondii. Eukaryot Cell 8: 877-887.
40. Mogi T, Kita K (2010) Diversity in mitochondrial metabolic pathways in parasitic protists
Plasmodium and Cryptosporidium. Parasitol Int 59: 305-312.
41. Woo YH, Ansari H, Otto TD, Klinger CM, Kolisko M, et al. (2015) Chromerid genomes
reveal the evolutionary path from photosynthetic algae to obligate intracellular parasites.
Elife 4: e06974.
42. McFadden DC, Tomavo S, Berry EA, Boothroyd JC (2000) Characterization of
cytochrome b from Toxoplasma gondii and Q(o) domain mutations as a mechanism of
atovaquone-resistance. Mol Biochem Parasitol 108: 1-12.
43. Baggish AL, Hill DR (2002) Antiparasitic agent atovaquone. Antimicrob Agents
Chemother 46: 1163-1173.
44. Pullman ME, Monroy GC (1963) A Naturally Occurring Inhibitor of Mitochondrial
Adenosine Triphosphatase. J Biol Chem 238: 3762-3769.
45. Krause F, Reifschneider NH, Goto S, Dencher NA (2005) Active oligomeric ATP
synthases in mammalian mitochondria. Biochem Biophys Res Commun 329: 583-590.
46. Zimmermann L, Stephens A, Nam SZ, Rau D, Kubler J, et al. (2017) A Completely
Reimplemented MPI Bioinformatics Toolkit with a New HHpred Server at its Core. J
Mol Biol.
47. Soding J (2005) Protein homology detection by HMM-HMM comparison. Bioinformatics
21: 951-960.
48. Kelley LA, Mezulis S, Yates CM, Wass MN, Sternberg MJ (2015) The Phyre2 web portal
for protein modeling, prediction and analysis. Nat Protoc 10: 845-858.
49. Claros MG, Vincens P (1996) Computational method to predict mitochondrially imported
proteins and their targeting sequences. Eur J Biochem 241: 779-786.
50. Sidik SM, Huet D, Ganesan SM, Huynh MH, Wang T, et al. (2016) A Genome-wide
CRISPR Screen in Toxoplasma Identifies Essential Apicomplexan Genes. Cell 166:
1423-1435 e1412.
51. Dunkley TP, Watson R, Griffin JL, Dupree P, Lilley KS (2004) Localization of organelle
proteins by isotope tagging (LOPIT). Mol Cell Proteomics 3: 1128-1134.
52. Behnke MS, Wootton JC, Lehmann MM, Radke JB, Lucas O, et al. (2010) Coordinated
progression through two subtranscriptomes underlies the tachyzoite cycle of Toxoplasma
gondii. PLoS One 5: e12354.
53. Bozdech Z, Llinas M, Pulliam BL, Wong ED, Zhu J, et al. (2003) The transcriptome of
the intraerythrocytic developmental cycle of Plasmodium falciparum. PLoS Biol 1: E5.
54. Allen RD, Schroeder CC, Fok AK (1989) An investigation of mitochondrial inner
membranes by rapid-freeze deep-etch techniques. J Cell Biol 108: 2233-2240.
55. Buzhynskyy N, Sens P, Prima V, Sturgis JN, Scheuring S (2007) Rows of ATP synthase
dimers in native mitochondrial inner membranes. Biophys J 93: 2870-2876.
56. Davies KM, Strauss M, Daum B, Kief JH, Osiewacz HD, et al. (2011) Macromolecular
organization of ATP synthase and complex I in whole mitochondria. Proc Natl Acad Sci
U S A 108: 14121-14126.
57. Bornhovd C, Vogel F, Neupert W, Reichert AS (2006) Mitochondrial membrane potential
is dependent on the oligomeric state of F1F0-ATP synthase supracomplexes. J Biol Chem
281: 13990-13998.

58. Sturm A, Mollard V, Cozijnsen A, Goodman CD, McFadden GI (2015) Mitochondrial
ATP synthase is dispensable in blood-stage Plasmodium berghei rodent malaria but
essential in the mosquito phase. Proc Natl Acad Sci U S A 112: 10216-10223.
59. Roos DS (1993) Primary structure of the dihydrofolate reductase-thymidylate synthase
gene from Toxoplasma gondii. J Biol Chem 268: 6269-6280.
60. Donald RG, Roos DS (1993) Stable molecular transformation of Toxoplasma gondii: a
selectable dihydrofolate reductase-thymidylate synthase marker based on drug-resistance
mutations in malaria. Proc Natl Acad Sci U S A 90: 11703-11707.
61. Kim K, Soldati D, Boothroyd JC (1993) Gene replacement in Toxoplasma gondii with
chloramphenicol acetyltransferase as selectable marker. Science 262: 911-914.
62. Huynh MH, Carruthers VB (2009) Tagging of endogenous genes in a Toxoplasma gondii
strain lacking Ku80. Eukaryot Cell 8: 530-539.
63. Roos DS, Donald RG, Morrissette NS, Moulton AL (1994) Molecular tools for genetic
dissection of the protozoan parasite Toxoplasma gondii. Methods Cell Biol 45: 27-63.
64. Bradford MM (1976) A rapid and sensitive method for the quantitation of microgram
quantities of protein utilizing the principle of protein-dye binding. Anal Biochem 72: 248-
254.
65. Wittig I, Braun HP, Schagger H (2006) Blue native PAGE. Nat Protoc 1: 418-428.
66. Nijtmans LG, Henderson NS, Holt IJ (2002) Blue Native electrophoresis to study
mitochondrial and other protein complexes. Methods 26: 327-334.
67. Allegretti M, Klusch N, Mills DJ, Vonck J, Kuhlbrandt W, et al. (2015) Horizontal
membrane-intrinsic alpha-helices in the stator a-subunit of an F-type ATP synthase.
Nature 521: 237-240.
68. Korwar AM, Vannuruswamy G, Jagadeeshaprasad MG, Jayaramaiah RH, Bhat S, et al.
(2015) Development of Diagnostic Fragment Ion Library for Glycated Peptides of Human
Serum Albumin: Targeted Quantification in Prediabetic, Diabetic, and Microalbuminuria
Plasma by Parallel Reaction Monitoring, SWATH, and MSE. Mol Cell Proteomics 14:
2150-2159.
69. Myles Hollander DAW, Eric Chicken (2013) Nonparametric Statistical Methods, 3rd
Edition: Wiley. 848 p.
70. Cohen J (1988) Statistical power analysis for the behavioral sciences. Hillsdale, N.J.: L.
Erlbaum Associates. xxi, 567 p. p.
71. Katoh K, Standley DM (2013) MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. Mol Biol Evol 30: 772-780.
72. Larkin MA, Blackshields G, Brown NP, Chenna R, McGettigan PA, et al. (2007) Clustal
W and Clustal X version 2.0. Bioinformatics 23: 2947-2948.
73. Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, et al. (2011) Fast, scalable
generation of high-quality protein multiple sequence alignments using Clustal Omega.
Mol Syst Biol 7: 539.
74. Robert X, Gouet P (2014) Deciphering key features in protein structures with the new
ENDscript server. Nucleic Acids Res 42: W320-324.

Main Figure Legends

Fig 1. Detail of F-type ATP synthase subunits missing in *T. gondii* and other apicomplexan parasites.

(A) Schematic representation of core subunit composition of F-type ATP synthase. The different subunits are color coded and annotated. Middle, the complete set of subunits of the core enzyme; left and right, subunits identified and not identified by *in silico* methods in apicomplexan genome. (B) Table showing the presence/absence of orthologs of yeast and bovine F-type ATP synthase in *Apicomplexa* and other alveolate species. +/-, presence/absence of ortholog; blue, diverged (no clear orthologs) functional equivalent; light blue, experimentally identified species-specific novel subunits (numeric values indicate the total number of such novel subunits). Species names: *Pfal*, *P. falciparum*; *Pviv*, *P. vivax*; *Tpar*, *T. parva*; *Bbov*, *B. bovis*; *Tgon*, *T. gondii* ; *Ncan*, *N. caninum*; *Cmur*, *C. muris*; *Cpar*, *C. parvum*; *Chom*, *C. hominis*; *Cvel*, *C. velia*; *Pmar*, *P. marinus*; *Tthe*, *T. thermophila*; *Scer*, *S. cerevisiae*; *Btau*, *B. taurus*; *Atha*, *A. thaliana*; *Crei*, *C. reinhardtii*; *Tbru*, *T. brucei*.

Fig 2. Endogenous tagging of *Tgatpβ* and *Tgatposcp* genes with YFP-HA tags.

(A) Schematic representation of the *Tgatpβ* (TGME49_261950) gene locus showing coding region in light blue and 3' UTR in grey. The hatched region shows the homology region (HR), which was cloned into a donor plasmid (dark blue) upstream of an in-frame coding region for YFP-HA tag. The donor plasmid also includes the DHFR cassette, which confers pyrimethamine resistance. Asterisk denotes the position of stop codon. The donor plasmid was linearized with the restriction enzyme BstZ17I to facilitate single crossover homologous recombination and the resulting modification of the gene locus was confirmed by PCR using the primers ConF and YFP_R. A similar strategy was used to tag the *Tgatposcp* gene locus. (B) and (C) Confirming the expression of the *Tg*ATPβ−YFP-HA and *Tg*ATPOSCP-YFP-HA by SDS-PAGE Western blotting (B) and microscopy (C) on respective clonal transgenic

lines. Gel lanes in (B): 1, Cell lysate from *Tg*ATPb-YFP-HA transgenic parasites; 2, Cell lysate

from *Tg*ATPOSCP-YFP-HA transgenic parasites; 3, Cell lysate from parental parasites; M,

Molecular weight size markers. Mitotracker was used to visualize the mitochondrion in (C). (D)

Functional validation of mitochondrial ATP synthesis in *wt* parental strain, and *Tgatpβ-YFP-*

HA and *Tgatposcp-YFP-HA* expressing transgenic parasites. All three strains exhibited

similar response to atovaquone treatment in the presence and absence of glucose, thereby

confirming that the YFP-HA tag had no effect on mitochondrial ATP synthesis.

Fig 3. Identifying the dimer and monomer forms of *T. gondii* F-type ATP synthase by

BNP analysis.

(A) Mitochondria lysates were prepared from tachyzoites stage *Tgatpβ-YFP-HA* transgenic

parasites were separated by BNP. Lane M, native molecular weight markers; lane A,

Coomassie blue staining of BNP gel; lane B, Western blotting of BNP gel using α-HA

antibodies. The dimer and monomer forms are indicated by arrows. The boxed region in lane

A corresponds to the excised gel piece, which was processed for LC-MS/MS analysis. (B)

In-gel ATPase activity assays following BNP separation confirms that the dimer and

monomer forms of F-type ATP synthase are functionally intact.

Fig 4. Partial purification of dimer and monomer forms of *T. gondii* F-type ATP

synthase by chromatography.

(A) Ion exchange (DEAE sepharose) separation of mitochondrial lysates prepared from

Tgatpβ-YFP-HA transgenic parasites. Absorbance at 280 nm (filled circles) and NaCl

concentration (open circles) are plotted for each fraction. Fractions 5 and 18 are marked with

arrows. Bottom panel shows SDS-PAGE Western blotting for fractions 5-18 to find out the

elution profile of *Tg*ATPβ-YFP-HA. (B) Size exclusion profile of the pooled fractions from

ion exchange chromatography. The Absorbance at 280 nm is plotted for each fraction. The

size exclusion column was calibrated with the following native markers; thyroglobulin (T,

660 kDa), ferritin (F, 440 kDa), conalbumin (C, 75 kDa), Ovalbumin (O, 45 kDa). Peak

elution volume for each marker is indicated by arrow. Fractions 1-3, 4-6 and 7-9 were pooled,

concentrated and subject to SDS-PAGE Western blotting to detect TgATP-β-YFP-HA as

shown in bottom panel.

Fig 5. Identification of novel _T. gondii_ F-type ATP synthase subunits from LC-MS/MS analysis.

(A) The Venn diagram shows shared identification of proteins following blue native page

(BNP), size exclusion chromatography (SEC) and immunoprecipitation (IP) sample

preparation. Total proteins identified in each technique is given within brackets outside the

Venn diagram. The numbers shown in white font and within brackets are the final set of

proteins assigned as subunit components of _T. gondii_ F-type ATP synthase. (B) List of all

genes identified in this work by mass spectrometry analysis. Gene ID and product description

details are from _Toxodb.org_ (release 36). The entries shown in bold indicate that the protein

was detect with high confidence from only BNP and IP samples. Asterisk indicates that the

corresponding peptides were detected in only one of the replicate runs for SEC sample. BNP

data is a combination of experiments done with both TgATPβ-YFP-HA and TgATPoscp-

YFP-HA expressing transgenic parasites. SEC and IP data are from _Tgatpβ-YFP-HA_ and

Tgatposcp-YFP-HA transgenic parasites respectively.

Fig 6. HHPred identification of novel F$_O$ subunit a based on conserved structural features.

(A) Representation of the pairwise sequence alignments generated by HHPred [46,47] for the

putative F$_O$ subunit a from _T. gondii_ and four other F$_O$ subunit a proteins for which the

structure is known. The table provides details of the amino acid length and a probability score

for the prediction from the hit alignments. The red lines indicated the 3 transmembrane

domains present in the *T. gondii* protein. (B) Protein sequence alignments and secondary

structure information were made using the Clustal Omega [73] and ESPript [74] softwares,

and only the C-terminal portion of respective proteins is shown. The names of alveolate

species are highlighted in blue. For the non-alveolate species included in the alignment, the

F_O subunit a is either readily identified from sequence (*Scer, Hsap, Atha, Ecol* and *Pten*), or

has been experimentally determined (*Crei* and *Poly*). Positions with similar amino acids are

highlighted in red in the alignment. The arginine and glutamine residues, highlighted in

yellow, are conserved in all species and important for function. The helices (α9, α10 and

α12) shown are from the structure of *Poly* F-type ATP synthase. Species names: *Tgon, T.*

gondii; *Hham, H. hammondi*; *Pfal, P. falciparum*; *Pviv, P. vivax*; *Tann, T. annulata*; *Bbov, B.*

bovis; *Cmur, C. muris*; *Cvel, C. velia*; *Crei, C. reinhardtii*; *Poly, Polytomella sp.*; *Scer, S.*

cerevisiae; *Hsap, H. sapiens*; *Atha, A. thaliana*; *Ecol, E. coli*; *Pden, P. denitrificans*.

Fig 7. Phylogenetic profile of the alveolate infrakingdom for all subunits mapped to *T.*

***gondii* F-type ATP synthase.**

Alveolate clades are highlighted with a grey background and their expected phylogenetic

relationship is indicated by a tree structure above. Grey and white boxes indicate the presence

and absence of the corresponding ortholog, respectively. The hatched boxes represent the

presence of the ortholog in *C. muris* only and absence in *C. parvum* and *C. hominis*. The table

on the left lists the gene ID for all ASAPs, along with the annotation, essentiality phenotypes

(phenotype score from CRISPR/Cas9 knockout [50]), and protein localization experiments

from an independent unpublished study (see text in Materials and Methods for details). The

key for localization annotation is given below the table.

Fig 8. Gene co-expression analysis for *T. gondii* and *P. falciparum* F-type ATP synthase

subunits.

The distribution of co-expression values - as measured by the Pearson correlation coefficients

– are plotted for the three gene pairs categories, as shown for *T. gondii* (A) and *P. falciparum*

(B). The statistical support from Mann-Whitney U p-values and Cohen's d values are shown

in the table within each plot. Blue (1), co-expression correlation between ATP synthase

genes; Orange (2), co-expression correlation between ATP synthase genes and non- ATP

synthase genes; Green (3), co-expression correlation between non- ATP synthase genes.

MWU, Mann-Whitney U p-values; Cohen's d, Cohen's d values.

A

F-type ATP Synthase
(core components)

B

F-type ATP synthase module	Sub-unit	*Pfal*	*Pviv*	*Tpar*	*Bbov*	*Tgon*	*Ncan*	*Cmur*	*Cpar*	*Chom*	*Cvel*	*Pmar*	*Tthe*	*Scer*	*Btau*	*Atha*	*Crei*	*Tbru*
F_1 Sector	α	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	β	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	γ	+	+	+	+	+	+	+	-	-	+	+	+	+	+	+	+	+
	ε	+	+	+	+	+	+	+	-	-	+	+	-	+	+	+	+	+
	δ	+	+	+	+	+	+	+	-	-	+	+	+	+	+	+	+	+
F_O Sector	c	+	+	+	+	+	+	+	-	-	+	+	+	+	+	+	+	+
	OSCP	+	+	+	+	+	+	+	-	-	+	+	+	+	+	+	+	+
	a	-	-	-	-	-	-	-	-	-	-	-	+	+	+	+	+	+
	b	-	-	-	-	-	-	-	-	-	-	-	-	+	+	+	-	+
	d	-	-	-	-	-	-	-	-	-	-	-	+	+	+	+	-	-
	f	-	-	-	-	-	-	-	-	-	-	-	-	+	+	+	-	-
	8/A6L	-	-	-	-	-	-	-	-	-	-	-	-	*	+	+	-	-
	h/f6	-	-	-	-	-	-	-	-	-	-	-	-	+	+	+	-	-
Other associated proteins	g	-	-	-	-	-	-	-	-	-	-	-	-	+	+	+	-	-
	IF1	-	-	-	-	-	-	-	-	-	-	-	-	+	+	+	-	-
	e	-	-	-	-	-	-	-	-	-	-	-	-	+	+	+	-	-
	j	-	-	-	-	-	-	-	-	-	-	-	-	+	-	-	-	-
	k	-	-	-	-	-	-	-	-	-	-	-	-	+	-	-	-	-
	AGP	-	-	-	-	-	-	-	-	-	-	-	-	-	+	-	-	-
	MLQ	-	-	-	-	-	-	-	-	-	-	-	-	-	+	-	-	-
Novel subunits													13				9	14

1000

1001

1002

1003

Salunke et. al., Fig 3

1004

1005

1006

1007

1008

1009

Salunke et. al., Fig 5

A

BNP (96)   SEC (64)

44   24   21

19
(17)

9
(9)   0

IP (29)

B

S. No.	Gene ID	Product description
1	TGME49_204400	F_1alpha subunit
2	TGME49_261950	F_1beta subunit
3	TGME49_226000	F_1delta subunit
4	TGME49_314820	F_1epsilon subunit
5	**TGME49_231910**	**F_1 gamma subunit**
6	TGME49_284540	F_0 OSCP subunit
7	**TGME49_310360**	**hypothetical protein**
8	TGME49_231410	hypothetical protein
9	TGME49_268830	hypothetical protein
10	TGME49_223040	hypothetical protein
11	TGME49_247410	hypothetical protein
12	**TGME49_260180**	**hypothetical protein***
13	**TGME49_218940**	**hypothetical protein**
14	TGME49_282180	hypothetical protein
15	TGME49_285510	hypothetical protein
16	TGME49_215610	hypothetical protein
17	**TGME49_290030**	**hypothetical protein**
18	**TGME49_310180**	**hypothetical protein**
19	**TGME49_214930**	**hypothetical protein**
20	TGME49_245450	hypothetical protein
21	**TGME49_208440**	**hypothetical protein**
22	TGME49_201800	hypothetical protein
23	TGME49_225730	membrane protein
24	TGME49_263080	hypothetical protein
25	TGME49_263990	hypothetical protein
26	**TGME49_270360**	**hypothetical protein**

1010

1011

1012

A

HHPred prediction for *TGME49_310360*

Hit	AA No.	Probability
5T40_K	196..265	64.3
6F36_M	215..308	37.7
5DN6_X	193..276	24.5
6B8H_a	154..243	23.3

B

```
                   α9                    α10
               ooooooooo   oooooooooooooooooooooooooooooooooooo
                 .           .           .           .           .
Tgon  133 FLRPTK----MASFAATFLLNFRFYFMYMARTTFQAVRPLLAFSVFGEVMKLVLATMSSG 188
Hham  133 FLRPAK----MASFAATFLLNFRFYFMYMARTTFQAVRPLLAFSVFGEVMKLVLATMSSG 188
Pfal   88 YVNPLD----MTKFAIKYILSYRFFFIYMARTTFQAVRPLMAFCVFGELMKLILATMTSG 143
Pviv   86 YVNPID----MTKFTIKYILSYRFFFIYMARTTFQAVRPLMAFCVFGELMKLILATMTSG 141
Tann  113 YVNPVS----IVKFTAKYLLSYRFFFIYMARTTFQAARPLLAFCVFGEIMKLVLANLSGG 168
Bbov   82 YVDPLS----MVKFASKYVLSLRFFFIYMARTTFQAVRPLLAFCVFGEIMKLILANISGG 137
Cmur  110 FIYFTD----ILNFTSSIFTNLREYITLLARMSFQAIRPLLAFSVLGEAVKIALALITGS 165
Cvel  304 KFSLNQ----FLSFGYGYVFSLRDIFIFAFRLIFQTGRPMMAVYIGGTLMKTVFSITLSY 359
Crei  217 F--PGHFIPGGTPWPMAFIFVPLETISYTFRAVSLGVRLWVNMLAGHTLLHILTGMALAL 274
Poly  204 F--PGHFIPGGTPWPMAFIFVPLETISYTFRAVSLGVRLWVNMLAGHTLLHILTGMALAL 261
Scer  151 F--FSLFVPAGTPLPLVPLLVIIETLSYFARAISLGLRLGSNILAGHLLMVILAGLTFNF 208
Hsap  124 A--LAHFLPQGTPTPLIPMLVIIETISLLIQPMALAVRLTANITAGHLLMHLIGSATLAM 181
Atha  276 F--FSFLLPAGVPLPLAPFLVLLELISYCFRALSLGIRLFANMMAGHSLVKILSGFAWTM 333
Ecol  173 GFTKELTLQPFNHWAFIPVNLILEGVSLLSKPVSLGLRLFGNMYAGELIFILIAGLLPWW 232
Pden  147 F--LGLFWVSSAPLALRPVLAVIELISYFVRPVSHSIRLAGNIMAGHAVIKVFAAFAAVA 204
                 .           .           :         *         .           .
```

```
                             α12
               oooooooooooooooooooooooooooooooooooooooooooo
                 .           .           .           .           .
Tgon  189 ----------------LFSFLFSFVLAFEVFYFFLQCYISY----TFLTMFFTVLF---- 224
Hham  189 ----------------LFSFLFSFVLAFEVFYFFLQCYISY----TFLTMFFTVLF---- 224
Pfal  143 ----------------VFAFFFSFVLAFEVLYFFLQCYISY----TFLTMFFDVMF---- 179
Pviv  142 ----------------VFAFFFSFVLAFEVLYFFLQCYISY----TFLTMFFDVMF---- 177
Tann  168 ----------------VPAYLFSFVLAFEVFYFFLQLYISY----TFLMMFFTVMF---- 204
Bbov  138 ----------------VPAFLFSFVLAFEVLYFFLQCYISY----TFLTMFFTVMF---- 173
Cmur  166 ----------------SIAFIFSFILAFEVFYFFLQCFISF----VFLSMFFKAIV---- 201
Cvel  360 ----------------SFLLFFSLIAGFEVFYFGLQCYIGF----VFLSFFSEGALM--- 396
Crei  275 P--FSLSFFAMVPATFAVACLLSALVGLEYLVAVLQSGVFSILSTVYVGEFNSVKLAGPL 332
Poly  262 P--FSLGFFSMVPATFGVCCLLSALVGLEYLVAVLQSGVFSILSTVYVGEFNHDKFIGPA 319
Scer  209 M--LINL-FTLVF-GFVPLAMILAIMMLEFAIGIIQGYVWAILTASYLKDAVYLH----- 259
Hsap  182 S--TINLPSTL-----IIFTILILLTILEIAVALIQAYVFTLLVSLYLHDNT-------- 226
Atha  334 L--CMNDIFYFIG-ALGPLFIVLALTGLELGVAILQAYVFTILICIYLNDAINLH----- 385
Ecol  233 SQWILNV-------------PWAIFHILIITLQAFIFMVLTIVYLSMASEEH----- 271
Pden  205 A--I----------APVSVVAITAMYGLEVLVCLIQAYVFTILTCVYLKDALHPAH---- 248
                          :.        :*   :         ::
```

1013

1014

1015

Salunke et. al., Fig 7

Gene ID	Subunit Annotation	CRISPR Pheno Score*	Mito. location	Haemosporida	Piroplasmida	Coccidia	Cryptosporidiidae	Chromerida	Dinoflagellata	Ciliophora	Out groups
TGME49_204400	F_1 alpha	-3.84	Mito#								
TGME49_261950	F_1 beta	-4.84	**Mito#**								
TGME49_226000	F_1 delta	-4.57	Mito#								
TGME49_314820	F_1 epsilon	-3.21	Mito#								
TGME49_231910	F_1 gamma	-3.94	Mito#								
TGME49_249720	F_0 c	-2.98	Mito								
TGME49_284540	OSCP	-3.94	**Mito#**								
TGME49_310360	ASAP-1 (F_0 a?)	-4.49	Mito								
TGME49_231410	ASAP-2 (F_0 b?)	-5.37	Mito*#								
TGME49_268830	ASAP-3 (F_0 d?)	-3.10	Mito#								
TGME49_223040	ASAP-4	-4.49	**N#**								
TGME49_247410	ASAP-5	-3.90	N*#								
TGME49_260180	ASAP-6	-4.07	Mito*#								
TGME49_218940	ASAP-7	-3.92	N*#								
TGME49_282180	ASAP-8	-2.86	**Mito#**								
TGME49_285510	ASAP-9	-1.87	N#								
TGME49_215610	ASAP-10	-3.55	Mito*#								
TGME49_290030	ASAP-11	-3.88	**Mito#**								
TGME49_310180	ASAP-12	-3.40	N#								
TGME49_214930	ASAP-13	-1.37	N#								
TGME49_245450	ASAP-14	-2.95	**Mito#**								
TGME49_208440	ASAP-15	-3.54	Mito#								
TGME49_201800	ASAP-16	-4.01	Mito#								
TGME49_225730	ASAP-17	-3.65	Mito#								
TGME49_263080	ASAP-18	-3.10	Mito#								
TGME49_263990	ASAP-19	0.22	N#								
TGME49_270360	ASAP-20	0.32	N#								

Key for location data
Mito = mitochondria localization signal present
N = no mitochondria localization signal
Bold = Localized in this study
* = Localized previously [55]
\# = unpublished evidence from Hyper-LOPIT experiment

1016

1017

1018

A

B

1019

1020

1021