Metastable XBP1u transmembrane domain mediates insertion into the ER membrane and intramembrane proteolysis by the signal peptide peptidase

Sara Suna Yücel[1,4], Walter Stelzer[2,4], Alessandra Lorenzoni[1], Manfred Wozny[3], Dieter Langosch[2,*], Marius K. Lemberg[1,*]

[1]Centre for Molecular Biology of Heidelberg University (ZMBH), DKFZ-ZMBH Alliance, Im Neuenheimer Feld 282, 69120 Heidelberg, Germany.

[2]Center for Integrated Protein Science Munich (CIPSM) at the Lehrstuhl Chemie der Biopolymere, Technical University of Munich, Weihenstephaner Berg 3, 85354 Freising, Germany.

[3]MassMap GmbH & Co. KG, Meichelbeckstr. 13a, 85356 Freising, Germany.

[4]These authors contributed equally to this work.

*Corresponding authors: m.lemberg@zmbh.uni-heidelberg.de; langosch@tum.de

Abstract

The unspliced *XBP1* mRNA encodes the transcriptionally inert variant of the unfolded protein response (UPR) transcription factor XBP1u. Besides acting as a negative regulator for UPR, XBP1u targets its mRNA-ribosome-nascent-chain-complex to the endoplasmic reticulum (ER) facilitating IRE1-mediated splicing. Yet, its membrane association is controversial. Here, we use cell-free translocation assays and living cells to define a moderately hydrophobic stretch in XBP1u that is sufficient for insertion into the ER membrane. Mutagenesis of this transmembrane (TM) region reveals residues that determine XBP1u for an ER-associated degradation route centered around the signal peptide peptidase (SPP). Furthermore, the impact of these mutations on TM helix dynamics was assessed by recording residue-specific amide exchange kinetics, where data was evaluated by a novel semi-automated method. As compared to the previously employed procedure, this method reduces the time needed for data processing by approximately an order of magnitude. Based on our results, we suggest that SPP-catalyzed intramembrane proteolysis of TM helices is not only determined by its conformational flexibility, but also by side chain interactions near the cleavage site with the enzyme's active site.

Introduction

The endoplasmic reticulum (ER) controls synthesis, folding and modification of almost one third of our proteome. Under changing physiological circumstances or proteotoxic stress, the folding capacity of the ER may face an imbalance that can lead to an accumulation of misfolded proteins. In order to keep the ER functional and restore ER homeostasis, ER stress activates an intracellular signaling pathway called the unfolded protein response (UPR) that among other factors transcriptionally upregulates folding chaperones and the ER-associated degradation (ERAD) machinery (Walter and Ron, 2011). Subsequently, ER folding capacity increases and terminally misfolded protein species are targeted by a diverse set of ERAD factors for proteasomal degradation (Christianson and Ye, 2014).

Chronic ER stress leads to UPR induced apoptosis, which emerges as a driver of several human diseases including diabetes and neurological disorders (Hetz and Papa, 2018).

In metazoans, three UPR signaling branches act in parallel, namely regulated intramembrane proteolysis of ATF6, translational control by the PERK kinase and an IRE1-mediated unconventional splicing of the *XBP1* mRNA (Walter and Ron, 2011). IRE1 is a membrane integral stress sensor that upon recognition of bulk amount of unfolded proteins in the ER lumen oligomerizes and activates its endonuclease activity (Li et al., 2010; Yoshida et al., 2001). Consequently, the *XBP1* mRNA is spliced (removing a 26-nt intron) leading to an open reading frame that encodes for the mature transcription factor XBP1s comprising an N-terminal basic leucine zipper (bZIP) dimerization domain followed by a C-terminal transcription activation domain. In contrast, the unspliced variant of *XBP1* mRNA encodes XBP1u, which consists of

a hydrophobic stretch in the C-terminal portion and lacks the transcription activation domain. Although short lived, XBP1u acts as a negative regulator of the UPR by targeting XBP1s and activated ATF6 for degradation and thereby fine-tuning the UPR (Tirosh et al., 2006; Yoshida et al., 2006; Yoshida et al., 2009). Moreover, XBP1u mediates a crosstalk with other signaling networks (Zhao et al., 2013). Interestingly, newly synthesized XBP1u targets its own mRNA-ribosome-nascent chain complex to the ER in order to facilitate its splicing by IRE1 (Yanagitani et al., 2009; Yanagitani et al., 2011). When emerging from the translating ribosome, the C-terminal hydrophobic stretch of XBP1u is recognized by the signal recognition particle (SRP) resulting in co-translational targeting to the Sec61 translocon (Kanda et al., 2016; Plumb et al., 2015). Different to insertion of canonical nascent signal peptides and signal anchor sequences (Mothes et al., 1994; Voorhees and Hegde, 2016), Kohno and co-workers suggest that XBP1u interacts with Sec61 and the ER membrane on the cytosolic side from where it may detach to reach the nucleus in an unprocessed form (Kanda et al., 2016; Yanagitani et al., 2009). Contrary, our previous research indicates that the XBP1u hydrophobic region is capable to insert into the ER membrane as a type II transmembrane (TM) domain, facing the ER lumen with its C-terminus (Chen et al., 2014). Furthermore, we revealed that XBP1u undergoes proteolytic cleavage by the signal peptide peptidase (SPP) that funnels it into the ERAD pathway (Chen et al., 2014). SPP belongs to a special group of enzymes, so-called intramembrane proteases, that cleaves TM anchors of a wide range of proteins to trigger their release from cellular membranes (Lemberg, 2011). Although mechanistically quite diverse, comprising aspartic proteases such as SPP and γ-secretase, rhomboid serine proteases, S2P metalloproteases and the glutamyl protease Rce1, all these proteases form membrane-embedded aqueous active sites and face

similar problems in substrate recognition, unwinding of substrate TM helices and cleavage (Langosch et al., 2015; Strisovsky, 2015; Wolfe, 2009).

While SPP was initially identified as the protease clearing signal peptides from the ER membrane (Lemberg and Martoglio, 2002; Weihofen et al., 2002), in recent years the repertoire of SPP substrates has been expanding as we and others revealed that SPP also cleaves certain *bona fide* TM domains (Boname et al., 2014; Chen et al., 2014; Hsu et al., 2015). This indicates that there are mechanistic parallels to SPP-like (SPPL) proteases, which cleave type II membrane proteins in the Golgi and the late secretory pathway (for review see Mentrup et al., 2017). However, to target ERAD substrates for degradation, SPP functionally interacts with the rhomboid pseudoprotease Derlin1 and the E3 ubiquitin ligase TRC8 forming a 500-kDa complex (Chen et al., 2014). While TRC8 ubiquitinates XBP1u to ensure its efficient turnover by the proteasome (Boname et al., 2014; Chen et al., 2014), Derlin1 acts as a substrate receptor for the luminal domain of XBP1u (Chen et al., 2014), obviating the need for a short tail required for SPP-catalyzed cleavage of signal peptides (Lemberg and Martoglio, 2002). Although the list of substrates and the physiological significance of SPP in the cell is expanding, its substrate recognition mechanism and cleavage determinants are still ill defined. Our previous analysis of signal peptides revealed that amino acid residues predicted to lower TM helix stability are required for efficient SPP-catalyzed intramembrane cleavage (Lemberg and Martoglio, 2002). However, it is not yet clear whether SPP recognizes a certain consensus sequence as has been described for bacterial rhomboid intramembrane serine proteases (Strisovsky et al., 2009). Here, we elucidate the recognition of XBP1u by the Sec61 translocon and show that the moderately hydrophobic segment of XBP1u is sufficient for

membrane insertion. However, different to most other TM segments, XBP1u is

rapidly processed by the intramembrane protease SPP defining it as a metastable TM

anchor. Moreover, a combination of *in vitro* translocation and cleavage assays, cell-

based assays and deuterium/hydrogen exchange (DHX) experiments broaden our

understanding of the effects of conformational dynamics in the TM helix and the

influence on substrate recognition and cleavage by SPP. Most significantly, we find

that conformational flexibility in the N-terminal portion of the TM segments is a

prerequisite, but SPP-catalyzed cleavage of *bona fide* TM helices is governed by

specific side chain interactions.

Results

In order to characterize the efficiency of membrane insertion of XBP1u and to define

requirements for its cleavage by SPP, we examined i) the role of its hydrophobic

region in a microsome translocation assay and cells and ii) correlated the impact of

various mutations within this hydrophobic domain on helix dynamics and

cleavability.

Functional interaction of XBP1u with the Sec61 translocon

Previously, ER targeting of XBP1u has been described to be in an SRP-dependent

pathway (Kanda et al., 2016). However, to which extend it inserts into the Sec61

translocon and whether it adopts a type II membrane anchor is controversially debated

(Chen et al., 2014; Kanda et al., 2016; Plumb et al., 2015). One attempt to rationalize

membrane insertion efficiency of TM segments is based on a thermodynamic scale

that had been experimentally determined by von Heijne and colleagues using a

glycosylation reporter protein in an *in vitro* translocation assay (Hessa et al., 2007).

Based on its apparent free energy (ΔG_{app}), the hydrophobic region of XBP1u is not

predicted to form a stable TM segment, when compared to the thermodynamically

favorable insertion of the *bona fide* TM helix of calnexin, to give an example (Figure

1A). However, by using this prediction algorithm, signal sequences also have

frequently positive, 'forbidden' values (see Figure 1A for examples) and were

excluded in the analysis by von Heijne (Hessa et al., 2007). Nonetheless, it is well

accepted that the Sec61 translocon mediates efficient insertion of signal peptides of

nascent peptide chains (Mothes et al., 1994; Voorhees and Hegde, 2016), indicating

that the translocon can also insert certain helices of low hydrophobicity into the

bilayer.

We therefore asked, whether the putative TM region of XBP1u interacts with the

Sec61 translocon in the manner of a nascent signal peptide. Although variable in

amino acid sequence, signal peptides have a common three-domain structure

comprised by a short positive-charged 'n-region', a hydrophobic 'h-region' of 7 to 15

residues and a polar 'c-region' including the site for signal peptidase cleavage

(Haeuptle et al., 1989). While commonly positioned at the N-terminus, there are also

internal signal sequences with n-regions in the size range of XBP1u's bZIP reported

(Kapp et al., 2009). With a low scoring signal peptidase cleavage site predicted by

SignalP 3.0 (Bendtsen et al., 2004) in the putative c-region between alanine-206 and

phenylalanine-207 (Figure 1B), the XBP1u hydrophobic region shows certain

characteristics of an internal signal sequence. Hence, we created a chimeric protein in

which we exchanged the signal sequence of pre-prolactin (pPrl) for this predicted

XBP1u TM domain and few N-terminal flanking residues (XBP1u-TMD-Prl) and

tested whether this XBP1u neo-signal peptide is sufficient to mediate ER targeting and translocation in a cell-free system (Figure 1B). To this end, mRNA either encoding pPrl wild-type (wt) or the XBP1u-TMD-Prl fusion were translated in reticulocyte lysate supplemented with [^{35}S]-labelled methionine and cysteine. Upon addition of ER-derived canine rough microsomes, signal sequences are inserted into the membrane leading to signal peptidase cleavage and liberation of the translocated polypeptide (Blobel and Dobberstein, 1975). Indeed, a faster migrating band corresponding to mature Prl was observed for both constructs. This product was resistant to proteinase K but was degraded when detergent was added in addition to solubilize the microsomal membrane (Figure 1B). Successful translocation indicates that in the context of the isolated XBP1u TM domain, signal peptidase cleaves the cryptic signal peptidase site at alanine-206. Moreover, these results show that the XBP1u TM domain alone is sufficient for productive interaction with Sec61 and subsequent translocation of a downstream fusion protein.

Significant fraction of XBP1u adopts membrane-spanning topology
Earlier *in vitro* studies of XBP1u targeting lead to the conclusion that its signal sequence has evolved to functionally interact with SRP but avoids efficient insertion into the membrane leading to a predominantly peripherally attached protein (Kanda et al., 2016; Plumb et al., 2015). Contrary, our previous topology analysis including *in vitro* and *in cellulo* assays has lead us to conclude that XBP1u inserts as a type II membrane protein that subsequently becomes turned over by an SPP-dependent ERAD pathway (Chen et al., 2014). One way to interfere with SPP-mediated degradation is to exchange hydrophilic TM residues by leucine residues (Lemberg and Martoglio, 2002). Earlier, we showed that exchanges to leucine (referred as mt1

and mt2, see Figure 1A) block SPP-catalyzed cleavage of XBP1u (Chen et al., 2014).

These mutations are predicted to favor insertion as a TM helix since an increase in

hydrophobicity increases the gain in the free energy ΔG_{app} upon Sec61 insertion

(Figure 1A), posing the question whether increased stability is primarily a

consequence of a changed biosynthesis pathway. Therefore, we tested these stable

XBP1u mutants in our cell free translocation assay (Figure 1B) and by a previously

used cell-based XBP1u-R232N glycosylation reporter construct (Figure 1C) (Chen et

al., 2014). While mutating the N-terminal portion of the XBP1u TM domain (mt1)

promotes efficient translocation of the Prl fusion protein (Figure 1B), exchange of a

C-terminal XBP1u TM motif (mt2) was not functional in this context (result not

shown), indicating that canonical N-terminal signal sequences and XBP1u follow

related but not identical targeting pathways. Despite that caveat, in the context of the

full-length XBP1u-R232N, both TM domain variants promoted efficient insertion as a

glycosylated type II membrane protein. When compared to XBP1u wt TM domain,

the steady state level of XBP1u and subsequently the glycosylation efficiency of both

TM mutants (mt1 and mt2) increases from approx. 50% to >80%. We speculate that

this reduced steady-state level of XBP1u wt may be a consequence of both, the pre-

emptive degradation of XBP1u that failed ER-targeting and insertion (Kang et al.,

2006) and of SPP-mediated turnover of the type II inserted pool (Chen et al., 2014).

However, a combination of our *in vitro* and cell-based assays demonstrates that

XBP1u wt inserts to a significant fraction (>50%) as a type II membrane protein and

differences by mutating the TM domain are primarily caused by blocking SPP-

triggered ERAD. This does not rule out that a certain fraction fails targeting or is

stalled as an insertion intermediate by the translation pause site (Yanagitani et al.,

2011).

Turnover of XBP1u is governed by its TM domain

Our previous analysis of XBP1u degradation revealed that the C-terminal luminal domain is recognized by the ERAD factor Derlin-1 priming it for the SPP-turnover without the need for a short tail (Chen et al., 2014). A short tail is otherwise required for SPP-catalyzed cleavage of signal peptides (Lemberg and Martoglio, 2002). Although we had reported that deleting the C-terminal portion drastically accelerates turnover and omits any requirements for Derlin-1, the TM determinants for SPP-catalyzed XBP1u cleavage had not been investigated in the isolated situation. We therefore created tail-anchored (TA) versions of the TM domain mutants by deleting the C-terminal 54 amino acids of XBP1u. To confirm ER targeting, transfected Hek293T cells were fixed and analyzed by immunofluorescence microscopy (Figure 2A). Next, XBP1u-TA mutants were tested for their degradation rate by cycloheximide chase experiments. Consistent with our previous report, the half-life of XBP1u-TA wt diminished to approx. 15 min compared to 60 min of the full-length XBP1u construct (Figure 2B). Contrary, XBP1u-TA mt1 and mt2 both were stable and no decay was observed for the entire chase (Figure 2B). Consistent results were obtained in a microsome-based *in vitro* assay using mt1 of the XBP1u-TMD-Prl fusion protein (Figure 2C). To test for signal peptide processing, we isolated microsomes from a translation/translocation assay using pPrl and XBP1u-TMD-Prl mRNA, respectively. Upon addition of microsomes, both signal sequences were cleaved by signal peptidase and the liberated signal peptides are anchored in the microsomal membrane similar to a membrane protein with a single TM region until they are released by SPP-catalyzed intramembrane proteolysis (Lemberg et al., 2001; Weihofen et al., 2000). To quantify this release, we isolated microsomes from a

reaction in presence of the SPP inhibitor (Z-LL)$_2$-ketone (Figure 2C) (Weihofen et al., 2000). As had been observed for pPrl, also for the liberated XBP1u neo-signal peptide a weak band in the low molecular range emerged that is stabilized upon treatment with (Z-LL)$_2$-ketone (Figure 2C). This product co-migrated with an *in vitro* translated reference peptide comprising the XBP1u neo-signal peptide. The corresponding signal peptide derived from XBP1u-TMD mt1, however, was only processed by approx. 30% compared to >60% for the XBP1u wt TM domain sequence (Figure 2C). This reduction of SPP processing efficiency is in the range previously observed for signal peptides lacking obvious helix-destabilizing residues (Lemberg and Martoglio, 2002). Overall, these results confirm that the XBP1u TM segments harbors features for SPP-catalyzed intramembrane proteolysis.

Increasing rigidity of TM helix prevents SPP-catalyzed cleavage
Next, we investigated the effects of XBP1u mutations on the helicity and the conformational flexibility of the TM domain by amide DHX kinetics. To this end, we created synthetic peptides and dissolved them in aqueous trifluoroethanol (TFE) that serves as a mimic of the aqueous environment within the catalytic cleft of an intramembrane protease and readily solubilizes the peptides (Pester et al., 2013). CD spectroscopy confirmed that the peptides form >60% helix in this solvent, with a small but significant increase in the helicity of mt1 (Figure S1A), as shown before (Chen et al., 2014). This ensures that the following measurements predominantly describe the properties of the helical state. The overall conformational flexibility of exhaustively (> 95%) deuterated peptides was assessed in the same solvent by recording DHX kinetics that reflect the stability of intrahelical amide hydrogen-bonds (H-bonds). At time point zero, where the labile deuterons bound to non-H-bonded

polar atoms have already exchanged for protons, ~22 deuterons remain (Figure 2D).

These remaining deuterons represent >90% of those 24 backbone amides that can at

least be partially protected from DHX by intrahelical H-bond formation. As

exemplified by the wt spectra shown in Figure S1B, a gradual shift in addition to the

lack of a bimodal shape of the isotopic distributions is diagnostic of transient local

unfolding events described by a kinetic model, known as EX2 kinetics, where

individual deuterons exchange in an uncorrelated fashion (Konermann et al., 2011).

Next, we investigated the leucine substitutions in the cleavage-deficient mt1 and mt2.

Leucine substitutions are predicted to stabilize intra-helical contacts between their

large and flexible side chains and the side chains of other residues within the helix

(Quint et al., 2010). Consistent with this, amide exchange of mt1 and mt2 is slower

than that of wt, indicating their lower average helix backbone flexibility (Figure 2D).

This refines our previous structural analysis solely relying on CD spectroscopy (Chen

et al., 2014) in revealing that mutagenizing either N-terminal or the C-terminal TM

residues to leucine render the TM helix more rigid.

Increasing TM helix flexibility has differential effects on cleavage

Results described above provide further evidence that XBP1u spans the ER

membrane with a metastable type II TM anchor that is rapidly processed by SPP to

target it for proteasomal degradation (Chen et al., 2014). Next, we wanted to

understand how increased flexibility of the TM helix modulates its cleavage by SPP.

To this end, we inserted a di-glycine motif in the N-terminal (mt3), central (mt4) or

C-terminal (mt5) portion of the TM domain (Figure 3A). Due to the absence of a side

chain, glycine destabilizes a TM helix by introducing a packing defect (Hogel et al.,

2018).

First, we assessed the topology of the di-glycine mutants via the XBP1u-R232N

glycosylation reporter. We observed that mt3 and mt5 are efficiently targeted to the

ER and adopt a type II topology, whereas only a neglectable fraction was

glycosylated in the case of mt4 (Figure 3B). Consistent with this,

immunofluorescence analysis revealed ER localization for mt3 and mt5, while for

mt4 only a minor pool co-localized with the ER marker and the predominant fraction

was detected in the nucleus (Figure S2A). Therefore, introducing the di-glycine motif

into the center of the TM helix has the strongest effect on membrane insertion, as also

reflected by a positive ΔG_{app} (Table S1), thus limiting the biological relevance of mt4.

Next, we determined the stability of these mutants in transfected Hek293T cells by

adding cycloheximide to block synthesis of new proteins and chase the pre-existing

fraction of XBP1u over time. Interestingly, mt5 was resistant to degradation whereas

mt3 showed degradation rates similar to XBP1u wt (Figure 3C). Likewise, the ER

targeting-deficient mt4 showed not significant stabilization (Figure S2B), indicating

that cytoplasmic/nucleoplasmic degradation pathways can also efficiently break down

XBP1u. On the other hand, mt3, which is ER resident (Figure S2A) and adopts a type

II topology (Figure 3B), is turned over at the same rate as the wt construct (Figure

3B). Its degradation is sensitive to (Z-LL)$_2$-ketone (Figure S2C), indicating that, like

XBP1u wt, mt3 is a substrate for SPP-dependent ERAD. Contrary, mt5 accumulates

in the ER and only fades away by a low-level, (Z-LL)$_2$-ketone-insensitive, alternative

degradation route (Figure S2A and C). Similarly, we observed inefficient SPP

processing of the mt5 neo-signal peptide in the *in vitro* translocation assay (Figure

3D). This corroborates that the C-terminal di-glycine motif prevents cleavage by SPP.

In order to detect a potential correlation between cleavability and helix flexibility, we determined amide exchange kinetics. As predicted, mt3, mt4 and mt5 exhibit faster overall DHX kinetics (Figure 4A). CD spectroscopy showed a similar overall helicity of the wt and these mutants (Figure S3A), demonstrating that DHX is a much more sensitive tool to detect changes in helix flexibility (Figure 4A). A close inspection of the kinetics reveals the following rank order of the glycine impact: mt4 > mt5 > mt3 (Figure 4A, inset); this rank order reflects the relative importance of the mutated positions for overall helix flexibility. To refine our analysis, we sought to correlate cleavability of our constructs to TM flexibility that was resolved at the single-residue level. To this end, we performed electron transfer dissociation (ETD) of our peptides in the gas phase after different periods of DHX. Using a novel software solution that accelerates ETD fragment analyses by an order of magnitude, we determined the DHX kinetics of individual backbone amides that is quantitated by the exchange rate constants k_{exp} shown in Figure S3B and Table S2. While only incomplete kinetics could be determined for several N- and C-terminal residues due to exchange near the helix termini, that is too rapid for measurement, the DHX kinetics of amides within the core of the XBP1u TM helix from isoleucine-186 to phenylalanine-207 is covered by sufficient numbers of data points for all investigated constructs (Figure S3C). At all these residues, the k_{exp} values are below the respective calculated chemical amide exchange rate constants k_{ch}, which reflects their participation in secondary structure, *viz.* helix. Exchange rate constants were then converted to the free energy changes associated with H-bond formation ΔG (Table S2). Figure 4B shows the distribution of ΔG values for the wt TM helix that increase from ~2 kcal/mol near the helix N-terminus to ~4.5 kcal/mol. These values are plausible, given that values of ~1 kcal/mol were previously reported for intrahelical H-bonds in water, which can

effectively compete with intrahelical H-bonding, ~4 kcal/mol for H-bonds of model

compounds in apolar solvent, and 6.6 kcal/mol for H-bonds in vacuo (Bowie, 2011).

To quantitate the effects of the mutations, we calculated the differences in H-bond

stability $\Delta\Delta G$ by subtracting wt ΔG values from the respective mutant values. The

results (Figure 4C) reveal that mutations to leucine in mt1 and mt2 stabilize local

amide H-bonds by up to ~3 kcal/mol; in case of mt1, stabilization extends to several

residues downstream of the mutations. By contrast, the di-glycine mutations in mt3

and mt5 destabilize by up to ~1 kcal/mol (Figure 4C). For mt4, which fails efficient

ER targeting, the extent of destabiliziation is much larger with up to ~4 kcal/mol

around that central di-glycine motif (Figure S3D). Taken together, these results show

that introducing leucine or glycine residues into the XBP1u TM domain has profound

effects on TM helix flexibility, but there is no clear correlation to cleavability by SPP.

Contrary to the initial proposal that primarily TM helix flexibility determines the

efficiency of SPP cleavage (Lemberg and Martoglio, 2004), mutations destabilizing

the TM domain (mt5) can also reduce SPP processing efficiency.

Single residues can affect cleavage

Since the di-glycine motif in XBP1u mt5 prohibits SPP-dependent degradation,

cleavage by SPP may not only depend on TM helix dynamics but also on substrate

recognition requiring the mutated residues. In order to study the importance of the

glutamine-199 and serine-200 that were simultaneously mutated in mt5, we generated

two single glycine mutants, namely mt6 (Q199G) and mt7 (S200G) (Figure 5A). We

confirmed ER targeting and topology of these single TM domain mutants by the

glycosylation reporter assay (Figure 5B) and tested their degradation rate by

cycloheximide chase experiments (Figure 5C). Interestingly, mt7 (S200G) is

stabilized similar as we see for mt5, while mt6 showed a degradation rate indistinguishable from XBP1u wt, suggesting that SPP recognizes the side chain of serine-200. We noted however, that several well-characterized SPP substrates such as the signal peptides of calreticulin or insulin do not have a serine in the appropriate position (Kronenberg-Versteeg et al., 2018; Lemberg and Martoglio, 2002). Despite that, we wondered whether this feature is important to foster processing of actual TM helices. Based on a recent proteomics screen, the TA protein heme oxygenase 1 (HO1) has been shown to be a native substrate for the SPP-dependent ERAD pathway (Boname et al., 2014), but its substrate determinants have not been analyzed yet. Interestingly, HO1 has a serine in the C-terminal portion of its TM domain (Figure 6A). To test the influence of such a putative substrate determinant, we mutated serine-275 to leucine, alanine or glycine, respectively and tested the stability of HO1 by cycloheximide chase (Figure 6A, B and S4). Moreover, we mutated two putative helix-destabilizing residues in the N-terminal TM domain portion, namely proline-266 and arginine-269 (Figure 6A). Interestingly, all mutants were significantly stabilized, suggesting that they are not cleaved by SPP (Figure 6B and S4). This indicates that like for XBP1u, dynamics of the N-terminal TM helix portion and a serine in the C-terminal half play a role in cleavage by SPP. For the HO1-S275A mutant a putative SPP-generated N-terminal cleavage fragment has been observed to localize to the nucleus in transfected HeLa cells (Hsu et al., 2015). Thus, we next asked whether these mutations also prevent generation of cleavage products. Due to a rapid proteasomal turnover, analyzing ectopically expressed HO1 we do not observe cleavage fragments generated by endogenous SPP (Boname et al., 2014). Because of tight coupling to the ERAD machinery, fragments are commonly rapidly degraded by the proteasome (Boname et al., 2014; Chen et al., 2014). However, for XBP1u we

previously observed that blocking the proteasome also suppressed SPP activity and only low amount of XBP1u cleavage fragments could be detected (Chen et al., 2014). The HO1 cleavage fragment, however, escapes to some extend complete proteasomal destruction (Hsu et al., 2015). To detect HO1 cleavage fragments without inhibition of the proteasome, we ectopically expressed increasing levels of SPP (Figure 6C). Next, we compared processing of HO1 wt with our mutant constructs revealing processing only at the highest SPP levels (Figure 6C). Although these results show that the serine is not a strict requirement for SPP-catalyzed processing when overexpressing SPP. However, taken together with the cycloheximide chase experiments, reduced processing efficiency in the presence of endogenous protease levels strongly indicate that SPP makes side chain-specific interactions to its substrates. We therefore set out to determine the site of SPP-catalyzed cleavage. Previous analysis of SPP *in vitro* and tissue culture cells point towards cleavage in the middle of the TM span (Chen et al., 2014; Hussy et al., 1996; Lemberg and Martoglio, 2002; Weihofen et al., 2000). Because of tight coupling to the ERAD machinery with SPP-catalyzed XBP1u cleavage we failed to isolate sufficient XBP1u cleavage fragment for cleavage site determination (results not shown). HO1, however, escapes to some extend complete proteasomal destruction and a much less "backlash" of proteasome inhibitors on SPP-trigged release has been observed (Boname et al., 2014; Hsu et al., 2015) making it the ideal substrate for cleavage site determination by MS analysis. To allow isolation of a short peptide comprising the carboxy-terminus generated by SPP, we co-expressed SPP with an HO1 TM domain GFP-fusion construct including an engineered TEV protease cleavage site followed by a FLAG-tag (Figure 6D). Subsequently, we immunoisolated the SPP-generated cleavage fragment from the cytosolic fraction with an GFP antibody and then digested it on the

beads with TEV protease. This leads to the release of 5 kDa-peptides having the FLAG-tag at their new N-terminus and C-terminally ending at the peptide bond where SPP cleavage occurred. For cleavage site determination, this peptide was immunoprecipitated and analyzed by MS. A major peak was observed with 5274.76 Da, which corresponds to cleavage after threonine-273 (Figure 6E). In addition, three less intense peaks were observed corresponding to peptides further shortened by one amino acid, respectively. These alternate C-termini may either originate from SPP cleavage at alternative sites or originate from trimming by cytoplasmic carboxypeptidases. However, we did not detect any cleavage fragment matching the previously identified tryptic HO1 TM domain peptides of 6 to 7 residue length that indicated cleavage after serine-275 (Hsu et al., 2015). The predominant fragment observed by our analysis of 5-kDa peptide generated by a tailored TEV protease site, which is predicted to be more accurate, strongly indicates that HO1 is predominantly cleaved at threonine-273. Taken together with the requirement for serine-275, this suggest that SPP shows a so far unanticipated substrate specificity in the P2' position.

Discussion

Here, we revisited XBP1u ER targeting and insertion. We show that the XBP1u hydrophobic region can function as a signal peptide and thus productively interacts with Sec61 leading to the subsequent translocation of a downstream protein domain. While previous work pointed towards a peripheral attachment of XBP1u to ER-derived microsomes (Kanda et al., 2016; Plumb et al., 2015), by using a cell-based glycosylation reporter construct we further show that at least half of the protein inserts as a type II membrane protein. Since glycosylation close to the C-terminal tail is

commonly not complete, the insertion rate of XBP1u as a type II membrane protein

may exceed 50%, thus corroborating results from our previous topology analysis

(Chen et al., 2014). By cell-based degradation assays, we show that membrane

integration of the XBP1u TM domain is metastable, prompting efficient recognition

by SPP and intramembrane cleavage. This is consistent with our previous analysis,

demonstrating that upon inhibition of SPP also endogenous XBP1u accumulates in

the ER (Chen et al., 2014). Interestingly, introducing a di-glycine motif in the C-

terminal portion of the TM domain of XBP1u does not interfere with ER targeting but

reduces SPP-triggered turnover. The kinetics of amide exchange reveals the expected

local increase of the TM helix flexibility. Site-specific mutagenesis of XBP1u and the

TA protein HO1, which is also a substrate for SPP (Boname et al., 2014), shows that a

single serine residue is required for efficient SPP-catalyzed cleavage. Over all, our

results suggest that like rhomboid intramembrane serine proteases (Strisovsky et al.,

2009) also SPP may recognize defined substrate sequence features.

Polar TM residues introduce functionality but affect ER targeting and helix stability

TM segments are 18 to 25 amino acid long α-helical peptides with a high content of

hydrophobic residues. Their partitioning into the lipid bilayer of a cellular membrane

via lateral exit from the Sec61 channel is thought to be driven by the hydrophobic

effect (Cymer et al., 2014). While N-terminal and internal signal sequences are

commonly targeted to Sec61 via SRP, thereby avoiding exposure of their hydrophobic

domain to the aqueous environment, TA proteins are targeted by the machinery of the

GET pathway (Hegde and Keenan, 2011). The thermodynamics of Sec61-mediated

insertion of TM segments is described by ΔG_{app} determined by von Heijne and

colleagues from measurements in a microsome-based system (Hessa et al., 2007).

This scale, however, reflects the situation of internal TM segments, rather than that of N-terminal signal sequences (Hessa et al., 2007). Here, we show that XBP1u's TM domain is recognized by the Sec61 translocon leading to successful translocation and glycosylation of the C-terminus of ~50% of the present protein, which is consistent with a predicted ΔG_{app} being close to zero (Figure 1A). Certain XBP1u mutants fail targeting. When fused to Prl as an N-terminal signal sequence, the XBP1u TM domain mt2, a mutationally rigidified TM helix variant, does not successfully insert. Consistent with this, N-terminal signal sequences frequently contain a helix-break-helix structure that is thought to favor loop-like insertion of the targeted nascent chain (van Klompenburg and de Kruijff, 1998). Contrary, mt4, where a di-glycine motif in the center of the TM domain region introduces helix flexibility, fails targeting. This is probably due to a failed interaction with SRP and/or ER insertion resulting from a positive ΔG_{app}. These results show that for signal anchor sequences there is a fine balance between the number of hydrophilic residues that are tolerated and requirements for ER targeting and insertion. Despite that, a plethora of important physiological functions is determined by polar amino acid residues in TM spans. While we may not know the full set of XBP1u molecular properties and functions, here we show that its TM segments serves as a membrane-integral degradation signal that drives its turnover by an SPP-dependent ERAD pathway. Analogously, charged TM residues are recognized as so-called "degrons" by the ERAD machinery to target orphan receptor subunits for degradation (Bonifacino et al., 1990; Fleig et al., 2012).

SPP-catalyzed cleavage is governed by conformational control and site-specific recognition

Since the first discovery of intramembrane proteolysis more than 20 years ago, an enigmatic question remaining is how a lipid-embedded helical substrate is specifically recognized by a membrane-integral protease for cleavage (Brown et al., 2000; Langosch et al., 2015; Strisovsky, 2015; Wolfe, 2009). Because for several intramembrane protease substrates, a strict requirement for amino acid residues with a low predicted TM helix preference has been observed (Chen et al., 2014; Lemberg and Martoglio, 2002; Moin and Urban, 2012; Urban and Freeman, 2003; Ye et al., 2000a), it is commonly believed that the translocation of the substrate TM helix from a peripheral exosite towards the protease active site is a rate-limiting step requiring a flexible helix (Lemberg and Martoglio, 2004). For example, TM domain mutations in the amyloid precursor protein, a substrate of γ-secretase, can lead to Alzheimer's disease and have an influence on processing. Such mutations have been proposed to change the global flexibility of the TM helix and thus to affect its positioning within the enzyme (Langosch et al., 2015; Stelzer et al., 2016). In addition, introducing glycine near the site where γ-secretase performs the first endoproteolytic cut, referred to as ε-site, facilitates cleavage (Fernandez et al., 2016) and solid-state NMR analysis revealed a helix-to-coil transition near the ε-site (Sato et al., 2009).

Here, we reveal an impact of the flexibility of the SPP substrate XBP1u TM helix by relating cell-based degradation assays to amide DHX kinetics. We observed that, mutating hydrophilic residues either in the N- or C-terminal half of the TM domain to leucine locally stabilize the TM helix, which positively correlates with a failure in SPP-catalyzed turnover. Interestingly, upon introducing di-glycine motifs at the

equivalent positions, we observed an increase in local helix dynamics as predicted, but no increase of cleavage efficiency. This is in line with our previous observation of flexibility requirement in the N-terminal TM domain portion (mt1) (Chen et al., 2014). However, the fact that the single-site mutation in mt7, but not that in mt6, mirrors the effect on cleavage seen with mt5 indicates that it is not helix flexibility but the primary structure that determines the turnover efficiency by SPP. Consistent with this model, we observed that in the TA protein HO1 a single serine-275, two positions downstream of the scissile peptide bond, was required for cleavage by SPP. Hence, it is attractive to speculate that the SPP active site binds its substrates with a serine in the P2' position. Although the exact cleavage site is not known for the XBP1u TM domain, comparison with reference peptides suggests processing between residues 194 and 199 (Chen et al., 2014), supporting the model that also here a serine residue is placed in the P2' position (see Figure 1 for XBP1u sequence). Slightly different results were observed with short signal peptides, where short h-regions completely lacking serine residues can be cleaved (Kronenberg-Versteeg et al., 2018; Lemberg and Martoglio, 2002). This suggest that there is no strict need for a particular side chain, but rather a combination of different factors help to accelerate SPP-catalyzed intramembrane cleavage. However, we showed that the neo-signal peptide derived from the XBP1u TM domain is not efficiently cleaved when the serine is replaced by a di-glycine motif (Figure 3D), highlighting the relevance of this feature also for signal peptide processing. Based on our results, we propose a model for substrate recognition by SPP (Figure 7). First, the helical substrate TM segment is recognized by a substrate-binding site on SPP that is remote from the active site, referred to as "exosite", as it has been suggested for other intramembrane proteases (Arutyunova et al., 2014; Fukumori and Steiner, 2016; Kornilova et al., 2005; Strisovsky et al., 2009).

We postulate that the substrate TM helix subsequently partially unwinds with the peptide comprising the scissile peptide backbone binding into a putative active site grove. Consistent with this, a recent study by resonance Raman spectroscopy revealed upon binding to an archaeal presenilin/SPP homologue (PSH) unwinding of the substrate helix into a β-strand geometry (Brown et al., 2018). Prior to local unwinding for cleavage, flexibility in the N-terminal half of the TM helix may be required for substrate translocation and its correct positioning (Fig 7A). Analogous, h-regions of signal peptides may have an intrinsic tendency to enter the SPP active site to minimize the energetically unfavored hydrophobic mismatch caused by their short span through the lipid bilayer (Figure 7B). While for short substrates this may be the only rate-limiting determinant, for more stable TM helices the equilibrium may be more on the side of the helical conformation and only a cognate recognition motif tightly docking into the active site may be efficiently bound and cleaved (Fig. 7A). Likewise, recent kinetic measurements using the purified archaeal PSH showed a preference for cleavage at a threonine at the scissile peptide bond (Naing et al., 2018). Although the mechanism of substrate selection may be different, in an intriguing parallel also rhomboid proteases combine recognition of a defined cleavage site motif (Strisovsky et al., 2009) with sampling TM domain dynamics (Akiyama and Maegawa, 2007; Moin and Urban, 2012; Strisovsky, 2015; Urban and Freeman, 2003). Overall, the emerging picture is that as it is known from the limited proteolysis by soluble proteases, also in the lipid bilayer a continuum of conformational control and site-specific substrate recognition governs protease selectivity. However, we note that for SPP this model remains speculative until the structure of an SPP-type protease with its substrate has been determined.

SPP-catalyzed cleavage acts as potent signal for degradation

Although prominent examples show that intramembrane proteases serve as a key trigger in activating membrane-tethered molecules, as observed for S2P-catalyzed release of the UPR transcription factor ATF6 (Ye et al., 2000b), there is growing evidence that a conserved role is in protein degradation (for review see (Avci and Lemberg, 2015)). Thereby, membrane anchors can be efficiently removed, e.g. leading to efficient degradation of XBP1u without any assistance of the AAA-ATPase p97 (Chen et al., 2014), which commonly pulls TM segments and luminal domains of ERAD substrates into the cytoplasm (Avci and Lemberg, 2015). In addition to a rapid and direct degradation route, it is attractive to speculate that the TM segment of XBP1u may sample specific properties of the surrounding lipid environment and therefore link its degradation rate to homeostasis control. Likewise, there is growing evidence that TM segments of the UPR sensors IRE1 and PERK are sensitive to membrane stress by sensing lipid perturbations (Kono et al., 2017). The influence of lipid composition on XBP1u TM helix dynamics and its recognition by the SPP-triggered ERAD are important questions remaining. Taken together with our earlier work on the nutrient regulated yeast SPP orthologue Ypf1 (Avci et al., 2014), proteolytic clipping of ERAD substrates emerges as an important evolutionary conserved principle in membrane protein abundance control. Further insights into the substrate spectrum of SPP and how its activity is regulated will likely reveal additional unexpected roles for this ERAD machinery.

Materials and Methods

Additional information on the materials and methods can be found in the

supplemental information.

Cell lines and transfection

Hek293T cells were grown in DMEM (Invitrogen) supplemented with 10% fetal

bovine serum (FBS) at 37°C in 5% CO_2. Transient transfections were performed

using 25 kDa linear polyethylenimine (Polysciences) (Durocher et al., 2002).

Typically, 1 μg of XBP1u plasmid and 300 ng of HO1 plasmid was used per well of a

6-well plate (see supplemental information for details on the plasmids). Total

transfected DNA (2 μg/well) was held constant by the addition of empty plasmid. If

not otherwise stated, cells were harvested 24 h after transfection. To inhibit the

proteasome cells were treated with 1 μM epoxomicin (Calbiochem) for 16 hours and

subsequently harvested for western blot analysis.

Cycloheximide chase experiments

Cycloheximide (100 μg/ml) chase was conducted 24 h after transfection of Hek293T

cells, and cell extracts were subjected to Western blot analysis. To inhibit SPP, 50 μM

(Z-LL)$_2$-ketone (Calbiochem) in dimethyl sulfoxide (DMSO) was added 2 h before

the addition of cycloheximide and the inhibitor was kept on the cells during the chase

experiment.

SDS-PAGE and western blotting

Proteins were solubilized in SDS sample buffer (50 mM Tris-Cl pH 6.8; 10 mM

EDTA, 5% glycerol, 2% SDS, 0.01% bromphenol blue) containing 5% β-

mercaptoethanol. All samples were incubated for 15 min at 65°C. For

deglycosylation, solubilized proteins were treated with EndoH (New England

Biolabs) according to the manufacturer's protocol. For western blot analysis, proteins

were separated by SDS-PAGE using Tris-glycine acrylamide gels and transferred to

PVDF membrane followed by enhanced chemiluminescence analysis (Pierce) to

detect the bound antibodies (see supplemental information for further information).

For detection the LAS-4000 system (Fuji) was used. Data shown are representative of

three independent experiments. For quantification, we used the FIJI (Schindelin et al.,

2012) and data acquired from the LAS-4000.

Microscopy

For immunofluorescence analysis, fixed cells adhered to glass coverslides were

analyzed using a Zeiss LSM confocal microscope as described in the supplemental

information.

***In vitro* transcription, translation/translocation, protease protection assay and
signal peptide processing**

pRK5-Prl or pRK5-XBP1uTMD-Prl plasmids were linearized with XhoI and

transcribed with the SP6 RNA polymerase at 42°C in the presence of 500 μM

m7G(5′)ppp(5′)G CAP analogue (New England Biolabs). mRNA was translated in

rabbit reticulocyte lysate (Promega) containing [^{35}S]-methionine/cysteine as described

before (Lemberg and Martoglio, 2002). Where indicated, nuclease-treated rough

microsomes prepared from dog pancreas were added (Martoglio et al., 1998). For

protease protection, the reactions were treated with 0.5 mg/ml proteinase K for 45 min

on ice. As a control, proteinase K was added in the presence of 1% Triton X-100.

Subsequently, 10 µg/ml PMSF was added, and samples were analyzed on 12% Tris-glycine acrylamide gels and autoradiography as described previously (Lemberg and Martoglio, 2002). To inhibit SPP processing in the signal peptide processing assay 10 µM (Z-LL)$_2$-ketone was added. Samples were incubated 20 min at 30°C. Microsomes were extracted with 500 mM KOAc and analyzed on Tris–Bicine acrylamide gels (15% T, 5% C, 8 M urea). Translation of the reference peptide containing the XBP1u-TMD was performed in wheat germ extract at 25°C for 30 min (Martoglio et al., 1998).

Determination of SPP-cleavage site

SPP cleavage site was determined by LC/MS analysis of the C-terminal peptide released from immunoprecipitated GFP-TEV-FLAG-HO1 cytoplasmic cleavage fragment by *in vitro* TEV digest as outlined in the supplemental information.

In silico **TM domain analysis**

To compare the XBP1u TM region, signal sequences and canonical TM segments we determined the free energy differences (ΔG_{app}) by the online ΔG prediction server (http://dgpred.cbr.su.se). To predict the presence and the site of signal peptide cleavage we used SignalP 3.0 (http://www.cbs.dtu.dk/services/SignalP-3.0) and TM domain mutants were evaluated using TMHMM (http://www.cbs.dtu.dk/services/TMHMM).

Peptide synthesis

Peptides were synthesized by Fmoc chemistry (PSL, Heidelberg, Germany) and purified by HPLC. They were > 90% pure as judged by mass spectrometry.

Concentrations were determined via UV spectroscopy using an extinction coefficient at 280 nm of 11 000 $M^{-1} \cdot cm^{-1}$.

Circular dichroism spectroscopy

For CD spectroscopy, peptides were dissolved at 50 µM in 80% (v/v) TFE, 20% (v/v) 10 mM Tris/HCl (respectively 5 mM PBS for mt5 and mt3), pH 7. CD spectra were obtained using a Jasco J-710 CD spectrometer from 190 nm to 260 nm in a 1.0 mm quartz cuvette at 20°C with a response of 1 s, a scan speed of 100 nm/min and a sensitivity of 100 mdeg/cm. Spectra were the signal-averaged accumulations of 10 scans with the baselines (corresponding to solvent) subtracted. Mean molar ellipticities were calculated and secondary structure contents estimated by deconvoluting the spectra using CDNN software (Bohm et al., 1992) with CDNN complex reference spectra.

DHX experiments

Detailed information on DHX exchange experiments and analysis of residue-specific amide exchange kinetics by the semi-automated software module *ETD Fragment Analyzer* that is part of the software suite MassMap® (MassMap GmbH & Co. KG, Freising, Germany) can be found in the supplemental online information.

Acknowledgements

We thank Dönem Avci for critical reading of the manuscript, Stefan Lichtenthaler for advice and Thomas Ruppert (ZMBH, MS facility) for MS-based cleavage site determination. This study was supported by the Deutsche Forschungsgemeinschaft (FOR2290) to DL (La699/20-1) and MKL (Le2749/1-1).

Authors Contribution

S.S.Y. designed and performed all cell-free translocation assays, most cell-based experiments and wrote the manuscript. W.S. performed all CD and DHX experiments and analyzed data. A.L. helped determining the SPP-cleavage site. M.W. developed the *ETD Fragment Analyzer* software. D.L. and M.K.L. guided the project, designed experiments, and wrote the manuscript.

Declaration of Interests

M.W. is founder and owner of MassMap GmbH & Co. KG, the company that commercializes the *ETD Fragment Analyzer*.

References

Akiyama, Y., and Maegawa, S. (2007). Sequence features of substrates required for cleavage by GlpG, an Escherichia coli rhomboid protease. Mol. Microbiol. *64*, 1028-1037.

Arutyunova, E., Panwar, P., Skiba, P.M., Gale, N., Mak, M.W., and Lemieux, M.J. (2014). Allosteric regulation of rhomboid intramembrane proteolysis. EMBO J *33*, 1869-1881.

Avci, D., Fuchs, S., Schrul, B., Fukumori, A., Breker, M., Frumkin, I., Chen, C.Y., Biniossek, M.L., Kremmer, E., Schilling, O.*, et al.* (2014). The Yeast ER-Intramembrane Protease Ypf1 Refines Nutrient Sensing by Regulating Transporter Abundance. Mol Cell *56*, 630-640.

Avci, D., and Lemberg, M.K. (2015). Clipping or Extracting: Two Ways to Membrane Protein Degradation. Trends Cell Biol *25*, 611-622.

Bendtsen, J.D., Nielsen, H., von Heijne, G., and Brunak, S. (2004). Improved prediction of signal peptides: SignalP 3.0. J Mol Biol *340*, 783-795.

Blobel, G., and Dobberstein, B. (1975). Transfer of proteins across membranes. I. Presence of proteolytically processed and unprocessed nascent immunoglobulin light chains on membrane-bound ribosomes of murine myeloma. J. Cell Biol. *67*, 835-851.

Bohm, G., Muhr, R., and Jaenicke, R. (1992). Quantitative analysis of protein far UV circular dichroism spectra by neural networks. Protein Eng *5*, 191-195.

Boname, J.M., Bloor, S., Wandel, M.P., Nathan, J.A., Antrobus, R., Dingwell, K.S., Thurston, T.L., Smith, D.L., Smith, J.C., Randow, F.*, et al.* (2014). Cleavage by signal peptide peptidase is required for the degradation of selected tail-anchored proteins. J Cell Biol *205*, 847-862.

Bonifacino, J.S., Suzuki, C.K., and Klausner, R.D. (1990). A peptide sequence confers retention and rapid degradation in the endoplasmic reticulum. Science *247*, 79-82.

Bowie, J.U. (2011). Membrane protein folding: how important are hydrogen bonds? Current opinion in structural biology *21*, 42-49.

Brown, M.C., Abdine, A., Chavez, J., Schaffner, A., Torres-Arancivia, C., Lada, B., JiJi, R.D., Osman, R., Cooley, J.W., and Ubarretxena-Belandia, I. (2018). Unwinding of the Substrate Transmembrane Helix in Intramembrane Proteolysis. Biophys J *114*, 1579-1589.

Brown, M.S., Ye, J., Rawson, R.B., and Goldstein, J.L. (2000). Regulated intramembrane proteolysis: a control mechanism conserved from bacteria to humans. Cell *100*, 391-398.

Chen, C., Malchus, N.S., Hehn, B., Stelzer, W., Avci, D., Langosch, D., and Lemberg, M.K. (2014). Signal peptide peptidase functions in ERAD to cleave the unfolded protein response regulator XBP1u. EMBO J *33*, 2492-2506.

Christianson, J.C., and Ye, Y. (2014). Cleaning up in the endoplasmic reticulum: ubiquitin in charge. Nat Struct Mol Biol *21*, 325-335.

Cymer, F., von Heijne, G., and White, S.H. (2014). Mechanisms of Integral Membrane Protein Insertion and Folding. J Mol Biol.

Durocher, Y., Perret, S., and Kamen, A. (2002). High-level and high-throughput recombinant protein production by transient transfection of suspension-growing human 293-EBNA1 cells. Nucleic Acids Res *30*, E9.

Fernandez, M.A., Biette, K.M., Dolios, G., Seth, D., Wang, R., and Wolfe, M.S. (2016). Transmembrane Substrate Determinants for gamma-Secretase Processing of APP CTFbeta. Biochemistry *55*, 5675-5688.

Fleig, L., Bergbold, N., Sahasrabudhe, P., Geiger, B., Kaltak, L., and Lemberg, M.K. (2012). Ubiquitin-Dependent Intramembrane Rhomboid Protease Promotes ERAD of Membrane Proteins. Mol Cell *47*, 558-569.

Fukumori, A., and Steiner, H. (2016). Substrate recruitment of gamma-secretase and mechanism of clinical presenilin mutations revealed by photoaffinity mapping. EMBO J *35*, 1628-1643.

Haeuptle, M.T., Flint, N., Gough, N.M., and Dobberstein, B. (1989). A tripartite structure of the signals that determine protein insertion into the endoplasmic reticulum membrane. J. Cell Biol. *108*, 1227-1236.

Hegde, R.S., and Keenan, R.J. (2011). Tail-anchored membrane protein insertion into the endoplasmic reticulum. Nat Rev Mol Cell Biol *12*, 787-798.

Hessa, T., Meindl-Beinker, N.M., Bernsel, A., Kim, H., Sato, Y., Lerch-Bader, M., Nilsson, I., White, S.H., and von Heijne, G. (2007). Molecular code for transmembrane-helix recognition by the Sec61 translocon. Nature *450*, 1026-1030.

Hetz, C., and Papa, F.R. (2018). The Unfolded Protein Response and Cell Fate Control. Mol Cell *69*, 169-181.

Hogel, P., Gotz, A., Kuhne, F., Ebert, M., Stelzer, W., Rand, K.D., Scharnagl, C., and Langosch, D. (2018). Glycine Perturbs Local and Global Conformational Flexibility of a Transmembrane Helix. Biochemistry *57*, 1326-1337.

Hsu, F.F., Yeh, C.T., Sun, Y.J., Chiang, M.T., Lan, W.M., Li, F.A., Lee, W.H., and Chau, L.Y. (2015). Signal peptide peptidase-mediated nuclear localization of heme oxygenase-1 promotes cancer cell proliferation and invasion independent of its enzymatic activity. Oncogene *34*, 2360-2370.

Hussy, P., Langen, H., Mous, J., and Jacobsen, H. (1996). Hepatitis C virus core protein: carboxy-terminal boundaries of two processed species suggest cleavage by a signal peptide peptidase. Virology *224*, 93-104.

Kanda, S., Yanagitani, K., Yokota, Y., Esaki, Y., and Kohno, K. (2016). Autonomous translational pausing is required for XBP1u mRNA recruitment to the ER via the SRP

pathway. Proceedings of the National Academy of Sciences of the United States of America *113*, E5886-E5895.

Kang, S.W., Rane, N.S., Kim, S.J., Garrison, J.L., Taunton, J., and Hegde, R.S. (2006). Substrate-specific translocational attenuation during ER stress defines a pre-emptive quality control pathway. Cell *127*, 999-1013.

Kapp, K., Schrempf, S., Lemberg, M.K., and Dobberstein, B. (2009). Post-targeting functions of signal peptides. In Protein Transport into the Endoplasmic Reticulum, R. Zimmermann, ed. (Austin: Landes Bioscience), pp. 1-16.

Konermann, L., Pan, J., and Liu, Y.H. (2011). Hydrogen exchange mass spectrometry for studying protein structure and dynamics. Chem Soc Rev *40*, 1224-1234.

Kono, N., Amin-Wetzel, N., and Ron, D. (2017). Generic membrane-spanning features endow IRE1alpha with responsiveness to membrane aberrancy. Mol Biol Cell *28*, 2318-2332.

Kornilova, A.Y., Bihel, F., Das, C., and Wolfe, M.S. (2005). The initial substrate-binding site of gamma-secretase is located on presenilin near the active site. Proceedings of the National Academy of Sciences of the United States of America *102*, 3230-3235.

Kronenberg-Versteeg, D., Eichmann, M., Russell, M.A., de Ru, A., Hehn, B., Yusuf, N., van Veelen, P.A., Richardson, S.J., Morgan, N.G., Lemberg, M.K.*, et al.* (2018). Molecular Pathways for Immune Recognition of Preproinsulin Signal Peptide in Type 1 Diabetes. Diabetes *67*, 687-696.

Langosch, D., Scharnagl, C., Steiner, H., and Lemberg, M.K. (2015). Understanding intramembrane proteolysis: from protein dynamics to reaction kinetics Trends Biochem Sci *40*, 318-327.

Lemberg, M.K. (2011). Intramembrane Proteolysis in Regulated Protein Trafficking. Traffic *12*, 1109-1118.

Lemberg, M.K., Bland, F.A., Weihofen, A., Braud, V.M., and Martoglio, B. (2001). Intramembrane proteolysis of signal peptides: an essential step in the generation of HLA-E epitopes. J. Immunol. *167*, 6441-6446.

Lemberg, M.K., and Martoglio, B. (2002). Requirements for signal peptide peptidase-catalyzed intramembrane proteolysis. Mol. Cell *10*, 735-744.

Lemberg, M.K., and Martoglio, B. (2004). On the mechanism of SPP-catalysed intramembrane proteolysis; conformational control of peptide bond hydrolysis in the plane of the membrane. FEBS Lett. *564*, 213-218.

Lermyte, F., Łącki, M., Valkenborg, D., Baggerman, G., Gambin, A., and Sobott, F. (2015). Understanding reaction pathways in top-down ETD by dissecting isotope distributions: A mammoth task. International Journal of Mass Spectrometry *390*, 146-154.

Li, H., Korennykh, A.V., Behrman, S.L., and Walter, P. (2010). Mammalian endoplasmic reticulum stress sensor IRE1 signals by dynamic clustering. Proceedings of the National Academy of Sciences of the United States of America *107*, 16113-16118.

Martoglio, B., Hauser, S., and Dobberstein, B. (1998). Cotranslational translocation of proteins into microsomes derived from the rough endoplasmic reticulum of mammalian cells. In Cell Biology: A Laboratory Handbook, J.E. Celis, ed. (San Diego: Academic Press), pp. 265-274.

Mentrup, T., Fluhrer, R., and Schroder, B. (2017). Latest emerging functions of SPP/SPPL intramembrane proteases. European journal of cell biology *96*, 372-382.

Moin, S.M., and Urban, S. (2012). Membrane immersion allows rhomboid proteases to achieve specificity by reading transmembrane segment dynamics. elife *1*, e00173.

Mothes, W., Prehn, S., and Rapoport, T.A. (1994). Systematic probing of the environment of a translocating secretory protein during translocation through the ER membrane. EMBO J *13*, 3973-3982.

Naing, S.H., Kalyoncu, S., Smalley, D.M., Kim, H., Tao, X., George, J.B., Jonke, A.P., Oliver, R.C., Urban, V.S., Torres, M.P.*, et al.* (2018). Both positional and chemical variables control in vitro proteolytic cleavage of a presenilin ortholog. J Biol Chem *293*, 4653-4663.

Pester, O., Gotz, A., Multhaup, G., Scharnagl, C., and Langosch, D. (2013). The cleavage domain of the amyloid precursor protein transmembrane helix does not exhibit above-average backbone dynamics. Chembiochem *14*, 1943-1948.

Plumb, R., Zhang, Z.R., Appathurai, S., and Mariappan, M. (2015). A functional link between the co-translational protein translocation pathway and the UPR. Elife *4*.

Poschner, B.C., Quint, S., Hofmann, M.W., and Langosch, D. (2009). Sequence-specific conformational dynamics of model transmembrane domains determines their membrane fusogenic function. J Mol Biol *386*, 733-741.

Quint, S., Widmaier, S., Minde, D., Hornburg, D., Langosch, D., and Scharnagl, C. (2010). Residue-specific side-chain packing determines the backbone dynamics of transmembrane model helices. Biophys J *99*, 2541-2549.

Rand, K.D., Pringle, S.D., Morris, M., and Brown, J.M. (2012). Site-specific analysis of gas-phase hydrogen/deuterium exchange of peptides and proteins by electron transfer dissociation. Anal Chem *84*, 1931-1940.

Sato, T., Tang, T.C., Reubins, G., Fei, J.Z., Fujimoto, T., Kienlen-Campard, P., Constantinescu, S.N., Octave, J.N., Aimoto, S., and Smith, S.O. (2009). A helix-to-coil transition at the epsilon-cut site in the transmembrane dimer of the amyloid precursor protein is required for proteolysis. Proceedings of the National Academy of Sciences of the United States of America *106*, 1421-1426.

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., *et al.* (2012). Fiji: an open-source platform for biological-image analysis. Nature methods *9*, 676-682.

Schrul, B., Kapp, K., Sinning, I., and Dobberstein, B. (2010). Signal peptide peptidase (SPP) assembles with substrates and misfolded membrane proteins into distinct oligomeric complexes. Biochem J *427*, 523-534.

Skinner, J.J., Lim, W.K., Bedard, S., Black, B.E., and Englander, S.W. (2012). Protein dynamics viewed by hydrogen exchange. Protein Sci *21*, 996-1005.

Snapp, E.L., Sharma, A., Lippincott-Schwartz, J., and Hegde, R.S. (2006). Monitoring chaperone engagement of substrates in the endoplasmic reticulum of live cells. Proceedings of the National Academy of Sciences of the United States of America *103*, 6536-6541.

Stelzer, W., Poschner, B.C., Stalz, H., Heck, A.J., and Langosch, D. (2008). Sequence-specific conformational flexibility of SNARE transmembrane helices probed by hydrogen/deuterium exchange. Biophys J *95*, 1326-1335.

Stelzer, W., Scharnag, C., Leurs. U, Rand, K., and Langosch, D. (2016). The Impact of the 'Austrian' Mutation of the Amyloid Precursor Protein Transmembrane Helix is Communicated to the Hinge Region. Chemistry Select *1*, 4408-4412.

Strisovsky, K. (2015). Why cells need intramembrane proteases - a mechanistic perspective. FEBS J.

Strisovsky, K., Sharpe, H.J., and Freeman, M. (2009). Sequence-specific intramembrane proteolysis: identification of a recognition motif in rhomboid substrates. Mol Cell *36*, 1048-1059.

Tirosh, B., Iwakoshi, N.N., Glimcher, L.H., and Ploegh, H.L. (2006). Rapid turnover of unspliced Xbp-1 as a factor that modulates the unfolded protein response. J Biol Chem *281*, 5852-5860.

Urban, S., and Freeman, M. (2003). Substrate specificity of rhomboid intramembrane proteases is governed by helix-breaking residues in the substrate transmembrane domain. Mol. Cell *11*, 1425-1134.

van Klompenburg, W., and de Kruijff, B. (1998). The role of anionic lipids in protein insertion and translocation in bacterial membranes. J Membr Biol *162*, 1-7.

Voorhees, R.M., and Hegde, R.S. (2016). Structure of the Sec61 channel opened by a signal sequence. Science *351*, 88-91.

Walter, P., and Ron, D. (2011). The unfolded protein response: from stress pathway to homeostatic regulation. Science *334*, 1081-1086.

Weihofen, A., Binns, K., Lemberg, M.K., Ashman, K., and Martoglio, B. (2002). Identification of signal peptide peptidase, a presenilin-type aspartic protease. Science *296*, 2215-2218.

Weihofen, A., Lemberg, M.K., Ploegh, H.L., Bogyo, M., and Martoglio, B. (2000). Release of signal peptide fragments into the cytosol requires cleavage in the transmembrane region by a protease activity that is specifically blocked by a novel cysteine protease inhibitor. J. Biol. Chem. *275*, 30951-30956.

Wolfe, M.S. (2009). Intramembrane proteolysis. Chem Rev *109*, 1599-1612.

Yanagitani, K., Imagawa, Y., Iwawaki, T., Hosoda, A., Saito, M., Kimata, Y., and Kohno, K. (2009). Cotranslational targeting of XBP1 protein to the membrane promotes cytoplasmic splicing of its own mRNA. Mol Cell *34*, 191-200.

Yanagitani, K., Kimata, Y., Kadokura, H., and Kohno, K. (2011). Translational pausing ensures membrane targeting and cytoplasmic splicing of XBP1u mRNA. Science *331*, 586-589.

Ye, J., Dave, U.P., Grishin, N.V., Goldstein, J.L., and Brown, M.S. (2000a). Asparagine-proline sequence within membrane-spanning segment of SREBP triggers intramembrane cleavage by site-2 protease. Proc. Natl. Acad. Sci. USA *97*, 5123-5128.

Ye, J., Rawson, R.B., Komuro, R., Chen, X., Dave, U.P., Prywes, R., Brown, M.S., and Goldstein, J.L. (2000b). ER stress induces cleavage of membrane-bound ATF6 by the same proteases that process SREBPs. Mol. Cell *6*, 1355-1364.

Yoshida, H., Matsui, T., Yamamoto, A., Okada, T., and Mori, K. (2001). XBP1 mRNA is induced by ATF6 and spliced by IRE1 in response to ER stress to produce a highly active transcription factor. Cell *107*, 881-891.

Yoshida, H., Oku, M., Suzuki, M., and Mori, K. (2006). pXBP1(U) encoded in XBP1 pre-mRNA negatively regulates unfolded protein response activator pXBP1(S) in mammalian ER stress response. J Cell Biol *172*, 565-575.

Yoshida, H., Uemura, A., and Mori, K. (2009). pXBP1(U), a negative regulator of the unfolded protein response activator pXBP1(S), targets ATF6 but not ATF4 in proteasome-mediated degradation. Cell Struct Funct *34*, 1-10.

Zhao, Y., Li, X., Cai, M.Y., Ma, K., Yang, J., Zhou, J., Fu, W., Wei, F.Z., Wang, L., Xie, D.*, et al.* (2013). XBP-1u suppresses autophagy by promoting the degradation of FoxO1 in cancer cells. Cell Res *23*, 491-507.

Figure Legends

Figure 1. XBP1u TM domain is sufficient for insertion into the ER membrane in a type II orientation.

(A) Predicted apparent free energy differences (ΔG_{app}) for Sec61 insertion of the XBP1u TM domain (TMD) and ER-targeting signal peptides (SP) compared to the TMD of calnexin (Hessa et al., 2007). Sequences taken for the calculation are underlined.

(B) Translocation and protease protection assay revealing that the XBP1u TM domain is sufficient to mediate ER insertion. *In vitro* translation of mRNA encoding either Prl or the XBP1u-TMD-Prl fusion proteins (wt and mt1) in the absence or presence of canine rough microsomes. Reactions were treated with Proteinase K and Triton X-100 as indicated and analyzed by SDS-PAGE and autoradiography. Filled arrow, pre-protein; open arrow, translocated Prl.

(C) XBP1u wt, mt1 and mt2 ER targeting and insertion as type II membrane protein assessed by Endo H-sensitivity of the glycoslylation mutant R232N (RN) (triangle). Hek293T cells are transfected with FLAG-tagged XBP1u-RN harboring wt, mt1 or mt2 TM domain and cell lysates are treated with EndoH. Glycosylation efficiencies are indicated respectively.

Figure 2. Turnover of XBP1u is governed by its TM domain.

(A) Immunofluorescent microscopy of transiently transfected Hek293T cells with FLAG-tagged tail-anchored (TA) XBP1u wt, mt1 and mt2 co-expressing the ER marker RFP-KDEL. TA-anchored XBP1u mt1 and mt2 are efficiently targeted to the ER, while TA-anchored XBP1u wt showed faint ER-staining and prominent co-localization with Hoechst-staining (shown in blue) for chromosomal DNA, suggesting

that SPP-generated cleavage fragments are either degraded by the cytoplasmic proteasome or accumulate in the nucleus. Scale bar, 10 μM.

(B) XBP1u-TA mt1 and mt2 are stable, whereas the wt construct is rapidly degraded as assessed cycloheximide (CHX) chase and western blotting (mean ± SEM, n=3). Quantification of the degradation kinetics of the full-length (FL) XBP1u is shown for comparison.

(C) Processing of signal peptides in microsomes. *In vitro* translation of mRNA encoding pPrl, XBP1u-TMD-Prl fusion proteins (wt and mt1) in the absence (lane 1), or presence of ER-derived microsomes and 1 μM of the SPP inhibitor $(Z-LL)_2$-ketone. Peptides were separated on 15% Tris-Bicine gels and visualized by phosphorimaging. Right panel shows quantification of signal peptide processing comparing relative amounts of signal peptide obtained from vehicle treated to SPP inhibitor treated sample (mean ± SEM, n=4). Ref, shows a reference peptide comprising the XBP1u-TMD neo-signal peptide; asterisk, putative dimer of *in vitro* translated reference signal peptide.

(D) Overall amide DHX kinetics of XBP1u wt TM domain compared to mt1 and mt2 (n ≥ 3, SEM are smaller than the sizes of the symbols). All used peptides contain N- and C-terminal KKK tags for better solubility and gas phase fragmentation.

Figure 3. Positional effect of di-glycine motifs on SPP-catalyzed turnover

(A) Sequences of XBP1u di-glycine TM domain mutants.

(B) XBP1u mt3 and mt5 insert into the ER as type II membrane protein whereas mt4 predominantly stays cytoplasmic. ER targeting and insertion efficiencies assessed by the R232N glycosylation reporter (RN).

(C) Degradation kinetics of XBP1u wt, mt3 and mt5 in transfected Hek293T cells as

assessed cycloheximide (CHX) chase and western blotting (mean ± SEM, n=3).

(D) *In vitro* signal peptide cleavage of XBP1u-TMD-Prl mt5. Quantification of signal peptide processing comparing relative amounts of signal peptide obtained from vehicle treated to SPP inhibitor treated sample (mean ± SEM, n=4). Quantification of XBP1u-TMD-Prl wt is shown for comparison (see Figure 2C).

Figure 4. Di-glycine mutants show local TM helix destabilization.

(A) Overall amide DHX kinetics of XBP1u wt compared to mt3, mt4 and mt5 recorded from the masses of triply charged ions ($n \geq 3$, SEM are smaller than the sizes of the symbols). The inset focuses on the initial 2 h of the incubation period.

(B) Free energy change ΔG of intrahelical amide H-bond formation.

(C) Differences in ΔG of wt and ΔG of mutant TM domains (ΔΔG). Values are only shown for those residues where sufficient data points could be collected for wt and the respective mutant (see Table S2). Wt residues in the mutant TM domain sequences are represented as dots. Error bars correspond to standard confidence intervals calculated from the standard errors of k_{exp}.

Figure 5. XBP1u turnover depends not only on global TM helix dynamics but also on residue specific features.

(A) Sequences of XBP1u single-residue TM domain mutants.

(B) ER targeting and topology of XBP1u mt6 and mt7 assessed by the R232N glycosylation reporter (RN).

(C) Degradation kinetics of XBP1u mt6 and mt7 in transfected Hek293T cells assessed by cycloheximide (CHX) chase and western blotting (mean ± SEM, n=3). Quantification of XBP1u wt is shown for comparison (see Figure 3C).

Figure 6. Serine residue facilitates SPP-catalyzed cleavage of TA proteins.

(A) Sequence of HO1 TM domain wt and mutants.

(B) Influence of proline-266, arginine-269 and serine-275 residues on HO1 stability assessed cycloheximide (CHX) chase and western blotting (mean ± SEM, n=3). See Figure S4 for representative western blots.

(C) Steady-state western blot analysis of HO1 wt and serine-275 mutants together with increasing concentrations of SPP to detect SPP-generated cleavage fragments.

(D) Schematic representation of GFP-TEV-FLAG-HO1-TMD, its cleavage by SPP and further *in vitro* processing with TEV protease to generate small peptides for MS analysis.

(E) Cleavage site mapping of HO1. The N-terminal fragment of the GFP-TEV-FLAG-HO1-TMD (as outlined in panel D) generated by ectopically expressed SPP was immunoprecipitated from the cytosol fraction after overnight treatment with 0.5 μM epoxomicin, digested with TEV protease and the resulting FLAG-tagged peptide was analyzed by MS. Cytosol from mock transfected cells was used as control. The inset shows high-resolution isotopic pattern of the 5274.76 Da peak.

Figure 7. Model for SPP substrate recognition.

(A) SPP binds helical TM spans of type II membrane proteins and TA proteins by a putative docking site (dark blue), which despite its predicted low affinity may serve as an "exosite" introducing the first layer of specificity. In order to expose the scissile peptide bond to the active site (bright color, "D" indicates catalytic residues), substrate TM helix has to partially unfold, a reaction that is facilitated by helix-destabilizing features (X). This equilibrium may be shifted to the open conformation

by site-specific recognition of substrate features such as serine (green circle) that is bound to the active site cleft via putative "subsites". Reaction rate may be determined by helix-dynamics (rate controlled) and the primary sequence (affinity driven).

(B) With a h-region in the range of 7 to 15 amino acids, TM-spans of signal peptides are commonly too short to form a regular TM helix and may directly unfold into the SPP active site to minimize the high energetic state of hydrophobic mismatch in the surrounding lipid bilayer. Subsequently, side-specific contacts to cleavage site region is not a strict requirement and cleavage is only rate controlled.

Figure S1. Examination of XBP1u mt1 and mt2 secondary structure, related to Figure 2.

(A) Helical content of XBP1u wt, mt1 and mt2 TM peptides determined by CD spectroscopy. CD spectra were recorded in 80% TFE/buffer, averaged, and secondary structure helix contents were calculated with CDNN (inset, n = 3, means ± SEM).

(B) Exemplary mass spectra of the triply charged XBP1u wt ion at different time points. The spectrum at t = 0 min was recorded after exchange under stop conditions (3 min incubation on ice at pH 2.5) where only very labile deuteriums exchange. Low intensity isotopic envelopes at calculated masses ~22 Da above the dominant envelopes likely originate from Na^+-adducts.

Figure S2. Positional effect of di-glycine motifs on ER targeting and SPP-catalyzed turnover, related to Figure 3.

(A) XBP1u mt3 and mt5 are efficiently targeted to the ER whereas mt4 predominantly localizes to the cytoplasm and nucleoplasm. Immunofluorescence microscopy of transiently transfected Hek293T cells co-expressing FLAG-tagged

XBP1u full-length mt3, mt4 and mt5 with the ER marker RFP-KDEL is shown.

Hoechst-staining staining is shown in blue. Scale bar, 10 μM.

(B) Cycloheximide (CHX) chase analysis to determine degradation rate of XBP1u-

mt4 (mean ± SEM, n=3). For comparison quantification of the degradation kinetics of

XBP1u wt is shown (see Fig. 3C).

(C) Cycloheximide (CHX) chase analysis to determine degradation rate of XBP1u

mt3 and mt5 in presence of 50 μM (Z-LL)$_2$-ketone (n=2); DMSO, vehicle control.

Figure S3. Di-glycine mutants show local TM helix destabilization, related to

Figure 4.

(A) Helical content of di-glycine mutants XBP1u mt3, mt4 and mt5 TM peptides

determined by CD spectroscopy. CD spectra were recorded in 80% TFE/buffer,

averaged, and secondary structure helix contents were calculated with CDNN (inset, n

= 3, means ± SEM). For comparison spectrum of XBP1u wt is shown (taken from

Figure S1A).

(B) Exchange rate constants k_{exp} of individual amide deuterons. The values of k_{exp}

(filled symbols, means ± SEM) were derived from exponential fits of the residue-

specific DHX kinetics (given in panel C) as obtained from ETD experiments. Only

those values are shown that are covered by sufficient data points. Empty symbols

represent the respective chemical exchange rate constant k_{ch} that reflects the DHX

kinetics in an unfolded state.

(C) Residue-specific DHX kinetics obtained after ETD. The calculated deuterium

contents D (mean values, n ≥ 3) of the respective amides are plotted against the log of

the exchange period t. The identities of the wt residues (in black), along with the

identities of the respective mutants (color coded) are given in the insets. Exponential

fits are depicted for those kinetics that were complete enough for calculating the exchange rate constants given in panel (B) and in Table S2.

(D) $\Delta\Delta G$ between wt and mt4 TM domain (compared to $\Delta\Delta G$ of XBP1u wt, see Fig. 4A).

Figure S4. Influence of proline-266, arginine-269 and serine-275 on stability of HO1, related to Figure 4.

FLAG-tagged HO1-wt, S275L, S275A, S275G as well as P266L and R269L were transfected in Hek293T cells and their stability was assessed by cycloheximide (CHX) chase and western blotting.

Table S1. ΔG_{app} of XBP1u TM domain glycine mutants, related to Figure 3 and Figure 5.

XBP1u	Sequence	ΔG_{app}
wt	ISPWILAVLTLQIQSLISCWAFWTTW	0,07
mt3	I**GG**WILAVLTLQIQSLISCWAFWTTW	-0,27
mt4	ISPWILAV**GG**LQIQSLISCWAFWTTW	0,766
mt5	ISPWILAVLTLQI**GG**LISCWAFWTTW	-1,222
mt6	ISPWILAVLTLQI**G**SLISCWAFWTTW	-0,986
mt7	ISPWILAVLTLQIQ**G**LISCWAFWTTW	-0,158

Table S2. Numerical values of k_{exp} and ΔG, related to Figure 4.

See separate file.

Supplemental Methods

Plasmids

For the cell-based assays, if not otherwise stated, all open reading frames were cloned into pcDNA3.1+ (Invitrogen). Plasmids encoding human XBP1u (harboring a silent mutation in the 3'-IRE1-splice-site) tagged with an N-terminal triple FLAG-tag and SPP with triple HA-tag inserted between residue 373 and the C-terminal KKXX ER-retention signal (SPP-HA) had been described previously (Chen et al., 2014). The XBP1u TM domain mutants mt1 (S187L, P188L, A192L), mt2 (Q197L, Q199L, S200L, S203L), mt3 (S187G, P188G), mt4 (L194G, T195G), mt5 (Q199G, S200S), mt6 (Q199G), mt7 (S200G) as well as the R232N glycosylation mutants were generated by Quick-Change site-directed mutagenesis (Stratagene). The deletion constructs i.e. the tail-anchored XBP1u mutants were generated by subcloning the respective region of the open reading frame (1-207). HO1 (full-length ORF Gateway clone 187935150) was cloned into pcDNA3.1 with an N-terminal triple FLAG-tag and the point mutations (P266A, R269L, S275L, S275A and S275G) were introduced by Quick-Change mutagenesis. For the *in vitro* experiments all constructs were cloned into pRK5. The construct coding bovine pPrl has been described previously (Schrul et al., 2010). The XBP1u-TMD-Prl, wt and mt1 were ordered as gene blocks (IDT) and cloned into pRK5. XBP1u-TMD-Prl mt4 was generated by Quick- Change site-directed mutagenesis (Stratagene) by using XBP1u-TMD-Prl wt as a template. In order to map the SPP-cleavage site in transfected cells, the open reading frame encoding residues 237-288 of HO1 was cloned into pEGFP-C1 (Invitrogen) proceeded by a linker contacting a TEV protease cleavage site and a single FLAG-tag (...ENLYFQ^GDYKDDDDKG...). The ER marker ssRFP-KDEL were previously described (Snapp et al., 2006).

Microscopy

For immunofluorescence detection of N-terminally FLAG-tagged XBP1u, transfected Hek293T cells were grown on glass cover slides and fixed in PBS containing 4% paraformaldehyde for 30 min. Permeabilization was performed in PBS containing 0.5% Triton X-100 for 10 min. After blocking cells for 30 min in PBS containing 20% FBS and 0.5% Tween-20, cells were incubated in PBS/FBS/Tween-20 with the primary antibody for 1 h at room temperature (see below for further information). Subsequently, cells were washed in PBS/FBS/Tween-20, incubated with fluorescent secondary antibody, washed, stained with Hoechst 33342 (Thermo Fisher Scientific) and mounted for analysis by confocal microscopy. For selective permeabilization of PFA-fixed cells fixed on glass cove confocal microscopy was performed on a Zeiss LSM confocal microscope. Images were taken with a Plan-APOCHROMAT 63×/1.4

Oil objective lens with a pinhole setting of 1.0 Airy unit. Image processing was performed using FIJI (Schindelin et al., 2012).

Antibodies

The following primary antibodies were used: mouse monoclonal anti-FLAG (M2, Sigma; WB and IF 1:1000), mouse monoclonal anti-HA (BioLegend; WB: 1:1000) mouse monoclonal anti-β-Actin (Sigma; WB 1:4000). The following secondary antibodies were used: Donkey-anti-mouse IgG –HRP conjugated (Dianova; WB 1:10000), Donkey-anti-mouse IgG (Alexa Fluor 488, Invitrogen; IF 1:2000).

Mass spectrometry-based mapping of SPP-cleavage site

Hek293T cells were transiently transfected with GFP-TEV-FLAG-HO1-TMD fusion construct and SPP-HA and treated 24 h post transfection with 0.5 μM epoxomicin for 16 h. Subsequently, cells were detached in cold PBS-EDTA and resuspended in hypotonic buffer (10 mM HEPES-KOH pH 7.4, 1.5 mM $MgCl_2$, 10 mM KCl, 0.5 mM DTT, 10 μg/ml phenylmethylsulfonyl fluoride (PMSF), and each of 10 μg/ml chymostatin, leupeptin, antipain and pepstatin). After 10 min incubation on ice, cells were lysed by passing five times through a 27-gauge needle. Cell lysates were cleared by centrifugation at 5,000 g for 5 min and 20,000 g for 30 min at 4°C and the obtained cytosolic fraction was pre-absorbed for 1 h on BSA-coupled sepharose beads. Anti-GFP immunoprecipitation was performed for 30 min at room temperature using a GFP-specific single chain antibody fragment coupled to sepharose beads as described (Fleig et al., 2012). Immunoprecipitates were washed three times in buffer WB (50 mM HEPES-KOH, pH 7.4, 150 mM NaCl, 2 mM $MgOAc_2$, 10% glycerol, 1 mM EGTA, 0.1% Triton X-100) followed by washing once in buffer TDB (50 mM Tris-HCl, pH8.0, 0.5 mM EDTA, 1 mM DTT, 5 μg/ml PMSF, 10% glycerol, 0.1% Triton X-100). Subsequently immunoprecipitated proteins were digested by adding 3 μl recombinant TEV protease to the beads suspended in buffer TDB over night at 20°C on a rotary shaker. Supernatant and buffer from three wash steps with PBS were pooled and eluted GFP-tagged peptides were first immunoabsorbed with GFP-specific single chain antibody beads, followed by immunoprecipitation of FLAG-tagged peptides with FLAG M2 agarose (Sigma). Agarose was washed two times with PBS followed by two times with ddH_2O. Peptides were eluted from the beads with 15 ul 1% TFA/10% acetonitrile. 5ul were analyzed by a nanoHPLC system (Ultimate RSLC 3000) coupled to an ESI QExactive mass spectrometer (Thermo Fisher). The

sample was loaded on a self-packed analytical column (75um x 250mm, Reprosil-Pur C-18-AQ 1.9um (Maisch GmbH)) and eluted with a flow rate of 300nl/min in a 60 min gradient of 3% to 40% buffer B (0.1% formic acid, acetonitrile). Buffer A was 0.1% formic acid. One survey scan (res: 120.000) was followed by 10 information dependent product ion scans.

Mass spectrometry and DHX experiments

Peptides were dissolved in 80% (v/v) d_1-HFIP/20% D_2O at a concentration of 300 μM and incubated at 37°C for 7 days, solvent was removed by vacuum centrifugation, and the pellet was re-dissolved in deuterated solvent (80% (v/v) d_1-TFE, 2 mM ND_4-acetate). This resulted in > 95% deuteration as determined by ESI-MS. Deuterated dithiothreitol (DTT) was prepared by repeatedly dissolving DTT in D_2O and lyophilizing overnight; the deuterated DTT was finally dissolved in D_2O. Deuterated DTT was added at 1 mM to prevent cysteine oxidation in the deuterated peptides. For measurements of overall DHX, the deuterated peptides were diluted 1:20 from a 100 μM stock solution into 80% (v/v) TFE, 2 mM NH_4-acetate, 1 mM tris(2-carboxyethyl)phosphine (TCEP), pH 5.0, and incubated for different time periods at 20.0°C after which DHX was quenched on ice and by adding (0.5% (v/v) formic acid (pH ≈2.5). For electron transfer dissociation (ETD), 300 μM stock solutions were diluted 1:20. Mass/charge (m/z) ratios were recorded after the indicated time periods by electron-spray-ionization mass spectrometry (ESI-MS) as described (Poschner et al., 2009) using a Synapt G2 HDMS (Waters Co.) with one scan/second and evaluated as described (Stelzer et al., 2008). For ETD-DHX, incubation times from 1 min to 3 days at pH 5.0 were chosen. An incubation time of 0.1 min at pH 5 was simulated by incubation for 1 min at pH 4 and an incubation time of 30 days at pH 5 was simulated by incubating for22 h at pH 6.5. ETD was conducted as described (Stelzer et al., 2016) at a flow rate of 3 μl/min using a cooled syringe. ETD was started by decreasing the wave height from 1.5 V to 0.2 V. Hydrogen scrambling was < 5-10%, as determined by the ammonia loss method (Rand et al., 2012). All experiments were done at least in triplicate.

ETD fragment spectra recorded over 10 min were combined and evaluated using the module *ETD Fragment Analyzer* of the software suite MassMap® (MassMap GmbH & Co. KG, Freising, Germany) that is based on GRAMS/AI (Thermo Fisher Scientific). The theory and the detailed evaluation procedure are described in the section "*Analysis of amide-exchange/ETD Data using the ETD Fragment Analyzer*" (see below). Briefly, the evaluation consisted of the following steps:

1) Determination of the c- and z-fragment ions to be included in the analysis as well as of the 'extra hydrogens' of the fragment ions from the ETD spectrum of a non-deuterated sample. 'Extra hydrogens' result from hydrogen transfer during fragmentation or from neutralization of precursor ion proton by electron transfer (Lermyte et al., 2015).

2) After DHX and ETD, the expected isotopic patterns of the fragment ions (determined in step 1) were calculated by considering the rapid exchange of labile deuterons linked to side-chain heteroatoms or the N- and C- terminus, charge carrying deuterons, and extra deuterons (determined in step 1).

3) The calculated isotopic patterns were compared to the experimental ones by determining the extents of amide deuteration.

4) To increase the precision, a smoothing and interpolation procedure was applied to the mean numbers of deuterons as a function of the fragment numbers by fitting second order polynomials to four consecutive data points. The extent of amide deuteration was then determined from the smoothed and interpolated mean deuteron numbers of fragment ions.

5) The numbers of amide deuterons as a function of exchange period D(t) as obtained in step 4) were used to compute the residue-specific first order exchange rate constant k_{exp} using Eq. (1) (which considers 5% remaining deuterated solvent)

$$D(t) = 0.95 * e^{(-k_{exp}*t)} + 0.05 \qquad (1)$$

6) The equilibrium constants K_{exp} were calculated from k_{exp} and the sequence specific chemical exchange rate constants k_{ch} based on Eq. (2) based on Linderstrom-Lang theory, assuming EX2 conditions and a predominant folded state (Skinner et al., 2012)

$$k_{exp} = (k_{op} \times k_{rc})/(k_{rc} + k_{op} + k_{cl}) \qquad (2)$$

where k_{op} and k_{cl} are the rate constants of H-bond opening and closing, respectively, and k_{rc} represents the chemical rate constant that was calculated by http://landing.foxchase.org/research/labs/roder/sphere/ (under the set conditions: D-to-H-exchange, reduced Cys, pH =5.0, T = 20.0°C).

7) The free energies ΔG required for H-bond opening were then calculated according to Eq. (3)

$$\Delta G = - RT \ln(k_{exp} / (k_{ch} - k_{exp})) \qquad (3)$$

It should be noted, that the ΔΔG values obtained with this procedure are an upper estimate of the true values since (i) the molarity of water in 80% (v/v) TFE solvent is only 20% of the bulk molarity used for the determination of the reference chemical exchange rates k_{ch}, and (ii) the hydration of residues in the hydrophobic core of a TMD is possibly reduced relative to bulk. Both factors likely reduce the chemical

exchange rate in our experiments. In addition, TFE might have an impact on the auto-ionization constant of water and the chemical exchange rate constants (Stelzer et al., 2016).

Analysis of amide-exchange/ETD Data using the ETD FRAGMENT ANALYZER
1. Definition of Terms and Abbreviations

Term	Meaning
Asymptotic D number	Relative deuteron abundance of exchanging hydrogens that is asymptotically approached for very long exchange periods. The limit equals the ratio of deuterated to hydrogenated solutions employed in exchange.
H/D exchange	Experiment where H is exchanged by D
D/H exchange	Experiment where D is exchanged by H
D number	For a fragment ion: mean number of deuterons occupying the locations of slowly exchanging amide hydrogens For an amide hydrogen: relative deuteron abundance of the amide hydrogen
Extra hydrogens	Hydrogens that either result from the transfer of hydrogens as part of the fragmentation process or result from the neutralization of charge carrying hydrogens of the precursor ions by means of electron transfer
Fast exchange	Exchange observed for exchangeable hydrogens other than amide hydrogens. Fast exchange is observed for four types of hydrogens: hydrogens linked to heteroatoms of the amino acid side chains, hydrogens linked to the heteroatoms of the functional groups of the N and C terminus, charge carrying hydrogens, extra hydrogens (for definition see above).
Isotopic number	Number counting the isotopic peaks of an isotopic cluster starting with 0 for the monoisotopic peak. The number is incremented by 1 for the addition of a single nominal mass unit (^{12}C replaced by ^{13}C, addition of a single extra hydrogen, ^{1}H replaced by ^{2}H).
Slow exchange	Exchange observed for amide hydrogens and slow enough to be followed experimentally

Abbreviation	Meaning
FWHM	Full width at half maximum height/intensity
i	Isotopic number
i_{max}	Maximum isotopic number considered for fragments including extra hydrogens
i_{limit}	Maximum isotopic number considered for the calculated isotopic pattern of a neutral fragment without extra hydrogens. The limitation of the isotopic numbers is employed in order to avoid the consideration of isotopic peaks whose intensity is too low to be of practical relevance.
I	Expected isotopic distribution of a neutral fragment without extra hydrogens
I_{xxx}, $I^{(k)}$	Different isotopic distributions employed for the analysis of experimental isotopic patterns (xxx = +j, fit, …; k = 1,2)
m_{av}	Average mass of a neutral fragment without extra hydrogens
m_{mi}	Monoisotopic mass of a neutral fragment without extra hydrogens
z	Charge number of a fragment
$z_{precursor}$	Charge number of the precursor ion

2.   *Definition of an amide-exchange/ETD project*

The pieces of information and the data needed for an amide-exchange/ETD project are as follows:

- Information sufficient for the calculation of the expected full width at half maximum intensity (FWHM) of isotopic peaks (type of mass analyzer, resolution function).

- Amino acid sequence of the peptide.

- If applicable, status of Cysteine residues (Cysteine residues reduced or alkylated by means of one of the common alkylation reagents).

- For all modified amino acids, positions and type of modification, e.g. acetylation of the N-terminus or amidation of the C-terminus.

- Type of the exchange experiment, i.e. D exchanged by H (below referred to as D/H exchange) or H exchanged by D (H/D exchange).

- Asymptotic D number depending on the ratio of deuterated to hydrogenated solutions employed in exchange.

In the case of a D/H exchange experiment, a fully (>95%) deuterated peptide dissolved in a fully deuterated medium ('starting solution') is mixed with a fully hydrogenated solution ('dilution solution'). If the volume of the starting solution and the volume of the dilution solution are abbreviated by V_D and V_H, respectively, the asymptotic D number is calculated as the quotient $V_D/(V_D+V_H)$. For H/D exchange, the asymptotic D number is calculated in exactly the same way. The only difference is the meaning of the two volume variables: V_H and V_D then stand for the volume of the starting solution (fully hydrogenated medium containing the peptide) and the volume of the dilution solution (fully deuterated medium without the peptide), respectively.

The term 'asymptotic D number' is based on the expectation that the degree of deuteration of the slowly exchanging amide hydrogens will asymptotically approach that number for very long exchange times. For the calculations of the isotopic distributions of the peptide and fragment ion clusters, it is assumed that the degree of deuteration of all fast exchanging hydrogens equals the asymptotic D number. The term 'fast exchanging hydrogens' subsumes all the non-amide exchanging hydrogens, e.g. the side chain hydrogens linked to heteroatoms.

In the formulas shown below, the asymptotic D number is abbreviated by $D_{asymptotic}$.

- Charge number of the precursor ion ($z_{precursor}$).

- Data sets with fragment scans acquired for different samples under exactly the same conditions, i.e. with exactly the same solvents and gases as well as with the same parameters of the ion source and the mass analyzer.

As part of the evaluation, sum mass spectra are calculated using either all the scans or the scans of a user-selected scan interval.

One of the samples, below referred to as 'non-deuterated', is the fully hydrogenated peptide dissolved in the fully hydrogenated medium. The other samples ('exchange samples') are mixtures of the starting solution and the dilution solution. The mixtures only differ by the exchange period, i.e. the length of the time interval ranging from the time of mixing to the time of mass spectrometric analysis.

3. *Evaluation of amide-exchange/ETD Project*

The evaluation is implemented as a particular module of the MassMap® software suite called *ETD Fragment Analyzer*. The module was designed in order to perform as many tasks as possible in an automated manner and, at the same time, to allow for full user control of all relevant decisions.

The evaluation of a single amide-exchange/ETD project consists of several steps that are described in the following sections.

3.1. *Evaluation of the non-deuterated sample*

The first step is the evaluation of the sum mass spectrum of the non-deuterated sample. It is the aim of this step to identify the fragment clusters and to determine the number of neutral hydrogens of the fragments. Neutral hydrogens either result from the transfer of hydrogens as part of the fragmentation process or result from the neutralization of charge carrying protons of the precursor ions by means of electron transfer. These neutral hydrogens will be referred to as 'extra hydrogens' below. The very first decision is made by the user who defines the types of fragment ions (x, x*, x° with x=a, b, c, x, z, y; * and °: ions with neutral loss of ammonia or water, respectively) to be investigated. For the charge numbers (z) that are also selected by the user, the program automatically analyzes the isotopic patterns of all possible fragment clusters. The analysis of the isotopic patterns is based on the count sums over the extraction intervals of the isotopic peaks. The isotopic numbers (i) of the peaks range from 0 (isotopic number of the monoisotopic peak of the fragment without extra hydrogen) to i_{max}. To determine i_{max}, the normalized isotopic pattern or distribution I of the fragment is approximated by a binomial distribution characterized by the effective number n_C of carbon atoms. The number n_C is calculated by means of the monoisotopic mass m_{mi} and the average mass m_{av} of the fragment in a way that $n_C \cdot (mc_{13} - mc_{12}) \cdot a_{13}$ equals the difference $m_{av} - m_{mi}$. The quantities mc_{13}, mc_{12} and a_{13} stand for the mass of ^{13}C, the mass of ^{12}C and the natural abundance of ^{13}C amounting to 0.011, respectively. Hence, the elements I(i) of the isotopic distribution I are calculated according to equation (1):

$$(1) \qquad I(i) = \frac{(n_C)!}{(n_C - i)! \cdot i!} \cdot (a_{13})^i \cdot (1 - a_{13})^{n_C - i}, \quad i = 0, 1, ..., n_C \,.$$

Only the peaks with isotopic numbers ranging from 0 to i_{limit} are considered. The upper limit of the isotopic numbers i_{limit} is the smallest natural number meeting the following criterion:

$$(1a) \qquad \sum_{i=0}^{i_{limit}} I(i) \geq 0.999 \,.$$

In order to take the extra hydrogens into account that are caused by the fragmentation process itself or by neutralized charge carrying protons of the precursor ion, the largest isotopic number i_{max} for which the count sum is calculated is determined by equation (1b):

$$(1b) \qquad i_{max} = i_{limit} + z_{precursor} + 2 \,.$$

The limits $mz_{lower,\,i}$ and $mz_{upper,\,i}$ of the extraction interval of an isotopic peak with isotopic number i is based on the FWHM of the isotopic peak defined by the settings of the mass analyzer. In the following, the center m/z of the isotopic peak is termed mz_i. The quantities mz_i, $mz_{lower,\,i}$ and $mz_{upper,\,i}$ are calculated in the following way:

$$(2a) \qquad mz_i = \left(m_{mi} + z \cdot m_{H^+} + i \cdot (m_{13} - m_{12}) \right) / z \,,$$
$$(2b) \qquad mz_{lower,i} = mz_i - \alpha \cdot FWHM \,,$$
$$(2c) \qquad mz_{upper,i} = mz_i + \alpha \cdot FWHM \,.$$

By default, the value of factor α amounts to 1. The value may be slightly changed by the user in order to avoid cross-talk from interfering peaks or in order to make sure that all the relevant counts of the isotopic peaks of the clusters are taken into account.

The count sum C(i) of the isotopic peak with isotopic number i is calculated as the sum of the intensities of the data points of the sum mass spectrum of the non-deuterated sample, whose m/z values are within the interval $[mz_{lower,\,i}\,;\,mz_{upper,\,i}]$.

In cases of a calibration error of the mass spectrometer, the user may select two peaks of the sum mass spectrum that are used for a linear recalibration of the m/z axis of the mass spectrum. The recalibration option is not restricted to the mass spectrum of the non-deuterated sample. Rather, it is available for all the sum mass spectra of an amide-exchange/ETD project.

As the last step of the evaluation of fragment clusters of the non-deuterated sample, multiple linear regression is performed in order to determine the distribution of extra hydrogens of the clusters. For that purpose, the contributions of the unchanged isotopic distribution I as calculated by equation (1) (corresponding to fragment ions without extra hydrogens) and of the shifted isotopic distributions I_{+j} with j=1, 2, …, $z_{precursor}+2$ are determined. The shifted distribution I_{+j} corresponds to the expected distribution of the fragment cluster with j extra hydrogens:

$$\text{(3a)} \quad I_{+j}(i) = \begin{cases} 0, & \text{if } (i < j) \lor (i > j + i_{\text{limit}}) \\ I(i-j), & \text{if } j \le i \le j + i_{\text{limit}} \end{cases},$$

with $i = 0, 1, ..., i_{\text{max}}$ and $j = 1, 2, ..., z_{\text{precursor}} + 2$.

To determine the contributions of the different distributions to the experimental distribution C (composed of the count sums C(i) with i=0, 1, ..., i_{max}), the coefficients c_j (j=0, 1, ..., $y_{\text{precursor}}$+2; I_{+0} = I) and the normalization factor a are calculated by minimizing the quantity χ^2 of equation (3b):

$$\text{(3b)} \quad \chi^2 = \sum_{i=0}^{i_{\text{max}}} \left(C(i) - a \cdot I_{\text{fit}}(i) \right)^2 \quad \text{with } I_{\text{fit}}(i) := \sum_{j=0}^{z_{\text{precursor}}+2} \left(c_j \cdot I_{+j}(i) \right) \text{ and } \sum_{j=0}^{z_{\text{precursor}}+2} c_j = 1.$$

I_{fit} denotes the normalized isotopic distribution fitted to the experimental distribution. The fitting procedure defined by equations (3a) and (3b) is performed automatically as soon as a cluster is activated. In order to eliminate mass spectrometric peaks that are most probably part of other clusters, peaks with mass deviations above a user-defined limit or with too high or too low intensities are excluded from the fit by means of an iterative procedure. The final decision, which peaks are to be included in the fit, is made by the user. In most cases, intense clusters well above the limit of detection do not need user action, whereas for clusters around that limit user action is frequently needed in order to get valid results.

3.2. Evaluation of the exchange samples

The second step is the evaluation of the sum mass spectra of the samples having undergone exchange. In principle, the procedure applied is the same as the procedure employed for the analysis of the clusters of the sum mass spectrum of the non-deuterated sample. The differences between spectra with and without prior exchange are due to the fact that quickly exchanging hydrogens as well as slowly exchanging hydrogens have to be taken into account. Fast exchange, i.e. exchange too fast to be followed experimentally, is to be considered for the extra hydrogens (for definition see the previous section), for the side chain hydrogens linked to heteroatoms, for the carboxyl and the amino hydrogens located at the N or C termini as well as for the charge carrying protons. The slowly exchanging hydrogens are non-hydrogen bonded amide hydrogens of the peptide backbone. The isotopic distribution of a cluster of an exchange sample is calculated from the isotopic distribution I_{fit} of the respective cluster of the non-deuterated sample by a series of successive convolution steps.

The first convolution steps are due to the fast exchange of the extra hydrogens. The extra hydrogens of the ion species with extra hydrogens are assumed to be fast exchanging. For these hydrogens, the probability for the incorporation of a D instead

of an H is assumed to be equal to the asymptotic D number $D_{asymptotic}$. Hence, the additional distribution to be considered for the ion species with j extra hydrogens is a binomial distribution denoted by $I_{extra,j}$:

$$(4a) \quad I_{extra,j}(i) = \frac{(j)!}{(j-i)! \cdot i!} \cdot (D_{asymptotic})^i \cdot (1 - D_{asymptotic})^{j-i}, \quad i = 0, 1, ..., j.$$

As a consequence, the fast exchanging extra hydrogens will change the distribution I_{fit} into the distribution $I^{(1)}$ that is calculated in the following way:

$$(4b) \quad I^{(1)} = \sum_{j=0}^{z_{precursor}+2} c_j \cdot \left(I_{+j} * I_{extra,j} \right).$$

The notation f*g stands for the convolution of the two discrete distributions f and g. If f and g have n_f+1 and n_g+1 elements with indices ranging from 0 to n_f and n_g, respectively, h=f*g constitutes a distribution with the $(n_f + n_g +1)$ elements h(i):

$$(5) \quad h(i) = \begin{cases} \sum_{m=0}^{i} f(i-m) \cdot g(m) & \text{for} & i \le n_g \\ \sum_{m=0}^{n_g} f(i-m) \cdot g(m) & \text{for} & n_g < i \le n_f \\ \sum_{m=i-n_f}^{n_g} f(i-m) \cdot g(m) & \text{for} & n_f < i \end{cases}$$

$$\text{with} \quad i = 0, 1, 2, ..., n_f + n_g .$$

In addition to the extra hydrogens that change the expected isotopic distribution from I_{fit} to $I^{(1)}$, the fast exchanging charge carrying protons and the fast exchanging hydrogens covalently linked to heteroatoms have to be considered.

If the number of these hydrogens is denoted by n_{fast_plus}, the distribution $I^{(2)}$, i.e. the distribution expected in the case of lacking slow exchange, is the convolution of the binomial distribution I_{fast_plus} of the n_{fast_plus} fast exchanging hydrogens with the distribution $I^{(1)}$ of equation (4b):

$$(6a) \quad I^{(2)} = I^{(1)} * I_{fast\_plus} ,$$

$$(6b) \quad I_{fast\_plus}(i) = \frac{(n_{fast\_plus})!}{(n_{fast\_plus} - i)! \cdot i!} \cdot (D_{asymptotic})^i \cdot (1 - D_{asymptotic})^{n_{fast\_plus} - i}, \quad i = 0, 1, ..., n_{fast\_plus} .$$

The final distribution that is to be compared with the experimental distribution C found in the mass spectrum of the exchange sample is the convolution of distribution $I^{(2)}$ with the distribution of the slowly exchanging hydrogens.

From experience, for a significant fraction of the experimental clusters, satisfying fits are only obtained by employing a superposition of two binomial distributions, i.e. a

bimodal binomial distribution for the distribution of the n_{slow} slowly exchanging hydrogens. Apart from n_{slow}, such a bimodal distribution is characterized by three numbers, namely the fraction f_α of the first of the partial distributions and the exchange probabilities α and β of the first and the second partial distribution, respectively. Thus, the calculation of the distribution I_{slow} of the n_{slow} slowly exchanging hydrogens is performed in the following way:

(7a) $\quad I_{slow} = f_\alpha \cdot I_\alpha + (1 - f_\alpha) \cdot I_\beta$ with

(7b) $\quad I_\alpha(i) = \dfrac{(n_{slow})!}{(n_{slow} - i)! \cdot i!} \cdot \alpha^i \cdot (1 - \alpha)^{n_{slow} - i}, \ \ i = 0, 1, ..., n_{slow}$ and

(7c) $\quad I_\beta(i) = \dfrac{(n_{slow})!}{(n_{slow} - i)! \cdot i!} \cdot \beta^i \cdot (1 - \beta)^{n_{slow} - i}, \ \ i = 0, 1, ..., n_{slow}$.

The convolution of the distribution $I^{(2)}$ and the distribution I_{slow} results in the distribution $I^{(3)}$ that is fitted to the distribution C made up by the count sums of the isotopic peaks of the fragment cluster of interest. The numbers f_α, α and β, all ranging between 0 and 1, as well as a normalization factor (defined as in equation (3b)) are obtained by a non-linear least squares fitting procedure. The mean number of deuterons $D_{cluster}$ occupying the positions of the slowly exchanging amide hydrogens of the cluster is calculated according to the following equation:

(8) $\quad D_{cluster} = n_{slow} \cdot \left(f_\alpha \cdot \alpha + (1 - f_\alpha) \cdot \beta \right).$

3.3. Determination of the D numbers of the slowly exchanging amide hydrogens

The third step consists in the determination of the D numbers of individual slowly exchanging amide hydrogens. For the calculation of the D number D_k of the amide hydrogen of amino acid residue k (k=1, 2 , …,n_{AS}; n_{AS}: number of amino acids of the peptide), suitable pairs of consecutive fragment ions of the same type are used. To be suitable for analysis, a given fragment ion (A) must contain a single additional amide hydrogen relative to another fragment ion (B). The D number D_k of that amide hydrogen is calculated as the difference $D_A - D_B$.

As an option, the D numbers of the clusters of a particular type of fragments can be smoothed and interpolated before they are used for the determination of the D numbers of the individual amide hydrogens. The smoothing and interpolation algorithm applied uses all quadruples of consecutive existing D numbers for the least squares fitting of polynomials of second degree. The D numbers calculated by the polynomials are used to replace the D numbers of the amide hydrogens that are calculated from the directly determined D numbers of the fragments. The smoothed D number of a particular amide hydrogen is the weighted average of all the

polynomial D numbers of the particular amide hydrogen. For a polynomial D number to be considered, the amino acid numbers used for the polynomial fit must include the particular amino acid number. The weighting factors decrease rapidly with the distance of the amino acid considered from the amino acid of the polynomial fit closest to the amino acid considered. It should be mentioned that the smoothing algorithm contains a routine for outlier detection.

The rationale of the smoothing and interpolation procedure described in the previous paragraph is the evidence that the D numbers of the fragments of a certain type constitute a smooth function of the fragment number. Due to the smoothing procedure, the precision of the finally employed D numbers of the amide hydrogens is significantly improved. In the case of an amide hydrogen whose D number cannot be determined directly because one of needed fragments is lacking or not suitable for analysis, the D number of that fragment is replaced by the result of the smoothing process. We believe that such interpolation is justified to complement the data by individual missing values, except where the missing D number of the amide hydrogen corresponds to an amino acid with a chemical exchange constant that is strongly deviant from the numbers of its neighbors. In this case, the experimental series of exchange rates may exhibit a discontinuity that should not be neglected by smoothing.

4.   *Evaluation of the Exchange Kinetics*

4.1.   *Kinetic model*
The widely accepted kinetic model corresponds to the following scheme:

$$(9) \qquad F_D \underset{k_-}{\overset{k_+}{\rightleftharpoons}} U_D \underset{k_{H \to D}}{\overset{k_{D \to H}}{\rightleftharpoons}} U_H \underset{k_+}{\overset{k_-}{\rightleftharpoons}} F_H \quad \text{with}$$

$F_{D,H}$:   amide hydrogen deuterated, hydrogenated and folded state,

$U_{D,H}$:   amide hydrogen deuterated, hydrogenated and unfolded state,

$k_{+,-}$:   rate constants for conversions between folded and unfolded state,

$k_{D \to H}$: rate constant for hydrogenation of U_D,

$k_{H \to D}$: rate constant for deuteration of U_H.

If there is no isotopic effect for the hydrogenation of U_D and the deuteration of U_H, the rate constant $k_{H \to D}$ and the rate constant $k_{D \to H}$ will be proportional to the ratio of deuterated to hydrogenated exchange solvent and to the ratio of hydrogenated to deuterated exchange solvent, respectively. Hence, with the chemical exchange rate k_{ch} (exchange rate constant for fully deuterated exchange solvent) and with the above defined $D_{asymptotic}$, the rate constants can be expressed as follows:

(10a)　$k_{H \to D} = k_{ch} \cdot D_{asymptotic}$,

(10b)　$k_{D \to H} = k_{ch} \cdot \left(1 - D_{asymptotic}\right)$.

If the rate constants k_+ and k_- for the local unfolding and folding of the peptide are much larger than the rate constant k_{ch}, the influence of $k_{H \to D}$ and $k_{D \to H}$ on the interconversion of F_D and U_D can be neglected. This means that the concentration ratio of F_D and U_D can be approximated by the quotient k_-/k_+, i.e. by the value obtained for the situation where the middle part of scheme (9) is absent. The same approximation applies for the interconversion of the hydrogenated species F_H and U_H.

To facilitate the solution of the dynamic system underlying the reaction scheme (9), some additional quantities are defined:

(11)　f_D and u_D:　concentration of the peptide with the deuterated amide hydrogen in its folded and unfolded state, respectively

　　　　f_H and u_H:　concentration of the peptide with the hydrogenated amide hydrogen in its folded and unfolded state, respectively

　　　　$c_D := f_D + u_D$:　concentration of the peptide with the deuterated amide hydrogen, irrespective of the state of folding

　　　　$c_H := f_H + u_H$:　concentration of the peptide with the hydrogenated amide hydrogen, irrespective of the state of folding

　　　　$c_P := c_D + c_H$:　total concentration of the peptide, irrespective of the state of the amide hydrogen and the state of folding

With these quantities and the assumptions outlined above, the quantities u_D and u_H needed for the formulation of the differential equation resulting from the reaction scheme (9) can be expressed by the constant overall peptide concentration c_P and the time dependent concentration c_D of the peptide with the deuterated amide hydrogen:

(12a)　$c_D = u_D + f_D = u_D \cdot \left(1 + \dfrac{k_-}{k_+}\right)$　　$\to$　　(12b)　$u_D = c_D \cdot \left(1 + \dfrac{k_-}{k_+}\right)^{-1}$,

(12c)　$c_P - c_D = c_H = u_H + f_H = u_H \cdot \left(1 + \dfrac{k_-}{k_+}\right)$　　$\to$　　(12d)　$u_H = \left(c_P - c_D\right) \cdot \left(1 + \dfrac{k_-}{k_+}\right)^{-1}$.

Together with the equations (10a) and (10b) and the equations (12b) and (12d), the reaction scheme (9) leads to a first order differential equation for c_D:

(13) $\quad \dfrac{d}{dt} c_D(t) = \dfrac{d}{dt} u_D(t) = \left(-k_{D \to H} \cdot u_D + k_{H \to D} \cdot u_H \right) =$

$$= k_{ch} \cdot \left(-\left(1 - D_{asymptotic}\right) \cdot c_D \cdot \left(1 + \dfrac{k_-}{k_+}\right)^{-1} + D_{asymptotic} \cdot \left(c_P - c_D\right) \cdot \left(1 + \dfrac{k_-}{k_+}\right)^{-1} \right) =$$

$$= \left(1 + \dfrac{k_-}{k_+}\right)^{-1} \cdot k_{ch} \cdot \left(\left(D_{asymptotic} - 1\right) \cdot c_D + D_{asymptotic} \cdot \left(c_P - c_D\right)\right) =$$

$$= \left(\dfrac{k_+}{k_+ + k_-}\right) \cdot k_{ch} \cdot \left(D_{asymptotic} \cdot c_P - c_D\right) = k_{exchange} \cdot \left(D_{asymptotic} \cdot c_P - c_D\right) \text{ with}$$

(13a) $\quad k_{exchange} := \left(\dfrac{k_+}{k_+ + k_-}\right) \cdot k_{ch}.$

The general solution of equation (13) is the function $c_D(t)$ that contains a constant B needed for adjusting the initial conditions (D/H exchange: $c_D(0) = c_P$, H/D exchange: $c_D(0) = 0$):

(14a) $\quad c_D(t) = D_{asymptotic} \cdot c_P + C \cdot e^{-k_{exchange} \cdot t}$,

$\quad\quad$ D / H exchange: $\quad\quad c_D(0) = c_P \to \quad C = (1 - D_{asymptotic}) \cdot c_P \to$

(14b) $\quad\quad\quad\quad\quad\quad\quad\quad \dfrac{c_D(t)}{c_P} = D_{asymptotic} + (1 - D_{asymptotic}) \cdot e^{-k_{exchange} \cdot t}$,

$\quad\quad$ H / D exchange: $\quad\quad c_D(0) = 0 \to \quad C = -D_{asymptotic} \cdot c_P \to$

(14c) $\quad\quad\quad\quad\quad\quad\quad\quad \dfrac{c_D(t)}{c_P} = D_{asymptotic} - D_{asymptotic} \cdot e^{-k_{exchange} \cdot t}$.

The quantity $k_{exchange}$ equals the rate constant that is experimentally determined by fitting the time dependent D numbers of the amide hydrogens to the functions (14b) and (14c).

4.2. Analysis of the exchange kinetics

The final step of the overall analysis is the analysis of the exchange kinetics. It is assumed that exchange data are available for N exchange periods t_n (n=1, 2, …, N). To analyze the exchange kinetics of the amide hydrogen of amino acid m, the first step consists in the determination of the mean numbers $D_{mean}(m,n)$ of the D numbers D_m obtained as described in section 3.3 for all the fragment types analyzed for time point n. In order to make the final results less dependent on outliers, the calculation of the mean D numbers may include outlier detection. The sensitivity of the outlier

detection is decided by the user. The routine employed for outlier detection is based on the analysis of the variance of the numbers, for which the mean is calculated.

The model employed for the analysis of the kinetics is outlined in the preceding section and predicts a first order reaction that leads from the starting value D_{start} of the D number to the asymptotic D number $D_{asymptotic}$. The experimental rate constant $k_{exp,m}$ for the exchange of amide hydrogen m is determined by a non-linear least squares fitting routine that minimizes the quantity χ^2 of equation (15):

$$(15) \quad \chi^2 = \sum_{n=1}^{N} \left(\chi_{m,n} \right)^2 \text{ with } \chi_{m,n} := D_{mean}(m,n) - D_{fit}(m,n) \text{ and}$$

$$D_{fit}(m,n) := D_{asymptotic} + \left(D_{start} - D_{asymptotic} \right) \cdot \exp\left(-k_{exp,m} \cdot t_n \right).$$

For H/D and D/H exchange, the number D_{start} amounts to 0 and 1, respectively. In case some of the numbers $D_{mean}(m,n)$ are not available due to missing fragments for certain time points, the fitting procedure is restricted to the time points with available D numbers. For the fitting to be performed, the number of time points with D numbers must not be smaller than three.

To estimate the standard error of the decimal logarithm of the experimental rate constant $k_{exp,m}$, the standard deviation σ_m is calculated for the residuals $\chi_{m,n}$ obtained for the best fit value of the rate constant. The standard deviation σ_m is then used to perform the fitting defined in (15) for additional 2N sets of D numbers:

Set 1-1:	$D_{mean}(m,1) + \sigma_m$	$D_{mean}(m,2)$	...	$D_{mean}(m,N-1)$	$D_{mean}(m,N)$
Set 1-2:	$D_{mean}(m,1) - \sigma_m$	$D_{mean}(m,2)$	...	$D_{mean}(m,N-1)$	$D_{mean}(m,N)$
...	...	...	...	...	...
Set n-1:	$D_{mean}(m,1)$	...	$D_{mean}(m,n) + \sigma_m$	...	$D_{mean}(m,N)$
Set n-2:	$D_{mean}(m,1)$	...	$D_{mean}(m,n) - \sigma_m$	...	$D_{mean}(m,N)$
...	...	...	...	...	...
Set N-1:	$D_{mean}(m,1)$	$D_{mean}(m,2)$	...	$D_{mean}(m,N-1)$	$D_{mean}(m,N) + \sigma_m$
Set N-2:	$D_{mean}(m,1)$	$D_{mean}(m,2)$	...	$D_{mean}(m,N-1)$	$D_{mean}(m,N) - \sigma_m$

The uncertainty $\Delta_{m,n}$ of the decimal logarithm of the rate constant $k_{exp,m}$ that is caused by the uncertainty σ_m of the numbers $D_{mean}(m,n)$ is estimated as the maximum of the two numbers $|\log[k_{exp,m}]-\log[k(\text{Set n-1})]|$ and $|\log[k_{exp,m}]-\log[k(\text{Set n-2})]|$. The quantities k(Set n-1) and k(Set n-2) stand for the best fit rate constants obtained with the sets of D-numbers Set n-1 and Set n-2, respectively. By linear error propagation, the standard error $\Delta\log(k_{exp,m})$ of the decimal logarithm of the experimental rate constant $k_{exp,m}$ is calculated in the following way:

$$(16) \quad \Delta \log\left(k_{exp,m}\right) = \sum_{n=1}^{N} \Delta_{m,n}.$$

In cases where $k_{ch,m}$ is available, the experimental rate constant $k_{exp,m}$ can be used to calculate the difference ΔG_m of the Gibbs Free Energy associated with the equilibrium between the effectively folded and the effectively unfolded situation for amide hydrogen m. If the rate constant of the unfolding reaction of the peptide, which leads to an exchange competent state of amide hydrogen m, is denoted by $k_{m,+}$, and if the rate constant of the reverse reaction that makes the amide hydrogen m not exchange competent is denoted by $k_{m,-}$, ΔG_m is given by:

$$(17) \quad \Delta G_m = -RT \cdot \ln\left(K_m\right) = -RT \cdot \ln\left(\frac{k_{m,+}}{k_{m,-}}\right).$$

According to equation (13a), the fraction $k_{m,+}/(k_{m,+} + k_{m,-})$ can be determined from the experimental rate constant $k_{exp,m}$ and the intrinsic rate constant $k_{ch,m}$:

$$(18) \quad \frac{k_{m,+}}{k_{m,+} + k_{m,-}} = \frac{k_{exp,m}}{k_{ch,m}}.$$

Thus, the fraction of rate constants needed for the calculation of ΔG_m by means of equation (17) can be calculated from the experimental rate constant $k_{exp,m}$ and the intrinsic rate constant $k_{ch,m}$:

$$(19) \quad \frac{k_{exp,m}}{k_{ch,m}} = \frac{k_{m,+}}{k_{m,+} + k_{m,-}} = \frac{\dfrac{k_{m,+}}{k_{m,-}}}{\dfrac{k_{m,+}}{k_{m,-}} + 1} \quad \rightarrow \quad \frac{k_{m,+}}{k_{m,-}} \cdot \left(1 - \frac{k_{exp,m}}{k_{ch,m}}\right) = \frac{k_{exp,m}}{k_{ch,m}} \quad \rightarrow$$

$$\rightarrow \quad \frac{k_{m,+}}{k_{m,-}} = \frac{k_{exp,m}}{k_{ch,m} - k_{exp,m}}.$$

The limits of the standard confidence interval of ΔG_m are calculated by means of the standard error $\Delta\log(k_{exp,m})$ of the decimal logarithm of $k_{exp,m}$:

$$(20a) \quad \Delta G_{m,min} = -RT \cdot \ln\left(\frac{k_{+,m}}{k_{-,m}}\right)_{max} \quad \text{with} \quad \left(\frac{k_{+,m}}{k_{-,m}}\right)_{max} = \frac{k_{exp,m} \cdot 10^{\Delta\log\left(k_{exp,m}\right)}}{k_{ch,m} - k_{exp,m} \cdot 10^{\Delta\log\left(k_{exp,m}\right)}},$$

$$(20b) \quad \Delta G_{m,max} = -RT \cdot \ln\left(\frac{k_{+,m}}{k_{-,m}}\right)_{min} \quad \text{with} \quad \left(\frac{k_{+,m}}{k_{-,m}}\right)_{min} = \frac{k_{exp,m} \cdot 10^{-\Delta\log\left(k_{exp,m}\right)}}{k_{ch,m} - k_{exp,m} \cdot 10^{-\Delta\log\left(k_{exp,m}\right)}}.$$

A

Protein	Sequence	ΔG_{app}
XBP1u wt TM region	...$\overset{186}{\text{I}}$SPWILAVLTLQIQSLISCWAFWTT$\overset{211}{\text{W}}$...	**+0.07**
XBP1u mt1 TM region	...$\overset{186}{\text{I}}$**LL**WIL**L**VLTLQIQSLISCWAFWTT$\overset{211}{\text{W}}$...	**-2.26**
XBP1u mt2 TM region	...$\overset{186}{\text{I}}$SPWILAVLTL**LILLLIL**CWAFWTT$\overset{211}{\text{W}}$...	**-5.76**
Prolactin signal peptide	...$\overset{5}{\text{G}}$SSQKGSRLLLLLVVSNLLLCQGVV$\overset{30}{\text{S}}$...	**+0.35**
Calnexin signal peptide	$\overset{1}{\text{M}}$EGKWLLCMLLVLGTAIVE$\overset{20}{\text{A}}$...	**+0.74**
Calnexin TMD	...$\overset{479}{\text{E}}$RPWLWVVYILTVALPVFLVILFC$\overset{503}{\text{C}}$...	**-2.26**

B

SPase

XBP1u TMD wt $\overset{1}{\text{M}}$AQLSPLQNISPWILAVLTLQIQSLISCWA–$\overset{30}{\text{F}}$...

C

R232N

wt-RN   mt1-RN   mt2-RN

49.7%   80.8%   81.2%

Yücel *et al.* Figure 4

A

```
wt   ...ISPWILAVLTLQIQSLISCWAFWTTW...
mt6  ...ISPWILAVLTLQIGSLISCWAFWTTW...
mt7  ...ISPWILAVLTLQIQGLISCWAFWTTW...
```

Yücel *et al.* Figure 7

A

B

Yücel *et al.* Supplementary Figure 4

Yücel et al. Table S2. Numerical values of k_{exp} and DG, related to Figure 4.

amide	wt log k [min⁻¹]	wt ΔG [kcal/mol]	amide	mt1 log k [min⁻¹]	mt1 ΔG [kcal/mol]	amide	mt2 log k [min⁻¹]	mt2 ΔG [kcal/mol]	amide	mt3 log k [min⁻¹]	mt3 ΔG [kcal/mol]	amide	mt4 log k [min⁻¹]	mt4 ΔG [kcal/mol]	amide	mt5 log k [min⁻¹]	mt5 ΔG [kcal/mol]
K	n.d.[2]		•	n.d.		•	n.d.		•	n.d.		•	n.d.		•	n.d.	
K	n.d.		•	n.d.		•	n.d.		•	n.d.		•	n.d.		•	n.d.	
I186	n.d.		•	n.d.		•	n.d.		•	n.d.		•	n.d.		•	n.d.	
S187	n.d.		L	n.d.		•	n.d.		G	n.d.		•	n.d.		•	n.d.	
P188	n.d.		L	n.d.		•	n.d.		G	n.d.		•	n.d.		•	n.d.	
W189	n.d.		•	-1.58 ± 0.27	2.4 ± 0.4	•	n.d.		•	n.d.		•	n.d.		•	n.d.	
I190	-1.15 ± 0.08[3]	1.5 ± 0.1[4]	•	-2.73 ± 0.15	3.6 ± 0.2	•	-1.28 ± 0.05	1.6 ± 0.1	•	-0.87 ± 0.13	1.0 ± 0.2	•	-0.75 ± 0.16	0.8 ± 0.3	•	-1.01 ± 0.23	1.2 ± 0.4
L191	-1.65 ± 0.07	2.2 ± 0.1	•	-3.26 ± 0.08	4.4 ± 0.1	•	-1.58 ± 0.04	2.1 ± 0.1	•	-1.19 ± 0.06	1.5 ± 0.1	•	-0.83 ± 0.08	1.0 ± 0.1	•	-1.39 ± 0.11	1.8 ± 0.2
A192	-1.98 ± 0.16	3.4 ± 0.2	L	-3.57 ± 0.24	4.8 ± 0.3	•	-1.78 ± 0.08	3.2 ± 0.1	•	-1.58 ± 0.12	2.9 ± 0.2	•	-0.91 ± 0.07	2.0 ± 0.1	•	-1.76 ± 0.12	3.2 ± 0.2
V193	-2.52 ± 0.23	3.5 ± 0.3	•	-3.68 ± 0.13	4.8 ± 0.2	•	-2.33 ± 0.08	3.3 ± 0.2	•	-1.97 ± 0.11	2.8 ± 0.2	G	-0.95 ± 0.10	1.4 ± 0.1	•	-2.20 ± 0.18	3.1 ± 0.2
L194	-2.98 ± 0.17	4.1 ± 0.2	•	-3.89 ± 0.72	5.3 ± 1.0	•	-2.94 ± 0.08	4.1 ± 0.1	•	-2.53 ± 0.18	3.5 ± 0.2	G	-0.79 ± 0.12	2.3 ± 0.2	•	-2.68 ± 0.30	3.7 ± 0.4
T195	-2.93 ± 0.10	4.6 ± 0.1	•	-3.72 ± 0.26	5.7 ± 0.3	•	-3.05 ± 0.08	4.8 ± 0.1	•	-2.53 ± 0.17	4.1 ± 0.2	G	-0.53 ± 0.11	2.4 ± 0.2	•	-2.46 ± 0.19	4.0 ± 0.2
L196	-2.69 ± 0.15	4.2 ± 0.2	•	-3.47 ± 0.14	5.2 ± 0.2	•	-3.33 ± 0.26	5.0 ± 0.3	•	-2.32 ± 0.24	3.7 ± 0.3	•	-0.11 ± 0.13	0.5 ± 0.3	•	-2.53 ± 0.40	4.0 ± 0.5
Q197	-2.44 ± 0.08	4.1 ± 0.1	•	-2.90 ± 0.28	4.8 ± 0.4	L	-3.86 ± 0.46	5.2 ± 0.6	•	-2.28 ± 0.20	3.9 ± 0.3	•	-0.69 ± 0.07	1.8 ± 0.1	•	-1.87 ± 0.09	3.4 ± 0.1
I198	-2.48 ± 0.09	3.7 ± 0.1	•	-2.77 ± 0.10	4.1 ± 0.1	•	-4.57 ± 0.28	6.0 ± 0.4	•	-2.36 ± 0.17	3.5 ± 0.2	•	-1.28 ± 0.11	2.1 ± 0.1	•	-1.91 ± 0.09	2.9 ± 0.1
Q199	-2.40 ± 0.12	4.1 ± 0.2	•	-2.77 ± 0.20	4.6 ± 0.3	L	-4.57 ± 0.20	6.1 ± 0.3	•	n.d.		•	-1.41 ± 0.11	2.7 ± 0.1	G	-1.39 ± 0.16	3.0 ± 0.2
S200	-2.08 ± 0.11	4.6 ± 0.1	•	-2.33 ± 0.16	5.0 ± 0.2	L	-4.61 ± 0.19	6.2 ± 0.3	•	-2.15 ± 0.13	4.7 ± 0.2	•	-1.28 ± 0.13	3.6 ± 0.2	G	-1.32 ± 0.20	3.4 ± 0.3
L201	-1.98 ± 0.18	3.4 ± 0.2	•	-2.03 ± 0.18	3.4 ± 0.2	•	-4.72 ± 0.18	6.4 ± 0.2	•	-1.79 ± 0.37	3.1 ± 0.5	•	-1.48 ± 0.14	2.7 ± 0.2	•	-1.12 ± 0.12	2.0 ± 0.2
I202	n.d.		•	n.d.		•	-3.98 ± 0.53	5.2 ± 0.7	•	n.d.		•	-1.74 ± 0.20	2.2 ± 0.3	•	-1.07 ± 0.16	1.2 ± 0.2
S203	n.d.		•	n.d.		L	n.d.		•	-1.16 ± 0.26	2.8 ± 0.4	•	n.d.		•	-0.70 ± 0.27	2.2 ± 0.4
C204	n.d.		•	n.d.		•	n.d.		•	n.d.		•	n.d.		•	-0.54 ± 0.11	2.9 ± 0.2
W205	n.d.		•	n.d.		•	n.d.		•	n.d.		•	n.d.		•	n.d.	
A206	-0.58 ± 0.24	1.7 ± 0.3	•	n.d.		•	n.d.		•	-0.59 ± 0.25	1.7 ± 0.4	•	0.01 ± 0.14	0.8 ± 0.2	•	-0.17 ± 0.51	1.1 ± 0.9
F207	-0.49 ± 0.10	1.4 ± 0.1	•	n.d.		•	n.d.		•	-0.41 ± 0.19	1.2 ± 0.3	•	-0.04 ± 0.18	0.6 ± 0.4	•	n.d.	
K	-0.04 ± 0.28	1.1 ± 0.5	•	n.d.		•	n.d.		•	-0.29 ± 0.11	0.5 ± 0.2	•	n.d.		•	0.03 ± 0.10	0.9 ± 0.2
K	n.d.		•	n.d.		•	n.d.		•	n.d.		•	n.d.		•	n.d.	
K	n.d.		•	n.d.		•	n.d.		•	n.d.		•	n.d.		•	n.d.	

[1] Identity of the TM domain peptide

[2] n.d. = not determined

[3] k_{exp} from exponential fit of means ± SE

[4] ΔG with confidence interval