Genomic insights into metabolism and phylogeography of the mesophilic

Thermotogae genus *Mesotoga*

Camilla L. Nesbø[1,2] , Rhianna Charchuk[1], Stephen M. J. Pollo[1], Karen Budwill[3], Ilya V.

Kublanov[4], Thomas H.A. Haverkamp[5,6] and Julia Foght[1]

1 Department of Biological Sciences, University of Alberta, Edmonton, AB, Canada

2 BioZone, Dept. of Chemical Engineering and Applied Chemistry, Wallberg Building,

University of Toronto, Toronto, ON, Canada.

3 InnoTech Alberta, 250 Karl Clark Road, Edmonton, Alberta, Canada T6N 1E4

4 Winogradsky Institute of Microbiology, Russian Academy of Sciences, Moscow,

Russia

5 Centre for Ecological and Evolutionary Synthesis, Department of Biosciences,

University of Oslo, Blindern, Oslo, Norway.

6 Norwegian Veterinary Institute, Oslo, Norway.

*Corresponding Authors: nesbo@ualberta.ca

Running title: Genome analysis of *Mesotoga*.

Key words: Thermotogae, subsurface, recombination, oil reservoir, phylogeny, sulfur
metabolism, hydrogenase, anaerobe.

Subject Category: Evolutionary Genetics

Abstract

The genus *Mesotoga*, the only described mesophilic Thermotogae lineage, is commonly

detected in low-temperature anaerobic hydrocarbon-rich environments. Besides

mesophily, *Mesotoga* displays lineage-specific phenotypes, such as no or little H_2

production, dependence on sulfur-compound reduction and ability to oxidize acetate,

which may influence its ecological role. We used comparative genomics of 18 *Mesotoga*

strains and a transcriptome of *M. prima* to investigate how life at moderate temperatures

affects phylogeography, and to interrogate the genomic features of its lineage-specific

metabolism. Phylogenetic analysis revealed three distinct *Mesotoga* lineages having

different geographic distributions patterns and high levels of intra-lineage recombination

but little gene-flow between lineages. All 16S rRNA genes from the genomes had > 99%

identity whereas average nucleotide identity among genomes was 90.6–98.6% within

groups and 80–86% between groups. Including data from metagenomes, phylogeographic

patterns reveled that geographical separation is more important for *Mesotoga* than their

thermophilic relatives, and we suggest its distribution is constrained by their strictly

anaerobic lifestyle. We propose that H_2 oxidation and thiosulfate reduction are

accomplished by a sulfide dehydrogenase and a hydrogenase-complex and that a

pyruvate:ferredoxin oxidoreductase acquired from *Clostridia* is responsible for oxidizing

acetate.

Introduction

The genus *Mesotoga* is the only characterized mesophilic lineage within the otherwise thermophilic bacterial phylum Thermotogae (Pollo *et al.*, 2015). *Mesotoga* spp. have been isolated from and detected in polluted marine sediments, low temperature oil reservoirs, and waste water treatment facilities (Nesbø *et al.*, 2010; 2012; Hania *et al.*, 2011; 2013), and are common in anaerobic methanogenic environments (Nesbø *et al.*, 2010) where they may be involved in syntrophic acetate degradation (Nobu *et al.*, 2015). The first described member of this genus, *Mesotoga prima* MesG1Ag4.2 (hereafter, *M. prima*), was isolated from a PCB-degrading enrichment culture inoculated with sediments from Baltimore Harbor, Maryland (USA) (Nesbø *et al.*, 2006; 2012). Sequencing the genomes of *M. prima* and the very closely related *M. prima* PhosAc3 (hereafter, PhosAc3) isolated in Tunisia (Hania *et al.*, 2015) revealed larger genomes than thermophilic Thermotogae, with more genes involved in regulatory functions and interactions with the environment (Zhaxybayeva *et al.*, 2012).

Genome size in Thermotogae inversely correlates with optimum growth temperature (Pollo *et al.*, 2015; Zhaxybayeva *et al.*, 2012). However, it is unclear how growth temperature affects other aspects of genome evolution including levels of homologous recombination. Hyperthermophilic *Thermotoga* display extremely high levels of homologous recombination, which could be a side effect of the need for DNA repair at high temperatures (Nesbø *et al.*, 2015). Nesbø *et al.* (2015) also found that *Thermotoga* spp. from similar environments have exchanged more genes than geographically close isolates from different environments. This has been observed for metagenomes in general, where geographical distance has little effect on gene sharing,

and gene composition is more strongly affected by ecology than proximity (Fondi *et al.*,

2016). Moreover, the phylogeography and patterns of gene flow among *Thermotoga*

genomes suggested that oil reservoirs were colonized from subsurface populations rather

than being buried with the sediments that mature into oil reservoirs (a corollary of the

paleosterilization hypothesis; Wilhelms *et al.*, 2001). Comparative genomic analyses of

mesophilic Thermotogae may shed light on the role of growth temperature on

recombination and phylogeography.

In addition to lower optimal growth temperature (37°C - 40°C), *Mesotoga*'s core

energy metabolism also differs from that of other characterized Thermotogae. For

instance, while growth of most Thermotogae is stimulated by adding sulfur-compounds

to the medium (Ravot *et al.*, 1995; Boileau *et al.*, 2016), reduction of sulfur compounds

appears to be essential for growth of *Mesotoga* in pure culture and they produce little or

no H_2 (Hania *et al.*, 2011; 2013; Fadhlaoui *et al.*, 2017).

Here we compare 18 *Mesotoga* genomes obtained from isolates and single cells

originating from six geographically different sites, including three low temperature

continental oil reservoirs, in order to elucidate genomic markers of metabolic differences

and to investigate how growth temperature influences phylogeography and prevalence of

recombination. We also compare our genomes to *Mesotoga* sequences available in

publicly available metagenomes. We infer that geographic separation has had a

significant influence on the phylogeography of *Mesotoga,* possibly due to selective

pressures of dispersal of strict anaerobes through aerobic environments. Finally, we

present a model that accounts for *Mesotoga*'s distinct sulfur-dependent metabolism

involving a hydrogenase complex.

Materials and Methods

Sources of genome sequences

Nine *Mesotoga* strains (BR, HF and BH designations) were isolated from oil reservoirs

and anaerobic sediments in Canada and the USA (Table 1). In addition, seven single cells

were physically selected from oil field fluids or oil sands enrichment cultures from

Canada or a contaminated aquifer in the USA (PW, NAPDC and TOLDC designations,

respectively) and amplified by PCR to produce single cell amplified genomes (SAGs).

Detailed descriptions of isolation procedures, DNA extraction, genome assembly and

annotation are provided in Supplementary Information.

To augment the strain genomes, 15 publicly available metagenomes containing

large numbers of *Mesotoga* spp. sequences were identified using blastn searches of IMG

(JGI; accessed February 2017) and SRA (NCBI; accessed December 2016) using *rpoB*

from *M. prima* as a probe and expected (exp.) set to $< e^{-50}$. For additional details on

search parameters and information on assembly of draft genomes from metagenomic

sequences or contigs see Supplementary Information.

Genome content and genome alignments

Shared genes were identified in IMG Version 4 (Markowitz *et al.*, 2014) using translated

proteins and 70% identity cut-off and exp. $< e^{-10}$, whereas 30% sequence identity cut-off

and exp. $< e^{-5}$ were used to identify strain-specific genes and for comparing *Mesotoga*

genomes to other Thermotogae genomes.

Pan-genome calculations were performed in Panseq (Laing *et al.*, 2010) using a

fragment size of 500 bp and 70% identity cutoff, and exp. $< e^{-10}$. The data matrices of

shared core single nucleotide polymorphisms (SNPs) and 500-bp fragments were

converted into uncorrected distances and visualized in SplitsTree 4 (Huson and Bryant,

2006) using NeighborNet clustering.

Whole genome alignments were carried out in MAUVE version 2.3.1 (Darling *et*

al., 2010) using automatically calculated seed weights and minimum Locally Collinear

Blocks (LCB) scores. LCB positions with gaps were removed and the edited LCB were

concatenated in Geneious v.10 (www.geneious.com). Average nucleotide identities

(ANI) were calculated at http://enve-omics.ce.gatech.edu/ani/ (Goris *et al.*, 2007).

Recombination detection

The relative rate of recombination to mutation within lineages, as well as the average

recombination tract length, were assessed using the LDhat package (McVean *et al.*, 2002;

Jolley, 2004) as described by Nesbø *et al.* (2015) on concatenated alignments (including

LCB > 10,000 bp) of genomes from the W and the A lineage separately. Recombinant

fragments between lineages were detected using LikeWind Version 1.0 (Archibald 2002)

on the concatenated MAUVE alignment (above), using a sliding window of 1000 bp with

100-bp increments.

RNAseq analysis

RNA isolation from a culture of *M. prima* (grown at 45ºC for 73 h in 0.5% yeast extract,

0.01 M thiosulfate and 0.5% xylose) and subsequent sequencing as one of five barcoded

libraries were performed as described by Pollo et al. (2017). RNAseq analysis was carried

out in CLC Genomics Workbench version 7.0.4 as described by Pollo *et al.*(2017).

H$_2$ and H$_2$S measurements

Standard gas chromatographic analysis of culture headspace gas was performed using an

Agilent CP4900 Micro Gas Chromatograph to detect H$_2$ production by the cultures, as

described in Supplementary Information. Dissolved sulfide concentrations were measured

using a VACUettes® Visual High Range Kit (Chemetrics), following the manufacturer's

recommendations.

Results and Discussion

Phylogenetic and genomic analyses of isolates and single cells reveal three distinct

Mesotoga **lineages**

We generated draft genomes for eight newly isolated *Mesotoga* strains from two Albertan

oil reservoirs (H and B), and one *Mesotoga* strain from a PCB-degrading enrichment

culture from Baltimore Harbor, Maryland (Table 1). Seven partial SAGs were obtained

from cells sorted from produced water from an Albertan oil reservoir (PW), a naphtha-

degrading enrichment culture inoculated with sediments from an Albertan oil sands

tailings pond (NAPDC), and a toluene-degrading enrichment culture inoculated with

sediments from a contaminated aquifer in Colorado (TOLDC). We also included in our

analyses the draft genome of PhosAc3, previously isolated in Tunisia (Hania *et al.*, 2015)

and the closed genome of *M. prima* (Nesbo *et al.*, 2012) from Baltimore Harbor.

The 16S rRNA genes of all 17 genomes had ≥99% identity to *M. prima*;

phylogenetic trees revealed three distinct lineages (Figure 1a). Genome networks based

on core SNPs also had topologies consistent with 16S rRNA gene phylogeny, with three

distinct lineages being evident (Figure 1b). Two lineages have a widespread geographical

distribution: the World lineage (W; found in all regions represented) and the lineage

found in locations in USA (US; found in Baltimore Harbor and Colorado). The Alberta

(A) lineage was observed in the Albertan samples only. Interestingly, *M. prima* has one

16S rRNA from the W lineage and one from the US lineage, suggesting one copy has

been acquired laterally.

Very little reticulate evolution was observed among the three *Mesotoga* lineages

(Figure 1b) and the A lineage in particular showed very little connection with the other

two groups, suggesting that the three lineages have evolved independently for a relatively

long time. The same pattern was observed for the pan-genome, with most lateral

connections occurring within groups (Figure 1c).

A high level of recombination was detected, with the majority (> 200) of

recombination events involving genomes from the same lineage (Figure S1). For the W

and A lineages, respectively, the average recombination tract length was estimated to be

36,000 – 56,000 bp and 17,000–23,000 bp; the population mutation rate (θ) was

estimated to be 0.022 and 0.013, and the population recombination rate (γ) to be 1.8 (1.5–

2.2) and 1.5 (1.3–1.7). The resulting high γ/θ ratios of ~82–115 indicate high levels of

recombination and are similar to estimates for *Thermotoga* spp. (Nesbø *et al.*, 2015).

Phylogenetic analysis identified 52 regions with recombination likely occurring

between lineages: 39 regions showed evidence of recombination between *Mesotoga* sp.

BH458 and the W lineage, eight regions suggested recombination between *Mesotoga* sp.

BH458 and the A lineage, and only five regions showed possible recombination between

A- and W-lineage genomes (Figure 2). The regions with recombination involving the A

lineage were relatively short (range 230–530 bp) and the sequences more divergent,

whereas several of the fragments involving the W lineage and *Mesotoga* sp. BH458 were

> 5 kb (average 3000 bp, range 260–20,000), suggesting that recombination events

between the W lineage and *Mesotoga* sp. BH458 are more recent than those involving the

A lineage. Very high levels of recombination were observed for a few genes. Among

these is Theba_0319 in *M. prima*, the fourth most highly expressed gene (Supplementary

Table S1) that encodes the OmpB protein (Petrus *et al.*, 2012), a major component of the

toga of Thermotogae. This high level of recombination is consistent with strong

diversifying selective pressure on surface proteins, commonly observed in pathogenic

bacteria (e.g., Wachter and Hill, 2016).

The observation of several recent recombination events between the W and US

lineages, which currently co-exist in at least one location (i.e., Baltimore Harbor),

demonstrate that recombination between lineages is possible. We therefore suggest that

the three *Mesotoga* lineages have evolved independently due to geographical, not genetic,

isolation. Although it may seem counterintuitive that mesophilic *Mesotoga* would be

more affected by geographical separation than hyperthermophilic *Thermotoga*, this may

be a consequence of their strictly anaerobic metabolism. Chakraborty *et al.* (2018)

showed that bacteria are dispersed out of deep hot subsurface oil reservoirs and into the

ocean through hydrocarbon seeps, and this might serve as a major route of migration

between these environments. Hyperthermophilic *Thermotoga* cells will be inactive in

cold aerobic ocean water, but mesophilic *Mesotoga* cells will be active at these

temperatures and therefore may quickly succumb to oxygen exposure, limiting viable

dispersal. In support of this observation, many *Mesotoga*-specific genes appear to be

involved in O_2 or H_2O_2 detoxification (see below).

***Mesotoga* lineages have conserved core genomes and diverse pan-genomes**

Comparative analysis of *Mesotoga* genomes revealed greater gene-content diversity than

was observed in hyperthermophilic Thermotogae (Nesbø *et al.*, 2015). The pan-genome

of the *Mesotoga* isolates and SAGs was estimated to be 7,452,537 bp with an accessory

genome of 5,664,475 bp; each genome contained a considerable amount of lineage-

specific DNA (Figure S1; see Supplementary Information for additional details of the

pan-genome and within-sample site diversity). In pairwise comparisons, the genomes

shared on average 76% of their genes (Supplementary Table S2). Genomes from each

lineage defined in Figure 1 shared, on average, 81% - 89% of their genes whereas in

pairwise comparisons *Thermotoga* spp. share > 90% of their genes (Nesbø *et al.*, 2015).

Comparing genomes from different lineages, the US lineage had an intermediate position,

sharing more genes with the A and W lineages: on average, genomes from A and W share

69% of genes, W and US share 73%, and A and US share 75% of their genes.

Comparing nucleotide divergence within the core genomes revealed 'species-level'

divergence among the three lineages detected (ANI < 86%), whereas ANI within the A

and W lineages was very high at 98.6% and 97.9%, respectively (Supplementary Table

S3). In comparison, ANI among *Thermotoga* genomes was 95.3% (Nesbø *et al.*, 2015).

Thus *Mesotoga* spp., particularly those from the W lineage, appears to have more

conserved core genomes and more diverse pan-genomes than their hyperthermophilic

relatives.

Nesbø *et al.* (2015) suggested that high levels of recombination may be

responsible for homogenizing the *Thermotoga* spp. genomes. However, since we

observed similar high levels of recombination within the *Mesotoga* W and A lineages,

additional forces must be responsible for the differences in pan-genome sizes. Perhaps

more cryptic niches are available in low-temperature than in high-temperature subsurface

environments (McInerney *et al.*, 2017), or *Mesotoga* may have larger effective population

sizes *in situ* than the hyperthermophiles (Andreani *et al.*, 2017). *Mesotoga* spp. might also

interact with viruses more frequently, which may drive population diversity (Koskella

and Brockhurst, 2014).

Phylogeographic patterns of the three *Mesotoga* lineages: isolation by distance,

range expansion and burial with isolation

To challenge our interim conclusions about genome structure and geographic distribution,

we expanded the *Mesotoga* sequence dataset by searching IMG/M (in JGI) and SRA (in

NCBI) databases for metagenomes containing *Mesotoga* spp. sequences. Fifteen

metagenomes containing sequences closely related to the *Mesotoga* genomes investigated

here were identified, arising from two environments already described (tailings pond and

oil reservoir in Alberta), as well as oil reservoirs, contaminated sediments, wastewaters

and hotspring sediments across the continental USA, and wastewaters in China (Table 2

and Supplementary Information).

Recent range expansion of the W-lineage: Blastn comparisons confirmed wide

distribution of the *Mesotoga* W lineage in wastewater treatment systems. A network

including *Mesotoga* contigs from three metagenomes dominated by W lineage sequences

(Long Beach, Boston and Hong Kong) revealed no geographical structuring, which,

together with high identity in core sequences (Figure S3a), suggests a recent range

expansion of this lineage (Choudoir *et al.*, 2017). Interestingly, W lineage *Mesotoga* have

been detected only at sites heavily influenced by human activities (e.g., drilling,

contamination), suggesting an anthropogenic role in their dispersal. One of the genes

specific to the W lineage (Theba_0620, Supplementary Information) is involved in

synthesis of poly-gamma-glutamate, which has been implicated in cell survival under

harsh conditions and may have contributed to this lineage's wide distribution.

Isolation by distance can explain the distribution of US-genomes: The

metagenome data greatly expanded the observed distribution of the US lineage. As

expected, metagenome IMG 15764 from Albertan oil reservoir E (the source of *Mesotoga*

sp. SC_PW1-3) contained sequences with high identity to the A lineage. However, it also

contained many sequences with high identity to the US lineage (Table 2), and sequence

binning yielded two *Mesotoga* metagenome-assembled genomes (MAGs): one most

similar to NA-genomes (Figure S3b) and one with a mix of sequences from the A lineage

and US lineage (not shown). The intermediate position of US between the A- and W-

lineages noted above may therefore be due to this lineage co-existing with both the W

and A lineages.

The network of NA-*Mesotoga* including the metagenomes in Table 2 revealed

three groups (Figure S3b) where metagenomes from New York and Blank Spring

(California) form a cluster that does not contain any genomes sequenced by us. The

remaining metagenomes clustered according to geography as well as by environment

type: the MAG assembled from oil reservoir E (Alberta), two MAGs from an Alaskan oil

reservoir, and the *Mesotoga* sequences from Alameda (California) clustered with

SC_TOLDC from Colorado (western North America), while the *Mesotoga* sequences

from New Jersey clustered with *Mesotoga* sp. BH458 from Baltimore Harbor (eastern

North America). We therefore suggest that the divergence patterns seen for this lineage

can be explained at least partly by an isolation-by-distance model.

Evolution of the A-lineage in isolation in North-American oil reservoirs: The

metagenome sequences revealed that the A lineage is not restricted to Alberta, nor is it

specific to oil reservoirs (Table 2), with substantial numbers of A-lineage sequences

detected in wastewater metagenomes. For this lineage, MAGs were available from the

same oil reservoir in Alaska as where we observed the NA-lineage (Hu *et al.*, 2016), an

anaerobic wastewater digester in Oakland (California), and one, assembled by us, from a

PCB-fed culture inoculated with sediments from Liangjiang River, China (Wang and He,

2013). Network analysis revealed that the genome from the Alaskan oil reservoir is most

similar to those from the Albertan oil reservoir B, whereas the genomes from China and

California show high similarity (> 99%) to each other and to *Mesotoga* sp. SC_NapDC

from a northern Albertan oil sands tailings pond (Figure S3c). In fact, these genomes

show the second highest level of pairwise identity among the A lineage genomes (Figure

S3d), suggesting a recent dispersal possibly due to human activities in the last decades.

The A lineage is more isolated from the other lineages (Figure 1 and 2), which

might suggest that this clade evolved in isolation since the formation of oil reservoir

sediments in Alberta 55–120 Ma (Head, 2014; Schaefer, 2005; Selby, 2005)). The high

similarity of the MAGs from the Alaskan oil field to the Albertan genomes and MAGs

from the A and NA lineages (Figure S3) could be due to these oil reservoir sediments

being laid down around the same time (~100 Ma Hu *et al.*, 2016). However, the position

of these MAG in the genome networks could also be explained by these oil reservoirs

being colonized by the same subsurface population, as suggested for *Thermotoga* spp.

(Nesbø *et al.*, 2015). Additional indigenous oil reservoir genomes are needed to resolve

this question.

Distinct metabolism in mesophilic Thermotogae

Phylogeographic analyses provided intriguing evidence that *Mesotoga*'s lower growth

temperature has influenced its distribution patterns. However, the *Mesotoga* genus also

displays additional lineage-specific phenotypes. We therefore examined the newly

available genomes for metabolic insights, which may be linked to *Mesotoga*'s lower

growth temperatures and may influence the role(s) *Mesotoga* play in their environments.

***Mesotoga*-specific genes:** Comparing the *Mesotoga* isolate genomes to other

Thermotogae genomes in IMG (Markowitz *et al.*, 2014) revealed 200 *M. prima* genes

found in all *Mesotoga* genomes (including the more distantly related *M. infera* not

included in the phylogenomic analyses), but in no other Thermotogae genomes. The

majority of these genes were hypothetical proteins (N=119, Supplementary Table S1).

When *Mesotoga*-specific genes with a predicted function were classified according to

Clusters of Orthologous Groups (COG) categories, the largest category was 'Amino Acid

metabolism and transport' with 11 genes, most of which were dipeptidases (COG4690,

N=6). Interestingly, amino acid metabolism genes were up-regulated during growth at

low temperatures in *Mesotoga*'s closest relative *Kosmotoga olearia* (Pollo *et al.*, 2017),

suggesting a role in low temperature adaptations.

Reducing equivalents and thiosulfate reduction: *Mesotoga*'s core metabolism

differs from that of other characterized Thermotogae. While growth of most Thermotogae

is stimulated by adding thiosulfate, sulfur, or other reduced sulfur compounds to the

medium (Ravot *et al.*, 1995; Boileau *et al.*, 2016), reduction of sulfur compounds appears

to be essential for growth of *Mesotoga* in pure culture (Hania *et al.*, 2011; 2013;

Fadhlaoui *et al.*, 2017). The first description of *M. prima* (Nesbø *et al.*, 2012) reported

that growth was only slightly stimulated by the presence of thiosulfate and sulfur.

However, here we observed growth of this isolate only with sulfur or thiosulfate

(Supplementary Table S4 and Table S5), confirming that this is a general trait of

Mesotoga spp. Additionally, while other Thermotogae produce H_2 (and H_2S if grown with

partially reduced sulfur compounds) *Mesotoga* spp. produce large amounts of H_2S and no

or little H_2 (Supplementary Table S5). Fadhlaoui *et al.*, (2017) suggested that *Mesotoga's*

inability to ferment sugars without sulfur compounds is due to the lack of a bifurcating

hydrogenase observed, for instance, in *Thermotoga* spp. However, *K. olearia* also lacks

this enzyme yet produces large amounts of H_2 while fermenting pyruvate by using a Fe-

hydrogenase (Kole_0172) (Pollo *et al.*, 2017).

To reconcile these observations with genomic data, transcriptome analysis was

performed using a culture of *M. prima* grown with 0.5% yeast extract, xylose and

thiosulfate. RNAseq analysis revealed high expression of Theba_0443 (RPKM of 3650;

Supplementary Tables S1 and S6) encoding a Fe-hydrogenase homologous to the one

used by *K. olearia* (Kole_0172). Hydrogenases are indeed essential in Thermotogae for

recycling of ferredoxins (Schut *et al.*, 2013); therefore, finding the same hydrogenase to

be highly expressed in *M. prima* and *K. olearia*, and conserved in all *Mesotoga* genomes

investigated here, suggests that *Mesotoga* possesses a mechanism relying on oxidized

sulfur compounds, efficiently converting all intracellularly produced H_2 to H_2S. Notably,

there was no change in the culture headspace gas H_2:N_2 ratio after incubating *Mesotoga*

spp. in a 1:9 H_2:N_2 atmosphere for > 5 months (Supplementary Table S5), suggesting that

Mesotoga neither produces nor takes up externally supplied H_2.

No homologs of characterized thiosulfate reductases were identified, although the

Mesotoga genomes carry homologs (Theba_0076; Theba_0077 in *M. prima*) of an

archaeal intracellular ferredoxin:NADP oxidoreductase (SudAB; Hagen *et al.*, 2000)

capable of acting as a sulfide dehydrogenase in the presence of elemental sulfur or

polysulfide (Figure 3). Both genes were transcribed at moderate levels in *M. prima* grown

with thiosulfate (RPKM 341 and 243, respectively), while the *K. olearia* homologs

(Kole_1827, Kole_1828) were highly expressed under similar conditions (RPKM > 1000,

(Pollo *et al.*, 2017). SudAB complexes, however, are not known to be involved in

thiosulfate reduction. This is probably due to an unfavorable $E°$= 82 mV for the reaction

when NADH acts as electron donor: $E°`$ [$S_2O_3^{2-}$/ $HS^-$ + $SO_3^{2-}$] = -402 mV and $E°`$

[NAD+/ NADH] = -320 mV. The $E°`$ of [Fd_{Ox} / Fd_{Red}] is similarly high at -390 mV.

Comparable endergonic reactions are catalyzed by the *Salmonella enterica* thiosulfate

reductase Phs by utilizing proton-motive force (Stoffels *et al.*, 2012). However, the

cytoplasmic SudAB complex cannot couple proton-motive force and reduction of an

external electron acceptor. Thus, neither NADH nor Fd_{Red} can function as electron donors

for thiosulfate reduction by *M. prima*. Instead molecular H_2 with $E°`$ [$2H^+ / H_2$] = -410

mV appears to be a thermodynamically preferable electron donor for thiosulfate reduction.

The only hydrogenase present in the *M. prima* genome is the highly expressed FeFe-

hydrogenase (Theba_0443), which usually is involved in Fd-dependent H_2 production

(Vignais and Billoud, 2007). A cluster of five highly transcribed genes (Theba_0461 –

0465, RPKM 1203-3697, Supplementary Tables S1 and S6) encode proteins homologous

to all subunits of the NADP-reducing hydrogenase Hnd of *Desulfovibrio fructosovorans*

(Nouailler *et al.*, 2006) except the catalytic subunit (HndD). These proteins may work

together with Theba_0443 to form a FeFe-hydrogenase complex (Figure 3). We

hypothesize that this complex is involved in intracellular synthesis of molecular hydrogen

for thiosulfate reduction by SudAB coupled to NADH oxidation (Fd_{Red}, Figure 3). The

Hnd genes have homologs in other Thermotogae, however, similar genomic context is

observed only in genomes of other *Mesotoga* and *Kosmotoga* spp. (Supplementary Table

S7).

Mesotoga cells require enzymes that re-oxidize Fd_{red} formed during sugar

oxidation. This might be carried out by either the NADP:ferredoxin oxidoreductase

complex (Mbx; Theba_1796-1808 in *M. prima*. Schut et al., 2013) or the Rnf ion-motive

electron transport complex (Theba_1343-1348; Müller *et al.*, 2008). Conserved motifs

(Mulkidjanian *et al.*, 2008) suggested a Na^+-translocating F-type ATP synthase operating

in *M. prima*. As a consequence, both Mbx and Rnf complexes are predicted to export Na^+

generating sodium- motive force instead of protein-motive force. Genes encoding Mbx

and Rnf show low and moderate expression (RPKM 37-88 and 236-478, respectively)

during growth on thiosulfate, and the expression values suggests that Rnf is the main

complex involved. The model shown in Figure 3 accounts for the observed dependence

of *M. prima* on sulfur or thiosulfate for growth, the lack of H_2 production, and involves

genes already implicated in hydrogen and sulfur metabolism. Importantly, however,

currently there are no known enzymes that couple H_2 oxidation and thiosulfate/sulfur

reduction. It is therefore possible that *M. prima* SudAB uses NADH as the electron donor

and it is much more effective than the hydrogenase which results in almost no H_2 as

growth product (Figure 3 panel C).

Alternatively, thiosulfate reduction coupled to H_2 oxidation (i.e., the postulated

role of SudAB; Figure 3a) may be performed solely by one of the highly-transcribed

hypothetical *Mesotoga* protein with no match to genes in *Kosmotoga* and other

Thermotogae, or in combination with SudAB (Figure 3 panel b). Several candidate genes

listed in Supplementary Table S6 encode proteins with unknown functions. Functional

studies of these genes, as well as the gene products shown in Figure 3, are needed to

assess their role, if any, in thiosulfate reduction. Additional genes that may be involved in

thiosulfate uptake and electron transfer are also discussed in Supplementary Information.

Interestingly, PhosAc3 and *M. infera* were found to reduce only elemental sulfur

(Hania *et al.*, 2015; 2013) whereas the strains isolated by us also reduce thiosulfate.

These differences may reflect selection during isolation; all the isolates obtained in the

current study were from enrichment cultures containing thiosulfate, whereas PhosAc3

and *M. infera* were enriched with sulfur. This suggests that the sulfur-compound-

preference may be a variable and flexible phenotype in *Mesotoga* populations.

Acetate and xylose utilization: Growth on acetate was reported for *Mesotoga*

PhosAc3 (Hania *et al.*, 2015), and we observed weak stimulation of growth of its close

relative *M. prima* on acetate (day 5-10 in Supplementary Figure S4 and Table S4). Nobu

et al. (2015) suggested that Ca. "*Mesotoga acetoxidans*", a MAG closely related to *M.*

infera, oxidizes acetate by using a novel pathway even though the genes comprising the

pathway are conserved in all Thermotogae genomes. Yet, this phenotype is uncommon

among Thermotogae and has been reported only for *Pseudothermotoga lettingae* (Balk,

2002). Instead, many Thermotogae are inhibited by acetate, including one of *Mesotoga*'s

closest relatives, *K. olearia* (Dipippo *et al.*, 2009). Our search for *Mesotoga*-specific

genes that may be responsible for their observed growth on acetate revealed candidate

gene encoding a bacterial homodimeric pyruvate:ferredoxin oxidoreductase (PFOR;

Theba_1954). Its close homologs were found only in *Kosmotoga pacifica* (L'Haridon *et*

al., 2013) and *Mesoaciditoga lauensis* (Reysenbach *et al.*, 2013). Unfortunately, the

description of these two species did not investigate growth on acetate. The *pfor* gene is

distantly related to the archaeal multi-subunit-type used by other Thermotogae (Ragsdale,

2003) and almost all its close homologs fall within the *Clostridia* (Supplementary Figure

S5). Genes having 97-99% identity to *pfor* from *M. infera*, and 83-85% identity to the *M.*

prima homolog, were found in both the metagenome and metatranscriptome published by

Nobu *et al.*; 2015) (locus tag JGI12104J13512_10052834 and

JGI11944J13513_10066464) but were not included in their model. We propose that

PFOR may work with the acetate kinase (Theba_0428 in *M. prima*) and

phosphotransacetylase (Theba_0782 in *M. prima*) found in all Thermotogae to enable

Mesotoga to grow on acetate. At high extracellular acetate concentrations we suggest that

PFOR shifts the balance favoring the production of pyruvate from acetyl-CoA (i.e. serves

as an acetate switch Wolfe, 2005).

M. prima grows optimally on xylose, a sugar fermented by many Thermotogae

(Bhandari and Gupta, 2014). The D-xylose utilization pathway is similar to that observed

in Firmicutes (Gu *et al.*, 2010)(Figure 3). Several possible xylulose kinases genes were

found co-localized with genes encoding xylosidases, sugar transporters and kinases,

suggesting their synergetic activities in xylan hydrolysis, xylose import and utilization.

Mesotoga-specific genes related to O_2 exposure*:* Several *Mesotoga*-specific

genes are predicted to be involved in oxygen-radical defense (Supplementary Table S6).

One of the most highly conserved genes across all the *Mesotoga* genomes (Theba_1553;

average pairwise identity 96.3%) shows similarity to peroxiredoxin and alkyl

hydroperoxide reductase domain-encoding genes. Moreover, a catalase gene

(Theba_0075) is found in all isolate genomes except those from oil reservoir H.

The higher abundance of oxygen-radical defense may be linked to the lower

growth temperatures of *Mesotoga* versus thermophilic Thermotogae. O_2 solubility in

water is greater and free radicals are stabilized at low temperatures, and organisms living

at low temperatures are therefore exposed to higher concentrations of reactive oxygen

species (Piette *et al.*, 2010). It should be noted that the transcriptome of *M. prima* grown

anaerobically revealed that some of the genes possibly involved in O_2 or H_2O_2 defense

(e.g., catalase) were highly expressed (top 5% of expressed genes; Supplementary Table

S1 and S6), suggesting that these genes may have additional or alternative functions

under anaerobic conditions. Further investigation is needed to clarify the transcriptional

responses of these genes and identify the targets of their enzymes.

Concluding remarks.

Our genomic analyses show that the lower growth temperature of *Mesotoga* spp.

compared to the hyperthermophilic *Thermotoga* likely has significantly influenced

Mesotoga phylogeography, with geographic separation having a greater influence than

genetic separation, possibly due to the damaging effects of oxygen exposure during

dispersal. Whether this is a general feature of strictly anaerobic organisms remains to be

resolved. There is also some indication of possible eco-type differentiation among the

Mesotoga lineages, with the US lineage being common in communities degrading

aromatic pollutants (PCB, tolulene) and the A lineage in hydrocarbon-impacted sites.

However, for both of these lineages, inspection of metagenomes revealed they are not

restricted to these environments.

The ecological role of *Mesotoga in situ* may differ from their thermophilic

relatives. For instance, hydrogen-producing *Thermotoga* spp. have been shown to grow in

syntrophy with hydrogenotrophic methanogens (e.g., Johnson *et al.*, 2005) but this is

likely not the case for *Mesotoga* that produce only trace amounts or no detectable H_2.

Supporting this proposal, we were unable to establish co-cultures of *M. prima* and a

hydrogenotrophic methanogen (not shown). Instead Fadhlaoui *et al.* (2017) showed that

Mesotoga spp. prefer to grow in syntrophy with hydrogenotrophic sulfate-reducing

bacteria. This, together with the ability to both produce and consume acetate, suggests

that *Mesotoga* will assume different environmental roles than their thermophilic relatives,

for instance by supporting the growth of sulfate reducers rather than methanogens. An

interesting question is whether they also grow syntrophically with other common

hydrogenotrophic organisms in their niches, such as organohalide-respiring

Dehalococcoides (e.g. Fagervold *et al.*, 2007). Finally, the large amounts of H_2S

produced by *Mesotoga* could have detrimental effects on oil reservoirs, production

facilities and pipelines where *Mesotoga* is commonly found. Monitoring the presence of

Mesotoga spp. in addition to the more commonly targeted sulfate reducers in these

industrial environments (Lee *et al.*, 1995) may be informative and valuable.

Acknowledgements

This work was supported by a Norwegian Research Council award (project no.
180444/V40) to C.L.N. and by a Genome Canada grant (Hydrocarbon Metagenomics
Project) to J.F. The work of IVK was supported by the Federal Agency for Scientific
Organiations of Russia. We thank Dr. Alexander Lebedinsky for constructive criticism
and helpful suggestions.

References

Andreani NA, Hesse E, Vos M. (2017). Prokaryote genome fluidity is dependent on
effective population size. *The ISME Journal* **11**: 1719–1721.

Balk M, Weijma J, Stams AJM. (2002). *Thermotoga lettingae* sp. nov., a novel
thermophilic, methanol-degrading bacterium isolated from a thermophilic anaerobic
reactor. *Int J Syst Evol Micr* **52**: 1361–1368.

Bhandari V, Gupta RS. (2014). The Phylum Thermotogae. In: *The Prokaryotes*. Springer,
Berlin, Heidelberg: Berlin, Heidelberg, pp 989–1015.

Boileau C, Auria R, Davidson S, Casalot L, Christen P, Liebgott P-P, *et al.* (2016).
Hydrogen production by the hyperthermophilic bacterium *Thermotoga maritima* part I:
effects of sulfured nutriments, with thiosulfate as model, on hydrogen production and
growth. *Biotechno Biofuels* **9**: 269.

Chakraborty A, Ellefson E, Li C, Gittins D, Brooks JM, Bernard BB, *et al.* (2018).
Thermophilic endospores associated with migrated thermogenic hydrocarbons in deep
Gulf of Mexico marine sediments. *The ISME Journal* 1–12.

Choudoir MJ, Panke-Buisse K, Andam CP, Buckley DH. (2017). Genome Surfing As
Driver of Microbial Genomic Diversity. *Trends Microbiol* 1–13.

Darling AE, Mau B, Perna NT. (2010). ProgressiveMauve: Multiple Genome Alignment

with Gene Gain, Loss and Rearrangement. *PLoS ONE* **5**: e11147.

Dipippo JL, Nesbø CL, Dahle H, Doolittle WF, Birkland N-K, Noll KM. (2009).
Kosmotoga olearia gen. nov., sp. nov., a thermophilic, anaerobic heterotroph isolated
from an oil production fluid. *Int J Syst Evol Micr* **59**: 2991–3000.

Eckford RE, Fedorak PM. (2002). Planktonic nitrate-reducing bacteria and sulfate-
reducing bacteria in some western Canadian oil field waters. *J Ind Microbiol Biotechnol*
29: 83–92.

Fadhlaoui K, Hania WB, Armougom F, Bartoli M, Fardeau M-L, Erauso G, *et al.* (2017).
Obligate sugar oxidation in *Mesotoga* spp., phylum Thermotogae, in the presence of
either elemental sulfur or hydrogenotrophic sulfate-reducers as electron acceptor. *Environ*
Microbiol **20**: 281–292.

Fagervold SK, May HD, Sowers KR. (2007). Microbial reductive dechlorination of
Aroclor 1260 in Baltimore Harbor sediment microcosms is catalyzed by three phylotypes
within the Phylum Chloroflexi. *Appl Environ Microbiol* **73**: 3009–3018.

Fondi M, Karkman A, Tamminen MV, Bosi E, Virta M, Fani R, *et al.* (2016). 'Every
Gene Is Everywhere but the Environment Selects': Global Geolocalization of Gene
Sharing in Environmental Samples through Network Analysis. *Genome Biol Evol* **8**:
1388–1400.

Fowler SJ, Dong X, Sensen CW, Suflita JM, Gieg LM. (2012). Methanogenic toluene
metabolism: community structure and intermediates. *Environ Microbiol* **14**: 754–764.

Gieg LM, Kolhatkar RV, McInerney MJ, Tanner RS, Harris SH, Sublette KL, *et al.*
(1999). Intrinsic bioremediation of petroleum hydrocarbons in a gas condensate-
contaminated aquifer. *Environ Sci Technol* **33**: 2550–2560.

Goris J, Konstantinidis KT, Klappenbach JA, Coenye T, Vandamme P, Tiedje JM. (2007).
DNA-DNA hybridization values and their relationship to whole-genome sequence
similarities. *Int J Syst Evol Micr* **57**: 81–91.

Gu Y, Ding Y, Ren C, Sun Z, Rodionov DA, Zhang W, *et al.* (2010). Reconstruction of
xylose utilization pathway and regulons in Firmicutes. *BMC Genomics* **11**: 255.

Hagen WR, Silva PJ, Amorim MA, Hagedoorn PL, Wassink H, Haaker H, *et al.* (2000).
Novel structure and redox chemistry of the prosthetic groups of the iron-sulfur
flavoprotein sulfide dehydrogenase from *Pyrococcus furiosus*; evidence for a [2Fe-2S]
cluster with Asp Cys). *JBIC* **5**: 527-534

Hania WB, Fadhlaoui K, Brochier-Armanet C, Persillon C, Postec A, Hamdi M, *et al.*
(2015). Draft genome sequence of *Mesotoga strain* PhosAC3, a mesophilic member of
the bacterial order Thermotogales, isolated from a digestor treating phosphogypsum in
Tunisia. *Stand Genomic Sci* **10**:12.

Hania WB, Ghodbane R, Postec A, Brochier-Armanet C, Hamdi M, Fardeau M-L, *et al.*
(2011). Cultivation of the first mesophilic representative ('mesotoga') within the order
Thermotogales. *Systematic and Applied Microbiology* **34**: 581–585.

Hania WB, Postec A, Aüllo T, Ranchou-Peyruse A, Erauso G, Brochier-Armanet C, *et al.*
(2013). *Mesotoga infera* sp. nov., a mesophilic member of the order Thermotogales,
isolated from an underground gas storage aquifer. *Int J Syst Evol Micr* **63**: 3003–3008.

Head IM, Gray ND, Larter SR. (2014). Life in the slow lane; biogeochemistry of
biodegraded petroleum containing reservoirs and implications for energy recovery and
carbon management. *Front Microbiol* **5**: Article 566.

Holoman TR, Elberson MA, Cutter LA, May HD, Sowers KR. (1998). Characterization
of a defined 2,3,5, 6-tetrachlorobiphenyl-ortho-dechlorinating microbial community by
comparative sequence analysis of genes coding for 16S rRNA. *Appl Environ Microbiol*
64: 3359–3367.

Hu P, Tom L, Singh A, Thomas BC, Baker BJ, Piceno YM, *et al.* (2016). Genome-
resolved metagenomic analysis reveals roles for Candidate Phyla and other microbial
community members in biogeochemical transformations in oil reservoirs. *mBio* **7**:
e01669–15–12.

Hulecki JC, Foght JM, Gray MR, Fedorak PM. (2009). Sulfide persistence in oil field
waters amended with nitrate and acetate. *J Ind Microbiol Biotechnol* **36**: 1499–1511.

Huson DH, Bryant D. (2006). Application of phylogenetic networks in evolutionary
studies. *Mol Biol Evol* **23**: 254–267.

Johnson MR, Conners SB, Montero CI, Chou CJ, Shockley KR, Kelly RM. (2005). The
Thermotoga maritima phenotype is impacted by syntrophic interaction with
Methanococcus jannaschii in hyperthermophilic coculture. *Appl Environ Microbiol* **72**:
811–818.

Jolley KA, Wilson DJ, Kriz P, McVean G, Martin MCJ (2004). The influence of
mutation, recombination, population history, and selection on patterns of genetic diversity
in *Neisseria meningitidis*. *Mol Biol Evol* **22**: 562–569.

Koskella B, Brockhurst MA. (2014). Bacteria-phage coevolution as a driver of ecological
and evolutionary processes in microbial communities. *FEMS Microbiol Rev* **38**: 916–931.

Krzywinski M, Schein J, Birol İ, Connors J, Gascoyne R, Horsman D, *et al.* (2009).
Circos: An information aesthetic for comparative genomics. *Genome Research* **19**: 1639–
1645.

Laing C, Buchanan C, Taboada EN, Zhang Y, Kropinski A, Villegas A, *et al.* (2010).
Pan-genome sequence analysis using Panseq: an online tool for the rapid analysis of core
and accessory genomic regions. *BMC Bioinformatics* **11**: 461.

Lee W, Lewandowski Z, Nielsen PH, Hamilton WA. (1995). Role of sulfate-reducing
bacteria in corrosion of mild steel: A review. *Biofouling* **8**: 165–194.

L'Haridon S, Jiang L, Alain K, Chalopin M, Rouxel O, Beauverger M, *et al.* (2013).
Kosmotoga pacifica sp. nov., a thermophilic chemoorganoheterotrophic bacterium
isolated from an East Pacific hydrothermal sediment. *Extremophiles* **18**: 81–88.

Markowitz VM, Chen I-MA, Palaniappan K, Chu K, Szeto E, Pillay M, *et al.* (2014).
IMG 4 version of the integrated microbial genomes comparative analysis system. *Nucleic*
Acids Res **42**: D560–7.

Martin DP, Lemey P, Lott M, Moulton V, Posada D, Lefeuvre P. (2010). RDP3: a
flexible and fast computer program for analyzing recombination. *Bioinformatics* **26**:
2462–2463.

McInerney JO, McNally A, O'Connell MJ. (2017). Why prokaryotes have pangenomes.
Nat Microbiol **2**: 1–5.

McVean G, Awadalla P, Fearnhead P. (2002). A Coalescent-Based method for detecting
and estimating recombination from gene sequences. *Genetics* **160**: 1231–1241.

Mulkidjanian AY, Galperin MY, Makarova KS, Wolf YI, Koonin EV. (2008).
Evolutionary primacy of sodium bioenergetics. *Biology Direct* **3**: 13.

Müller V, Imkamp F, Biegel E, Schmidt S, Dilling S. (2008). Discovery of a
Ferredoxin:NAD+-Oxidoreductase (Rnf) in *Acetobacterium woodii*. *Ann NY Acad Sci*
1125: 137–146.

Nesbø CL, Bradnan DM, Adebusuyi A, Dlutek M, Petrus AK, Foght J, *et al.* (2012).
Mesotoga prima gen. nov., sp. nov., the first described mesophilic species of the
Thermotogales. *Extremophiles* **16**: 387–393.

Nesbø CL, Dlutek M, Zhaxybayeva O, Doolittle WF. (2006). Evidence for existence of
'mesotogas,' members of the order Thermotogales adapted to low-temperature
environments. *Appl Environ Microbiol* **72**: 5061–5068.

Nesbø CL, Kumaraswamy R, Dlutek M, Doolittle WF, Foght JM. (2010). Searching for
mesophilic Thermotogales bacteria: 'mesotogas' in the wild. *Appl Environ Microbiol* **76**:
4896–4900.

Nesbø CL, S Swithers K, Dahle H, Haverkamp THA, Birkeland N-K, Sokolova T, *et al.*
(2015). Evidence for extensive gene flow and *Thermotoga* subpopulations in subsurface
and marine environments. *The ISME Journal* **9**: 1532–1542.

Nobu MK, Narihiro T, Rinke C, Kamagata Y, Tringe SG, Woyke T, *et al.* (2015).
Microbial dark matter ecogenomics reveals complex synergistic networks in a
methanogenic bioreactor. *The ISME Journal*. **9**: 1710-1722.

Nouailler M, Morelli X, Bornet O, Chetrit B, Dermoun Z, Guerlesquin F. (2006).
Solution structure of HndAc: a thioredoxin-like domain involved in the NADP-reducing
hydrogenase complex. *Protein Sci* **15**: 1369–1378.

Petrus AK, Swithers KS, Ranjit C, Wu S, Brewer HM, Gogarten JP, *et al.* (2012). Genes
for the major structural components of Thermotogales species' togas revealed by
proteomic and evolutionary analyses of OmpA and OmpB homologs. *PLoS ONE* **7**:
e40236.

Piette F, D'Amico S, Struvay C, Mazzucchelli G, Renaut J, Tutino ML, *et al.* (2010).
Proteomics of life at low temperatures: trigger factor is the primary chaperone in the
Antarctic bacterium *Pseudoalteromonas haloplanktis* TAC125. *Mol Microbiol* **76**: 120–
132.

Pollo SMJ, Adebusuyi AA, Straub TJ, Foght JM, Zhaxybayeva O, Nesbø CL. (2017).
Genomic insights into temperature-dependent transcriptional responses of Kosmotoga
olearia, a deep-biosphere bacterium that can grow from 20 to 79 °C. *Extremophiles* **21**:
963–979.

Pollo SMJ, Zhaxybayeva O, Nesbø CL. (2015). Insights into thermoadaptation and the
evolution of mesophily from the bacterial phylum *Thermotogae*. *Can J Microbiol* **61**:
655–670.

Ragsdale SW. (2003). Pyruvate Ferredoxin Oxidoreductase and Its Radical Intermediate
†. *Chem Rev* **103**: 2333–2346.

Ravot G, Ollivier B, Magot M, Patel BKC, Fardeau ML, Garcia J-L. (1995). Thiosulfate
reduction, an important physiological feature shared by members of the order
Thermotogales. *Appl Environ Microbiol* **61**: 2053–2055.

Reysenbach A-L, Liu Y, Lindgren AR, Wagner ID, Sislak CD, Mets A, *et al.* (2013).
Mesoaciditoga lauensis gen. nov., sp. nov., a moderately thermoacidophilic member of
the order Thermotogales from a deep-sea hydrothermal vent. *Int J Syst Evol Micr* **63**:
4724–4729.

Schaefer BF. (2005). GEOCHEMISTRY: When Do Rocks Become Oil? *Science* **308**:
1267–1268.

Schut GJ, Boyd ES, Peters JW, Adams MWW. (2013). The modular respiratory
complexes involved in hydrogen and sulfur metabolism by heterotrophic
hyperthermophilic archaea and their evolutionary implications. *FEMS Microbiol Rev* **37**:
182–203.

Selby D. (2005). Direct radiometric dating of hydrocarbon deposits using Rhenium-
Osmium isotopes. *Science* **308**: 1293–1295.

Stoffels L, Krehenbrink M, Berks BC, Unden G. (2012). Thiosulfate reduction in
Salmonella enterica is driven by the proton motive force. *J Bact* **194**: 475–485.

Tan B, Jane Fowler S, Laban NA, Dong X, Sensen CW, Foght J, *et al.* (2015).
Comparative analysis of metagenomes from three methanogenic hydrocarbon-degrading
enrichment cultures with 41 environmental samples. *The ISME Journal* **9**: 2028–2045.

Vignais PM, Billoud B. (2007). Occurrence, classification, and biological function of
hydrogenases: an overview. *Chem Rev* **107**: 4206–4272.

Voordouw G, Grigoryan AA, Lambo A, Lin S, Park HS, Jack TR, *et al.* (2009). Sulfide
remediation by pulsed injection of Nitrate into a low temperature Canadian heavy oil
reservoir. *Environ Sci Technol* **43**: 9512–9518.

Wachter J, Hill S. (2016). Positive selection pressure drives variation on the surface-
exposed variable proteins of the pathogenic *Neisseria*. *PLoS ONE* **11**: e0161348.

Wang S, He J. (2013). Phylogenetically distinct bacteria involve extensive dechlorination
of Aroclor 1260 in sediment-free cultures. *PLoS ONE* **8**: e59178.

Wilhelms A, Larter SR, Head I, Farrimond P, di-Primio R, Zwach C. (2001).
Biodegradation of oil in uplifted basins prevented by deep-burial sterilization. *Nature*
411: 1034–1037.

Wolfe AJ. (2005). The acetate switch. *Microbiology and Molecular Biology Reviews* **69**:
12–50.

Zhaxybayeva O, Swithers KS, Foght J, Green AG, Bruce D, Detter C, *et al.* (2012).
Genome Sequence of the mesophilic thermotogales bacterium *Mesotoga prima* MesG1.
Ag. 4.2 reveals the largest thermotogales genome to date. *Genome Biol Evol* **4**: 700–708.

Figure legends

Figure 1. Phylogenetic relationships among *Mesotoga* genomes based on (a)

16SrRNA genes, (b) core SNPs and (c) presence/absence of shared 500-bp genomic

fragments. The 16S rRNA maximum likelihood phylogeny was estimated using RAxML

in Geneious v 10. For networks shown in (b) and (c), data were obtained using PanSeq

(Laing *et al.*, 2010). Core SNPs in (b) were required to be present in 14 of 18 genomes

(including SAGs), and genomic fragments were considered shared if they were at least

70% identical. The network in (c) was constructed using only genomes from isolates;

shared fragments were required to be present in all 9 genomes and be at least 70%

identical in sequence. Networks were calculated in SplitsTree using NeighborNet

algorithm (Huson and Bryant, 2006) from uncorrected distances. The three lineages are

indicated by W, US and A.

Figure 2. Visualization of recombination events detected among *Mesotoga* genomes

from different lineages. The genomes are color-coded according to lineage and arranged

on the circumference of the circle; W lineage, blue; US lineage, orange; A lineage, green.

The recombination events with predicted donor and recipient are shown as lines

connecting the two genomes and the locations of recombined regions, where line color

reflects the donor lineage and the width of the line is proportional to the length of the

recombinant region. Predicted events were required to be significant in at least 3 of 4

algorithms in RDP (Martin *et al.*, 2010). The diagram was generated using Circos

Version circos-0.69 (Krzywinski *et al.*, 2009).

Figure 3. Model of energy generation pathway in *Mesotoga prima* during growth on

xylose and thiosulfate. Glucose and xylose poly- and oligosaccharides are hydrolyzed by

various intracellular and interstitial glycosidases (GHs). Glucose oxidation occurs via the

glycolytic Embden-Meyerhof-Parnas pathway, whereas xylose is utilized via xylose

isomerase (*XylA*, Theba_1394), xylulose kinase (*XylB*, Theba_1395, Theba_2230, Theba

2429, Theba 2518, Theba 2544, Theba 2588), ribulose phosphate 3-epimerase

(Theba_0639) and enzymes of the pentose-phosphate pathway. Specifically, xylose

isomerase converts D-xylose to D-xylulose, which is phosphorylated by the set of

xylulose kinases to D-xylulose 5-phosphate, and further to ribulose 5-phosphate by the

ribulose-phosphate 3-epimerase. Both xylulose 5-phosphate and ribulose 5-phosphate

produced by this pathway are common metabolic intermediate in the pentose-phosphate

pathway. The xylose isomerase Theba_1394 was among the most highly transcribed

genes during cultivation of *M. prima* on xylose and thiosulfate (Supplementary Table S5).

Acetyl-CoA formation occurs by means of pyruvate-ferredoxin oxidoreductase (PFOR,

Theba_1954). In the possible case of growth on acetate, its activation occurs by means of

acetate kinase (ACKA, Theba_0428) and phosphotransacetylase (PTA, Theba_0782),

acting in reverse. The model includes gene products hypothesized to be involved in

thiosulfate reduction. $\underline{Na^+}$ refers to Na+ ions involved in generating the generating the

sodium motive force. A: The FeFe hydrogenase (Theba_0443 and Theba_0461 – 0465)

reduces NADH to form H_2, which is used as an electron donor for thiosulfate reduction

catalyzed by SudAB (Theba_0076, Theba_0077). Mbx (Theba_1796-1808) and/or Rnf

(Theba_1343-1348) complexes provide additional NADH along with the oxidation of

excessive reducing equivalents (Fd_{red}) and generation of a sodium-motive force. B and C:

other possible scenarios of H_2 oxidation and thiosulfate reduction.

Table 1. List of genomes analyzed. All genomes, except those of *Mesotoga prima, M. prima* PhosAc3 and *Mesotoga infera,* were sequenced as part of the current study.

Name and Source	Short Name	Genome Size	% GC	Ref. for description of sample site / accession no. in Genbank	% completeness of SAG[a]
Isolates					
Produced water from oil field B near Brooks, Alberta, Canada[b]				(Hulecki *et al.*, 2009)	
Mesotoga sp. Brooks.08.YT.4.2.5.1[c]	BR5.1	2,957,195	45.9	AYTX01000000	
Mesotoga sp. Brooks.08.YT.4.2.5.2	BR5.2	2,953,308	45.9	JPGZ00000000	
Mesotoga sp. Brooks.08.YT.4.2.5.4[c]	BR5.4	3,002,147	45.9	ATCT01000000	
Mesotoga sp. Brooks.08.YT.105.5.1	BR105.1	2,992,699	45.9	AYTW01000000	
Mesotoga sp. Brooks.08.YT.105.6.4	BR105.4	3,205,299	45.9	JWIM00000000	
Free water knockout fluids from oil field H near Stettler, Alberta[d]				(Eckford and Fedorak, 2002)	
Mesotoga sp. HF07.pep.5.2[c]	HF5.2	2,838,813	45.3	JFHJ01000000	
Mesotoga sp. HF07.pep.5.3	HF5.3	2,934,282	45.3	AYTV01000000	
Mesotoga sp. HF07.pep.5.4	HF5.4	2,968,642	45.3	JFHM01000000	
Sediments from Baltimore Harbour, Maryland, USA				(Holoman *et al.*, 1998)	
Mesotoga prima MesG1.Ag.4.2[e]	*M.prima*	2,974,229	45.5	NC_017934	
Mesotoga sp. BH458.6.3.2.1[f]	BH458	3,234,409	45.7	JFHL01000000	
Wastewater treatment plant, Tunisia				(Hania *et al.*, 2011)	
Mesotoga prima PhosAc3	PhosAc3	3,108,267	45.2	NZ_CARH01000000	

(continued)

Genomes assembled from single cell amplified genomes (SAGs)

Produced water from oil field E near Medicine Hat, Alberta[g]				(Voordouw *et al.*, 2009)	
Mesotoga sp. 3PWK154PWL11	SC_PW.1	876,625	46.8	JMRN01000000	21%
Mesotoga sp. 3PWM13N19	SC_PW.2	1,886,634	45.8	JMRM01000000	78%
Mesotoga sp. 4PWA21	SC_PW.3	1,541,163	45.9	JMQL01000000	34%
Oil sands tailings pond sediments near Fort McMurray, Alberta				(Tan *et al.*, 2015)	
Mesotoga sp. NapDC	SC_NapDC	1,885,291	45.8	JNFM01000000	88%
Mesotoga sp. NapDC2	SC_NapDC2	1,337,305	45.6	JQSC01000000	53%
Mesotoga sp. NapDC3	SC_NapDC3	1,828,922	45.6	JWIP00000000	66%
Contaminated aquifer fluids from Colorado, USA				(Fowler *et al.*, 2012; Gieg *et al.*, 1999)	
Mesotoga sp. TolDC[h]	SC_TOLDC	2,257,992	46.1	AYSI01000000	74%

[a] Completeness of single cell genomes was calculated based on HMM hits to 119 single copy marker genes (see Supplementary Material).

[b] Belongs geologically to the Glauconitic formation.

[c] These genomes were sequenced using IonTorrent PGM; all other genomes and single cells were sequenced using Illumina MiSeq

[d] Belongs geologically to the Upper Mannville Group – Cretaceous age.

[e] The *M. prima* genome was sequenced by Zhaxybayeva *et al.* (2012).

[f] This is a sister-culture of the enrichment culture that yielded *M. prima* MesG1.Ag.4.2 (Nesbø *et al.*, 2006; 2012).

[g] Belongs geologically to the Western Canadian Sedimentary Basin.

[h] In addition to the SAG sequences, 39 *Mesotoga* fosmid clones prepared from the same culture were included in the assembly (http://hmp.ucalgary.ca/HMP/metagenomes/isolates.html).

Table 2. *Mesotoga* sequences recovered from publicly available metagenomes. For the sequences obtained from the IMG (JGI) database, only sequences classified as Thermotogae were downloaded. The predominant *Mesotoga* lineage in each metagenome is shown in boldface. Metagenomes dominated by sequences similar to *Mesotoga infera* were not included.

Metagenome origin[a]	IMG, Genbank or SRA accession nos.	*Mesotoga* sequences (no. contigs)	*Mesotoga* sequences with best match to lineage A, W or US (no. contigs)	% average pairwise identity (range)
Alberta				
MLSB tailings pond	IMG: 26785	491,657 bp (1,707)	**A: 485,638 bp (1,694)**	**99.8% (98.5-100%)**
			W: 2,455 bp (8)	93.9% (88.7-99.6)
			US: 634 bp (2)	86.4% (89.7-95%)
Oil reservoir E	IMG: 15764	5,190,293 bp (4,833)	**A: 3,137,228 bp (3,259)**	**98.6% (71.3-100%)**
			W: 195,159 bp (317)	94.5% (79.7-100%)
			US: 1,693,057 bp (1,308)	**95.3% (82.2-100%)**
USA				
California				
Alameda Naval Air Station: Soil contaminated with Chloroethene	IMG: 5776	407,588 bp (453)	A: 17,955 bp (23)	92% (85.3-99.5%)
			W: 47,017 bp (61)	93.4% (83-99.1%)
			US: 297,011 bp (360)	**94.2%(83.6-100%)**
Blank Spring hotspring sediment	IMG: 94476	4,060,664 bp (2,037)	**A: 920,911 bp (924)**	**91.6% (64.3-100%)**
			W: 226,161 bp (176)	85.4% (64.3-99.1%)
			US: 2,025,524 bp (825)	**89.3% (66.1-99.6 %)**
Long Beach Municipal wastewater AD_UKC097	IMG: 89744	2,339,863 bp (1,810)	**A: 390,020 bp (588)**	**96.1% (65-100%)**
			W: 710,377 bp (914)	**94.1% (64.3-100%)**
			US: 202,508 bp (207)	88.4% (63.6-100%)

(continued)

Illinois				
Decatur municipal wastewater AD_UKC034[b]	IMG: 89745	4,031,397 bp (2,218)	**A: 2,085,927 bp (1,162)** **W: 820,163 bp (588)** US: 271,756 bp (260)	**96.5% (65.1-100%)** **92.7% (64.6-100%)** 89.0% (65.0-100%)
New York Sulfidogenic MTBE-NYH[c]	IMG: 62988	5,195,092 bp[i] (1,727)	**A: 421,552 bp (261)** W: 209,518 bp (218) **US:1,825,787 bp (450)**	**85.8% (63.8-99.2%)** 84.6% (64.5-99.8%) **89.2% (65.9-100%)**
New Jersey Methanogenic MTBE-AKM[d]	IMG: 62224	2,464,953 bp (4,206)	A: 31,663 bp (104) W: 122,392 bp (320) **US: 1,154,398 bp (2,706)**	87.2% (66.8-100%) 93.8% (65.0-100%) **98.0% (66.5-100%)**
Sulfidogenic MTBE-AKS2[e]	IMG: 62223	4,971,880 (3,215)	A: 152,359 bp (276) W: 240,489 bp (307) **US: 2,665,479 bp (1,575)**	86.3% (65.9-99.3%) 86.6% (65.6-100%) **94.3% (65.8-100%)**
Boston Wastewater AD_UKC077[f]	IMG: 89805	5,651,655[i] (1,386)	A: 149,486 bp (212) **W: 1,379,334 bp (326)** US: 141,168 bp (150)	80.2% (63.4-100%) **88.2% (63.7-100%)** 77.6% (64.4-99.7%)
Hong Kong Wastewater AD_UKC109[f] 2015-03-06	IMG: 89888	4,071,190 bp (4,096)	A: 415,727 bp (615) **W: 1,189,447 bp (2,476)** US: 632,042 bp(902)	92.8% (64.7-100%) **93.6% (64.3-100%)** 91.9% (63-100%)
Wastewater AD_UKC119[f] 2015-01-26	IMG: 89894	5,462,335 bp (2,499)	**A: 2,347,962 bp (853)** W: 586,387 bp (497) **US: 1,521,681(1,071)**	**95.7% (65.4-100%)** 89.4% (64.3-100%) **93.0% (63.9-100%)**

(continued)

Metagenome-assembled genomes (MAGs)

Alaska Oil reservoir LGGP01[g]	Genbank: GCA_001508515	1,712,609 bp (440)	**A: 1,470,927 bp (384)** W: 80,31 bp (30) US: 47,775 bp (21)	**98.5% (71.3-100%)** 92.3% (70.0-99.6%) 88.9% (71.9-100%)
LGGH01[g]	GCA_001508435	1,225,111 bp (267)	A: 48,008 bp (13) W: 63,609 bp (21) **US: 1,009,462 bp (233)**	89.9% (78.8-99.6%) 92.5% (80.4-100%) **94.5% (80.9-99.6)**
LGGW01[g]	GCA_001509115	1,622,264 bp (264)	A: 85,486 bp (20) W: 104,756 bp (25) **US: 1,139,439 (211)**	87.3% (67.8-99.6%) 92.3% (82.6-99.9%) **94.5% (83.6-99.8%)**
California Anaerobic digester in Oakland[h]	IMG: 81407 (Unclassified Thermotogales bacterium Bin 13)	3,480,910 bp (395)	**A: 2,287,852 bp (247)** W: 109,011 bp (48) US: 118,885 bp (33)	**93.6% (64.5-100%)** 83.1% (64.4-100%) 79.1% (65.8-96.5)
China PCB-fed mixed *Dehalococcoides* culture CG1 from sand and silt near Liangjiang River[i]	SRA: SRX392467	2,727,841bp (379)	**A: 2,226,7034 bp (345)** W: 22,520 bp (14) US: 40,010 bp (20	**98.0% (84.6-100%)** 90.2% (68.2-97.4%) 90.7% (78.2-99.8%)

---

a) *Mesotoga* sequences were identified by performing blastn searches using a database containing all the *Mesotoga* spp. genomes listed in Table 1. We used word size =11 and expected = e^{-10}. Sequences with matches were then sorted according to the *Mesotoga* lineage with the best match (A, W or US; see main text).

b) Nine metagenomes and three transcriptomes are available from the same system; only one was selected as representative.

c) Sulfidogenic MTBE-degrading enrichment culture microbial communities from New York harbour sediments

d) Methanogenic MTBE-degrading enrichment culture microbial communities from Arthur Kill sediments

e) Sulfidogenic MTBE-degrading enrichment culture microbial communities from Arthur Kill sediments. Two very similar metagenomes are available (Figure S3b); only one is included here.
f) Wastewater treatment anaerobic digesters. Additional metagenomes with similar *Mesotoga* lineage composition are available from this site and the ones included here were chosen as representatives.
g) Metagenome assembled genomes from (Hu *et al.*, 2016). LGGP01 and LGGH01 were from oil reservoir sample SB1 and LGGW01was from oil reservoir sample SB2.
h) Sludge microbial communities from wastewater, phosphite and CO_2-enriched.
i) Genome extracted and assembled by us

a 16S rRNA

Mprima 16S RNA2 ●
TOLDC □
BH458 ●

US

HF52 ■
HF54 ■
PhosAc3 ●
HF53 ■
Mprima 16S RNA1 ●

W

NapDC3 ☆
NapDC2 ☆
BR51 ★
BR52 ★
BR54 ★
BR10551 ★
PW1 ✦
PW3 ✦
NapDC ☆
B10564 ★
PW2 ✦

A

0.02

b Core SNPs

W
Mprima ●
HF53 ■
HF52 ■
HF54 ■
PhosAc3 ●

US
BH458 ●
TOLDC □

0.1

Baltimore Harbor, Maryland ●
Contaminated Aquifer, Colorado □
Wastewater treatment, Tunisia ●
FWKO tank Oil reservoir H, Alberta ■
Produced water Oil reservoir B, Alberta ★
Produced water Oil reservoir E, Alberta ✦
Tailing pond storing waste from oil sands, Alberta ☆

PW3 ✦
B10564 ★
BR10551 ★
BR52 PW1 ★
PW2 ✦
NapDC BR51 ☆★
NapDC2 BR54 ☆★
NapDC3 ☆

A

c Pangenome

PhosAc3 ●
HF54 ■
BH458 ●

US

W
HF52 ■
0.01
HF53 ■
Mprima ●

B10564 ★
BR10551 ★
BR52 ★

A