The iron-dependent repressor YtgR regulates the tryptophan

salvage pathway through a bipartite mechanism of transcriptional

control in *Chlamydia trachomatis*

Nick D. Pokorzynski[1], Amanda J. Brinkworth[1] and Rey A. Carabeo[1]

[1]Center for Reproductive Biology, School of Molecular Biosciences, College of

Veterinary Medicine, Washington State University, Pullman, WA, USA, 99164.

Corresponding Author:

Rey A. Carabeo

1770 NE Stadium Way, Biotechnology & Life Sciences Bldg., Washington State

University

Pullman, WA, 99164

(509) 335-7788

rey.carabeo@wsu.edu

Keywords: stress response, transcription, tryptophan biosynthesis, iron-dependent

regulation, repression

Abstract (Word Limit:150, Word Count:150)

During infection, pathogens are starved of essential nutrients such as iron and

tryptophan by host immune effectors. Without conserved global stress response

regulators, how the obligate intracellular bacterium *Chlamydia trachomatis* arrives at a

physiologically similar "persistent" state in response to starvation of either nutrient

remains unclear. Here, we report on the iron-dependent regulation of the *trpRBA*

tryptophan salvage pathway in *C. trachomatis.* Iron starvation specifically induces *trpBA*

expression from a novel promoter element within an intergenic region flanked by *trpR*

and *trpB*. YtgR, the only known iron-dependent regulator in *Chlamydia*, can bind to the

trpRBA intergenic region upstream of the alternative *trpBA* promoter to repress

transcription. Simultaneously, YtgR binding promotes the termination of transcripts from

the primary promoter upstream of *trpR*. This is the first description of an iron-dependent

mechanism regulating prokaryotic tryptophan biosynthesis that may indicate the

existence of novel approaches to gene regulation and stress response in *Chlamydia.*

Introduction

Nutrient acquisition is critical for the success of pathogenic bacteria. Many pathogenic bacteria must siphon nutrients from their hosts, such as nucleotides, amino acids and biometals (Brown, Palmer, & Whiteley, 2008; Eisenreich, Dandekar, Heesemann, & Goebel, 2010; Ray, Marteyn, Sansonetti, & Tang, 2009; Skaar, 2010). This common feature among pathogens renders them susceptible to nutrient limitation strategies associated with the host immune response (Hood & Skaar, 2012). Counteractively, bacterial pathogens have evolved sophisticated molecular mechanisms to respond to nutrient deprivation, involving increasingly complex and sophisticated nutrient-sensing regulatory networks. These stress response mechanisms are essential for pathogens to avoid clearance by the immune system. By delineating their function at the molecular level, we can better target aspects of the host-pathogen interface suitable for therapeutic manipulation. However, stress responses in the obligate intracellular bacterium *Chlamydia trachomatis* are relatively poorly characterized, leaving unanswered many fundamental questions about the biology of this pathogen.

Chlamydia trachomatis is the leading cause of bacterial sexually transmitted infections (STIs) and infection-derived preventable blindness worldwide (CDC, 2017; Newman et al., 2015; H. R. Taylor, Burton, Haddad, West, & Wright, 2014). Genital infections of chlamydia disproportionately affect women and are associated with serious sequelae in the female reproductive tract such as tubal factor infertility (Hafner, 2015). Chlamydiae are Gram-negative bacterial parasites that develop within a pathogen-specified membrane-bound organelle termed the inclusion (Moore & Ouellette, 2014). Chlamydial development is uniquely characterized by a biphasic interconversion of an

infectious elementary body (EB) with a non-infectious, but replicative reticulate body

(RB) (AbdelRahman & Belland, 2005). An obligate intracellular lifestyle has led to

reductive genome evolution across chlamydial species; Chlamydiae have retained

genes uniquely required for their survival, but have become nutritionally dependent on

their hosts by discarding many metabolism-related genes (Clarke, 2011). Of note, *C.*

trachomatis does not possess genes necessary for eliciting a stringent response to

nutrient starvation (*e.g. relA, spoT*), suggesting that this pathogen may utilize novel

mechanisms to respond to nutrient stress (Stephens et al., 1998).

It is well established that in response to various stressors, Chlamydiae deviate

from their normal developmental program to initiate an aberrant developmental state,

termed "persistence" (Wyrick, 2010). This persistent state is distinguished by the

presence of viable, but non-cultivable, abnormally enlarged chlamydial organisms that

display dysregulated gene expression. Importantly, *Chlamydia* can be reactivated from

persistence by abatement of the stress condition. As such, chlamydial persistence at

least superficially resembles a global stress response mechanism. Yet the molecular

underpinnings of this phenotype are poorly understood, with most published studies

focusing on the molecular and metabolic character of the aberrant, persistent form. It is

therefore unclear to what extent primary stress responses contribute to the global

persistent phenotype in *Chlamydia*.

The best described inducer of persistence is the pro-inflammatory cytokine

interferon-gamma (IFN-γ). The bacteriostatic effect of IFN-γ has been primarily

attributed to host cell tryptophan (Trp) catabolism, an amino acid for which *C.*

trachomatis is auxotrophic (Byrne, Lehmann, & Landry, 1986; Fehlner-Gardiner et al.,

2002; M. W. Taylor & Feng, 1991). Following IFN-γ stimulation, infected host cells up-

regulate expression of indoleamine-2,3-dioxygenase (IDO1), which catabolizes Trp to

N-formylkynurenine via cleavage of the indole ring (Macchiarulo, Camaioni, Nuti, &

Pellicciari, 2009). *C. trachomatis* cannot recycle kynurenines, unlike some other

chlamydial species (Wood, Roshick, & McClarty, 2004)*,* and thus IFN-γ stimulation

effectively results in Trp starvation to *C. trachomatis*. The primary regulatory response

to Trp starvation in *C. trachomatis* is mediated by a TrpR ortholog, whose Trp-

dependent binding to cognate promoter elements represses transcription (Akers & Tan,

2006; Carlson, Wood, Roshick, Caldwell, & McClarty, 2006). This mechanism of

regulatory control is presumably limited in *C. trachomatis*, as homologs of genes

regulated by TrpR in other bacteria (*e.g. trpF, aroH, aroL*) have not been shown to

respond to Trp limitation (Wood et al., 2003).

In many Gram-negative bacteria, such as *Escherichia coli*, *trpR* is monocistronic

and distal to the Trp biosynthetic operon. In *C. trachomatis,* TrpR is encoded in an

operon, *trpRBA*, which also contains the Trp synthase α- and β- subunits (TrpA and

TrpB, respectively), and possesses a 351 base-pair (bp) intergenic region (IGR) that

separates *trpR* from *trpBA*. The functional significance of the *trpRBA* IGR is poorly

characterized. While a putative TrpR operator sequence was identified in the IGR

overlapping an alternative transcriptional origin for *trpBA* (Carlson et al., 2006), TrpR

binding was not shown (Akers & Tan, 2006). Based on *in silico* predictions, an

attenuator sequence has been annotated within the *trpRBA* IGR (Merino & Yanofsky,

2005), but this has not been thoroughly validated experimentally. Regardless, the IGR is

>99% conserved at the nucleotide sequence level across ocular, genital and

lymphogranuloma venereum (LGV) serovars of *C. trachomatis*, indicating functional

importance (Carlson, Porcella, Mcclarty, & Caldwell, 2005; Seth-Smith et al., 2009;

Stephens et al., 1998; Thomson et al., 2008). Therefore, outside of TrpR-mediated

repression, the complete detail of *trpRBA* regulation remains poorly elucidated and

previous reports have indicated the possibility of more complex mechanisms of

regulation (Brinkworth, Wildung, & Carabeo, 2018).

In evaluating alternative regulatory modes of the *trpRBA* operon, an interesting

consideration is the pleiotropic effects induced by IFN-γ stimulation of infected cells.

IFN-γ is involved in many processes that limit iron and other essential biometals to

intracellular pathogens as a component of host nutritional immunity (Cassat & Skaar,

2013; Hood & Skaar, 2012). *Chlamydia* have a strict iron dependence for normal

development, evidenced by the onset of persistence following prolonged iron limitation

(Raulston, 1997). Importantly, *Chlamydia* presumably acquire iron via vesicular

interactions between the chlamydial inclusion and slow-recycling transferrin (Tf)-

containing endosomes (Ouellette & Carabeo, 2010). IFN-γ is known to down-regulate

transferrin receptor (TfR) expression in both monocytes and epithelial cells with

replicative consequences for resident intracellular bacteria (T. F. Byrd & Horwitz, 1993;

T. Byrd & Horwitz, 1989; Igietseme, Ananaba, Candal, Lyn, & Black, 1998; Nairz et al.,

2008). However, iron homeostasis in *Chlamydia* is poorly understood due to the lack of

functionally characterized homologs to iron acquisition machinery that are highly

conserved in other bacteria (Pokorzynski, Thompson, & Carabeo, 2017). Only the

ytgABCD operon, encoding a metal permease, has been clearly linked to iron

acquisition (J. D. Miller, Sal, Schell, Whittimore, & Raulston, 2009). Intriguingly, the

YtgC (CTL0325) open reading frame (ORF) encodes a N-terminal permease domain

fused to a C-terminal DtxR-like repressor domain, annotated YtgR (Akers, HoDac,

Lathrop, & Tan, 2011; Thompson, Nicod, Malcolm, Grieshaber, & Carabeo, 2012). YtgR

is cleaved from the permease domain during infection and functions as an iron-

dependent transcriptional repressor to autoregulate the expression of its own operon

(Thompson et al., 2012). YtgR represents the only identified iron-dependent

transcriptional regulator in *Chlamydia.* Whether YtgR maintains a more diverse

transcriptional regulon beyond the *ytgABCD* operon has not yet been addressed and

remains an intriguing question in the context of immune-mediated iron limitation to *C.*

trachomatis.

Consistent with the highly reduced capacity of the chlamydial genome, it is likely

that *C. trachomatis* has a limited ability to tailor a specific response to each individual

stress. In the absence of identifiable homologs for most global stress response

regulators in *C. trachomatis*, we hypothesized that primary stress responses to

pleiotropic insults may involve mechanisms of regulatory integration, whereby important

molecular pathways are co-regulated by stress-responsive transcription factors such

that they can be utilized across multiple host-mediated stresses. Here, we report on the

unique iron-dependent regulation of the *trpRBA* operon in *Chlamydia trachomatis.* We

propose a model of iron-dependent transcriptional regulation of *trpRBA* mediated by the

repressor YtgR binding specifically to the IGR, which may have implications for how *C.*

trachomatis responds to immunological and environmental insults. Such a mechanism

of iron-dependent regulation of Trp biosynthesis has not been previously described in

any other prokaryote and adds to the catalog of regulatory models for Trp biosynthetic

operons in bacteria. Further, it reveals a highly dynamic mode of regulatory integration

within the *trpRBA* operon, employing bipartite control at the transcription initiation and

termination steps.

Results

Brief iron limitation via 2,2-bipyridyl treatment yields iron-starved, but non-

persistent *Chlamydia trachomatis.* To identify possible instances of regulatory

integration between iron and Trp starvation in *C. trachomatis*, we optimized a stress

response condition that preceded the development of a characteristically persistent

phenotype. We reasoned that in order to effectively identify regulatory integration, we

would need to investigate the bacterium under stressed, but not aberrant, growth

conditions such that we could distinguish primary stress responses from abnormal

growth. To specifically investigate the possible contribution of iron limitation to a broader

immunological (*e.g.* IFN-γ-mediated) stress, we utilized the membrane-permeable iron

chelator 2,2-bipyridyl (Bpdl), which has the advantage of rapidly and homogeneously

starving *C. trachomatis* of iron (Thompson & Carabeo, 2011). We chose to starve *C.*

trachomatis serovar L2 of iron starting at 12 hrs post-infection (hpi), or roughly at the

beginning of mid-cycle growth. At this point the chlamydial organisms represent a

uniform population of replicative RBs that are fully competent, both transcriptionally and

translationally, to respond to stress. We treated infected HeLa cell cultures with 100 μM

Bpdl or mock for either 6 or 12 hours (hrs) to determine a condition sufficient to limit iron

to *C. trachomatis* without inducing hallmark persistent phenotypes. We stained infected

cells seeded on glass coverslips with convalescent human sera and analyzed

chlamydial inclusion morphology under both Bpdl- and mock-treated conditions by laser

point-scanning confocal microscopy (Figure 1A). Following 6 hrs of BpdI treatment,

chlamydial inclusions were largely indistinguishable from mock-treated inclusions,

containing a homogeneous population of larger organisms, consistent with RBs in mid-

cycle growth. However, by 12 hrs of BpdI treatment, the inclusions began to display

signs of aberrant growth: they were perceptibly smaller, more comparable in size to 18

hpi, and contained noticeably fewer organisms, perhaps indicating a delay in RB-to-EB

differentiation. These observations were consistent with our subsequent analysis of

genome replication by quantitative PCR (qPCR; Figure 1B.) At 6 hrs of BpdI treatment,

there was no statistically distinguishable difference in genome copy number when

compared to the equivalent mock-treated time-point. However, by 12 hrs of treatment,

genome copy number was significantly reduced 4.7-fold in the BpdI-treated group

relative to mock-treatment (p = 0.0033). We then assayed the transcript expression of

two markers for persistence by reverse transcription quantitative PCR (RT-qPCR): the

early gene *euo*, encoding a transcriptional repressor of late-cycle genes (Figure 1C),

and the adhesin *omcB,* which is expressed late in the developmental cycle (Figure 1D).

Characteristic persistence would display elevated *euo* expression late into infection, and

suppressed *omcB* expression throughout development. We observed that at 6 hrs of

BpdI treatment, there was no statistically distinguishable difference in either *euo* or

omcB expression when compared to the mock-treatment. Still at 12 hrs of BpdI

treatment, *euo* expression was unchanged. However, *omcB* expression was

significantly induced following 12 hrs of BpdI-treatment (p = 0.00015). This was

unexpected, but we note that *omcB* expression has been shown to vary between

chlamydial serovars and species when starved for iron (Pokorzynski et al., 2017).

Collectively, these data indicated that 6 hrs of BpdI treatment was a more suitable time-

point at which to monitor iron-limited stress responses.

We additionally assayed these same metrics following 6 or 12 hrs of Trp

starvation by culturing cells in either Trp-replete or Trp-deplete DMEM-F12 media

supplemented with fetal bovine serum (FBS) pre-dialyzed to remove amino acids. We

observed no discernable change in inclusion morphology out to 12 hrs of Trp starvation

(Figure 1 – Figure Supplement 1A), but genome copy numbers were significantly

reduced 2.7-fold at this time-point (p = 0.00612; Figure 1 – Figure Supplement 1B). The

transcript expression of *euo* (Figure 1 – Figure Supplement 1C) and *omcB* (Figure 1 –

Figure Supplement 1D) did not significantly change at either treatment duration, but Trp-

depletion did result in a 2.0-fold reduction in *omcB* expression, consistent with a more

characteristic persistent phenotype. These data therefore also indicated that 6 hrs of

treatment would be ideal to monitor non-persistent responses to Trp limitation.

Figure 1. Brief iron limitation via 2,2-bipyridyl treatment precedes the onset of characteristic chlamydial persistence. (*A*) *C. trachomatis* L2-infected HeLa cells were fixed and stained with convalescent human sera to image inclusion morphology by confocal microscopy following BpdI treatment at the indicated times post-infection. Arrowheads indicate inclusions with visibly fewer organisms in the 12-hour BpdI-treated condition. Figure shows representative experiment of three biological replicates. Scale bar = 25 μm, Inset scale bar = 5 μm. (*B*) Genomic DNA (gDNA) was harvested from infected HeLa cells at the indicated times post-infection under iron-replete (blue) and -deplete (red) conditions. Chlamydial genome copy number was quantified by qPCR. Chlamydial genome replication is stalled following 12 hours of BpdI treatment, but not 6. N=2. (*C*) Total RNA was harvested from infected HeLa cells at the indicated times post-infection under iron-replete (teal) and -deplete (orange) conditions. The transcript abundance of hallmark persistence genes *euo* and (*D*) *omcB* were quantified by RT-qPCR and normalized against genome copy number. Only at 12 hours of BpdI treatment is *omcB* expression significantly affected. N=3 for 12+6, N=2 for 12+12. Statistical significance was determined by One-Way ANOVA followed by post-hoc pairwise *t*-tests with Bonferroni's correction for multiple comparisons. * = $p < 0.05$, ** = $p < 0.01$, *** = $p < 0.005$.

We next sought to determine whether our brief 6-hr BpdI treatment was sufficient

to elicit a transcriptional iron starvation phenotype. We chose to analyze the expression

of three previously identified iron-regulated transcripts, *ytgA* (Figure 2A), *ahpC* (Figure

2B) and *devB* (Figure 2C), by RT-qPCR under BpdI- and mock-treated conditions (Dill,

Dessus-Babus, & Raulston, 2009; Thompson & Carabeo, 2011). In addition, we

analyzed the expression of one non-iron-regulated transcript, *dnaB* (Figure 2D), as a

negative control (Brinkworth et al., 2018). Following 6 hrs of BpdI treatment, we

observed that the transcript expression of the periplasmic iron-binding protein *ytgA* was

significantly elevated 1.75-fold relative to the equivalent mock-treated time-point (p =

0.0052). However, we did not observe induction of *ytgA* transcript expression relative to

the 12 hpi time-point. We distinguish here between *elevated* and *induced* transcript

expression, as chlamydial gene expression is highly developmentally regulated. Thus, it

can be more informative to monitor longitudinal expression of genes, *i.e.* their induction,

as opposed to elevation relative to an equivalent control time-point, which may simply

represent a stall in development. While we did not observe induction of *ytgA*, which

would be more consistent with an iron-starved phenotype, we reason that this is a

consequence of the brief treatment period, and that longer iron starvation would

produce a more robust induction of iron-regulated transcripts. Note that the identification

of *ytgA* as iron-regulated has only been previously observed following extended periods

of iron chelation (J. D. Miller et al., 2009; Raulston et al., 2007; Thompson & Carabeo,

2011). Similarly, we observed that the transcript expression of the thioredoxin *ahpC* was

significantly elevated 2.15-fold relative to the equivalent mock-treated time-point (p =

0.038) but was not induced relative to the 12 hpi time-point. The transcript expression of

devB, encoding a 6-phosphogluconolactonase involved in the pentose phosphate

pathway, was not observed to significantly respond to our brief iron limitation condition,

suggesting that it is not a component of the primary iron starvation stress response in *C.*

trachomatis. As expected, the transcript expression of *dnaB*, a replicative DNA helicase,

was not altered by our iron starvation condition, consistent with its presumably iron-

independent regulation (Brinkworth et al., 2018). Overall, these data confirmed that our

6-hr BpdI treatment condition was suitable to produce a mild iron starvation phenotype

at the transcriptional level, thus facilitating our investigation of iron-dependent regulatory

integration.

Figure 2. Brief iron limitation condition produces mild iron-starved transcriptional
phenotype. (*A*) Total RNA and gDNA was harvested from infected HeLa cells at the
indicated times post-infection under iron-replete (teal) and -deplete (orange) conditions.
The transcript abundance of iron-regulated *ytgA*, (*B*) *ahpC*, (*C*) *devB* and (*D*) non-iron
regulated *dnaB* were quantified by RT-qPCR and normalized against genome copy
number. The transcript expression of *ytgA* and *ahpC* were significantly elevated
following 6-hour BpdI treatment, indicative or iron starvation to *C. trachomatis.* N=3.
Statistical significance was determined by One-Way ANOVA followed by post-hoc
pairwise *t*-tests with Bonferroni's correction for multiple comparisons. * = $p < 0.05$, ** = p
< 0.01, *** = $p < 0.005$.

Transcript expression of the *trpRBA* operon is differentially regulated by iron in

Chlamydia trachomatis. Upon identifying an iron limitation condition that produced a

relevant transcriptional phenotype while avoiding the onset of persistent development,

we aimed to investigate whether the immediate response to iron starvation in *C.*

trachomatis would result in the consistent induction of pathways unrelated to iron

utilization/acquisition, but nevertheless important for surviving immunological stress.

The truncated Trp biosynthetic operon, *trpRBA* (Figure 3A), has been repeatedly linked

to the ability of genital and LGV serovars (D-K and L1-3, respectively) of *C. trachomatis*

to counter IFN-γ-mediated stress. This is due to the capacity of the chlamydial Trp

synthase in these serovars to catalyze the β synthase reaction, *i.e.* the condensation of

indole to the amino acid serine to form Trp (Fehlner-Gardiner et al., 2002). In the

presence of exogenous indole, *C. trachomatis* is therefore able to biosynthesize Trp

such that it can prevent the development of IFN-γ-mediated persistence.

Correspondingly, the expression of *trpRBA* is highly induced following IFN-γ stimulation

of infected cells (Belland et al., 2003; Østergaard et al., 2016). These data have

historically implicated Trp starvation as the primary mechanism by which persistence

develops in *C. trachomatis* following exposure to IFN-γ. However, these studies have

routinely depended on prolonged treatment conditions that monitor the terminal effect of

persistent development, as opposed to the immediate molecular events which may

have important roles in the developmental fate of *Chlamydia.* As such, these studies

may have missed the contribution of other IFN-γ-stimulated insults such as iron

limitation.

To decouple Trp limitation from iron limitation and assess their relative

contribution to regulating a critical pathway for responding to IFN-γ-mediated stress, we

monitored the transcript expression of the *trpRBA* operon under brief Trp or iron

starvation by RT-qPCR. When starved for Trp for 6 hrs, we observed that the

expression of *trpR*, *trpB* and *trpA* were all significantly induced greater than 10.5-fold

relative to 12 hpi (p = 0.00077, 0.025 and 9.7e-5, respectively; Figure 3B). All three

ORFs were also significantly elevated relative to the equivalent mock-treated time-point

(p = 0.00076, 0.025 and 9.7e-5, respectively). This result was surprising with respect to

the relative immediacy of operon induction in response to our Trp starvation protocol,

confirming the relevant Trp-starved transcriptional phenotype. To induce Trp-deprived

persistence in *C. trachomatis*, many laboratories rely on compounded techniques of

IFN-γ pre-treatment to deplete host Trp pools in conjunction with culturing in Trp-

depleted media, among other strategies. While the phenotypic end-point differs here, it

is nonetheless interesting to note that only 6 hrs of media replacement is sufficient to

markedly up-regulate *trpRBA* expression. This suggests that *C. trachomatis* has a

highly attuned sensitivity to even moderate changes in Trp levels.

We next performed the same RT-qPCR analysis on the expression of the *trpRBA*

operon in response to 6 hrs of iron limitation via Bpdl treatment (Figure 3C). While we

observed that the transcript expression of all three ORFs was significantly elevated at

least 2.1-fold relative to the equivalent mock-treated time-point (p = 0.015, 0.00098 and

0.0062, respectively), we made the intriguing observation that only the expression of

trpB and *trpA* was significantly induced relative to 12 hpi (p = 0.00383 and 0.0195,

respectively). The significant induction of *trpBA* expression, but not *trpR* expression,

suggested that *trpBA* are specifically regulated by iron availability. This result is

consistent with a recent survey of the iron-regulated transcriptome in *C. trachomatis* by

RNA-Sequencing, which also reported that iron-starved *Chlamydia* specifically up-

regulate *trpBA* expression in the absence of altered *trpR* expression (Brinkworth et al.,

2018). Our results expand on this finding by providing a more detailed investigation into

the specific profile of this differential regulation of *trpRBA* in response to iron

deprivation. Taken together, these findings demonstrate that an important stress

response pathway, the *trpRBA* operon, is regulated by the availability of both Trp and

iron, consistent with the notion that the pathway may be cooperatively regulated to

respond to various stress conditions. Notably, iron-dependent regulation of Trp

biosynthesis has not been previously documented in other prokaryotes.

Figure 3. Expression of the *trpRBA* operon in *C. trachomatis* is differentially regulated
by brief iron limitation. (*A*) Cartoon depiction of the *trpRBA* operon (drawn to scale) with
the primary transcriptional start site upstream of *trpR* annotated. (*B*) Total RNA and
gDNA were harvested from infected HeLa cells at the indicated times post-infection
under Trp-replete (black) and -deplete (red) conditions. The transcript expression of
trpRBA operon was quantified by RT-qPCR and normalized against genome copy
number. All three ORFs are significantly induced relative to 12 hpi following Trp
starvation. N=2. (*C*) Total RNA and gDNA were harvested from infected HeLa cells at
the indicated times post-infection under iron-replete (blue) and -deplete (red) conditions.
The transcript expression of *trpRBA* operon was quantified by RT-qPCR and normalized
against genome copy number. Only *trpB* and *trpA* expression was significantly induced
relative to 12 hpi. N=3. Statistical significance was determined by One-Way ANOVA
followed by post-hoc pairwise *t*-tests with Bonferroni's correction for multiple
comparisons. * = $p < 0.05$, ** = $p < 0.01$, *** = $p < 0.005$.

Specific iron-regulated expression of *trpBA* originates from a novel alternative

transcriptional start site within the *trpRBA* intergenic region. We hypothesized that

the specific iron-related induction of *trpBA* expression relative to *trpR* expression may

be attributable to an iron-regulated alternative transcriptional start site (alt. T<u>S</u>S)

downstream of the *trpR* ORF. Indeed, a previous study reported the presence of an alt.

TSS in the *trpRBA* IGR, located 214 nucleotides upstream of the *trpB* translation start

position (Carlson et al., 2006). However, a parallel study could not identify a TrpR

binding site in the *trpRBA* IGR (Akers & Tan, 2006). We reasoned that a similar alt. TSS

may exist in the IGR that controlled the iron-dependent expression of *trpBA.* We

therefore performed <u>R</u>apid <u>A</u>mplification of <u>5</u>'-<u>c</u>DNA <u>E</u>nds (5'-RACE) on RNA isolated

from *C. trachomatis* L2-infected HeLa cells using the SMARTer 5'/3' RACE Kit workflow

(Takara Bio). Given the low expression of the *trpRBA* operon during normal

development, we utilized two sequential gene-specific amplification steps (nested 5'-

RACE) to identify 5' cDNA ends in the *trpRBA* operon. These nested RACE conditions

resulted in amplification that was specific to infected-cells (Figure 4 – Figure

Supplement 1A). Using this approach, we analyzed four conditions: 12 hpi, 18 hpi, 12

hpi + 6 hrs of Bpdl treatment, and 12 hpi + 6 hrs of Trp-depletion (Figure 4A). We

observed three RACE products that migrated with an apparent size of 1.5, 1.1 and 1.0

kilobases (kb). At 12 and 18 hpi, all three RACE products exhibited low abundance,

even following the nested PCR amplification. This observation was consistent with the

expectation that the expression of the *trpRBA* operon is very low under normal, iron and

Trp-replete conditions. However, we note that the 6-hr difference in development did

appear to alter the representation of the 5' cDNA ends, which may suggest a stage-

specific promoter utilization within the *trpRBA* operon. In our Trp starvation condition,

we observed an apparent increase in the abundance of the 1.5 kb RACE product, which

was therefore presumed to represent the primary TSS upstream of *trpR*, at nucleotide

position 511,389 (*C. trachomatis* L2 434/Bu). Interestingly, the 1.0 kb product displayed

a very similar apparent enrichment following BpdI treatment, suggesting that this RACE

product represented a specifically iron-regulated TSS. Both the 1.5 and 1.0 kb RACE

products were detectable in the Trp-depleted and iron-depleted conditions, respectively,

during the primary RACE amplification, consistent with their induction under these

conditions (Figure 4 – Figure Supplement 1B).

If iron depletion was inducing *trpBA* expression independent of *trpR,* we

reasoned that we would observe specific enrichment of *trpB* sequences in our 5'-RACE

cDNA samples relative to *trpR* sequences. We again utilized RT-qPCR to quantify the

abundance of *trpB* transcripts relative to *trpR* transcripts in the 5'-RACE total RNA

samples (Figure 4B). In agreement with our model, only under iron starved conditions

did we observe a significant enrichment of *trpB* relative to *trpR* ($p < 0.01$). Additionally,

we observed that at 12 and 18 hpi in iron-replete conditions, the ratio of *trpB* to *trpR* was

approximately 1.0, suggesting non-preferential basal expression across the three

putative TSSs. Another factor contributing to this ratio is the synthesis of the full-length

trpRBA polycistron. In support of this, the *trpB* to *trpR* ratio remained near 1.0 under the

Trp-starved condition, which would be expected during transcription read-through of the

whole operon. The apparent lack of preferential promoter utilization as described above

could be attributed to the relatively low basal expression of the operon at 12 and 18 hpi

under Trp- and iron-replete conditions, thus precluding quantitative detection of

differential promoter utilization in this assay.

476   To determine the specific location of the 5' cDNA ends within the *trpRBA* operon,

we isolated the 5'-RACE products across all conditions by gel extraction and cloned the

products into the pRACE vector supplied by the manufacturer. We then sequenced the

ligated inserts and BLASTed the sequences against the *C. trachomatis* L2 434/Bu

genome to identify the location of the 5'-most nucleotides (Figure 4C). These data are

displayed as a statistical approximation of the genomic regions most likely to be

represented by the respective 5'-RACE products in both histogram (semi-continuous)

and density plot (continuous) format (See Supplementary File 1 for a description of all

mapped 5'-RACE products). As expected, the 1.5 kb product mapped in a distinct and

tightly grouped peak near the previously annotated *trpR* TSS, with the mean and modal

nucleotide being 511,388 and 511,389, respectively (Figure 4 – Figure Supplement 2A).

Surprisingly, we found that neither the 1.1 or 1.0 kb RACE product mapped to the

previously reported alt. TSS in the *trpRBA* IGR, at position 511,826. Instead, we

observed that the 1.1 kb product mapped on average to nucleotide position 511,878,

with the modal nucleotide being found at 511,898 (Figure 4 – Figure Supplement 2B).

The 1.0 kb product mapped with a mean nucleotide position of 512,013, with the modal

nucleotide being 512,005 (Figure 4 – Figure Supplement 2C), only 35 bases upstream

of the *trpB* coding sequence. Interestingly, the 1.0 kb product mapped to a region of the

trpRBA IGR flanked by consensus σ^{66} -10 and -35 promoter elements, found at

positions 512,020-5 and 511,992-7, respectively (Ricci, Ratti, & Scarlato, 1995). These

data collectively pointed toward the 1.0 kb 5'-RACE product representing a novel, iron-

regulated alt. TSS and *bona fide* σ^{66}-dependent promoter element that allows for the

specific iron-dependent expression of *trpBA*.

Figure 4. Iron-dependent induction of *trpBA* expression initiates within the *trpRBA* intergenic region from a novel alternative transcriptional start site. (*A*) Total RNA was harvested from infected HeLa cells at the indicated times post-infection to examine iron-dependent and Trp-dependent changes in the 5'-cDNA profile of the *trpRBA* operon by Rapid Amplification of 5' cDNA Ends (5'-RACE). RACE products were separated on an agarose gel, revealing three distinct and specific bands with apparent sizes of 1.5, 1.1 and 1.0 kb. Trp depletion led to the apparent enrichment of the 1.5 kb product, while BpdI treatment produced a similarly enriched 1.0 kb RACE product. Figure shows representative experiment of three biological replicates. (*B*) To confirm that iron-dependent induction of *trpBA* could originate from alternative transcription initiation, RT-qPCR was performed on 5'-RACE total RNA to quantify the abundance of *trpB* transcripts relative to *trpR.* Only under iron-limited conditions were *trpB* transcripts enriched relative to *trpR*. N=3. Statistical significance determined by One-way ANOVA followed by post-hoc pairwise *t*-tests. * = $p < 0.05$, ** = $p < 0.01$, *** = $p < 0.005$. (*C*) The nucleotide position of the 5' cDNA ends generated from RACE were mapped to the *C. trachomatis* L2 434/Bu genome by nucleotide BLAST. Figure displays histogram (semi-continuous; bin width=20) and overlaid density plot (continuous) distribution of 5' nucleotide positions generated from each 5'-RACE product. The dotted line represents the weighted mean of the distribution, as indicated by the integer value above each line. The identified alt. TSSs are depicted on the *trpRBA* operon (drawn to scale) above the plot. N=3.

YtgR specifically binds to the *trpRBA* intergenic region in an operator-dependent

manner to repress transcription of *trpBA*. As the only known iron-dependent

transcriptional regulator in *Chlamydia*, we hypothesized that YtgR may regulate the iron-

dependent expression of *trpBA* from the putative promoter element we characterized by

5'-RACE. Using bioinformatic sequence analysis, we investigated whether the *trpRBA*

IGR contained a candidate YtgR operator sequence. By local sequence alignment of

the putative YtgR operator sequence (Akers et al., 2011) and the *trpRBA* IGR, we

identified a high-identity alignment (76.9% identity) covering 67% of the putative

operator sequence (Figure 5A). Interestingly, this alignment mapped to the previously

identified palindrome suspected to have operator functionality (Carlson et al., 2006). By

global sequence alignment of the YtgR operator to the palindromic sequence, an

alignment identical to the local alignment was observed, which still displayed relatively

high sequence identity (43.5% identity). We hypothesized that this sequence functioned

as an YtgR operator, despite being located 184 bp upstream of the *trpBA* alt. TSS.

To investigate the ability of YtgR to bind and repress transcription from the

putative *trpBA* promoter, we implemented a heterologous two-plasmid assay that

reports on YtgR repressor activity as a function of β-galactosidase expression

(Thompson et al., 2012). In brief, a candidate DNA promoter element was cloned into

the pCCT expression vector between an arabinose-inducible pBAD promoter and the

reporter gene *lacZ*. This plasmid was co-transformed into BL21 (DE3) *E. coli* along with

an IPTG-inducible pET151 expression vector with (pET151-YtgR) or without (pET151-

EV) the C-terminal 139 amino acid residues of CTL0325 (YtgC). Note that we have

previously demonstrated that this region is a functional iron-dependent repressor

domain (Thompson et al., 2012). To verify the functionality of this assay, we determined

whether ectopic YtgR expression could repress pCCT reporter gene expression in the

presence of three candidate DNA elements: a no-insert empty vector (pCCT-EV), the

putative promoter element for *C. trachomatis lpdA* (pCCT-*lpdA*), and the promoter

region of the *ytgABCD* operon (pCCT-*ytgABCD*; Figure 5B). As expected, from the

pCCT-EV reporter construct, ectopic YtgR expression did not significantly reduce the

activity of β-galactosidase. Additionally, reporter gene expression from pCCT-*lpdA*,

containing the promoter of iron-regulated *lpdA* (Brinkworth et al., 2018), which is not

known to be YtgR-regulated, was not affected by ectopic expression of YtgR. This

demonstrated that the assay can discriminate between the promoter elements of iron-

regulated genes and *bona fide* YtgR targeted promoters. Indeed, in the presence of

pCCT-*ytgABCD*, induction of YtgR expression produced a significant decrease in β-

galactosidase activity (p = 0.03868) consistent with its previously reported auto-

regulation of this promoter (Thompson et al., 2012).

Using this same assay, we then inserted into the pCCT reporter plasmid 1) the

trpR promoter element (pCCT-*trpR*), 2) the putative *trpBA* promoter element

represented by the IGR (pCCT-*trpBA*), and 3) the same putative *trpBA* promoter

element with a mutated YtgR operator sequence that was diminished for both

palindromicity and A-T richness, two typical features of prokaryotic promoter elements

(pCCT-*trpBA*ΔOperator; Figure 5C) (Schmitt, 2002; Tao, Boydt, & Murphy, 1992). When

YtgR was ectopically expressed in the pCCT-*trpR* background, we observed no

statistically distinguishable change in β-galactosidase activity. However, in the pCCT-

trpBA background, ectopic YtgR expression significantly reduced β-galactosidase

activity at levels similar to those observed in the pCCT-*ytgABCD* background (p =

0.01219). This suggested that YtgR was capable of binding to the *trpBA* promoter

element specifically. Interestingly, this repression phenotype was abrogated in the

pCCT-*trpBA*∆Operator background, where we observed no statistically meaningful

difference in β-galactosidase activity. We subsequently addressed whether the region of

the *trpRBA* IGR containing the YtgR operator site was sufficient to confer YtgR

repression in this assay (Figure 5 – Figure Supplement 1). We cloned three fragments

of the *trpRBA* IGR into the pCCT reporter plasmid: the first fragment represented the 5'-

end of the IGR containing the operator site at the 3'-end (pCCT-IGR1), the second

fragment represented a central region of the IGR containing the operator site at the 5'-

end (pCCT-IGR2), and the third fragment represented the 3'-end of the IGR and did not

contain the operator site (pCCT-IGR3). Surprisingly, we observed that none of these

fragments alone were capable of producing a significant repression phenotype in our

reporter system. This finding indicated that while the operator site was necessary for

YtgR repression, it alone was not sufficient. Together, these data indicated that YtgR

could bind to the *trpBA* promoter element and that this binding was dependent upon an

intact AT-rich palindromic sequence, likely representing an YtgR operator, but that

further structural elements in the *trpRBA* IGR may be necessary for repression.

Nonetheless, we demonstrated the existence of a functional YtgR binding site that

conferred iron-dependent transcriptional regulation to *trpBA*, independent of the major

trpR promoter.

Figure 5. Ectopically expressed YtgR domain is capable of binding the putative *trpBA* promoter element in an operator-specific manner to repress transcription in a heterologous system. (*A*) Identification of putative YtgR operator sequence by local and global nucleotide sequence alignment using EMBOSS Water and Needle algorithms, respectively, to align the previously identified YtgR operator to both the *trpRBA* IGR and palindromic candidate sequence. The palindrome was then mutated in our YtgR repression assay as depicted to abolish palindromicity and AT-richness. (*B*) Ectopic expression of YtgR significantly represses β-galactosidase activity only from the promoter of its own operon, *ytgABCD*, and not from an empty vector or another iron-regulated but presumably non-YtgR targeted promoter, *lpdA*. N = 2 or 3. (*C*) Expression of recombinant YtgR represses β-galactosidase activity from the putative *trpBA* promoter element, but not the *trpR* promoter, and this repression is dependent on the unaltered operator sequence identified in Fig. 5A. N = 2 or 3. Statistical significance determined by two-sided unpaired Student's *t*-test with Welch's correction for unequal variance. * = $p < 0.05$, ** = $p < 0.01$, *** = $p < 0.005$.

Transcripts initiated at the primary *trpR* promoter terminate at the YtgR operator site. We hypothesized that YtgR binding at the *trpRBA* YtgR operator site may disadvantage the processivity of RNAP reading-through the IGR from the upstream *trpR* promoter. Similar systems of RNAP read-through blockage have been reported; the transcription factor Reb1p "roadblocks" RNAPII transcription read-through in yeast by promoting RNAP pausing and subsequent degradation (Colin et al., 2014). To investigate this question, we first returned to RNA-Sequencing data we generated to define the immediate iron-dependent transcriptional regulon in *C. trachomatis* (Brinkworth et al., 2018). Using data obtained from *C. trachomatis*-infected HeLa cells at 12 hpi + 6h mock or BpdI treatment, we mapped the sequenced reads in batch across three biological replicates to the *C. trachomatis* L2 434/Bu genome (NC_010287) which we modified to include annotations for non-operonic IGRs. This coverage map indicated that under BpdI-treated conditions, a higher proportion of reads mapped to the *trpRBA* IGR (IGR_trpB) relative to mock-treatment (Figure 6A). However, a notable increase in reads mapping to the upstream *trpR* CDS was not observed, suggesting that under iron replete conditions, transcripts originating from the primary *trpR* promoter may be terminated before reading through the IGR. Note that this is consistent with the original observation that *trpR* is not differentially expressed under iron-limited conditions in this RNA-Seq dataset (Brinkworth et al., 2018). We additionally investigated the abundance of reads mapping to the IGRs upstream of *euo* (IGR_euo; not iron-regulated, not YtgR-regulated) and *lpdA* (IGR_lpdA; iron-regulated, not YtgR-regulated), and were unable to observe a similar increase in read coverage at these IGRs following BpdI treatment, indicating that the increased coverage at the *trpRBA* IGR was specific (Figure 6 –

Figure Supplement 1A-B). The absolute number of reads mapping to the *trpRBA* IGR

under these conditions was very low relative to either upstream or downstream CDS,

implying that the terminated transcript species are rare. We therefore turned to more

sensitive and quantitative methods to interrogate possible transcript termination within

the *trpRBA* IGR.

To identify transcription termination sites (T<u>T</u>Ss) in the *trpRBA* operon in *C.*

trachomatis, we utilized 3'-RACE to map the 3'-ends of transcripts using gene-specific

primers within the *trpR* CDS (Figure 6B; bottom). We again utilized two RACE

amplification cycles to generate distinct, specific bands suitable for isolation and

sequencing (Figure 6 – Figure Supplement 2B-C). By gel electrophoresis of the 3'-

RACE products, we observed the appearance of four distinct bands that migrated with

an apparent size of 0.55, 0.45, 0.40 and 0.20 kb. In our Trp-depleted condition, we

observed only a very weak amplification of the 2.5 – 3 kb full-length *trpRBA* message by

3'-RACE (Figure 6 – Figure Supplement 2A). However, we did observe it across all

replicates. To confirm that the full-length product was specific to the Trp-deplete

treatment, we amplified the *trpRBA* operon by RT-PCR from the 3'-RACE total RNA

(Figure 6B; top). As expected, only in the Trp-deplete sample did we observe robust

amplification of the full-length *trpRBA* message. We note however that image contrast

adjustment reveals a very weak band present in all experimental samples. In

accordance with the RNA-Sequencing data, 3'-RACE demonstrated the presence of

unique transcription termination events in the *trpRBA* operon.

To identify the specific TTS locations, we gel extracted the four distinct 3'-RACE

bands across all conditions and cloned them into the pRACE sequencing vector as was

done for the 5'-RACE experiments. We then sequenced the inserted RACE products

and mapped them to the *C. trachomatis* L2 434/Bu genome (Figure 6B). This revealed a

highly dynamic TTS landscape within the *trpRBA* IGR, which has not previously been

investigated (For a full description of mapped 3'-RACE products, see Supplementary

File 2). The 0.20 kb RACE product mapped to the 3'-end of the *trpR* CDS, with a mean

nucleotide position of 511,665 and a modal nucleotide position of 511,667 (Figure 6 –

Figure Supplement 3A). Contrastingly, the other three 3'-RACE products did not map in

such a way so as to produce specific, unambiguous modal peaks. Instead, their

distribution was broader and more even, with only a few nucleotide positions mapping

more than once. Accordingly, the 0.45 kb product mapped with an average nucleotide

position of 511,889, just downstream of the 1.1 kb 5'-RACE product (Figure 6 – Figure

Supplement 3C), while the 0.55 kb product mapped with an average nucleotide position

of 511,986, upstream of the 1.0 kb 5'-RACE product (Figure 6 – Figure Supplement

3D). Interestingly, the 0.40 kb product mapped to a region directly overlapping the

putative YtgR operator site, with a mean nucleotide position of 511,811 (Figure 6 –

Figure Supplement 3B). We therefore reasoned that this putative TTS may have an

iron-dependent function.

Figure 6. Transcription from the primary *trpR* promoter terminates in the *trpRBA* IGR region, notably at the YtgR operator site. (*A*) Coverage map of RNA-Sequencing reads mapped to the *C. trachomatis* L2/434 Bu genome (NC_010287) edited to contain annotations for non-operonic IGRs. Read coverage at the *trpRBA* IGR (IGR_trpB) is increased following Bpdl treatment, but *trpR* read coverage is not similarly increased. (*B*) Total RNA was harvested from *C. trachomatis*-infected HeLa cells to analyze 3'-cDNA landscape downstream of the *trpR* promoter. The top panel displays representative RT-PCR of full-length *trpRBA* message across experimental conditions (NTC = No Template Control). Bottom panel depicts electrophoresed 3'-RACE products and estimated sizes. Intensity plot to the right of image was generated using the Fiji Dynamic ROI Profiler plugin to monitor intensity across the 18 hpi condition. Note the presence of four distinct peaks, corresponding to each 3'-RACE product. N=3. (*C*) 3'-RACE products were sequenced and mapped to *C. trachomatis* L2 434/Bu genome by nucleotide BLAST. Figure displays histogram (semi-continuous; bin width=20) and overlaid density plot (continuous) distribution of 3' nucleotide positions generated from each 3'-RACE product. The dotted line represents the weighted mean of the distribution, as indicated by the integer value above each line. The identified alt. TTSs are depicted on the *trpRBA* operon (drawn to scale) above the plot. The 0.40 kb RACE product mapped to a region overlapping the predicted YtgR operator site. N=3.

Iron limitation promotes transcription read-through at the YtgR operator site.

Given the observation that transcripts terminated at the YtgR operator site, we

hypothesized that YtgR binding may promote transcription termination at this locus.

Conversely, we hypothesized that inactivating YtgR DNA-binding by BpdI treatment

would allow transcription to read through the YtgR operator site. To quantitatively

analyze the possibility that iron-depletion, and thus dissociation of YtgR, may facilitate

transcription read-through at the operator site, we utilized RT-qPCR to monitor the

abundance of various amplicons across the *trpRBA* operon (Figure 7A). We quantified

these data in relation to a "read-through" normalization amplicon that, based on 5'- and

3'-RACE data, should only be represented when the full-length *trpRBA* message is

transcribed (Figure 7A). The representation of a specific mRNA species relative to the

full-length transcript should therefore be interpretable through a simple ratio of the

experimental amplicon to the "read-through" amplicon. If an mRNA species is poorly

represented relative to the full-length transcript, the ratio should be approximately 1.0.

Conversely, if that species is over-represented relative to the full-length transcript, the

ratio should exceed 1.0 (Figure 7B; left). Therefore, as each amplicon is increasingly

represented as a part of the full-length transcript, the ratio of the specific amplicon to the

normalization amplicon should approach 1.0 (Figure 7B; right).

We first analyzed an amplicon from nucleotide 511,416 – 531 to monitor the

relative abundance of transcript species associated with transcription initiating at the

trpR promoter (Figure 7C). We observed that the representation of this amplicon was

not significantly altered following iron limitation relative to 12 hpi, suggesting that the

depletion of iron was not affecting initiation of transcription at the *trpR* promoter.

Interestingly, at 18 hpi, the representation ratio of this amplicon significantly shifted

further away from 1.0 (p = 0.00358), indicating that at 18 hpi this amplicon is

represented less as a component of read-through transcription relative to 12 hpi. As

expected, under Trp-deplete conditions, the representation ratio shifted significantly

closer to 1.0 (p = 0.00064), consistent with read-through transcription of the full-length

trpRBA message.

We then preformed the same analysis on an amplicon from nucleotide 511,639 –

764, immediately upstream of the TTS at the YtgR operator site to monitor condition-

dependent read-through at this site (Figure 7D). We again observed that at 18 hpi, the

representation ratio was significantly increased (p = 0.01046), and following Trp-

depletion, the ratio was significantly decreased (p = 0.00023), as expected. Notably,

and consistent with our hypothesis, we observed that the representation ratio of this

amplicon was also significantly closer to 1.0 following iron limitation (p = 0.00407),

suggesting that transcription read-through was increased at this site under iron limited

conditions. Indeed, if YtgR is dissociating from the operator site during iron depletion, a

greater proportion of transcripts would be expected to read-through this locus.

Finally, we analyzed an amplicon from nucleotide 513,856 – 968, at the very 3'-

end of *trpA* to asses changes in terminal transcription under our experimental conditions

(Figure 7E). At 18 hpi, we observed a significant increase in the representation ratio of

this amplicon (p = 0.00476), which is likely attributable to both basal levels of alternative

transcription from the IGR as well as poor transcription read-through of the full-length

message. Following 6 hrs of BpdI treatment, we also observed a significant increase in

the representation ratio of this amplicon (p = 0.01510), which supports the finding that

trpBA is being preferentially transcribed under this condition, distinct from the full-length

trpRBA transcript. We were only able to detect a marginal decrease in the

representation of this amplicon under Trp-depleted conditions (p = 0.07942), which may

suggest that the very 3'-end of *trpRBA* is under-represented relative to our

normalization amplicon, which falls within the middle of the operon. In fact, recent work

has reported on the relatively poor representation of 3'-end mRNAs in *Chlamydia*

(Ouellette, Rueden, & Rucks, 2016). In sum, this set of experiments provides evidence

that iron-depletion specifically alters the representation of particular mRNA species

across the *trpRBA* operon. Additionally, they implicate iron-dependent YtgR DNA-

binding as the mediator of these effects. By alleviating YtgR repression via iron

depletion, transcription is allowed to proceed through the operator site, albeit at basal

levels. Concomitantly, transcription is specifically activated at the downstream alt. TSS

for *trpBA*.

Figure 7. Iron limitation promotes transcription read-through at the YtgR operator site. (*A*) Cartoon depiction of amplicons analyzed in these experiments across the *trpRBA* operon, with the identified TSSs and YtgR operator site anotated. Drawn to scale. (*B*) Schematic representation of RT-qPCR analysis. On the left, a biological interpretation of the ratio used to determine the relative transcript abundance is provided. When a termination product (red) is poorly represented, *e.g.* read-through is high, the ratio of specific amplicon (blue) to full-length transcript (grey) should be close to 1.0. When the termination product is abundant, *e.g.* read-through is low, then the ratio should exceed 1.0. On the right, a graphical demonstration of this concept is provided using mock data. The dotted line represents the theoretical asymptote at a value of 1.0. (*C*) Analysis of transcription read-through by RT-qPCR was performed on 3'-RACE total RNA at three distinct loci across the *trpRBA* operon representing upstream transcription initiation (511,416-531), (*D*) YtgR operator site termination (511,639-764) and (*E*) terminal *trpBA* transcription (513,856-968). Abundance of each amplicon was normalized to a region (Read-through) predicted to be transcribed only as a part of the full-length product based on 5' and 3'-RACE data (511,792-512,080). Thus, the ratio of each amplicon to the normalization amplicon represents the proportion of that amplicon encoded as part of the full-length transcript. At the YtgR operator termination site, iron limitation reduces the ratio relative to 12 hpi, suggesting that transcription read-through increases at this site under this condition. Statistical significance determined by One-way ANOVA followed by post-hoc pairwise *t*-tests. * = $p < 0.05$, ** = $p < 0.01$, *** = $p < 0.005$.

Figure 8. Model for proposed mechanism of iron-dependent YtgR-mediated regulation of *trpRBA* expression. Iron is imported through the YtgABCD ABC-type metal permease complex. YtgR is cleaved from the YtgCR permease-repressor fusion protein. In the presence of sufficient iron, holo-YtgR can bind to the *trpRBA* IGR to both terminate basal transcription from the primary *trpR* promoter and repress transcription initiation at the alternative *trpBA* promoter. Iron depletion inactivates YtgR DNA-binding, thus promoting read-through of basal transcription from the *trpR* promoter while also inducing transcription at the downstream *trpBA* promoter.

Discussion

In this study, we provide a mechanistic explanation for the specific iron-limited

induction of *trpBA* expression mediated by the repressor YtgR, representing a novel

instance of integrated stress adaptation in *Chlamydia*. Utilizing an infected-epithelial cell

culture model, we identified a previously undescribed iron-regulated promoter element

within the *trpRBA* IGR responsible for the iron-limited induction of *trpBA* expression

independent of *trpR.* Using *in silico*, biochemical and chemical-genetic methods, we

demonstrate that YtgR binds the *trpRBA* IGR to regulate iron-dependent *trpBA*

expression. Importantly, transcriptional repression in our heterologous system was

shown to be dependent on an unaltered operator sequence that bears significant

homology to the previously defined operator element in the *ytg* promoter. Previously

published reports have demonstrated that YtgR is capable of directly binding DNA

sequences containing this operator *in vitro* (Akers et al., 2011; Thompson et al., 2012).

Furthermore, our infected-cell culture studies revealed that transcripts originating from

the primary *trpR* promoter terminate within the IGR, notably at the putative YtgR

operator site, and that transcription read-through at this locus is iron-dependent. Thus,

we propose that YtgR regulates *trpBA* expression at two levels: repression of the *trpBA*

promoter and premature termination of the major transcript generated from the *trpR*

promoter (Figure 8). This is the first time an iron-dependent mode of regulation has

been shown to control the expression of tryptophan biosynthesis in prokaryotes, which

reflects the uniquely specialized nature of *C. trachomatis*.

While we demonstrate here that iron-dependent *trpBA* expression originates from

a novel promoter element immediately upstream of the *trpB* CDS, this is not the first

description of an alt. TSS within the *trpRBA* IGR. Carlson, *et al*. identified an alt. TSS

within the IGR which they suggested was responsible for *trpBA* expression (Carlson et

al., 2006). In these studies, we were unable to confirm the presence of the previously

identified alt. TSS by 5'-RACE. This is likely because Carlson, *et al*. examined the

presence of transcript origins following 24 hrs of Trp starvation whereas here we

monitored immediate responses to stress following only 6 hrs of treatment. Prolonged

Trp depletion would result in a more homogeneously stressed population of chlamydial

organisms that may exhibit the same preferential utilization of the promoter identified by

Carlson, *et al*., the detection of which is precluded in a more heterogeneous, transiently-

stressed population. This may explain the observation of multiple T(S/T)Ss across the

trpRBA operon in our studies. However, the contribution of such a Trp-dependent alt.

TSS as identified by Carlson *et al.* to the general stress response of *C. trachomatis*

remains unclear given its association with presumably abnormal organisms. Does

utilization of this alt. TSS indicate abnormal growth or a *bona fide* stress adaptation?

Moreover, Akers & Tan were unable to verify TrpR binding to the *trpRBA* IGR by EMSA,

suggesting that some other Trp-dependent mechanism may control transcription from

this site (Akers & Tan, 2006). Ultimately, our approach of investigating more immediate

responses to stress revealed previously unreported mechanisms functioning to regulate

Trp biosynthesis in *C. trachomatis*, underscoring the value of transient as opposed to

sustained induction of stress.

Another mechanism of regulation reported to control the chlamydial *trpRBA*

operon is Trp-dependent transcription attenuation. Based on sequence analysis, a

leader peptide has been annotated within the *trpRBA* IGR (Merino & Yanofsky, 2005).

Presumably, this functions analogously to the attenuator in the *E. coli trpEDCBA*

operon; Trp starvation causes ribosome stalling at sites of enriched Trp codons such

that specific RNA secondary structures form to facilitate RNAP read-thru of downstream

sequences – in this case, *trpBA* (Yanofsky, 1981). However, robust experimental

evidence to support the existence of attenuation in *C. trachomatis* is lacking. To date,

the only experimental evidence that supports this model was reported by Carlson, *et al*.,

who demonstrated that in a TrpR-mutant genetic background, an additional increase in

trpBA expression could be observed following 24 hr Trp-depletion. (Carlson et al.,

2006). However, this could be attributable to an alternative Trp-dependent, *but TrpR-*

independent mechanism controlling *trpBA* expression at the alt. TSS identified by

Carlson, *et al*. None of the data presented here point conclusively to the existence of a

Trp-dependent attenuator. The additional termination sites identified in our 3'-RACE

assay may represent termination events mediated by an attenuator, but without more

specific analysis utilizing mutated sequences we cannot attribute attenuator function to

those termination sites.

Interestingly, in *Bacillus subtilis,* Trp-dependent attenuation of transcription takes

on a form markedly different from that in *E. coli*. Whereas attenuation functions in *cis* for

the *E. coli trp* operon, *B. subtilis* utilize a multimeric <u>T</u>ryptophan-activated <u>R</u>NA-binding

<u>A</u>ttenuation <u>P</u>rotein, TRAP, which functions in *trans* to bind *trp* operon RNA under Trp-

replete conditions, promoting transcription termination and inhibiting translation

(Gollnick, Babitzke, Antson, & Yanofsky, 2005). This interaction is antagonized by anti-

TRAP in the absence of charged tRNA^{Trp}, leading to increased expression of TRAP

regulated genes. We suggest that YtgR may represent the first instance of a separate

and distinct clade of attenuation mechanisms: iron-dependent *trans*-attenuation. This

mechanism may function independently of specific RNA secondary structure, relying

instead on steric blockage of RNAP processivity, but ultimately producing a similar

result. Possible regulation of translation remains to be explored. The recent

development of new genetic tools to alter chromosomal sequences and generate

conditional knockouts in *C. trachomatis* should enable a more detailed analysis of

trpRBA regulation, including possible *trans*-attenuation (Mueller, Wolf, & Fields, 2016;

Ouellette, 2018).

As a Trp auxotroph, what might be the biological significance of iron-dependent

YtgR regulation of the *trpRBA* operon in *C. trachomatis*? We have already noted the

possibility that iron-dependent *trpBA* regulation in *C. trachomatis* may enable the

induction of a similar response to both Trp and iron starvation, stimuli likely mediated by

IFN-γ *in vivo*. This mechanism also presents the opportunity for *C. trachomatis* to

respond similarly to distinct *sequential* stresses, where a particular stress may prime the

pathogen to better cope with subsequent stresses. To reach the female upper genital

tract (UGT), where most significant pathology is identified following infection with *C.*

trachomatis, the pathogen must first navigate the lower genital tract (LGT). *Chlamydia*

infections of the female LGT are associated with bacterial vaginosis (BV), which is

characterized by obligate and facultative anaerobe colonization, some of which produce

indole (Sasaki-Imamura, Yoshida, Suwabe, Yoshimura, & Kato, 2011; Ziklo, Huston,

Hocking, & Timms, 2016). This provides *C. trachomatis* with the necessary substrate to

salvage tryptophan via TrpBA. Interestingly, the LGT is also likely an iron-limited

environment. Pathogen colonization and BV both increase the concentration of mucosal

lactoferrin (Lf), an iron-binding glycoprotein, which can starve pathogens for iron (Spear

et al., 2011; Valenti et al., 2018). Lf expression is additionally hormone-regulated, and

thus the LGT may normally experience periods of iron limitation (Cohen, Britigan,

French, & Bean, 1987; Kelver et al., 1996). Moreover, the expression of TfR is

constrained to the basal cells of the LGT stratified squamous epithelium (Lloyd,

O'Dowd, Driver, & Tee, 1984), which likely restricts necessary Tf-bound iron from *C.*

trachomatis infecting the accessible upper layers of the stratified epithelia (Nogueira,

Braun, & Carabeo, 2017; Ouellette & Carabeo, 2010).

For *C. trachomatis*, iron limitation may therefore serve as a critical signal in the

LGT, inducing the expression of *trpBA* such that Trp is stockpiled from available indole,

allowing the pathogen to counteract impending IFN-γ-mediated Trp starvation. We

suggest the possibility that iron limitation in the LGT may be a significant predictor of

successful pathogen colonization in the UGT. Unfortunately, testing these hypotheses in

cell culture models of infection presents a significant challenge. Evaluating rescue of

chlamydial growth in the presence of indole to specifically assess the iron-dependent

role of *trpBA* requires simultaneous Trp and iron depletion. The former ensures indole

utilization by the bacteria, and the latter de-represses YtgR-regulated *trpBA* expression.

In theory, this is feasible, but in practice the combined stress rapidly induces aberrant

development, muddying results obtained from such studies (data not shown). Ideally,

genetic approaches could be employed to distinguish the regulatory effects of YtgR

independent of TrpR. However, the genetic manipulation of *trans*-acting factors (*e.g.*

YtgR) will presumably have unpredictable off-target effects. Genetically altering *cis*-

acting factors – such as operator sequences – is more feasible, but at present we lack

the information necessary to rationally mutate these sequences in *C. trachomatis* to

interrogate these questions. The tight regulatory coordination at both the transcription

initiation and termination steps would likely mean any mutation in the *cis*-acting

sequences would affect both processes indiscriminately. Furthermore, *in vivo* infection

models present challenges: attempting to answer these questions will likely require the

use of *in vivo* non-human primate studies, as mouse models of *Chlamydia*-infection do

not recapitulate immune-mediated Trp starvation (Nelson et al., 2005). Ultimately, these

limitations do not undermine the biological significance of an iron-dependent mode of

regulating Trp salvage, given the critical role played by this pathway during infection.

Finally, and of note, the expression of the ribonucleotide diphosphate-reductase-

encoding *nrdAB* was also recently shown to be iron-regulated in *C. trachomatis*

(Brinkworth et al., 2018). The regulation of *nrdAB* is known to be mediated by the

presumably deoxyribonucleotide-dependent transcriptional repressor NrdR, encoded

distal to the *nrdAB* locus (Case, Akers, & Tan, 2011). As NrdR activity is not known to

be modulated by iron availability, this raises the intriguing possibility that here too a

unique iron-dependent mechanism of regulation may integrate the chlamydial stress

response to promote a unified response across various stress conditions. Future studies

may require more metabolomics-based approaches to thoroughly dissect the integration

of these stress responses, as transcriptome analyses alone often miss broader,

pathway-oriented metabolic coordination. Ultimately, these studies point towards a need

to carefully re-evaluate the molecular stress response in *Chlamydia*, with greater

emphasis on the use of targeted approaches and treatment protocols that induce stress,

but not persistence. We anticipate that the rapid progress of the field in recent years will

continue to catalyze exciting and important discoveries regarding the fundamental

biology of *Chlamydia*.

Materials and Methods

Eukaryotic Cell Culture and Chlamydial Infections
Human cervical epithelial adenocarcinoma HeLa (ATCC® CCL-2) cells were cultured at
37° C with 5% atmospheric CO_2 in Dulbecco's Modified Eagle Medium (DMEM)
supplemented with 10 μg/mL gentamicin, 2 mM L-glutamine, and 10% (*v/v*) filter
sterilized fetal bovine serum (FBS). For all experiments, HeLa cells were cultured
between passage numbers 4 and 16. *Chlamydia trachomatis* serovar L2 (434/Bu) was
originally obtained from Dr. Ted Hackstadt (Rocky Mountain National Laboratory,
NIAID). Chlamydial EBs were isolated from infected HeLa cells at 36-40 hours post-
infection (hpi) and purified by density gradient centrifugation essentially as described
(Caldwell, Kromhout, & Schachter, 1981).
1035
For the infection of 6-well tissue culture plates, HeLa cells cultured to 80-90%
confluency were first washed with pre-warmed Hanks Buffered Saline Solution (HBSS)
prior to the monolayer being overlaid with inoculum (un-supplemented DMEM) at the
indicated multiplicity of infection (MOI). Tissue culture plates were then centrifuged at 4°
C with a speed of 1000 RPM (Eppendorf 5810 R table top centrifuge, A-4-81 rotor) for 5
minutes to synchronize the infection. Inoculum was aspirated and cells were washed
again with pre-warmed HBSS prior to the media being replaced with pre-warmed
complete DMEM. Infected cultures were then returned to the tissue culture incubator
until the indicated times post-infection. This procedure was replicated exactly for the
infection of 24-well tissue culture plates.
1046
Iron Starvation
Chlamydia trachomatis L2-infected HeLa cell cultures were starved for iron by
supplementation of the media with the iron chelator 2,2-bipyridyl (Bpdl; Sigma Aldrich,
St. Louis, MO, USA; CAS: 366-18-7) essentially as described (Thompson & Carabeo,
2011). Briefly, at the indicated times post-infection, infected cell cultures were washed
with pre-warmed HBSS prior to the addition of complete DMEM (mock) or complete
DMEM supplemented with 100 μM Bpdl. Infected cell cultures were returned to the
incubator for the indicated treatment periods. Bpdl was prepared as a 100 mM stock
solution in 100% ethanol and stored at -20° C for no longer than 6 months.
1056
Tryptophan Starvation
Chlamydia trachomatis L2-infected HeLa cell cultures were starved for tryptophan by
replacement of complete DMEM with tryptophan-depleted medium. In brief, Tryptophan-
replete or –deplete DMEM-F12 (U.S. Biological Life Sciences, Salem, MA, USA)
powder media was prepared following manufacture instructions and supplemented with
10% (*v/v*) filter-sterilized FBS which had been previously dialyzed 16-20h at 4° C in PBS
in a 10 kDa MWCO dialysis cassette. Media was then further supplemented with 10
μg/mL gentamicin. At the indicated times post-infection, complete DMEM was aspirated

and wells were washed with pre-warmed HBSS prior to the addition of tryptophan-
replete or –deplete medium. Infected cell cultures were returned to the incubator for the
indicated treatment periods.
1068

1069 **Cloning**
1070 All constructs were cloned using standard molecular cloning techniques, *e.g.* restriction
1071 enzyme, homology-directed, etc. All primers and plasmids used in this study can be
1072 found in Supplementary file 3 and 4, respectively.
1073

Immunofluorescent Confocal Microscopy
At the indicated times post-infection, *C. trachomatis* L2-infected HeLa cell cultures
seeded on glass coverslips in 24-well tissue cultures plates were first washed with pre-
warmed HBSS prior to fixation with 4% paraformaldehyde (PFA) in phosphate buffered
saline (PBS) for 20 minutes at RT ° C. Fixation solution was aspirated and wells were
washed with PBS prior to permeabilization with 0.2% Triton X-100 in PBS for 5 minutes
at RT° C. Permeabilization solution was then decanted and cells were washed with
PBS. The coverslips were blocked for 30 minutes with 1% bovine serum albumin (BSA)
in PBS at RT° C. To stain for *Chlamydia*, coverslips were washed with PBS and PBS
supplemented with 1% BSA and 1:500 convalescent human sera was added to wells
and incubated at RT° C for 1 hour with rocking. Primary antibody solution was decanted
and coverslips were again washed with PBS. Goat anti-human Alexa-647 (Invitrogen,
ThermoFisher Scientific, Waltham, MA, USA) diluted 1:1000 in PBS with 1% BSA was
then added to the wells and incubated in the dark for another hour at RT° C with
rocking. Secondary antibody solution was then decanted, coverslips were washed again
with PBS and coverslips were either immediately mounted on microscopy slides using
ImmuMount (ThermoFisher Scientific) or VectaShield H-1000 (Vector Laboratories,
Burlingame, CA, USA) or stored in the dark at 4° C until mounting. All images were
acquired on a Leica TCS SP8 laser scanning confocal microscope, using identical
settings, in the Integrative Physiology and Neuroscience Advanced Imaging Center at
Washington State University. All images are Z-projections and were processed in Fiji
(Schindelin et al., 2012) and Adobe Creative Suite identically for each comparative time-
point.
1097

Nucleic Acid Preparation
RNA was harvested from *C. trachomatis*-infected HeLa cell monolayers by scraping 3
wells of a 6-well plate in ice-cold Trizol® Reagent (ThermoFisher Scientific). Samples
were then pooled and split into two technical replicates (RT-qPCR) or kept as one
biological replicate (RACE). Trizol®-extracted samples were then thoroughly vortexed
with a 100 μL volume of Zirconia beads prior to chloroform extraction. 100% ethanol
was added to the aqueous phase and RNA was isolated using the Ambion®
RiboPure™ RNA Purification kit for bacteria following manufacturer instructions
(ThermoFisher Scientific). DNA was removed from RNA samples using the Invitrogen
DNA-*free*™ DNA Removal Kit following manufacturer instructions (ThermoFisher
Scientific). RNA was stored at -20° C until further use.
1109

cDNA was generated using either SuperScript® IV Reverse Transcriptase (RT-qPCR;
ThermoFisher Scientific) or SMARTScribe™ Reverse Transcriptase (RACE and RACE-
specific qRT-PCR); Takara Bio, Kusatsu, Shiga Prefecture, Japan) essentially as
described by the respective manufacturers. For cDNA generated for RT-qPCR, 650 ng
of total RNA was used as a template in a 20 μL total reaction volume. For every RT
reaction, a "no-RT" control, generated from 350 ng of total RNA template in a 10 μL total
volume, was included. For 5'-RACE, cDNA was generated from 250 ng of total RNA
using random primers in a 10 μL total volume and further processed in the RACE
workflow. cDNA was stored at -20° C.

1119

gDNA was harvested from *C. trachomatis*-infected HeLa cell monolayers by scraping 3
wells of a 6-well plate in ice-cold PBS + 10% Proteinase K (ThermoFisher Scientific).
Samples were then pooled and split into two technical replicates for analysis of genome
copy number by qPCR. gDNA was isolated using the DNeasy Blood and Tissue Kit
following manufacturer protocols (QIAGEN, Hilden, Germany). gDNA was stored at -20°
C until further use.

1126

Reverse Transcription Quantitative Polymerase Chain Reaction (RT-qPCR)
cDNA (or gDNA in qPCR), prepared as described above, was diluted 1:10 or 1:100 in
nuclease-free H_2O depending on the experimental condition being assayed (*e.g.*
treatment, point in development cycle, etc.). On ice, 3.3 μL of diluted sample was added
to 79 μL of PowerUp™ SYBR® Green Master Mix (ThermoFisher Scientific) with
specific qPCR primers diluted to 500 nM. From this master mix, each experimental
sample was assayed in triplicate 25 μL reactions. Assays were run on an Applied
Biosystems 7300 Real Time PCR System with cycling conditions as follows: Stage 1:
50.0° C for 2 min, 1 rep. Stage 2: 95.0° C for 10 min, 1 rep. Stage 3: 95.0° C for 15 sec,
40 reps. Stage 4: 60.0° C for 1 min, 1 rep. Primers were subjected to dissociation curve
analysis to ensure that a single product was generated. For each primer set, a standard
curve was generated using purified *C. trachomatis* L2 gDNA from EB preparations
diluted from 2 x 10^{-3} to 2 x10^0 ng per reaction. Ct values generated from each
experimental reaction were then fit to standard curves (satisfying an efficiency of
95±5%) for the respective primer pair and from the calculated ng quantities, transcript or
genome copy number was calculated as follows:

1143

$$1144 \quad Genome\ copy\ number\ (\frac{genome\ copies}{ng\ total\ gDNA}) = \frac{ng\ genome\ \times\ df}{ng\ total\ gDNA} \times \frac{892,000\ copies}{ng\ DNA}$$

1145

$$1146 \quad Transcript\ copy\ number\ \left(\frac{transcript\ copies}{\frac{genome\ copies}{ng\ total\ gDNA}}\right) = \frac{ng\ transcript\ \times\ df}{\frac{genome\ copies}{ng\ total\ gDNA}} \times \frac{892,000\ copies}{ng\ DNA}$$

1147

Where *df* = dilution factor and the number of copies/ng DNA is calculated based on the
size of the *C. trachomatis* L2 genome assuming that the molar mass per base pair is
650 (g/mol)/bp (note that this value should be the same for any single-copy ORF on the
genome). All quantifications of genome copy number were determined using the *ahpC*

qPCR primer set. Values from replicate assays were averaged, and values from
replicate RNA/gDNA isolations were averaged to obtain the mean and standard
deviation for one biological replicate. For some experiments, to account for batch effects
across biological replicates, data was transformed such that the mean of all samples in
each replicate was identical. In some instances, batch correcting generated negative
values, and in this case data sets were scaled such that the lowest value equaled 1.0.
HeLa cells were infected at an MOI of 2 for all RT-qPCR studies.
1159

5' Rapid Amplification of cDNA Ends (5'-RACE)
All RACE studies were performed using the SMARTer® RACE 5'/3' Kit (Takara Bio). To
observe 5'-RACE products from the *trpRBA* operon, a "nested" RACE protocol was
used as outlined in the SMARTer® RACE 5'/3' Kit user manual. Briefly, 1.25-2.5 μL of
cDNA generated for RACE was added to a 25 μL reaction volume and run in a thermal
cycler for 40 cycles using the touch-down PCR conditions described by the
manufacturer. In brief, 5 cycles were run at an annealing temperature of both 72° C and
70° C prior to 30 cycles run with an annealing temperature of 68° C. Following this
primary amplification, the RACE products were diluted 1:50 in Tricine-EDTA Buffer
supplied by the manufacturer, and 2.5 μL of diluted primary RACE product was added
to a 25 μL reaction volume and subjected to another 20 cycles of nested PCR, as
described by the manufacturer, using primers designed within the amplicon of the
primary RACE products. Samples were electrophoresed on a 2% agarose gel for
visualization and analysis. HeLa cells were infected at a MOI of 5 for all RACE studies.
1174

3' Rapid Amplification of cDNA Ends (3'-RACE)
3'-RACE studies were performed essentially identical to 5'-RACE with the exception
that total RNA was subjected to poly(A) tailing with a Poly(A) Polymerase following
manufacturer instructions (New England Biolabs, Ipswich, MA, USA). In brief, at least
3.5 μg of total RNA was incubated at 37° C with Poly(A) Polymerase in reaction buffer
supplemented with ATP and murine RNase Inhibitor (New England Biolabs) for 30
minutes prior to heat-inactivation at 65° C for 20 minutes. RNA was re-isolated through
an RNA clean-up filter cartridge (Ambion, ThermoFisher Scientific). A total of 125 ng of
poly(A)-tailed total RNA was then used to generate 3'-RACE ready cDNA in a 10 μL
reaction volume following manufacturer instructions. Primary and nested RACE was
performed using 3'-RACE gene-specific primers following the same protocol for
amplification described for 5'-RACE, with the exception that the extension time was
adjusted to accommodate amplification of the full ~3 kb *trpRBA* polycistronic message.
1188

Mapping of 5'/3'-RACE Products
5'-RACE products generated from either primary or nested RACE reactions were
excised from the agarose gel and DNA was isolated using the NucleoSpin Gel and PCR
Clean-up kit (Macherey-Nagel, Takara Bio). The isolated RACE products were then
cloned into the pRACE vector supplied in the SMARTer® RACE 5'/3' Kit using the In-
Fusion HD cloning kit (Takara Bio). Ligated vectors were transformed into chemically
competent Stellar™ *E. coli* cells by heat shock. Transformed bacteria were plated on LB
agar containing 50 μg/mL carbenicillin and incubated overnight at 37° C. Colonies were
selected and screened for relevant inserts by PCR. Positive colonies were cultured

overnight at 37° C in LB liquid broth containing 50 μg/mL carbenicillin and plasmids
were isolated using the QIAprep Spin Miniprep kit (QIAGEN). Inserts were then
sequenced by Eurofins Genomics using the default M13 Reverse sequencing primer.
Returned sequencing data was aligned to the *C. trachomatis* L2 (434/Bu) genome
(NCBI Accession: NC_010287) by BLAST and the most 5' aligned nucleotide was
considered the 5' end of the insert. In the case of 3'-RACE data, the reverse
complement sequence was first generated prior to alignment. Grouping of individual
products was determined 1.) by clusters being greater than 30 nucleotides apart and 2.)
by the specific RACE band that the alignment was derived from. These two criteria were
not both satisfied in all cases and in those cases criteria 1.) was favored.
1208
Sequence Alignments
All *C. trachomatis* L2 434/Bu genome sequences were obtained from NCBI Accession
NC_010287. Global pairwise sequence alignments were made using the EMBOSS
Needle algorithm. Alignment parameters were set as follows: Matrix: DNAfull, Gap
Open: 20, Gap Extend: 0.8, Output Format: pair, End Gap Penalty: True, End Gap
Open: 10, End Gap Extend: 0.5. These conditions were sufficient to replicate the
previously published alignment between the putative YtgR operator sequence and the
TroR operator (Akers et al., 2011). Local pairwise sequence alignments were made
using the EMBOSS Water algorithm. The putative YtgR operator was aligned to the
entire 348 bp intergenic region of the *trpRBA* operon (*C. trachomatis* L2 [434/Bu]
genome position 511,692-512,039). The alignment parameters were set as follows:
Matrix: DNAfull, Gap Open:10, Gap Extend: 0.5, Output Format: pair. These are the
default conditions and were chosen to remove bias from the alignment results.
1222
Two-Plasmid Reporter Assay
The YtgR-binding reporter assay was performed essentially as described, with minor
modifications (Thompson et al., 2012). Promoter regions of interest were amplified from
the *C. trachomatis* L2 (434/Bu) genome by PCR using the indicated primer sets, which
included KpnI restriction endonuclease sites at the 5' and 3' ends of the promoter
amplicon. The amplified fragments and the pCCT-EV plasmid were then KpnI-digested
and the promoters ligated into the vector using T4 or Quick Ligase (New England
BioLabs). Insert directionality was confirmed by directional colony PCR and positive
clones were sequence verified. pCCT-*trpBA*ΔOperator was cloned by amplifying two
fragments of the pCCT-*trpBA* vector with one ~60mer primer containing the bases to be
substituted for each fragment. Thus, the whole vector was split into two half-fragments
containing the substituted bases. The two fragments were then cloned back together
using In-Fusion Homology-Directed cloning (Takara Bio) to yield the final vector.
Electrocompetent BL21(DE3) *E. coli* (Sigma Aldrich) were co-transformed by
electroporation with the pCCT reporter plasmid and the pET151 expression vector (-EV
or –YtgR) and plated on double selective LB agar containing 50 μg/mL carbenicillin and
15 μg/mL tetracycline. Prior to plating of transformed cells, 50 μL of 40 mg/mL X-Gal in
DMSO (EMD Millipore, Burlington, MA, USA) was applied to the plate for colorimetric
determination of β-galactosidase expression. Transformants were incubated overnight
at 37° C. The following evening, blue colonies from each experimental condition were
selected and cultured overnight in LB liquid broth containing 0.2% (*w/v*) D-glucose (for

catabolite repression of expression vectors), 50 μg/mL carbenicillin and 15 μg/mL
tetracycline. Cultures were incubated overnight at 37° C. The following morning,
overnight cultures were spun down to remove glucose-containing media and sub-
cultured in LB liquid broth medium containing 50 μM FeSO$_4$, 50 μg/mL carbenicillin and
15 μg/mL tetracycline to an OD$_{600}$ of 0.45. Cultures were incubated for 1 hour at 37° C
and sub-cultured a second time in the same media to an OD$_{600}$ of 0.1. Cultures were
returned to the incubator for another hour prior to the addition of 500 μM isopropyl β-D-
1-thiogalactopyranoside (IPTG) to induce pET151 expression from the *lac* promoter.
Cultures were incubated another hour prior to the addition of 0.2% L-arabinose to
induce *lacZ* expression from the *araBAD* promoter. Cultures were incubated a final 2
hours prior to the collection of a 0.1 mL volume of cells for assaying β-galactosidase
activity by the Miller Assay (J. H. Miller, 1972). Cell pellets were stored at -80° C prior to
being assayed. To assay β-galactosidase activity, cell pellets were first re-suspended in
Z-buffer (pH 7.0, 60 mM Na$_2$HPO$_4$, 40 mM NaH$_2$PO$_4$, 10 mM KCl, 1 mM MgSO$_4$ and 2.7
μL/mL β-mercaptoethanol). 50 μL of 0.1% SDS and 100 μL of chloroform were then
added to each sample prior to thorough vortexing. Samples were equilibrated for 5
minutes at 30° C and 200 μL of 4 mg/mL ortho-nitrophenyl-β-galactoside (ONPG)
prepared in Phosphate Buffer (pH 7.0, 60 mM Na$_2$HPO$_4$, 40 mM NaH$_2$PO$_4$) were added
to the samples to initiate the reaction. Reactions were stopped by the addition of 500 μL
1 M Na$_2$CO$_3$. Absorbance was measured on a FLUOStar Optima plate reader (BMG
Labtech, Offenburg, Germany) at 420 nm and Miller Units were calculated as:
1265

$$1000 \times \frac{Abs_{420}}{t \times v \times OD_{final}}$$

1267
Where t = reaction time, v = volume of cells and OD_{final} = OD$_{600}$ at the time of sample
collection. It was empirically determined that the subtraction of absorbance at 550 nm
had a negligible effect on the calculated value. A blank sample lacking cells was
included in each experimental batch and used as a reference for absorbance. For each
experimental condition, three independent co-transformed colonies were assayed in
technical triplicate. In some instances, significantly high Miller Unit outliers were
excluded by Grubb's Test (p < 0.05) under the assumption that extreme *lacZ* expression
may reflect plasmid copy number or reporter gene expression issues.
1276
1277 RNA-Sequencing
RNA-Sequencing experiments were performed as described in their original publication
(Brinkworth et al., 2018). Coverage maps were generated by mapping all reads across
three biological replicates to a single reference file in CLC Genomics Workbench v11.
To facilitate easy analysis of IGR boundaries, the *C. trachomatis* L2 434/Bu genome
(Accession: NC_010287) was modified to contain annotations for non-operonic
intergenic regions, and this genome was used as the reference for read mapping. Read
mapping was performed using default settings in CLC Genomics Workbench. Data
aggregation in the Reads track was set to aggregate above 1bp.
1286
1287 Graphs and Statistical Analysis

All graphs were generated using the ggplot2 package (Wickham, 2009) in R Studio, in
Excel and/or in the Adobe Creative Suite. All line plots and bar graphs represent the
mean ± one standard deviation unless otherwise noted. All box and whisker plots
represent the distribution of data between the 1st and 3rd quartile range within the box,
while the whiskers represent data within 1.5 interquartile ranges of the 1st or 3rd quartile.
Extreme values outside this range are plotted as open circles. The 2nd quartile (median)
is plotted as a black line within the box. Histogram plots were generated with a bin width
of 20 and are plotted on a density scale. The overlaid density plots represent a
statistical approximation of the data over a continuous scale. All statistical analyses
were carried out in R Studio. All statistical computations were performed on the mean
values of independent biological replicates calculated from the indicated number of
respective technical replicates. For single pairwise comparisons, a two-sided unpaired
Student's t-test with Welch's correction for unequal variance was used to determine
statistical significance. For multiple pairwise comparisons, a One-Way Analysis of
Variance (ANOVA) was conducted to identify significant differences within groups. If a
significant difference was detected, then the indicated post-hoc pairwise test was used
to identify the location of specific statistical differences. A p-value less than 0.05 was
considered statistically significant. For all figures, * = $p < 0.05$, ** = $p < 0.01$, and *** = p
< 0.005.
1307

Acknowledgements. We thank Liam Caven, Korinn Murphy and Matthew Romero for

critical review of this manuscript; Dr. Christopher C. Thompson for the establishment of

the *E. coli* YtgR reporter system and generation of the pCCT construct; Dr. Scot P.

Ouellette for support, critical feedback and advice; and Dr. David Dewitt for expert

advice, training and maintenance of equipment in the IPN Advanced Imaging Center.

This work was supported by NIH grants R01-AI065545 to R.A.C.; 1F31AI136295 and

5T32GM008336 to N.D.P.; N.D.P. was also supported by an Achievement Reward for

College Scientists (ARCS; Seattle Chapter) Fellowship.

References

AbdelRahman, Y. M., & Belland, R. J. (2005). The chlamydial developmental cycle.

FEMS Microbiology Reviews, *29*(5), 949–959.

http://doi.org/10.1016/j.femsre.2005.03.002

Akers, J. C., HoDac, H., Lathrop, R. H., & Tan, M. (2011). Identification and functional

analysis of CT069 as a novel transcriptional regulator in Chlamydia. *Journal of*

Bacteriology, *193*(22), 6123–6131. http://doi.org/10.1128/JB.05976-11

Akers, J. C., & Tan, M. (2006). Molecular mechanism of tryptophan-dependent

transcriptional regulation in Chlamydia trachomatis. *Journal of Bacteriology*,

188(12), 4236–4243. http://doi.org/10.1128/JB.01660-05

Belland, R. J., Nelson, D. E., Virok, D., Crane, D. D., Hogan, D., Sturdevant, D., …

Caldwell, H. D. (2003). Transcriptome analysis of chlamydial growth during IFN-

gamma-mediated persistence and reactivation. *Proceedings of the National*

Academy of Sciences of the United States of America, *100*(26), 15971–15976.

http://doi.org/10.1073/pnas.2535394100

Brinkworth, A. J., Wildung, M. R., & Carabeo, R. A. (2018). Genomewide Transcriptional

Responses of Iron-Starved Chlamydia trachomatis Reveal Prioritization of

Metabolic Precursor Synthesis over Protein Translation. *MSystems*, *3*(1), e00184-

17.

Brown, S. A., Palmer, K. L., & Whiteley, M. (2008). Revisiting the host as a growth

medium. *Nature Reviews Microbiology*, *6*(9), 657–666.

http://doi.org/10.1038/nrmicro1955

Byrd, T. F., & Horwitz, M. A. (1993). Regulation of transferrin receptor expression and

ferritin content in human mononuclear phagocytes: Coordinate upregulation by iron

transferrin and downregulation by interferon gamma. *Journal of Clinical*

Investigation, *91*(3), 969–976. http://doi.org/10.1172/JCI116318

Byrd, T., & Horwitz, M. A. (1989). Interferon gamma-activated human monocytes

downregulate transferrin receptors and inhibit the intracellular multiplication of

Legionella pneumophila by limiting the availability of iron. *Journal of Clinical Investigation*, *83*(5), 1457–1465. http://doi.org/10.1172/JCI114038

Byrne, G. I., Lehmann, L. K., & Landry, G. J. (1986). Induction of tryptophan catabolism is the mechanism for gamma-interferon-mediated inhibition of intracellular Chlamydia psittaci replication in T24 cells. *Infection and Immunity*, *53*(2), 347–351.

Caldwell, H. D., Kromhout, J., & Schachter, J. (1981). Purification and partial characterization of the major outer membrane protein of Chlamydia trachomatis. *Infection and Immunity*, *31*(3), 1161–1176.

Carlson, J. H., Porcella, S. F., Mcclarty, G., & Caldwell, H. D. (2005). Comparative Genomic Analysis of Chlamydia trachomatis Oculotropic and Genitotropic Strains. *Infection and Immunity*, *73*(10), 6407–6418. http://doi.org/10.1128/IAI.73.10.6407

Carlson, J. H., Wood, H., Roshick, C., Caldwell, H. D., & McClarty, G. (2006). In vivo and in vitro studies of Chlamydia trachomatis TrpR:DNA interactions. *Molecular Microbiology*, *59*(6), 1678–1691. http://doi.org/10.1111/j.1365-2958.2006.05045.x

Case, E. D. R., Akers, J. C., & Tan, M. (2011). CT406 encodes a chlamydial ortholog of NrdR, a repressor of ribonucleotide reductase. *Journal of Bacteriology*, *193*(17), 4396–4404. http://doi.org/10.1128/JB.00294-11

Cassat, J. E., & Skaar, E. P. (2013). Iron in Infection and Immunity. *Cell Host and Microbe*, *13*(5), 509–519. http://doi.org/10.1016/j.chom.2013.04.010

CDC. (2017). *2016 Sexually Transmitted Diseases Surveillance - Chlamydia*. Retrieved from https://www.cdc.gov/std/stats16/chlamydia.htm

Clarke, I. N. (2011). Evolution of Chlamydia trachomatis. *Annals of the New York Academy of Sciences*, *1230*, 11–18. http://doi.org/10.1111/j.1749-

6632.2011.06194.x

Cohen, M., Britigan, B., French, M., & Bean, K. (1987). Preliminary observations on

lactoferrin secretion in human vaginal mucus: variation during the menstrual cycle,

evidence of hormonal regulation, and implications for infection with Neisseria

gonorrhoeae. *American Journal of Obstetrics and Gynecology*, *157*(5), 1122–1125.

Colin, J., Candelli, T., Porrua, O., Boulay, J., Zhu, C., Lacroute, F., … Libri, D. (2014).

Roadblock termination by reb1p restricts cryptic and readthrough transcription.

Molecular Cell, *56*(5), 667–680. http://doi.org/10.1016/j.molcel.2014.10.026

Dill, B. D., Dessus-Babus, S., & Raulston, J. E. (2009). Identification of iron-responsive

proteins expressed by Chlamydia trachomatis reticulate bodies during intracellular

growth. *Microbiology*, *155*(1), 210–219. http://doi.org/10.1099/mic.0.022731-0

Eisenreich, W., Dandekar, T., Heesemann, J., & Goebel, W. (2010). Carbon metabolism

of intracellular bacterial pathogens and possible links to virulence. *Nature Reviews*

Microbiology, *8*(6), 401–412. http://doi.org/10.1038/nrmicro2351

Fehlner-Gardiner, C., Roshick, C., Carlson, J. H., Hughes, S., Belland, R. J., Caldwell,

H. D., & McClarty, G. (2002). Molecular basis defining human Chlamydia

trachomatis tissue tropism: A possible role for tryptophan synthase. *Journal of*

Biological Chemistry, *277*(30), 26893–26903.

http://doi.org/10.1074/jbc.M203937200

Gollnick, P., Babitzke, P., Antson, A., & Yanofsky, C. (2005). Complexity in regulation of

tryptophan biosynthesis in Bacillus subtilis. *Annual Review of Genetics*,

39(October), 47–68. http://doi.org/10.1146/annurev.genet.39.073003.093745

Hafner, L. M. (2015). Pathogenesis of fallopian tube damage caused by Chlamydia

trachomatis infections. *Contraception*, *92*(2), 108–115.

http://doi.org/10.1016/j.contraception.2015.01.004

Hood, M. I., & Skaar, E. P. (2012). Nutritional immunity: Transition metals at the

pathogen-host interface. *Nature Reviews Microbiology*, *10*(8), 525–537.

http://doi.org/10.1038/nrmicro2836

Igietseme, J. U., Ananaba, G. A., Candal, D. H., Lyn, D., & Black, C. M. (1998). Immune

control of Chlamydial growth in the human epithelial cell line RT4 involves multiple

mechanisms that include nitric oxide induction, tryptophan catabolism and iron

deprivation. *Microbiology and Immunology*, *42*(9), 617–625.

Kelver, M. E., Kaul, A., Nowicki, B., Findley, W. E., Hutchens, T. W., & Nagamani, M.

(1996). Estrogen regulation of lactoferrin expression in human endometrium.

American Journal of Reproductive Immunology, *36*(5), 243–247.

Lloyd, J. M., O'Dowd, T., Driver, M., & Tee, D. E. (1984). Demonstration of an epitope of

the transferrin receptor in human cervical epithelium--a potentially useful cell

marker. *Journal of Clinical Pathology*, *37*(2), 131–135. Retrieved from

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Ci

tation&list_uids=6198338

Macchiarulo, A., Camaioni, E., Nuti, R., & Pellicciari, R. (2009). Highlights at the gate of

tryptophan catabolism : a review on the mechanisms of activation and regulation of

indoleamine 2,3-dioxygenase (IDO), a novel target in cancer disease. *Amino Acids*,

37, 219–229. http://doi.org/10.1007/s00726-008-0137-3

Merino, E., & Yanofsky, C. (2005). Transcription attenuation: A highly conserved

regulatory strategy used by bacteria. *Trends in Genetics*, *21*(5), 260–264.

http://doi.org/10.1016/j.tig.2005.03.002

Miller, J. D., Sal, M. S., Schell, M., Whittimore, J. D., & Raulston, J. E. (2009).

Chlamydia trachomatis YtgA is an iron-binding periplasmic protein induced by iron

restriction. *Microbiology*, *155*(9), 2884–2894. http://doi.org/10.1099/mic.0.030247-0

Miller, J. H. (1972). Assay of β-galactosidase. In *Experiments in Molecular Genetics*.

Cold Spring Harbor Laboratory Press.

Moore, E. R., & Ouellette, S. P. (2014). Reconceptualizing the chlamydial inclusion as a

pathogen-specified parasitic organelle: an expanded role for Inc proteins. *Frontiers*

in Cellular and Infeciton Microbiology, *4*(October), 1–10.

http://doi.org/10.3389/fcimb.2014.00157

Mueller, K. E., Wolf, K., & Fields, K. A. (2016). Gene Deletion by Fluorescence-

Reported Allelic Exchange Mutagenesis in Chlamydia trachomatis. *MBio*, *7*(1), 1–9.

http://doi.org/10.1128/mBio.01817-15.Invited

Nairz, M., Fritsche, G., Brunner, P., Talasz, H., Hantke, K., & Weiss, G. (2008).

Interferon-γ limits the availability of iron for intramacrophage Salmonella

typhimurium. *European Journal of Immunology*, *38*(7), 1923–1936.

http://doi.org/10.1002/eji.200738056

Nelson, D. E., Virok, D. P., Wood, H., Roshick, C., Johnson, R. M., Whitmire, W. M., …

Caldwell, H. D. (2005). Chlamydial IFN-gamma immune evasion is linked to host

infection tropism. *Proceedings of the National Academy of Sciences of the United*

States of America, *102*(30), 10658–63. http://doi.org/10.1073/pnas.0504198102

Newman, L., Rowley, J., Hoorn, S. Vander, Wijesooriya, N. S., Unemo, M., Low, N., …

Temmerman, M. (2015). Global Estimates of the Prevalence and Incidence of Four

Curable Sexually Transmitted Infections in 2012 Based on Systematic Review and

Global Reporting. *PLoS ONE*, *10*(12), 1–17.

http://doi.org/10.1371/journal.pone.0143304

Nogueira, A. T., Braun, K. M., & Carabeo, R. A. (2017). Characterization of the Growth

of Chlamydia trachomatis in In Vitro-Generated Stratified Epithelium. *Frontiers in*

Cellular and Infection Microbiology, *7*(October), 1–16.

http://doi.org/10.3389/fcimb.2017.00438

Østergaard, O., Follmann, F., Olsen, A. W., Heegaard, N. H., Andersen, P., &

Rosenkrands, I. (2016). Quantitative Protein Profiling of Chlamydia trachomatis

Growth Forms Reveals Defense Strategies Against Tryptophan Starvation.

Molecular & Cellular Proteomics : MCP, *15*(12), 3540–3550.

http://doi.org/10.1074/mcp.M116.061986

Ouellette, S. P. (2018). Feasibility of a Conditional Knockout System for Chlamydia

Based on CRISPR Interference. *Frontiers in Cellular and Infection Microbiology*,

8(February). http://doi.org/10.3389/fcimb.2018.00059

Ouellette, S. P., & Carabeo, R. A. (2010). A functional slow recycling pathway of

transferrin is required for growth of Chlamydia. *Frontiers in Microbiology*, *1*(OCT),

1–12. http://doi.org/10.3389/fmicb.2010.00112

Ouellette, S. P., Rueden, K. J., & Rucks, E. A. (2016). Tryptophan codon-dependent

transcription in chlamydia pneumoniae during gamma interferon-mediated

tryptophan limitation. *Infection and Immunity*, *84*(9), 2703–2713.

http://doi.org/10.1128/IAI.00377-16

Pokorzynski, N. D., Thompson, C. C., & Carabeo, R. A. (2017). Ironing Out the

Unconventional Mechanisms of Iron Acquisition and Gene Regulation in

Chlamydia. *Frontiers in Cellular and Infection Microbiology*, *7*(September), 1–19.

http://doi.org/10.3389/fcimb.2017.00394

Raulston, J. E. (1997). Response of Chlamydia trachomatis serovar E to iron restriction

in vitro and evidence for iron-regulated chlamydial proteins. *Infection and Immunity*,

65(11), 4539–4547.

Raulston, J. E., Miller, J. D., Davis, C. H., Schell, M., Baldwin, A., Ferguson, K., & Lane,

H. (2007). Identification of an iron-responsive protein that is antigenic in patients

with Chlamydia trachomatis genital infections. *FEMS Immunology and Medical*

Microbiology, *51*(3), 569–576. http://doi.org/10.1111/j.1574-695X.2007.00336.x

Ray, K., Marteyn, B., Sansonetti, P. J., & Tang, C. M. (2009). Life on the inside: The

intracellular lifestyle of cytosolic bacteria. *Nature Reviews Microbiology*, *7*(5), 333–

340. http://doi.org/10.1038/nrmicro2112

Ricci, S., Ratti, G., & Scarlato, V. (1995). Transcriptional regulation in the Chlamydia

trachomatis pCT plasmid. *Gene, 154*(1), 93–98. http://doi.org/10.1016/0378-

1119(94)00825-D

Sasaki-Imamura, T., Yoshida, Y., Suwabe, K., Yoshimura, F., & Kato, H. (2011).

Molecular basis of indole production catalyzed by tryptophanase in the genus

Prevotella. *FEMS Microbiology Letters*, *322*(1), 51–59.

http://doi.org/10.1111/j.1574-6968.2011.02329.x

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., …

Cardona, A. (2012). Fiji: An open-source platform for biological-image analysis.

Nature Methods, *9*(7), 676–682. http://doi.org/10.1038/nmeth.2019

Schmitt, M. P. (2002). Analysis of a DtxR-Like Metalloregulatory Protein, MntR, from

Corynebacterium diphtheriae That Controls Expression of an ABC Metal

Transporter by an Mn(2+)-Dependent Mechanism. *Journal of Bacteriology*, *184*(24),

6882–6892. http://doi.org/10.1128/JB.184.24.6882

Seth-Smith, H. M. B., Harris, S. R., Persson, K., Marsh, P., Barron, A., Bignell, A., …

Clarke, I. N. (2009). Co-evolution of genomes and plasmids within Chlamydia

trachomatis and the emergence in Sweden of a new variant strain. *BMC Genomics*,

10, 239. http://doi.org/10.1186/1471-2164-10-239

Skaar, E. P. (2010). The battle for iron between bacterial pathogens and their vertebrate

hosts. *PLoS Pathogens*, *6*(8), 1–2. http://doi.org/10.1371/journal.ppat.1000949

Spear, G. T., Kendrick, S. R., Chen, H. Y., Thomas, T. T., Bahk, M., Balderas, R., …

Landay, A. L. (2011). Multiplex immunoassay of lower genital tract mucosal fluid

from women attending an urban STD clinic shows broadly increased IL1ß and

lactoferrin. *PLoS ONE*, *6*(5), 1–7. http://doi.org/10.1371/journal.pone.0019560

Stephens, R. S., Kalman, S., Lammel, C., Fan, J., Marathe, R., Aravind, L., …

Gunderson, J. H. (1998). Genome sequence of an obligate intracellular pathogen of

humans: Chlamydia trachomatis. *Science*, *282*(5389), 754–759.

http://doi.org/10.1126/science.282.5389.754

Tao, X., Boydt, J., & Murphy, J. R. (1992). Specific binding of the diphtheria tox

regulatory element DtxR to the tox operator requires divalent heavy metal ions and

a 9-base-pair interrupted palindromic sequence. *Proceedings of the National*

Academy of Sciences of the United States of America, *89*(13), 5897–5901.

Taylor, H. R., Burton, M. J., Haddad, D., West, S., & Wright, H. (2014). Trachoma. *The*

Lancet, *384*(9960), 2142–2152. http://doi.org/10.1016/S0140-6736(13)62182-0

Taylor, M. W., & Feng, G. S. (1991). Relationship between interferon-gamma,

indoleamine 2,3-dioxygenase, and tryptophan catabolism. *FASEB J.*, *5*(11), 2516–

2522. Retrieved from http://www.fasebj.org/cgi/reprint/5/11/2516.pdf

Thompson, C. C., & Carabeo, R. A. (2011). An optimal method of iron starvation of the

obligate intracellular pathogen, Chlamydia trachomatis. *Frontiers in Microbiology*,

2(20). http://doi.org/10.3389/fmicb.2011.00020

Thompson, C. C., Nicod, S. S., Malcolm, D. S., Grieshaber, S. S., & Carabeo, R. A.

(2012). Cleavage of a putative metal permease in Chlamydia trachomatis yields an

iron-dependent transcriptional repressor. *Proceedings of the National Academy of*

Sciences of the United States of America, *109*(26), 10546–51.

http://doi.org/10.1073/pnas.1201398109

Thomson, N. R., Holden, M. T. G., Carder, C., Lennard, N., Lockey, S. J., Marsh, P., …

Clarke, I. N. (2008). Chlamydia trachomatis : Genome sequence analysis of

lymphogranuloma venereum isolates. *Genome Research*, *18*(1), 161–171.

http://doi.org/10.1101/gr.7020108.

Valenti, P., Rosa, L., Capobianco, D., Lepanto, M. S., Schiavi, E., Cutone, A., …

Mastromarino, P. (2018). Role of Lactobacilli and Lactoferrin in the Mucosal

Cervicovaginal Defense. *Frontiers in Immunology*, *9*(March), 1–14.

http://doi.org/10.3389/fimmu.2018.00376

Wickham, H. (2009). *ggplot2: Elegant Graphics for Data Analysis.* Springer-Verlag New

York.

Wood, H., Fehlner-Gardner, C., Berry, J., Fischer, E., Graham, B., Hackstadt, T., …

McClarty, G. (2003). Regulation of tryptophan synthase gene expression in

Chlamydia trachomatis. *Molecular Microbiology*, *49*(5), 1347–1359.

http://doi.org/10.1046/j.1365-2958.2003.03638.x

Wood, H., Roshick, C., & McClarty, G. (2004). Tryptophan recycling is responsible for

the interferon-? resistance of Chlamydia psittaci GPIC in indoleamine dioxygenase-

expressing host cells. *Molecular Microbiology*, *52*(3), 903–916.

http://doi.org/10.1111/j.1365-2958.2004.04029.x

Wyrick, P. B. (2010). Chlamydia trachomatis Persistence In Vitro: An Overview. *The*

Journal of Infectious Diseases, *201*, 88–95. http://doi.org/10.1086/652394

Yanofsky, C. (1981). Attenuation in the control of expression of bacterial operons.

Nature, *289*(5800), 751–758. http://doi.org/10.1038/289751a0

Ziklo, N., Huston, W. M., Hocking, J. S., & Timms, P. (2016). Chlamydia trachomatis

Genital Tract Infections: When Host Immune Response and the Microbiome

Collide. *Trends in Microbiology*, *24*(9), 750–765.

http://doi.org/10.1016/j.tim.2016.05.007

1543

Author Contributions: N.D.P. and R.A.C. wrote the manuscript; A.J.B. and R.A.C.

designed and analyzed data for the RNA-Sequencing experiments; A.J.B.

performed RNA-Seq experiments; N.D.P. and R.A.C. re-analyzed the RNA-

Sequencing data for this publication; N.D.P. and R.A.C. designed all other

experiments; N.D.P. performed all other experiments; N.D.P. and R.A.C. analyzed

and interpreted the data for all other experiments.

1550

Supplementary Figures

Figure 1 – Figure Supplement 1. Brief media-defined tryptophan limitation does not produce characteristically persistent *Chlamydia.* (*A*) *C. trachomatis* L2-infected HeLa cells were fixed and stained with convalescent human sera to image inclusion morphology by confocal microscopy following tryptophan limitation at the indicated times post-infection. Figure shows representative experiment of two biological replicates. Scale bar = 15 μm, Inset scale bar = 5 μm. (*B*) Genomic DNA (gDNA) was harvested from infected HeLa cells at the indicated times post-infection under tryptophan-replete (black) and -deplete (red) conditions. Chlamydial genome copy number was quantified by qPCR. Chlamydial genome replication is stalled following 12 hours of tryptophan limitation, but not 6. N=2. (*C*) Total RNA was harvested from infected HeLa cells at the indicated times post-infection under tryptophan-replete (black) and -deplete (red) conditions. The transcript abundance of hallmark persistence genes *euo* and (*D*) *omcB* were quantified by RT-qPCR and normalized against genome copy number. No period of tryptophan limitation significantly impacted *euo* or *omcB* expression. N=2. Statistical significance was determined by One-Way ANOVA followed by post-hoc pairwise *t*-tests with Bonferroni's correction for multiple comparisons.

Figure 4 – Figure Supplement 1. 5'-RACE conditions produce *Chlamydia*-specific products that are amplified in primary RACE. (*A*) Total RNA harvested from mock-infected HeLa cells was processed for 5'-RACE in parallel with infected samples to determine specificity of amplified products to *Chlamydia*-infected cells. No RACE products were detected in the mock-infected sample. (*B*) Primary products amplified from 5'-RACE were electrophoresed on an agarose gel and visualized. Weak bands corresponding to those detected by nested 5'-RACE were observed, with the noted relative abundance of the 1.0 and 1.5 kb products in the Bpdl-treated and Trp-deplete conditions, respectively.

1692 **Figure 4 – Figure Supplement 2.** Mapping of 5'-RACE products at the individual
1693 nucleotide level. (*A*) Mapping of the 1.5 kb 5'-RACE product. (*B*) Mapping of the 1.1 kb
1694 5'-RACE product. (*C*) Mapping of the 1.0 kb 5'-RACE product.

Figure 5 – Figure Supplement 1. Truncated fragments of the *trpRBA* IGR are not sufficient to confer YtgR repression phenotype, regardless of the presence of the putative operator site. The cartoon depiction of the *trpRBA* operon indicates the position of the IGR fragments cloned into each of the reporter vectors (*i.e.* pCCT-IGR1 through IGR3). No statistically meaningful differences were detected in the β-galactosidase activity in the presence of any of the IGR fragments.

Figure 6 – Figure Supplement 1. RNA-Sequencing coverage map of reads mapping to the intergenic regions upstream of *euo* and *lpdA*. (*A*) Coverage map for the IGR upstream of *euo*, a gene that is neither iron-regulated nor YtgR-regulated. (*B*) Coverage map for the IGR upstream of *lpdA*, a gene that is iron-regulated but not YtgR-regulated. Neither coverage map indicates an increase in reads mapping to these intergenic regions, suggesting that the effect observed at the *trpRBA* IGR is specific.

Figure 6 – Figure Supplement 2. 3'-RACE conditions produce *Chlamydia*-specific
products that are amplified in primary RACE. (*A*) Contrast-adjusted image to show weak
amplification of ~3.0 kb full-length *trpRBA* transcript in nested 3'-RACE reaction. (*B*)
Total RNA harvested from mock-infected HeLa cells was processed for 3'-RACE in
parallel with infected samples to determine specificity of amplified products to
Chlamydia-infected cells. No RACE products were detected in the mock-infected
sample. (*C*) Primary products amplified from 3'-RACE were electrophoresed on an
agarose gel and visualized. Weak bands that were relatively non-specific were detected
in the 3'-RACE primary amplification, emphasizing the utility of producing enhanced
specificity in the nested amplification.

Figure 6 – Figure Supplement 3. Mapping of the 3'-RACE products at the individual nucleotide level. (*A*) Mapping of the 0.20 kb 3'-RACE product. (*B*) Mapping of the 0.40 kb 3'-RACE product. (*C*) Mapping of the 0.45 kb 3'-RACE product. (*D*) Mapping of the 0.55 kb 3'-RACE product.