The tryptophan salvage pathway is dynamically regulated by the iron-dependent repressor YtgR in *Chlamydia trachomatis*

Nick D. Pokorzynski[1] and Rey A. Carabeo[1]

[1]Center for Reproductive Biology, School of Molecular Biosciences, College of Veterinary Medicine, Washington State University, Pullman, WA, USA, 99164.

Corresponding Author:

Rey A. Carabeo

1770 NE Stadium Way, Biotechnology & Life Sciences Bldg., Washington State University

Pullman, WA, 99164

(509) 335-7788

rey.carabeo@wsu.edu

Keywords: stress response, transcription, tryptophan biosynthesis, iron-dependent regulation, repression

Abstract

Mammalian hosts restrict cellular nutrient availability to starve invading pathogens and successfully clear an infection by a process termed "nutritional immunity". For the obligate intracellular pathogen *Chlamydia trachomatis*, nutritional immunity likely encompasses the simultaneous limitation of the amino acid tryptophan and the essential biometal iron. Unlike other model bacteria, *C. trachomatis* lacks many global stress response systems – such as "stringent response" homologs – adapted to these host insults. However, a physiological response by *Chlamydia* that is common to both stresses is the development of an aberrant, "persistent" state, suggesting that tryptophan and iron starvation trigger a coordinated developmental program. Here, we report that the *trpRBA* operon for tryptophan salvage in *C. trachomatis* serovar L2 is regulated at the transcriptional level by iron. The expression of the tryptophan synthase encoding genes, *trpBA*, is induced following iron limitation while that of the repressor *trpR* is not. We show that this specific induction of *trpBA* expression initiates from a novel promoter element within an intergenic region flanked by *trpR* and *trpB*. YtgR, a DtxR-homolog and the only known iron-dependent transcriptional regulator in *Chlamydia*, can bind to the *trpRBA* intergenic region upstream of the alternative *trpBA* promoter to repress transcription. This binding also likely attenuates transcription from the primary promoter upstream of *trpR* by blocking RNA polymerase read-through. These data illustrate a dynamic and integrated method of regulating tryptophan biosynthesis in an iron-dependent manner, which has not been described in any other prokaryote, underscoring the uniqueness of *Chlamydia*.

Significance Statement

Genital serovars of the obligate intracellular parasite *Chlamydia trachomatis* are the leading cause of bacterial sexually-transmitted infections globally. Proliferation of *C. trachomatis* is likely controlled by simultaneous immunological and environmental restriction of critical nutrients such as tryptophan and iron. However, our understanding of the immediate chlamydial responses to these stimuli is poorly defined. We utilized expression of the stress-responsive *trpRBA* operon in *C. trachomatis* L2 as a proxy for regulatory integration between iron and tryptophan limitation. We identified a unique iron-dependent regulatory mechanism for *trpRBA* in *C. trachomatis*, mediated by the transcriptional repressor YtgR. This distinguishes *Chlamydia* from other bacteria which regulate tryptophan biosynthesis strictly by tryptophan-dependent mechanisms, highlighting a distinct evolutionary adaptation in *C. trachomatis* to integrate stress responses.

/body

Introduction

Nutrient acquisition is critical for the success of pathogenic bacteria. Many

pathogenic bacteria must siphon nutrients from their hosts, such as nucleotides, amino

acids and biometals (1–4). This common feature among pathogens renders them

susceptible to nutrient limitation strategies associated with the host immune response

(5). Counteractively, bacterial pathogens have evolved sophisticated molecular

mechanisms to counter nutrient deprivation, involving increasingly complex and

sophisticated nutrient-sensing regulatory networks. These stress response mechanisms

are essential for pathogens to avoid clearance by the immune system. By delineating

their function at the molecular level, we can better target aspects of the host-pathogen

interface suitable for therapeutic manipulation. However, stress responses in the

obligate intracellular bacterium *Chlamydia trachomatis* are relatively poorly

characterized, leaving unanswered many fundamental questions about the biology of

this pathogen.

Chlamydia trachomatis is the leading cause of bacterial sexually transmitted

infections (STIs) and infection-derived blindness worldwide (6–8). Genital infections of

chlamydia are associated with serious sequelae in the female reproductive tract such as

tubal factor infertility (9). Chlamydiae are Gram-negative bacterial parasites that develop

within a pathogen-specified membrane-bound organelle termed the inclusion (10).

Chlamydial development is uniquely characterized by a biphasic interconversion of an

infectious elementary body (EB) with a non-infectious, but replicative reticulate body

(RB) (11). An obligate intracellular lifestyle has led to reductive genome evolution

across chlamydial species; Chlamydiae have retained genes uniquely required for their

survival, but have become nutritionally dependent on their hosts by discarding many

metabolism-related genes (12). Of note, *C. trachomatis* does not possess genes

necessary for eliciting a stringent response to nutrient starvation (*e.g. relA, spoT*),

suggesting that this pathogen may utilize novel mechanisms to respond to stress (13).

It is well established, however, that in response to various stressors, Chlamydiae

deviate from their normal developmental program to initiate an aberrant developmental

state, termed "persistence" (14). This persistent state is distinguished by the presence

of viable, but non-cultivable, abnormally enlarged chlamydial organisms that display

dysregulated gene expression. Importantly, *Chlamydia* can be reactivated from

persistence by abatement of the stress condition. As such, chlamydial persistence at

least superficially resembles a global stress response mechanism. Yet the molecular

underpinnings of this phenotype are poorly understood, with most published studies

focusing on the molecular and metabolic character of the aberrant, persistent form. It is

therefore unclear to what extent primary stress responses contribute to the global

persistent phenotype in *Chlamydia*.

The best described inducer of persistence is the pro-inflammatory cytokine

interferon-gamma (IFN-γ). The bacteriostatic effect of IFN-γ has been primarily

attributed to host cell tryptophan (Trp) catabolism, for which *C. trachomatis* is

auxotrophic (15–17). Following IFN-γ stimulation, infected host cells up-regulate

expression of indoleamine-2,3-dioxygenase (IDO1), which catabolizes Trp to *N*-

formylkynurenine via cleavage of the indole ring (18). *C. trachomatis* cannot recycle

kynurenines, unlike some other chlamydial species (19)*,* and thus IFN-γ stimulation

effectively results in Trp starvation to *C. trachomatis*. The primary response to Trp

starvation in *C. trachomatis* is mediated by a TrpR ortholog, whose Trp-dependent

binding to cognate promoter elements represses transcription (20, 21). This

mechanism of regulatory control is presumably limited in *C. trachomatis*, as homologs of

genes regulated by TrpR in other bacteria (*e.g. trpF, aroH, aroL*) have not been shown

to respond to Trp limitation (22).

In many Gram-negative bacteria, such as *Escherichia coli*, *trpR* is monocistronic

and distal to the Trp biosynthetic operon. In *C. trachomatis,* TrpR is encoded in an

operon, *trpRBA*, which also contains the Trp synthase α- and β- subunits (TrpA and

TrpB, respectively), and possesses a 351 base-pair (bp) intergenic region (IGR) that

separates *trpR* from *trpBA*. The functional significance of the *trpRBA* IGR is poorly

characterized; while a putative operator sequence was identified overlapping an

alternative transcriptional origin for *trpBA* (20), the IGR was not shown to be bound by

TrpR (21). Based on *in silico* predictions, an attenuator sequence has been annotated

within the *trpRBA* IGR (23), but this has not been thoroughly validated experimentally.

Regardless, the IGR is more than 99% conserved at the nucleotide sequence level

across ocular, genital and lymphogranuloma venereum (LGV) serovars of *C.*

trachomatis, indicating functional importance (Fig. S1). Therefore, relative to other

model bacteria, the regulation of the *trpRBA* operon remains poorly elucidated.

In evaluating alternative regulatory modes of the *trpRBA* operon, an interesting

consideration is the pleiotropic effects induced by IFN-γ stimulation of infected cells.

IFN-γ is involved in many processes that limit iron and other essential biometals to

intracellular pathogens as a component of host nutritional immunity (5, 24). *Chlamydia*

have a strict iron dependence for normal development, evidenced by the onset of

persistence following prolonged iron limitation (25). Importantly, *Chlamydia* presumably

acquire iron via vesicular interactions between the chlamydial inclusion and slow-

recycling transferrin (Tf)-containing endosomes (26). IFN-γ is known to down-regulate

transferrin receptor (TfR) expression in both monocytes and epithelial cells with

replicative consequences for resident intracellular bacteria (27–30). However, iron

homeostasis in *Chlamydia* is poorly understood, due to the lack of functionally

characterized homologs to iron acquisition machinery that are highly conserved in other

bacteria (31). Only one operon, represented by *ytgABCD*, has been clearly linked to iron

acquisition. The periplasmic metal-binding protein YtgA displays a specific binding

affinity for ferric iron over other divalent cations and likely transports iron from the outer

membrane to an ABC-3 type inner membrane metal permease formed by the YtgBCD

complex (32). Intriguingly, the YtgC (CTL0323) open reading frame (ORF) encodes a N-

terminal permease domain fused to a C-terminal DtxR-like repressor domain, annotated

YtgR (33, 34). YtgR is cleaved from the permease domain during infection and functions

as an iron-dependent transcriptional repressor to autoregulate the expression of its own

operon (34). YtgR represents the only identified iron-dependent transcriptional regulator

in *Chlamydia.* Whether YtgR maintains a more diverse transcriptional regulon beyond

the *ytgABCD* operon has not yet been addressed and remains an intriguing question in

the context of immune-mediated iron limitation to *C. trachomatis*. Crucially,

simultaneous IFN-γ-mediated iron and Trp starvation raise the possibility that *C.*

trachomatis could have evolved an integrated response to multiple stresses.

Consistent with the highly reduced capacity of the chlamydial genome, it is likely

that *C. trachomatis* has a limited ability to tailor a specific response to each individual

stress. In the absence of identifiable homologs for most global stress response

regulators in *C. trachomatis*, we hypothesized that primary stress responses to

pleiotropic insults may involve mechanisms of regulatory integration, whereby important

molecular pathways are co-regulated by stress-responsive transcription factors such

that they can be utilized across multiple stress conditions simultaneously induced by the

host. Here, we report on the unique iron-dependent regulation of the *trpRBA* operon in

Chlamydia trachomatis. We propose a model of iron-dependent transcriptional

regulation of *trpRBA* mediated by the repressor YtgR binding specifically to the IGR,

which may have implications for how *C. trachomatis* responds to immunological and

environmental insults. Such a mechanism of iron-dependent regulation of Trp

biosynthesis has not been previously described in any other prokaryote and adds to the

catalog of regulatory models for Trp biosynthetic operons in bacteria. Further, it reveals

a highly dynamic mode of regulatory integration within the *trpRBA* operon, exemplifying

the importance of this pathway to chlamydial stress response.

Results

Brief iron limitation via 2,2-bipyridyl treatment yields iron-starved, but non-

persistent *Chlamydia trachomatis.* To identify possible instances of regulatory

integration between iron and Trp starvation in *C. trachomatis*, we optimized a stress

response condition that preceded the development of a characteristically persistent

phenotype. We reasoned that in order to effectively identify regulatory integration, we

would need to investigate the bacterium under stressed, but not aberrant, growth

conditions such that we could distinguish primary stress responses from abnormal

growth. To specifically investigate the possible contribution of iron limitation to a broader

immunological (*e.g.* IFN-γ-mediated) stress, we utilized the membrane-permeable iron

chelator 2,2-bipyridyl (Bpdl), which has the advantage of rapidly and homogeneously

starving *C. trachomatis* of iron (35). We chose to starve *C. trachomatis* serovar L2 of

iron starting at 12 hrs post-infection (hpi), or roughly at the beginning of mid-cycle

growth. At this point the chlamydial organisms represent a uniform population of

replicative RBs that are fully competent, both transcriptionally and translationally, to

respond to stress. We treated infected HeLa cell cultures with 100 μM Bpdl or mock for

either 6 or 12 hours (hrs) to determine a condition sufficient to limit iron to *C.*

trachomatis without inducing hallmark persistent phenotypes. We stained infected cells

seeded on glass coverslips with convalescent human sera and analyzed chlamydial

inclusion morphology under both Bpdl- and mock-treated conditions by laser point-

scanning confocal microscopy (Fig. 1A). Following 6 hrs of Bpdl treatment, chlamydial

inclusions were largely indistinguishable from mock-treated inclusions, containing a

homogeneous population of larger organisms, consistent with RBs in mid-cycle growth.

However, by 12 hrs of Bpdl treatment, the inclusions began to display signs of aberrant

growth: they were perceptibly smaller, more comparable in size to 18 hpi, and contained

noticeably fewer organisms, perhaps indicating a delay in RB-to-EB differentiation.

These observations were consistent with our subsequent analysis of genome replication

by quantitative PCR (qPCR; Fig. 1B.) At 6 hrs of Bpdl treatment, there was no

statistically distinguishable difference in genome copy number when compared to the

equivalent mock-treated time-point. However, by 12 hrs of treatment, genome copy

number was significantly reduced 4.7-fold in the Bpdl-treated group relative to mock-

treatment ($p = 0.0033$). We then assayed the transcript expression of two markers for

persistence by reverse transcription quantitative PCR (RT-qPCR): the early gene *euo*,

encoding a transcriptional repressor of late-cycle genes (Fig.1C), and the adhesin

omcB, which is expressed late in the developmental cycle (Fig. 1D). Characteristic

persistence would feature sustained high *euo* expression late into infection, and

suppressed *omcB* expression throughout development. We observed that at 6 hrs of

Bpdl treatment, there was no statistically distinguishable difference in either *euo* or

omcB expression when compared to the mock-treatment. Still at 12 hrs of Bpdl

treatment, *euo* expression was unchanged. However, *omcB* expression was

significantly induced following 12 hrs of Bpdl-treatment ($p = 0.00015$). This was

unexpected, but we note that *omcB* expression has been shown to vary between

chlamydial serovars and species when starved for iron (31). Collectively, these data

indicated that 6 hrs of Bpdl treatment was a more suitable time-point at which to monitor

iron-limited stress responses.

We additionally assayed these same metrics following 6 or 12 hrs of Trp

starvation by culturing cells in either Trp-replete or Trp-deplete DMEM-F12 media

supplemented with fetal bovine serum (FBS) pre-dialyzed to remove amino acids. We

observed no discernable change in inclusion morphology out to 12 hrs of Trp starvation

(Fig. S2A), but genome copy numbers were significantly reduced 2.7-fold at this time-

point ($p = 0.00612$; Fig. S2B). The transcript expression of *euo* (Fig. S2C) and *omcB*

(Fig. S2D) did not significantly change at either treatment duration, but Trp-depletion did

result in a 2.0-fold reduction in *omcB* expression, consistent with a more characteristic

persistent phenotype. These data therefore also indicated that 6 hrs of treatment would

be ideal to monitor non-persistent responses to Trp limitation.

We next sought to determine whether our brief 6-hr BpdI treatment was sufficient

to elicit a transcriptional iron starvation phenotype. We chose to analyze the expression

of three previously identified iron-regulated transcripts, *ytgA* (Fig. 2A), *ahpC* (Fig. 2B)

and *devB* (Fig. 2C), by RT-qPCR under BpdI- and mock-treated conditions (35, 36). In

addition, we analyzed the expression of one non-iron-regulated transcript, *dnaB* (Fig.

2D), as a negative control (37). Following 6 hrs of BpdI treatment, we observed that the

transcript expression of the periplasmic iron-binding protein *ytgA* was significantly

elevated 1.75-fold relative to the equivalent mock-treated time-point (p = 0.0052).

However, we did not observe induction of *ytgA* transcript expression relative to the 12

hpi time-point. We distinguish here between *elevated* and *induced* transcript expression,

as chlamydial gene expression is highly developmentally regulated. Thus, it can be

more informative to monitor longitudinal expression of genes, *i.e.* their induction, as

opposed to elevation relative to an equivalent control time-point, which may simply

represent a stall in development. While we did not observe induction of *ytgA*, which

would be more consistent with an iron-starved phenotype, we reason that this is a

consequence of the brief treatment period, and that longer iron starvation would

produce a more robust induction of iron-regulated transcripts. Note that the identification

of *ytgA* as iron-regulated has only been previously observed following extended periods

of iron chelation (32, 35, 38). Similarly, we observed that the transcript expression of the

thioredoxin *ahpC* was significantly elevated 2.15-fold relative to the equivalent mock-

treated time-point (p = 0.038) but was not induced relative to the 12 hpi time-point. The

transcript expression of *devB*, encoding a 6-phosphogluconolactonase involved in the

pentose phosphate pathway, was not observed to significantly respond to our brief iron

limitation condition, suggesting that it is not a component of the primary iron starvation

stress response in *C. trachomatis.* As expected, the transcript expression of *dnaB*, a

replicative DNA helicase, was not altered by our iron starvation condition, consistent

with its presumably iron-independent regulation (37). Overall, these data confirmed that

our 6-hr BpdI treatment condition was suitable to produce a mild iron starvation

phenotype at the transcriptional level, thus facilitating our investigation of iron-

dependent regulatory integration.

Transcript expression of the *trpRBA* operon is differentially regulated by iron in

Chlamydia trachomatis. Upon identifying an iron limitation condition that produced a

relevant transcriptional phenotype while avoiding the onset of persistent development,

we aimed to investigate whether the immediate response to iron starvation in *C.*

trachomatis would result in the consistent induction of pathways unrelated to iron

utilization/acquisition, but nevertheless important for surviving immunological stress.

The truncated Trp biosynthetic operon, *trpRBA* (Fig. 3A), has been repeatedly linked to

the ability of genital and LGV serovars (D-K and L1-3, respectively) of *C. trachomatis* to

counter IFN-γ-mediated stress. This is due to the capacity of the chlamydial Trp

synthase in these serovars to catalyze the β synthase reaction, *i.e.* the condensation of

indole to the amino acid serine to form Trp (17). In the presence of exogenous indole,

C. trachomatis is therefore able to biosynthesize Trp such that it can prevent the

development of IFN-γ-mediated persistence. Correspondingly, the expression of *trpRBA*

is highly induced following IFN-γ stimulation of infected cells (39, 40). These data have

historically implicated Trp starvation as the primary mechanism by which persistence

develops in *C. trachomatis* following exposure to IFN-γ. However, these studies have

routinely depended on prolonged treatment conditions that monitor the terminal effect of

persistent development, as opposed to the immediate molecular events which may

have important roles in the developmental fate of *Chlamydia.* As such, these studies

may have missed the contribution of other IFN-γ-stimulated insults such as iron

limitation.

To decouple Trp limitation from iron limitation and assess their relative

contribution to regulating a critical pathway for responding to IFN-γ-mediated stress, we

monitored the transcript expression of the *trpRBA* operon under brief Trp or iron

starvation by RT-qPCR. When starved for Trp for 6 hrs, we observed that the

expression of *trpR*, *trpB* and *trpA* were all significantly induced greater than 10.5-fold

relative to 12 hpi (p = 0.00077, 0.025 and 9.7e-5, respectively; Fig. 3B). All three ORFs

were also significantly elevated relative to the equivalent mock-treated time-point (p =

0.00076, 0.025 and 9.7e-5, respectively). This result was surprising with respect to the

relative immediacy of operon induction in response to our Trp starvation protocol,

confirming the relevant Trp-starved transcriptional phenotype. To induce Trp-deprived

persistence in *C. trachomatis*, many laboratories rely on compounded techniques of

IFN-γ pre-treatment to deplete host Trp pools in conjunction with culturing in Trp-

depleted media, among other strategies. While the phenotypic end-point differs here, it

is nonetheless interesting to note that only 6 hrs of media replacement is sufficient to

markedly up-regulate *trpRBA* expression. This suggests that *C. trachomatis* has a

highly attuned sensitivity to even moderate changes in Trp levels.

We next performed the same RT-qPCR analysis on the expression of the *trpRBA*

operon in response to 6 hrs of iron limitation via Bpdl treatment (Fig. 3C). While we

observed that the transcript expression of all three ORFs was significantly elevated at

least 2.1-fold relative to the equivalent mock-treated time-point ($p = 0.015$, 0.00098 and

0.0062, respectively), we made the intriguing observation that only the expression *trpB*

and *trpA* was significantly induced relative to 12 hpi ($p = 0.00383$ and 0.0195,

respectively). The significant induction of *trpBA* expression, but not *trpR* expression,

suggested that *trpBA* are specifically regulated by iron availability. This result is

consistent with a recent survey of the iron-regulated transcriptome in *C. trachomatis* by

RNA sequencing, which also reported that iron-starved *Chlamydia* specifically up-

regulate *trpBA* expression in the absence of altered *trpR* expression (37). Our results

expand on this finding by providing a more detailed investigation into the specific profile

of this differential regulation of *trpRBA* in response to iron deprivation. Taken together,

these findings demonstrated that an important stress response pathway, the *trpRBA*

operon, is regulated by the availability of both Trp and iron, consistent with the notion

that the pathway may be cooperatively regulated to respond to various stress

conditions. Notably, iron-dependent regulation of Trp biosynthesis has not been

previously documented in other prokaryotes.

Specific iron-regulated expression of *trpBA* originates from a novel alternative

transcriptional start site within the *trpRBA* intergenic region. We hypothesized that

the specific iron-related induction of *trpBA* expression relative to *trpR* expression may

be attributable to an iron-regulated alternative transcriptional start site (alt. TSS)

downstream of the *trpR* ORF. Indeed, a previous study reported the presence of an alt.

TSS in the *trpRBA* IGR, located 214 nucleotides upstream of the *trpB* translation start

position (20). However, a parallel study could not identify a TrpR binding site in the

trpRBA IGR (21). We reasoned that a similar alt. TSS may exist in the IGR that

controlled the iron-dependent expression of *trpBA.* We therefore performed Rapid

Amplification of 5'-cDNA Ends (5'-RACE) on RNA isolated from *C. trachomatis* L2-

infected HeLa cells using the SMARTer 5'/3' RACE Kit workflow (Takara Bio). Given the

low expression of the *trpRBA* operon during normal development, we utilized two

sequential gene-specific amplification steps (nested 5'-RACE) to identify 5' cDNA ends

in the *trpRBA* operon. These nested RACE conditions resulted in amplification that was

specific to infected-cells (Fig. S3A). Using this approach, we analyzed four conditions:

12 hpi, 18 hpi, 12 hpi + 6 hrs of BpdI treatment, and 12 hpi + 6 hrs of Trp-depletion (Fig.

4A). We observed three RACE products that migrated with an apparent size of 1.5, 1.1

and 1.0 kilobases (kb). At 12 and 18 hpi, all three RACE products exhibited low

abundance, even following the nested PCR amplification. This observation was

consistent with the expectation that the expression of the *trpRBA* operon is very low

under normal, iron and Trp-replete conditions. However, we note that the 6-hr difference

in development did appear to alter the representation of the 5' cDNA ends, which may

suggest a stage-specific promoter utilization within the *trpRBA* operon. In our Trp

starvation condition, we observed an apparent increase in the abundance of the 1.5 kb

RACE product, which was therefore presumed to represent the primary TSS upstream

of *trpR*, at nucleotide position 511,389 (*C. trachomatis* L2 434/Bu). Interestingly, the 1.0

kb product displayed a very similar apparent enrichment following BpdI treatment,

suggesting that this RACE product represented a specifically iron-regulated TSS. Both

the 1.5 and 1.0 kb RACE products were detectable in the Trp-depleted and iron-

depleted conditions, respectively, during the primary RACE amplification, consistent

with their induction under these conditions (Fig. S3B).

If iron depletion was inducing *trpBA* expression independent of *trpR,* we

reasoned that we would observe specific enrichment of *trpB* sequences in our 5'-RACE

cDNA samples relative to *trpR* sequences. We again utilized RT-qPCR to quantify the

abundance of *trpB* transcripts relative to *trpR* transcripts in the 5'-RACE cDNA samples

(Fig. 4B). In agreement with our model, only under iron starved conditions did we

observe a significant enrichment of *trpB* relative to *trpR* ($p < 0.01$). Additionally, we

observed that at 12 and 18 hpi in iron-replete conditions, the ratio of *trpB* to *trpR* was

approximately 1.0, suggesting non-preferential basal expression across the three

putative TSSs. Another factor contributing to this ratio is the synthesis of the full-length

trpRBA polycistron. In support of this, the *trpB* to *trpR* ratio remained near 1.0 under the

Trp-starved condition, which would be expected during transcription read-through of the

whole operon. The apparent lack of preferential promoter utilization as described above

could be attributed to the relatively low basal expression of the operon at 12 and 18 hpi

under Trp- and iron-replete conditions, thus precluding quantitative detection of

differential promoter utilization in this assay.

To determine the specific location of the 5' cDNA ends within the *trpRBA* operon,

we isolated the 5'-RACE products across all conditions by gel extraction and cloned the

products into the pRACE vector supplied by the manufacturer. We then sequenced the

ligated inserts and BLASTed the sequences against the *C. trachomatis* L2 434/Bu

genome to identify the location of the 5'-most nucleotides (Fig. 4C). These data are

displayed as a statistical approximation of the genomic regions most likely to be

represented by the respective 5'-RACE products in both histogram (semi-continuous)

and density plot (continuous) format (See Dataset S1 for a description of all mapped 5'-

RACE products). As expected, the 1.5 kb product mapped in a distinct and tightly

grouped peak near the previously annotated *trpR* TSS, with the mean and modal

nucleotide being 511,388 and 511,389, respectively (Fig. S4A). Surprisingly, we found

that neither the 1.1 or 1.0 kb RACE product mapped to the previously reported alt. TSS

in the *trpRBA* IGR, at position 511,826. Instead, we observed that the 1.1 kb product

mapped on average to nucleotide position 511,878, with the modal nucleotide being

found at 511,898 (Fig. S4B). The 1.0 kb product mapped with a mean nucleotide

position of 512,013, with the modal nucleotide being 512,005 (Fig. S4C), only 35 bases

upstream of the *trpB* coding sequence. Interestingly, the 1.0 kb product mapped to a

region of the *trpRBA* IGR flanked by consensus σ^{66} -10 and -35 promoter elements,

found at positions 512,020-5 and 511,992-7, respectively (41). These data collectively

pointed toward the 1.0 kb 5'-RACE product representing a novel, iron-regulated alt. TSS

and *bona fide* σ^{66}-dependent promoter element that allows for the specific iron-

dependent expression of *trpBA*.

YtgR specifically binds to the *trpRBA* intergenic region in an operator-dependent

manner to repress transcription of *trpBA*. As the only known iron-dependent

transcriptional regulator in *Chlamydia*, we hypothesized that YtgR may regulate the iron-

dependent expression of *trpBA* from the putative promoter element we characterized by

5'-RACE. Using bioinformatic sequence analysis, we investigated whether the *trpRBA*

IGR contained a candidate YtgR operator sequence. By local sequence alignment of

the putative YtgR operator sequence (33) and the *trpRBA* IGR, we identified a high-

identity alignment (76.9% identity) covering 67% of the putative operator sequence (Fig.

5A). Interestingly, this alignment mapped to the previously identified palindrome

suspected to have operator functionality (20). By global sequence alignment of the YtgR

operator to the palindromic sequence, an alignment identical to the local alignment was

observed, which still displayed relatively high sequence identity (43.5% identity). We

hypothesized that this sequence functioned as an YtgR operator, despite being located

184 bp upstream of the *trpBA* alt. TSS.

To investigate the ability of YtgR to bind and repress transcription from the

putative *trpBA* promoter, we implemented a heterologous two-plasmid assay that

reports on YtgR repressor activity as a function of β-galactosidase expression (14). In

brief, a candidate DNA promoter element was cloned into the pCCT expression vector

between an arabinose-inducible pBAD promoter and the reporter gene *lacZ*. This

plasmid was co-transformed into BL21 (DE3) *E. coli* along with an IPTG-inducible

pET151 expression vector with (pET151-YtgR) or without (pET151-EV) the C-terminal

139 amino acid residues of CTL0323 (YtgC). Note that we have previously

demonstrated that this region is a functional iron-dependent repressor domain (34). To

verify the functionality of this assay, we determined whether ectopic YtgR expression

could repress pCCT reporter gene expression in the presence of three candidate DNA

elements: a no-insert empty vector (pCCT-EV), the putative promoter element for *C.*

trachomatis lpdA (pCCT-*lpdA*), and the promoter region of the *ytgABCD* operon (pCCT-

ytgABCD; Fig. 5B). As expected, from the pCCT-EV reporter construct, ectopic YtgR

expression did not significantly reduce the activity of β-galactosidase. Additionally,

reporter gene expression from pCCT-*lpdA*, containing the promoter of iron-regulated

lpdA (37), which is not known to be YtgR-regulated, was not affected by ectopic

expression of YtgR. This demonstrated that the assay can discriminate between the

promoter elements of iron-regulated genes and *bona fide* YtgR targeted promoters.

Indeed, in the presence of pCCT-*ytgABCD*, induction of YtgR expression produced a

significant decrease in β-galactosidase activity (p = 0.03868) consistent with its

previously reported auto-regulation of this promoter (34).

Using this same assay, we then inserted into the pCCT reporter plasmid 1) the

trpR promoter element (pCCT-*trpR*), 2) the putative *trpBA* promoter element

represented by the IGR (pCCT-*trpBA*), and 3) the same putative *trpBA* promoter

element with a mutated YtgR operator sequence that was diminished for both

palindromicity and A-T richness, two typical features of prokaryotic promoter elements

(pCCT-*trpBA*ΔOperator; Fig. 5C) (42, 43). When YtgR was ectopically expressed in the

pCCT-*trpR* background, we observed no statistically distinguishable change in β-

galactosidase activity. However, in the pCCT-*trpBA* background, ectopic YtgR

expression significantly reduced β-galactosidase activity at levels similar to those

observed in the pCCT-*ytgABCD* background (p = 0.01219). This suggested that YtgR

was capable of binding to the *trpBA* promoter element specifically. Interestingly, this

repression phenotype was abrogated in the pCCT-*trpBA*ΔOperator background, where

we observed no statistically meaningful difference in β-galactosidase activity. We

subsequently addressed whether the region of the *trpRBA* IGR containing the YtgR

operator site was sufficient to confer YtgR repression in this assay (Fig. S5). Therefore,

we cloned three fragments of the *trpRBA* IGR into the pCCT reporter plasmid: the first

fragment represented the 5'-end of the IGR containing the operator site at the 3'-end

(pCCT-IGR1), the second fragment represented a central region of the IGR containing

the operator site at the 5'-end (pCCT-IGR2), and the third fragment represented the 3'-

end of the IGR and did not contain the operator site (pCCT-IGR3). Surprisingly, we

observed that none of these fragments alone were capable of producing a significant

repression phenotype in our reporter system. This finding indicated that while the

operator site was necessary for YtgR repression, it alone was not sufficient. Together,

these data indicated that YtgR could bind to the *trpBA* promoter element and that this

binding was dependent upon an intact AT-rich palindromic sequence, likely representing

an YtgR operator, but that further structural elements in the *trpRBA* IGR may be

necessary for repression. Nonetheless, we demonstrated the existence of a functional

YtgR binding site that conferred iron-dependent transcriptional regulation to *trpBA*,

independent of the major *trpR* promoter.

Iron limitation promotes transcription read-through at the *trpRBA* YtgR operator

site. If YtgR binds to the operator sequence within the intergenic region under iron-

replete conditions, does it influence RNA polymerase (RNAP) read-through from the

major *trpR* promoter to synthesize the polycistronic *trpRBA* mRNA? We hypothesized

that the presence of YtgR at the *trpRBA* operator may disadvantage the processivity of

RNAP initiating transcription at the upstream *trpR* promoter. Similar systems of RNAP

read-through blockage have been reported; the transcription factor Reb1p "roadblocks"

RNAPII transcription read-through in yeast by a mechanism of RNAP pausing and

subsequent labelling for degradation (44). To investigate this question, we first identified

transcription termination sites (TTSs) in the *trpRBA* operon in *C. trachomatis.* We

utilized 3'-RACE to map the 3'-ends of transcripts using gene-specific primers within the

trpR CDS (Fig. 6A; bottom). We again utilized two RACE amplification cycles to

generate distinct, specific bands suitable for isolation and sequencing (Fig. S6B-C). By

gel electrophoresis of the 3'-RACE products, we observed the appearance of four

distinct bands that migrated with an apparent size of 0.55, 0.45, 0.40 and 0.20 kb. In our

Trp-depleted condition, we observed only a very weak amplification of the 2.5 – 3 kb

full-length *trpRBA* message by 3'-RACE (Fig. S6A). However, we did observe it across

all replicates. To confirm that the full-length product was relatively specific to the Trp-

deplete treatment, we amplified the *trpRBA* operon by RT-PCR from the 3'-RACE cDNA

(Fig. 6A; top). As expected, only in the Trp-deplete sample did we observe robust

amplification of the full-length *trpRBA* message. We note however that image contrast

adjustment reveals a very weak band present in all experimental samples.

To identify the specific TTS locations, we gel extracted the four distinct 3'-RACE

bands across all conditions and cloned them into the pRACE sequencing vector as was

done for the 5'-RACE experiments. We then sequenced the inserted RACE products

and mapped them to the *C. trachomatis* L2 434/Bu genome (Fig. 6B). This revealed a

highly dynamic TTS landscape contained almost exclusively within the *trpRBA* IGR,

which has not previously been investigated (For a full description of mapped 3'-RACE

products, see Dataset S2). The 0.20 kb RACE product mapped tightly to the 3'-end of

the *trpR* CDS, with a mean nucleotide position of 511,665 and a modal nucleotide

position of 511,667 (Fig. S7A). Contrastingly, the other three RACE products did not

map in such a way so as to produce specific, unambiguous modal peaks. Instead, their

distribution was broader and more even, with only a few nucleotide positions mapping

more than once. Accordingly, the 0.45 kb product mapped with an average nucleotide

position of 511,889, just downstream of the 1.1 kb 5'-RACE product (Fig. S7C), while

the 0.55 kb product mapped with an average nucleotide position of 511,986, upstream

of the 1.0 kb 5'-RACE product (Fig. S7D). Interestingly, the 0.40 kb product mapped to

a region directly overlapping the putative YtgR operator site, with a mean nucleotide

position of 511,811 (Fig. S7B). We therefore reasoned that this putative TTS may have

an iron-dependent function.

We next aimed to quantitatively analyze the possibility that iron-depletion, and

thus dissociation of YtgR from this region, may facilitate transcription read-through at

the operator site. Working from the 3'-RACE generated cDNAs, we utilized RT-qPCR to

monitor the abundance of various amplicons across the *trpRBA* operon in relation to a

"read-through" normalization amplicon that should only be represented when the full-

length *trpRBA* message is transcribed (Fig. 6C). Therefore, as each amplicon is

increasingly represented as a portion of the full-length, read-through transcript, the

representation ratio of the specific amplicon to the normalization amplicon should

approach 1.0. We first analyzed an amplicon from nucleotide 511,416 – 531 to monitor

transcript species associated with transcription initiating at the *trpR* promoter. We

observed that the representation of this amplicon was not significantly altered following

iron limitation relative to 12 hpi, suggesting that the depletion of iron was not affecting

initiation of transcription at the *trpR* promoter. Interestingly, at 18 hpi, the representation

ratio of this amplicon significantly shifted further away from 1.0 (p = 0.00358), indicating

that at 18 hpi this amplicon is represented less as a component of read-through

transcription relative to 12 hpi. As expected, under Trp-deplete conditions, the

representation ratio shifted significantly closer to 1.0 ($p = 0.00064$), consistent with read-

through transcription of the full-length *trpRBA* message.

We then preformed the same analysis on an amplicon from nucleotide 511,639 –

764, immediately upstream of the TTS at the YtgR operator site. We again observed

that at 18 hpi, the representation ratio was significantly increased ($p = 0.01046$), and

following Trp-depletion, the ratio was significantly decreased ($p = 0.00023$), as

expected. Notably, and consistent with our hypothesis, we observed that the

representation ratio of this amplicon was also significantly closer to 1.0 following iron

limitation ($p = 0.00407$), suggesting that transcription read-through was increased at this

site under iron limited conditions. If YtgR is dissociating from the operator site during

iron depletion, a greater proportion of transcripts would be expected to read-through this

locus.

Finally, we analyzed an amplicon from nucleotide 513,856 – 968, at the very 3'-

end of *trpA* to monitor terminal transcription under our experimental conditions. At 18

hpi, we observed a significant increase in the representation ratio of this amplicon ($p =$

0.00476), which is likely attributable to both basal levels of alternative transcription from

the IGR as well as poor transcription read-through of the full-length message. Following

6 hrs of BpdI treatment, we also observed a significant increase in the representation

ratio of this amplicon ($p = 0.01510$), which supports the finding that *trpBA* is being

preferentially transcribed under this condition, distinct from the full-length *trpRBA*

transcript. We were only able to detect a marginal decrease in the representation of this

amplicon under Trp-depleted conditions ($p = 0.07942$), which may suggest that the very

3'-end of *trpRBA* is relatively under-represented than our normalization amplicon, which

falls within the middle of the operon. In fact, recent work has reported on the relatively

poor representation of 3'-end mRNAs in *Chlamydia* (45). In sum, this set of experiments

provides evidence that iron-depletion specifically alters the representation of particular

mRNA species across the *trpRBA* operon. Additionally, they implicate iron-dependent

YtgR DNA-binding as the mediator of these effects. By alleviating YtgR repression via

iron depletion, transcription is allowed to proceed through the operator site, albeit at

basal levels. Concomitantly, transcription is specifically activated at the downstream alt.

TSS for *trpBA*.

Discussion

In this study, we report a mechanism of stress adaptation that integrates

responses to iron and Trp starvation. Specifically, we demonstrated that the *trpRBA*

operon is transcriptionally regulated by the iron-dependent repressor YtgR. We

determined that iron-dependent expression of *trpBA* initiates from a novel internal

promoter in an IGR of the *trpRBA* operon and that this IGR also contains an YtgR

operator site necessary to confer a transcriptional repression phenotype. We suggest

that the dual promoter configuration of *trpRBA* presents the opportunity for YtgR to

block transcription read-through from the *trpR* promoter; transcripts terminate at the

YtgR operator site and iron depletion facilitates read-through of the operon at this locus.

This is the first time an iron-dependent mode of regulation has been shown to control

the expression of tryptophan biosynthesis in prokaryotes, which is a reflection of the

uniquely specialized nature of *C. trachomatis*.

The distance separating the YtgR operator site from the *trpBA* alt. TSS indicate

the involvement of a regulatory mechanism more complex than simple steric hindrance

of RNAP by YtgR. One possible explanation is that YtgR functions similarly to other

prokaryotic transcription factors that repress "at a distance" by a mechanism of DNA-

bending (46, 47). In this scenario, YtgR binding simultaneously to an additional operator

site may facilitate a conformational bend in the double-helix DNA such that RNAP no

longer has access to the alternative *trpBA* promoter site. This would be consistent with

the observation that a truncated *trpRBA* IGR containing the candidate YtgR operator is

insufficient to confer transcriptional repression. The topological alteration induced by

DNA-bending could also feasibly contribute to diminished RNAP read-through from the

trpR promoter. *In silico* identification of additional putative YtgR operator sites was

unsuccessful, which could be due to the lack of enough validated binding sites to

generate a robust consensus sequence. Thus, a more unbiased approach (*e.g.* ChIP-

Sequencing) will be required to identify additional YtgR binding events. We also note

that the YtgR operator upstream of *ytgA*, while only 21 bp from the predicted -35

element, rests within a 660-bp IGR, raising the possibility that other cryptic YtgR

operators are present in this sequence. Another possibility is that YtgR functions in

concert with additional transcription factors more proximal to the *trpBA* promoter

elements. In *E. coli*, the repressor Fis binds 135 bp upstream of the *nir* promoter TSS,

controlling co-activation of *nir* expression by proximally-bound Fnr and NarL/NarP (48).

While we have no evidence to suggest other transcription factors are controlling *trpBA*

expression, this does not preclude the possibility of their involvement.

While we demonstrate here that iron-dependent *trpBA* expression originates from

a novel promoter element immediately upstream of the *trpB* CDS, this is not the first

description of an alt. TSS within the *trpRBA* IGR. Carlson, et al. identified an alt. TSS

within the IGR which they suggested was responsible for *trpBA* expression (20).

Interestingly, this TSS was observed to originate from within the palindromic sequence

that we have identified here as a functional YtgR operator site. In these studies, we

were unable to confirm the presence of the previously identified alt. TSS by 5'-RACE.

This is likely because Carlson, et al. examined the presence of transcript origins

following 24 hrs of Trp starvation whereas here we monitored immediate responses to

stress following only 6 hrs of treatment. Prolonged Trp depletion would result in a more

homogeneously stressed population of chlamydial organisms that may exhibit the same

preferential utilization of the promoter identified by Carlson, et al. Population

heterogeneity in response to brief stress may explain the observation of multiple

T(S/T)Ss across the *trpRBA* operon in our studies. However, the contribution of such a

Trp-dependent alt. TSS to the general stress response of *C. trachomatis* remains

unclear. Akers & Tan were unable to verify TrpR binding to the *trpRBA* IGR by EMSA,

suggesting that some other Trp-dependent mechanism may control transcription from

this site (21). Ultimately, our approach of investigating more immediate responses to

stress revealed previously unreported mechanisms functioning to regulate Trp

biosynthesis in *C. trachomatis*, underscoring the value of transient as opposed to

sustained induction of stress.

Another mechanism of regulation reported to control the chlamydial *trpRBA*

operon is Trp-dependent transcription attenuation. Based on sequence analysis, a

leader peptide has been annotated within the *trpRBA* IGR (23). Presumably, this

functions analogously to the attenuator in the *E. coli trpEDCBA* operon; Trp starvation

causes ribosome stalling at sites of enriched Trp codons such that specific RNA

secondary structures form to facilitate RNAP read-thru of downstream sequences – in

this case, *trpBA* (49). However, robust experimental evidence to support the existence

of attenuation in *C. trachomatis* is lacking. To date, the only experimental evidence that

supports this model was reported by Carlson, et al., who demonstrated that transcript

expression of *trpBA* is increased following 24 hr Trp-depletion in a *trpR*-mutant strain of

C. trachomatis, suggesting that an additional level of Trp-dependent regulation controls

trpBA expression (20). However, this could be attributable to an alternative Trp-

dependent mechanism controlling *trpBA* expression at the alt. TSS identified by

Carlson, et al. None of the data presented here point conclusively to the existence of a

Trp-dependent attenuator; while we acknowledge that the additional termination sites

identified in our 3'-RACE assay may represent termination events mediated by an

attenuator, without more specific analysis utilizing mutated sequences we cannot draw

definitive conclusions about the functional relevance of those termination sites.

Additionally, it is unlikely that we would be able to observe Trp-dependent attenuation

under our brief stress conditions given that attenuation has a much higher tolerance for

Trp-depletion than TrpR-mediated transcriptional repression in *E. coli* (50).

Interestingly, in *Bacillus subtilis,* Trp-dependent attenuation of transcription takes

on a form markedly different from that in *E. coli*. Whereas attenuation functions in *cis* for

the *E. coli trp* operon, *B. subtilis* utilize a multimeric <u>T</u>ryptophan-activated <u>R</u>NA-binding

<u>A</u>ttenuation <u>P</u>rotein, TRAP, which functions in *trans* to bind *trp* operon RNA under Trp-

replete conditions, promoting transcription termination and inhibiting translation (51).

This interaction is antagonized by anti-TRAP in the absence of charged tRNA^{Trp}, leading

to increased expression of TRAP regulated genes. We suggest that YtgR may

represent the first instance of a separate and distinct clade of attenuation mechanisms:

iron-dependent *trans*-attenuation. We have demonstrated that transcription terminates

in the *trpRBA* operon at the YtgR operator site, and that read-thru of the operon is

facilitated under iron-deplete conditions, which is consistent with the idea that relief of

iron-activated YtgR DNA-binding at this site would permit RNAP to pass through the

YtgR operator site. This mechanism may function independently of specific RNA

secondary structure, relying instead on steric blockage of RNAP processivity, but

ultimately producing a similar result. Possible regulation of translation remains to be

explored. The recent development of new genetic tools to alter chromosomal

sequences and generate conditional knockouts in *C. trachomatis* should enable a more

detailed analysis of *trpRBA* regulation, including possible *trans*-attenuation (52, 53).

As a Trp auxotroph, what might be the biological significance of iron-dependent

YtgR regulation of the *trpRBA* operon in *C. trachomatis*? We have already noted the

possibility that iron-dependent *trpBA* regulation in *C. trachomatis* may enable a

response to simultaneous Trp and iron starvation, such as that likely mediated by IFN-γ.

However, this mechanism also presents the opportunity for *C. trachomatis* to respond

similarly to distinct *sequential* stresses, where a particular stress primes the pathogen to

better cope with subsequent stresses. To reach the female upper genital tract (UGT),

where most significant pathology is identified following infection with *C. trachomatis*, the

pathogen must first navigate the lower genital tract (LGT). *Chlamydia* infections of the

female LGT are associated with bacterial vaginosis (BV), which is characterized by

obligate and facultative anaerobe colonization, some of which catalyze the production of

indole (54, 55). This provides *C. trachomatis* with the necessary substrate to salvage

tryptophan via TrpBA. Interestingly, the LGT is also likely an iron-limited environment.

Pathogen colonization and BV both increase the concentration of mucosal lactoferrin

(Lf), an iron-binding glycoprotein, which can starve pathogens for iron (56, 57). Lf

expression is additionally hormone-regulated, and thus the LGT may normally

experience periods of iron limitation (58, 59). Moreover, the expression of TfR is

constrained to the basal cells of the LGT stratified squamous epithelium (60), which

likely restricts iron from *C. trachomatis* infecting the accessible upper layers of the

stratified epithelia. In fact, it was recently demonstrated that *C. trachomatis*

development is attenuated in the terminally differentiated layers of an *in vitro*-generated

stratified squamous epithelium (61). Collectively, LGT conditions that favor *C.*

trachomatis infection may be marked by concomitant iron limitation and indole

accessibility. For *C. trachomatis*, iron limitation may therefore serve as a critical signal in

the LGT, inducing the expression of *trpBA* such that Trp is stockpiled from available

indole, allowing the pathogen to counteract oncoming IFN-γ-mediated Trp starvation.

We propose the possibility that iron limitation in the LGT may be a significant predictor

of pathogen colonization in the UGT.

Finally, and of note, the expression of the ribonucleotide diphosphate-reductase

encoding *nrdAB* was also recently shown to be iron-regulated in *C. trachomatis* (37).

The regulation of *nrdAB* is known to be mediated by the presumably

deoxyribonucleotide-dependent transcriptional repressor NrdR, encoded distal to the

nrdAB locus (62). As NrdR activity is not known to be modulated by iron availability, this

raises the intriguing possibility that here too a unique iron-dependent mechanism of

regulation may integrate the chlamydial stress response to promote a unified response

across various stress conditions. Future studies may require more metabolomics-based

approaches to thoroughly dissect the integration of these stress responses, as

transcriptome analyses alone often miss broader, pathway-oriented metabolic

coordination. Ultimately, these studies point towards a need to carefully re-evaluate the

molecular stress response in *Chlamydia*, using more targeted approaches to answer

more specific questions. We anticipate that the rapid progress of the field in recent

years will continue to catalyze exciting and important discoveries regarding the

fundamental biology of *Chlamydia*.

Materials and Methods

Please refer to the *SI Appendix* for a complete and detailed description of all

experimental reagents and methodology. For all infections, *Chlamydia trachomatis* LGV

serological variant type II was used to infect human cervical epithelial adenocarcinoma

HeLa cells at a multiplicity of infection of 2 or 5, depending on experiment. Indirect

immunofluorescent confocal microscopy experiments were performed on a Leica TCS

SP8 laser scanning confocal microscope in the Integrative Physiology and

Neuroscience Advanced Imaging Center at Washington State University. RT-qPCR and

qPCR experiments were performed essentially as described (35, 37). Transcript

abundance was normalized to genome copy number for all RT-qPCR analyses. RACE

experiments were conducted using the SMARTer® RACE 5'/3' Kit (Takara Bio) with

minor modifications as noted in *SI Materials and Methods*. RACE products were

isolated by gel extraction using the Macherey-Nagel Nucleospin PCR/gel clean-up kit

(Takara Bio) and sent to Eurofins Genomics, LLC for sequencing. Sequenced RACE

products were mapped to the *C. trachomatis* L2 434/Bu genome (NCBI Accession:

NC_010287) by nucleotide BLAST on the NCBI server. The *E. coli* YtgR reporter assay

was performed essentially as described (34) with minor modifications as indicated in *SI*

Materials and Methods. Briefly, BL21 (DE3) *E. coli* were co-transformed with the

indicated pCCT and pET vectors and clonal populations were selected on double-

selective media. Clones were cultured in double-selective media and recombinant YtgR

expression was induced by the addition of IPTG prior to induction of *lacZ* expression by

the addition of L-arabinose. Cell lysates were collected and β-galactosidase activity was

measured by the Miller Assay (63). All plasmids and primers used in this study are listed

in Tables S1 and S2, respectively.

Acknowledgements. We thank Dr. Amanda Brinkworth, Liam Caven, Korinn Murphy

and Matthew Romero for critical review of this manuscript; Christopher C. Thompson for

the establishment of the *E. coli* YtgR reporter system and generation of the pCCT

construct; Dr. David Dewitt for expert advice, training and maintenance of equipment in

the IPN Advanced Imaging Center. This work was supported by NIH grants 5RO1-

AI065545-07 to R.A.C.; 1F31AI136295-01A1 and 5T32GM008336-29 to N.D.P.; N.D.P.

was also supported by an Achievement Reward for College Scientists (ARCS; Seattle

Chapter) Fellowship.

References

1.   Skaar EP (2010) The battle for iron between bacterial pathogens and their

vertebrate hosts. *PLoS Pathog* 6(8):1–2.

2.   Eisenreich W, Dandekar T, Heesemann J, Goebel W (2010) Carbon metabolism

of intracellular bacterial pathogens and possible links to virulence. *Nat Rev*

Microbiol 8(6):401–412.

3.   Brown SA, Palmer KL, Whiteley M (2008) Revisiting the host as a growth

714   medium. *Nat Rev Microbiol* 6(9):657–666.

715   4.   Ray K, Marteyn B, Sansonetti PJ, Tang CM (2009) Life on the inside: The

716   intracellular lifestyle of cytosolic bacteria. *Nat Rev Microbiol* 7(5):333–340.

717   5.   Hood MI, Skaar EP (2012) Nutritional immunity: Transition metals at the

718   pathogen-host interface. *Nat Rev Microbiol* 10(8):525–537.

719   6.   CDC (2017) *2016 Sexually Transmitted Diseases Surveillance - Chlamydia*

720   Available at: https://www.cdc.gov/std/stats16/chlamydia.htm.

721   7.   Taylor HR, Burton MJ, Haddad D, West S, Wright H (2014) Trachoma. *Lancet*

722   384(9960):2142–2152.

723   8.   Newman L, et al. (2015) Global Estimates of the Prevalence and Incidence of

724   Four Curable Sexually Transmitted Infections in 2012 Based on Systematic

725   Review and Global Reporting. *PLoS One* 10(12):1–17.

726   9.   Hafner LM (2015) Pathogenesis of fallopian tube damage caused by Chlamydia

727   trachomatis infections. *Contraception* 92(2):108–115.

728   10.   Moore ER, Ouellette SP (2014) Reconceptualizing the chlamydial inclusion as a

729   pathogen-specified parasitic organelle: an expanded role for Inc proteins. *Front*

730   *Cell Infecit Microbiol* 4(October):1–10.

731   11.   AbdelRahman YM, Belland RJ (2005) The chlamydial developmental cycle.

732   *FEMS Microbiol Rev* 29(5):949–959.

733   12.   Clarke IN (2011) Evolution of Chlamydia trachomatis. *Ann N Y Acad Sci* 1230:11–

734   18.

13.   Stephens RS, et al. (1998) Genome sequence of an obligate intracellular

pathogen of humans: Chlamydia trachomatis. *Science (80-)* 282(5389):754–759.

14.   Wyrick PB (2010) Chlamydia trachomatis Persistence In Vitro: An Overview. *J*

Infect Dis 201:88–95.

15.   Byrne GI, Lehmann LK, Landry GJ (1986) Induction of tryptophan catabolism is

the mechanism for gamma-interferon-mediated inhibition of intracellular

Chlamydia psittaci replication in T24 cells. *Infect Immun* 53(2):347–351.

16.   Taylor MW, Feng GS (1991) Relationship between interferon-gamma,

indoleamine 2,3-dioxygenase, and tryptophan catabolism. *FASEB J* 5(11):2516–

2522.

17.   Fehlner-Gardiner C, et al. (2002) Molecular basis defining human Chlamydia

trachomatis tissue tropism: A possible role for tryptophan synthase. *J Biol Chem*

277(30):26893–26903.

18.   Macchiarulo A, Camaioni E, Nuti R, Pellicciari R (2009) Highlights at the gate of

tryptophan catabolism□: a review on the mechanisms of activation and regulation

of indoleamine 2,3-dioxygenase (IDO), a novel target in cancer disease. *Amino*

Acids 37:219–229.

19.   Wood H, Roshick C, McClarty G (2004) Tryptophan recycling is responsible for

the interferon-? resistance of Chlamydia psittaci GPIC in indoleamine

dioxygenase-expressing host cells. *Mol Microbiol* 52(3):903–916.

20.   Carlson JH, Wood H, Roshick C, Caldwell HD, McClarty G (2006) In vivo and in

vitro studies of Chlamydia trachomatis TrpR:DNA interactions. *Mol Microbiol*

59(6):1678–1691.

21.   Akers JC, Tan M (2006) Molecular mechanism of tryptophan-dependent

transcriptional regulation in Chlamydia trachomatis. *J Bacteriol* 188(12):4236–

4243.

22. Wood H, et al. (2003) Regulation of tryptophan synthase gene expression in

Chlamydia trachomatis. *Mol Microbiol* 49(5):1347–1359.

23. Merino E, Yanofsky C (2005) Transcription attenuation: A highly conserved

regulatory strategy used by bacteria. *Trends Genet* 21(5):260–264.

24. Cassat JE, Skaar EP (2013) Iron in Infection and Immunity. *Cell Host Microbe*

13(5):509–519.

25. Raulston JE (1997) Response of Chlamydia trachomatis serovar E to iron

restriction in vitro and evidence for iron-regulated chlamydial proteins. *Infect*

Immun 65(11):4539–4547.

26. Ouellette SP, Carabeo RA (2010) A functional slow recycling pathway of

transferrin is required for growth of Chlamydia. *Front Microbiol* 1(OCT):1–12.

27. Igietseme JU, Ananaba GA, Candal DH, Lyn D, Black CM (1998) Immune control

of Chlamydial growth in the human epithelial cell line RT4 involves multiple

mechanisms that include nitric oxide induction, tryptophan catabolism and iron

deprivation. *Microbiol Immunol* 42(9):617–625.

28. Nairz M, et al. (2008) Interferon-γ limits the availability of iron for intramacrophage

Salmonella typhimurium. *Eur J Immunol* 38(7):1923–1936.

29. Byrd T, Horwitz MA (1989) Interferon gamma-activated human monocytes

downregulate transferrin receptors and inhibit the intracellular multiplication of

Legionella pneumophila by limiting the availability of iron. *J Clin Invest*

83(5):1457–1465.

30.   Byrd TF, Horwitz MA (1993) Regulation of transferrin receptor expression and

ferritin content in human mononuclear phagocytes: Coordinate upregulation by

iron transferrin and downregulation by interferon gamma. *J Clin Invest* 91(3):969–

976.

31.   Pokorzynski ND, Thompson CC, Carabeo RA (2017) Ironing Out the

Unconventional Mechanisms of Iron Acquisition and Gene Regulation in

Chlamydia. *Front Cell Infect Microbiol* 7(September):1–19.

32.   Miller JD, Sal MS, Schell M, Whittimore JD, Raulston JE (2009) Chlamydia

trachomatis YtgA is an iron-binding periplasmic protein induced by iron restriction.

Microbiology 155(9):2884–2894.

33.   Akers JC, HoDac H, Lathrop RH, Tan M (2011) Identification and functional

analysis of CT069 as a novel transcriptional regulator in Chlamydia. *J Bacteriol*

193(22):6123–6131.

34.   Thompson CC, Nicod SS, Malcolm DS, Grieshaber SS, Carabeo RA (2012)

Cleavage of a putative metal permease in Chlamydia trachomatis yields an iron-

dependent transcriptional repressor. *Proc Natl Acad Sci U S A* 109(26):10546–51.

35.   Thompson CC, Carabeo RA (2011) An optimal method of iron starvation of the

obligate intracellular pathogen, Chlamydia trachomatis. *Front Microbiol* 2(20).

doi:10.3389/fmicb.2011.00020.

36.   Dill BD, Dessus-Babus S, Raulston JE (2009) Identification of iron-responsive

proteins expressed by Chlamydia trachomatis reticulate bodies during intracellular

growth. *Microbiology* 155(1):210–219.

37.   Brinkworth AJ, Wildung MR, Carabeo RA (2018) Genomewide Transcriptional

Responses of Iron-Starved Chlamydia trachomatis Reveal Prioritization of

Metabolic Precursor Synthesis over Protein Translation. *mSystems* 3(1):e00184-

17.

38.   Raulston JE, et al. (2007) Identification of an iron-responsive protein that is

antigenic in patients with Chlamydia trachomatis genital infections. *FEMS*

Immunol Med Microbiol 51(3):569–576.

39.   Belland RJ, et al. (2003) Transcriptome analysis of chlamydial growth during IFN-

gamma-mediated persistence and reactivation. *Proc Natl Acad Sci U S A*

100(26):15971–15976.

40.   Østergaard O, et al. (2016) Quantitative Protein Profiling of Chlamydia

trachomatis Growth Forms Reveals Defense Strategies Against Tryptophan

Starvation. *Mol Cell Proteomics* 15(12):3540–3550.

41.   Ricci S, Ratti G, Scarlato V (1995) Transcriptional regulation in the Chlamydia

trachomatis pCT plasmid. *Gene* 154(1):93–98.

42.   Tao X, Boydt J, Murphy JR (1992) Specific binding of the diphtheria tox regulatory

element DtxR to the tox operator requires divalent heavy metal ions and a 9-base-

pair interrupted palindromic sequence. *Proc Natl Acad Sci U S A* 89(13):5897–

5901.

43.   Schmitt MP (2002) Analysis of a DtxR-Like Metalloregulatory Protein, MntR, from

Corynebacterium diphtheriae That Controls Expression of an ABC Metal

Transporter by an Mn(2+)-Dependent Mechanism. *J Bacteriol* 184(24):6882–

6892.

44.   Colin J, et al. (2014) Roadblock termination by reb1p restricts cryptic and

readthrough transcription. *Mol Cell* 56(5):667–680.

45.   Ouellette SP, Rueden KJ, Rucks EA (2016) Tryptophan codon-dependent transcription in chlamydia pneumoniae during gamma interferon-mediated tryptophan limitation. *Infect Immun* 84(9):2703–2713.

46.   Gralla JD (1989) Bacterial gene regulation from distant DNA sites. *Cell* 57(2):193–195.

47.   Sankar A (1989) Multipartite genetic control elements: communication by DNA loop. *Annu Rev Genet* 23(53):227–250.

48.   Wu HC, Tyson KL, Cole JA, Busby SJW (1998) Regulation of transcription initiation at the Escherichia coli nir operon promoter: A new mechanism to account for co-dependence on two transcription factors. *Mol Microbiol* 27(2):493–505.

49.   Yanofsky C (1981) Attenuation in the control of expression of bacterial operons. *Nature* 289(5800):751–758.

50.   Yanofsky C, Kelley RL, Horn V (1984) Repression is relieved before attenuation in the trp operon of Escherichia coli as tryptophan starvation becomes increasingly severe. *J Bacteriol* 158(3):1018–1024.

51.   Gollnick P, Babitzke P, Antson A, Yanofsky C (2005) Complexity in regulation of tryptophan biosynthesis in Bacillus subtilis. *Annu Rev Genet* 39(October):47–68.

52.   Mueller KE, Wolf K, Fields KA (2016) Gene Deletion by Fluorescence-Reported Allelic Exchange Mutagenesis in Chlamydia trachomatis. 7(1):1–9.

53.   Ouellette SP (2018) Feasibility of a Conditional Knockout System for Chlamydia Based on CRISPR Interference. *Front Cell Infect Microbiol* 8(February).

doi:10.3389/fcimb.2018.00059.

54.   Sasaki-Imamura T, Yoshida Y, Suwabe K, Yoshimura F, Kato H (2011) Molecular

basis of indole production catalyzed by tryptophanase in the genus Prevotella.

FEMS Microbiol Lett 322(1):51–59.

55.   Ziklo N, Huston WM, Hocking JS, Timms P (2016) Chlamydia trachomatis Genital

Tract Infections: When Host Immune Response and the Microbiome Collide.

Trends Microbiol 24(9):750–765.

56.   Valenti P, et al. (2018) Role of Lactobacilli and Lactoferrin in the Mucosal

Cervicovaginal Defense. *Front Immunol* 9(March):1–14.

57.   Spear GT, et al. (2011) Multiplex immunoassay of lower genital tract mucosal fluid

from women attending an urban STD clinic shows broadly increased IL1ß and

lactoferrin. *PLoS One* 6(5):1–7.

58.   Kelver ME, et al. (1996) Estrogen regulation of lactoferrin expression in human

endometrium. *Am J Reprod Immunol* 36(5):243–247.

59.   Cohen M, Britigan B, French M, Bean K (1987) Preliminary observations on

lactoferrin secretion in human vaginal mucus: variation during the menstrual

cycle, evidence of hormonal regulation, and implications for infection with

Neisseria gonorrhoeae. *Am J Obstet Gynecol* 157(5):1122–1125.

60.   Lloyd JM, O'Dowd T, Driver M, Tee DE (1984) Demonstration of an epitope of the

transferrin receptor in human cervical epithelium--a potentially useful cell marker.

J Clin Pathol 37(2):131–135.

61.   Nogueira AT, Braun KM, Carabeo RA (2017) Characterization of the Growth of

Chlamydia trachomatis in In Vitro-Generated Stratified Epithelium. *Front Cell*

Infect Microbiol 7(October):1–16.

62.   Case EDR, Akers JC, Tan M (2011) CT406 encodes a chlamydial ortholog of

NrdR, a repressor of ribonucleotide reductase. *J Bacteriol* 193(17):4396–4404.

63.   Miller JH (1972) Assay of β-galactosidase. *Experiments in Molecular Genetics*

(Cold Spring Harbor Laboratory Press).

Figure Legends

Fig. 1. Brief iron limitation via 2,2-bipyridyl treatment precedes the onset of

characteristic chlamydial persistence. (*A*) *C. trachomatis* L2-infected HeLa cells were

fixed and stained with convalescent human sera to image inclusion morphology by

confocal microscopy following Bpdl treatment at the indicated times post-infection.

Arrowheads indicate inclusions with visibly fewer organisms in the 12-hour Bpdl-treated

condition. Figure shows representative experiment of three biological replicates. (*B*)

Genomic DNA (gDNA) was harvested from infected HeLa cells at the indicated times

post-infection under iron-replete (blue) and -deplete (red) conditions. Chlamydial

genome copy number was quantified by qPCR. Chlamydial genome replication is

stalled following 12 hours of Bpdl treatment, but not 6. N=2. (*C*) Total RNA was

harvested from infected HeLa cells at the indicated times post-infection under iron-

replete (teal) and -deplete (orange) conditions. The transcript abundance of hallmark

persistence genes *euo* and (*D*) *omcB* were quantified by RT-qPCR and normalized

against genome copy number. Only at 12 hours of Bpdl treatment is *omcB* expression

significantly affected. N=3 for 12+6, N=2 for 12+12. Statistical significance was

determined by One-Way ANOVA followed by post-hoc pairwise *t*-tests with Bonferroni's

correction for multiple comparisons. * = $p < 0.05$, ** = $p < 0.01$, *** = $p < 0.005$.

Fig. 2. Brief iron limitation condition produces mild iron-starved transcriptional

phenotype. (*A*) Total RNA and gDNA was harvested from infected HeLa cells at the

indicated times post-infection under iron-replete (teal) and -deplete (orange) conditions.

The transcript abundance of iron-regulated *ytgA*, (*B*) *ahpC,* (*C*) *devB* and (*D*) non-iron

regulated *dnaB* were quantified by RT-qPCR and normalized against genome copy

number. The transcript expression of *ytgA* and *ahpC* were significantly elevated

following 6-hour BpdI treatment, indicative or iron starvation to *C. trachomatis.* N=3.

Statistical significance was determined by One-Way ANOVA followed by post-hoc

pairwise *t*-tests with Bonferroni's correction for multiple comparisons. * = $p < 0.05$, ** = p

< 0.01, *** = $p < 0.005$.

Fig. 3. Expression of the *trpRBA* operon in *C. trachomatis* is non-uniformly regulated by

brief iron limitation. (*A*) Cartoon depiction of the *trpRBA* operon (drawn to scale) with the

primary transcriptional start site upstream of *trpR* annotated. (*B*) Total RNA and gDNA

were harvested from infected HeLa cells at the indicated times post-infection under Trp-

replete (black) and -deplete (red) conditions. The transcript expression of *trpRBA*

operon was quantified by RT-qPCR and normalized against genome copy number. All

three ORFs are significantly induced relative to 12 hpi following Trp starvation. N=2. (*C*)

Total RNA and gDNA were harvested from infected HeLa cells at the indicated times

post-infection under iron-replete (blue) and -deplete (red) conditions. The transcript

expression of *trpRBA* operon was quantified by RT-qPCR and normalized against

genome copy number. Only *trpB* and *trpA* expression was significantly induced relative

to 12 hpi. N=3. Statistical significance was determined by One-Way ANOVA followed by

post-hoc pairwise *t*-tests with Bonferroni's correction for multiple comparisons. * = $p <$

0.05, ** = $p < 0.01$, *** = $p < 0.005$.

Fig. 4. Iron-dependent induction of *trpBA* expression initiates within the *trpRBA*

intergenic region from a novel alternative transcriptional start site. (*A*) Total RNA was

harvested from infected HeLa cells at the indicated times post-infection to examine iron-

dependent and Trp-dependent changes in the 5'-cDNA profile of the *trpRBA* operon by

Rapid Amplification of 5' cDNA Ends (5'-RACE). RACE products were separated on an

agarose gel, revealing three distinct and specific bands with apparent sizes of 1.5, 1.1

and 1.0 kb. Trp depletion led to the apparent enrichment of the 1.5 kb product, while

Bpdl treatment produced a similarly enriched 1.0 kb RACE product. Figure shows

representative experiment of three biological replicates. (*B*) To confirm that iron-

dependent induction of *trpBA* could originate from alternative transcription initiation, RT-

qPCR was performed on 5'-RACE cDNA to quantify the abundance of *trpB* transcripts

relative to *trpR*. Only under iron-limited conditions were *trpB* transcripts enriched relative

to *trpR*. N=3. Statistical significance determined by One-way ANOVA followed by post-

hoc pairwise *t*-tests. * = $p < 0.05$, ** = $p < 0.01$, *** = $p < 0.005$. (*C*) The nucleotide

position of the 5' cDNA ends generated from RACE were mapped to the *C. trachomatis*

L2 434/Bu genome by nucleotide BLAST. Figure displays histogram (semi-continuous;

bin width=20) and overlaid density plot (continuous) distribution of 5' nucleotide

positions generated from each 5'-RACE product. The dotted line represents the

weighted mean of the distribution, as indicated by the integer value above each line.

The identified alt. TSSs are depicted on the *trpRBA* operon (drawn to scale) above the

plot. N=3.

Fig. 5. Ectopically expressed YtgR domain is capable of binding the putative *trpBA*

promoter element in an operator-specific manner and repress transcription in a

heterologous system. (*A*) Identification of putative YtgR operator sequence by local and

global nucleotide sequence alignment using EMBOSS Water and Needle algorithms,

respectively, to align the previously identified YtgR operator to both the *trpRBA* IGR and

palindromic candidate sequence. The palindrome was then mutated in our YtgR

repression assay as depicted to abolish palindromicity and AT-richness. (*B*) Ectopic

expression of YtgR significantly represses β-galactosidase activity only from the

promoter of its own operon, *ytgABCD*, and not from an empty vector or another iron-

regulated but presumably non-YtgR targeted promoter, *lpdA*. $3 \leq N \geq 2$. (*C*) Expression

of recombinant YtgR represses β-galactosidase activity from the putative *trpBA*

promoter element, but not the *trpR* promoter, and this repression is dependent on the

unaltered operator sequence identified in Fig. 5A. $3 \leq N \geq 2$. Statistical significance

determined by two-sided unpaired Student's *t*-test with Welch's correction for unequal

variance. $* = p < 0.05$, $** = p < 0.01$, $*** = p < 0.005$.

Fig. 6. Transcription termination at *trpRBA* YtgR operator site coincides with iron-

dependent transcription read-through. (*A*) Total RNA was harvested from *C.*

trachomatis-infected HeLa cells to analyze 3'-cDNA landscape downstream of *trpR*

promoter. The top panel displays representative RT-PCR of full-length *trpRBA* message

across experimental conditions (NTC = No Template Control). Bottom panel depicts

electrophoresed 3'-RACE products and estimated sizes. N=3. (*B*) 3'-RACE products

were sequenced and mapped to *C. trachomatis* L2 434/Bu genome by nucleotide

BLAST. Figure displays histogram (semi-continuous; bin width=20) and overlaid density

plot (continuous) distribution of 3' nucleotide positions generated from each 3'-RACE

product. The dotted line represents the weighted mean of the distribution, as indicated

by the integer value above each line. The identified alt. TTSs are depicted on the

trpRBA operon (drawn to scale) above the plot. The 0.40 kb RACE product mapped to a

region overlapping the predicted YtgR operator site. N=3. (*C*) Analysis of transcription

read-through by RT-qPCR was performed on 3'-RACE cDNAs at three distinct loci

across the *trpRBA* operon representing upstream transcription initiation (511,416-531),

YtgR operator site termination (511,639-764) and terminal *trpBA* transcription (513,856-

968). Abundance of each amplicon was normalized to a region (Read-through)

predicted to be transcribed only as a part of the full-length product based on 5' and 3'-

RACE data (511,792-512,080). Thus, the ratio of each amplicon to the normalization

amplicon represents the proportion of that amplicon encoded as part of the full-length

transcript, approaching one as the two more closely coincide. At the YtgR operator

termination site, iron limitation reduces the ratio relative to 12 hpi, suggesting that

transcription read-through increases at this site under this condition. The transcription

initiation ratio is unaffected by iron limitation, while the terminal *trpBA* amplicon is

increased, consistent with alternative transcription from the alt. TSS. Statistical

significance determined by One-way ANOVA followed by post-hoc pairwise *t*-tests. * = *p*

< 0.05, ** = *p* < 0.01, *** = *p* < 0.005.

Author Contributions: N.D.P. and R.A.C. wrote the manuscript; N.D.P. and R.A.C.

designed the experiments; N.D.P. performed the experiments; N.D.P. and R.A.C.

analyzed and interpreted the data.