1

The Repertoire of Mutational Signatures in Human Cancer

Ludmil B Alexandrov[1]*, Jaegil Kim[2]*, Nicholas J Haradhvala[2,3]*, Mi Ni Huang[4]*, Alvin WT Ng[4], Yang Wu[4], Arnoud Boot[4], Kyle R Covington[5], Dmitry A Gordenin[6], Erik N Bergstrom[1], S M Ashiqul Islam[1], Nuria Lopez-Bigas[7,8,9], Leszek J Klimczak[10], John R McPherson[4], Sandro Morganella[11], Radhakrishnan Sabarinathan[7,8], David A Wheeler[5], Ville Mustonen[12], the PCAWG Mutational Signatures Working Group, Gad Getz[2,3,13,14]**, Steven G Rozen[4]**§, Michael R Stratton[11]**§

[1]Department of Cellular and Molecular Medicine and Department of Bioengineering and Moores Cancer Center, University of California, La Jolla, San Diego, CA, USA

[2]Broad Institute, Cambridge, MA, USA

[3]Center for Cancer Research, Massachusetts General Hospital, Boston, MA 02129, USA

[4]Centre for Computational Biology and Programme in Cancer & Stem Cell Biology, Duke NUS Medical School, Singapore

[5]Human Genome Sequencing Center, and Dan L. Duncan Cancer Center, Baylor College of Medicine, Houston, TX, USA

[6]Genome Integrity and Structural Biology Laboratory, National Institute of Environmental Health Sciences, US National Institutes of Health, Research Triangle Park, NC, USA

[7]Institute for Research in Biomedicine (IRB Barcelona), The Barcelona Institute of Science and Technology, Baldiri Reixac, 10, 08028 Barcelona, Spain

[8]Research Program on Biomedical Informatics, Universitat Pompeu Fabra, Barcelona, Spain

[9]Institució Catalana de Recerca i Estudis Avançats (ICREA), Barcelona, Spain

[10]Integrative Bioinformatics Support Group, National Institute of Environmental Health Sciences, US National Institutes of Health, Research Triangle Park, NC, USA

[11]Wellcome Sanger Institute, Wellcome Genome Campus, Hinxton, Cambridge CB10 1SA, UK

[12]Organismal and Evolutionary Biology Research Programme, Department of Computer Science, Institute of Biotechnology, University of Helsinki, Helsinki, Finland

[13]Department of Pathology, Massachusetts General Hospital, Boston, MA 02114, USA

[14]Harvard Medical School, Boston, MA 02215, USA

*Denotes equal contribution
**Denotes equal supervisory contribution

§Correspondence and requests for materials should be addressed to Steven G Rozen (steverozen@gmail.com) and Michael R Stratton (mrs@sanger.ac.uk).

ABSTRACT
Somatic mutations in cancer genomes are caused by multiple mutational processes each of
which generates a characteristic mutational signature. Using 84,729,690 somatic mutations
from 4,645 whole cancer genome and 19,184 exome sequences encompassing most cancer
types we characterised 49 single base substitution, 11 doublet base substitution, four
clustered base substitution, and 17 small insertion and deletion mutational signatures. The
substantial dataset size compared to previous analyses enabled discovery of new signatures,
separation of overlapping signatures and decomposition of signatures into components that
may represent associated, but distinct, DNA damage, repair and/or replication mechanisms.
Estimation of the contribution of each signature to the mutational catalogues of individual
cancer genomes revealed associations with exogenous and endogenous exposures and
defective DNA maintenance processes. However, many signatures are of unknown cause.
This analysis provides a systematic perspective on the repertoire of mutational processes
contributing to the development of human cancer including a comprehensive reference set
of mutational signatures in human cancer.
55
56

INTRODUCTION

Somatic mutations in cancer genomes are caused by mutational processes of both exogenous and endogenous origins that have operated during the cell lineage between the fertilised egg and the cancer cell[1]. Each mutational process may involve components of DNA damage/modification, DNA repair and DNA replication, any of which may be normal or abnormal, and generates a characteristic mutational signature that may incorporate base substitutions, small insertions and deletions, genome rearrangements, and chromosome copy number changes[2]. The catalogue of mutations from an individual cancer genome may have been generated by multiple mutational processes and thus incorporates multiple superimposed mutational signatures. Therefore, in order to systematically characterise the mutational processes contributing to cancer, mathematical methods have been developed that can be used to *(i)* decipher mutational signatures from a set of somatic mutational catalogues, *(ii)* estimate the numbers of mutations attributable to each signature in each sample, and *(iii)* annotate each mutation class in each tumour with the probability of arising from each signature[3-15].

Previous studies of multiple cancer types identified >30 single base substitution signatures, some of known but many of unknown aetiologies, some ubiquitous and others rare, some part of normal cell biology and others associated with abnormal exposures or operative during neoplastic progression[6,16-27]. Six genome rearrangement signatures have also been identified in breast cancer[18] and further patterns of rearrangements have been described[13,28-30]. However, analysis of other mutation classes has been relatively limited[31,17,18,32,33].

Thus far, mutational signature analysis has predominantly used cancer exome sequences. However, the many fold greater numbers of somatic mutations in whole-genome sequences provide substantially increased power for signature decomposition, enabling better separation of partially correlated signatures and extraction of signatures that contribute relatively small numbers of mutations. Furthermore, technical artefacts and differences in sequencing technologies and mutation calling algorithms can themselves generate mutational signatures. Therefore, the uniformly processed and highly curated sets of all classes of somatic mutations from the 2,780 cancer genome sequences of the Pan Cancer Analysis of Whole-Genomes (PCAWG) project[34,35], combined with almost all other cancer genomes and exomes for which suitable mutational catalogues are publicly available, https://www.synapse.org/#!Synapse:syn11801788, presents a notable opportunity to establish the repertoire of mutational signatures and to determine their activities across the range of cancer types.

RESULTS
Cancer genomes and somatic mutations

Somatic mutational catalogues from 23,829 samples of most cancer types, including the 2,780 highly curated PCAWG whole-genomes[34,35], 1,865 additional whole-genomes and 19,184 exomes were studied. From these, 79,793,266 somatic single base substitutions, 814,191 doublet base substitutions and 4,122,233 small insertions and deletions (indels) were analysed for mutational signatures, ~10–fold more mutations than any previous study (https://www.synapse.org/#!Synapse:syn11801889)[4,36].

104
To enable mutational signature analysis classifications were developed for each type of
mutation. For single base substitutions, the primary classification comprised 96 classes
constituted by the six base substitutions C>A, C>G, C>T, T>A, T>C, T>G (in which the
mutated base is represented by the pyrimidine of the Watson-Crick base pair) plus the
flanking 5' and 3' bases (https://cancer.sanger.ac.uk/cosmic/signatures/SBS). In some
analyses, two flanking bases 5' and 3' to the mutated base were considered (generating
1,536 classes) or mutations within transcribed genome regions were selected and classified
according to whether the pyrimidine of the mutated base pair fell on the transcribed or
untranscribed strand (192 classes). A classification was also derived for doublet base
substitutions (78 classes, https://cancer.sanger.ac.uk/cosmic/signatures/DBS). Indels were
classified as deletions or insertions and, when of a single base, as C or T and according to the
length of the mononucleotide repeat tract in which they occurred. Longer indels were
classified as occurring at repeats or with overlapping microhomology at deletion
boundaries, and according to the size of indel, repeat, and microhomology (83 classes,
https://cancer.sanger.ac.uk/cosmic/signatures/ID).
120
Mutational signature analysis
The mutational catalogues from the 2,780 PCAWG whole-genome, 1,865 additional whole-
genome, and 19,184 exome sequences of cancer were analysed separately
(https://doi.org/10.7303/syn11801889)[34,35]. For each of these catalogue sets, signature
extraction was conducted using methods based on nonnegative matrix factorisation
(NMF)[3,6] on each cancer type individually and also on all cancer types together. Analyses
were carried out separately for single base substitutions (SBS signatures), doublet base
substitutions (DBS signatures) and indels (ID signatures) and also for the three mutation
types together (1697 mutation classes if the 1536 classes of SBS in pentanucleotide context
was employed) generating composite signatures.
131
Mutational signatures were extracted using two independently developed NMF-based
methods: *(i)* SigProfiler, a further elaborated version of the framework used to generate the
signatures shown in the previous version of the COSMIC compendium of mutational
signatures (COSMICv2)[3,18,36-38], and *(ii)* SignatureAnalyzer, based on a Bayesian variant of
NMF used in several previous publications[6,15,39,40]. NMF determines both the signature
profiles and the contributions of each signature to each cancer genome as part of its
factorization of the input matrix of mutation spectra. However, given a substantial number
of signatures and/or heterogeneous mutation burdens across samples, it is possible to
reconstruct the mutations observed in a particular sample in multiple ways, often with very
small and/or biologically implausible contributions from many signatures. Therefore, each
method developed a separate procedure to estimate the contributions of signatures to each
sample (Methods).
144
We tested SignatureAnalyzer and SigProfiler on 11 sets of synthetic data, encompassing a
total of 64,400 synthetic samples, in which known signature profiles were used to generate
catalogues of synthetic mutational spectra. Both approaches performed well in re-extracting
the known signatures in realistically complex data. The tests highlighted the importance of,
and challenges in, selecting the number of signatures, because extracted signatures
discordant from the known input usually arose from difficulty in selecting the correct

number of signatures. Thus, these tests confirmed that use of NMF-based approaches to
extract signatures is not a purely algorithmic process. Instead, signature extraction requires
human judgement that considers all of the available data, including evidence from
experimental delineation of mutational signatures and the literature on DNA damage and
repair, and prior evidence of biological plausibility. In addition, signature extraction requires
human-guided sensitivity analysis to confirm that extractions from different groupings of
tumours yield essentially the same signatures. These types of evidence and techniques were
used in the determination of the signature profiles reported here. The findings we report
from tests on synthetic data are consistent with results regarding NMF, and the related
areas of probabilistic topic modelling and latent Dirichlet allocation, in multiple problem
domains[41-43]. It is widely understood that the choice of the number of latent variables (for
our purposes, the number of mutational signatures) is rarely amenable to complete
automation. In further simulations, we also found that mutation catalogues from whole
genomes allowed substantially better signature extraction than the much smaller catalogues
from whole exomes and that signature extraction on whole genome data from half as many
tumours would have supported inferior signature extraction. See Methods for further
details; all results are at https://doi.org/10.7303/syn18497223 and a summary can be found
at https://doi.org/10.7303/syn18511087.1.
169
The results of SigProfiler and SignatureAnalyzer exhibited many similarities, and we assigned
the same identifiers to similar signatures extracted by the two methods
https://www.synapse.org/#!Synapse:syn12016215. However, there were also noteworthy
differences. The number of SBS signatures found in low mutation burden tumours in the
PCAWG set (94.4% of cases that harbour 47% of mutations) was similar: 31 by SigProfiler
and 35 by SignatureAnalyzer. The number of additional SBS signatures extracted from
hyper-mutated PCAWG samples (5.6% of cases and 53% of mutations), however, was
different: 13 by SigProfiler and 25 by SignatureAnalyzer. There were also differences in SBS
signature profiles, including among signatures found in low mutation burden cases. The
latter primarily involved "flat", relatively featureless signatures, which are mathematically
challenging to deconvolute. Finally, there were differences in signature attributions to
individual samples. In general, SignatureAnalyzer used more signatures to reconstruct the
mutational profiles (Extended Data Figure 1,
https://www.synapse.org/#!Synapse:syn12169204,
https://www.synapse.org/#!Synapse:syn12177011) and the attribution to flat signatures
was different, with SigProfiler assigning mutations to SBS5 and SBS40 and SignatureAnalyzer
using combinations of multiple signatures (Extended Data Figure 2ab,
https://www.synapse.org/#!Synapse:syn12169204). The DBS and ID signatures were
generally similar between the two methods (Extended Data Figure 2cd). These comparisons
provide a useful perspective on both the consistency and variability of signature extraction
and attribution depending on the methodology used.
191
The final sets of reference mutational signatures were determined from the PCAWG analysis
supplemented by additional signatures from the other datasets. SBS signatures using the 96
mutation classification were supported by the outcomes of analyses using the 192 and 1536
mutation classifications, the existence of individual cancer samples dominated by a
particular signature, and, where available, prior experimental evidence for certain
mutational signatures (Methods,

https://doi.org/10.7303/syn12025148, https://doi.org/10.7303/syn12009645, COSMIC at
https://cancer.sanger.ac.uk/cosmic/signatures). Each signature was allocated a number
consistent with, and extending, the COSMICv2 annotation[37]. Some previous signatures split
into multiple constituent signatures and these were numbered as before but with additional
letter suffixes (e.g., single SBS17 split into signatures SBS17a and SBS17b). DNA sequencing
and analysis artefacts also generate mutational signatures, and we indicate which signatures
are possible artefacts (https://www.synapse.org/#!Synapse:syn12009767) but do not
present them below. However, future studies employing this signature set as a reference
may consider utilizing artefact signatures for data quality control. The results of both
SignatureAnalyzer and SigProfiler were used throughout the research reported here.
However, for brevity and for continuity with the signature set previously displayed in
COSMIC[37], which has been widely used as a reference, SigProfiler results are outlined below
and SignatureAnalyzer results are provided at (Extended Data Figures 3,4,
https://www.synapse.org/#!Synapse:syn11738307).
212
Single base substitution (SBS) mutational signatures
There were substantial differences in numbers of SBSs between samples (ranging from
hundreds to millions) and between cancer types, as previously observed[44] (Figure 1). In total,
67 SBS mutational signatures were extracted, of which 49 were considered to be likely of
biological origin (Figure 2, Methods, https://cancer.sanger.ac.uk/cosmic/signatures/SBS/).
Except for SBS25, all mutational signatures reported in COSMICv2 (i.e.,
https://cancer.sanger.ac.uk/cosmic/signatures_v2) [4,23,37] were confirmed in the new set of
analyses (median cosine similarity between the newly derived signatures and those on
COSMICv2: 0.95, excluding "split" signatures which are discussed below; range 0.74 to
0.9996 https://www.synapse.org/#!Synapse:syn12016215). SBS14, SBS16, and SBS20
changed the most; for explanation, see https://cancer.sanger.ac.uk/cosmic/signatures/SBS/.
SBS25 was previously found only in cell lines derived from Hodgkin lymphomas, at least one
of which had been previously treated with chemotherapy, and, to our knowledge, no data
from primary cancers of this type are currently available. The newly derived signatures
showed much improved separation from each other and hence more distinct signature
profiles, presumably due to the substantially increased statistical power of this analysis
(online Methods section *Better separation compared to COSMICv2 signatures*).
230
Thirteen new likely real SBS signatures compared to the set previously described in
COSMICv2[37] were extracted (excluding those that are the consequence of signature
splitting). Some were in cancers with a previously unanalysed exogenous exposure (SBS42),
some were in chemotherapy treated samples which have often been excluded from
previous studies (SBS31, SBS32, SBS35) and some were rare and hence absent by chance
from previous analyses (SBS36, SBS44). Others were more common, but contributed
relatively few mutations to individual cancer genomes, or were similar to previously
discovered signatures and thus not isolated from datasets based predominantly on cancer
exome sequences (e.g., SBS38, SBS39, SBS40). Notably, SBS40 was extracted from kidney
cancer in which it appears to be required for optimal reconstruction of mutational
catalogues. It is a relatively featureless ("flat") signature, with similarity to SBS5 and other
flat signatures, and this may account for it only clearly emerging now with the availability of
whole cancer genomes. SBS40 may contribute to other cancer types but its similarity to
SBS5 renders this uncertain and larger datasets will be required to clarify the extent of its

activity. For some new signatures there were plausible underlying aetiologies (Figure 3,
Extended Data Figures 4,5): SBS31 and SBS35, prior platinum compound chemotherapy[45];
SBS32, prior azathioprine therapy; SBS36, inactivating germline or somatic mutations in
MUTYH which encodes a component of the base excision repair machinery[46,47]; SBS38,
additional effects of ultraviolet light (UV) exposure; SBS42, occupational exposure to
haloalkanes[27]; SBS44, defective DNA mismatch repair due to MLH1 inactivation[48]. SBS33,
SBS34, SBS37, SBS39, SBS40, and SBS41 are of unknown cause.
252
Three previously characterised base substitution signatures (SBS7, SBS10, SBS17) split into
multiple constituent signatures (Figure 2). We previously regarded SBS7 as a single signature
composed predominantly of C>T at C$\underline{C}$N and T$\underline{C}$N trinucleotides (the mutated base is
underlined) together with many fewer T>N mutations. It was found in malignant melanomas
and squamous skin carcinomas and is likely due to UV induced pyrimidine dimer formation
followed by translesion DNA synthesis by error-prone polymerases which predominantly
insert adenine opposite damaged bases. With the larger dataset now available, SBS7 has
decomposed into four constituent signatures: SBS7a consisting mainly of C>T at T$\underline{C}$N; SBS7b
consisting of C>T mainly at C$\underline{C}$N and to a lesser extent at T$\underline{C}$N; SBS7c and SBS7d, which
constituted relatively minor components of the previous SBS7 and consist predominantly of
T>A at N$\underline{T}$T and T>C at N$\underline{T}$T respectively[49]. Splitting of a mutational signature likely reflects
the existence of multiple distinct mutational processes, initiated by the same exposure,
which have closely, but not perfectly, correlated activities. For example, the constituent
signatures of SBS7 are probably all initiated by UV-induced DNA damage. SBS7a and SBS7b
may reflect different dipyrimidine photoproducts whereas SBS7c and SBS7d may be due to
low frequencies of misincorporation by translesion polymerases of T and G opposite
thymines in pyrimidine dimers rather than the more frequent and non-mutagenic A.
Splitting of SBS10 and SBS17 is described at
https://cancer.sanger.ac.uk/cosmic/signatures/SBS/.
272
Several base substitution signatures showed transcriptional strand bias
(https://www.synapse.org/#!Synapse:syn12009767). Transcriptional strand bias is often
attributable to transcription coupled nucleotide excision repair (TC-NER) acting on DNA
damaged by exogenous exposures which cause covalently bound bulky adducts or
crosslinking to other bases and consequent distortion of the helical structure. This results in
stalling of RNA polymerase and hence recruitment of the TC-NER machinery. An excess of
DNA damage on untranscribed compared to transcribed strands of genes may also
contribute to transcriptional strand bias[50]. Both mechanisms, however, result in more
mutations of a damaged base on the untranscribed compared to the transcribed strands of
genes. Assuming that either or both are responsible for the observed transcriptional strand
biases (which may not always be the case), DNA damage to cytosine (SBS7a, SBS7b),
guanine (SBS4, SBS8, SBS19, SBS23, SBS24, SBS31, SBS32, SBS35, SBS42), thymine (SBS7c,
SBS7d, SBS21, SBS26, SBS33) and adenine (SBS5, SBS12, SBS16, SBS22, SBS25) may underlie
these mutational signatures (see https://cancer.sanger.ac.uk/cosmic/signatures/SBS/ for
plots of strand bias). Although the likely underlying DNA damaging agents are known for
SBS4 (tobacco mutagens), SBS7a, SBS7b, SBS7c, SBS7d (UV), SBS22 (aristolochic acid), SBS24
(aflatoxin), SBS25 (prior chemotherapy), SBS31 and SBS35 (platinum compounds), SBS32
(azathioprine), and SBS42 (haloalkanes), the causes of the remainder are unknown. Indeed,
some signatures showing transcriptional strand bias are associated with defective DNA

mismatch repair (SBS21 and SBS26) and it is conceivable that, for these, exogenous DNA
damage is not involved. The extent of transcriptional strand bias appears to differ in
different sectors of the genome. For example, consideration of the whole transcribed
genome showed absent or minimal transcriptional strand bias in the APOBEC related SBS2
and SBS13 and in the defective polymerase epsilon proof-reading related SBS10a. However,
consideration of exons alone showed clear evidence of transcriptional strand bias in these
signatures   (https://cancer.sanger.ac.uk/cosmic/signatures/SBS/).   The   mechanism(s)
underlying this amplification of transcriptional strand bias in exons is unknown and appears
to be signature specific, since there is minimal difference in the extent of transcriptional
strand bias between exons and other transcribed regions for other signatures (for example,
SBS4 and SBS22).

Employing the single base substitution classification of 1536 mutation types, which uses the
pentanucleotide sequence context two bases 5' and two bases 3' to each mutated base,
yielded a set of signatures largely consistent with that based on substitutions in
trinucleotide context alone. Notably, however, the pentanucleotide context enabled the
extraction of two forms of both SBS2 and SBS13, one with mainly a pyrimidine (C or T) and
the other with a purine (A or G) at the -2 base (the second base 5' to the mutated cytosine).
These may represent the activities of the cytidine deaminases APOBEC3A and APOBEC3B,
respectively[51]. If so, APOBEC3A accounts for many more mutations than APOBEC3B in
cancers with high APOBEC activity. Several other signatures showed non-random sequence
contexts at +2 and -2 positions. In particular, the -2 bases in SBS17a and SBS17b and the -2
and +2 bases in SBS9 were predominantly A and T. In general, however, sequence context
effects were much stronger for bases immediately 5' and 3' to the mutated bases.

SBS signatures showed substantial variation in the numbers of cancer types and cancer
samples in which they were found, ranging from SBS1 and SBS5 which were present in
almost every cancer type and almost every cancer sample, to SBS23 which was only
observed in a small subset of liver cancers (Figure 3). The numbers of mutations per cancer
sample attributed to each signature also varied greatly, from a few tens of mutations for
SBS1 to millions of mutations for SBS10b. Almost all individual cancer samples exhibited
multiple signatures, with a mode of three signatures per sample in the PCAWG set
(https://www.synapse.org/#!Synapse:syn12169204). The assigned signatures reconstruct
well the mutational spectra of the cancer samples (in PCAWG samples, median cosine
similarity 0.97; 96.3% of samples with cosine similarity >0.90) (illustrative examples are
shown in Figure 4).

Clustered single base substitution mutational signatures
Some mutational processes generate mutations that cluster in small regions of the genome.
The relatively limited number of mutations generated by such processes, compared to those
acting genome-wide, may result in failure to detect their signatures by standard methods.
To obviate this problem, we first identified clustered mutations in each genome and
analysed these separately (Methods). Four main signatures associated with clustered
mutations were identified (Figure 2) and were consistent with previous reports[15,16,32]. Two
found in multiple cancer types were similar to single base substitution SBS2 and SBS13,
which have been attributed to APOBEC enzyme activity (mostly APOBEC3B) and represent
foci of *kataegis*[17,32,52]. Two additional clustered mutational signatures, one characterised by

C>T and C>G mutations at (A|G)C(C|T) trinucleotides[53] and the other T>A and T>C
mutations at (A|T)T(A|T) were found in lymphoid neoplasms and likely represent direct and
indirect consequences of activation induced cytidine deaminase (AID) mutagenesis and
translesion DNA synthesis by error-prone polymerases (SBS84 and SBS85 respectively)[15].
The possibility that further processes may generate clustered mutations is not excluded.
344
345
Doublet base substitution (DBS) mutational signatures
Tandem doublet, triplet, quadruplet, quintuplet, and sextuplet base substitutions
(https://www.synapse.org/#!Synapse:syn11801938,
https://www.synapse.org/#!Synapse:syn11726620) at immediately adjacent bases were
observed at ~1% the prevalence of single base substitutions. In most cancer genomes, the
observed number of DBSs was considerably higher than expected from random adjacency of
SBSs (https://www.synapse.org/#!Synapse:syn12177057) indicating the existence of
commonly occurring, single mutagenic events that cause substitutions at neighbouring
bases. There was substantial variation in the number of DBSs, ranging from zero to 20,818 in
a sample. Across cancer types, the numbers of DBSs were generally proportional to the
numbers of SBSs in that cancer type (Figure 1). However, colorectal adenocarcinomas had
significantly fewer DBSs than expected, and lung cancers and melanomas had more
(Extended Data Table 1). The large dataset analysed here allowed, for the first time,
systematic analysis of DBS and indel signatures (described below). Eleven DBS signatures
were extracted (Figure 2). Of these, to our knowledge, only two have been previously
reported[33] evidencing further the value of the large numbers of mutations from whole
genome data.
363
DBS1 was characterised almost exclusively by CC>TT mutations (Figure 2), contributed 100s-
10,000s of mutations in malignant melanomas (Figure 3) with SBS7a and SBS7b. DBS1
exhibited transcriptional strand bias consistent with damage to cytosines
(https://www.synapse.org/#!Synapse:syn12177063). CC>TT mutations associated with UV
induced DNA damage are well established in the literature, were previously reported in
melanomas, and are thought to be due to generation of pyrimidine dimers and subsequent
error-prone translesion DNA synthesis by polymerases that introduce adenines opposite the
damaged bases[33,54].
372
Reanalysis after exclusion of malignant melanomas and other cancers with evidence of UV
exposure still yielded a signature (termed DBS11) characterised predominantly by CC>TT
mutations and smaller numbers of other doublet base substitutions at CC and TC which
contributed 10s of mutations to many samples of multiple cancer types (Figures 2 and 3).
DBS11 was associated with SBS2 which is due to APOBEC activity. Thus, APOBEC activity may
also generate DBS11, although the mechanism by which it induces doublet base
substitutions is not well understood.
380
DBS2 was composed predominantly of CC>AA mutations, with smaller numbers of CC>AG
and CC>AT mutations, and contributed 100s-1000s of mutations in lung adenocarcinoma,
lung squamous and head and neck squamous carcinomas, which are often caused by
tobacco smoking, which has been reported previously (Figures 2 and 3)[33]. DBS2 showed
transcriptional   strand   bias   indicative   of   guanine   damage

(https://www.synapse.org/#!Synapse:syn12177064) and was associated with SBS4 which is
caused by tobacco smoke exposure. It is likely, therefore, that DBS2 can be a consequence
of DNA damage by tobacco smoke mutagens.
389
Analysis of each cancer type separately, however, revealed a signature very similar to DBS2
contributing 100s of mutations to liver cancers and 10s of mutations to cancers of other
types without evidence of tobacco smoke exposure. A pattern closely resembling DBS2 and
characterised predominantly by CC>AA mutations, together with smaller contributions of
CC>AG and CC>AT, dominates DBSs in normal mouse cells and is particularly frequent in the
liver[55]. The nature of the mutational processes underlying these doublet signatures in
smoking-unrelated human cancers and in normal mice is unknown. However, acetaldehyde
exposure in experimental systems generates a mutational signature characterised primarily
by CC>AA and lower burdens of CC>AG and CC>AT mutations together with C>A single base
substitutions[56]. Acetaldehyde is an oxidation product of alcohol and a constituent of
cigarette smoke. The role of acetaldehyde, and perhaps other aldehydes, in generating
DBS2, whether associated with tobacco smoking, alcohol consumption or in non-exposed
cells, merits further investigation[57].
403
DBS3, DBS7, DBS8 and DBS10 showed 100s-1000s of mutations in rare colorectal, stomach
and oesophageal cancers some of which showed evidence of defective DNA mismatch
repair (DBS7, DBS10) or polymerase epsilon exonuclease domain mutations (DBS3)
generating hypermutator phenotypes (Figures 2, 3). DBS5 was found in cancers previously
exposed to platinum chemotherapy and is associated with SBS31 and SBS35. The remaining
DBS signatures are of uncertain cause.
410

Small insertion and deletion (ID) mutational signatures
Indels were usually present at ~10% the frequency of base substitutions (Figure 1). There
was substantial variation between cancer genomes in numbers of indels, even when cancers
with evidence of defective DNA mismatch repair were excluded. Overall, the numbers of
deletions and insertions were similar, but there was variation between cancer types with
some showing more deletions and others more insertions of various subtypes (Figure 1).
Seventeen indel mutational signatures were extracted (Figure 2).
418
Indel signature 1 (ID1) was composed predominantly of insertions of thymine and ID2 of
deletions of thymine, both at long (≥5) thymine mononucleotide repeats (Figure 2). 10s to
100s of mutations of both signatures were found in the large majority of most cancer types
but were particularly common in colorectal, stomach, endometrial and oesophageal cancers
and in diffuse large B cell lymphoma (Figure 3). Most of these cancers are likely to be DNA
mismatch repair proficient on the basis of the relatively limited numbers of indels and
absence of the SBS signatures (SBS6, SBS14, SBS15, SBS20, SBS21, SBS26, and SBS44)
associated with DNA mismatch repair deficiency. Together, ID1 and ID2 accounted for 97%
and 45% of indels in hypermutated and non-hypermutated cancer genomes, respectively
(Extended Data Table 2), and both signatures have also been found in non-neoplastic cells[58].
They are likely due to the intrinsic tendency to slippage during DNA replication of long
mononucleotide tracts. However, the mechanistic basis for separation into two signatures,
one presumably due to slippage of the nascent strand (ID1) and the other the template

strand (ID2) is unclear. Similarly, the substantial differences in their mutation frequencies
between cancer types are not well understood.

ID3 was characterised predominantly by deletions of cytosine at short (≤5bp long)
mononucleotide cytosine repeats and exhibited 100s of mutations in tobacco smoking
associated cancers of the lung and head and neck (Figures 2 and 3). There was
transcriptional strand bias of mutations, with more guanine deletions than cytosine
deletions on the untranscribed strands of genes, compatible with TC-NER of adducted
guanine   (https://www.synapse.org/#!Synapse:syn12177065,
https://www.synapse.org/#!Synapse:syn12177066). The numbers of ID3 mutations in
cancer samples positively correlated with the numbers of SBS4 and DBS2 mutations, both of
which have been associated with tobacco smoking (Extended Data Figure 6). It is therefore
likely that DNA damage by components of tobacco smoke underlie ID3 but the
mechanism(s) by which indels are generated is unclear.

ID13 was characterised predominantly by deletions of thymine at thymine-thymine
dinucleotides and exhibited large numbers of mutations in malignant melanomas of the skin
(Figures 2 and 3). The numbers of ID13 mutations correlated with the numbers of SBS7a,
SBS7b and DBS1 mutations, which have been attributed to DNA damage induced by UV
(Extended Data Figure 6). It is, however, notable that a similar mutation of the other
pyrimidine, i.e., deletion of cytosine at cytosine-cytosine dinucleotides, does not feature
strongly in ID13, perhaps reflecting the predominance of thymine compared to cytosine
dimers induced by UV[59]. The mechanism(s) underlying thymine deletion is unclear.

ID6 and ID8 were both characterised predominantly by deletions ≥5bp (Figure 2). ID6
exhibited overlapping microhomology at deletion boundaries with a mode of 2bp and often
longer stretches. This signature was correlated with SBS3 which has been attributed to
defective homologous recombination based repair (Extended Data Figure 6). By contrast,
ID8 deletions showed shorter or no microhomology at deletion boundaries, with a mode of
1bp, and did not strongly correlate with SBS3 mutations (Figures 2 and 3). These patterns of
deletion may be characteristic of DNA double strand break repair by non-homologous
recombination based end-joining mechanisms, and if so, suggest that at least two distinct
forms of end-joining mechanism are operative in human cancer[60].

A small fraction of cancers exhibited very large numbers of ID1 and ID2 mutations (>10,000)
(Figure 3, https://cancer.sanger.ac.uk/cosmic/signatures/ID). These were usually
accompanied by SBS6, SBS14, SBS15, SBS20, SBS21, SBS26 and/or SBS44 which are
associated with DNA mismatch repair deficiency, sometimes combined with POLE or POLD1
proofreading deficiency (SBS14, SBS20)[40]. Occasional cases with these signatures
additionally showed large numbers of ID7 indels
(https://www.synapse.org/#!Synapse:syn11738668). In addition, rare samples showed large
numbers of either ID4, ID11, ID14, ID15, ID16 or ID17 mutations but did not show ID1 and
ID2 mutations or the single base substitution signatures usually associated with DNA
mismatch repair deficiency. The mechanisms underlying these signatures are unknown.

Composite mutational signatures

In the analyses described above mutational signatures were extracted for each mutation
type separately. However, mutational processes in nature generate composite signatures
that may include SBSs, DBSs, IDs, genome rearrangements and chromosome number
changes. We therefore also extracted signatures using combined catalogues of SBSs, DBSs,
and IDs (257 mutation subclasses or 1697 if the 1536 classification of single base
substitutions was used). Fifty-two composite signatures were extracted.
484
A composite signature with components similar to SBS4, DBS2 (characterised predominantly
by CC>AA mutations) and ID3 (characterised predominantly by deletion of cytosine at short
runs of cytosines) was found mainly in lung cancers, suggesting that it is the consequence of
tobacco smoke exposure (Extended Data Figure 7). Similarly, composite signatures with
components similar to SBS7a, SBS7b, DBS1 (characterised predominantly by CC>TT
mutations) and ID13 (characterised predominantly by deletion of thymine at thymine–
thymine dinucleotides) were found in skin cancers and are thus likely due to UV induced
DNA damage (Extended Data Figure 7). A further composite signature in breast and ovarian
cancers included features of SBS3 and ID6 combined with ID8 (deletions >5bp with varying
degrees of overlapping microhomology) and is likely associated with defective homologous
recombination based repair (Extended Data Figure 7). In these composite signatures
attributions of the constituent SBS, DBS and ID signatures extracted independently in the
main analyses were correlated with each other, adding support to the existence of the
composite signatures (Extended Data Figure 6). Various forms of defective DNA mismatch
repair were also associated with multiple SBS, DBS and ID signatures.
500
Correlations with age
A positive correlation between age of cancer diagnosis and the number of mutations
attributable to a signature suggests that the mutational process underlying the signature
has been operative, at a more or less constant rate, throughout the cell lineage from
fertilized egg to cancer cell, and thus in normal cells from which that cancer type
develops[4,61]. Confirming previous reports, the numbers of SBS1 and SBS5 mutations
correlated with age, exhibiting different rates in different tissue types (q-values in
https://www.synapse.org/#!Synapse:syn12030687,
https://www.synapse.org/#!Synapse:syn20317940
https://www.synapse.org/#!Synapse:syn12217988). In addition, SBS40 correlated with age
in multiple cancer types. However, given the similarity in signature profile between SBS5
and SBS40 the possibility of misattribution between these signatures cannot currently be
excluded. The numbers of DBSs and IDs were much lower than the numbers of SBSs and the
numbers of samples in which DBS and ID signatures could be attributed were also lower.
Nevertheless, DBS2 and DBS4 correlated with age and, consistent with the interpretation of
activity in normal cells, the profiles of DBS2 and DBS4 together closely resemble the
spectrum of DBS mutations found in normal mouse cells[55]. Neither DBS2 nor DBS4,
however, was clearly correlated with an SBS or ID signature that correlates with age. ID1,
ID2, ID5 and ID8 showed correlations with age in multiple tissues. ID1 and ID2 indels are
likely due to slippage at poly T repeats during DNA replication and correlated with the
number of SBS1 substitutions. SBS1 has previously been proposed to reflect the number of
mitoses a cell has experienced and thus SBS1, ID1 and ID2 may all be generated during DNA
replication at mitosis[4]. The number of ID5 mutations correlated with the number of SBS40
mutations and thus the mutational processes underlying these two age-correlated

signatures may also harbour common components. ID8 is predominantly composed of
deletions >5bp with no or 1bp of microhomology at their boundaries. These are likely due to
DNA double strand breaks which have not been repaired by homologous recombination
based mechanisms, but instead by a non-homologous-end joining mechanism. The features
of ID8 resemble those of some ionising radiation associated mutations and this may,
therefore, be an underlying aetiological factor[62]. Taken together, the results indicate that
multiple mutational processes operate in normal cells.
532
533
DISCUSSION
Cancers arise as a result of somatic mutations. Mutational signature analysis therefore
provides important insights into cancer development through comprehensive
characterisation of the underlying mutational processes. There are, however, important
constraints, limitations and assumptions in the analytic frameworks we have used that
should be recognised. Although designed to reflect the mutational consequences of
recurrent mutational processes, mutational signatures extracted from sample sets in which
multiple mutational processes are operative remain mathematical approximations, with
profiles that can be influenced by the mathematical approach used and by additional
factors, such as the other mutational processes present. For conceptual and practical
simplicity, we have assumed that there is a single signature associated with each mutational
process and have provided an average reference signature to represent it. However, we do
not discount the possibility that further nuances and variations of signature profiles exist,
for example between different tissues. Moreover, although the extent of separation
between partially correlated signatures has been improved in this analysis, some signatures
may still represent combinations of constituent signatures. Contributions from each
signature to the burden of mutations in each sample have been estimated. However, with
increasing numbers of signatures and multiple orders of magnitude differences in mutation
burdens from certain signatures, prior knowledge can help to avoid biologically implausible
results. Thus, further development of methods for deciphering mutational signatures and
attribution of mutations is warranted and this needs to be supplemented by signatures
derived from experimental systems in which the causes of the mutations are known. The
numbers of DBSs, clustered substitutions, IDs and genome rearrangements (reported in ref.
[30]) are small compared to single base substitutions. Thus, larger datasets may be required to
robustly characterise their mutational signatures. Nevertheless, the results outlined here
indicate that signatures with many similarities and some differences can be found by
different mathematical approaches, and that these are confirmed in many different ways,
including experimentally elucidated signatures[22,31,45,48,49,61,63-69] and the observation of
tumours dominated by a single signature
(https://www.synapse.org/#!Synapse:syn12016215).
564
Prior reports have provided only a relatively limited examination of doublet and indel
mutational spectra and, to the best of our knowledge, no previous comprehensive analysis
of doublet and indel mutational signatures has been performed. Here, we provide the first
systematic analysis of these mutation types by considering 83 mutational subtypes for
indels and 78 mutational subtypes for doublets. This analysis also includes almost all publicly
available exome and whole-genome cancer sequences, amounting in aggregate to 23,829
cancers of most cancer types. Some rare or geographically restricted signatures may not

have been captured and signatures of therapeutic mutagenic exposures have not been
exhaustively explored. Nevertheless, it is likely that a substantial proportion of the naturally-
occurring mutational signatures found in human cancer have now been described. This
comprehensive repertoire provides a foundation for future research into *(i)* geographical
and temporal differences in cancer incidence to elucidate underlying differences in
aetiology, *(ii)* the mutational processes and signatures present in normal tissues and caused
by non-neoplastic disease states, *(iii)* clinical and public health applications of signatures as
indicators of sensitivity to therapeutics and past exposure to mutagens, and *(iv)* mechanistic
understanding of the mutational processes underlying carcinogenesis.
581

ACKNOWLEDGEMENTS
The results here are partly based on data generated by the TCGA Research Network
(http://cancergenome.nih.gov/) and the ICGC/TCGA Pan-cancer Analysis of Whole Genomes
Network. This work was supported by Wellcome grant reference 206194 (M.R.S.), Singapore
National Medical Research Council grants NMRC/CIRG/1422/2015 and MOH-000032/MOH-
CIRG18may-0004 and the Singapore Ministry of Health via the Duke-NUS Signature Research
Programmes (M.N.H., A.W.T.N., Y.W., A.B., S.G.R.), US National Institute of Health
Intramural Research Program Project Z1AES103266 (D.A.G.), the European Research Council
Consolidator Grant 682398 (N.L.-B.), US National Cancer Institute U24CA143843 (D.A.W.),
and Cancer Research UK Grand Challenge Award C98/A24032 (E.N.B., S.M.A.I., L.B.A.,
M.R.S.). G.G and J.K were partially supported by the National Cancer Institute grants
U24CA210999 and U24CA143845. G.G. was partially supported by the Paul C. Zamecnick,
MD, Chair in Oncology at the Massachusetts General Hospital Cancer Center. N.J.H and G.G
were partially supported by G.G's start up funds at MGH.

596
597
598

Figure legends.
Figure 1. **Mutation burdens of single base substitutions, doublet base substitutions and**
small insertions and deletions for the 2,780 PCAWG tumours. Each sample is displayed
according to its tumour type. Tumour types are ordered according to the median number of
single base substitutions. The numbers of cases of each tumour type are shown. The
proportions of each mutation subclass in each sample are shown as coloured bar charts.
605
Figure 2. Profiles of single base substitution, doublet base substitution and small insertion
and deletion mutational signatures. The subclassifications of each mutation type (single
base substitutions, 96 subtypes; doublet base substitutions, 78 subtypes; indels, 83
subtypes) are described in the main text. Magnified versions of signatures SBS4, DBS2 and
ID3 (which are all associated with tobacco smoking) are shown to illustrate the positions of
each mutation subtype on each plot.
612
Figure 3. The number of mutations contributed by each mutational signature to the 2,780
PCAWG tumours. The numbers of mutations attributed are shown by cancer type. The size
of each dot represents the proportion of samples of each tumour type that show the
mutational signature. The colour of each dot represents the median mutation burden of the
signature in samples which show the signature. Contributions are shown for single base
substitution, doublet base substitution and indel mutational signatures separately.
Contributions of composite signatures to the PCAWG cancers and single base substitution
signatures to the complete set of cancer samples analysed are shown in Supplementary
information.
622
623 **Figure 4. Illustrative examples of mutational spectra of individual cancer samples.** A breast
624 cancer, a lung cancer, and a malignant melanoma and their contributory single base
625 substitution, doublet base substitution, and small insertion and deletion mutational
626 signatures.
627

628 **Online Methods**
629
Principles and strategy of mutational signature analysis adopted in this report
Conceptual principles.
• Multiple mutational processes generate the somatic mutations present in each
individual human cancer.
• Each mutational process generates a particular pattern of somatic mutations known
as a mutational signature.
• Each mutational process may incorporate a component of DNA
damage/modification, DNA repair and DNA replication, each of which may be part of
normal or abnormal cell biology. Differences in any of the three components may
result in a different mutational signature, thus, by definition, constituting a distinct
mutational process.
• Multiple mutational processes operating continuously or intermittently during the
cell lineage from the fertilised egg to the cancer cell may contribute to the aggregate
set of mutations found in the cancer cell. Thus, the catalogue of somatic mutations
from a single cancer sample often includes mutations of many different mutational
signatures.
646
647 *Aims of the study.*
648 • To decipher the mutational signatures present in essentially the full set of whole
649 genome and exome sequenced human cancers from which data is currently available
650 and subsequently to estimate the contributions of each signature to each cancer
651 genome.
652
Approach used.
• Several mathematical approaches have been used to deconvolute/extract the
mutational signatures present in a set of mutational catalogues[3,6,7,9,14-16,39,70-72]. They
are all based on the premise that different mutational processes (and thus their
signatures) contribute to different extents to different samples within the set.
• Two independently developed methods based on NMF (SigProfiler and
SignatureAnalyzer) were applied separately to the sets of mutational catalogues. By
using two methods we aimed to provide perspective on the impact different
methodologies can have on numbers of signatures generated, signature profiles and
attributions. The two methods are described in detail below and the code for both is
available (https://www.synapse.org/#!Synapse:syn11801488). Results from the two
methods have been compared (https://www.synapse.org/#!Synapse:syn12177006).
• Briefly, SigProfiler employs an elaboration of previously presented approaches for
signature extraction and for attribution of mutation counts to mutational signatures
in individual tumours[3,4,18,36].
• Briefly, SignatureAnalyzer employs a Bayesian variant of NMF[6,15,39]. This method
enables inferences for the number of signatures through the automatic relevance
determination technique and delivers highly interpretable and sparse
representations for both signature profiles and attributions at a balance between
data fitting and model complexity.

• The methods that SigProfiler and SignatureAnalyzer use for determining the number
of extracted signatures are presented in the detailed descriptions of each of these
methods, below.
• Both methods assume that the spectra of individual tumours can be represented as
linear combinations of signatures. Thus, if the combination of two simultaneously
operating mutational processes were to create a signature profile that is not a linear
combination of the two, both SigProfiler and SignatureAnalyzer would extract this as
a separate signature. We believe this is the case for SBS20, which appears to be due
to the simultaneous operation of *POLD1* mutation and mismatch repair deficiency.
682

Role of NMF in extraction and attribution of mutational signatures.
• NMF is the approximate representation of a nonnegative matrix V, in this case the
observed mutational spectra (or profiles) of a set of tumors, as the product of two
usually smaller nonnegative matrices, W and H, which are the signatures and the
attributions respectively.
• In our experience, however, calculating a single NMF is rarely sufficient to allow
confident extraction and attribution of signatures that reflect the underlying
biological mutational processes. There are two main reasons for this:
o The profiles of extracted signatures can vary substantially depending on the
692   tumour samples present in V. For example, this may be especially evident
693   when some tumors in V have high numbers of mutations (e.g., samples due
694   to UV exposure or DNA mismatch repair deficiency), while others have low
695   numbers. In situations such as this, signatures due to highly mutagenic
696   processes sometimes capture mutations from other processes and also
697   "bleed" into other signatures.
698   o With multiple potentially similar signatures operating, there are multiple
699   possible and reasonably accurate reconstruction solutions for each tumour,
700   often with many small and/or biologically implausible contributions.
701   • To address these challenges two key additional analytic features have been
702   incorporated into our analyses:
703   o Both SigProfiler and SignatureAnalyzer carried out multiple NMFs on
704   different subsets of tumours for signature extraction, and indeed, each
705   signature extraction by SigProfiler entails 1024 NMFs with different random
706   initial conditions. We describe below how we selected representative
707   mutational signature profiles.
708   o Both SigProfiler and SignatureAnalyzer developed a process of attributing
709   signature activities to tumours that is separate from the process of extracting
710   (discovering) the signatures.
711   • The use of multiple extractions to support confidence in results:
712   o SignatureAnalyzer, carried out the main extraction procedure on (1) the
713   majority of the PCAWG tumours excluding certain highly mutated tumours
714   and (2) the melanomas, microsatellite-instable tumours, and a single
715   temozolomide-exposed   tumour
716   (https://www.synapse.org/#!Synapse:syn11738314).
717   o SigProfiler extracted signatures from

- Separate extraction of SBS, DBS, and ID signatures from all PCAWG whole-genomes together (the main source of the reference mutational signature).
- Separate extraction of SBS, DBS, and ID signatures from PCAWG whole-genomes with each tumour type examined by itself.
- Extraction of SBS signatures from all non-PCAWG whole-genomes together.
- Extraction of SBS signatures from non-PCAWG whole-genomes with each tumour type examined by itself.
- Separate extraction of SBS and ID signatures from all TCGA exomes together.
- Separate extraction of SBS and ID signatures from TCGA exomes with each tumour type examined by itself.
- Separate extraction of SBS and ID signatures from all non-TCGA exomes together.
- Separate extraction of SBS and ID signatures from non-TCGA exomes with each tumour type examined by itself.

This allowed the extraction of signatures that were not present in the PCAWG tumours (e.g., SBS42, which has been attributed to haloalkane exposure and seen only in whole exome sequencing data). It also served as an important validation, as extraction of similar signatures from single tumour types and other sample sets supports the correctness of the signature extracted from the PCAWG samples (https://www.synapse.org/#!Synapse:syn12016215).

- Signature extraction from each tumour type (or from some other subset of cancers) separately has the advantages of:
    - Usually including fewer (and different) mutational signatures in each tumour type sample set than in the set of all cancers together and thus fewer (and different) opportunities for inter-signature interference.
    - Allowing multiple independent opportunities for extraction of a signature that is present in multiple tumour types, and thus of obtaining validation/confirmation of the signature's existence and profile.
    - Allowing extraction of a signature that may (for a number of reasons) fail to be extracted in analysis of all tumour types together.
    - Providing primary evidence for the existence of the signature in each tumour type.
    - Allowing separation of highly mutated cancer types/samples from cancer types/samples with low mutation burdens.
- Signature extraction from multiple tumour types together has the advantages of:
    - Usually including more samples with a particular signature than in each individual cancer type and thus being better powered to separate a signature from other partially correlated signatures and/or from signatures with similar profiles.

763   ▪ Providing a single profile for a signature rather than the multiple
764   slightly different profiles which emerge from extraction of each
765   tumour type separately.

• The profiles of the mutational signatures extracted from cancer are highly variable.
They range from some that have contributions from mutations of all subtypes in the
mutation classification ("flat" or "featureless" signatures, e.g., SBS5 and SBS40) to
others that are essentially defined by mutations at only one (or a small number) of
the mutation subtypes (e.g., signatures SBS2, SBS13, SBS10a and SBS10b). There
appears to be less concordance between the results of SigProfiler and
SignatureAnalyzer for flat signatures than for signatures with distinct features
indicating that generally, these may be more difficult to accurately extract and
distinguish from each other. However, there is experimental support for the
existence of SBS5 and SBS3[61,68].

• We represented each signature as a single reference. This selection of a single
reference signature does not exclude the possibility that signature profiles may show
nuances and further complexity and may vary in different contexts (e.g., in different
tissues). The rationale for selecting a single reference signature was the view that
this would be a level of granularity useful to most researchers. For those with
specialised interests in particular mutational processes and their components, we
also provided the signatures extracted from individual tumour types, comprising
PCAWG and non-PCAWG genomes and exomes
(https://www.synapse.org/#!Synapse:syn12025142).

• Attribution of signatures to cancer samples:
o The reference signatures from SigProfiler and SignatureAnalyzer were used to
estimate the number of mutations due to each signature in each tumour
(https://www.synapse.org/#!Synapse:syn11804065).
o SigProfiler and SignatureAnalyzer differ in their approaches for attributing
signatures. However, both incorporate a set of rules based on prior
knowledge and biological plausibility, and incorporate techniques to
encourage sparsity in the number of signatures attributed to a given tumour.
o Sparsity (limiting the numbers of signatures and limiting the numbers of
signatures attributed to each cancer sample) is an important concept and
feature of both SigProfiler and SignatureAnalyzer (both in signature
extraction and attribution). Our prior beliefs are that *(i)* there is a limited set
of significantly contributing mutational processes (and hence a limited set of
mutational signatures) operating to generate somatic mutations across all
cancers and *(ii)* that a limited set of mutational processes contribute to
individual cancer genomes (as opposed to all mutational signatures
contributing to all samples). Our aim in discovering mutational signatures is
to reflect the underlying biological processes and to attribute them
appropriately. It is not a mathematical exercise in which the main objective
and priority is to minimize the difference between $W \times H$ and the original
spectra in V. Indeed, if the latter was the main aim, for 96 mutation classes a
set of 96 signatures each constituted entirely of mutations in just one class
(and therefore ignoring sparsity), will always provide error free
reconstruction but will provide absolutely no information about underlying
mutational processes.

810
Presentation of the results of signature extraction and attribution from SigProfiler and
SignatureAnalyzer.
• The results (signatures and attributions) of the two methods have been presented
separately. We have done this in preference to combining them. We have handled
the two outputs in this way because we believe that this provides a simpler
conceptual and technical basis on which the research community can understand
the results, can employ the methods in future and can compare results with those
shown in this paper. We also do not have a basis for believing that a
combined/averaged/overlapping single result set is a better representation of the
natural truth than either of the two result sets individually and do not have a well-
founded and simple technical approach for combining them. We have, however,
provided comparisons of the outputs.
• For brevity and for continuity with previous publications, the results from SigProfiler,
a further elaborated version of previously described approaches[3,4,18,36] that
generated the 30 signatures previously shown in COSMICv2[37], are shown in the main
manuscript, and the results from SignatureAnalyzer in supplementary data
(https://www.synapse.org/#!Synapse:syn11738307).
• Nomenclature of signatures is based on and extends the nomenclature previously
used in COSMIC (COSMICv2, https://cancer.sanger.ac.uk/cosmic/signatures_v2)[37].
• Both methods analysed each mutation type (SBSs, DBSs and IDs) separately and also
together as a composite signature. In future, however, SigProfiler will usually use the
separately extracted single base substitution, indel and doublet base substitution
signatures as its standard. This generally facilitates portability, and comparison of
signature profiles with those from a variety of sample sets including targeted
sequences, exomes etc.
• SBS signatures reported in Supplementary Data include possible artefacts
(https://cancer.sanger.ac.uk/cosmic/signatures/SBS/ and see below).
838
Quality control: annotating signatures as likely real or a possible artefact
• Sequencing artefacts and differences in analysis pipelines can also generate
mutational signatures. We have annotated which signatures are likely real or
"possible artefact".
• There are multiple reasons for believing a signature reflects a biological mutational
signature rather than an artefact.
o The input data supporting the signature seem correct: key mutational
features of the putative signature look real in a mapped-read browser such as
Integrative Genomics Viewer (IGV,
https://software.broadinstitute.org/software/igv/), or characteristic mutations
are experimentally confirmed in the tumour and normal samples. Inspection
in a mapped read browser is especially important in checking for possible
problems in potentially new signatures arising in datasets other than the
highly scrutinized and checked PCAWG and TCGA sets. Features associated
with experimental, mapping, or other computational artefacts include strong
preference for the first read, very low variant allele fractions, variants in
regions of low germ-line sequencing coverage, variants found near indels in

856		low-complexity regions, variants from a signature only found in one
857		sequencing centre etc.
858	o	The 96-mutation profile and additional features (e.g., strand asymmetry,
859		association with replication timing), are known to result from a particular
860		process in experimental systems. Examples: UV, polymerase epsilon
861		proofreading deficiency, aristolochic acid and cisplatin exposure.
862	o	The putative signature is broadly consistent with previous biochemical
863		knowledge of mutational processes (e.g., preference for G adducts in
864		aflatoxin).
865	o	The putative signature dominates the spectra of some tumours (column J of
866		https://www.synapse.org/#!Synapse:syn12016215).
867	o	The putative mutational signature is consistently deciphered from multiple
868		independent datasets; this indicates that the signatures is either a common
869		sequencing artefact or something real.
870	o	The putative signature correlates with known or suspected mutational
871		exposures, endogenous processes, or repair defects, especially if some of
872		those exposures/processes/repair defects result in overwhelming mutational
873		spectra. Examples: melanoma / fair skin / UV exposure, POLE mutations,
874		MMR deficiency and APOBEC germ line variants.
875	o	The putative signature correlates with other clinical characteristics, such as
876		age at diagnosis (examples SBS1 and SBS5) or tobacco smoking (SBS4).
877	o	The mutational signature exhibits a strong transcriptional strand bias; it is
878		hard to imagine an artefact with transcriptional strand bias.
879	o	The putative signature shows association with other genomic features, such
880		as microindels in homopolymers, replication strand, replication timing, or
881		nucleosome occupancy.

Cancer sample sets on which different analyses have been conducted.

• Because PCAWG genomes are of high quality with respect to the calling of all
mutation types, all our analyses (all types of signature extraction and all types of
signature attribution) have been conducted on the 2,780 PCAWG genomes.

• SigProfiler also extracted SBS signatures from the non-PCAWG whole genomes,
TCGA exomes, and non-TCGA exomes and attributed SBS signatures to them.

• ID signatures have been extracted and attributed to PCAWG genomes and to a
subset of TCGA exomes with large numbers of indels (the latter SigProfiler only). We
have not done this for indels in non-PCAWG whole genome sequences and non-
TCGA exomes *(i)* because of the unknown and variable accuracy and standardisation
of indel mutation calls from different groups generating the data, *(ii)* because in
some cases no indel calls were provided by the data generator and *(iii)* because for
exomes in most cases there would be very few mutations.

• DBS signatures have been extracted and attributed to PCAWG genomes only. We
have not done this for the other categories of samples because of the unknown and
variable quality of the mutation calls, the possibility that filters introduced for quality
control might deliberately exclude doublet mutations, and the small numbers of
doublet mutations in exomes.

• Consistent with the above, composite mutational signatures have only been
extracted and attributed for PCAWG genomes.

Splitting of mutational signatures.

- Certain previously existing single signatures have split into multiple constituent signatures in this analysis. This is likely due to the existence of multiple, partially correlated mutational processes with the same initiating factor (for example, UV exposure) but subsequent differences in underlying mechanisms which differ in intensity in different tissues or other contexts. A previous example of this for which we have allocated different signature numbers is the split of the usually co-occurring but independently varying consequences of APOBEC mutagenesis into signatures SBS2 and SBS13 (https://cancer.sanger.ac.uk/cosmic/signatures/SBS/).
- Depending on the extent of correlation of the two signatures, and the available dataset/statistical power such signatures may manifest as a single signature, overlapping partially separated signatures or as two separate signatures.
- We are aware that splitting of signatures can also be a mathematical artefact. However, we have used multiple extractions to confirm and validate signature splits and applied the principle of sparsity to limit artefactual splits (https://cancer.sanger.ac.uk/cosmic/signatures/SBS/).

Better separation compared to COSMICv2 signatures

As described in the manuscript, all mutational signatures previously reported on COSMIC were confirmed in the new set of analyses with median cosine similarity of 0.95. However, the separation between the COSMICv2 mutational signatures (https://cancer.sanger.ac.uk/cosmic/signatures_v2) is much worse compared to the separation between the PCAWG mutational signatures. One can easily discern this by visual examination of signature profiles. For example, in COSMICv2, signatures 5 and 16 have a cosine similarity of 0.90, thus making them hard to distinguish from one another. In contrast, in the current PCAWG analysis, SBS5 and SBS16 have a cosine similarity of 0.65. This allows unambiguously assigning SBS5 and SBS16 to different samples. In the PCAWG analysis, the larger number of samples has allowed reducing the bleeding between signatures and has given more unique and easily distinguishable signatures. One can evaluate the overall separation of a set of mutational signatures by examining the distribution of cosine similarities between the signatures in the set. The COSMICv2 signatures have a median cosine similarity between the signatures in COSMICv2 of 0.238. In contrast, the PCAWG signatures have a much lower median cosine similarity between the signatures in PCAWG of 0.098. This 2-fold reduction in similarity is highly statistically significant (p-value: 9.1×10^{-25}) and indicates a better separation between the signatures in the current PCAWG analysis.

Correlations of mutational signature activity with age

Prior to evaluating the association between age and the activity of a mutational signatures, all outliers for both age and numbers of mutations attributed to a signature in a cancer type were removed from the data. Outlier was defined as any value outside three standard deviations from the mean value. A robust linear regression model that estimates the slope of the line and whether this slope is significantly different from zero (F-test; p-value<0.05) was performed using the MATLAB function robustfit (https://www.mathworks.com/help/stats/robustfit.html) with default parameters. The p-values yielded from the F-tests were corrected using the Benjamini-Hochberg procedure for

950 false discovery rate. Results are at https://www.synapse.org/#!Synapse:syn12030687 and
951 https://www.synapse.org/#!Synapse:syn20317940.
952
SigProfiler overview
SigProfiler incorporates two distinct steps for identification of mutational signatures based
on the previously described methodology[3,4,18,36]. The first step, SigProfilerExtraction,
encompasses a hierarchical *de novo* extraction of mutational signatures based on somatic
mutations and their immediate sequence context, while the second step,
SigProfilerAttribution, focuses on accurately estimating the number of somatic mutations
associated with each extracted mutational signature in each sample.
960
SigProfilerExtraction
(Note: This phase is termed SigProfiler in the MATLAB code and SigProfilerExtractor in
Python). The hierarchical *de novo* extraction approach is an extension of our previous
framework for analysis of mutational signatures (Extended Data Figure 8a)[3,18]. Briefly, for a
given set of mutational catalogues, the previously developed algorithm was hierarchically
applied to an input matrix $M \in \mathbb{R}_+^{K \times G}$ of non-negative integers with dimension $K \times G$,
where K is the number of mutation types and G is the number of samples. This previously
described algorithm deciphers a minimal set of mutational signatures that optimally
explains the proportion of each mutation type and estimates the contribution of each
signature to each sample. The algorithm uses multiple NMFs to identify the matrix of
mutational signatures, $P \in \mathbb{R}_+^{K \times N}$, and the matrix of the activities of these signatures, $E \in$
$\mathbb{R}_+^{N \times G}$, as previously described[3]. The unknown number of signatures, N, is determined by
human assessment of the stability and accuracy of solutions for a range of values for N, as
described[3]. The identification of M and P is done by minimizing the generalized Kullback-
Leibler divergence:
976

$$\min_{\substack{P \in \mathbb{R}_+^{(K,N)} \\ E \in \mathbb{R}_+^{(N,G)}}} \sum_{ij} (M_{ij} \log \frac{M_{ij}}{\widehat{M}_{ij}} - M_{ij} + \widehat{M}_{ij}),$$

977
where $\widehat{M} \in \mathbb{R}_+^{K \times G}$ is the unnormalized approximation of M, *i.e.*, $\widehat{M} = P \times E$. The
framework is applied hierarchically to increase its ability to find mutational signatures
generating few mutations or present in few samples. In detail, after application to a
matrix M containing the original samples, the accuracy of reconstructing the mutational
spectrum of each sample with the extracted mutational signatures is evaluated. Samples
that are well-reconstructed are removed, after which the framework is applied to the
remaining sub-matrix of M.
985
Transcriptional strand bias associated with mutational signatures was assessed by applying
SigProfilerExtraction to catalogues of in-transcript mutations that capture strand
information (192 mutations classes, https://www.synapse.org/#!Synapse:syn12026195).
These 192-class signatures were collapsed to strand-invariant 96-class signatures and
compared to the signatures extracted from the 96-class data, revealing very high cosine
similarities (median 0.90, column F in https://www.synapse.org/#!Synapse:syn12016215).
992

SigProfilerAttribution (single sample attribution)
(Note: This phase is termed SigProfilerSingleSample in both the MATLAB and Python code).
After signatures are discovered by SigProfilerExtraction, another procedure,
SigProfilerAttribution, estimates their contributions to individual samples. For each
examined sample, $C \in \mathbb{R}_+^{K \times 1}$, the estimation algorithm involves finding the minimum of the
Frobenius norm of a constrained function (see below for constraints) for a set of vectors
$S_{i=1..q} \in$ Q, where Q is a (not necessarily proper) subset of the set of mutational signatures,
P, ie, Q $\subseteq$ P.
1001

$$min \left\| \vec{C} - \sum_{r=1}^{q} (\overrightarrow{S_r} \times E_r) \right\|_F^2 \tag{1}$$

1002
In equation (1), $\vec{C}$ and each $\overrightarrow{S_r}$ are vectors of K nonnegative components reflecting,
respectively, the mutational spectrum of a sample and the *r-th* reference mutational
signature. All mutational signatures, $\overrightarrow{S_r}$, were identified in the SigProfilerExtraction step.
Each E_r is unknown scalar reflecting the number of mutations contributed by signature $\overrightarrow{S_r}$ in
the mutational spectrum $\vec{C}$. The minimization of equation (1) is always performed under
two additional constraints: (i) $E_r \geq 0$ and (ii) $\left\| \vec{C} \right\|_1 \geq E_r$; The constrained minimization of
equation (1) is performed using a nonlinear convex optimization programming solver using
the interior-point algorithm[73].
1011
1012   SigProfilerAttribution follows a multistep process, wherein equation (1) is minimized
1013   multiple times with additional constraints (Extended Data Figure 8b).
1014
In the first phase, the subset Q contains all signatures that were found by
SigProfilerExtraction in the same cancer type as the examined sample. Furthermore,
signatures violating biologically meaningful constraints based on transcriptional strand bias
and/or total number of somatic mutations are excluded from the set Q
(https://www.synapse.org/#!Synapse:syn12177009). Further, any $\overrightarrow{S_r} \times E_r$ for which the
cosine similarity between $\hat{C}$ and $\vec{C}$ is $\leq$ 0.01 are sequentially removed, where $\hat{C} =$
$\sum_{r=1}^{q} (\overrightarrow{S_r} \times E_r)$. Let T be the final set of signatures attributed to the sample at the end of
the first phase.
1023
In the second phase, equation (1) is minimized by sequentially allowing each signature,
$S_r \in$ P\Q,to be added provided that it increases the cosine similarity between $\hat{C}$ and $\vec{C}$
by >0.05. During this second phase, several additional biological conditions are enforced: *(i)*
signatures SBS1 and SBS5 are allowed in all samples, *(ii)* if one connected SBS signature is
found in a sample than another one is also allowed in the sample (e.g., if SBS17a is found in
a sample then SBS17b is allowed in the sample).
1030
1031
1032   *SignatureAnalyzer overview*
1033   SignatureAnalyzer employs a Bayesian variant of NMF that infers the number of signatures
1034   through the automatic relevance determination technique and delivers highly interpretable

1035 and sparse representations for both signature profiles and attributions that strike a balance
1036 between data fitting and model complexity. Please see references [6,15,39] for more details.
1037
SignatureAnalyzer signature extraction
In 2,780 PCAWG samples, we applied a two-step signature extraction strategy using 1536
penta-nucleotide contexts for SBSs, 83 ID features, and 78 DBS features. In addition to
separate extraction of SBS, ID, and DBS signatures, we performed a "COMPOSITE" signature
extraction based on all 1697 features (1536 SBS + 78 DBS + 83 ID). For SBSs, the 1536 SBS
COMPOSITE signatures are preferred, and for DBSs and IDs, the separately extracted
signatures are preferred.
In step 1 of the two-step extraction process, global signature extraction was performed for
the low mutation burden samples (n = 2,624). These excluded hyper-mutated tumours:
those with putative polymerase epsilon (POLE) defects or mismatch repair defects
(microsatellite instable tumours - MSI), skin tumours (which had intense UV mutagenesis),
and one tumour with temozolomide (TMZ) exposure. Because SignatureAnalyzer's
underlying algorithm performs a stochastic search, different runs can produce different
results. In step 1 we ran SignatureAnalyzer 10 times and selected the solution with the
highest posterior probability. In step 2, additional signatures unique to hyper-mutated
samples were extracted (again selecting the highest posterior probability over 10 runs),
while allowing all signatures found in the low mutation burden-samples to explain some of
the spectra of hyper-mutated samples. This approach was designed to minimize a well-
known "signature bleeding" effect or a bias of hyper- or ultra-mutated samples on the
signature extraction. In addition, this approach provided information about which
signatures are unique to the hyper-mutated samples which is later used when attributing
signatures to samples.
1060
SignatureAnalyzer signature attribution
A similar strategy was used for signature attribution; we performed a separate attribution
process for low- and hyper-mutated samples in all COMPOSITE, SBS, DBS, and ID signatures.
For downstream analyses, we preferred to use the COMPOSITE attributions for SBSs and the
separately calculated attributions for DBSs and IDs. Signature attribution in low-mutation
burden samples was performed separately in each tumour type (e.g., Biliary-AdenoCA,
Bladder-TCC, Bone-Osteosarc, etc.). Attribution was also performed separately in the
combined MSI (n=39), POLE (n=9), skin melanoma (n=107), and TMZ-exposed samples
(https://www.synapse.org/#!Synapse:syn11738314). In both groups, signature availability
(i.e., which signatures were active or not) was primarily inferred through the automatic
relevance determination process applied to the activity matrix H only, while fixing the
signature matrix, W. The attribution in low-mutation burden samples was performed using
only signatures found in the step 1 of the signature extraction. Two additional rules were
applied in SBS signature attribution to enforce biological plausibility and minimize a
signature bleeding: *(i)* allow signature SBS4 (smoking signature) only in lung and head and
neck cases; *(ii)* allow signature SBS11 (TMZ signature) in a single GBM sample. This was
enforced by introducing a binary, signature-by-sample, signature indicator matrix Z (1 -
allowed and 0 - not allowed), which was multiplied by the H matrix in every multiplication
update of H. No additional rules were applied to ID or DBS signature attributions, except
that signatures found in hyper-mutated samples were not allowed in low-mutation burden
samples.

1082
Tests on Synthetic Data
Our goal was to evaluate SignatureAnalyzer (SA) and SigProfiler (SP) on realistic synthetic
data. We operationally defined "realistic" as corresponding to either SA's or SP's analysis of
the PCAWG genome data. SA's reference signature profiles were based on "COMPOSITE"
signatures, consisting of 1536 strand-agnostic single base substitutions (SBSs) in
pentanucleotide context, 78 doublet base substitutions and 83 types of small insertions and
deletions, for a total of 1,697 mutation types. SP's reference analysis was based on strand-
agnostic single base substitutions in the context of one 5' and one 3' base; we term this
"SBS96" data. For each test, we generated two sets of "realistic" data: *SP-realistic*, based on
SP's reference signatures and attributions, and *SA-realistic*, based on SA's reference
signatures and attributions, as well as two other types of data that involved using SA profiles
with SP attributions and vice versa.
1095
1096 ***Generating synthetic data – overview.*** For tests (i) through (x) below, Synthetic data for
1097 sets of synthetic tumours of a given cancer type, *t*, were generated based on three
1098 parameters that were in turn based on the observed statistics for each signature, *s*, in
1099 cancer type *t*:
1100
1101 π, the proportion of tumours of cancer type *t* with signature *s*
1102
1103 μ, the mean of $\log_{10}$ of the number of *s* mutations across those tumours of type *t* that have
1104 signature *s*
1105
1106 σ, the standard deviation of $\log_{10}$ of the numbers of *s* mutations across those *t* tumours that
1107 have *s*
1108
To generate synthetic data,
(i) the proportion of tumours affected by *s* was drawn from the binomial distribution based
on π,
(ii) the number of mutations due to *s* in an affected tumour was drawn from a normal
distribution based on μ and σ.
The code used to generate the synthetic data and summarize SignatureAnalyzer and
SigProfiler results is open-source and freely available as the SynSig package:
https://github.com/steverozen/SynSig/tree/v0.2.0.
Description of each suite of synthetic data sets
1118
1119 ***i.* Synthetic pancreatic adenocarcinoma (1,000 spectra).**
1120 https://doi.org/10.7303/syn18500212.1
1121
***ii.* 2,700 synthetic whole-genome mutational spectra – 300 spectra from each of 9 cancer**
types. These spectra consist of 300 synthetic spectra from each of the following cancer
types: bladder transitional cell carcinoma, oesophageal adenocarcinoma, breast
adenocarcinoma, lung squamous cell carcinoma, renal cell carcinoma, ovarian
adenocarcinoma, osteosarcoma, cervical adenocarcinoma, and stomach adenocarcinoma.
https://doi.org/10.7303/syn18500213.1

1128

iii. **Mutational spectra generated from combinations of flat, relatively featureless**
mutational signatures -- version 1, 1000 synthetic tumours comprised of 500 synthetic
Kidney-RCCs (high prevalence and mutation load from SBS5 and SBS40 signatures) and 500
synthetic ovarian adenocarcinomas (high prevalence of and mutation load from SBS3). This
data set embodies tumours with high prevalence of the main flat signatures, SBS3, SBS5,
and SBS40, in a realistic context.
https://doi.org/10.7303/syn18500214.1

1136

iv. **Mutational spectra generated from combinations of flat, relatively featureless**
mutational signatures -- version 2, 1000 synthetic spectra all constructed entirely from
SBS3, SBS5, and SBS40, using mutational loads modelled on kidney-RCC (SBS5 and SBS40)
and ovarian adenocarcinoma (SBS3). Most synthetic spectra have contributions from all
three signatures.
https://doi.org/10.7303/syn18500215.1

1143

1144 *v.* **Mutational spectra generated from signatures with overlapping and potentially**
1145 **interfering profiles - version 1.** 500 synthetic bladder transitional cell carcinomas (high in
1146 SBS2 and SBS13) and 500 synthetic skin melanomas (high in SBS7a,b,c,d). The potential
1147 interference is between SBS2 (mainly C > T) and SBS7a,b (mainly C > T).
1148 https://doi.org/10.7303/syn18500217.1

1149

vi. **Mutational spectra generated from signatures with overlapping and potentially**
interfering profiles - version 2. 1000 synthetic tumours composed from SBS2 and
SBS7a,b. Mutational load distributions were drawn from bladder transitional cell carcinoma
(SBS2) and skin melanoma (SBS7a,b). Most spectra contain both signatures. The potential
interference is between SBS2 (mainly C > T) and SBS7a,b (mainly C > T).
https://doi.org/10.7303/syn18500216.1

1156

vii. Mutational spectra generated from combinations of signatures conferring high and
low mutation burdens. Based on 500 synthetic non-hypermutated tumours (parameters for
SBS1 and SBS5 estimated from colorectal and uterine adenocarcinomas) and 500 hyper-
mutated tumours (parameters for SBS26 and SBS44 estimated from hypermutated
colorectal and uterine adenocarcinomas). High and low mutation burden tumours are
segregated for SignatureAnalyzer (which analyses low mutation burden tumours first, then
high-burden tumours). SigProfiler analyses all tumours together.
https://doi.org/10.7303/syn18500218.1
https://doi.org/10.7303/syn18500219.1
https://doi.org/10.7303/syn18500216.1

1167

1168 **viii. A set of 30 random 96-feature mutational signature profiles and a set of 30 random**
1169 **1697-feature signature profiles (mimicking COMPOSITE signatures, which have 1697**
1170 **mutation types).** Each of these are used in two types of exposures, one with more (mean
1171 ~15.6) signatures per tumour and one with fewer (mean ~4) signatures per tumour.
1172 https://doi.org/10.7303/syn18500221.1

1173

1174 **ix. 2,700 whole-exome mutational spectra consisting of 300 synthetic spectra from each of**
1175 **9 different cancer types.** This test data set was generated from **test ii** by reducing the
1176 number of mutations of each type by 0.013 (approximately ratio of mutation counts
1177 between whole exome and whole genome mutational spectra).
1178 https://doi.org/10.7303/syn18909829.4
1179
1180 *Summary of findings*: Both SA and SP extracted substantially fewer signatures in this test
1181 than in **test ii**. In particular:
1182
SA: SA extracted only 18 signatures from the SA-realistic whole-exome data in this suite,
compared to the 40 signatures it extracted from the corresponding whole-genome synthetic
data in **test ii** and compared to the 39 ground-truth signatures in the synthetic spectra. The
average cosine similarity between ground-truth and extracted signatures for the synthetic
exome data was 0.863, compared to 0.968 for the signatures extracted from the whole-
genome spectra in **test ii**.
1189
1190 **SP**: SP extracted only 8 signatures from the SP-realistic whole-exome data in this suite,
1191 compared to the 19 it extracted from the whole-genome data in **test ii** and the 21 ground-
1192 truth signatures in the synthetic spectra. The average cosine similarity between ground-
1193 truth and extracted signatures for the synthetic exome data was 0.825, compared to 0.965
1194 for the signatures extracted from the whole-genome spectra in **test ii**.
1195
1196 **x. 1,350 synthetic whole-genome mutational spectra: 150 spectra from each of 9 cancer**
1197 **types.** This test data set consisted of every other tumour from **test ii**.
1198
Summary of findings: On the SA-realistic synthetic data, SA extracted fewer signatures in
this data set than in **test ii**, and in fact the number of signatures extracted was closer to the
ground truth and the cosine similarities were there higher. SA over-split in the
corresponding set of 2,700 tumours, and we speculate that SA's tendency to over-split
signatures is partly dependent on the number of input spectra, with the result that
extraction on 1,350 led to less over-splitting. SP extracted fewer signatures on this data set
than on **test ii**. In particular:
1206
1207 **SA**: SA extracted 38 signatures from the SA-realistic data in this suite, compared to the 40
1208 signatures it extracted from the 2,700 whole-genome spectra in **test ii** and compared to the
1209 39 ground-truth signatures. The average cosine similarity between ground-truth and
1210 extracted signatures for 1,350 genomes was 0.979 compared to 0.968 for the signatures
1211 extracted from the 2,700 whole-genome spectra in **test ii**.
1212
1213 **SP**: SP extracted 16 signatures from the SP-realistic data in this suite, compared to the 19
1214 signatures it extracted from the 2,700 whole-genome spectra in **test ii** and the 21 ground-
1215 truth signatures. The average cosine similarity between ground-truth and extracted
1216 signatures for the 1,350 spectra was 0.939 compared to 0.965 for the signatures extracted
1217 from the 2,700 spectra in **test ii**.
1218
1219 ***xi. Extraction of signatures from exome subsets of PCAWG mutational spectra.*** Our
1220 objective was to further test whether availability of mutations from whole-genome

1221 mutational spectra, as opposed to whole-exome spectra, enabled us to extract larger
1222 numbers of more accurate mutational signature profiles. In this test, we extracted
1223 signatures from mutational spectra that were based on only the exome regions of the actual
1224 PCAWG tumours (rather than on the purely synthetic data in **test ix**). The input data and
1225 extraction results are at https://doi.org/10.7303/syn18818766. We next summarize our
1226 findings for each of the SBS, DBS, and ID mutational signatures.

1227

1228 *xi-1 SBS signatures.* SignatureAnalyzer on COMPOSITE mutational classes (1536 SBS in
1229 pentanucleotide context plus DBS and ID) extracted 12 mutational signature profiles from
1230 the whole-exome data, none of which strongly resembled any of the 58 signatures it
1231 extracted from the whole-genome data. However, some signatures were unions or splits of
1232 the signatures extracted from the whole genome data. For example, WI was a union of the
1233 APOBEC signatures BI_COMPOSITE_SBS2_P and BI_COMPOSITE_SBS13_P. More broadly,
1234 somewhat recognizable SBS portions of the signatures were combined with the DBS and ID
1235 portions of the signatures in difficult-to-interpret combinations. We believe that SBS
1236 mutation counts were too low when spread across 1536 mutational classes to support
1237 robust mutational signature extraction.

1238

1239 SigProfiler on 96 SBS mutational classes extracted 17 mutational signature profiles from the
1240 exome data, compared to 48 that it extracted from the whole-genome data. The median
1241 cosine similarity of the exome-extracted signature profiles to the mutational signature
1242 profiles extracted from the whole genome data was 0.94. An outlier was SBS-E-2, which was
1243 a union of SBS2 and SBS13 (which tend to co-occur).

1244

1245 *xi-2 DBS signatures.* SignatureAnalyzer extracted 2 DBS signatures from the whole-exome
1246 data, compared to 15 DBS signatures that it extracted from the full whole genome data. One
1247 exome-extracted signature was essentially identical to BI_DBS1 (consisting almost entirely
1248 of CC > TT mutations), and one somewhat similar to BI_DBS2 (mostly CC > AA) but with
1249 many other mutational classes in addition.

1250

1251 SigProfiler extracted 3 DBS signatures from the whole-exome data, compared to the 11 DBS
1252 signatures that it extracted from the whole genome data. The exome-extracted signatures
1253 were good approximations of DBS1, DBS2, and DBS10 (cosine similarities 1, 0.93, and 0.98).

1254

1255 *xi-3 ID signatures.* SignatureAnalyzer extracted 4 ID signatures from the whole-exome data,
1256 compared to 29 ID signatures extracted from the whole-genome data. It extracted close
1257 approximations of BI_ID1_P and BI_ID2_P with cosine similarities 0.97 and 0.94. These are
1258 insertions (signature W.3) and deletions (signature W.1) of T:A in poly T:A.
1259 SignatureAnalyzer extracted 2 additional signatures. One of these (W.4) was a version of
1260 BI_ID4_P with several mutational classes absent. The other (W.2) appeared to be a union of
1261 many of the remaining ID signatures.

1262

1263 SigProfiler extracted 6 ID signatures from the whole-exome data, compared to the 17 ID
1264 signatures that it extracted from the whole genome data. Signatures ID-E-1, ID-E-2, ID-E-3,
1265 and ID-E-4 were good approximations of ID1, ID2, ID3, and ID4, respectively. An additional
1266 signature, ID-E-5, was approximately a union of ID6 and ID8. The remaining signature, ID-E-6
1267 was a partial version (deletions in C homopolymers only) of ID7.

1268

1269 ***Detailed Summary of Results (including links to input synthetic data sets and the signature***
1270 ***profiles extracted);*** https://doi.org/10.7303/syn18497223 provides a table with the number
1271 of signatures extracted by SigProfiler and SignatureAnalyzer for each synthetic data set and
1272 the cosine similarities to the input ground-truth signatures. See above for overall
1273 interpretation of the results.

1274

1275 ***Data Availability***
1276 Data are available at https://www.synapse.org/#!Synapse:syn11726601/wiki/513478. All
1277 figures and extended data figures have associated raw data.

1278

Code Availability
SigProfiler is available both as a MATLAB framework and as a Python package. In both cases,
SigProfiler is fully functional, free, and open-source tool distributed under the permissive 2-
Clause BSD License. SigProfiler in MATLAB can be downloaded from:
https://www.mathworks.com/matlabcentral/fileexchange/38724-sigprofiler
SigProfiler in Python can be downloaded from:
https://github.com/AlexandrovLab/SigProfilerExtractor. SignatureAnalyzer code is available at
https://www.synapse.org/#!Synapse:syn11801492. The code used to generate the synthetic data
and summarize SignatureAnalyzer and SigProfiler results is open-source and freely available as the
SynSig package: https://github.com/steverozen/SynSig/tree/v0.2.0 under the GPL3 license.
1289

1290 References
1291

1   Stratton, M. R., Campbell, P. J. & Futreal, P. A. The cancer genome. *Nature* **458**, 719-
724, doi:10.1038/nature07943 (2009).
2   Alexandrov, L. B. & Stratton, M. R. Mutational signatures: the patterns of somatic
mutations hidden in cancer genomes. *Current opinion in genetics & development* **24**,
52-60 (2014).
3   Alexandrov, L. B., Nik-Zainal, S., Wedge, D. C., Campbell, P. J. & Stratton, M. R.
Deciphering signatures of mutational processes operative in human cancer. *Cell Rep*
3, 246-259 (2013).
4   Alexandrov, L. B. *et al.* Clock-like mutational processes in human somatic cells. *Nat*
Genet **47**, 1402-1407, doi:10.1038/ng.3441 (2015).
5   Morganella, S. *et al.* The topography of mutational processes in breast cancer
genomes. *Nat Commun* **7**, 11383, doi:10.1038/ncomms11383 (2016).
6   Kim, J. *et al.* Somatic ERCC2 mutations are associated with a distinct genomic
signature in urothelial tumors. *Nat Genet* **48**, 600-606, doi:10.1038/ng.3557 (2016).
7   Fischer, A., Illingworth, C. J., Campbell, P. J. & Mustonen, V. EMu: probabilistic
inference of mutational processes and their localization in the cancer genome.
Genome Biol **14**, R39, doi:10.1186/gb-2013-14-4-r39 (2013).
8   Roberts, N. hdp (hierarchical Dirichelet process) R package.
https://github.com/nicolaroberts/hdp (2015).
9   Gehring, J. S., Fischer, B., Lawrence, M. & Huber, W. SomaticSignatures: inferring
mutational signatures from single-nucleotide variants. *Bioinformatics* **31**, 3673-3675
(2015).
10   Shiraishi, Y., Tremmel, G., Miyano, S. & Stephens, M. A simple model-based
approach to inferring and visualizing cancer mutation signatures. *PLoS genetics* **11**,
e1005657 (2015).
11   Rosales, R. A., Drummond, R. D., Valieris, R., Dias-Neto, E. & da Silva, I. T. signeR: an
empirical Bayesian approach to mutational signature discovery. *Bioinformatics* **33**, 8-
16, doi:10.1093/bioinformatics/btw572 (2017).
12   Ardin, M. *et al.* MutSpec: a Galaxy toolbox for streamlined analyses of somatic
mutation spectra in human and mouse cancer genomes. *BMC bioinformatics* **17**, 170
(2016).
13   Funnell, T. *et al.* Integrated single-nucleotide and structural variation signatures of
DNA-repair deficient human cancers. *bioRxiv*, doi:10.1101/267500 (2018).
14   Blokzijl, F., Janssen, R., van Boxtel, R. & Cuppen, E. MutationalPatterns:
comprehensive genome-wide analysis of mutational processes. *Genome medicine*
10, 33 (2018).
15   Kasar, S. *et al.* Whole-genome sequencing reveals activation-induced cytidine
deaminase signatures during indolent chronic lymphocytic leukaemia evolution. *Nat*
Commun **6**, 8866, doi:10.1038/ncomms9866 (2015).
16   Alexandrov, L. B. *et al.* Signatures of mutational processes in human cancer. *Nature*
500, 415-421 (2013).
17   Nik-Zainal, S. *et al.* Mutational processes molding the genomes of 21 breast cancers.
Cell **149**, 979-993 (2012).
18   Nik-Zainal, S. *et al.* Landscape of somatic mutations in 560 breast cancer whole-
genome sequences. *Nature* **534**, 47-54, doi:10.1038/nature17676 (2016).

19   Petljak, M. & Alexandrov, L. B. Understanding mutagenesis through delineation of
mutational signatures in human cancer. *Carcinogenesis* **37**, 531-540,
doi:10.1093/carcin/bgw055 (2016).
20   Schulze, K. *et al.* Exome sequencing of hepatocellular carcinomas identifies new
mutational signatures and potential therapeutic targets. *Nat Genet* **47**, 505-511,
doi:10.1038/ng.3252 (2015).
21   Poon, S. L. *et al.* Mutation signatures implicate aristolochic acid in bladder cancer
development. *Genome Med* **7**, 38, doi:10.1186/s13073-015-0161-3 (2015).
22   Poon, S. L. *et al.* Genome-wide mutational signatures of aristolochic acid and its
application as a screening tool. *Sci Transl Med* **5**, 197ra101,
doi:10.1126/scitranslmed.3006086 (2013).
23   Alexandrov, L. B. Understanding the origins of human cancer. *Science* **350**, 1175,
doi:10.1126/science.aad7363 (2015).
24   Hayward, N. K. *et al.* Whole-genome landscapes of major melanoma subtypes.
Nature **545**, 175-180, doi:10.1038/nature22071 (2017).
25   Polak, P. *et al.* A mutational signature reveals alterations underlying deficient
homologous recombination repair in breast cancer. *Nat Genet* **49**, 1476-1486,
doi:10.1038/ng.3934 (2017).
26   Merlevede, J. *et al.* Mutation allele burden remains unchanged in chronic
myelomonocytic leukaemia responding to hypomethylating agents. *Nat Commun* **7**,
10767, doi:10.1038/ncomms10767 (2016).
27   Mimaki, S. *et al.* Hypermutation and unique mutational signatures of occupational
cholangiocarcinoma in printing workers exposed to haloalkanes. *Carcinogenesis* **37**,
817-826, doi:10.1093/carcin/bgw066 (2016).
28   Drier, Y. *et al.* Somatic rearrangements across cancer reveal classes of samples with
distinct patterns of DNA breakage and rearrangement-induced hypermutability.
Genome Res **23**, 228-235, doi:10.1101/gr.141382.112 (2013).
29   Stephens, P. J. *et al.* Massive genomic rearrangement acquired in a single
catastrophic event during cancer development. *Cell* **144**, 27-40,
doi:10.1016/j.cell.2010.11.055 (2011).
30   Li, Y. *et al.* Patterns of structural variation in human cancer. *bioRxiv*,
doi:10.1101/181339 (2017).
31   Meier, B. *et al.* C. elegans whole-genome sequencing reveals mutational signatures
related to carcinogens and DNA repair deficiency. *Genome research* **24**, 1624-1636
(2014).
32   Supek, F. & Lehner, B. Clustered Mutation Signatures Reveal that Error-Prone DNA
Repair Targets Mutations to Active Genes. *Cell* **170**, 534-547 e523,
doi:10.1016/j.cell.2017.07.003 (2017).
33   Chen, J. M., Férec, C. & Cooper, D. N. Patterns and mutational signatures of tandem
base substitutions causing human inherited disease. *Human mutation* **34**, 1119-1130
(2013).
34   Campbell, P. J. *et al.* Pan-cancer analysis of whole genomes. *Preprint on bioRxiv,*
https://doi.org/10.1101/162784 (2017).
35   Yung, C. K. *et al.* Large-scale uniform analysis of cancer whole genomes in multiple
computing environments. *BioRxiv*, 161638 (2017).
36   WTSI Mutational Signature Framework,
http://www.mathworks.com/matlabcentral/fileexchange/38724 (2013).

37   Wellcome Trust Sanger Institute. *COSMIC, Catalog of Somatic Mutations in Cancer -*
Signatures of Mutational Processes in Human Cancer,
<http://cancer.sanger.ac.uk/cosmic/signatures> (2017).

38   Forbes, S. A. *et al.* The Catalogue of Somatic Mutations in Cancer (COSMIC). *Curr*
Protoc Hum Genet **Chapter 10**, Unit 10 11, doi:10.1002/0471142905.hg1011s57
(2008).

39   Tan, V. Y. & Févotte, C. Automatic relevance determination in nonnegative matrix
factorization with the/spl beta/-divergence. *IEEE Transactions on Pattern Analysis*
and Machine Intelligence **35**, 1592-1605 (2013).

40   Haradhvala, N. J. *et al.* Distinct mutational signatures characterize concurrent loss of
polymerase proofreading and mismatch repair. *Nature Communications* **9**, 1746,
doi:10.1038/s41467-018-04002-4 (2018).

41   Cichocki, A., Zdunek, R., Phan, A. H. & Amari, S.-i. *Nonnegative matrix and tensor*
factorizations: applications to exploratory multi-way data analysis and blind source
separation. (John Wiley & Sons, 2009).

42   Devarajan, K. Nonnegative matrix factorization: an analytical and interpretive tool in
computational biology. *PLoS computational biology* **4**, e1000029 (2008).

43   Blei, D., Carin, L. & Dunson, D. Probabilistic Topic Models: A focus on graphical model
design and applications to document and image analysis. *IEEE signal processing*
magazine **27**, 55 (2010).

44   Lawrence, M. S. *et al.* Mutational heterogeneity in cancer and the search for new
cancer-associated genes. *Nature* **499**, 214-218, doi:10.1038/nature12213 (2013).

45   Boot, A. *et al.* In-depth characterization of the cisplatin mutational signature in
human cell lines and in esophageal and liver tumors. *Genome Res* **28**, 654-665
doi:10.1101/gr.230219.117 (2018).

46   Viel, A. *et al.* A Specific Mutational Signature Associated with DNA 8-Oxoguanine
Persistence in MUTYH-defective Colorectal Cancer. *EBioMedicine* **20**, 39-49,
doi:10.1016/j.ebiom.2017.04.022 (2017).

47   Pilati, C. *et al.* Mutational signature analysis identifies MUTYH deficiency in colorectal
cancers and adrenocortical carcinomas. *J Pathol* **242**, 10-15, doi:10.1002/path.4880
(2017).

48   Drost, J. *et al.* Use of CRISPR-modified human stem cell organoids to study the origin
of mutational signatures in cancer. *Science* **358**, 234-238 (2017).

49   Saini, N. *et al.* The Impact of Environmental and Endogenous Damage on Somatic
Mutation Load in Human Skin Fibroblasts. *PLoS Genetics* **12**, e1006385,
doi:10.1371/journal.pgen.1006385 (2016).

50   Haradhvala, N. J. *et al.* Mutational Strand Asymmetries in Cancer Genomes Reveal
Mechanisms of DNA Damage and Repair. *Cell* **164**, 538-549,
doi:10.1016/j.cell.2015.12.050 (2016).

51   Chan, K. *et al.* An APOBEC3A hypermutation signature is distinguishable from the
signature of background mutagenesis by APOBEC3B in human cancers. *Nat Genet* **47**,
1067-1072, doi:10.1038/ng.3378 (2015).

52   Roberts, S. A. *et al.* Clustered mutations in yeast and in human cancers can arise
from damaged long single-strand DNA regions. *Molecular cell* **46**, 424-435 (2012).

53   Kasar, S. & Brown, J. R. Mutational landscape and underlying mutational processes in
chronic lymphocytic leukemia. *Mol Cell Oncol* **3**, e1157667,
doi:10.1080/23723556.2016.1157667 (2016).

54 Brash, D. E. UV Signature Mutations. *Photochemistry and Photobiology* **91**, 15-26,
doi:doi:10.1111/php.12377 (2015).
55 Hill, K. A., Wang, J., Farwell, K. D. & Sommer, S. S. Spontaneous tandem-base
mutations (TBM) show dramatic tissue, age, pattern and spectrum specificity. *Mutat*
Res **534**, 173-186 (2003).
56 Matsuda, T., Kawanishi, M., Yagi, T., Matsui, S. & Takebe, H. Specific tandem GG to
TT base substitutions induced by acetaldehyde are due to intra-strand crosslinks
between adjacent guanine bases. *Nucleic Acids Res* **26**, 1769-1774 (1998).
57 Garaycoechea, J. I. *et al.* Alcohol and endogenous aldehydes damage chromosomes
and mutate stem cells. *Nature* (2018).
58 Rahbari, R. *et al.* Timing, rates and spectra of human germline mutation. *Nat Genet*
48, 126-133, doi:10.1038/ng.3469 (2016).
59 Pfeifer, G. P. Formation and processing of UV photoproducts: effects of DNA
sequence and chromatin environment. *Photochemistry and photobiology* **65**, 270-
283 (1997).
60 Ceccaldi, R., Rondinelli, B. & D'Andrea, A. D. Repair pathway choices and
consequences at the double-strand break. *Trends in cell biology* **26**, 52-64 (2016).
61 Blokzijl, F. *et al.* Tissue-specific mutation accumulation in human adult stem cells
during life. *Nature* **538**, 260-264, doi:10.1038/nature19768 (2016).
62 Behjati, S. *et al.* Mutational signatures of ionizing radiation in second malignancies.
Nature communications **7**, 12605 (2016).
63 Huang, M. N. *et al.* Genome-Scale Mutational Signatures of Aflatoxin In Cells, Mice
And Human Tumors. *Genome Research* **27**, 1475-1486, doi:10.1101/gr.220038.116
(2017).
64 Nik-Zainal, S. *et al.* The genome as a record of environmental exposure. *Mutagenesis*
30, 763-770, doi:10.1093/mutage/gev073 (2015).
65 Olivier, M. *et al.* Modelling mutational landscapes of human cancers in vitro. *Sci Rep*
4, 4482, doi:10.1038/srep04482 (2014).
66 Szikriszt, B. *et al.* A comprehensive survey of the mutagenic impact of common
cancer cytotoxics. *Genome biology* **17**, 99 (2016).
67 Zhivagui, M. *et al.* Experimental analysis of exome-scale mutational signature of
glycidamide, the reactive metabolite of acrylamide. *bioRxiv*, 254664 (2018).
68 Zámborszky, J. *et al.* Loss of BRCA1 or BRCA2 markedly increases the rate of base
substitution mutagenesis and has distinct effects on genomic deletions. *Oncogene*
36, 746 (2017).
69 Zou, X. *et al.* Validating the concept of mutational signatures with isogenic cell
models. *Nature communications* **9**, 1744 (2018).
70 Roberts, N. D. *Patterns of somatic genome rearrangement in human cancer*,
University of Cambridge, (2018).
71 Rosales, R. A., Drummond, R. D., Valieris, R., Dias-Neto, E. & da Silva, I. T. signeR: an
empirical Bayesian approach to mutational signature discovery. *Bioinformatics* **33**, 8-
16 (2016).
72 Gori, K. & Baez-Ortega, A. sigfit: flexible Bayesian inference of mutational signatures.
bioRxiv, 372896 (2018).
73 Byrd, R. H., Hribar, M. E. & Nocedal, J. An interior point algorithm for large-scale
nonlinear programming. *SIAM Journal on Optimization* **9**, 877-900 (1999).

1477

Extended Data Figure and Table Legends

Extended Data Figure 1. Histogram of number of signatures attributed in each of 2,780 PCAWG samples by SigProfiler and SignatureAnalyzer. Hypermutated tumours and melanomas (156) are listed at https://www.synapse.org/#!Synapse:syn11738314.

Extended Data Figure 2. Comparisons between SigProfiler and SignatureAnalyzer results. Comparison of the attributions for corresponding SigProfiler *(a)* and SignatureAnalyzer *(b)* signatures. Each of the SBS signatures extracted by SigProfiler and SignatureAnalyzer was paired with the signature of highest cosine similarity in the extraction by the other method (if one with >0.85 cosine similarity exists). The first column of the plot corresponds to the fraction of mutations assigned by one method (summed across samples and mutation types) that were also assigned by the other method. The remaining mutations were then re-distributed to the other signatures in the extraction, weighted by their relative probabilities of having been generated by each signature, and the resulting fraction of mutations is plotted. Signatures on the x-axis are only shown if they contribute at least 0.1 fraction of mutations to at least one signature on the y-axis. Cosine similarities between SigProfiler and SignatureAnalyzer DBS *(c)* and ID *(d)* signatures. Brown nodes represent SigProfiler signatures; green nodes represent SignatureAnalyzer signatures. Matches with cosine similarities > 0.8 are show as edges, with the width of the edge indicate the strength of the similarity. The locations of the nodes have no significance. Signatures with no matches of > 0.8 cosine similarity are show below. Note that SigProfiler ID15 and ID17 were extracted from data that were not analysed by SignatureAnalyzer. Suffixes 'P' and 'S' on SignatureAnalyzer signature names indicate (1) signatures extracted from non-hypermutated, non-melanoma tumours and (2) hypermutated and melanoma tumours, respectively.

Extended Data Figure 3. SignatureAnalyzer reference signatures. See legend of main text Figure 2.

Extended Data Figure 4. The number of SBS mutations attributed to each mutational signature for each cancer type over the 2,780 PCAWG tumours by SignatureAnalyzer. See main text Figure 3 for explanation.

Extended Data Figure 5. The number of SBS mutations attributed to each mutational signature to each cancer type over the complete set of 23,829 cancer samples analysed by SigProfiler. See main text Figure 3 for explanation.

Extended Data Figure 6. Associations of between SBS, DBS, and ID signature activities for SigProfiler *(a)* and SignatureAnalyzer *(b)*. Each node represents an SBS (light green), DBS (dark green) or ID (black) signature. Any two signatures with sample attributions that significantly correlated with R^2 > 0.3 (SigProfiler) or > 0.5 (SignatureAnalyzer) are connected by edges. Edge widths are proportional to the strength of the correlation. Signatures with no significant correlation to any other signature above the relevant threshold are not shown. Signature locations are fit for display purposes only and do not indicate similarity.

Extended Data Figure 7. Mutational signatures extracted from the composite feature set consisting of SBSs in pentanucleotide context, DBSs, and IDs. For each of the four composite mutational signatures shown, the top panel is the SBS signature collapsed to 96 SBS classes, the middle panel is the co-extracted DBS signature, and the lower panel is the co-extracted ID signature. Note the similarities between the DBS portion of Composite 4 and DBS2, between the ID portion of Composite 4 and ID3, and other similarities noted in the figure.

Extended Data Figure 8. SigProfiler signature extraction *(a)* **and attribution** *(b).* See Methods for description.

Extended Data Table 1. The number of DBSs is proportional to the number of SBSs with the exception of a few cancer types (ColoRect-AdenoCA, Lung-AdenoCA, Lung-SCC, Skin-Melanoma) analysed by the following linear regression (computed by an R function call):
$$glm(DBS.counts \sim SBS.counts + Cancer.Types).$$

Extended Data Table 2. Numbers of insertion/deletion mutations due to ID1, ID2, and all other ID signatures in hypermutators and non-hypermutators.

Fig1

Single Base Substitution

Fig 2 part 2

SBS1 SBS2 SBS3 SBS4
SBS5 SBS6 SBS7a SBS7b
SBS7c SBS7d SBS8 SBS9
SBS10a SBS10b SBS11 SBS12
SBS13 SBS14 SBS15 SBS16
SBS17a SBS17b SBS18 SBS19
SBS20 SBS21 SBS22 SBS23
SBS24 SBS25 SBS26 SBS28
SBS29 SBS30 SBS31 SBS32
SBS33 SBS34 SBS35 SBS36
SBS37 SBS38 SBS39 SBS40
SBS41 SBS42 SBS44 SBS84
SBS85

Doublet Base Substitution

DBS1 DBS2 DBS3 DBS4
DBS5 DBS6 DBS7 DBS8
DBS9 DBS10 DBS11

Insertion and Deletion

ID1 ID2 ID3 ID4
ID5 ID6 ID7 ID8
ID9 ID10 ID11 ID12
ID13 ID14 ID15 ID16
ID17

Fig 3

Fig 4

Extended Data Fig 1

SignatureAnalyzer reference SBS signatures

Extended data Fig 3

SignatureAnalyzer reference DBS signatures

SignatureAnalyzer reference ID signatures

Extended Data Fig 4

Extended Data Fig 5

a **SigProfiler**

b **SignatureAnalyzer** Extended Data Fig 6

a Extraction of mutational signatures

Step A (Apply the approach to a set of samples D; initially D contains all samples, i.e., D=M)
Described in detail in (Alexandrov et al., Cell Rep. 2013;3(1):246-59).

Repeat $N = 1 ... (G − 1)$

Step B (Solution evaluation and re-iteration)
Extracted mutational signatures and their activities to individual samples are saved into a set *S*. The activity of any signature that does not increase the cosine similarity of a sample with more 0.01 was removed from the sample (i.e., assigned a value of zero). ***Step A*** is repeated for all samples for which the identified signatures do not explain their patterns (cosine similarity <0.95). The algorithm continues to the ***step C*** when ***step A*** cannot find any stable signatures.

Step C (Clustering of mutational signatures)
Hierarchical consensus clustering was applied to the set S to derive the consensus mutational signatures across the set of samples *M.*

b Attribution of activities of mutational signatures in samples

Complete set of mutational signatures

S1: Examine an individual sample

S2: Consider signatures extracted from this tumour type

S3: Apply rules for mutational signatures

Signature rules (only SBSs)
1) Exclude signatures if they lack the appropriate statistically significant strand bias: applied to signatures 4, 8, 7a/b/c/d, 11, 12, 16, 22, 23, 24, 25, 27, 31, 32, 33, 35, 42
2) Exclude 10a/b if sample has < 10^5 SBSs in WGS
3) Exclude 6, 14,15, 20, 21, 26 if sample has < 10^4 SBSs in WGS

1) Remove least contributing signature if removal reduces cosine similarity < 0.01
2) "Connected signature inclusion rules" for SBSs; see Methods

S4: Evaluate sample with all **N** remaining signatures

S5: Exclude each of the **N** signatures and evaluate the sample with **N-1** signatures

S6: Evaluate sample with the remaining **M** signatures

1) For SBS, add SBS1 and SBS5 if addition improves cosine similarity
2) For DBS, ID, and SBS signatures other than 1 or 5, add most contributing signature if addition increases cosine similarity > 0.05
3) "Connected signature inclusion rules" for SBSs; see Methods

S8: Include each of the remaining signatures from the global set of signatures and evaluate the sample with **M+1** signatures

S7: Consider all other signatures

S9: Output the results

Extended Data Table 1. The number of DBSs is proportional to the number of SBSs with the exception of a few cancer types (ColoRect-AdenoCA, Lung-AdenoCA, Lung-SCC, Skin-Melanoma), analysed by the following linear regression (computed by an R function call):   glm(DBS.counts ~ SBS.counts + Cancer.Types)

	Estimate	Std.Error	t value	Pr(>|t|)	
(Intercept)	5.61E+00	8.76E+01	0.064	0.9489	
SBS.counts	3.74E-03	1.25E-04	29.841	<2.00E-16	***
Bladder-TCC	1.32E+01	1.39E+02	0.095	0.92432	
Bone-Osteosarc	2.18E+00	1.21E+02	0.018	0.98567	
Bone-Other	-2.81E+00	1.33E+02	-0.021	0.9831	
Breast	5.32E+00	9.44E+01	0.056	0.95511	
Cervix	-1.06E+01	1.45E+02	-0.073	0.94185	
CNS-GBM	-2.81E+01	1.19E+02	-0.236	0.81352	
CNS-Medullo	-7.04E+00	9.75E+01	-0.072	0.94239	
CNS-Oligo	-1.03E+01	1.50E+02	-0.069	0.94539	
CNS-PiloAstro	-5.87E+00	1.03E+02	-0.057	0.95467	
ColoRect-AdenoCA	-4.11E+02	1.12E+02	-3.667	0.00025	***
Eso-AdenoCA	-1.56E+01	1.02E+02	-0.153	0.87838	
Head-SCC	5.27E+01	1.11E+02	0.474	0.63541	
Kidney-ChRCC	-3.14E+00	1.17E+02	-0.027	0.97857	
Kidney-RCC	5.61E+01	9.76E+01	0.574	0.56584	
Liver-HCC	7.82E+01	9.21E+01	0.849	0.39575	
Lung-AdenoCA	5.02E+02	1.21E+02	4.136	3.63E-05	***
Lung-SCC	5.85E+02	1.15E+02	5.078	4.08E-07	***
Lymph-BNHL	1.04E+01	1.01E+02	0.103	0.91765	
Lymph-CLL	-4.30E+00	1.02E+02	-0.042	0.96655	
Myeloid-AML	-1.89E+00	1.79E+02	-0.011	0.99156	
Myeloid-MDS/MPN	-7.43E+00	1.10E+02	-0.067	0.94622	
Ovary-AdenoCA	3.59E+01	1.00E+02	0.358	0.72023	
Panc-AdenoCA	-8.34E-01	9.37E+01	-0.009	0.99289	
Panc-Endocrine	-5.70E+00	1.04E+02	-0.055	0.95628	
Prost-AdenoCA	2.52E+00	9.27E+01	0.027	0.97831	
Skin-Melanoma	1.67E+03	1.02E+02	16.47	<2.00E-16	***
SoftTissue-Leiomyo	5.98E+00	1.60E+02	0.037	0.97016	
SoftTissue-Liposarc	7.77E+00	1.48E+02	0.053	0.95804	
Stomach-AdenoCA	-3.04E+01	1.06E+02	-0.287	0.77417	
Thy-AdenoCA	-4.80E+00	1.15E+02	-0.042	0.96676	
Uterus-AdenoCA	-1.25E+02	1.14E+02	-1.096	0.27304	

Extended Data Table 2. Numbers of insertion/deletion mutations due to ID1, ID2, and all other ID signatures combined, in hypermutators and non-hypermutators

Signature	Hypermutators		Non-hypermutators		All Tumours	
	Count	Fraction	Count	Fraction	Count	Fraction
ID1	593,935	0.236	399,633	0.276	993,568	0.250
ID2	1,838,867	0.730	252,893	0.174	2,091,760	0.527
ID1+ID2	2,432,802	0.966	652,526	0.450	3,085,328	0.777
Other ID signatures	85,038	0.034	797,964	0.550	883,002	0.223
Total	2,517,840	1	1,450,490	1	3,968,330	1