Learning the sequence of influenza A genome assembly during viral replication using point

process models and fluorescence in situ hybridization

Timothy D. Majarian,[1,2*] Robert F. Murphy,[1-4#] and Seema S. Lakdawala[5#]

[1]Computational Biology Department, [2]Department of Biological Sciences, [3]Department of

Biomedical Engineering, [4]Machine Learning Department, Carnegie Mellon University

[5]Department of Microbiology and Molecular Genetics , University of Pittsburgh School of

Medicine

*Present address: Broad Institute of MIT and Harvard

#Corresponding authors: murphy@cmu.edu, Lakdawala@pitt.edu

Abstract

Within influenza virus infected cells, viral genomic RNA are selectively packed into progeny

virions, which predominantly contain a single copy of 8 viral RNA segments. Intersegmental

RNA-RNA interactions are thought to mediate selective packaging of each viral

ribonucleoprotein complex (vRNP). Clear evidence of a specific interaction network culminating

in the full genomic set has yet to be identified. Using multi-color fluorescence *in situ*

hybridization to visualize four vRNP segments within a single cell, we developed image-based

models of vRNP-vRNP spatial dependence. These models were used to construct likely

sequences of vRNP associations resulting in the full genomic set. Our results support the notion

that selective packaging occurs during cytoplasmic transport and identifies the formation of

multiple distinct vRNP sub-complexes that likely form as intermediate steps toward full

genomic inclusion into a progeny virion. The methods employed demonstrate a statistically

driven, model based approach applicable to other interaction and assembly problems.

Author Summary

Influenza virus consists of eight viral ribonucleoproteins (vRNPs) that are assembled by infected

cells to produce new virions. The process by which all eight vRNPs are assembled is not yet

understood. We therefore used images from a previous study in which up to four vRNPs had

been visualized in the same cell to construct spatial point process models that measure how

well the subcellular distribution of one vRNP can be predicted from one or more other vRNPs.

We used the likelihood of these models as an estimate of the extent of association between

vRNPs and thereby constructed likely sequences of vRNP assembly that would produce full

virions. Our work identifies the formation of multiple distinct vRNP sub-complexes that likely

form as intermediate steps toward production of a virion. The results may be of use in

designing strategies to interfere with virus assembly. We also anticipate that the approach may

be useful for studying other assembly processes, especially for complexes with modest affinities

and more components than can be visualized simultaneously.

Introduction

Influenza A virus, part of the *orthomyxoviridae* family, has a segmented genome of eight distinct viral RNA segments coding at least 11 major proteins and several auxiliary peptides. Most notable of the 11 viral proteins are hemagglutinin (HA) and neurominidase (NA), the canonical *H* and *N* in influenza strain designation. A segmented genome offers potential evolutionary advantages during viral replication, in the form of reassortment. The exchange of genomic material between two distinct viral strains in a co-infected cell often causes a shift within the genome, greatly increasing viral genetic diversity. In only the last century, influenza pandemics of 1957 (H2N2), 1968 (H3N2), and 2009 (H1N1) have emerged through reassortment of at least two viruses [1]. Segmented genomes also come with an inherent mechanistic challenge: ensuring that progeny receive a complete set of genomic segments. Random packaging is possible but comes at the cost of producing an overwhelming majority of progeny that are not viable [2]. Evidence suggests that there exists an active mechanism of selective packaging ensuring progeny viability through efficient and accurate genomic packaging [3, 4].

Within a virion, the viral genome is organized into individual viral ribonucleoprotein (vRNP) complexes, composed of the viral RNA, virally encoded nucleoprotein (NP), and a heterotrimeric polymerase complex made up of PB1, PB2 and PA. The vRNP structure is known classically as a helical panhandle, where the RNA is wrapped around NP with RNA bases exposed and the 3' and 5' ends associated with the polymerase complex [3]. We have recently demonstrated that the structure of vRNP is more complex than the classically depicted "beads-

on-a-string" schematic. The NP and viral RNA associate in a non-uniform manner, where regions

of the RNA are unbound by NP and capable of forming complex structures [5]. Macro-

organization of vRNPs within the viral capsid show tight packaging, organized in a '7+1'

orientation, a single vRNP center shaft surrounded by seven others (4-6).

Given the need to package one copy of all eight vRNP segments, there is strong evidence for

the existence of a selective packaging mechanism prior to budding [3, 6-9]. Intersegmental

RNA-RNA interactions have been proposed to mediate selective packaging of all eight influenza

vRNP segments. The 5' and 3' regions of each segment have been implicated to be essential for

efficient packaging and may be critical for the proposed RNA-RNA interactions [3, 10-13].

Previous studies using *in vitro* transcribed viral RNA have successfully demonstrated RNA-RNA

binding. These studies also show that a set of eight reproducible interactions can suffice to

form a full genome complex [14-16]. However, these studies were performed on viral RNA in

the absence of NP which would alter the regions of viral RNA accessible for the proposed RNA-

RNA interactions. Recent *in vivo* experiments have also pointed towards RNA-RNA interaction

as the driving mechanism for selective packaging [17]. While there is consensus of its existence,

a clear network of segment interactions leading to a full genome complex has yet to be

elucidated.

We and others have previously used fluorescence *in situ* hybridization (FISH) studies to visualize

the intracellular localization of multiple vRNP segments during a productive viral infection [18,

19]. These studies, performed eight hours post infection to capture an initial infection cycle and

reduce complication of cytoplasmic vRNP from a subsequent infection, have shown that distinct

vRNP segments colocalize within the cytoplasm. Based on our previous studies, we proposed a

model whereby vRNP segments are exported from the nucleus, the site of vRNP synthesis, as

subcomplexes that undergo further assembly en route to the plasma membrane through

dynamic fusion or colocalization events [19]. We also performed a series of binary

colocalization comparisons, which would postulate a simple linear vRNP interaction network.

However, no such network was identified, suggesting the presence of a more complex vRNP

interaction network, including higher order subcomplexes as intermediate steps [19].

The lack of a method to image all eight vRNP segments within a single infected cell has limited

our ability to examine the precise spatial relationship between segments. In this study, we

developed a model based approach, rooted in point process theory, to quantify vRNP spatial

dependence as a metric for RNA-RNA interaction using multi-color FISH images. A spatial point

process model is a statistical model that captures the probability that an event will happen at

any position within a given geometry. The probability can depend on the spatial distributions

of other events or objects, often referred to as covariates or factors. In this context, an event is

the observance of fluorescent signal (corresponding to one or more vRNPs) at a given location.

We have previously used point process modeling to construct generative models reflecting the

spatial dependencies between various punctate cell organelles and other cellular structures,

such as the cell and nuclear membranes, microtubules and the endoplasmic reticulum [20, 21].

Related analytical methods have been used to analyze clustering of molecules reflecting spatial

dependencies between copies of the same structure [22, 23] Here we use point process models

to present a statistically rigorous analysis of intracellular influenza A genome assembly

dynamics from a spatial perspective during a productive viral infection.

For an imaged cell, the locations of each vRNP segment can be represented as "realizations" of

an underlying probability density specific to that vRNP, calculated over the cytoplasm, and

dependent on the spatial proximity to various other structures within the cell. That is, at any

given position within the cell, there exists a probability of observing a single segment, of a

particular identity, that depends on the positions of cellular organelles and other vRNP

segments.   By modeling the locational densities of each vRNP, we can therefore learn a

dependency network from which likely vRNP interactions are implied.

FISH imaging yields many distinct observations of individual cells, each with multiple vRNP

patterns observed. These point patterns, when viewed as realizations of point processes, allow

for statistical learning of spatial dependencies over many replicates for each vRNP segment. By

defining a set of covariates as the minimum distances between a given segment and other

segments observed within the same cell, we produce a set of models that describe the spatial

relationship between distinct segments. The extent to which two (or more) vRNPs are

dependent upon each other (e.g., likely to be found at nearby positions) is reflected in the

likelihood that images of those vRNPs would have been produced by a learned model: high

likelihoods signify that the positions of a given vRNP can be predicted well from the others in

that model. Model likelihoods can then be seen as a metric of vRNP interaction, with higher

likelihoods indicating more probable vRNP association. These likelihoods can then be used to

construct the most probable sequence of interactions to form the full genome with methods

borrowed from evolutionary tree construction [24].

Results:

Imaging influenza virus infected cells

To investigate the spatial dependence between vRNP segments, we utilized images from a

previous study of infected cells stained for different combinations of vRNP segments using four-

color FISH at eight hours post infection [19]. The data set included 14 different probe

combinations that covered all pairwise vRNP associations and 32 out of 56 possible triple vRNP

combinations (Supplemental Table 1 and 2). Prior to initiating pattern analysis, each image was

processed to remove background, isolate individual cells, and find punctate vRNP segments.

Figure 1a and b presents representative raw and segmented images with detected points.

Assessing randomness of the spatial distributions of vRNP

We began by determining the extent to which the individual vRNPs could be considered

randomly distributed throughout the cell, since our search for spatial dependencies between

different vRNPs would be illogical if their positions were all random. We defined the null

hypothesis as complete spatial randomness, as would be exhibited by a homogenous Poisson

process, and estimated a p-value for this hypothesis using Monte Carlo methods. Briefly, for a

given vRNP in a particular cell, we started by segmenting the cytoplasm and counting the

number of points observed within. We then simulated (generated) many random point

configurations, of the same number of points, within the segmented cytoplasm. We calculated

a test statistic for each random configuration (see Methods) and used these to assign a p-value

to the test statistic for the observed pattern for that cell; these were averaged for each vRNP.

As shown in Table 1, every vRNP segment showed significant deviation (at the $p < 0.05$ level),

with low variance, from a spatially random distribution. This indicates the presence of a spatial

trend and some dependency on cellular structures for their location.

Modeling cell and nuclear membrane dependence

Given that the localizations of vRNPs are not random, we sought to determine if some

relationship to the cell and nuclear membranes could explain the spatial trend of vRNP

segments. Three covariates were defined for each position in the cell (minimum distance to the

cell membrane, minimum distance to the nuclear membrane, and the minimum ratio of cell to

nuclear distance for each point) and models were created for each vRNP using each of the

covariates separately. We estimated the accuracy of each model by cross validated likelihood

calculation (see Methods), where values closer to 0 are more likely. It is important to note that,

because actual likelihoods were calculated, they can be compared between models of different

complexity.

Dependency on the cell membrane produced the lowest likelihood (worst) models relative to all

others (Table 2) for all vRNP segments. Congregation of vRNP to the apical cell membrane is a

potential skewing factor, since there will be a high concentration of all segments at this cellular

location. The many observed points within close proximity to the membrane potentially caused

low fitted probability of observing other points closer to the nucleus. Cell shape differences

combined with lower resolution in the z-dimension may have also influenced these models.

For most vRNPs, models incorporating nuclear distance showed higher likelihoods than those

using cell membrane distance (Table 2). Some vRNP segments, such as PB2, PB1, PA and NS,

showed greater dependency on nuclear distance than the other segments. The ratio of cell

membrane to nuclear distance generally produced models better than cell membrane alone,

but they were worse than nuclear alone in all cases except NA (Table 2). Thus nuclear distance

provides the most cogent model for dependence of vRNP localization on a cellular structure.

Inter-segment dependence in pairs, triplets, and quadruplets

Every pair of vRNP segments was observed in at least three cells, allowing for modeling of

vRNP-vRNP dependence among all segment pairs. For each pair of vRNP segments, two models

were trained, switching the dependent (primary) and independent (secondary) patterns. Since

the minimum inter-pattern distance measure is not commutative, the two models need not be

the same. More simply, a given vRNP location may depend on its proximity to another vRNP,

but the converse is not necessarily true. This can be observed when one vRNP segment is

found in two distinct locations, and a second vRNP segment is found in only one of those

locations. The second vRNP segment would be observed to have a high likelihood for being

predicted from the first, but the first would not have a high likelihood of being predicted from

the second.

In all cases, pairwise inter-pattern distance models (Table 3) for a given vRNP had higher

likelihoods than those produced for that vRNP with only the cell and/or nuclear membrane

factors (Table 2). Some pairs showed higher likelihoods than others, such as PB1-NP, PB1-M,

PA-NP, HA-PB1, and NP-PB1. These pairings also showed relatively low variance in cross-

validated likelihood estimates (Supplemental Table 3). Many of the highest likelihood vRNP

pairs are observed between segments encoding the viral polymerase components (PB2, PB1, PA

and NP). Given the dependence between these proteins both structurally and functionally [25-

27], it is possible that these segments evolved a dependence upon each other to ensure their

joint packaging in a virion (a possibility that would require extensive additional work to

examine). Figure 2 highlights the pairwise models in Table 4 by illustrating vRNP pairs that are

below 0.23 log likelihood. Overall, vRNPs showed the greatest dependence on PA and PB1, with

the most connections between these two segments and the others. The least dependence was

observed for NA and NS vRNP segments.

Extending the pairwise models, nuclear distance was added as another covariate. Minor

improvements in model likelihood were seen in some models (Table 4) with others showing

decreased likelihood. Overall there was an increase in the variance of our estimates

(Supplemental Table 4). Overfitting to nuclear distance may be the cause of this increased

variance.

Thirty-two of 56 possible triplet vRNP segment complexes (referred to as triplets) were present

within the dataset analyzed and could therefore be modeled within the same cell. This process

allows for modeling of complex vRNP relations containing more than two vRNP segments.

Three models were trained for each observed triplet, one for each vRNP depending on the

other two. The highest likelihood model was chosen. Model likelihoods tended to be higher

than that of either pair model for a given vRNP (Table 5). We observed that the most likely

triplet models for each vRNP included four of the five most likely vRNP pairings observed in pair

likelihood models (Table 3), (namely PB1-NP, PA-NP, HA-PB1, and NP-PB1). However, not all of

the most likely pairwise associations (e.g., PB1-M) were observed in the most likely triplet

models, highlighting that models based solely on pairwise comparisons might not accurately

represent larger order vRNP complexes. Model likelihoods for all triplets observed can be found

in Supplemental Table 5.

A small subset of all possible four vRNP segment complexes (referred to as quadruplets) were

contained within the dataset and models for predicting one of the four from the other three

were also constructed. Again, considerable overlap is seen between the most likely quadruplet

models and both triplet and pair models (Table 6), such as PB2-PB1-PA and PB1-PA. The

consistency in vRNP composition between observed likely quadruplet, triples, and pairs further

validates the spatial dependence of vRNP segments and the presence of vRNP subcomplexes as

influenza A assembly intermediates. Model likelihoods for all quadruplets observed can be

found in Supplemental Table 6.

One of our major goals was to attempt to extrapolate higher order relationships from the pair,

triple and quadruplet models. To project how accurately this might be done, we compared our

observed triple and quadruple models with predicted models calculated from any of the four

pair models (vRNP pairs (Table 3), vRNP pairs + nucleus (Table 4), vRNP pairs +cellular

membrane, and vRNP pairs +nuclear/cell membrane ratio) (Figure 3a). We also compared

observed quadruplet likelihoods to those predicted using both the pair and triple models,

similar to the pair models any of the 4 models including the various cellular features were

included (Figure 3b). Most of our observed triples and quadruplets could be predicted with

modest accuracy from the pair models but the overall r^2 value was only 0.302. This was due to

inaccuracy in predicting the triple and quadruple clusters containing the PB2-NA pair with

either NS, PB1 or both. These combinations were predicted to occur much more often than

was observed in the experimental data sets. The pairwise likelihood of NA with PB2 is >-0.6,

which is consistent with the observed likelihood and may be the driving force behind the score

compared to the other pairwise interactions. Analysis of the predicted triples and quadruples

after removing the PB2-NA containing clusters resulted in a r^2 value increases to 0.642 (we

cannot justify this removal on a statistical basis but rather on the observation that the

combinations with the poorest predictions all involved the same pair). Predicting quadruples

from pairs and triples resulted in a r^2 value of 0.450, and with removal of the same quadruple it

rose to 0.714. Overall, these data support the notion that pairwise point process models can be

used to predict higher order complexes with moderate accuracy, and that prediction accuracy

increases if higher order models are used.

Interaction network construction

Given these results, we sought to create a model of the set of vRNP interactions (complexes)

that occur at all steps in the formation of a complete viral genome. To do this, we represented

all possible interactions between vRNP segments as a weighted, directed, acyclic graph with

nodes labeled by each unique, unordered set of vRNP segments. A path from all single nodes to

the root represents a set of vRNP interactions resulting in a full genome complex (see

Methods). For illustration, if all eight vRNP segments directly formed a complete set without

forming any intermediate complexes (a highly unlikely scenario), this would be represented by

weights of one for the connection between each vRNP and the complete vRNP supramolecular

complex containing all eight segments and weights of zero for all other complexes. Such a

scenario would result in a tree with the root connected to each of the leaves

In our analysis, we estimated the weight for each possible intermediate vRNP complex

assembly subcomplex to contain at least two and up to eight segments based upon either the

observed or predicted likelihoods (see Methods). Dynamic programming, an efficient method

for recursively finding the highest scoring path to a given node, was then used to find the most

probable path within the graph using the highest likelihood trained model for a given nodes

from those with or without cell and/or nuclear features (Figure 4a). The resultant tree produces

two distinct clusters HA, M, NA, NS (hereafter referred to as C1) and PB1, PB2, NP (C2) that

merge with PA as a final step towards forming the full genome. This network suggests three

steps to form a complete set of all eight vRNP segments: 1) formation of C1, 2) formation of C2

and 3) addition of PA to C1 and C2 to form the final product.

To assess whether a particular vRNP localization dataset was driving this network, we

performed a series of network analyses with data excluded. First, the contribution of observed

quadruplets and triplets was assessed by excluding observed quadruplets (Figure 4b) and

without observed quadruplets and triplets (Figure 4c). Interestingly, exclusion of all observed

quadruplets did not drastically alter the network, with both C1 and C2 still being present,

although formation of C1 required two steps. However, networks built only with observed pairs

(excluding triplets and quadruplets) resulted in a quite different network lacking larger

subcomplexes (Figure 4c), reflecting the fact that pairs were only able to make approximate

estimates of higher order interactions

Given that the dynamic programming approach yields only a single most likely tree, we sought

to assess the robustness of the inferred tree to potential inaccuracy in the estimated

dependencies. We therefore assessed the stability of the tree by adding various amounts of

random noise to the estimated likelihoods (see Methods). Surprisingly, the addition of noise of

up to a quarter of the mean log-likelihood had no effect on the network found, yielding the

same network (Figure 4a). As noise levels increased to half of the mean log-likelihood, one

other tree was generated (Figure 4f). This demonstrates the stability limit of the construction

but notably this revised tree still contains clusters that resemble the original. If the original tree

was based on biased likelihoods from imaging artifacts, the addition of noise would have

resulted in an altered tree network at lower noise levels and with higher frequency, but since

this did not occur, we are confident in the accuracy of our original likelihood estimations.

Finally, to exclude the possibility that datasets directly measuring C1 and C2 (reactions L and B,

respectively) were biasing the network towards inclusion of those sets of vRNPs, we performed

a network analysis of a dataset where the measured likelihoods of these two clusters was

excluded. Removal of data derived from images of the sets of vRNPs in C1 and C2 (Figure 4e)

yielded networks that resemble Figure 4a, indicating that even in the absence of data from

these clusters, similar clusters will form during assembly. Removal of each cluster alone was

recapitulated by lower order interactions. Removal of C1 [HA, M, NA, NS] data resulted in the

same tree presented in figure 4b, where HA, M, NA, NS form in two steps from a triplet and a

single vRNP segment. Exclusion of C2 [NP, PB1, PB2} result in a tree similar to figure 4d, where

C2 cluster forms from a pair and a single segment (Figure 4d). Removal of data was also

combined with the addition of noise to further test stability. Predominantly, the same networks

were generated with and without noise. As noise levels increased in the exclusion

constructions, a tree with HA, M, NS, *PB1* in place of C1 and NP, *PA*, PB2 in place of C2, where

PB1 and PA were not observed in the C1 or C2 original clusters respectively (Figure 4f).

Together, these constructed networks suggest a primary interaction scheme that can be further

experimentally tested and the importance of triplet and quadruplet observations in predicting

higher order vRNP complexes. Other, very low frequency trees that were generated in the

presence of high levels of noise are located in the reproducible research archive.

Discussion:

Influenza A vRNP segments selectively assemble within an infected cell to produce progeny

virions containing one copy of all eight segments. The mechanism driving selective assembly is

still largely unknown, but RNA-RNA interaction between the segments has been proposed [14-

16]. In this study, we have modeled the *in vivo* spatial dependencies of influenza A vRNP

segments using multi-color fluorescent *in situ* images to generate possible vRNP-vRNP

interaction networks and propose a new perspective on genome packaging during viral

replication. By utilizing a rigorous statistical framework, we have extended the efforts of

previous groups and present a novel method for construction of vRNP interaction networks

based on their precise spatial information within actively infected cells. Using spatial proximity

as a proxy for physical interaction, point process modeling in this study suggests clear spatial

dependencies between certain vRNP segments, and confirms our previous observations of

subcomplex formation during cytoplasmic transport (18).

Previous studies using *in vitro* transcribed RNA suggest multiple interactions between vRNP

segments are expected [3, 6, 14, 16, 28]. Similarly, the modeled likelihoods show potentially

multiple interactions for each vRNP segment. Electron tomography studies have revealed a

conserved '7+1' supramolecular structure within the viral interior, where a center shaft is

surrounded by seven remaining segments [3, 6, 28], demonstrating an ordered process in

genome assembly.

Our results suggest two candidates for the center shaft or 'master segment': PA and PB1. From

the six constructed networks, PA is the most variable in that it is least often paired early in

assembly but, with the addition of noise and exclusion of data, may interact with both primary

observed clusters. This would be expected if a segment was evenly dependent on most others,

a possible signature of a core segment. PB1 also behaves similarly but to a lesser degree. Both

PB1 and PA show the highest average pairwise model likelihood over all other segments.

A few vRNP pairs were seen often in the most likely pairwise, triplet, and quadruplet models

and in the final constructed network. PB1-PB2 and HA-M, show commutative relationships,

appearing in one of the top most likely triplet and quadruplet models for each vRNP within the

pair. Some non-commutative dependencies were also seen, most notably in NP depending on

PB2 in both triplet and quadruplet models with little PB2-NP dependence.

By using dynamic programming, we were able to generate the most likely interaction network

based on all models. While this network was surprising robust to the addition of noise to the

model likelihoods, subtle variations in the vRNP interaction network were observed with the

exclusion of higher order datasets, suggesting that there may exist multiple interaction

pathways all acting at once in a single cell. Some of the variation in our results may also be due

to the presence of some lower specificity vRNA-vRNA interactions also occurring during

genome assembly. This would explain some of the highly likely pairwise interactions observed

that were not included in the final interaction network, such as PB1 and HA. In addition, vRNP

networks could also change over the course of a viral infection when packaging fidelity may be

compromised [29]. Our current data set only considers one time point, eight hours post

infection, which represents the time of initial virion release from infected cells, and would miss

alternate networks present at later time points. Therefore, the analysis presented here may

only capture a snapshot of a dynamic assembly process that could change throughout an

infection. A system with multiple possible interaction networks could be potentially

advantageous in both ensuring packaging of a full genome in viral particles and in future

reassortment events. For example, reassortment of one vRNA segment during a coinfection

may decrease the binding efficacy necessary for a single interaction network but a second

network may serve to rescue viral viability while also increasing genetic diversity.

With this, we also point out the consistent presence of the PB2, PB1 and NP cluster. Since these

three proteins function together, with PA, to form the viral polymerase, which promote vRNA

replication and transcription, this cluster is potentially important. Viruses capable of packaging

complementary polymerase segments into progeny virions will have a fitness advantage

compared to viruses with polymerase mismatches. This phenotype has been observed in

experimental reassortment experiments between 2009 H1N1 pandemic and seasonal H3N2

viruses[25, 27].

More generally, the computational pipeline presented may function as a useful tool in

elucidating spatial dependency over a wide range of biological phenomena observable through

microscopy. As we have demonstrated, point process models can capture key aspects of point

distributions while retaining a basis in probability. Network construction synthesizes the results

derived through modeling into a cogent, most-likely set of spatial interactions or dependencies.

Future work in spatial modeling of influenza A packaging should incorporate a temporal

dimension through live cell imaging. Point process models have been adapted for

spatiotemporal modeling [30]. Incorporation of intracellular markers important for influenza A

vRNP transport may increase the accuracy of models. Influenza A vRNP transport from the

nucleus, the site of vRNP synthesis, to the plasma membrane is a complex process utilizing a

variety of host proteins [31]. Rab11A-containing vesicles are thought to be the primary mode of

transport although there is evidence that a Rab11A-independent mechanism exists [32-34]. A

recent study has implicated the ER to mediating transport of vRNP segments through anchored

Rab11A proteins [35]. Novel methods in fluorescent multiplexing [36] present an exciting

opportunity for observing many cellular structures which will provide a holistic image the

spatial dependence of vRNP segments upon subcellular structures and to each other.

Methods:

Multi-color fluorescent in situ hybridization images

This study used previously published multi-color FISH images from [19] that were generated at

the National Institutes of Health. Briefly, MDCK cells were infected with recombinant

WSN/1933 H1N1 for 8 hours and then fixed and stained with FISH probes, obtained from

Biosearch technologies, against four distinct vRNP segments. DAPI was included as well to label

DNA. Multi-color FISH samples were imaged on an Leica SP5 white light laser to ensure spectral

separation of the five colors (Dapi, Alexa 488, Quasar 570, Cal Fluor Red 590, Quasar 670). The

specificity of the probes and spectral separation between fluorophores was previously

confirmed [19].

Image preprocessing, cell segmentation and point detection

All image preprocessing was performed in MATLAB, v. 2015a. Prior to point detection and

segmentation, image noise was removed by convolution with a Gaussian mask. Each image

channel was denoised individually. As described in [19], each image was captured with 0.17 um

z-step size spanning the entire cell volume, defined for each cell using both the nuclear and

FISH staining to define the apical cell membrane. Each image was captured with a pixel size of

~50x50x168 nm.

Individual vRNP segments were identified by finding connected areas of signal within the

denoised image. The center of each object was taken as the maximum intensity pixel within,

yielding a set of point coordinates. Note that due to the thickness of the z sections, the

apparent distance between points may be an underestimate of the true distance, but this effect

is expected to average out when considering many points.

Since there was no fluorescent tag for cell membrane components, we estimated the cell

boundary using the vRNP images. We assumed that fluorescent signal would be denser within

the cell cytoplasm than in the surrounding area with cell-cell junctions subtly defined by

'valleys', curves of low signal density whose normal vectors point towards increasing density.

The gradient of point density over the image was then used to segment individual cells from

both the background and each other by the mean-shift algorithm [37]. Hand segmentation was

also used to ensure accurate cell membrane segmentation. In most cases, the convex hull of all

points within an identified cell region then defined the cell membrane. The nucleus was

segmented through simple thresholding and smoothing.

Point processes and patterns

Within each cell, the vRNP segments form a point pattern, $\mathbf{x} = \{x_1, x_2, ..., x_n\}$, where n is the

number of observed points and x_i is the 3-dimensional coordinate vector for point i. The point

pattern is defined over a bounded region, W, the segmented cell cytoplasm. $\mathbf{x}$ is then viewed as

a "realization" (an output) of some random point process $\mathbf{X}$, the generating distribution for all

patterns of the particular vRNP identity. The process $\mathbf{X}$ then represents the culmination of

biological factors that determines vRNP segment location, eg. nuclear export, directed

transport over the microtubule network, inhibition by other organelles, etc.

To model $\mathbf{X}$, we first define its locational density, f, a function over the cytoplasm, where $f(u)$ is

the probability of observing a point at position $\mathbf{u}$. The simplest model for this density is the

Homogenous Poisson, characterized by complete randomness over space, or uniform

probability:

$$f(\mathbf{x}|n) = \frac{1}{Z}\lambda^n \qquad\qquad (1)$$

where λ determines point density, and Z a normalizing constant.   In the point process

literature, λ is referred to as *intensity*, but we refer to it as point density to avoid confusion

with fluorescence intensity.

Characterizing spatial randomness

466 Hypothesis testing for spatial randomness, comparing an observed pattern to an

467 expectation under a Poisson assumption, was used as an initial motivation for further

468 modeling. Ripley's K-function describes the number of neighboring points within a given

469 distance (r) of each observed point in a pattern, a measure of clustering or inhibition of

470 points within space. Under a Poisson assumption,

$$K_{poi}(r) = \pi r^2 \qquad (2)$$

474 The expected K-function value can then be used to measure the difference between a given

475 pattern and that of a Homogenous Poisson with an equal number of points. To assess this

476 for observed point patterns, we used the test statistic

$$T = \int_0^{r_{max}} \left(\widehat{K}(r) - K_{poi}(r) \right)^2 dr \qquad (3)$$

480 where $\widehat{K}(r)$ is the estimated K-function for a given pattern and r a radius defining the point

481 neighborhood. To provide a background distribution for this statistic for a given cell geometry,

482 we generated 100 samples from a Homogenous Poisson process defined over its cell cytoplasm

483 and calculated the test statistic. We then obtained a p-value for the hypothesis that an

484 observed distribution was drawn from a homogenous Poisson process:

$$p = \frac{1 + \sum_{i=1}^{m} I(t_{obs} \geq t_i)}{m+1} \tag{4}$$

where m is the number of samples drawn. The p-values were averaged over all cells for a given

vRNP.

Modeling spatial dependence on cellular structures

An Inhomogenous Poisson Process, in its simplest form, is characterized by a spatially

dependent locational density. Per point "factors" (e.g., distance to cell membrane) are used to

determine the probability of a point occurring at a given location. As above, let **X** be some point

process with realization **x**, and define $s(\mathbf{x})$, an $n \times k$ matrix where each $s_{i,j}$ is some factor j that

can be calculated for each point x_i. The Inhomogenous Poisson point density function is:

$$\lambda(x_i; \theta) = \left(exp(\theta^T s(x_i)) \right)$$

500   where θ is the k-dimensional parameter vector. The likelihood of the model given point data is

501   then defined by:

$$l(\mathbf{x}; \theta) = \left(\prod_{i=1}^{n} \lambda(x_i) \right) exp \left(- \int_W \lambda(u) du \right)$$

To quantify vRNP dependence on cellular structures, we defined the first factor (f_1) as the

distance to the nearest point on the nuclear membrane, the distance to the nearest point on

the cell membrane, or the ratio of cell to nuclear distance.

Modeling spatial dependence between vRNPs

The second factor, minimum inter-pattern point distance, was used to explore whether there is

vRNP-vRNP interaction between vRNPs of different types. For each observed vRNP segment of

type l, and some other vRNP point pattern of type j, the minimum inter-pattern point distance

is:

$$f_2\left(x_i^l \mid l \neq t\right) = \min_j d\left(x_i^l, x_j^t\right)$$

For each vRNP type, we use a 'one depends on all' schema. That is, each model with the same

set of vRNPs is not equivalent but exclusively represents a single vRNP pattern (the primary

pattern) depending on others (the secondary patterns). Models were constructed for every

unique pair, triplet, and quadruplet of vRNP types.

For each model, all instances of its unique components were gathered, all images with tags for

every vRNP in the model. Parameters of the model were then fit for each instance using the

maximum pseudolikelihood [38]. The basic idea is that a given set of parameters can be used to

calculate a value that is proportional to the likelihood (which is referred to as a

pseudolikelihood) of observing a point at a given position, and the parameters can then be

adjusted to give the highest total pseudolikelihood for available observations. Hold one out

cross validation was used to assess the quality of fit over all instances yielding cross-validated

pseudolikelihood estimated parameters and variances. These fitted parameters were then used

to convert the pseudolikelihood to a true model likelihood by estimating the proportionality

constant as previously described (19).

There is some variation in the number of images in which different vRNPs were visualized

(Supplemental Table 1). This ranged from 4 for PB1 to 9 for M (the others were present 6 or 7

times). As pointed out by a reviewer, this difference raises the possibility that a vRNP that was

imaged less frequently might be found to be underrepresented in the extent to which other

segments were found to depend upon it. As seen in Table 3, this did not turn out to be the

case, as PB1 and M showed similar average likelihoods and more vRNPs showed strong

dependence on PB1 than on the others. Similarly, PB1 was not underrepresented in high

scoring triples (Table 5).

Interaction network assembly

We formulate the task of finding the most probable set of vRNA interactions yielding a full

genome as a graph problem. First, let the set of nodes in the graph be every possible

combination of vRNAs ranging in size from a single vRNA to all 8 segments. Directed edges in

the graph connect subset to superset nodes. We weight each edge of the graph by the

likelihood of the most likely model (chosen from models with only vRNP dependencies, and

those with additional dependencies on either nuclear distance, cell membrane distance, or cell

to nuclear distance ratio) that results in the superset and contains the subset as a primary or

secondary segment (Figure 1). Edges that represent models that have not been observed

remain unweighted.

From this graph, a set of interactions yielding the full genome is a tree with leaves as the vRNA

singletons and root as the full 8-mer of vRNA segments. In these trees, we require each node to

have at least two incoming edges, barring the leaf nodes. The set of incoming edges to a given

node must emanate from nodes whose labels, when combined, exactly equal the given node. A

set of incoming edges essentially 'combines' nodes, directly corresponding to the physical event

of vRNP segments binding. The most likely set of interactions is also the most highly weighted

tree. The problem of finding the most probable interaction network then reduces to finding the

highest weighted tree in the graph. This problem can easily be solved through dynamic

programming within a paradigm often used in evolutionary tree construction [24].

The likelihoods of models involving multiple vRNP segments can be taken as a measure of

spatial dependence of primary segments on secondary segments. vRNP segments that truly

interact are expected to be highly dependent either unidirectionally or bidirectionally. The cross

validated model likelihoods can be used to weight groups of edges in the graph. Since the graph

also has edges that were not observed in actual images (eg: any complex of more than 4

vRNAs), composite or inferred likelihoods were generated through recursion in a dynamic

programming scheme. For each unobserved complex, we took the most likely set of observed

models that could generate the complex.

For a given unobserved complex N = {N_1,N_2,N_3,...},

$$L(N) = \max(L(N_1|N_2,N_3,...),\ L(N_2|N_1,N_3,...),\ ...\ )$$

Then for some likelihood in the above, $L(N_1|N_2)$ where $N_1 = (n^1{}_1,\ ...\ n^1{}_k)$ and $N_2 = (n^2{}_1,\ ...\ n^2{}_l)$,

$|N_1| = k$, $|N_2| = l$,

$$L\big((N_1|N_2)\big) = \sum_{i=1}^{k} \underset{N_j^2 \in all\ unique\ sets\ of\ N_2}{\text{average}} (L(n_i^1|N_j^2))$$

Interaction network stability analysis

To assess the stability of the generated interaction network, two procedures of perturbation

were performed: introduction of noise and removing certain high likelihood models. We

simulated noise as a normal random distribution centered at 0 with varying standard

deviations, termed the noise levels. Under increasing noise levels, each observed model

likelihood was amended with a noise value drawn from this distribution prior to interaction

network construction. For each noise level, we simulated 100 trials and tallied the proportion of

output trees that contained each possible edge. Trees that align with the non-noisy assembly

signal stability while highly variable trees signal instability.

Reproducible research archive

All images, derived data and source code will be available upon publication at

http://murphylab.cbd.cmu.edu/software.

REFERENCES

<div align="center">**Bibliography**</div>
602

1.   Morens DM, Taubenberger JK. Pandemic influenza: certain uncertainties. Rev Med
Virol. 2011;21(5):262-84. Epub 2011/06/28. doi: 10.1002/rmv.689. PubMed PMID:
21706672.
2.   Sidorenko Y, Reichl U. Structured model of influenza virus replication in MDCK cells.
Biotechnol Bioeng. 2004;88(1):1-14. doi: 10.1002/bit.20096. PubMed PMID: 15384040.
3.   Hutchinson EC, von Kirchbach JC, Gog JR, Digard P. Genome packaging in influenza A
virus. J Gen Virol. 2010;91(Pt 2):313-28. Epub 2009/12/04. doi: 10.1099/vir.0.017608-0.
PubMed PMID: 19955561.
4.   Nakatsu S, Sagara H, Sakai-Tagawa Y, Sugaya N, Noda T, Kawaoka Y. Complete and
Incomplete Genome Packaging of Influenza A and B Viruses. MBio. 2016;7(5). doi:
10.1128/mBio.01248-16. PubMed PMID: 27601575; PubMed Central PMCID:
PMCPMC5013298.
5.   Lee N, Le Sage V, Nanni AV, Snyder DJ, Cooper VS, Lakdawala SS. Genome-wide
analysis of influenza viral RNA and nucleoprotein association. Nucleic Acids Res.
2017;45(15):8968-77. doi: 10.1093/nar/gkx584. PubMed PMID: 28911100.
6.   Noda T, Sagara H, Yen A, Takada A, Kida H, Cheng RH, et al. Architecture of
ribonucleoprotein complexes in influenza A virus particles. Nature. 2006;439(7075):490-2.
Epub 2006/01/27. doi: 10.1038/nature04378. PubMed PMID: 16437116.

7.   Chou YY, Vafabakhsh R, Doganay S, Gao Q, Ha T, Palese P. One influenza virus
particle packages eight unique viral RNAs as shown by FISH analysis. Proc Natl Acad Sci U S
A. 2012;109(23):9101-6. Epub 2012/05/02. doi: 10.1073/pnas.1206069109. PubMed
PMID: 22547828; PubMed Central PMCID: PMC3384215.
8.   Noda T, Kawaoka Y. Packaging of influenza virus genome: robustness of selection.
Proc Natl Acad Sci U S A. 2012;109(23):8797-8. Epub 2012/05/23. doi:
10.1073/pnas.1206736109. PubMed PMID: 22615368; PubMed Central PMCID:
PMC3384147.
9.   Isel C, Munier S, Naffakh N. Experimental Approaches to Study Genome Packaging of
Influenza A Viruses. Viruses. 2016;8(8). doi: 10.3390/v8080218. PubMed PMID:
27517951; PubMed Central PMCID: PMCPMC4997580.
10.   Fujii Y, Goto H, Watanabe T, Yoshida T, Kawaoka Y. Selective incorporation of
influenza virus RNA segments into virions. Proc Natl Acad Sci U S A. 2003;100(4):2002-7.
doi: 10.1073/pnas.0437772100. PubMed PMID: 12574509; PubMed Central PMCID:
PMCPMC149948.
11.   Marsh GA, Rabadan R, Levine AJ, Palese P. Highly conserved regions of influenza a
virus polymerase gene segments are critical for efficient viral RNA packaging. J Virol.
2008;82(5):2295-304. doi: 10.1128/JVI.02267-07. PubMed PMID: 18094182; PubMed
Central PMCID: PMCPMC2258914.
12.   Essere B, Yver M, Gavazzi C, Terrier O, Isel C, Fournier E, et al. Critical role of
segment-specific packaging signals in genetic reassortment of influenza A viruses. Proc Natl
Acad Sci U S A. 2013;110(40):E3840-8. doi: 10.1073/pnas.1308649110. PubMed PMID:
24043788; PubMed Central PMCID: PMCPMC3791739.
13.   Gerber M, Isel C, Moules V, Marquet R. Selective packaging of the influenza A
genome and consequences for genetic reassortment. Trends Microbiol. 2014;22(8):446-55.
doi: 10.1016/j.tim.2014.04.001. PubMed PMID: 24798745.
14.   Fournier E, Moules V, Essere B, Paillart JC, Sirbat JD, Cavalier A, et al. Interaction
network linking the human H3N2 influenza A virus genomic RNA segments. Vaccine.
2012;30(51):7359-67. Epub 2012/10/16. doi: 10.1016/j.vaccine.2012.09.079. PubMed
PMID: 23063835.
15.   Fournier E, Moules V, Essere B, Paillart JC, Sirbat JD, Isel C, et al. A supramolecular
assembly formed by influenza A virus genomic RNA segments. Nucleic Acids Res.
2012;40(5):2197-209. Epub 2011/11/15. doi: 10.1093/nar/gkr985. PubMed PMID:
22075989; PubMed Central PMCID: PMC3300030.
16.   Gavazzi C, Isel C, Fournier E, Moules V, Cavalier A, Thomas D, et al. An in vitro
network of intermolecular interactions between viral RNA segments of an avian H5N2
influenza A virus: comparison with a human H3N2 virus. Nucleic Acids Res.
2013;41(2):1241-54. Epub 2012/12/12. doi: 10.1093/nar/gks1181. PubMed PMID:
23221636; PubMed Central PMCID: PMC3553942.
17.   Gilbertson B, Zheng T, Gerber M, Printz-Schweigert A, Ong C, Marquet R, et al.
Influenza NA and PB1 Gene Segments Interact during the Formation of Viral Progeny:
Localization of the Binding Region within the PB1 Gene. Viruses. 2016;8(8). doi:
10.3390/v8080238. PubMed PMID: 27556479; PubMed Central PMCID: PMCPMC4997600.
18.   Chou YY, Heaton NS, Gao Q, Palese P, Singer R, Lionnet T. Colocalization of different
influenza viral RNA segments in the cytoplasm before viral budding as shown by single-
molecule sensitivity FISH analysis. PLoS Pathog. 2013;9(5):e1003358. Epub 2013/05/15.

doi: 10.1371/journal.ppat.1003358. PubMed PMID: 23671419; PubMed Central PMCID:
PMC3649991.
19.   Lakdawala SS, Wu Y, Wawrzusin P, Kabat J, Broadbent AJ, Lamirande EW, et al.
Influenza a virus assembly intermediates fuse in the cytoplasm. PLoS Pathog.
2014;10(3):e1003971. doi: 10.1371/journal.ppat.1003971. PubMed PMID: 24603687;
PubMed Central PMCID: PMCPMC3946384.
20.   Johnson GR, Li J, Shariff A, Rohde GK, Murphy RF. Automated Learning of Subcellular
Variation among Punctate Protein Patterns and a Generative Model of Their Relation to
Microtubules. PLoS Comput Biol. 2015;11(12):e1004614. doi:
10.1371/journal.pcbi.1004614. PubMed PMID: 26624011; PubMed Central PMCID:
PMCPMC4704559.
21.   Li Y, Majarian TD, Naik AW, Johnson GR, Murphy RF. Point process models for
localization and interdependence of punctate cellular structures. Cytometry A.
2016;89(7):633-43. doi: 10.1002/cyto.a.22873. PubMed PMID: 27327612.
22.   Griffie J, Shlomovich L, Williamson DJ, Shannon M, Aaron J, Khuon S, et al. 3D
Bayesian cluster analysis of super-resolution data reveals LAT recruitment to the T cell
synapse. Sci Rep. 2017;7(1):4077. doi: 10.1038/s41598-017-04450-w. PubMed PMID:
28642595; PubMed Central PMCID: PMCPMC5481387.
23.   Yunta ML, Lagache T, Santi-Rocca J, Bastin P, Olivo-Marin JC, editors. A statistical
analysis of spatial clustering along cell filaments using Ripley's K function. 2014 IEEE 11th
International Symposium on Biomedical Imaging (ISBI); 2014 April 29 2014-May 2 2014.
24.   Thompson JD, Higgins DG, Gibson TJ. CLUSTAL W: improving the sensitivity of
progressive multiple sequence alignment through sequence weighting, position-specific
gap penalties and weight matrix choice. Nucleic Acids Res. 1994;22(22):4673-80.
25.   Li C, Hatta M, Watanabe S, Neumann G, Kawaoka Y. Compatibility among polymerase
subunit proteins is a restricting factor in reassortment between equine H7N7 and human
H3N2 influenza viruses. J Virol. 2008;82(23):11880-8. doi: 10.1128/JVI.01445-08. PubMed
PMID: 18815312; PubMed Central PMCID: PMCPMC2583690.
26.   Te Velthuis AJ, Fodor E. Influenza virus RNA polymerase: insights into the
mechanisms of viral RNA synthesis. Nat Rev Microbiol. 2016;14(8):479-93. doi:
10.1038/nrmicro.2016.87. PubMed PMID: 27396566; PubMed Central PMCID:
PMCPMC4966622.
27.   Phipps KL, Marshall N, Tao H, Danzy S, Onuoha N, Steel J, et al. Seasonal H3N2 and
2009 Pandemic H1N1 Influenza A Viruses Reassort Efficiently but Produce Attenuated
Progeny. J Virol. 2017;91(17). doi: 10.1128/JVI.00830-17. PubMed PMID: 28637755;
PubMed Central PMCID: PMCPMC5553182.
28.   Noda T, Sugita Y, Aoyama K, Hirase A, Kawakami E, Miyazawa A, et al. Three-
dimensional analysis of ribonucleoprotein complexes in influenza A virus. Nat Commun.
2012;3:639. Epub 2012/01/26. doi: 10.1038/ncomms1647. PubMed PMID: 22273677;
PubMed Central PMCID: PMC3272569.
29.   Brooke CB. Population Diversity and Collective Interactions during Influenza Virus
Infection. J Virol. 2017;91(22). Epub 2017/09/01. doi: 10.1128/JVI.01164-17. PubMed
PMID: 28855247; PubMed Central PMCID: PMCPMC5660503.
30.   Illian JB, Soerbye, S, Rue, H & Hendrichsen, D. Using INLA to fit a complex point
process model with temporally varying effects – a case study Journal of Environmental
Statistics. 2012;3(7).

31.   Lakdawala SS, Fodor E, Subbarao K. Moving On Out: Transport and Packaging of
Influenza Viral RNA into Virions. Annu Rev Virol. 2016;3(1):411-27. doi: 10.1146/annurev-
virology-110615-042345. PubMed PMID: 27741407.
32.   Bruce EA, Digard P, Stuart AD. The Rab11 pathway is required for influenza A virus
budding and filament formation. J Virol. 2010;84(12):5848-59. Epub 2010/04/02. doi:
10.1128/JVI.00307-10. PubMed PMID: 20357086; PubMed Central PMCID: PMC2876627.
33.   Eisfeld AJ, Kawakami E, Watanabe T, Neumann G, Kawaoka Y. RAB11A is essential
for transport of the influenza virus genome to the plasma membrane. J Virol.
2011;85(13):6117-26. Epub 2011/04/29. doi: 10.1128/JVI.00378-11. PubMed PMID:
21525351; PubMed Central PMCID: PMC3126513.
34.   Nturibi E, Bhagwat AR, Coburn S, Myerburg MM, Lakdawala SS. Intracellular
Colocalization of Influenza Viral RNA and Rab11A is Dependent upon Microtubule
Filaments. J Virol. 2017;91(19). doi: 10.1128/JVI.01179-17. PubMed PMID: 28724771.
35.   de Castro Martin IF, Fournier G, Sachse M, Pizarro-Cerda J, Risco C, Naffakh N.
Influenza virus genome reaches the plasma membrane via a modified endoplasmic
reticulum and Rab11-dependent vesicles. Nat Commun. 2017;8(1):1396. doi:
10.1038/s41467-017-01557-6. PubMed PMID: 29123131; PubMed Central PMCID:
PMCPMC5680169.
36.   Lin JR, Fallahi-Sichani M, Sorger PK. Highly multiplexed imaging of single cells using
a high-throughput cyclic immunofluorescence method. Nat Commun. 2015;6:8390. doi:
10.1038/ncomms9390. PubMed PMID: 26399630; PubMed Central PMCID:
PMCPMC4587398.
37.   Comaniciu D, Meer P. Mean shift: A robust approach toward feature space analysis.
IEEE Transactions on pattern analysis and machine intelligence. 2002;24(5):603-19.
38.   Baddeley A, Turner R. Practical maximum pseudolikelihood for spatial point
patterns. Australian & New Zealand Journal of Statistics. 2000;42(3):283-322.
739
740

Figure 1: Illustration of the analysis pipeline. (a) Unprocessed fluorescence channels for cells FISH tagged for four vRNA segments and stained with DAPI. (b) Preprocessed image channels after filtering, nuclear and cell membrane segmentation, and point identification. The 3D images in (a) and (b) are displayed as maximum value projections. (c) Factors are calculated for each channel in each image, model parameters are fit, and a likelihood generated. (d) Dynamic programming is used to generate a most likely set of interactions yielding a full genome. Edges are possible interactions, solid lines indicate edges that are part of the most likely set of interactions. In this toy example, the most likely set of interactions is A and B binding to form AB, and then C binding to AB.

754 Figure 2: Pairwise interactions learned from MDCK cell images. Each vRNA segment is arranged
755 around the perimeter. Bands connect vRNA segments that interact with high likelihood. The
756 position of the base of each band indicates the independent (secondary) variable and the larger
757 the width of the base the more likely the interaction. Only interactions whose log likelihoods
758 are below 0.23 from Table 4 are shown.
759

b.

Figure 3: Comparison of composite and observed model likelihoods for triplet and quadruplet
models. To ensure that the recursion in our dynamic programming scheme accurately predicts
model likelihoods, we compared triplet and quadruplet observed likelihoods to those
generated from only pairwise models (that include all four possible models: vRNP pairs alone,
vRNP pairs + nucleus, vRNP pairs + cell membrane, vRNP pairs + nuclear/cell membrane ratio)
(a). We also compared observed quadruplet likelihoods to those generated from pairwise and
triplet models (using all 4 possible models) (b). Ideally, all points would fall on the red line,
matching observed and predicted values perfectly. The cluster in the lower left corner
represent interactions that are predicted to occur with high likelihood but in reality, do not. All
points within this cluster contain the same four vRNA segments (PB1, PB2, NA, NS). Exclusion of
this cluster from our analysis increases the association between predicted and observed triplets
and quadruplets.
774

Figure 4: Constructed interaction networks. Using dynamic programming, we constructed the single most likely set of interactions yielding a full genome. First, we used all model likelihoods with and without nuclear and cell features (a) to build a master tree. Hotter edge colors correspond to more likely interactions. Since the images contain only a subset of the possible triplets and quadruplet, we assessed how much this affected the master tree by removing all quadruplets (b) or both quadruplets and triplets (c). To test the stability of the master tree, trees were constructed with random noise added to each observed model likelihood before performing the dynamic programing. Multiple noise levels were tested up to 1/2 of the mean model likelihood with the noise level being the standard deviation of a Gaussian centered at 0. For each level of noise, network construction was performed 100 times. Noise up to 1/4 mean likelihood always resulted in (a). Increasing noise levels yielded (a) on 40% to 60% of constructions. Given the two observed clusters in the master tree, we next asked whether the composite likelihood function could recapture their interactions without actually observing them. Removing all models involving HA, M, NS, NA (C1) and NP, PB1, PB2 (C2) yielded (d). Adding noise up to 1/4 mean likelihood and removing both clusters also generated (d). Removal with higher levels of noise generated (e) for 27% of constructions. In all stability analyses with noise greater than 1/4 mean likelihood, (f) was generated with frequency between 5% and 15%.

796   Table 1: Monte Carlo p-values and variances for the hypothesis that the spatial distribution of
797   each vRNA segment is distributed by a homogeneous Poisson process.

	Mean p-value	Variance
PB2	0.0034	0.00011
PB1	0.00524	0.00034
PA	0.00467	0.0002
HA	0.00671	0.00015
NP	0.01185	0.00036
NA	0.00249	0.00003
M	0.00005	0
NS	0.00306	0.00007

798
799
800   Table 2: Cross validated negative log-likelihoods for models dependent on distance to different
801   reference locations.
802

	Cell Model	Nuclear Model	Cell+Nuclear Model
PB2	2.5547	0.4153	0.9158
PB1	2.3908	0.408	1.2574
PA	2.4228	0.4643	1.2352
HA	2.443	2.3928	4.396
NP	2.4885	2.4584	4.5447
NA	2.5019	2.4372	2.1754
M	2.4871	2.1294	3.0799
NS	2.3906	0.6666	1.4624

803
804
805
806   Table 3: Cross validated negative log-likelihoods for pairwise interaction models. The row label
807   indicates the independent (secondary) variable, the vRNP upon which a model predicting the
808   column label (the dependent or primary variable) was constructed.
809

	PB2	PB1	PA	HA	NP	NA	M	NS
PB2		0.3705	0.2694	0.282	0.3146	0.5605	0.2146	0.3926
PB1	0.2398		0.2018	0.2044	0.1186	0.4015	0.1701	0.2905
PA	0.2276	0.2217		0.2284	0.1502	0.3222	0.2816	0.2315
HA	0.3334	0.1266	0.1838		0.2787	0.232	0.2623	0.2845
NP	0.3438	0.1805	0.1942	0.2895		0.2529	0.275	0.3411
NA	0.3191	0.324	0.3093	0.2947	0.2742		0.2747	0.2982
M	0.3287	0.2236	0.2832	0.2803	0.2975	0.2897		0.2848
NS	0.3427	0.2888	0.2292	0.2838	0.325	0.3436	0.2841	

810
811
812

813 Table 4: Cross validated negative log-likelihoods for pairwise interaction models that also
814 include nuclear distance. The row label indicates the independent (secondary) variable, the
815 vRNP upon which a model predicting the column label (the dependent or primary variable) was
816 constructed.
817

	PB2	PB1	PA	HA	NP	NA	M	NS
PB2		0.3817	0.2694	0.2833	0.3145	0.5643	0.2166	0.3977
PB1	0.2427		0.2023	0.2191	0.1215	0.4188	0.1714	0.3097
PA	0.2278	0.2211		0.661	0.1504	0.3238	0.4967	0.5378
HA	0.336	0.13	0.9138		1.944	1.7048	1.2324	0.5447
NP	0.3084	0.1746	0.1949	0.2095		1.6062	1.6495	0.3523
NA	0.3166	0.3317	0.3114	1.7801	1.4877		1.0989	0.3
M	0.33	0.2246	0.5374	1.2838	1.7105	1.1884		0.4153
NS	0.3454	0.2816	0.6259	0.4561	1.7998	0.3466	0.4368	

818
819

Table 5: Cross validated negative log-likelihoods for the top two triple models with each vRNA segment as the primary pattern. Pairwise model negative log-likelihoods for the members of each model are shown in the right columns.

RNA	CV Likelihood	PB2	PB1	PA	HA	NP	NA	M	NS
PB2									
PB2 M PB1	0.114		0.382					0.217	
PB2 HA M	0.162				0.283			0.217	
PB1									
PB1 PA PB2	0.062	0.243		0.202					
PB1 NP PB2	0.071	0.243				0.122			
PA									
PA PB2 PB1	0.113	0.228	0.222						
PA PB2 NP	0.121	0.228				0.15			
HA									
HA M PB1	0.059		0.13					1.232	
HA PB1 NS	0.111		0.13						0.545
NP									
NP PB1 PB2	0.141	0.344	0.181						
NP PB2 PA	0.145	0.344		0.195					
NA									
NA HA NP	0.225				1.78	1.488			
NA HA NS	0.246				1.78				0.3
M									
M PB1 NS	0.201		0.225						0.415
M HA PB1	0.201		0.225		1.284				
NS									
NS M PB1	0.169		0.289					0.437	
NS HA PB1	0.184		0.289		0.456				

826 Table 6: Cross validated negative log-likelihoods for the top two quadruple models with each
827 vRNA segment as the primary pattern. Pairwise model negative log-likelihoods for the members
828 of each model are shown in the right columns.
829

RNA	CV Likelihood	PB2	PB1	PA	HA	NP	NA	M	NS
PB2									
PB2 NS PB1 NA	0.571		0.382				0.564		0.398
PB2 NP PA PB1	0.258		0.382	0.269		0.315			
PB1									
PB1 NS NA PB2	0.347	0.243					0.419		0.31
PB1 PA M PB2	0.16	0.243		0.202				0.171	
PA									
PA NS M HA	0.655			0.438	0.661			0.497	0.538
PA NA M NS	0.272			0.438			0.324	0.497	0.538
HA									
HA PA M NS	0.869			0.914				1.232	0.545
HA NS PB2 M	0.317	0.336						1.232	0.545
NP									
NP HA M NA	0.233				2.095		1.606	1.65	
NP PB2 PA PB1	0.139	0.344	0.181	0.195					
NA									
NA PA NS M	0.272			0.311				1.099	0.3
NA PB1 PB2 NS	0.268	0.319	0.332						0.3
M									
M NS HA PA	0.7			0.537	1.284				0.415
M PA NS NA	0.336			0.537			1.188		0.415
NS									
NS HA PA M	0.705			0.626	0.456			0.437	
NS PB2 M HA	0.340	0.343			0.288			0.289	

830
831
832

833 Supp. Table 1: FISH imaging probe sets. The leftmost column is the experiment label with the
834 right 4 columns showing which vRNA segments were tagged. Each experiment also used DAPI
835 staining.

Reaction	Probe 1	Probe 2	Probe 3	Probe 4
A	M	HA	PB2	NS
B	PB1	PB2	PA	NP
C	PB1	M	HA	NS
D	M	PA	HA	NS
E	M	HA	NA	NP
F	PB1	PB2	NA	NS
G		PA	NA	
H			NP	NS
I	PB1	M	PB2	PA
J		M	NA	NP
K		PB2	HA	NP
L	M	HA	NA	NS
M	M	NA	HA	PA
N	M	PB2	NP	
Rxn 1	M	PA	NA	NS

836
837
838

839　Supp. Table 2: Observed triplets within the reaction set. 32 of 56 possible triplets were
840　observed.
841

Observed triplets	Reaction
PB2, PB1, PA	B, I
PB2, PB1, NP	B
PB2, PB1, NA	F
PB2, PB1, M	I
Pb2, PB1, NS	F
PB2, PA, NP	B
PB2, PA, M	I
PB2, HA, NP	K
PB2, HA, M	A
PB2, HA, NS	A
PB2, NP, NS	N
PB2, NA, NS	F
PB2, M, NS	A
PB1, PA, NP	B
PB1, PA, M	I
PB1, HA, M	C
Pb1, HA, NS	C
PB1, NA, NS	F
PB1, M, NS	C
PA, HA, NA	M
PA, HA, M	D, M
PA, HA, NS	D
PA, NA, M	M, Rxn 1
PA, NA, NS	Rxn 1
PA, M, NS	Rxn 1
HA, NP, NA	E
HA, NP, M	E
HA, NA, M	E, L, M
HA, NA, NS	L
HA, M, NS	L
NP, NA, M	E, J
NA, M, NS	L, Rxn 1

842
843
844

845 Supp. Table 3: Cross validated log-variances for pairwise interaction models shown in Table 3.
846
847 (large)
848
849 Supp. Table 4: Variance in the cross validated negative log-likelihoods for pairwise interaction
850 models with cell-to-nuclear distance ratio shown in Table 4. The row label indicates the
851 independent variable, the vRNP upon which a model predicting the column label (dependent
852 variable) was constructed.
853

	PB2	PB1	PA	HA	NP	NA	M	NS
PB2		0.0552	0.0072	0.0310	0.0086	0.0028	0.0025	0.0919
PB1	0.0172		0.0025	0.0012	0.0010	0.0002	0.0012	0.0067
PA	0.0091	0.0055		0.0054	0.0035	0.0061	0.0055	0.0103
HA	0.0175	0.0018	0.0011		0.0052	0.0064	0.0119	0.0146
NP	0.0489	0.0006	0.0015	0.0029		0.0060	0.0079	0.0003
NA	0.0001	0.0002	0.0023	0.0561	0.0143		0.0066	0.0025
M	0.0146	0.0022	0.0064	0.0146	0.0104	0.0037		0.0068
NS	0.0033	0.0068	0.0083	0.0038	0.0004	0.0011	0.0042	

854
855
856 Supp. Table 5: Cross validated log-likelihoods for all observed triplets.
857
858 (large)
859
860 Supp. Table 6: Cross validated log-likelihoods for all observed quadruplets.
861
862 (large)
863
864
865