1   **A Ca^{2+}-regulated deAMPylation switch in human and bacterial FIC proteins.**

3   Simon Veyron[1], Giulia Oliva[2,3], Monica Rolando[2,3], Carmen Buchrieser[2,3],

4   Gérald Peyroche[1], Jacqueline Cherfils[1]

[1]Laboratoire de Biologie et Pharmacologie Appliquée, CNRS and Ecole normale supérieure

Paris-Saclay, Cachan, France

[2] Institut Pasteur, Biologie des Bactéries Intracellulaires, Paris, France

[3] CNRS, UMR 3525, Paris, France

Correspondence to :

Jacqueline Cherfils

Laboratoire de Biologie et Pharmacologie Appliquée,

CNRS and Ecole normale supérieure Paris-Saclay,

61 Avenue du Président Wilson

94235 Cachan CEDEX, France

jacqueline.cherfils@ens-paris-saclay.fr

Abstract

FIC proteins regulate molecular processes from bacteria to humans by catalyzing post-

translational modifications (PTM), the most frequent being the addition of AMP using ATP as

a cofactor, a reaction coined AMPylation. In a large group of AMPylating FIC proteins,

which includes single-domain bacterial FIC proteins and animal FICD/HYPE, AMPylation is

intrinsically repressed by a structurally conserved glutamate. Curiously, FICD departs from

previously characterized bacterial FIC proteins in that it acts bifunctionally to AMPylate and

deAMPylate its target, the endoplasmic reticulum BiP/GRP78 chaperone. BiP is a key

component of the unfolded protein response (UPR), and is AMPylated under normal

conditions where its activity is low, while its activation correlates with its deAMPylation.

Currently, a direct signal regulating AMPylation efficiency in bacterial and animal FIC

proteins has not been identified. Here, we addressed this question for a FIC protein from the

bacterial pathogen *Enterococcus faecalis* (EfFIC) and for human FICD. We discover that

EfFIC catalyzes both AMPylation and deAMPylation within the same active site, suggesting

that the conserved glutamate is the signature of AMPylation/deAMPylation bifunctionality.

Crystal structures and PTM assays identify a multi-position metal switch implemented by the

glutamate, whereby EfFIC uses Mg^{2+} and Ca^{2+} to control AMPylation and deAMPylation

differentially without conformational change. Remarkably, we find that variations in Ca^{2+}

levels also tune deAMPylation of BiP by human FICD. Together, our results identify metals

as diffusible signals that can regulate bifunctional FIC proteins directly, and they suggest that

FICD has features of an enzymatic sensor of Ca^{2+} depletion, a hallmark of the UPR.

Introduction

In less than a decade, FIC proteins have emerged as a large family of enzymes controling the

activity of target proteins by post-translationally modifying them with phosphate-containing

compounds (reviewed in [1,2,3,4]). These proteins are characterized by the presence of a

conserved FIC domain, which carries out the post-translational modification (PTM) of a Tyr,

Ser or Thr residue in a target protein [5,6,7,8,9,10,11]. The most frequent PTM reaction catalyzed

by FIC enzymes is the addition of AMP using ATP as a cofactor, coined AMPylation or

adenylylation. This PTM activity was originally discovered in toxins from bacterial

intracellular pathogens [12] and was later identified in the toxin component of bacterial

toxin/antitoxins modules (e.g. Bartonella VbhT/VbhA, [7]) and in various other bacterial FIC

proteins of unknown functions, including single-domain (e.g. Neisseria FIC [7]) and larger (e.g.

Clostridium FIC [11]) FIC proteins. Metazoans possess a single FIC protein, FICD/HYPE

(FICD hereafter), which controls the reversible AMPylation of BiP/GRP78, an Hsp70

chaperone localized in the endoplasmic reticulum (ER) [13,14,15]. BiP is a major component of

the unfolded protein response (UPR), a major pathway whereby cells respond to ER stress

(reviewed in [16]). AMPylation of BiP reduces its affinity for unfolded protein clients [17] and

inversely correlates with the burden of unfolded proteins [15]. In a recent twist, FICD was

shown to act bifunctionally in both AMPylation and deAMPylation of BiP and to carry out

deAMPylation as its primary enzymatic activity in vitro [18].

All PTM reactions catalyzed by FIC proteins use a motif of conserved sequence for catalysis,

the FIC motif, which carries an invariant histidine that is critical for nucleophilic attack of the

cofactor by the target residue, and an acidic residue (aspartate or glutamate) that binds an

Mg^{2+} ion to stabilize the negative charges of the cofactor phosphates at the transition state

(reviewed in [1,2,3,4]). A large group of FIC proteins also features a structurally conserved

glutamate that represses AMPylation by protruding into the catalytic site from either N-

terminal elements, as in human FICD [10], *Clostridum difficile* FIC [11] or *Shewanella oneidensis*

FIC [19], or from a C-terminal α-helix as in single-domain FIC proteins from *Neisseria*

meningitidis [7] and *Helicobacter pylori* (PDB 2F6S). Crystal structures showed that this

glutamate prevents ATP from binding in a catalytically-competent conformation [7,11], while its

mutation into glycine creates space for the γ-phosphate of ATP [7] and has been consistently

shown to increase AMPylation activities *in vitro* and in cells (reviewed in [20]). These

observations led to propose that this conserved glutamate represses AMPylation by impairing

the utilization of ATP as a donor for AMP, hence that it must be displaced to allow productive

binding of ATP [7]. In *N. meningitidis* FIC (NmFIC), upregulation of the AMPylation activity

was proposed to occur upon changes in the toxin concentration [21]. In this scheme, NmFIC is

in an inactive tetrameric state at high concentration, which is further stabilized by ATP, while

its dilution promotes its conversion into a monomeric state, leading to activation by

displacement of the inhibitory glutamate followed by autoAMPylations that reinforce

AMPylation efficiency [21]. Whether a similar mechanism applies to other glutamate-bearing

FIC proteins has not been investigated. Remarkably, in bifunctional human FICD, the

autoinhibitory glutamate is critical for deAMPYlation of the BiP chaperone [18], a reaction

whereby FICD contributes to the activation of BiP under ER stress conditions. Currently, a

signal that represses the deAMPylation activity of FICD under normal ER conditions has not

been identified. Together, these intriguing observations raise the question of whether

displacement of the autoinhibitory glutamate is the sole mode of regulation of glutamate-

containing AMPylating FIC proteins, and call for investigation of diffusible signals directly

able to release autoinhibition in glutamate-bearing AMPylating FIC proteins and to regulate

deAMPylation in FICD.

In this study, we addressed this question by combining structural analysis and PTM assays of

a single-domain FIC protein from *Enterococcus faecalis* (EfFIC), which carries an

autoinhibitory glutamate in C-terminus, and of human FICD. Enteroccoci are commensals of

the gastrointestinal tract that become pathogenic outside of the gut and cause difficult-to-treat

infections in the hospital due to acquisition and transmission of antibiotic resistance [22,23]. We

discover that EfFIC has both AMPylation and deAMPylation activities borne by the same

active site. Furthermore, the conserved glutamate implements a Ca^{2+}-controlled metal switch

that tunes the balance between these activities without conformational change. Finally, we

show that Ca^{2+} is an inhibitor of deAMPylation of BiP by human FICD. Our findings predict

that the presence of a glutamate is a signature for AMPylation/deAMPylation bifunctionality

in FIC proteins from bacteria to human, and they identify metals as diffusible signals that can

directly modulate the activity of glutamate-bearing FIC proteins by binding into the active

site. Importantly, they suggest that FICD is an enzymatic Ca^{2+} sensor which responds to the

drop in Ca^{2+} concentration, a hallmark of ER stress, with implications in diseases that ilvolve

the UPR and for therapeutic strategies.

Results.
113
114   *EfFIC is an AMPylator.*
115
Enterococcus faecalis FIC belongs to class III FIC proteins, which are comprised of a single
FIC domain and carry an autoinhibitory glutamate in their C-terminal α-helix. We determined
crystal structures of unbound, phosphate-bound, AMP-bound and ATPγS-bound wild-type
EfFIC (EfFIC^{WT}) and of unbound and sulfate-bound EfFIC carrying a mutation of the
catalytic histidine into an alanine (EfFIC^{H111A}) (**Table 1 and Table S1**). These structures
were obtained in different space groups, yielding 32 independent copies of the EfFIC
monomer with various environments in the crystal. All EfFIC monomers resemble closely to
each other and to structures of other class III FIC proteins (**Figure 1A**). Notably, the C-
terminal α-helix that bears the inhibitory glutamate shows no tendency for structural
flexibility, even in subunits that are free of intersubunit contacts in the crystal. The glutamate
has the same conformation as in other glutamate-bearing FIC protein structures (**Figure 1B**)
and is stabilized by intramolecular interactions and, when present, by interactions with the
nucleotide cofactor (**Figure 1C**). Two crystal structures were obtained in co-crystallization
with a non-hydrolyzable ATP analog (ATPγS), for which well-defined electron density was
observed for the ADP moiety (**Figure S1A**). The positions of the α and β phosphates of
ATPγS in these structures depart markedly from those seen in ATP bound unproductively to
wild type NmFIC [7], or bound non-canonically to CdFIC [11] (**Figure 1D**). In contrast, they
superpose well to cofactors bound in a position competent for PTM transfer [7,8] (**Figure 1E**).
This observation prompted us to assess whether EfFIC is competent for AMPylation, using
autoAMPylation as a convenient proxy in the absence of a known physiological target
(reviewed in [20]). Using [α-^{32}P]-ATP and autoradiography to measure the formation of
AMPylated EfFIC (denoted ^{AMP*}EfFIC^{WT}), we observed that EfFIC^{WT} has conspicuous
autoAMPylation activity in the presence of Mg^{2+} (**Figure 1F**). Mutation of the inhibitory
glutamate into glycine (E190G) increased AMPylation, indicating that AMPylation of
EfFIC^{WT} is not optimal (**Figure 1F**). We conclude from these experiments that wild-type
EfFIC has canonical features of an AMPylating FIC enzyme, and that the inhibitory glutamate
mitigates this activity.
143
144

EfFIC is a deAMPylator in the presence of Ca²⁺

To gain further insight into the enzymatic properties of EfFIC, we solved the crystal structure

of EfFIC bound to AMP (EfFIC^WT-AMP) (**Table 1 and Table S1**). AMP superposes exactly

to the AMP moiety of AMPylated CDC42 in complex with the FIC2 domain of the IbpA

toxin [5] (**Figure 2A**). Electron-rich density was observed next to AMP in the active site,

corresponding to a calcium ion present in the crystallization solution to the exclusion of all

other metal ions (**Figure S1B**). Ca²⁺ has 6 coordinations with distances in the expected 2.1-

2.9 Å range, arranged with heptahedral geometry in which one ligand, which would be

located opposite to one phosphate oxygen, is missing. It interacts with the phosphate of AMP,

with the acidic residue in the FIC motif (Glu115), and with the inhibitory glutamate (Glu190)

through a water molecule (**Figure 2B**). The position of Ca²⁺ in the EfFIC^WT-AMP structure

differs from that of Mg²⁺ observed in other FIC protein structures in complex with ATP

(**Figure 2C**), raising the intriguing issue that it may have an alternative catalytic function.

Inspired by the recent observation that animal FICD proteins have deAMPylation enzymatic

activity [18,24], we analyzed whether EfFIC would have deAMPylation activity in the presence

of Ca²⁺. Remarkably, addition of Ca²⁺ to EfFIC^WT that had been previously autoAMPylated in

the presence of Mg²⁺ and [α-³²P]-ATP induced conspicuous deAMPylation (**Figure 2D**).

In the above setup, the AMPylation and deAMPylation activities are potentially acting

concurrently. To characterize the deAMPylation reaction selectively, the hyperactive

EfFIC^E190G mutant was autoAMPylated in the presence of Mg²⁺, purified to remove ATP, PPi

and Mg²⁺ such that no AMPylation remains possible, then its deAMPylation was triggered by

addition of EfFIC^WT or EfFIC mutants in the presence of Ca²⁺. The level of AMPylated EfFIC

(denoted ^AMP-FAMEfFIC) was quantified by fluorescence using ATP-FAM, an ATP analog

fluorescently labeled on the adenine base. Robust deAMPylation was observed upon addition

of EfFIC^WT and Ca²⁺ (**Figure 2E**, EfFIC^WT panel). No spontaneous deAMPylation of ^AMP-

^FAMEfFIC^E190G was observed in the absence of EfFIC^WT (**Figure 2E**, control panel), indicating

that the deAMPylation reaction occurs in trans. We used this deAMPylation setup to identify

residues critical for deAMPylation (**Figure 2E**, mutant panels). Mutation of the catalytic

histidine (H111A) and of the metal-binding acidic residue in the FIC motif (E115A) impaired

deAMPylation of ^AMP-FAMEfFIC^E190G. Likewise, EfFIC^E190G, which carries the mutation of the

inhibitory glutamate, was unable to catalyze deAMPylation, consistent with the absence of

spontaneous deAMPylation in the assay. We conclude from these experiments that EfFIC is a

bifunctional enzyme, that AMPylation and deAMPylation are borne by the same active site,

and that the inhibitory glutamate is involved in the deAMPylation reaction.

The AMPylation and deAMPylation reactions are differentially regulated by metals.

The above results raise the issue of the nature of signals acting on the bifunctional active site

of EfFIC to regulate AMPylation/deAMPylation alternation. Previous work showed that

AMPylation of *Escherichia coli* DNA gyrase by NmFIC, which shares 56% sequence identity

with EfFIC, was highly sensitive to the toxin concentration, with a sharp drop of activity

above 250 µM [21]. We used purified $^{AMP-FAM}$EfFIC^{E190G} to analyze whether the deAMPylation

activity of EfFIC^{WT} would be similarly inhibited by increasing concentrations of EfFIC^{WT} (1-

2000 nM). As shown in **Figure 3A**, deAMPylation increased with EfFIC^{WT} concentration,

indicating that this reaction is not adversely affected by EfFIC concentration. Alternatively,

we reasoned that the distinct electrochemical properties of Ca^{2+} and Mg^{2+} (reviewed in [25])

may allow them to support AMPylation and deAMPylation differentially. Remarkably, Ca^{2+}

was unable to support AMPylation, contrary to Mg^{2+} (**Figure 3B**). In contrast, both Mg^{2+} and

Ca^{2+} supported potent deAMPylation (**Figure 3C, left panel**). Importantly, mutation of the

inhibitory glutamate eliminated the ability of EfFIC to use Ca^{2+} for deAMPylation, while the

mutant retained partial deAMPylation in the presence of Mg^{2+} (**Figure 3C, right panel**). To

understand how Ca^{2+} affects AMPylation and deAMPylation differentially, we determined the

crystal structure of EfFIC^{WT}-ATPγS-Ca^{2+}. In this structure, Ca^{2+} is heptacoordinated to the α-

and β-phosphates of ATPγS (of which again only the ADP moiety is visible), to the inhibitory

glutamate and to 4 water molecules (**Figure 3D**). Strikingly, Ca^{2+} is shifted by 3.2 Å from

Mg^{2+} bound to NmFIC in an equivalent configuration [7] (**Figure 3E**) and it lacks an interaction

with the acidic residue from the FIC motif, which explains why it inhibits AMPylation.

Together, these observations point towards a 3-position metal switch differentially operated

by Mg^{2+} and Ca^{2+}, which includes a position competent for AMPylation, a position that

inhibits AMPylation and a position competent for deAMPylation. To test this hypothesis, we

measured the apparent AMPylation efficiency of EfFIC^{WT} at different Mg^{2+}/Ca^{2+} ratio. As

shown in **Figure 3F**, AMPylation is prominent when Mg^{2+} exceeds Ca^{2+}, while Ca^{2+} in excess

over Mg^{2+} favors deAMPylation. We conclude from these experiments that competition

between Mg^{2+} and Ca^{2+} regulates the balance between AMPylation and deAMPylation

efficiencies and that this regulatory metal switch is implemented by differential usage of the

inhibitory glutamate and the acidic residue in the FIC motif for binding metals.

Ca²⁺ tunes deAMPylation of the BiP chaperone by human FICD.

DeAMPylation of the BiP chaperone has been recently identified as the primary in vitro

activity of human FICD [18], which features a glutamate structurally equivalent to the inhibitory

glutamate in EfFIC (see **Figure 1B**, [10]) that is critical for deAMPylation [18]. We analyzed

whether, as observed in EfFIC, Mg^{2+} and Ca^{2+} metals could also affect FICD activity, using

fluorescent ATP-FAM to monitor BiP AMPylation. No measurable AMPylation of BiP by

FICD^{WT} was observed, neither with Mg^{2+} nor Ca^{2+}, although FICD^{WT} itself showed weak

autoAMPylation in the presence of both metals (**Figure 4A**). Alternatively, we used

FICD^{E234G}, in which the conserved glutamate is mutated to glycine, to produce AMPylated

BiP. Remarkably, while purified ^{AMP-FAM}BIP was efficiently deAMPylated by FICD^{WT} in the

presence of Mg^{2+}, no deAMPylation was measured in the presence of Ca^{2+} (**Figure 4B**). To

determine whether FICD does not bind Ca^{2+} or is unable to use it for deAMPylation, we

carried out an Mg^{2+}/Ca^{2+} competition experiment in which FICD^{WT} and purified ^{AMP-FAM}BiP

were incubated at increasing Ca^{2+} concentration and a fixed Mg^{2+} concentration. As shown in

Figure 4C, deAMPylation efficiency decreased as the Ca^{2+}/Mg^{2+} ratio increased, suggesting

that Ca^{2+} inhibits deAMPylation by competing with Mg^{2+}. We conclude from these

experiments that Ca^{2+} binds to FICD in an inhibitory manner, which allows it to tune the

deAMPylation efficiency of FICD towards the BiP chaperone.

Discussion

In this study, we sought after a diffusible signal able to regulate directly the large group of

AMPylating FIC proteins in which the AMPylation activity is repressed by a conserved

glutamate within the active site. Combining crystallography and PTM assays, we first show

that bacterial EfFIC is a bifunctional enzyme that encodes AMPylating and deAMPylating

activities and that both reactions use the same active site. Next, we discover that the balance

between these opposing activities is controlled by a metal switch, in which each reaction is

differentially supported and inhibited by Mg^{2+} and Ca^{2+} in a manner that the Mg^{2+}/Ca^{2+} ratio

determines the net AMPylation level. Furthermore, we identify the inhibitory glutamate and

the acidic residue in the FIC motif as residues essential for the metal switch. Finally, we show

that deAMPylation of the endoplasmic reticulum BiP chaperone by human FICD is also

dependent on the Ca^{2+}/Mg^{2+} ratio, with high Ca^{2+} concentration inhibiting deAMPylation.

The identification of a potent deAMPylation activity in a bacterial FIC protein (this study)

and in human FICD [18], which depends on a structurally equivalent glutamate in these

otherwise remotely related FIC proteins, leads us to propose that the conserved glutamate is a

signature of the ability of FIC proteins to catalyze both AMPylation and deAMPylation. Our

data allow to delineate the catalytic basis for this bifunctionality, in which catalytic residues

are shared by the AMPylation and deAMPylation reactions but have different roles in

catalysis. In the AMPylation reaction, the invariant histidine in the FIC motif activates the

acceptor hydroxyl of a target protein by attracting a proton and the acidic residue (Asp or Glu)

in the FIC motif binds a metal that stabilizes the phosphates of the ATP cofactor at the

transition state (reviewed in [2,3]). Based on the observations that the α- and β-phosphates of

ATP bind with canonical positions in wild-type EfFIC and that AMPylation is potentiated by

mutation of the glutamate in various FIC proteins, we propose that the primary role of the

glutamate in AMPylation is to mitigate the efficiency of this reaction in the presence of Mg^{2+},

rather than to fully repress it, possibly in order to match AMPylation and deAMPylation

efficiencies. In the deAMPylation mechanism depicted in **Figure 5** (see also discussion in

Supplementary data and Figure S2), the conserved glutamate activates a water molecule for

nucleophilic attack of the phosphorus of the AMP moiety, and the invariant histidine

generates the free hydroxyl group in the protein residue by giving up a proton, as also

proposed in [18]. The nucleophilic water molecule is readily identified as the missing ligand in

the coordination sphere of Ca^{2+} in our EfFIC-AMP-Ca^{2+} structure, where it would be

precisely positioned for in-line nucleophilic attack (**see Figure 2B**). Interestingly, the metal is

bound to both the acidic residue of the FIC motif and the conserved glutamate in this

structure, which positions it to stabilize negative charges in the AMPylated substrate that

develop at the transition state.

Importantly, our data identify for the first time a diffusible signal able to modulate the activity

of glutamate-bearing FIC proteins. A remarkable feature in the above bifunctional mechanism

is that both reactions can be adversely regulated by a second metal that competes with the

catalytic metal. In EfFIC, we observed that Ca^{2+} binds to ATP in a position that is shifted with

respect to the canonical AMPylation Mg^{2+}-binding site, resulting in decreased AMPylation. In

a related scenario, Ca^{2+} competes with Mg^{2+} in the deAMPylation reaction catalyzed by

FICD, thereby decreasing deAMPylation. This multi-position metal switch constitutes a new

paradigm in bifunctional enzyme regulation, in which the relative affinities of specific metals

for the AMPylation and deAMPylation configurations tip the balance towards opposing

activities within the same active site. Future studies are now needed to determine the

bifunctionality spectrum of glutamate-bearing FIC proteins resulting from variations in metal

specificities and affinities. In addition, observations by us and others that FICD has distinct

AMPylation and deAMPylation patterns towards itself and BiP suggests that the protein

substrate influences the AMPylation/deAMPylation balance through still unknown

mechanisms, which will have to be investigated. Another important question for future studies

is how other levels of regulation, such as autoAMPylations and changes in oligomerization

that have been described for a close homolog of EfFIC [21], combine with the intrinsic metal

switch identified in this study.

Our findings predict that bacterial glutamate-containing FIC proteins are bifunctional

enzymes able to switch between AMPylation and deAMPylation activities upon changes in

metal homeostasis. An appealing hypothesis is thus that glutamate-bearing FIC proteins

function as toxin/antitoxin (TA) modules, in which the toxin and antitoxin activities would be

encoded within the same active site under the control of regulatory metals. They would

therefore differ from type III TA modules, in which the toxin is repressed by direct interaction

with the antitoxin, or type IV TA modules, in which the effects of the toxin are reverted by

the antitoxin without direct interaction (reviewed in [26]). It is interesting to note that the

antitoxin component of TA modules such as Bartonella VbhT/VbhA features a structurally

equivalent glutamate that points into the catalytic site of the toxin [7], raising the question of

whether there may exist conditions where the antitoxin glutamate implements deAMPylation

within the TA complex. The physiological conditions that involve variations in metals

homeostasis to activate the switch between opposing activities in these FIC proteins, and how

individual bacteria exploit the bifunctionality of their FIC proteins within their ecological

niche or in infections will be important issues to address in future studies.

Finally, our discovery that deAMPylation of the BiP chaperone by human FICD is modulated

by changes in Ca^{2+} concentration raises important questions with respect to the regulation and

role of FICD in endoplasmic reticulum (ER) functions. The ER is the major organelle

involved in Ca^{2+} homeostasis (reviewed in [27-29]), and the total Ca^{2+} concentration in the ER is

as high as 2-3 mM under resting conditions [30]. Disruption of Ca^{2+} homeostasis, such as

depletion of the ER Ca^{2+} store induced by the drug thapsigargin, swiftly alters protein folding

processes and activates the unfolded protein response [31]. Large Ca^{2+} fluctuations are therefore

considered as major determinants of ER stress responses (reviewed in [28,29]). On the other

hand, there is currently no evidence that the Mg^{2+} concentration changes significantly in the

ER over time (reviewed in [32]). Thus, while keeping in mind that Ca^{2+} and Mg^{2+} concentrations

and local gradients in the ER have remained difficult to determine with accuracy, an

essentially constant Mg^{2+} concentration and Ca^{2+} fluctuations in the 10μM-3mM range, such

as those used in our vitro assays, are plausibly encountered in the ER as it experiences

transition from resting to stress conditions. Recently, FICD has been demonstrated to

stimulate the activity of the BiP chaperone in response to an increase in the unfolded protein

load [13-15,24], and this activation was correlated to deAMPylation of BiP by FICD [18]. It is thus

tempting to speculate that inhibition of FICD deAMPylation activity by high Ca^{2+}, which we

observe *in vitro*, reflects its inhibition under ER homeostasis, where Ca^{2+} concentration is

high and BiP activity is low. Conversely, depletion of Ca^{2+} induces ER stress and triggers the

UPR. Depletion of Ca^{2+} may thus release inhibition of FICD deAMPylation activity, leading

to efficient deAMPylation of BiP and up-regulation of its activity, which is a key feature of

the UPR. We propose that FICD is has features of an enzymatic sensor of Ca^{2+} in the ER,

which could allow it to function as an integrator between Ca^{2+} homeostasis in the ER and the

BiP-mediated unfolded protein response.

In conclusion, we have identified Ca^{2+} as a diffusable signal that modulates the intrinsic

enzymatic activity of glutamate-bearing FIC proteins directly and tips the balance between

AMPylation and deAMPylation reactions without a conformational change in the enzyme.

Future studies are now needed to investigate how the metal switch of glutamate-bearing FIC

protein activities is exploited in bacterial stress, and, in the case of animal FICD, its role in the

unfolded protein response and its crosstalks with Ca^{2+}-controled processes in the ER. The

detrimental effects of prolonged UPR in a myriad of diseases (reviewed in [33,34]) makes new

druggable targets highly desirable (reviewed in [35]). Given the established pivotal role of the

BiP chaperone (reviewed in [16]) and of calcium (reviewed in [29]) in this process, the Ca^{2+}-

regulated activity of FICD uncovered in this study may constitute a new druggable target.

Figure Legends

Figure 1 : Structural basis for the AMPylation activty of EfFIC

A : Structure of the EfFIC monomer showing the FIC motif (pink), the C-terminal α-helix

bearing the inhibitory glutamate (orange) and the β-hairpin predicted to bind protein

substrates (cyan). The ADP moiety of ATPγS is shown in sticks.

B : The inhibitory glutamate from EfFIC^{WT} (orange) is structurally equivalent to the

glutamate found in the C-terminus of NmFIC ([7], PDB 2G03) and in the N-terminus of

Bacteroides BtFIC (PDB 3CUC), *Clostridium* CdFIC ([11], PDB 4X2E) and of human

FICD/HYPE ([10], PDB 4U04). Superpositions are done on the structurally highly conserved

FIC motif.

C : Interactions of the inhibitory glutamate with the active site of ATPγS-bound EfFIC^{WT}.

Hydrogen bonds are depicted by dotted lines.

D : The positions of the α- and β-phosphates of ATPγS bound to EfFIC^{WT} (yellow) diverge

from those of ATPγS bound to NmFIC^{WT} (cyan, [7], PDB 3S6A) and of ATP bound to CdFIC

(blue, [11], S31A/E35A mutant, PDB 4X2D) in a non-canonical conformations. Note that only

the ADP moiety of ATPγS is visible in the EfFIC^{WT} and NmFIC^{WT} crystal structures.

E : The α- and β-phosphates of ATPγS bound to EfFIC^{WT} (yellow) superpose well to those of

ATP bound to NmFIC (purple, [7], E186G mutant, PDB 3ZLM) and of CDP-choline bound to

AnkX (green, [8], H229A mutant, PDB 4BET) in a PTM-competent conformation.

F : AutoAMPylation of EfFIC^{WT} and EfFIC^{E190G}. The level of AMPylated proteins (indicated

as ^{AMP*}EfFIC^{WT}) was measured by autoradiography using radioactive [α-^{32}P]-ATP in the

presence of 100 nM Mg^{2+}. The reaction was carried out for one hour for EfFIC^{WT} and five

minutes for EfFIC^{E190G}. The total amount of EfFIC^{WT} measured by Coomassie staining in the

same sample is shown.

Figure 2 : EfFIC is a deAMPylator in the presence of Ca^{2+}

A : AMP bound to EfFIC^{WT} superposes with the AMP moiety of AMPylated CDC42 in

complex with the FIC protein IbpA ([5], PDB 4ITR). Superposition was carried out based on

the FIC motif.

B : Close-up view of Ca^{2+} bound to EfFIC^{WT}-AMP. The incomplete heptagonal coordination

sphere of Ca^{2+} is indicated by dotted lines. Water molecules are shown as red spheres. The

proposed position of the water molecule responsible for in-line nucleophilic attack is
indicated in grey.

C: Ca^{2+} bound to $EfFIC^{WT}$-AMP is shifted by 1.3 Å with respect to Mg^{2+} bound to $NmFIC^{E/G}$-
AMPPNP. The superposition is done on the FIC motif.

D : $EfFIC^{WT}$ has deAMPylation activity in the presence of Ca^{2+}. $EfFIC^{WT}$ was
autoAMPylated with $[\alpha\text{-}^{32}P]$-ATP in the presence 100 nM Mg^{2+} for one hour, then the sample
was incubated for 60 minutes with EDTA alone (1mM) (lane "-") or 5, 30 and 60 minutes
with EDTA (1mM) and an excess of Ca^{2+} (10 mM) (lanes "+"). AMPylation levels were
analyzed by autoradiography (upper panel). The total amount of $EfFIC^{WT}$ in each sample was
measured by Coomassie staining (lower panel).

E : Key residues of the AMPylation active site are required for deAMPylation. $EfFIC^{E190G}$
was auto-AMPylated for one hour in the presence of fluorescently-labeled ATP-FAM and
Mg^{2+}, then $^{AMP\text{-}FAM}EfFIC^{E190G}$ was purified to remove Mg^{2+}, PPi and ATP-FAM.
DeAMPylation was then triggered by addition of wild-type or mutant EfFIC as indicated in
the presence or not of 1mM Ca^{2+}. AMPylation levels (indicated as $^{AMP\text{-}FAM}EfFIC^{E190G}$) after
one hour incubation were analyzed by fluorescence (upper panel). The total amount of EfFIC
proteins (indicated as $EfFIC^{total}$) in each sample was measured by Coomassie staining (lower
panel).

Figure 3 : The Mg^{2+}/Ca^{2+} ratio tunes the balance between EfFIC AMPylation and de-
AMPylation activities

A : DeAMPylation is not inhibited at high EfFIC concentration. Purified $^{AMP\text{-}FAM}EfFIC^{E190G}$
was incubated with increasing concentrations of $EfFIC^{WT}$ for one hour in the presence of
100µM Ca^{2+}. AMPylation levels were measured by fluorescence as in **Figure 2D**.

B : $EfFIC^{WT}$ uses Mg^{2+} but not Ca^{2+} for autoAMPylation. AutoAMPylation was carried out at
1mM Ca^{2+} or Mg^{2+} for one hour and was measured by autoradiography as in **Figure 1F**.

C : The conserved glutamate of EfFIC is required for the usage of both Mg^{2+} and Ca^{2+} for
deAMPylation. Experiments were carried out for one hour as in **Figure 3A** in the presence of
0.5mM EDTA and 3mM of Mg^{2+} or Ca^{2+}. Quantification of deAMPylation efficiencies is
shown on the right panel.

D: Close-up view of Ca^{2+} bound to $EfFIC^{WT}$-ATPγS. The heptagonal coordination sphere of
Ca^{2+} is indicated by dotted lines. Water molecules are shown as red spheres.

E : Ca^{2+} bound to EfFIC^{WT}-ATPγS is shifted with respect to Mg^{2+} bound to AMPylation-

competent NmFIC$^{E/G}$-AMPPNP (PDF 2ZLM).

F: The net AMPylation efficiency of EfFIC^{WT} is controled by the Mg^{2+}/Ca^{2+} ratio. EfFIC^{WT}

was incubated for one hour in the presence of ATP-FAM at fixed Mg^{2+} concentration (1mM)

and increasing concentrations of Ca^{2+}. $^{AMP-FAM}$EfFIC levels were measured by fluorescence.

The steady-state AMPylation efficiency, which results from the relative AMPylation and

deAMPylation rates, is expressed as the percentage of the maximal AMPylation level

obtained with Mg^{2+} alone. All measurements have p-values <0.05 with respect to the

experiment containing Mg^{2+} alone. The total amount of EfFIC^{WT} measured by Coomassie

staining in each sample is shown.

Figure 4: DeAMPylation of BiP by FICD requires Mg^{2+} and is inhibited by Ca^{2+}.

A: FICD^{WT} does not AMPylate the BiP chaperone. Reactions were carried out for one hour in

presence of 1mM Mg^{2+} or 1mM Ca^{2+}. The level of AMPylated proteins (indicated as $^{AMP-}$

^{FAM}FICD^{WT} and $^{AMP-FAM}$BIP) was measured by fluorescence. The total amount of proteins

measured by Coomassie staining in the same sample is shown.

B: Ca^{2+} inhibits deAMPylation of BiP by FICD/HYPE^{WT}. BiP was first AMPylated for one

hour by the hyperactive FICD^{E234G} mutant in the presence of 100 μM Mg^{2+}, then $^{AMP-FAM}$BIP

was purified to remove Mg^{2+}, PPi and ATP-FAM. DeAMPylation of $^{AMP-FAM}$BIP was then

triggered by addition of wild-type or mutant FICD as indicated in the presence of 1mM Mg^{2+}

or 1mM Ca^{2+}. $^{AMP-FAM}$BiP levels were measured by fluorescence. The total amount of BiP

measured by Coomassie staining in each sample is shown.

C: The net deAMPylation efficiency of FICD^{WT} is tuned by the Mg^{2+}/Ca^{2+} ratio.

DeAMPylation was carried out as in **Figure 4B** using a fixed Mg^{2+} concentration (200μM)

and 0, 1, 2 or 3mM Ca^{2+}. AMPylation levels were measured by fluorescence, normalized to

the fluorescence intensity of FICD and expressed as the percentage of maximal

deAMPylation level obtained in the absence of Ca^{2+}. All data have p-values <0.05 with

respect to the control in the absence of Ca^{2+}. The total amount of BiP measured by Coomassie

staining in each sample is shown.

Figure 5: Model of the Ca^{2+}-assisted deAMPylation catalytic mechanism.

In this model, the regulatory glutamate (Glu190 in EfFIC) attracts a proton from a water

molecule coordinating the metallic cation (1) to activate it for nucleophilic attack of its

oxygen on the phosphorus of AMP moiety in the AMPylated substrate (2). The positive

charge provided by Ca^{2+} increases electrophilicity of the phosphorus and stabilizes the

negative charge of the intermediate (2 and 3). The intermediate harboring a pentavalent

phosphorus then rearranges, leading to the breaking of the phosphor-ester bond, which is

elicited by the capture of a proton provided by the catalytic histidine (4). R : AMPylated

protein.

Material and Methods.

Protein cloning, expression and purification

The codon-optimized gene encoding full-length *Enterococcus faecalis* EfFIC with an N-

terminal 6-histidine tag was from GeneArt Gene Synthesis (ThermoFisher Scientific) and

cloned into a pET22b(+) vector. The codon-optimized gene encoding human FICD/HYPE

(residues 45-459) carrying an N-terminal 6-His tag followed by SUMO tag was from GeneArt

Gene Synthesis and cloned into a pET151/D-TOPO vector (ThermoFisher Scientific). All

mutations were performed with the QuickChange II mutagenesis kit (Agilent). *Mus musculus*

BIP in pUJ4 plasmid is a kind gift from Ronald Melki (CNRS, Gif-sur-Yvette). All constructs

were verified by sequencing (GATC). All EfFIC constructs were expressed in *E. coli* BL21

(DE3) pLysS in LB medium. Overexpression was induced overnight with 0.5 mM IPTG at

20°C. Bacterial cultures were centrifuged for 40 min at 4000g. Bacterial pellets were

resuspended in lysis buffer (50 mM Tris-HCl pH 8.0, 150 mM NaCl, 5% glycerol,

0.25 mg/mL lysozyme) containing a protease inhibitor cocktail, disrupted at 125 psi using a

high pressure cell disrupter and centrifuged 30 min at 22000g. The cleared lysate supernatant

was loaded on a Ni-NTA affinity chromatography column (HisTrap FF, GE Healthcare) and

eluted with 250 mM imidazole. Purification was polished by gel filtration on a Superdex 200

16/600 column (GE Healthcare) equilibrated with storage buffer (50 mM Tris-HCl pH 8.0,

150 mM NaCl). Wild-type and mutant FICD/HYPE were expressed and purified as EfFIC,

except that the lysis buffer was complemented with 1mM DTT and 0.02% Triton X-100 and

other buffers with 1mM DTT. To remove the SUMO tag, FICD/HYPE was incubated with

SUMO protease (ThermoFischer) at 1/100 weight/weight ratio during 1 hour at room

temperature. The cleaved fraction was separated by affinity chromatography (HisTrap FF, GE

Healthcare) and further purified by gel filtration on a Superdex 200 10/300 column (GE

Healthcare) equilibrated with storage buffer (50 mM Tris pH 8.0, 150 mM NaCl, 1mM DTT,

5% glycerol). Mouse BIP was expressed and purified essentially as EfFIC.

Crystallization and structure determination

A summary of the crystal structures determined in this study is in **Table 1**. Proteins were

crystallized using a TTP Labtech's Mosquito LCP crystallization robot and crystallization

screens (Jena Bioscience and Quiagen). Conditions leading to crystals were subsequently

optimized. Diffraction data sets were recorded at synchrotron SOLEIL and ESRF. Datasets

were processed using XDS [36], xdsme (https://github.com/legrandp/xdsme) or autoProc [37].

Structures were solved by molecular replacement and refined with the Phenix suite [38] or

Buster (Bricogne G., Blanc E., Brandl M., Flensburg C., Keller P., Paciorek W.,Roversi P,

Sharff A., Smart O.S., Vonrhein C., Womack T.O. (2017). BUSTER version 2.10.2.

Cambridge, United Kingdom: Global Phasing Ltd.). Models were build using Coot [39].

Softwares used in this project were curated by SBGrid [40]. Crystallization conditions, data

collection statistics and refinement statistics are given in **Table S1**. All structures have been

deposited with the Protein Data Bank (PDB codes in **Table S1**).

AMPylation and deAMPylation assays

AMPylation and deAMPylation autoradiography assays were carried out using the following

protocols. For AMPylation reactions, 8 μg of purified proteins were mixed with 10 μCi [α-

^{32}P] ATP (Perkin Elmer) in a buffer containing 50mM Tris-HCl pH 7.4, 150 mM NaCl and

0.1 mM MgCl$_2$. Reactions were incubated for 1 h at 30 °C, then stopped with reducing SDS

sample buffer and boiling for 5 min. For deAMPylation, AMPylation was performed as

above, then 1 mM EDTA was added with or without 10 mM CaCl$_2$. Proteins were resolved by

SDS-PAGE and AMPylation was revealed by autoradiography.

EfFIC AMPylation and deAMPylation fluorescence assays were carried out using the

following protocols. AMPylation was carried out using a fluorescent ATP analog modified by

N^6-(6-Amino)hexyl on the adenine base (ATP-FAM, Jena Bioscience). AMPylated proteins

were obtained by incubation for one hour at 30 °C in 50mM Tris pH 8.0, 150 mM NaCl,

0.1 mM MgCl$_2$ and an equimolar amount of ATP-FAM. Before deAMPylation reactions, the

buffer was exchanged to 50 mM Tris-HCl pH 8.0 and 150 mM NaCl by 5 cycles of

dilution/concentration on a Vivaspin-500 with a cut-off of 10kDa (Sartorius), resulting in a

final dilution of ATP-FAM, MgCl$_2$ and PPi by about 10^5 times. DeAMPylation reactions

were carried out using 3 μM of AMPylated protein and 6 μM of freshly purified EfFIC

proteins (except for the experiment depicted in Figure 3A) in a buffer containing 50mM Tris-
HCl pH 8.0 and 150 mM NaCl, for 1h at 30°C. Reactions were stopped by addition of
reducing SDS sample buffer and boiling for 5 min. Proteins were resolved by SDS-PAGE and
modification by AMP-FAM was revealed by fluorescence using green channel (excitation:
488 nm, emission: 526 nm) on a Chemidoc XR+ Imaging System (BioRad).

FICD AMPylation and deAMPylation fluorescence assays were carried out using the
following protocols. AMPylation was carried out using fluorescent ATP-FAM. AMPylated
BIP was obtained by incubation for one hour at 30 °C in 50mM Tris pH 8.0, 150 mM NaCl,
0.1 mM $MgCl_2$, 2µM $FICD^{E234G}$ and an equimolar amount of ATP-FAM. Before
deAMPylation reactions, the buffer was exchanged with 50mM Tris-HCl pH 8.0 and 150 mM
NaCl by 5 cycles of dilution/concentration on a Vivaspin-500 with a cut-off of 50kDa
(Sartorius), resulting in a final dilution of ATP-FAM, $MgCl_2$ and PPi produced by the
reaction of about 10^5 times. DeAMPylation reactions were carried using 2 µg of AMPylated
protein and 4 µg of freshly purified $FICD^{WT}$ in a buffer containing 50mM Tris pH 8.0 and
150 mM NaCl, for 1h at 30°C. Reactions were stopped and ATP-FAM modification revealed
as EfFIC. Quantification of AMP-FAM levels was done using (ImageLab, BioRad).

All experiments were done at least in triplicate, except kinetics in Figure 3A that were done in
duplicate.

Acknowledgements.

This work was supported by grants from the Fondation pour la Recherche Médicale and from
the Agence Nationale pour la Recherche to J.C. and by grants n°ANR-10-LABX-62-IBEID
and from the Fondation pour la Recherche Médicale to C.B. S.V. was supported by a PhD
grant from the DIM MALINF and G.O. by a stipend from the Pasteur-Paris University
International PhD program. We are grateful to the scientific teams at the PX1 and PX2
beamlines at the SOLEIL synchrotron (Gif-sur-Yvette, France) and from the ID29, ID30-A3
and ID30B beamlines at the European Synchrotron Research Facility (ESRF, Grenoble,
France) for their expertise and advice. We thank Pascale Serror (INRA, Jouy-en-Josas,
France) and Philippe Glaser (Institut Pasteur) for discussions and the members of the Cherfils
lab for help and shared expertise.

Table 1 – Summary of crystal structures of EfFIC determined in this study. All crystals

have cell angles of α, β and γ = 90°. Crystallographic statistics are given in **Table S1**. * : only

the ADP moiety is visible.

Form	Space group	Cell parameters a, b, c (Å)	Ligand	Resolution	PDB
WT	P4$_1$2$_1$2	65.13, 65.13, 248.06	PO_4^{2-}	2.29	6ER8
WT	I222	121.54, 131.00, 136.94	/	2.40	5NV5
WT	P4$_1$2$_1$2	64.98 64.98 246.24	AMP – Ca^{2+}	2.35	6EP0
WT	P4$_3$2$_1$2	125.35 125.35 362.8	ATPγS* – Ca^{2+}	2.15	6EP2
WT	P4$_3$2$_1$2	87.84 87.84 364.94	ATPγS*	1.93	6EP5
H118A	P2$_1$22$_1$	76.67 77.11 103.15	/	2.60	5NWF
H118A	I222	121.93 131.16 136.71	SO_4^{2-}	2.20	5NVQ

547   **References**

549   1.   Kinch, L.N., Yarbrough, M.L., Orth, K. & Grishin, N.V. Fido, a novel AMPylation
550   domain common to fic, doc, and AvrB. *PLoS One* **4**, e5818 (2009).

551   2.   Garcia-Pino, A., Zenkin, N. & Loris, R. The many faces of Fic: structural and
552   functional aspects of Fic enzymes. *Trends Biochem Sci* **39**, 121-9 (2014).

553   3.   Roy, C.R. & Cherfils, J. Structure and function of Fic proteins. *Nat Rev Microbiol* **13**,
554   631-40 (2015).

555   4.   Harms, A., Stanger, F.V. & Dehio, C. Biological Diversity and Molecular Plasticity of
556   FIC Domain Proteins. *Annu Rev Microbiol* **70**, 341-60 (2016).

557   5.   Xiao, J., Worby, C.A., Mattoo, S., Sankaran, B. & Dixon, J.E. Structural basis of Fic-
558   mediated adenylylation. *Nat Struct Mol Biol* **17**, 1004-10 (2010).

559   6.   Luong, P. et al. Kinetic and structural insights into the mechanism of AMPylation by
560   VopS Fic domain. *J Biol Chem* **285**, 20155-63 (2010).

561   7.   Engel, P. et al. Adenylylation control by intra- or intermolecular active-site
562   obstruction in Fic proteins. *Nature* **482**, 107-10 (2012).

563   8.   Campanacci, V., Mukherjee, S., Roy, C.R. & Cherfils, J. Structure of the Legionella
564   effector AnkX reveals the mechanism of phosphocholine transfer by the FIC domain.
565   *EMBO J* **32**, 1469-77 (2013).

566   9.   Castro-Roa, D. et al. The Fic protein Doc uses an inverted substrate to phosphorylate
567   and inactivate EF-Tu. *Nat Chem Biol* **9**, 811-7 (2013).

568   10.   Bunney, T.D. et al. Crystal structure of the human, FIC-domain containing protein
569   HYPE and implications for its functions. *Structure* **22**, 1831-43 (2014).

570   11.   Dedic, E. et al. A Novel Fic (Filamentation Induced by cAMP) Protein from
571   Clostridium   difficile   Reveals   an   Inhibitory   Motif-independent
572   Adenylylation/AMPylation Mechanism. *J Biol Chem* **291**, 13286-300 (2016).

573   12.   Yarbrough, M.L. et al. AMPylation of Rho GTPases by Vibrio VopS disrupts effector
574   binding and downstream signaling. *Science* **323**, 269-72 (2009).

575   13.   Sanyal, A. et al. A novel link between Fic (filamentation induced by cAMP)-mediated
576   adenylylation/AMPylation and the unfolded protein response. *J Biol Chem* **290**, 8482-
577   99 (2015).

578   14.   Ham, H. et al. Unfolded protein response-regulated Drosophila Fic (dFic) protein
579   reversibly AMPylates BiP chaperone during endoplasmic reticulum homeostasis. *J*
580   *Biol Chem* **289**, 36059-69 (2014).

15.   Preissler, S. et al. AMPylation matches BiP activity to client protein load in the
endoplasmic reticulum. *Elife* **4**, e12621 (2015).

16.   Zhao, L. & Ackerman, S.L. Endoplasmic reticulum stress in health and disease. *Curr*
Opin Cell Biol **18**, 444-52 (2006).

17.   Preissler, S. et al. AMPylation targets the rate-limiting step of BiP's ATPase cycle for
its functional inactivation. *Elife* **6**(2017).

18.   Preissler, S., Rato, C., Perera, L.A., Saudek, V. & Ron, D. FICD acts bifunctionally to
AMPylate and de-AMPylate the endoplasmic reticulum chaperone BiP. *Nat Struct*
Mol Biol **24**, 23-29 (2017).

19.   Das, D. et al. Crystal structure of the Fic (Filamentation induced by cAMP) family
protein SO4266 (gi|24375750) from Shewanella oneidensis MR-1 at 1.6 A resolution.
Proteins **75**, 264-71 (2009).

20.   Veyron, S., Peyroche, G. & Cherfils, J. FIC proteins: from bacteria to humans and
back again. *Pathog Dis* **76**(2018).

21.   Stanger, F.V. et al. Intrinsic regulation of FIC-domain AMP-transferases by
oligomerization and automodification. *Proc Natl Acad Sci U S A* **113**, E529-37 (2016).

22.   Hegstad, K., Mikalsen, T., Coque, T.M., Werner, G. & Sundsfjord, A. Mobile genetic
elements and their contribution to the emergence of antimicrobial resistant
Enterococcus faecalis and Enterococcus faecium. *Clin Microbiol Infect* **16**, 541-54
(2010).

23.   Lebreton, F. et al. Tracing the Enterococci from Paleozoic Origins to the Hospital.
Cell **169**, 849-861 e13 (2017).

24.   Casey, A.K. et al. Fic-mediated deAMPylation is not dependent on homodimerization
and rescues toxic AMPylation in flies. *J Biol Chem* **292**, 21193-21204 (2017).

25.   Carafoli, E. & Krebs, J. Why Calcium? How Calcium Became the Best
Communicator. *J Biol Chem* **291**, 20849-20857 (2016).

26.   Harms, A., Brodersen, D.E., Mitarai, N. & Gerdes, K. Toxins, Targets, and Triggers:
An Overview of Toxin-Antitoxin Biology. *Mol Cell* **70**, 768-784 (2018).

27.   Bravo, R. et al. Endoplasmic reticulum and the unfolded protein response: dynamics
and metabolic integration. *Int Rev Cell Mol Biol* **301**, 215-90 (2013).

28.   Krebs, J., Agellon, L.B. & Michalak, M. Ca(2+) homeostasis and endoplasmic
reticulum (ER) stress: An integrated view of calcium signaling. *Biochem Biophys Res*
Commun **460**, 114-21 (2015).

29. Carreras-Sureda, A., Pihan, P. & Hetz, C. Calcium signaling at the endoplasmic
reticulum: fine-tuning stress responses. *Cell Calcium* **70**, 24-31 (2018).

30. Montero, M. et al. Monitoring dynamic changes in free Ca2+ concentration in the
endoplasmic reticulum of intact cells. *EMBO J* **14**, 5467-75 (1995).

31. Wong, W.L., Brostrom, M.A., Kuznetsov, G., Gmitter-Yellen, D. & Brostrom, C.O.
Inhibition of protein synthesis and early protein processing by thapsigargin in cultured
cells. *Biochem J* **289 (Pt 1)**, 71-9 (1993).

32. Romani, A.M. Cellular magnesium homeostasis. *Arch Biochem Biophys* **512**, 1-23
(2011).

33. Xu, C., Bailly-Maitre, B. & Reed, J.C. Endoplasmic reticulum stress: cell life and
death decisions. *J Clin Invest* **115**, 2656-64 (2005).

34. Morris, G. et al. The Endoplasmic Reticulum Stress Response in Neuroprogressive
Diseases: Emerging Pathophysiological Role and Translational Implications. *Mol*
Neurobiol (2018).

35. Maly, D.J. & Papa, F.R. Druggable sensors of the unfolded protein response. *Nat*
Chem Biol **10**, 892-901 (2014).

36. Kabsch, W. Integration, scaling, space-group assignment and post-refinement. *Acta*
Crystallogr D Biol Crystallogr **66**, 133-44 (2010).

37. Vonrhein, C. et al. Data processing and analysis with the autoPROC toolbox. *Acta*
Crystallogr D Biol Crystallogr **67**, 293-302 (2011).

38. Adams, P.D. et al. PHENIX: a comprehensive Python-based system for
macromolecular structure solution. *Acta Crystallogr D Biol Crystallogr* **66**, 213-21
(2010).

39. Emsley, P. & Cowtan, K. Coot: model-building tools for molecular graphics. *Acta*
crystallographica. Section D, Biological crystallography **60**, 2126-32 (2004).

40. Morin, A. et al. Collaboration gets the most out of software. *Elife* **2**, e01456 (2013).

Figure 1

A

^{AMP}Tyr

AMP

B

E115

Ca^{2+}

AMP

E190

C

Mg^{2+}

Ca^{2+}

NmFIC$^{E/G}$-AMPPNP-Mg^{2+}
EfFIC^{WT}-AMP-Ca^{2+}

D

	60'	5'	30'	60'
Ca^{2+}	−	+	+	+

^{AMP*}EfFIC^{WT}

EfFIC^{WT}

E

	Control		WT		H111A		E115A		E190G	
Ca^{2+}	−	+	−	+	−	+	−	+	−	+

$^{AMP-FAM}$EfFIC^{E190G}

EfFIC^{total}

Figure 2

A

[EfFIC^WT] (nM)   1   10   25   50   100   250   500   1000   2000

^AMP-FAM-EfFIC^E190G

EfFIC^total

B

Mg^2+   Ca^2+

^AMP*EfFIC^WT

EfFIC^WT

C

EfFIC^WT   EfFIC^E190G

EDTA   Mg^2+   Ca^2+   EDTA   Mg^2+   Ca^2+

^AMP-FAMEfFIC^E190G

EfFIC^total

Mg^2   Ca^2+

% deAMPylation

EfFIC^WT   EfFIC^E190G

D

E115

ATPγS

Ca^2+

E190

E

Mg^2+

Ca^2+

NmFIC^E/G-AMPPNP-Mg^2+
EfFIC^WT-ATPγS-Ca^2+

F

Mg^2+:Ca^2+ (mM)   1:0   1:1   1:2   1:3   1:4   1:5

^AMP-FAMEfFIC^WT

EfFIC^WT

% AMPylation

Figure 3

A

Mg²⁺ Ca²⁺

ᴬᴹᴾ⁻ᶠᴬᴹBIP

ᴬᴹᴾ⁻ᶠᴬᴹFICDᵂᵀ

BIP

FICDᵂᵀ

B

FICDᵂᵀ

0   Mg²⁺ Ca²⁺ Mg²⁺ Ca²⁺

ᴬᴹᴾ⁻ᶠᴬᴹBIP

BIP

C

Mg²⁺:Ca²⁺ ratio 1:0   1:5   1:10   1:15

ᴬᴹᴾ⁻ᶠᴬᴹBIP

BIP

% deAMPylation

Figure 4

Figure 5