Complex-centric proteome profiling by SEC-SWATH-MS

Moritz Heusel[1,2,6], Isabell Bludau[1,3,6], George Rosenberger[1], Robin Hafen[1,4], Max Frank[1], Amir Banaei-Esfahani [1,3], Ben C. Collins[1], Matthias Gstaiger[1,*], Ruedi Aebersold[1,5,*].

Institute of Molecular Systems Biology, Department of Biology, ETH Zurich, Zurich

PhD program in Molecular and Translational Biomedicine of the Competence Center Personalized Medicine UZH/ETH, Zurich, Switzerland

PhD Program in Systems Biology, Life Science Zurich Graduate School, University of Zurich and ETH Zurich, Zurich, Switzerland

Department of Computer Science, ETH Zurich

Faculty of Science, University of Zurich, Zurich, Switzerland.

Authors contributed equally

* Corresponding author

Character count: 56.070

Highlights

- Introduction of the concept of complex-centric proteome profiling

- Development of *CCprofiler*, a software framework for complex-centric data analysis

- Detection and quantification of subunit distribution of 462 distinct protein complexes containing 2127 proteins from a SEC-SWATH-MS dataset of HEK293 cells, and identification of novel complex variants such as assembly intermediates

- Statistical target-decoy model to estimate accurate false discovery rates for complexes quantified by complex-centric analysis

- *SECexplorer*, an online platform to support custom complex-centric exploration of SEC-SWATH-MS datasets.

Abstract

Proteins are major effectors and regulators of biological processes that can elicit multiple functions depending on their interaction with other proteins. The organization of proteins into macromolecular complexes and their quantitative distribution across these complexes is, therefore, of great biological and clinical significance.

In this paper we describe an integrated experimental and computational technique to quantify hundreds of protein complexes in a single operation. The method consists of size exclusion chromatography (SEC) to fractionate native protein complexes, SWATH/DIA mass spectrometry to precisely quantify the proteins in each SEC fraction and the computational framework *CCprofiler* to detect and quantify protein complexes by error-controlled, complex-centric analysis using prior information from generic protein interaction maps.

Our analysis of the HEK293 cell line proteome delineates 462 complexes composed of 2127 protein subunits. The technique identifies novel subcomplexes and assembly intermediates of central regulatory complexes while assessing the quantitative subunit distribution across them. We make the toolset *CCprofiler* freely accessible, and provide a web platform, *SECexplorer*, for custom exploration of the HEK293 proteome modularity.

Introduction

Molecular life science research over the last decades has been transformed by technological advances that aim at exploring biological processes as complex systems of interacting molecules. A range of high throughput technologies to analyze genomes, transcriptomes, metabolomes and proteomes now provide accurate molecular inventories of biological samples at high throughput. Yet, the notion of a modular biology[1] states that for the definition of the functional state of a cell the organization of cellular molecules into functional modules is as important as the composition of the respective "omes". This notion has been supported by decades of research into the structure and function of specific macromolecular complexes but the task to systematically probe the organization of biomolecules in the cell has remained technologically challenging. Among all macromolecular modules those containing or consisting of proteins are particularly functionally important because they catalyze and control the vast majority of biochemical functions and constantly adapt to and determine the state of the cell.

For high throughput analytical techniques to generate data sets that are quantitative, reproducible and contain low error rates it has frequently been useful to use prior information to guide the acquisition or analysis of the respective data[2]. For mass spectrometry-based proteomics, the concept of peptide-centric analysis[3] uses reference fragment ion spectra as prior information to detect and quantify proteolytic peptides in complex samples as surrogates for their corresponding proteins. Peptide-centric analyses have been implemented at a moderate level of multiplexing (tens to few hundred proteins) via selected reaction monitoring (SRM[4]) and parallel reaction monitoring (PRM[5]). More recently, massively parallel data independent analysis strategies (DIA) exemplified by SWATH-MS have been developed that reproducibly quantify tens of thousands of peptides from single sample injections into a mass spectrometer[6–8]. In this manuscript we describe and implement the concept of complex centric analysis. It is intended to systematically detect protein complexes in biological samples and to quantify the distribution of proteins across protein complex instances. Complex-centric analysis uses generic protein interaction information as prior information and conceptually extends the principles of peptide-centric analysis to the level of protein complexes.

Complex-centric proteome profiling consists of the robust and proven technique of size exclusion chromatography (SEC) to fractionate native protein complexes, SWATH/DIA mass spectrometry to precisely and reproducibly quantify proteins across SEC fractions and a new computational analysis strategy implemented in *CCprofiler*. *CCprofiler* carries out fast and automated detection of protein complexes in datasets of quantitative protein maps from consecutive SEC fractions and controls error rates by means of a target-decoy based statistical model. It uses prior information from generic protein interaction maps to detect and quantify protein complexes in the sample. Complex-centric

protein profiling is a new implementation of the general concept of protein correlation profiling[9–11]

that distinguishes itself from earlier implementations[12–14] by the following: i) the use of SWATH-MS

for the data generation provides complete protein elution profiles for each detected protein at

quantitative accuracy and a wide dynamic range supporting the quantification of even minor

components of the proteome, ii) the development of a statistical model in *CCprofiler* that uses a

target/decoy model to calculate a FDR for detected complexes, and iii) the use of prior information

from generic protein interaction maps to reduce the erroneous assignment of co-eluting proteins to

a complex.

A range of generic protein complex compendia have been generated by different approaches that

can be used as prior information for complex-centric analysis. They include: i) the CORUM reference

database of complexes[15] generated by curating results from classical biochemical and biophysical

analyses of protein complexes. CORUM presently contains 1753 distinct models of human complexes

consisting of 2532 proteins; ii) The BioPlex network[16] and related protein interaction databases,

generated by the mass spectrometric identification of proteins co-purifying with affinity tagged

"bait" proteins (AP-MS). BioPlex v1.0 describes 23,744 interactions among 7,688 proteins identified

as interactors of 2594 bait proteins; iii) the STRING database[17], an organism-wide protein-protein

interaction network generated by the computational integration of multiple lines of evidence for

physical and functional associations. STRING (v10) contains 383,626 high confidence interactions

(score ≥ 900) among 10,248 human proteins, and iv) protein complex databases generated by

correlation profiling of extensive chromatographic co-fractionation of native complexes, followed by

DDA mass-spectrometry [12–14]. In combination, these interaction compendia constitute an extensive,

yet incomplete representation of the organization of the (human) proteome into functional

complexes and thus provide an essential resource for the implementation of the complex centric

analysis strategy that is supported by the computational framework *CCprofiler*.

We benchmark the method, including the *CCprofiler* algorithm, against a manually curated set of

protein complexes and evaluate its complex identification performance against a reference method

consisting of multidimensional co-fractionation of native extracts and DDA of individual fractions[12].

The results demonstrate high performance of the *CCprofiler* algorithm in relation to manual

benchmarking, with observed true positive rates of up to 91 % (high quality signals) at an FDR of 5 %.

The data further shows superior performance of the complex-centric approach in recalling protein

complexes compared to the reference method, achieved at a significantly reduced experimental

effort (81 vs. 1,163 fractions analyzed by LC-MS/MS). We applied the complex-centric proteome

profiling strategy to quantify complexes in a native extract from HEK293 cells in exponential growth

state. The results indicate that 55 % of the protein mass is present in the form of complexes that

distribute across distinct states of complex formation. The data indicated quantitative complex

signals for 462 cellular assemblies if prior knowledge from the CORUM, BioPlex and StringDB

reference databases was used and the results were cumulatively integrated. The utility of quantifying

the distribution of specific proteins across different resolved submodules is exemplified by the

identification of previously unknown substructures of cellular effector complexes such as the

proteasome. Finally, we describe and provide access to *SECexplorer*, an interactive online platform

for customized expert interpretation of quantitative co-fractionation protein profiles generated by

SEC-SWATH-MS. We expect that the complex centric analysis method, the SEC-SWATH dataset

representing the organization of the proteome of the cycling HEK 293 cell line and the computational

tools to explore the data will find wide application in life science research.

Results

Principles and main features of complex-centric proteome analysis.

We describe an integrated mass spectrometric and computational method to systematically quantify the modular organization of the proteome. The method is schematically illustrated in Figure 1A and consists of five consecutive steps. First, complexes are isolated from a biological sample under mild conditions that retain their native form and fractionated according to their hydrodynamic radius via high-resolution size exclusion chromatography (SEC). Second, collected, consecutive fractions are subjected to bottom-up mass spectrometric analysis using SWATH/DIA mass spectrometry. Collectively, the thus generated 81 SWATH/DIA maps constitute the dataset that will ultimately be explored by complex-centric analysis of protein SEC elution profiles (Step 5). To accurately quantify protein elution along the SEC chromatographic fractions, peptides are identified and quantified from the composite SWATH/DIA data set in step three by peptide-centric analysis[7,18,19]. Specifically, peptide query parameters for tens of thousands of peptides are generated from a reference spectral library and systematically queried across the dataset to quantify each target peptide in each fraction. Fourth, *CCprofiler* is used to infer quantitative protein elution profiles from the peptide elution profiles across SEC fractions (for details on peptide detection and protein inference along the chromatographic fractions see Figure S1). Fifth, the protein SEC elution profiles are explored via complex-centric analysis using *CCprofiler* along with prior protein interaction information, to detect distinct protein modules and to determine the likelihood that each detected module is correctly identified. Specifically, the complex-centric analysis of *CCprofiler* in steps four and five entails (Fig 1B) (i) protein quantification, (ii) target complex query set generation based on prior protein connectivity information, (iii) the generation of corresponding decoy complex query sets used for downstream error estimation, (iv) detection of complex component subunit co-elution signals along SEC fractions, (v) decoy-based generation of a null model and according error estimation and, (vi) compilation of the results into a report detailing unique, chromatographically resolved instances of complexes and the distribution of shared protein subunits across them (For details, see extended experimental procedures).

Benchmarking and performance assessment

We evaluated the performance of the described complex-centric analysis method, (i) by benchmarking the *CCprofiler* algorithm and error model against a manually curated reference data set, (ii) by comparing its performance with the performance of a reference method consisting of multidimensional co-fractionation of native complexes and the proteomic analysis of 1,163 fractions by data dependent mass spectrometry[12], and (iii) by demonstrating increased sensitivity for complex

detection as a result of the improved consistency of quantification of SWATH/DIA compared to data-

dependent acquisition-based mass spectrometry (Figure 2).

Using the data generated from the HEK293 cell line proteome, we first benchmarked the automated

performance of complex-centric analysis and FDR estimation by *CCprofiler* against a manually curated

reference set (Figure 2A). The manual reference set was generated by manually testing protein

complexes reported in the CORUM knowledgebase[15] for evidence of complete or partial co-elution

signals among the protein level SEC chromatograms of the respective protein subunits (criteria: ≥ 2

proteins show at least one chromatographic co-elution peak, as judged by visual expert inspection).

Taking the manually annotated co-elution signals as ground truth, a false discovery rate (FDR) of the

complex detection in *CCprofiler* could be estimated based on the number of automatically detected

complex signals that were not confirmed by the manual reference set (manual FDR, also see

extended experimental procedures). This manual reference based FDR was compared to the

independent FDR estimation by the target-decoy approach (decoy FDR), demonstrating that the

target-decoy model provides accurate or slightly conservative error estimates of the algorithm

(Figure 2A). To further evaluate the sensitivity of *CCprofiler*, we tested the recall of manually

detected signals by the automated analysis. At 5% decoy-estimated FDR, the automated *CCprofiler*

analyses recalled 91% of the high-confidence manual signals and 76% of all manually annotated

signals (Figure 2A and Figures S2A/B).

Second, we compared the performance of the complex-centric analysis method with the

performance of a reference *de novo* complex analysis method implemented by Havugimana et al.

which is based on multidimensional fractionation of native complexes isolated from HEK293 and

HeLa cells[12] (Figure 2B). As a metric, we evaluated the ability of either method to recall complexes

reported in the CORUM knowledgebase which consists of a total of 1753 non-redundant complexes.

We considered a complex as recalled if at least 50 % of its CORUM annotated protein subunits were

stated as part of a reported complex by either method (For details, see extended experimental

procedures). The comparison comprised all 622 reported complexes from Havugimana et al. with

unknown error rate, compared to the collapsed set of 462 complexes derived from complex-centric

analysis of our HEK293 SEC-SWATH-MS dataset based on prior information from CORUM, BioPlex

and StringDB, each independently filtered for 5% FDR. The results show that the complex-centric

analysis method recalled 731 complexes from 81 fractions generated by single dimension SEC,

compared to 561 complexes recalled from 1,163 fractions by the multidimensional reference method

[12] (Figure 2B). The results also show a large agreement (438) between the recovered CORUM

complexes. However, both datasets also uniquely recall parts of the CORUM complexes (123

complexes were uniquely confirmed by Havugimana et al. and 293 by our workflow, respectively).

Due to a lack of ground truth in terms of the set of biologically present protein complexes in each

respective dataset, ultimate conclusions on the correctness of each set of reported complexes

remains challenging. However, these results demonstrate that a single-fractionation mass

spectrometry dataset analyzed by complex-centric proteome profiling can retrieve comparable, if not

more comprehensive information on the protein complex landscape, as compared to previous

multidimensional fractionation efforts including a fourteen times higher number of sample injections

coupled to *de novo* complex analysis.

Third, to assess the contribution of SWATH/DIA quantification to the favorable recall results of the

complex-centric proteome profiling workflow, we compared results obtained by SWATH/DIA based

protein quantification with those obtained by MS1 signal integration or spectral counting when the

same samples were analyzed by DDA. To generate the DDA dataset, aliquots of the peptide samples

of the 81 SEC fractions analyzed by SWATH/DIA were also analyzed by data-dependent acquisition on

the same TOF model 5600 mass spectrometer that was also used for SWATH/DIA acquisition. Results

are shown in Figure 2C. At a respective protein-level FDR control of 1 %, SWATH/DIA quantifies 4916

proteins across the SEC fractions ($\geq$ 2 independent proteotypic peptides), whereas the DDA data only

covers 4176 proteins when analyzed by MS1 quantification based on the top2 intensity sum, and

4497 proteins when quantified by spectral counting (for details on the respective data analysis

strategies see extended experimental procedures). To further assess the differences between DIA

and DDA quantification, we next analyzed the three datasets with respect to the consistency of

protein detection and quantification along consecutive SEC fractions (Figure 2C, left panel). The

results indicate that SWATH/DIA detects and quantifies a substantially higher number of proteins in

three or more consecutive fractions compared to DDA based analyses. Next, the precision of

quantification as judged by global correlation among quantitative profiles of peptides originating

from the same parent protein was compared between SWATH/DIA and DDA quantification, showing

favorable quantification precision for SWATH/DIA (Figure 2C, middle panel). Finally, the DIA and DDA

datasets were compared by their performance in detecting protein complexes (Figure 2C). At 5%

controlled FDR, complex-centric analysis provides co-elution evidence for 621 versus 298 and 343 of

the CORUM set of query complexes from quantitative data from SWATH-MS2, DDA-MS1 and DDA-

spectral counting, respectively. Overall, these results demonstrate the favorable quantitative

characteristics of SWATH/DIA data compared to DDA based analyses of data acquired on the same

Triple TOF model 5600 mass spectrometer (also see Figure S2C).

The presented results demonstrate that automated complex-centric analysis by *CCprofiler* allows

protein complex detection at a high sensitivity compared to manual inspection and that the system

provides an accurate decoy model for FDR estimation. The data further suggests that complex-

centric proteome profiling achieves competitive complex detection performance of the overall

workflow with only 81 LC-MS/MS measurements compared to a significantly larger scale

multidimensional fractionation experiment. Furthermore, our comparative analysis attests

SWATH/DIA more consistent and precise quantification when compared to DDA-based strategies and

largely increased sensitivity in targeted, complex-centric profiling under strict error rate control.

Complex-centric analysis of the HEK293 proteome: Insights into proteome modularity

We applied the complex-centric proteome profiling method to study the modularity of the HEK293

cell line proteome. Specifically, we first used the quantitative capacity of the method to estimate the

fraction of the observed proteome that was, under the extraction conditions used, part of protein

complexes as opposed to being present in monomeric form. Second, we tested the ability of the

method to conclusively confirm the presence of specific complexes in the sample and third, we

assessed the capability of the method to quantify the distribution of specific proteins across different

complexes.

Complex assembly state of the HEK293 proteome

To globally assess the state of assembly of the HEK293 proteome under the extraction and SEC

conditions used, we quantified for each of the 4916 proteins identified in the dataset (see above and

extended experimental procedures) the proportion that was detected in assembled or monomeric

state, respectively. To assign a protein signal to either state, we first calibrated a molecular weight

scale of proteins expected in each SEC fraction using a reference set of proteins with known

molecular weight (Figure S3). We then applied this scale to all detected proteins. We assigned

proteins to an assembled state if they eluted from the SEC column at an apparent molecular weight

that was minimally two times higher than the molecular weight indicated by the molecular weight

scale (Figure 3A). To assess the distribution of proteins across distinct molecular weight regions,

indicative of different assembly states as described above, we performed a protein-centric analysis of

the 58,792 peptide-level chromatograms (Figure 3A, compare Figure 1A, Step 4). Our analysis

identifies 5503 elution peaks for 4065 proteins (Supplemental Table 1), with no defined elution peaks

observable from the remaining 851 proteins. Of these, 2668 proteins (66 %) were observed in at least

one assembled state, whereas 1397 proteins (34%) were detected only in monomeric state, based on

the criteria used (Figure 3B). Of the 4065 proteins, 1103 proteins (27 %) eluted in more than one

peak and up to 6 elution peaks per protein were detected (Figure 3C). Proteins that were detected in

multiple assembled states were enriched in proteasome components, ribosomal proteins and

chaperones (Figure 3D). We further estimated the total protein mass that was detected in assembled

vs. monomeric state by integrating the total MS signals observed for proteins assigned to assembled

or monomeric states. The results show that 55 % of the detected protein mass was in assembled

state (Figure 3B).

Overall, these results indicate that a substantial fraction of the HEK293 proteome was detected in an

assembled state, both in terms of distinct protein elution peaks and protein mass (Figure 3B). The

results further demonstrate the capability of the method to quantify the distribution of proteins that

are part of different distinct complex assemblies (Figure 3C).

Complex-centric detection and quantification of complexes

As a next step we used the complex-centric workflow to confirm the presence of specific complexes

in the HEK293 cell sample. The query complexes were predicted from the CORUM, BioPlex and

StringDB reference databases of protein interactions, respectively, and the predictions were tested

by *CCProfiler* using the 4916 protein SEC elution profiles detected in the dataset (Figure 4 and

compare Figure 1A, Step 5). At a FDR of 5 % computed by the target-decoy model of *CCProfiler*,

complex-centric analysis confirmed 621, 1052 and 1795 of the tested query complexes from the

three respective input databases (For details, see extended experimental procedures and

Supplemental Table 2). Notably, *CCprofiler* was able to confidently detect complexes consisting of the

whole set of proteins predicted from the respective reference databases as well as complex signals

comprising only a subset of the reference proteins, thus supporting the quantification of fully and

partially assembled complexes. Up to this point in the analysis workflow, each protein complex signal

detected by *CCprofiler* is directly linked to one specific protein complex query in the prior

information dataset, derived from either CORUM, BioPlex or StringDB. However, some of the

subunits in each complex query might overlap with other complex queries. One simple example

would be that complex query A consists of subunits WXYZ and complex query B consists of subunits

VXYZ. If only XYZ are detected as a co-elution group in the data, they will, until this point, be reported

for both complex query A and B. In order to retrieve truly unique signals, the reported complex

signals can finally be collapsed based on a strategy that considers (i) subunit composition and (ii)

resolution in the chromatographic dimension. Taking the simple example from above, the signal

collapsing step will merge the two features from complex query A (XYZ discovered from querying

WXYZ) and B (XYZ discovered from querying VXYZ) to one unique protein signal XYZ that is

independent from the original complex queries (For more details, see extended experimental

procedures and Figure S4). According to this strategy, our integrated analysis across the three sets of

complex queries identifies 462 unique protein-protein complex signals (Figure 4).

Complex-centric detection of complex variants

The results above established the capacity of complex-centric profiling to detect and quantify subunit

distribution across complexes that are resolved by SEC and contain common proteins. We therefore

tested whether this capacity allowed us to detect novel protein modules of potential functional

significance. Among the 621 complex models that were confirmed by CCProfiler following predictions

from the CORUM database, 286 (46%) provided evidence of proteins common to two (152) and up to

5 or more (27) distinct, chromatographically separated complex instances (Figure 5A). For example,

the protein subunit fractionation profiles of the octameric COP9 signalosome complex, a central

regulator of E3 ligase activity and turnover, delineate both the CSN holo-complex consisting of all

eight subunits and a sub-complex consisting of subunits CSN1, CSN3 and CSN8 (Figure 5A&B). The

critical role of CSN proteins in regulating the ubiquitin-proteasome system and cellular homeostasis

has sparked great interest in the analysis in modules with variable subunit composition, and in

mechanisms that regulate their activity [20]. CSN proteins have also been linked to cancerogenesis[21–23].

Both CSN assemblies detected in the HEK293 dataset elute with apparent molecular weights in

accordance with a 1:1 stoichiometry. Further, the proteins CSN1/3/8 of the lower molecular weight

complex form a connected sub-module within the CSN holo-complex structure[24] (Figure 5C). The

occurrence of the distinct CSN1/3/8 complex detected in this study is consistent with protein

chromatographic data generated by co-fractionation-MS/MS in two other laboratories. Wan et al.

fractionated mild lysates of HEK293 cells by heparin ion exchange chromatography[25] followed by MS

analysis. Also in this separation dimension that is orthogonal to SEC, the quantitative MS profiles

show distinct co-fractionation of CSN1, CSN3 and CSN8 (Figure 5D, upper two panels). Larance et al.

fractionated mild lysates of U2OS cells by SEC and the quantitative profiles of the CSN subunits also

display distinct co-elution of CSN1, CSN3 and CSN8 at reduced molecular weight (Figure 5D, lower

panel). While the data of both research groups generally support the model of CSN1/3/8 as a distinct

cellular assembly, neither of them reported it as distinct from CSN holo-complex, likely owed to

limited resolution of the experimental data and the pairwise-interaction-focused analysis workflows

employed. Our findings suggest a potential functional role for the CSN sub-complex CSN1/3/8. We

therefore tested whether CSN1, CSN3 and CSN8 could stably assemble independent of the remaining

CSN components. We co-expressed human CSN1, CSN3 and CSN8 in insect cells, whereby CSN8 was

added with an N-terminal Strep(II)-tag and CSN1 & CSN3 were expressed with and N-terminal His6-

tags to facilitate reciprocal purification of the complex. The thus purified samples were analyzed by

SDS-PAGE and resulting banding patterns confirmed the formation of a stable trimer CSN1/3/8 in the

absence of the other CSN subunits that constitute the holo-complex (Figure 5D). Together, these

results support the finding from the complex-centric identification of the CSN1/3/8 complex as a

distinct substructure of the human COP9 Signalosome.

As a further example for the discovery of sub-complexes of a large holo-complex, complex-centric

proteome profiling detected six variant signals from the subunit chromatograms of the 26S

proteasome (Figure 6A). Two of the six variants represent known complexes, (i) the full 26S assembly

and (ii) the 20S core particle (Figure 6A). The remaining four co-elution signals point towards complex

variants of lower apparent molecular weight compared to the 26S and 20S particles (apex fractions

39, 40, 42 and 46, ~ 107-222 kDa) that consist predominantly of α and β subunits of the 20S core

particle. These reported complex variants point towards (iii) a β subunit assembly of β2, β3 and β7 at

fraction 39, (iv) a distinct assembly of α subunits α2 and α6 at fraction 40, (v) an assembly

intermediate of the seven α subunits α1, α2, α3, α4, α5, α6 and α7 at fraction 42, and (vi) a β6 and

proteasome regulatory subunit 8 assembly at fraction 46. To evaluate whether the observed signals

represent products of disassembly or complex biogenesis intermediates, we manually extended the

automated analysis of CCProfiler by additionally aligning the quantitative protein traces of the

chaperones known to be involved in 20S maturation with the respective complex subunits[26](Figure

6B). Strikingly, the distinctive co-elution of the early-stage specific chaperone PSMG3/PSMG4 dimer,

constitutive chaperone PSMG1/PSMG2 dimer and the late-stage specific proteasome maturation

factor POMP allowed us to classify the detected complex variants as early and late stage

intermediates of 20S biogenesis (Figure 6B). Notably, a systematic manual analysis of the

quantitative distribution of the proteasome and chaperone subunits across the detected complex

variants suggests the α1/α3/α4/α5/α7 and α1-7/β2/β3/β6/β7 complexes, respectively, as the

predominant early and late assembly intermediates on the path to 20S assembly, as assigned by

defined co-elution and inferred interaction with the chaperones specifically involved in early

(PSMG3/PSMG4 dimer) stages or late stages (POMP) of 20S proteasome biogenesis (Figure 6C).

Although the automated workflow could not fully resolve and explain the data, it successfully

pointed towards a distinct assembly of the alpha subunits (signal v) from the beta subunits (signal iii),

as well as the differential behavior of α2 and α6 compared to the other α subunits (signal iv). No

underlying biology could be determined for signal vi.

Together, these findings demonstrate the capacity of complex-centric profiling to derive models of

distinct variants of the queried complexes. These models can be reinforced by extending automated

analyses by the alignment of additional proteins' quantitative profiles followed by manual inspection.

SECexplorer – An interactive platform for complex-centric exploration of the HEK293 proteome

analyzed by SEC-SWATH-MS

To support customized, expert-driven and in-depth analyses of protein co-fractionation profiles

recorded by SEC-SWATH-MS of the HEK293 cell line, we set up the web platform *SECexplorer*.

SECexplorer enables visualization and interactive browsing of protein fractionation profiles of user-

defined sets of proteins. Users can perform multiple tasks, including (i) testing of novel predicted

models on complex formation between candidate proteins or (ii) interrogating the profile sets of

known modules for evidence pointing towards new variants or (iii) manual refinement and extension

of results obtained from automated complex-centric profiling, for example by extending the set of

automatically detected complex components with additional proteins e.g. derived from the literature

or from interaction network context. Analyses are assisted by the *CCprofiler* algorithm suggesting

distinct co-elution signals and calculating their expected to apparent molecular weight mismatch,

among other metrics, in order to speed up data interpretation by expert users. *SECexplorer* can be

accessed at *https://sec-explorer.ethz.ch/* (Figure 7A). As an example for the use of *SECexplorer*, we

followed up on the peak shoulder at elevated molecular weight observed in the CSN holo-complex

co-elution signal (Compare Figure 5B). Overlaying the elution profiles of known components of a E3-

CRL substrate of the COP9 Signalosome[27] revealed defined co-elution in the peak shoulder range,

supporting the detection of a likely E3-CRL-bound subpopulation of CSN holo-complexes (Figure 7B).

To derive a quantitative signal in the situation of only partial chromatographic resolution we

employed a Gaussian deconvolution mixture model, suggesting a substrate-bound fraction of CSN

holo-complex of 25 ± 3 % across the 8 subunits (Figures 7C and S5).

Discussion

In this paper we describe complex-centric proteome profiling, an integrated experimental and

computational approach to detect and quantify protein complexes isolated from their natural source,

to generate new insights into the modular organization of proteomes.

The need to systematically analyze the organization of the proteome arises from the notion of a

modular biology proposed by Hartwell et al.[1]. It essentially states that biochemical functions are for

the most part catalyzed and controlled by functional modules, most frequently protein complexes,

and that (genomic) perturbation of complexes results in perturbed biochemical functions and

potentially in disease phenotypes. The notion of a modular biology thus extends the pioneering work

of Pauling et al. on defining sickle cell anemia as a molecular disease[28] to the proteome level. Protein

complexes and protein-protein interactions have been studied extensively by a wide range of

techniques and have led to compendia of complexes[15,16,29] and maps of protein interaction

networks[16,30,31]. These compendia have in common that they describe generic, usually static

instances of complexes and interactions[32–34]. To distinguish between different biochemical states of a

cell, it is also essential to determine qualitative and quantitative differences in functional modules in

different samples. To date this has been attempted by two broad approaches. The first is based on

microscopic methods including FRET[35] which provide outstanding resolution and precision of steric

proximity but are labor intensive and focused on one to a few interactions at a time. The second is

based on a mass spectrometric approach referred to as correlation profiling[36] in which samples of

native modules are separated into a set of fractions and the protein contents of each fraction is

determined by quantitative mass spectrometry. The association of a protein to a specific module is

then asserted by the consistency of the quantitative pattern of the protein in question with other

proteins of the same module[37]. Initially used to define the composition of the specific modules such

as the large RNA polymerase II preinitiation complex[37] and the human centrosome [38], correlation

profiling has also been employed to broadly assign protein localization to different subcellular

compartments[39–41]and the scope has been extended towards systematically interrogating protein-

protein complexes by correlating protein patterns in fractions obtained from different biochemical

fractionation methods[9,11,42]. Such studies have used different native complex separation methods

including SEC, IEX, density gradient centrifugation and blue native gels[9,11,42]. The scientific scope has

extended to the analysis of cells of different species, culminating in the description of hundreds of

complexes in a single, albeit massive experiment[25]. Correlation profiling, therefore has the potential

to determine the quantity and composition of hundreds of protein modules in a single operation.

In the present paper we describe a conceptual and technical advance in the field of correlation

profiling. As a conceptual advance we introduce the principle of complex centric analysis. It is

inspired by the peptide-centric analysis concept employed for the specific and sensitive detection of

peptides from proteomic samples in targeted proteomic approaches, such as SWATH/DIA[6], and

extends the use of prior information for the analysis of proteins to the level of protein complexes.

Similar to peptide-centric analysis of SWATH/DIA data, high selectivity and sensitivity is achieved by

focusing the analysis on analytes conceivably expected in the sample when querying the protein co-

fractionation data for candidate protein complexes that are inferred from reference protein

interaction maps. Thereby, prior information significantly constrains complex inference from co-

fractionation profiling data and thus adds specificity and the possibility to develop a target-decoy

model to assess the reliability of the obtained results. Furthermore, complex identifications are

directly linked to quantitative chromatographic signals, a central feature of targeted proteomics

approaches. As technical advances we demonstrate the benefits of SWATH/DIA for the analysis of

the sequential SEC fractions, introduce a freely accessible computational framework *CCprofiler* and

provide a tool facilitating the exploitation of complex-centric data, *SECexplorer* (Figure 7).

In combination, these technical and conceptual developments provide the following advances to the

field of correlation profiling. First, the preferable quantitative performance of SWATH/DIA provides

more complete and consistent sampling of the eluting proteome, resulting in fewer gaps and noise in

the recorded profiles. This results in deeper insights into modular proteome organization, including

the detectability of low abundance complex intermediates. Second, the use of prior information

reduces false positive assignments of complex co-membership due to coincidental co-elution of

proteins that do in reality not interact. Third, the *CCprofiler* pipeline introduces the first statistical

target-decoy model to tightly control error rates in the inference of complexes from co-fractionation

profiling experiments and represents a comprehensive, open-source platform to support complex-

centric profiling of proteomes, irrespective of the fractionation method used. Fourth, the efficiency

of information retrieval and thus overall method throughput is drastically increased when compared

to current co-fractionation based complex analyses, generating comprehensive and accurate

assessments of proteome arrangement from an order of magnitude less LC-MS experiments than

necessitated earlier. Together, these advances transform the SWATH/DIA-based complex-centric

proteome profiling into a robust, generally applicable technique supported by a freely accessible

computational framework.

We applied complex-centric profiling to a native protein extract from exponentially growing Hek293

cells. Collectively, the results demonstrate the superior performance of the technique compared to

the state of the art and provide new biological insights, as follows. The analysis establishes estimates

for the overall assembly state of a human proteome - 55 % of inferred protein mass and two thirds

(66 %) of the observed protein species appear engaged in higher order assemblies; a lower-boundary

estimate given inevitable losses of associations in the experimental procedure. Besides detecting

cumulatively 462 cellular complexes upon targeted analysis, the method in many instances resolves

distinct variants of the expected complexes, such as sub-complexes that elute independently from

the chromatographic column. While sub-complex signals may originate from artefactual disruption of

cellular complexes, we demonstrate in two cases how orthogonal pieces of evidence can build

confidence in the biological relevance of substructures assigned from defined subunit co-elution.

First, we identified a new complex CSN1/3/8 as a sub-complex of the COP Signalosome (CSN)

holocomplex that elicits crucial regulatory functions towards E3 ligase complexes and the ubiquitin

proteasome system[43]. It is tempting to speculate that a putative function of the CSN1/3/8 sub-

complex could be the negative regulation of CSN holocomplex activity, due to the fact that the sub-

complex incorporates the subunit CSN1 which is involved in substrate recognition[27], but does not

contain the catalytically active CSN5 subunit. CSN5 embodies the de-neddylation activity to the CSN

holocomplex[27]. CSN1/3/8 may potentially sequester neddylated E3 CRLs from CSN-mediated de-

neddylation and thus affect their lifetimes and overall activity profiles. In a second example,

complex-centric analysis in combination with manual refinement identified early and late assembly

intermediates on the path towards the 20S proteasome particle. Strikingly, the early and late

intermediary complexes assigned (early: α1/α3/α4/α5/α7, late: α1-7/β2/β3/β6/β7) collide with

current models of the temporal order of subunit assembly[44,45] (for a graphical summary, see Figure

6B, lower panel). Current models entail early α-ring intermediates lacking subunits α3 and α4[46]. In

contrast, our model suggests assembly of pre-α-ring intermediates composed of subunits α4, α7 α5

α1 and α3 (forming a connected substructure of the α-ring in this order[47]) that lacks subunits α2 and

α6. These join thereafter to complete the α-ring, under involvement of the chaperone

POMP/hUmp1. Current models further suggest that ordered β-ring assembly scaffolded by α-rings in

the sequence of β2, β3, β4, β5, β6, β1 and lastly β7[44,45] help overcome a POMP-dependent

checkpoint for dimerization into the mature 20S particle[48]. The detection of late assembly

intermediate α1-7/β2/β3/β6/β7 in our data suggests an alternate sequence of assembly with early

incorporation of subunit β7 and dimerization after the recruitment of subunits β1, β4 and β5.

These insights into complex biogenesis could prove valuable, for example, in the design of future

therapeutic strategies aiming to counteract elevated proteasome expression and activity that has

been associated with cancer pathobiology[49]. This is exemplified by current attempts to target

proteasomal activity via the chaperone POMP[50] (Figure 6C). We expect that the data generated by

complex-centric proteome profiling will lead to the discovery of other instances of characteristic

protein complexes and sub-complexes and thus trigger research into their functional roles.

Despite the advances and benefits of complex-centric proteome profiling by SEC-SWATH-MS, the

method has a number of limitations. i) The balance of stability of complexes and extractability in

native form. Inevitably, associations are lost in the experimental procedure, most notably upon

dilution imposed during lysis and subsequent size exclusion chromatography, reducing protein

concentration by ca. five orders of magnitude from the cellular environment (ca. 300 mg/ml[51]) to the

conditions on the SEC column (ca. 0.06 mg/ml). Consequently, complex detectability is limited by

thermodynamic stability and despite best efforts towards minimizing complex disintegration (fast

processing in the cold and analyte adsorption-free chromatography), thermodynamically labile

interactions, particularly those with fast off-rates, are likely inaccessible by correlation profiling

methods, including complex-centric proteome profiling. While first studies have evaluated chemical

crosslinking as means to stabilize cellular modules for chromatographic analyses[52], it remains an

open challenge to identify uniformly beneficial crosslinking reagents and reaction conditions that

yield optimal balance between stabilization of biologically relevant structures and artefactual cross

linking across the full range of protein expression in the cell[53].

Furthermore, complex-centric proteome profiling is limited to the scope of the prior knowledge on

protein association employed. However, continued efforts to map cellular protein association space[54]

and computational integration of multiple lines of experimental evidence[29] will continually improve

the quality and completeness of the prior knowledge useable as input to targeted, complex-centric

analyses. Extended reference protein interaction maps will support near-complete mapping of the

complexes detectable in co-fractionation experimental data in the near future, supported by

scalability of the target-decoy statistical model. That being said, the statistical model itself is limited

to the assignment of an FDR on the evidence of detection of defined complexes in the complex query

set. Future improvements could potentially support a robust statistical model covering also post-

processing steps, such as collapsing of detected features across multiple complex query sets to

unique co-elution signals.

SEC-SWATH-MS accelerates the mapping of cellular complexes. Whereas the method yields a similar

coverage of complexes compared to state-of-the art at over fourteen times less LC-MS injections, it

still required 81 fractions to be analyzed at 2 h gradient time per fraction, culminating in 162 h of net

MS acquisition time. This fact limits the scope for cohort studies. However, this issue may well be

alleviated soon, given anticipated improvements SWATH/DIA sample throughput with minimal loss

of protein coverage that seem achievable because in SWATH/DIA acquisition the number of analytes

quantified does much less strongly depended on gradient length than is the case for DDA

acquisition. As a consequence of the high quantitative quality of the data, the increased efficiency of

the method and the error model this study lays the foundation to conclusively quantify changes of

proteome organization as a function of cell state. Ultimately, extensions of our workflow will support

the detection of subtle re-arrangements within proteomes that occur in response to perturbation or

along central biological processes. Such insights will help foster our understanding of the importance

of higher order organization of the parts to convey plasticity and regulation to cellular systems.

Author contributions

Conceptualization, R.A., M.G., B.C.C. and M.H.; Methodology, M.H., I.B., G.R.; Software, I.B., M.H.,

G.R., R.H., M.F. and A.B.E.; Validation, M.H. and I.B.; Formal Analysis, M.H. and I.B.; Investigation,

M.H.; Resources, M.G. and R.A.; Data Curation, I.B. and M.H.; Writing – Original Draft, M.H. and R.A.;

Writing – Review & Editing, I.B., M.F., G.R., R.H., A.B.E., B.C.C., M.G. and R.A.; Completion Submitted

Manuscript, M.H., I.B. and R.A.; Visualization, M.H., I.B. and R.H.; Supervision, B.C.C., M.G. and R.A.;

Project Administration, M.H.; Funding Acquisition, R.A.

Acknowledgments

We thank all Aebersold and Gstaiger lab members for helpful discussions, and with special emphasis

Betty Friedrich, Audrey van Drogen, Peter Blattmann, Hannes L. Roest, Ludovic Gillet and Yansheng

Liu. We further thank Prof. Nicolas Thomä, Dr. Lingaraju Manjappa of the Friedrich Miescher Institute

for Biomedical Research as well as Dr. Martin Renatus and Arnaud Decock of the Novartis Institutes

for Biomedical Research for materials and guidance in COP9 signalosome subunit co-expression and

co-purification experiments. We would also like to thank the Scientific IT Support (ID SIS) of ETH

Zurich for support and maintenance of the lab-internal computing infrastructure (iPortal) and

specifically Uwe Schmitt for his help with the SECexplorer setup. The project was supported by the

SystemsX.ch projects PhosphoNetX PPM and project TbX to R.A., and the European Research Council

(ERC-20140AdG 670821 to R.A.). M.H. was supported by a grant from Institut Mérieux. B.C.C. was

supported by a Swiss National Science Foundation Ambizione grant (PZ00P3_161435). A.B.E. was

supported by the National Institutes of Health project Omics4TB Disease Progression (U19

AI106761). I.B. was supported by the Swiss National Science Foundation (grant no. 31003A_166435).

Conflict of Interests

517 The authors declare that they have no conflict of interest.

Experimental Procedures

Preparation of native HEK293 proteome and fractions for MS analysis

HEK293 cells were obtained from ATCC and cultured in DMEM containing 10 % FCS and 50 µg/mL penicillin/streptomycin to 80 % confluency. Ca. 7e7 cells were mildly lysed by freeze-thawing into 0.5 % NP-40 detergent- and protease and phosphatase inhibitor containing buffer, essentially as described[55], albeit without the addition of Avidin. Lysates were cleared by 15 minutes of ultracentrifugation (100,000×g, 4 °C) and buffer was exchanged to SEC buffer (50 mM HEPES pH 7.5, 150 mM NaCl) over 30 kDa molecular weight cut-off membrane at a ratio of 1:50 and concentrated to 25-30 mg/ml (as judged by OD280). After 5 min of centrifugation at 16.9 ×g, 4 °C, the supernatant was directly subjected to fractionation on a Yarra-SEC-4000 column (300×7.8 mm, pore size 500 Å, particle size 3 µm, Phenomenex, CA, USA). Per SEC run, 1 mg native proteome (by OD280) was injected and fractionated at 500 µl/min flow rate at 4 °C, collecting fractions at 0.19 min per fraction from 10 to 28 minutes post-injection, fractions 3-83 of which were considered relevant proteome elution range and considered for further analysis with fractionation index 1-81. The fractions collected from two consecutive SEC fractionations of the same extract (2×1 mg) were pooled for subsequent bottom-up proteomic analysis. Apparent molecular weight per fraction was log-linearly calibrated based on column performance check protein mix analyzed prior and after each experimental replicate (AL0-3042, Phenomenex, CA, USA). An aliquot of the unfractionated mild proteome extract was included in peptide sample preparation and LC-MS analysis. Proteins were proteolyzed to peptide level by trypsin digestion (Promega V5111) in the presence of 1 % sodium deoxycholate (Sigma-Aldrich D6750), reduced, alkylated and de-salted on C18 reversed phase (96-Well MACROSpin Plate, The Nest Group, MA, USA) and each sample was supplemented with equal amounts of internal retention time calibration peptides (iRT kit, Biognosys, CH).

Baculoviral co-expression and co-purification

Sf21VM Cells were maintained in ExCell420 Medium in Erlenmeyer culture flasks shaking at 27.5 °C. Human COP9 Signalosome subunits bearing N-terminal Strep(II) or His6 tags were co-expressed by co-infection of Sf21VM cells with three baculoviral vectors obtained from Lingaraju et al.[24]. After 48 h, cells were mildly lysed and COP9 signalosome subunits and complexes differentially affinity-purified on StrepTactin and Ni-NTA-coated magnetic beads (Qiagen) followed by bead boiling in SDS loading buffer and subunit detection via SDS PAGE and InstantBlue staining (Expedeon). Subunits were identified by size and in reference to individual expression and in-gel detection.

MS analysis

LC-MS analysis of peptide samples was performed in both DDA and SWATH/DIA acquisition mode on an AB Sciex TripleTOF 5600+ instrument (AB Sciex, MA, USA), side-by-side per sample, sliding from early to late-eluting fractions. On-line reversed phase chromatography fractionated peptide samples delivering at 300 nL/min flow a 120-min gradient from 2–35% buffer B (0.1% formic acid, 90% acetonitrile) in buffer A (0.1 % formic acid, 2% acetonitrile) on a self-packed picoFrit emitter packed with 20 cm column bed of 3 μm 200-Å Magic C18 AQ stationary phase, essentially as described[6,55]. In data-dependent acquisition (DDA), MS1 survey spectra were acquired for the range of 360–1,460 m/z with a 500 ms fill time cap. The top 20 most intense precursors of charge state 2–5 were selected for CID fragmentation and MS2 spectra were collected for the range of 50–2,000 m/z, with 100 ms fill time cap and dynamic exclusion of precursor ions from reselection for 15 s, essentially as described[55].

Data-independent acquisition (SWATH/DIA) mass spectrometry was performed using an updated scheme of 64 variably sized precursor co-isolation windows optimized for human cell lysate MS signal density (SWATH® 2.0, essentially as described[56]). SWATH cycles (64 x 50 ms accumulation time) were interspersed by MS1 survey scans for the range of 360–1,460 m/z with a 250 ms fill time cap, resulting in an overall period cycle time of 3498 milliseconds. The MS2 mass range was set to 200 – 2000 m/z.

Data processing

DDA-MS data were processed using the MaxQuant software package (version 1.5.3.17) with the human canonical SwissProt reference database (build Aug-2014), standard parameters and variable methionine oxidation and N-terminal acetylation enabled. Match between runs was enabled to facilitate ID transfer and more consistent MS1 quantification (from and to) between adjacent fractions. Raw peptide MS1 intensities of individual peptide precursor signals were further considered.

The SWATH/DIA-MS data were analyzed via targeted, peptide-centric analysis, querying > 200,000 precursors based on the combined human assay library (CAL) in the SWATH fragment ion chromatograms, using a modified OpenSWATH[7], PyProphet[57,58] and TRIC[18] workflow and the iPortal framework[59] to recover precursors at an experiment-wide assay-level (TRIC target) FDR of 5 %. Precursor-level results were summed per peptide and further filtered on chromatography-informed scores, controlling the false discovery rate on protein level to below 1 %, employing the simple target-decoy method as implemented in *CCprofiler*. The protein-level fraction of false targets was estimated using R/SWATH2stats[60]. Specific filtering rules were, first, eliminating all values part of

consecutive identification stretches below length three and, second, exclusion of peptides based on

their quantitative fractionation pattern's average dissimilarity to those of sibling peptides (originating

from the same parent protein, discarding peptides with average sibling peptide correlation

coefficient (spc) below 0.316 to achieve an estimated FDR of < 1 % among the remaining 4958

proteins (4916 of which quantifiable with ≥2 proteotypic peptides for downstream analyses).

Protein-centric analysis was performed within the *CCprofiler* framework, detecting protein elution

signals among peptide SEC chromatogram sets grouped by parent SwissProt protein identifier (For

details, see extended experimental procedures).

Complex centric analysis was performed within the *CCprofiler* framework, detecting complex elution

signals among protein SEC chromatogram sets, defined by reference databases CORUM, BioPlex and

StringDB (Compare Figure 1 and Figure 3). Protein level chromatograms were generated by summing

the two highest intensity proteotypic peptide chromatograms. Complex queries as well as decoy

complex queries were formulated from the reference databases by splitting variants and re-merging

redundant human CORUM complexes (N = 1753 query groups) and by partitioning networks into

query groups of one seed protein including all direct interactors at the confidence thresholds (BioPlex

interaction probability ≥ 0.75 and StringDB interaction score ≥ 900). Decoy complex queries of

matching number and size were formulated in reference to the input connectivity information,

ensuring that decoy complex queries do not entail interacting subsets. Co-elution signals were then

detected using the signal processing algorithm of *CCprofiler* in grid search mode over a range of

processing parameter sets. Parameters and results were then selected to ensure precision of ≥ 95 %

among the protein peak group signals, followed by summarizing signals to unique groups at defined

chromatographic retention times, chromatographically resolved complexes and complex variants.

Generation of true positive reference sets of co-elution signals

To guide algorithm development and to evaluate extendibility of the target-decoy approach to the

level of protein complexes, we manually curated protein subunit co-elution signals among the 1753

protein chromatogram sets defined by the CORUM complexes.

A more comprehensive and detailed protocol for all data processing and analysis steps is provided in

the extended experimental procedures. The mass spectrometry proteomics data have been

deposited to the ProteomeXchange Consortium (http://proteomecentral.proteomexchange.org) via

the PRIDE partner repository[61] with the dataset identifier PXD007038. The *CCprofiler* package is

freely available on GitHub (https://github.com/CCprofiler/CCprofiler/) including a vignette describing

the main functionalities and usage of the software (Supplemental item 1).

References

1. Hartwell, L. H., Hopfield, J. J., Leibler, S. & Murray, A. W. From molecular to modular cell biology. *Nature* **402,** C47-52 (1999).

2. Ahrens, C. H., Brunner, E., Qeli, E., Basler, K. & Aebersold, R. Generating and navigating proteome maps using mass spectrometry. *Nat. Rev. Mol. Cell Biol.* **11,** 789–801 (2010).

3. Ting, Y. S. *et al.* Peptide-Centric Proteome Analysis: An Alternative Strategy for the Analysis of Tandem Mass Spectrometry Data. *Mol. Cell. Proteomics* **14,** 2301–2307 (2015).

4. Picotti, P. & Aebersold, R. Selected reaction monitoring-based proteomics: workflows, potential, pitfalls and future directions. *Nat. Methods* **9,** 555–66 (2012).

5. Bourmaud, A., Gallien, S. & Domon, B. Parallel reaction monitoring using quadrupole-Orbitrap mass spectrometer: Principle and applications. *Proteomics* **16,** 2146–2159 (2016).

6. Gillet, L. C. *et al.* Targeted data extraction of the MS/MS spectra generated by data-independent acquisition: a new concept for consistent and accurate proteome analysis. *Mol. Cell. Proteomics* **11,** O111.016717 (2012).

7. Röst, H. L. *et al.* OpenSWATH enables automated, targeted analysis of data-independent acquisition MS data. *Nat. Biotechnol.* **32,** 219–23 (2014).

8. Navarro, P. *et al.* A multicenter study benchmarks software tools for label-free proteome quantification. *Nat. Biotechnol.* **34,** 1130–1136 (2016).

9. Liu, X., Yang, W., Gao, Q. & Regnier, F. Toward chromatographic analysis of interacting protein networks. *J. Chromatogr. A* **1178,** 24–32 (2008).

10. Dong, M. *et al.* A 'tagless' strategy for identification of stable protein complexes genome-wide by multidimensional orthogonal chromatographic separation and iTRAQ reagent tracking. *J. Proteome Res.* **7,** 1836–49 (2008).

11. Rudashevskaya, E. L., Sickmann, A. & Markoutsa, S. Global profiling of protein complexes: current approaches and their perspective in biomedical research. *Expert Rev. Proteomics* **13,** 951–964 (2016).

12. Havugimana, P. C. *et al.* A census of human soluble protein complexes. *Cell* **150,** 1068–81 (2012).

13. Kristensen, A. R. & Foster, L. J. in *Methods in molecular biology (Clifton, N.J.)* **1188,** 263–270 (2014).

14. Kirkwood, K. J., Ahmad, Y., Larance, M. & Lamond, a I. Characterization of native protein
complexes and protein isoform variation using size-fractionation-based quantitative
proteomics. *Mol Cell Proteomics* **12,** 3851–3873 (2013).

15. Ruepp, A. *et al.* CORUM: the comprehensive resource of mammalian protein complexes--
2009. *Nucleic Acids Res.* **38,** D497-501 (2010).

16. Huttlin, E. L. *et al.* The BioPlex Network: A Systematic Exploration of the Human Interactome.
Cell **162,** 425–440 (2015).

17. Franceschini, A. *et al.* STRING v9.1: protein-protein interaction networks, with increased
coverage and integration. *Nucleic Acids Res.* **41,** D808-15 (2013).

18. Röst, H. L. *et al.* TRIC: an automated alignment strategy for reproducible protein
quantification in targeted proteomics. *Nat. Methods* **13,** 777–83 (2016).

19. Rosenberger, G. *et al.* A repository of assays to quantify 10,000 human proteins by SWATH-
MS. *Sci. data* **1,** 140031 (2014).

20. Dubiel, D., Rockel, B., Naumann, M. & Dubiel, W. Diversity of COP9 signalosome structures
and functional consequences. *FEBS Lett.* **589,** 2507–13 (2015).

21. Lee, M.-H., Zhao, R., Phan, L. & Yeung, S.-C. J. Roles of COP9 signalosome in cancer. *Cell Cycle*
10, 3057–3066 (2011).

22. Gummlich, L., Rabien, A., Jung, K. & Dubiel, W. Deregulation of the COP9 signalosome–cullin-
RING ubiquitin-ligase pathway: Mechanisms and roles in urological cancers. *Int. J. Biochem.*
Cell Biol. **45,** 1327–1337 (2013).

23. Chen, J. *et al.* CSN6 drives carcinogenesis by positively regulating Myc stability. *Nat. Commun.*
5, 5384 (2014).

24. Lingaraju, G. M. *et al.* Crystal structure of the human COP9 signalosome. *Nature* **512,** 161–5
(2014).

25. Wan, C. *et al.* Panorama of ancient metazoan macromolecular complexes. *Nature* (2015).
doi:10.1038/nature14877

26. Hirano, Y. *et al.* A heterodimeric complex that promotes the assembly of mammalian 20S
proteasomes. *Nature* **437,** 1381–5 (2005).

27. Cavadini, S. *et al.* Cullin–RING ubiquitin E3 ligase regulation by the COP9 signalosome. *Nature*
531, 598–603 (2016).

28. Pauling, L., Itano, H. A., Singer, S. J. & Wells, I. C. Sickle Cell Anemia, a Molecular Disease.
Science **110,** 543–548

29. Drew, K. *et al.* Integration of over 9,000 mass spectrometry experiments builds a global map
of human protein complexes. *Mol. Syst. Biol.* **13,** 932 (2017).

30. Rolland, T. *et al.* A Proteome-Scale Map of the Human Interactome Network. *Cell* **159,** 1212–
1226 (2014).

31. Szklarczyk, D. *et al.* STRING v10: protein–protein interaction networks, integrated over the
tree of life. *Nucleic Acids Res.* **43,** D447–D452 (2015).

32. Gstaiger, M. & Aebersold, R. Genotype–phenotype relationships in light of a modular protein
interaction landscape. *Mol. Biosyst.* **9,** 1064 (2013).

33. Havugimana, P. C., Hu, P. & Emili, A. Protein complexes, big data, machine learning and
integrative proteomics: lessons learned over a decade of systematic analysis of protein
interaction networks. *Expert Rev. Proteomics* **14,** 845–855 (2017).

34. Mehta, V. & Trinkle-Mulcahy, L. Recent advances in large-scale protein interactome mapping.
F1000Research **5,** (2016).

35. Song, Y., Madahar, V. & Liao, J. Development of FRET Assay into Quantitative and High-
throughput Screening Technology Platforms for Protein–Protein Interactions. *Ann. Biomed.*
Eng. **39,** 1224–1234 (2011).

36. Foster, L. J. *et al.* A mammalian organelle map by protein correlation profiling. *Cell* **125,** 187–
99 (2006).

37. Ranish, J. A. *et al.* The study of macromolecular complexes by quantitative proteomics. *Nat.*
Genet. **33,** 349–355 (2003).

38. Andersen, J. S. *et al.* Proteomic characterization of the human centrosome by protein
correlation profiling. *Nature* **426,** 570–4 (2003).

39. Foster, L. J. *et al.* A mammalian organelle map by protein correlation profiling. *Cell* **125,** 187–
99 (2006).

40. Dunkley, T. P. J. *et al.* Mapping the Arabidopsis organelle proteome. *Proc. Natl. Acad. Sci.* **103,**
6518–6523 (2006).

41. Yan, W., Aebersold, R. & Raines, E. W. Evolution of organelle-associated protein profiling. *J.*
Proteomics **72,** 4–11 (2009).

42.   Dong, M. *et al.* A 'tagless' strategy for identification of stable protein complexes genome-wide
by multidimensional orthogonal chromatographic separation and iTRAQ reagent Tracking. *J.*
Proteome Res. **7,** 1836–1849 (2008).

43.   Dubiel, D., Rockel, B., Naumann, M. & Dubiel, W. Diversity of COP9 signalosome structures
and functional consequences. *FEBS Lett.* **589,** 2507–13 (2015).

44.   Im, E. & Chung, K. C. Precise assembly and regulation of 26S proteasome and correlation
between proteasome dysfunction and neurodegenerative diseases. *BMB Rep.* **49,** 459–73
(2016).

45.   Hirano, Y. *et al.* Dissecting beta-ring assembly pathway of the mammalian 20S proteasome.
EMBO J. **27,** 2204–13 (2008).

46.   Hirano, Y. *et al.* A heterodimeric complex that promotes the assembly of mammalian 20S
proteasomes. *Nature* **437,** 1381–1385 (2005).

47.   Huang, X., Luan, B., Wu, J. & Shi, Y. An atomic structure of the human 26S proteasome. *Nat.*
Struct. Mol. Biol. **23,** 778–785 (2016).

48.   Li, X., Kusmierczyk, A. R., Wong, P., Emili, A. & Hochstrasser, M. beta-Subunit appendages
promote 20S proteasome assembly by overcoming an Ump1-dependent checkpoint. *EMBO J.*
26, 2339–49 (2007).

49.   Voutsadakis, I. A. Proteasome expression and activity in cancer and cancer stem cells. *Tumour*
Biol. **39,** 1010428317692248 (2017).

50.   Goldberg, A. L., Zhao, J. & Collins, G. A. Blocking Cancer Growth with Less POMP or
Proteasomes. *Mol. Cell* **59,** 143–5 (2015).

51.   Milo, R. What is the total number of protein molecules per cell volume? A call to rethink some
published values. *Bioessays* **35,** 1050–5 (2013).

52.   Larance, M. *et al.* Global Membrane Protein Interactome Analysis using *In vivo* Crosslinking
and Mass Spectrometry-based Protein Correlation Profiling. *Mol. Cell. Proteomics* **15,** 2476–
2490 (2016).

53.   Leitner, A., Faini, M., Stengel, F. & Aebersold, R. Crosslinking and Mass Spectrometry: An
Integrated Technology to Understand the Structure and Function of Molecular Machines.
Trends Biochem. Sci. **41,** 20–32 (2016).

54.   Huttlin, E. L. *et al.* Architecture of the human interactome defines protein communities and

disease networks. *Nature* (2017). doi:10.1038/nature22366

55.   Collins, B. C. *et al.* Quantifying protein interaction dynamics by SWATH mass spectrometry:

application to the 14-3-3 system. *Nat. Methods* **10,** 1246–53 (2013).

56.   Collins, B. C. *et al.* Multi-laboratory assessment of reproducibility, qualitative and quantitative

performance of SWATH-mass spectrometry. *bioRxiv* (2016). at

<http://biorxiv.org/content/early/2016/09/14/074567>

57.   Reiter, L. *et al.* mProphet: automated data processing and statistical validation for large-scale

SRM experiments. *Nat. Methods* **8,** 430–5 (2011).

58.   Teleman, J. *et al.* DIANA--algorithmic improvements for analysis of data-independent

acquisition MS data. *Bioinformatics* **31,** 555–62 (2015).

59.   Kunszt, P. *et al.* iPortal: the swiss grid proteomics portal: Requirements and new features

based on experience and usability considerations. *Concurr. Comput. Pract. Exp.* **27,** 433–445

(2015).

60.   Blattmann, P., Heusel, M. & Aebersold, R. SWATH2stats: An R/Bioconductor Package to

Process and Convert Quantitative SWATH-MS Proteomics Data for Downstream Analysis

Tools. *PLoS One* **11,** e0153160 (2016).

61.   Vizcaíno, J. A. *et al.* The Proteomics Identifications (PRIDE) database and associated tools:

status in 2013. *Nucleic Acids Res.* **41,** D1063–D1069 (2012).

Figure 1: Scheme of complex-centric proteome profiling by SEC-SWATH-MS

Figure 1: Scheme of complex-centric proteome profiling by SEC-SWATH-MS. **A** Workflow to quantify cellular complexes in five steps, extending the targeted analysis concept from peptide-centric interpretation of SWATH-MS data to the levels of protein and protein complex detection from size exclusion chromatographic fractions (also see Figure S1). **B** Specific steps of targeted, complex-centric analysis of co-fractionation data in the *CCprofiler* package.

Figure 2: Benchmarking and performance assessment of complex-centric proteome profiling.

Figure 2: Benchmarking and performance assessment of complex-centric proteome profiling. **A** Benchmark of *CCprofiler* algorithm and error model in reference to a manually curated reference set of signals displays conservative decoy-based FDR control and high sensitivity, recalling 91 % of high quality co-elution signals (also see extended experimental procedures and Figure S2A/B). **B** Assessment of complex identification performance for the overall complex-centric profiling workflow employing CORUM, BioPlex and StringDB proteome connectivity priors, based on CORUM complex recovery in comparison to multidimensional co-fractionation and 1,176 LCMS-run-based complexome map (See Figure 4 and extended experimental procedures). **C** Comparison of SWATH-MS-based quantification to DDA-MS-based strategies (MS1 XIC and spectral counting) with regard to consistency (as judged based upon protein-level SEC chromatogram robustness towards increasing requirements on the number of consecutive detections) and precision (judged based on the correlation between sibling peptide SEC chromatograms) of quantification and overall performance in error-controlled complex-centric query of CORUM complexes in the respective protein level chromatogram sets (also see Figure S2C).

Figure 3: Detection of protein elution via protein-centric analysis.

Figure 3: Detection of protein elution via protein-centric analysis.

Figure 3: Detection of protein elution via protein-centric analysis. A Peptide-level SEC chromatograms are grouped by UniprotKB identifier to detect co-elution signals indicative of protein elution ranges/peaks. Based on external size calibration of the apparent analyte molecular weight per SEC fraction, signals can be attributed to likely assembled or monomeric state (also see Figure S3). For TCPE, three distinct elution signals, numbered 1-3, are detected, two in the assembled and one in the monomer elution range. **B** Global statistics of protein signal attribution to assembled or monomeric state. The majority of proteins (66 %) and protein mass (55 %) appear in assembled state in SEC-SWATH-MS. **C** Proteins are observed eluting in 1-6 distinct peaks and with a wide range of apparent vs. monomeric molecular weight ratios (distributions, left panels). The molecular weight ratios of the three peaks detected for TCPE (displayed in A) are indicated. Many of the proteins eluting in a single peak (top panel and bar) appear assembled. Proteins eluting multiple times (27 % of the proteins) do so preferentially in the assembled range, suggesting frequent participation in multiple differently sized macromolecular assemblies (lower panels and pie chart). For a list of all detected protein peaks, see Supplemental Table 1. **D** Proteins observed in multiple assembled peaks (n = 659) are enriched in components of the proteasome and other known large complex assemblies.

Figure 4: Complex-centric profiling of the HEK293 proteome by targeted query of CORUM, BioPlex and StringDB.

Figure 4: Complex-centric profiling of the HEK293 proteome by targeted query of CORUM, BioPlex and StringDB. Schematic overview of the targeted, complex-centric analysis of the protein-level co-fractionation map recorded in SEC-SWATH-MS via *CCprofiler*. The three-tiered analysis is centered on complex hypotheses (i.e. groups of proteins queried for co-elution in the SEC data) obtained from CORUM or formulated from BioPlex and StringDB. At complex hypothesis FDR controlled to 5 % via the decoy-based error model, co-elution evidence is confidently detected for 621, 1052 and 1795 (representing 35.4, 13.8 and 19.2 %) of the queried CORUM-, BioPlex- and StringDB-derived hypotheses, respectively. Heterogeneity and redundancy within and across the different hypothesis sets translates to the co-elution signals retrieved, which, pieced together by collapsing on composition and SEC elution fraction, identify 462 distinct, chromatographically resolved co-elution groups representative of distinct complexes or equisized families of complexes (also see Figure S4). For a list of all detected complex signals, see Supplemental Table 2.

Figure 5: Complex-centric detection of COP9 Signalosome variant CSN1/3/8.

Figure 5: Complex-centric detection of COP9 Signalosome variant CSN1/3/8. A For nearly half the CORUM complex hypotheses queried, two or more distinct subunit co-elution signals were detected (See extended experimental procedures). **B** Among the four distinct co-elution signals detected from the eight canonical CSN subunits' chromatograms (here with CSN7A, not CSN7B) are two distinct signals delineating **C** distinct co-elution of holo-CSN (observed at the expected fraction 35) and Mini-CSN CSN1/3/8 (observed eluting offset only one fraction late, F45, of the expected fraction, F44). Expected fractions are estimated from the cumulative sum of one copy per component and external size calibration. Coloring adapted to highlight subversion components and their partitioning across holo- and sub-complex. CSN1/3/8 interact and form a submodule within the CSN holo-complex structure (PDB accession 4D10). **D** CSN1/3/8 display distinct fractionation patterns in co-fractionation experiments performed in other laboratories, specifically in orthogonal ion exchange fractionation of HEK293 lysates (Wan et al. 2015, upper panels) and size exclusion chromatographic fractionation of U2OS lysates (Larance et al. 2016, lower panel), in line with the CSN1/3/8 as distinct entity.

Figure 6: Complex-centric detection of 20S proteasome assembly intermediates

Figure 6: Complex-centric detection of 20S proteasome assembly intermediates.

Figure 6: Complex-centric detection of 20S proteasome assembly intermediates. A Protein-level SEC chromatograms of the 22 canonical 26S proteasome subunits. Vertical black lines indicate the apexes of six distinct co-elution signals detected in complex-centric scoring; two of which represent well-known co-occurring variants, the full 26S (i) and the 20S (ii) particle devoid the 19S lid and ATPase (Indicated by structural models, PDB accession 5GJR) and four of which, composed of predominantly 20S α and β subunits, appear at reduced size (222 – 107 kDa, fractions 39, 40, 42 and 46). **B** Zoom in to chromatograms of 20S components in full and reduced MW range and in the context of chaperones known to be involved in assembly according to the current model of 20S biogenesis (lower panel, assembled after Saeki & Tanaka, 2012[38] and PDB accession 5GJR), colored by protein class. Reduced MW species are classified into early and late assembly intermediates (as opposed to artifacts of disassembly) by defined co-elution of early assembly chaperone PSMG3/PSMG4 dimer, late assembly chaperone proteasome maturation protein POMP and constitutive chaperone PSMG1/2 dimer. **C** Subunit mass distribution across early and late assembly intermediate elution ranges suggests predominant components of the intermediary species accumulating in HEK293 cells.

Figure 7: SECexplorer tool for customized interrogation of SEC-SWATH elution profiles.

Figure 7: SECexplorer tool for customized interrogation of SEC-SWATH elution profiles. A SECexplorer web-interface for querying custom protein sets for co-elution behavior in the SEC-SWATH-MS data, viewing chromatograms for interpretation and with algorithmic assistance. **B** Zoom in to high MW peak shoulder of holo-COP9 Signalosome (compare Figure 5), where defined co-elution signals of CSN substrate components CUL4A and DDB2 suggest the partial resolution of substrate-bound and free pools of CSN holo-complex. **C** Estimation of the fraction of holo-CSN in the likely substrate-bound pool versus the free pool, with eight measurements along the eight subunits and based on Gaussian deconvolution of two signals underlying the observed peak and shoulder (also see Figure S5).