The return of the trophic chain: fundamental vs realized interactions in a simple arthropod

food web

Inmaculada Torres-Campos[1], Sara Magalhães[2], Jordi Moya-Laraño[2], Marta Montserrat[1]

[1]Instituto de Hortofruticultura Subtropical y Mediterránea "La Mayora", Universidad

de Málaga, Consejo Superior de Investigaciones Científicas (IHSM-UMA-CSIC),

Avda Dr. Weinberg s/n, Algarrobo-Costa, 29750 Málaga, Spain.

[2]cE3c: Centre for Ecology, Evolution and Environmental Changes, Faculdade de Ciências,

Universidade de Lisboa, Campo Grande, Edifício C2, 1749-016 Lisboa, Portugal.

[3]Estación Experimental de Zonas Áridas – CSIC, Carretera de Sacramento s/n, La Cañada de

San Urbano, 04120 Almería, Spain.

Abstract

The mathematical exploration of small assemblages of interacting species (community modules) has proven key to understand emergent properties of ecological communities. These models use differential equations to study pairwise relations between species. However, as community modules become more complex one may wonder whether all potential interactions are effectively realized. Here, we use community modules to experimentally explore whether the number of trophic links among species increases as another aspect of complexity increases; i.e., the number of species that are known to feed on each other in pair-wise trials. To this aim we use a simple mite community present in avocado orchards (*Persea americana*) composed of two predators (*Euseius stipulatus* and *Neoseiulus californicus*), one herbivore as shared prey (*Oligonychus perseae*), and pollen of *Carpobrotus edulis* as alternative food, with the potential for (intraguild) predation and (apparent) competition to be expressed. Using a series of controls, we could assess whether the presence of one species affected the numbers of another, or its conversion of food into offspring. We found that increasing the number of potential interactions did not result in realized more complex community modules. Instead, all communities were reduced to one or two linear trophic chains. Our results show that trophic links assumed to occur when species are confronted in pairs do not necessarily occur when other components of the community are present. Consequently, food web complexity in terms of connectance may be erroneously over-estimated in theoretical community modules that are parameterized based on pair-wise interactions observed when alternative prey is not present.

Introduction

Community ecology has traditionally viewed trophic interactions as linear chains, with an upper level controlling the densities of the level immediately below, generating a trophic cascade (Hairston *et al.* 1960; Oksanen *et al.* 1981; Carpenter *et al.* 1985). However, it is by now accepted that most communities do not follow this pattern as organisms are imbedded in complex food webs, blurring the notion of a trophic guild and the notion that widespread omnivory would destabilize food webs (Polis & Holt 1992; Polis & Strong 1996). Food webs can be decomposed into community modules (Holt 1997), and among those, intraguild predation, in which two consumers not only compete for a shared resource but also engage in predator-prey interactions (Polis *et al.* 1989), and apparent competition, in which two non-competing prey share a common predator (Holt 1977) are the most common (Bascompte & Melián 2005).

Whether and how often species engage in intraguild predation or apparent competition strongly affects community stability and composition. Indeed, theory predicts that intraguild predation destabilizes communities (Holt & Polis 1997). Some factors may, however, reduce such instability by promoting species coexistence, which generally occurs via a reduction in the strength of intraguild predation. For example, stability is enhanced if predators engage into intraguild predation only when competition for the shared prey is high (Křivan 2000), or the shared prey is less profitable than the intraguild prey (Křivan & Diehl 2005). Other studies suggest that the inclusion of habitat structure (Janssen *et al.* 2007), inducible defences (Kratina *et al.* 2010; Nakazawa *et al.* 2010) or temporal refuges (Amarasekare 2008) increases system stability, although this may depend on which species use refuges (Liu & Zhang 2013). Moreover, stage structure in the intraguild prey promotes stability, either by providing a stage refuge (Mylius *et al.* 2001; Rudolf & Armstrong 2008) or by inducing ontogenetic niche shifts in the predator (Hin *et al.* 2011). However, in all cases, the models still predict that overall stability is lower than that of a simple trophic chain. This lack of stability is corroborated by empirical laboratory studies (Diehl & Feißel 2000; Montserrat *et al.* 2008b), but runs counter the ubiquity of intraguild predation and trophic level omnivory in natural systems (Bascompte & Melián 2005; Gagnon *et al.* 2011). The occurrence of intraguild predation is also expected to lead to a less efficient control of the shared prey populations (Mylius *et al.* 2001). This is especially important in biological control systems, in which the introduction of several biocontrol agents may in fact reduce pest

control. Curiously, empirical studies, mostly stemming from such systems, show that intraguild predation may have variable effects on populations of the shared prey (e.g., Rosenheim *et al.* 1995; Janssen *et al.* 2006; Vance-Chalcraft *et al.* 2007; Messelink & Janssen 2014). This discrepancy between theory and data suggests that some assumptions of theoretical models are not met in natural systems.

Models of intraguild predation consider that the intraguild predator, the intraguild prey and the shared prey form an isolated community module. Although this certainly increases the tractability of models and of their respective experimental tests, it is a significant simplification of real ecosystems. To bring models closer to real systems, while maintaining tractability, researchers have tested how the incorporation of an alternative food source affected stability (Heithaus 2001; Daugherty *et al.* 2007; Holt & Huxel 2007; Rudolf 2007). The general prediction is that providing alternative food to the intraguild prey leads to wider parameter regions of species coexistence, whereas alternative food for the intraguild predator destabilizes the community (Daugherty *et al.* 2007; Holt & Huxel 2007). However, in the latter case, if the quality of the alternative food is high enough, then the intraguild predator switches to feeding on the alternative food, whereas the intraguild prey feeds on the shared prey (Daugherty *et al.* 2007; Holt & Huxel 2007). This again promotes coexistence, by bringing the community structure closer to two linear food chains. Therefore, it is crucial to determine on which resources do species in a community actually feed upon.

Here, we explore how pairwise interactions between species are modified by the inclusion of other species in a simple community. Our baseline hypothesis is that increasing an aspect of complexity in the community module, i.e., the richness of species that are known to interact when no alternative food is provided will increase the number of realized links in the more complex community (Box 1A). We mimicked different community modules of increasing complexity using a community composed of two predatory mite species as intraguild predators (*Euseius stipulatus* and *Neoseiulus californicus*, Acari: Phytoseiidae), one species of herbivore mite as the shared prey (*Oligonychus perseae*, Acari: Tetranychidae), and pollen (of several anemophilous species deposited on leaves) as alternative food (González-Fernández *et al.* 2009), all of which occur in the leaves of crops of avocado (*Persea americana*) in Southestern Spain. From pairwise experimental designs we know that predator-herbivore trophic interactions are stronger when *N. californicus* is the predator

than when it is *E. stipulatus* (González-Fernández et al. 2009); that pollen is an optimal food

source for *E. stipulatus* but not for *N. californicus* (Ferragut et al. 1987; González-Fernández

et al. 2009); and that *E. stipulatus* and *N. californicus* engage in size-dependent predator-

prey interactions (Abad-Moyano *et al.* 2010). This knowledge was used to build predictions

on realized trophic links occurring in our system across community modules of increasing

complexity (Box 1B), that is: *i*) in "trophic chain" community configurations, both predator

species will interact with the herbivore (Box 1B, a.1.1. and a.1.2.); *ii*) in "apparent

competition" community configurations, only *E. stipulatus* will interact with both the

herbivore and pollen (Box 1B, b.1.1. and b.1.2.); *iii*) in "intraguild predation" community

configurations, both IG-predator species will interact with the IG-prey and the herbivore

(Box 1B, c.1.1. and c.1.2.); and iv) in "Intraguild predation and apparent competition"

community configurations, only adults and juveniles of *E. stipulatus* will establish trophic

links with pollen (Box 1B, d.1.1. and d.1.2.). These predictions were tested through a series

of experimental treatments in which we assessed what interactions were realized within

each community module, by measuring IG-prey/herbivore mortality and how predation

translates into predator fecundity as a result of the interactions that were occurring.

Specifically, we examined a) whether (IG)predators feed on each prey type; b) whether

predation of (IG)predators on one prey type is affected by the presence of the other; c)

whether predation of (IG)predators on both prey, and of IG-prey on the herbivore, is

affected by the presence of alternative food; d) whether the presence of alternative food

affects predation of (IG)predators on the two types of prey when they are together; e) yield

egg production produced by feeding on each prey type in (IG)predators; and f) whether egg-

production is additive when (IG)predators have more than one food type available.

Material and Methods

129   All cultures and experiments were done in a climate chamber at 25±1ºC, 65±5% RH and

130   16:8h L:D (Light:Dark).

Mite cultures

132   Cultures of the predatory mite *E. stipulatus* were started in 2007 from ca. 300 individuals

133   collected from avocado trees located in the experimental station of "La Mayora". Rearing

134   units consisted of three bean plants (*Phaseolus vulgaris* L.) with 6-10 leaves, positioned

vertically, with the stems in contact with sponges (*ca.* 30 x 20 x 5 cm) covered with cotton wool and a plastic sheet (27 x 17 cm), and placed inside water-containing trays (8 L, 42.5 x 26 x 7.5 cm). The plant roots were in contact with the water, and the aerial parts were touching each other, forming a tent-like three-dimensional structure, where individuals could easily walk from one plant to the other. Cotton threads were placed on the leaves, to serve as oviposition sites for mite females. Mites were fed *ad libitum* twice a week with pollen of *Carpobrotus edulis* (cat´s claw) spread on leaves with a fine brush. *Euseius stipulatus* is able to develop and reproduce on this food source (Ferragut *et al.* 1987). Every three weeks, new rearings were made by transferring, from old rearings to a new unit, leaves with mites and the cotton threads filled with eggs. The culture was found to be contaminated a few times with *Tyrophagus* spp., a detritivorous mite species. In such instances, instead of moving entire leaves, adult *E. stipulatus* females (ca. 300) were collected individually and transferred to the new rearing unit.

The *N. californicus* population was obtained from Koppert Biological Systems S.L. in bottles of 1000 individuals (Spical®). Colonies were kept on detached bean leaves infested with *Tetranychus urticae* that were placed on top of inverted flower-pots (20 cm Ø) inside water-containing trays.

The herbivore *Oligonychus perseae* was not maintained in a laboratory culture due to technical difficulties in maintaining detached avocado leaves. They were thus collected from the field on a regular basis from avocado orchards located in the experimental station of "La Mayora".

Pollen of *C. edulis* was obtained from flowers collected in the experimental station. Stamens dried in a stove at 37ºC for 48h, then sieved (350 µm).

Community modules

Experimental arenas to test the outcome of community modules consisted of avocado leaf discs placed in petri dishes with a circular hole in the middle, as described in Guzmán *et al.* (2016a). Briefly, a hole (6.5 cm Ø) was cut in a petri dish (9 cm Ø) turned upside down and filled with an avocado leaf disc (7.5 cm Ø), with the borders glued to a clay ring. Inside the petri dish, wet cotton wool ensured enough humidity to keep leaves turgid. Petri dishes were then sealed with parafilm®. To prevent individuals from escaping, a ring of Tanglefoot® was applied along the outer margin of the leaf disc. Female and juvenile predators used in

the experiments were 10-14 and 2-3 days old since egg hatching, respectively. Cohorts of *E.*

stipulatus were made by transferring with a fine brush 400 eggs from the rearings to 2-3

bean leaves placed on top of sponges (30 x 20 x 5 cm, approx.) covered with cotton wool,

inside water-containing trays (3.5 L), and with pollen of *C. edulis* as food. Because it was

difficult to obtain eggs of *N. californicus* from the rearings, cohorts of this species were

made by transferring 100 females to 2-3 bean leaves infested with *Tetranychus urticae*

placed on containers similar to those used for the cultures. After allowing *N. californicus*

females to feed and oviposit for 48 h, they were removed with a fine brush. Before the

experiments, gravid predator females were randomly taken from the cohorts, and starved

for 16 h in experimental containers similar to those above. Starvation was done to

standardize hunger among individuals, and to ensure that egg production in tested females

was not obtained from food ingested prior to the experiment. Predator juveniles were taken

from the cohorts when needed. Arenas containing the herbivore were done as follows: Ten

females of *O. perseae* were let to build nests and lay eggs on experimental arenas during 4

days. The number of nests and eggs laid per nest on each arena was counted at the onset of

the experiment. Pollen in arenas assigned to treatments with alternative food was supplied

ad libitum, using a fine brush.

We performed experiments using two 'community blocks', depending on the identity

of the top predator (*N. californicus* or *E. stipulatus*). Increased complexity in each of the two

community blocks was mimicked through the combination of the presence / absence of 4

factors: predator/IG-predator, IG-prey, herbivore and alternative food. This resulted in the

community modules depicted in the X-axis of figures 1 and 2. These modules were: <u>Trophic</u>

<u>chain</u>: either one *E. stipulatus* or *N. californicus* female was introduced in arenas containing

10 females of *O. perseae* (treatment # 1 in Fig 1 and 2). Arenas containing either one *E.*

stipulatus or one *N. californicus* female without herbivores (treatment # 2), and containing

10 *O. perseae* females without predators (treatment # 3) were done as controls for predator

oviposition rate and prey natural mortality, respectively. <u>Apparent competition</u>: arenas

consisted of one female of either *E. stipulatus* or *N. californicus*, 10 females of *O. perseae*,

and pollen of *C. edulis* supplied *ad libitum* (treatment # 4)*.* Similar arenas but without the

herbivores (treatment # 5) were made as controls for oviposition rates of predators on

pollen only, and without the IG-predator (treatment # 6) to assess potential effects of pollen

on the survival of the herbivore. <u>Intraguild predation</u>: Because IGP is usually associated with

size differences between contestants IG-predators and IG-prey consisted of adult females and heterospecific juveniles, respectively. Arenas consisted of 10 *O. perseae* females, either one *E. stipulatus* or *N. californicus* female, acting as the IG-predators, and 10 heterospecific juveniles, acting as the IG-prey (treatment # 7). Additionally, several control treatments were done to evaluate: the predation/mortality rate of *O. perseae* in the presence of IG-prey but not of IG-predator (treatment # 8); the mortality rate of IG-prey in the absence of both IG-predator and prey (treatment # 9), and in the presence of IG-predator but not of herbivores (treatment # 10). <u>Intraguild predation - Apparent competition</u>: Arenas consisted of 10 *O. perseae* females, either one *E. stipulatus* or *N. californicus* female, acting as the IG-predators, 10 heterospecific juveniles, acting as the IG-prey, and pollen of *C. edulis* as alternative food, supplied *ad libitum* (treatment # 11). Similar arenas to those above but i) without IG-predators (treatment # 12), ii) without herbivores (treatment # 13), and iii) without IG-predators and herbivores (treatment # 14), were done to evaluate predation of IG-prey on the herbivore in the presence of pollen, predation of IG-predators on IG-prey in the presence of pollen, and mortality of IG-prey in the presence of pollen, respectively.

Twenty-four hours later, the number of dead herbivores/IG-prey (predation/mortality rate), and the number of eggs laid by predators/IG-predators (oviposition rate), were recorded. Each treatment was replicated between 10 to 18 times.

Data analyses

Analyses were done separately for communities where either *E. stipulatus* or *N. californicus* acted as the top-predator. Mortality rates of herbivores and IG-prey, and rates of oviposition of IG-predators, were analysed using Generalized Lineal Models (GLM) assuming a Poisson distribution and a Log link function, as no overdispersion of the data was detected. All the analyses were 3 full-factorial designs (see below). We followed a backward elimination procedure as follows: when the interaction among the three explanatory variables was not significant (and the model had higher AIC), this interaction was removed from the model. Subsequently, the same procedure was followed for second order interactions. GLM analyses were performed using the computer environment R (R Core Team 2017). After significance of general models, additional software (package "contrast") was used to perform planned comparisons to address specific questions (see Results). When

specific sets of data were used in multiple comparisons, their significance was corrected

using the sequential Bonferroni method.

Mortality of *O. perseae* females was analysed using data from treatments containing

this species. The model included the presence/absence of IG-predators, IG-prey and

alternative food as explanatory variables, as well as their interactions.

IG-prey mortality was analysed using data from treatments containing predator

juveniles. The full model included the presence/absence of the IG-predator, the herbivore

and alternative food as explanatory variables, as well as their interactions.

Oviposition rates were analysed using data from treatments containing IG-predators.

The full model included the presence/absence of the herbivore, the IG prey and alternative

food as explanatory variables, as well as their interactions.

Results

Communities with E. stipulatus as the (IG-)predator

Mortality rates of the herbivore and of the IG-prey were affected by all three main factors

(Table 1a and b) except pollen in the case of the IG-prey (Table 1b), and all the double

interactions except that between IG-predator and pollen, in the case of the herbivore (Table

1a), and that between the herbivore and pollen, in the case of the IG-prey (Table 1b).

Planned comparisons revealed a) that *E. stipulatus* preyed on *O. perseae* (t_{81} = 2.74, *P* =

0.0076; Fig 1a, compare bars 1 and 3) and marginally on *N. californicus* juveniles (t_{80} = -2.01,

P = 0.048, not significant after Bonferroni correction; Fig 1b, compare bar 9 to 10) when

each prey was offered alone; b) that adding IG-prey increased mortality of *O. perseae* (t_{81} = -

2.26, *P* = 0.026; Fig 1a, compare bar 1 to 7), but adding *O. perseae* did not influence

mortality of the IG-prey (t_{80} = -0.31, *P* = 0.755; Fig 1b, compare bar 10 to 7); c) that the

presence of pollen yielded a drastic reduction in predation of IG-predators on both the

herbivore (t_{81} = 2.99, *P* = 0.0037; Fig 1a, compare bar 1 to 4) and the IG-prey (t_{80} = 3.91, *P* <<

0.001; Fig 1b, compare bar 10 to 13); d) that when both prey were available, the presence of

pollen did not affect herbivore mortality (t_{81} = 0.88, *P* = 0.379; Fig 1a, compare bar 7 to 11),

but did lead to lower IG-prey mortality (t_{80} = 3.58, *P* << 0.001; Fig 1b, compare bar 7 to 11).

Oviposition rates of *E. stipulatus* were only affected by the presence of pollen (main

factor Pollen, Table 1c). However, further planned comparisons revealed that while feeding

on the herbivore yielded some egg production (t_{96} = 2.19, P = 0.021; Fig 1c, compare bar 1 to

2), feeding on IG-prey did not (t_{96} = -1,13, P = 0.259; Fig 1c, compare bar 10 to 2).

Communities with N. californicus as the (IG-)predator

Herbivore mortality was affected by all second order interactions (Table 2a), and that of the

IG-prey (juveniles of *E. stipulatus*) was only affected by the presence of pollen (Table 2b).

Paired comparisons revealed that a) *N. californicus* preyed on *O. perseae* (t_{90} = 3.32, P =

0.013; Fig 2a, compare bar 3 to 1) but not on *E. stipulatus* juveniles (t_{86} = -1.35, P = 0.182; Fig

2b, compare bar 9 to 10), when each prey was offered alone; b) adding IG-prey reduced

mortality of *O. perseae* (t_{90} = 2.56, P = 0.012; Fig 2a, compare bar 1 to 7), but adding *O.*

perseae did not change mortality of the IG-prey (t_{86} = -0.93, P = 0.353; Fig 2b, compare bar

10 to 7); c) the presence of pollen did not affect mortality of either *O. perseae* (t_{90} = -0.43, P

= 0.669; Fig 2a, compare bar 1 to 4) or the IG-prey (t_{86} = 1.80, P = 0.075; Fig 2b, compare bar

10 to 13); d) when both types of prey were available, the presence of pollen led to a

significant increase in mortality of *O.perseae* (t_{90} = -3.65, P << 0.001; Fig 2a, compare bar 7

to 11), but a significant decrease of mortality in *E. stipulatus* juveniles (t_{86} = 2.04, P = 0.044;

Fig 2b, compare bar 7 to 11).

Oviposition rates of *N. californicus* were affected by the main factor Herbivore and

the interaction between the IG-prey and pollen (Table 2c). Indeed, paired comparisons

revealed that e) eggs were produced when *N. californicus* was offered the herbivore alone

(t_{104} = 2.45, P = 0.016; Fig 2c, compare bar 1 to 2), but not when they were on arenas with

either the IG-prey (t_{104} = 0.01, P = 0.992; Fig 2c, compare bar 10 to 2) or pollen (t_{104} = -0.15,

P = 0.884; Fig 2c, compare bar 5 to 2) alone. Moreover, in the presence of the herbivore,

rates of oviposition were not influenced by the presence of pollen (t_{104} = -0.93, P = 0.352; Fig

2c, compare bar 1 to 4), but dramatically decreased in the presence of the IG-prey (t_{104} =

2.39, P = 0.019; Fig 2c, compare bar 1 to 7). However, when pollen was added to the system

with both prey types, IG-predators resumed oviposition to its maximum (t_{104} = -2.36, P =

0.020; Fig 2c, compare bar 7 to 11).

Discussion

In this study, we tested the effect of community structure on the realized interactions

within a community of predatory and herbivorous mites. Because in our system the

intraguild predator is the largest individual within a pair (as in most systems), we created

communities in which adults belonged to one species and juveniles to the other, then

inverted the species-stage identity in another set of communities. We then measured

predation and oviposition in communities with all possible combinations of the presence of

shared prey, the intraguild prey, the intraguild predator and alternative food. We show that

adding species to a community increases the number of potential interactions, but not

necessarily their occurrence. Indeed, despite the potential for apparent competition and

intraguild predation in modules with several species, all modules could be described by

linear food chains (Box 1C).

Although *N. californicus* killed more *O. perseae* females per day, oviposition rates

were similar between predators. This is in line with the finding that *E. stipulatus* can only

forage on mobile *O. perseae* mites when they wander outside nests, whereas *N. californicus*

can penetrate inside nests and forage on all the individuals residing within (González-

Fernández *et al.* 2009). This suggests that *E. stipulatus* is the most efficient predator

converting prey into eggs, but that *N. californicus* is more efficient at reducing herbivore

populations. Which of these strategies is best for biological control will depend on the

ecological condition: if bursts of prey are confined in time, it may be more efficient to select

a biocontrol agent that feeds more, as in "inundative" biocontrol strategies, whereas

controlling and keeping resident populations at low levels may be best achieved with a

predator with a strong numerical response, as in "innoculative" biocontrol strategies (Van

Driesche *et al.* 2007).

Moreover, unlike *N. californicus*, *E. stipulatus* fed and oviposited on pollen. This may

allow the latter to remain in the field for longer periods, as actually observed in field surveys

(González-Fernández *et al.* 2009). Such temporal segregation may facilitate the presence of

the two predators in the same fields (Otto *et al.* 2008). Moreover, our results revealed

asymmetry in the intraguild predation between. *E. stipulatus* and *N. californicus*, with adults

of the latter preying upon juveniles of the latter, but not the reverse. Because *N. californicus*

is likely the best competitor for the shared prey (González-Fernández *et al.* 2009),

coexistence between predators is thus possible in this system (Holt & Polis 1997).

The simultaneous presence of the two predators is likely to have little effect upon

the densities of the shared prey. Indeed, whereas adding adults of *E. stipulatus* to an arena

with *N. californicus* juveniles results in higher prey densities as compared to the presence of

N. californicus alone, the reverse is not true when adding juveniles of *E. stipulatus* to an

arena with adult *N. californicus*. Thus, the net effect of these interactions upon prey density

is probably negligible. This is corroborated by field studies (Montserrat *et al.* 2013).

However, the presence of alternative food (i.e. pollen) contributed to reduce trophic

interactions between predator species resulting in community configurations that could

enhance pest control. Thus, supplying alternative and preferred food to the IG-predator is

probably detrimental to populations of *O. perseae*. Again, this finding mirrors field data

(Montserrat *et al.* 2013).

By combining data of mortality and oviposition at different community structures,

we could recover who eats whom in a relatively complex food web. Although this approach

is very powerful, it does have its limitations. Indeed, it assumes additive effects of

conversion efficiencies of pairwise interactions. For example, if feeding on a prey item

allows predators to better convert the food provided by another prey, this cannot be

detected in our approach (i.e., indirect effects on conversion efficiency). Furthermore, it

may be largely unfeasible to extend this approach to more complex food webs, although it is

becoming clear that we need to know how food is transformed into predator offspring in

order to fully understand food webs in nature (Neutel & Thorne 2014). Indeed, such full-

factorial studies are extremely rare in the literature (but see Schmitz & Sokol-Hessner 2002;

Otto *et al.* 2008).

Food web ecology predicts that species of the highest trophic levels engage in

trophic interactions on more than one food source when these are available. Here, we show

that *E. stipulatus* feeds on the herbivore, *O. perseae*, on the intraguild prey, *N. californicus*,

and on the alternative food, pollen, when each of these are presented alone. However, in

the presence of pollen *E. stipulatus* reduces predation rates on both prey species. This may

be explained by the fact that pollen is the most profitable food for this species, as found

here and in other studies (Ferragut *et al.* 1987; McMurtry & Croft 1997; Bouras &

Papadoulis 2005; González-Fernández *et al.* 2009). Similarly, *N. californicus* adults and

juveniles ceased foraging on other food sources in presence of the herbivores. These results

suggest that realized interactions hinge on the presence of the most profitable food source.

In presence of the optimal food source for each of the two secondary consumers,

communities tended to melt down to become two simple trophic chains. Indeed, in the

most complex communities studied here, with all 5 species present, the presence of the

optimal food originated the split of the community into two trophic chains, one with *E.*
stipulatus feeding on pollen and the other with *N. californicus* feeding on the herbivore (Box
1 d), compare d.1.1. and d.1.2. with d.2.1. and d.2.2.).

Another factor that contributed to the linearization of the food web was that, when
both the IG-prey and the shared prey were together, *E. stipulatus* preyed mainly on the IG-
prey. Indeed, mortality of *O. perseae* in presence of the IG-prey, *N. californicus*, was not
affected by the presence of *E. stipulatus*. Furthermore, mortality of *N. californicus* juveniles
was significantly higher in treatments with presence of the IG-predator, compared to the
control without them. This suggests that mortality in the herbivore was mainly inflicted by
juveniles of *N. californicus*, and that *E. stipulatus* females preyed preferentially on juveniles
of *N. californicus*. This could be explained by *E. stipulatus* having no access to *O. perseae*
eggs or females located inside the nests (Montserrat *et al.* 2008a; González-Fernández *et al.*
2009), which leads to higher encounter rates between *E. stipulatus* and *N. californicus* than
between *E. stipulatus* and *O. perseae*. *Neoseiulus californicus* juveniles, however, can
penetrate *O. perseae* nests, and thus may feed on them. Therefore, the realized community
was that of a 4-level trophic chain (Box 1, c.2.1.). In the other community block, when *N.*
californicus acted as the IG-predator, mortality of *O. perseae* females was similar in all
communities with the IG-prey, irrespective of the presence of IG-predators. Furthermore,
mortality of IG-prey did not differ between treatments with and without the IG-predator,
indicating that *N. californicus* females did not forage on *E. stipulatus* juveniles. These results
suggest that, in presence of IG-prey (juveniles of *E. stipulatus*), the IG-predator ceased to
forage on either herbivore or IG-prey. Thus, the realized community was that of a trophic
chain composed of the IG-prey, the herbivore and the plant, with the IG-predator not
interacting at all (Box 1, c.2.2.). This can be explained by *N. californicus* females avoiding
foraging on a patch where its offspring (future) IG-predator is also there. In line with this,
Abad-Moyano *et al.* (2010) reported that the presence of *E. stipulatus* immatures exerted
non-lethal IG-effects on *N. californicus* females, causing daily oviposition to decrease over
time despite the availability of the shared prey was kept constant. In any case, here, the
trophic links are again linear, with *N. californicus* being excluded from the realized
community (Box 1, c.2.2.). Together, our results show that none of the complex
communities was actually realized, they were all trophic chains.

Our results presented may contribute to solving the paradox of stability in systems
with intraguild predation. Indeed, theoretical models exploring persistence in three-species
communities with IGP find a rather limited room for coexistence of IG-predator and IG-prey
(e.g. Mylius *et al.* 2001), but field observations show that IGP is actually widespread (Polis
1991). Our results suggest that IGP in some systems might actually be occasional, as
predators will tend to forage on the most profitable food, which generally is not the IG prey
(Polis *et al.* 1989). In line with this, some natural systems have shown that communities with
IGP actually show dynamics that are compatible with linear food chains, rather than with
IGP (Borer *et al.* 2003). Therefore, predators may coexist because they rarely engage in IGP,
and complexity may be over-estimated (Magalhães *et al.* 2005). In line with this, Hiltunen *et*
al. (2014) found long-term cycling dynamics when modelling a three-species planktonic food
web, with interaction strength between IG-predator and IG-prey set to be much weaker to
that between IG-predator and the shared resource.

It is becoming increasingly clear that connectance, that is, the number of realized
interactions in a food web, is generally much lower than the number of potential
interactions (Beckerman *et al.* 2006). Identifying trophic links in food webs, however, is not
a simple task. Molecular methods are useful to process field data and they deliver reliable
information on who eats whom, but such tools only provide semi-quantitative estimates of
predation, and they are expensive (Birkhofer *et al.* 2017). Another possible approach to
measure connectance is by observations in the field (Dunne *et al.* 2002; Tylianakis *et al.*
2007; Carnicer *et al.* 2009; Lazzaro *et al.* 2009; Plein *et al.* 2013; Baiser *et al.* 2016; Lemos-
Costa *et al.* 2016). Although this approach allows including a high number of species in the
observations, it suffers from two main shortfalls: (a) it is generally only possible to
undertake it in systems with two trophic levels in which one are primary producers (but see
Bukovinszky *et al.* 2008; Neutel & Thorne 2014), or in systems where trophic interactions
are detectable long after the actual events, as in parasitoid/host interactions or in the direct
study of gut contents; and (b) it does not account for how foraging on a given resource
translates into consumer offspring (but see Bukovinszky *et al.* 2008; Vázquez *et al.* 2015).
Observations in controlled experimental settings, in contrast, deliver quantitative estimates
of predation rates and concomitant offspring production, especially when trophic links, and
their strength, are estimated by confronting pairs of species. Alternatively, modelling
complex systems provide relative estimates on interaction strengths that go beyond pair-

wise interactions (Moya-Laraño *et al.* 2012; Moya-Laraño *et al.* 2014). Yet, one-on-one

approaches may ignore emergent indirect effects of having several species together

(Wootton 1994; Dambacher & Ramos-Jiliberto 2007). For instance, *Cancer productus*, a crab

native to the pacific Northwest, consumes equal amounts of native oysters and of invasive

drill oysters when each type of prey is offered alone, but when they are offered together

crabs interact with the native oyster species only (Grason & Miner 2012). Therefore, if

trophic links are not evaluated in presence of all species in the community, one may reach

erroneous conclusions on the strength of the interaction (Guzmán *et al.* 2016b; Fonseca *et*

al. 2017) and overestimate connectance in food webs. In this work, through the addition of

species we created a series of communities of increased number of species and increased

number of potential interactions (links) among them, both direct (trophic) or indirect

(competition, apparent competition). However, our experimental results revealed that all

communities ended up becoming a sum of one or more trophic chains (Box 1C). These

results are consistent with the idea that the fundamental trophic niche of a species (i.e., the

food items that the species is able to feed on is larger than the realized one (i.e., the food

items that the species feeds on when they are all present simultaneously – (Hutchinson

1961)). Our results suggest that some food webs may be less complex than previously

thought in terms on the frequency and strength of IGP. Unravelling realized food webs, i.e.

interaction strengths across different nodes and trophic levels, including indirect effects,

may be thus key to understanding these ecological networks and their persistence.

Acknowledgements

The authors are indebted to Rosa María Sahún Logroño for her valuable help. This work was

financed by the Spanish Ministry of Science and Innovation (CGL2015-66192-R and

AGL2011-30538-C03-03). I.T.C. was recipient of a grant from the Spanish National Research

Council (CSIC, ref: JAE-PRE 041). The animals used for the research of this publication are

not test animals in the legal sense. The authors have no conflict of interest to declare.

References

Abad-Moyano, R., Urbaneja, A. & Schausberger, P. (2010). Intraguild interactions between

Euseius stipulatus and the candidate biocontrol agents of *Tetranychus urticae* in

Spanish clementine orchards: *Phytoseiulus persimilis* and *Neoseiulus californicus*.
Experimental and Applied Acarology, 50, 23-34.

Amarasekare, P. (2008). Coexistence of intraguild predators and prey in resource-rich
environments. *Ecology*, 89, 2786-2797.

Baiser, B., Elhesha, R. & Kahveci, T. (2016). Motifs in the assembly of food web networks.
Oikos, 125, 480-491.

Bascompte, J. & Melián, C.J. (2005). Simple trophic modules for complex food webs.
Ecology, 86, 2868-2873.

Beckerman, A.P., Petchey, O.L. & Warren, P.H. (2006). Foraging biology predicts food web
complexity. *Proceedings of the National Academy of Sciences*, 103, 13745-13749.

Birkhofer, K., Bylund, H., Dalin, P., Ferlian, O., Gagic, V., Hambäck, P.A. *et al.* (2017).
Methods to identify the prey of invertebrate predators in terrestrial field studies.
Ecology and Evolution, 7, 1942-1953.

Borer, E.T., Briggs, C.J., Murdoch, W.W. & Swarbrick, S.L. (2003). Testing intraguild predation
theory in a field system: does numerical dominance shift along a gradient of
productivity? *Ecology Letters*, 6, 929-935.

Bouras, S.L. & Papadoulis, G.T. (2005). Influence of selected fruit tree pollen on life history
of *Euseius stipulatus* (Acari: Phytoseiidae). *Experimental and Applied Acarology*, 36,
1-14.

Bukovinszky, T., van Veen, F.F., Jongema, Y. & Dicke, M. (2008). Direct and indirect effects of
resource quality on food web structure. *Science*, 319, 804-807.

Carnicer, J., Jordano, P. & Melián, C.J. (2009). The temporal dynamics of resource use by
frugivorous birds: a network approach. *Ecology*, 90, 1958-1970.

Carpenter, S.R., Kitchell, J.F. & Hodgson, J.R. (1985). Cascading trophic interactions and lake
productivity. *BioScience*, 35, 634-639.

Dambacher, J.M. & Ramos-Jiliberto, R. (2007). Understanding and predicting effects of
modified interactions through a qualitative analysis of community structure. *The*
Quarterly Review of Biology, 82, 227-250.

Daugherty, M.P., Harmon, J.P. & Briggs, C.J. (2007). Trophic supplements to intraguild
predation. *Oikos*, 116, 662-677.

Diehl, S. & Feißel, M. (2000). Effects of enrichment on three-level food chains with
omnivory. *The American Naturalist*, 155, 200-218.

Dunne, J.A., Williams, R.J. & Martinez, N.D. (2002). Food-web structure and network theory:
the role of connectance and size. *Proceedings of the National Academy of Sciences*,
99, 12917-12922.

Ferragut, F., Garcia-Mari, F., Costa-Comelles, J. & Laborda, R. (1987). Influence of food and
temperature on development and oviposition of *Euseius stipulatus* and
Typhlodromus phialatus (Acari: Phytoseiidae). *Experimental and Applied Acarology*,
3, 317-329.

Fonseca, M.M., Montserrat, M., Guzmán, C., Torres-Campos, I., Pallini, A. & Janssen, A.
(2017). How to evaluate the potential occurrence of intraguild predation.
Experimental and Applied Acarology, 72, 103-114.

Gagnon, A.-E., Heimpel, G.E. & Brodeur, J. (2011). The ubiquity of intraguild predation
among predatory arthropods. *PLoS ONE*, 6, e28061.

González-Fernández, J., De la Peña, F., Hormaza, J.I., Boyero, J., Vela, J., Wong, E. *et al.*
(2009). Alternative food improves the combined effect of an omnivore and a
predator on biological pest control. A case study in avocado orchards. *Bulletin of
Entomological Research*, 99, 433-444.

Grason, E.W. & Miner, B.G. (2012). Preference alters consumptive effects of predators: top-
down effects of a native crab on a system of native and introduced prey. *PloS ONE*, 7,
e51322.

Guzmán, C., Aguilar-Fenollosa, E., Sahún, R.M., Boyero, J.R., Vela, J.M., Wong, E. *et al.*
(2016a). Temperature-specific competition in predatory mites: Implications for
biological pest control in a changing climate. *Agriculture, Ecosystems and
Environment*, 216, 89-97.

Guzmán, C., Sahún, R. & Montserrat, M. (2016b). Intraguild predation between phytoseiid
mite species might not be so common. *Experimental and Applied Acarology*, 68, 441-
453.

Hairston, N.G., Smith, F.E. & Slobodkin, L.B. (1960). Community structure, population
control, and competition. *The American Naturalist*, 94, 421-425.

Heithaus, M.R. (2001). Habitat selection by predators and prey in communities with
asymmetrical intraguild predation. *Oikos*, 92, 542-554.

Hiltunen, T., Ellner, S.P., Hooker, G., Jones, L.E. & Hairston Jr, N.G. (2014). Eco-evolutionary
dynamics in a three-species food web with intraguild predation: intriguingly
complex. In: *Advances in Ecological Research*. Elsevier, pp. 41-73.

Hin, V., Schellekens, T., Persson, L. & de Roos, A.M. (2011). Coexistence of predator and
prey in intraguild predation systems with ontogenetic niche shifts. *The American*
Naturalist, 178, 701-714.

Holt, R. (1997). Community modules. In: *Multitrophic interactions in Terrestrial Systems*
(eds. Gange, A & Brown, V). Blackwell Science London, UK, pp. 333-350.

Holt, R.D. (1977). Predation, apparent competition, and the structure of prey communities.
Theoretical Population Biology, 12, 197-229.

Holt, R.D. & Huxel, G.R. (2007). Alternative prey and the dynamics of intraguild predation:
theoretical perspectives. *Ecology*, 88, 2706-2712.

Holt, R.D. & Polis, G.A. (1997). A theoretical framework for intraguild predation. *The*
American Naturalist, 149, 745-764.

Hutchinson, G.E. (1961). The paradox of the plankton. *The American Naturalist*, 95, 137-145.

Janssen, A., Montserrat, M., HilleRisLambers, R., Roos, A.M., Pallini, A. & Sabelis, M.W.
(2006). Intraguild predation usually does not disrupt biological control. In: *Trophic*
and Guild in Biological Interactions Control (eds. Brodeur, J & Boivin, G). Springer
Netherlands Dordrecht, pp. 21-44.

Janssen, A., Sabelis, M.W., Magalha es, S., Montserrat, M. & van der Hammen, T. (2007).
Habitat structure affects intraguild predation. *Ecology*, 88, 2713-2719.

Kratina, P., Hammill, E. & Anholt, B.R. (2010). Stronger inducible defences enhance
persistence of intraguild prey. *Journal of Animal Ecology*, 79, 993-999.

Křivan, V. (2000). Optimal intraguild foraging and population stability. *Theoretical*
Population Biology, 58, 79-94.

Křivan, V. & Diehl, S. (2005). Adaptive omnivory and species coexistence in tri-trophic food
webs. *Theoretical Population Biology*, 67, 85-99.

Lazzaro, X., Lacroix, G., Gauzens, B., Gignoux, J. & Legendre, S. (2009). Predator foraging
behaviour drives food-web topological structure. *Journal of Animal Ecology*, 78,
1307-1317.

Lemos-Costa, P., Pires, M.M., Araújo, M.S., Aguiar, M.A. & Guimarães, P.R. (2016). Network
analyses support the role of prey preferences in shaping resource use patterns
within five animal populations. *Oikos*, 125, 492-501.

Liu, Z. & Zhang, F. (2013). Species coexistence of communities with intraguild predation: The
role of refuges used by the resource and the intraguild prey. *Biosystems*, 114, 25-30.

Magalhães, S., Tudorache, C., Montserrat, M., Van Maanen, R., Sabelis, M.W. & Janssen, A.
(2005). Diet of intraguild predators affects antipredator behavior in intraguild prey.
Behavioral Ecology, 16, 364-370.

McMurtry, J. & Croft, B. (1997). Life-styles of phytoseiid mites and their roles in biological
control. *Annual Review of Entomology*, 42, 291-321.

Messelink, G.J. & Janssen, A. (2014). Increased control of thrips and aphids in greenhouses
with two species of generalist predatory bugs involved in intraguild predation.
Biological Control, 79, 1-7.

Montserrat, M., de la Peña, F., Hormaza, J. & González-Fernández, J. (2008a). How do
Neoseiulus californicus (Acari: Phytoseiidae) females penetrate densely webbed
spider mite nests? *Experimental and Applied Acarology*, 44, 101-106.

Montserrat, M., Guzmán, C., Sahún, R., Belda, J. & Hormaza, J. (2013). Pollen supply
promotes, but high temperatures demote, predatory mite abundance in avocado
orchards. *Agriculture, Ecosystems & Environment*, 164, 155-161.

Montserrat, M., Magalhães, S., Sabelis, M.W., De Roos, A.M. & Janssen, A. (2008b). Patterns
of exclusion in an intraguild predator–prey system depend on initial conditions.
Journal of Animal Ecology, 77, 624-630.

Moya-Laraño, J., Bilbao-Castro, J.R., Barrionuevo, G., Ruiz-Lupión, D., Casado, L.G.,
Montserrat, M. *et al.* (2014). Eco-evolutionary spatial dynamics: rapid evolution and
isolation explain food web persistence. *Advances in Ecological Research*, 50, 75-144.

Moya-Laraño, J., Verdeny-Vilalta, O., Rowntree, J., Melguizo-Ruiz, N., Montserrat, M. &
Laiolo, P. (2012). Climate Change and Eco-Evolutionary Dynamics in Food Webs.
Advances in Ecological Research, 47, 1-80.

Mylius, S.D., Klumpers, K., de Roos, A.M. & Persson, L. (2001). Impact of intraguild predation
and stage structure on simple communities along a productivity gradient. *The
American Naturalist*, 158, 259-276.

Nakazawa, T., Miki, T. & Namba, T. (2010). Influence of predator-specific defense adaptation
on intraguild predation. *Oikos*, 119, 418-427.

Neutel, A.M. & Thorne, M.A. (2014). Interaction strengths in balanced carbon cycles and the
absence of a relation between ecosystem complexity and stability. *Ecology letters*,
17, 651-661.

Oksanen, L., Fretwell, S.D., Arruda, J. & Niemela, P. (1981). Exploitation ecosystems in
gradients of primary productivity. *The American Naturalist*, 118:2, 240-261.

Otto, S.B., Berlow, E.L., Rank, N.E., Smiley, J. & Brose, U. (2008). Predator diversity and
identity drive interaction strength and trophic cascades in a food web. *Ecology*, 89,
134-144.

Plein, M., Längsfeld, L., Neuschulz, E.L., Schultheiß, C., Ingmann, L., Töpfer, T. *et al.* (2013).
Constant properties of plant–frugivore networks despite fluctuations in fruit and bird
communities in space and time. *Ecology*, 94, 1296-1306.

Polis, G.A. (1991). Complex trophic interactions in deserts: an empirical critique of food-web
theory. *The American naturalist*, 138, 123-155.

Polis, G.A. & Holt, R.D. (1992). Intraguild predation: the dynamics of complex trophic
interactions. *Trends in Ecology & Evolution*, 7, 151-154.

Polis, G.A., Myers, C.A. & Holt, R.D. (1989). The ecology and evolution of intraguild
predation: potential competitors that eat each other. *Annual Review of Ecology and*
Systematics, 20, 297-330.

Polis, G.A. & Strong, D.R. (1996). Food web complexity and community dynamics. *The*
American Naturalist, 147, 813-846.

R Core Team (2017). R: A Language and Environment for Statistical Computing.
https://www.R-project.org/

Rosenheim, J.A., Kaya, H., Ehler, L., Marois, J.J. & Jaffee, B. (1995). Intraguild predation
among biological-control agents: theory and evidence. *Biological Control*, 5, 303-335.

Rudolf, V.H. (2007). The interaction of cannibalism and omnivory: consequences for
community dynamics. *Ecology*, 88, 2697-2705.

Rudolf, V.H.W. & Armstrong, J. (2008). Emergent impacts of cannibalism and size refuges in
prey on intraguild predation systems. *Oecologia*, 157, 675-686.

Schmitz, O.J. & Sokol-Hessner, L. (2002). Linearity in the aggregate effects of multiple
predators in a food web. *Ecology Letters*, 5, 168-172.

Tylianakis, J.M., Tscharntke, T. & Lewis, O.T. (2007). Habitat modification alters the structure
of tropical host–parasitoid food webs. *Nature*, 445, 202.

Van Driesche, R., Hoddle, M. & Center, T.D. (2007). Control de plagas y malezas por
enemigas naturales. US Department of Agriculture, US Forest Service, Forest Health
Technology Enterprise Team.

Vance-Chalcraft, H.D., Rosenheim, J.A., Vonesh, J.R., Osenberg, C.W. & Sih, A. (2007). The
influence of intraguild predation on prey suppression and prey release: a meta-
analysis. *Ecology*, 88, 2689-2696.

Vázquez, D.P., Ramos-Jiliberto, R., Urbani, P. & Valdovinos, F.S. (2015). A conceptual
framework for studying the strength of plant–animal mutualistic interactions.
Ecology letters, 18, 385-400.

Wootton, J.T. (1994). Putting the pieces together: testing the independence of interactions
among organisms. *Ecology*, 75, 1544-1551.

Table 1. Results of Generalized Lineal Models applied to a) herbivore mortality rates, b) IG-prey (juveniles of *N. californicus*) mortality rates, and c) (IG-)predator (females of *E. stipulatus*) oviposition rates. All the analyses were 3 full-factorial designs. When interactions among the three explanatory variables were not significant, and if the new model yielded a lower AIC, they were removed from the model. Subsequently, the same procedure was followed for double interactions. These cases are shown in the table as NS.

a)	Herbivore mortality rates	Estimate	Std. Error	z value	Pr(>|z|)
	Intercept	-1.755	0.712	-2.466	0.014
	IG-predator (1)	2.212	0.732	3.021	0.002
	IG-prey (2)	2.932	0.729	4.023	<0.001
	Pollen (3)	-1.851	0.609	-3.040	<0.001
	IG-predator * IG-prey	-2.302	0.756	-3.047	0.002
	IG-predator * Pollen	NS			
	IG-prey * Pollen	1.573	0.639	2.466	.014
	(1) * (2) * (3)	NS			

b)	IG-prey mortality rates	Estimate	Std. Error	z value	Pr(>|z|)
	Intercept	0.513	0.238	2.156	0.031
	IG-predator (1)	0.591	0.273	2.163	0.030
	Herbivore (2)	-1.624	0.496	-3.276	0.001
	Pollen (3)	-0.392	0.359	-1.091	0.275
	IG-predator * Herbivore	1.552	0.511	3.037	0.002
	IG-predator * Pollen	-1.705	0.517	-3.300	<0.001
	Herbivore * Pollen	0.749	0.520	1.439	0.150
	(1) * (2) * (3)	NS			

c)	IG-predator oviposition rates	Estimate	Std. Error	z value	Pr(>|z|)
	Intercept	-0.843	0.245	-3.443	<0.001
	IG-prey (1)	-0.194	0.220	-0.882	0.378
	Herbivore (2)	0.220	0.216	1.018	0.308
	Pollen (3)	1.104	0.235	4.703	<0.001
	IG-prey * Herbivore	NS			
	IG-prey * Pollen	NS			
	Herbivore * Pollen	NS			
	(1) * (2) * (3)	NS*			

Table 2. Results of Generalized Lineal Models applied to a) herbivore mortality rates, b) IG-prey (juveniles of *E. stipulatus*) mortality rates, and c) (IG-)predator (females of *N. californicus*) oviposition rates. All the analyses were 3 full-factorial designs. When interactions among the three explanatory variables were not significant, and if the new model yielded a lower AIC, they were removed from the model. Subsequently, the same procedure was followed for double interactions. These cases are shown in the table as NS.

a) Herbivore mortality rates	Estimate	Std. Error	z value	Pr(>|z|)
Intercept	-1.954	0.722	-2.707	0.007
IG-predator (1)	2.997	0.729	4.109	<0.001
IG-prey (2)	2.184	0.746	2.927	0.003
Pollen (3)	-0.888	0.499	-1.782	0.075
IG-predator * IG-prey	-2.825	0.764	-3.699	<0.001
IG-predator * Pollen	0.999	0.460	2.175	0.030
IG-prey * Pollen	0.791	0.325	2.436	0.015
(1) * (2) * (3)	NS*			

b) IG-prey mortality rates	Estimate	Std. Error	z value	Pr(>|z|)
Intercept	-0.4855	0.3035	-1.600	0.110
IG-predator (1)	0.6150	0.3152	1.951	0.051
Herbivore (2)	-0.3174	0.2851	-1.114	0.265
Pollen (3)	-1.1505	0.3416	-3.368	<0.001
IG-predator * Herbivore	NS*			
IG-predator * Pollen	NS*			
Herbivore * Pollen	NS*			
(1) * (2) * (3)	NS*			

c) IG-predator oviposition rates	Estimate	Std. Error	z value	Pr(>|z|)
Intercept	-2.7430	0.6172	-4.444	<0.001
IG-prey (1)	-2.5550	1.0378	-2.462	0.014
Herbivore (2)	2.5174	0.5989	4.204	<0.001
Pollen (3)	0.3476	0.3685	0.943	0.346
IG-prey * Herbivore	NS*			
IG-prey * Pollen	2.2175	1.1041	2.008	0.045
Herbivore * Pollen	NS*			
(1) * (2) * (3)	NS*			

Box 1. **A: Fundamental community modules** included in this study. a) trophic chain, b) apparent competition, c) intraguild predation, and d) intraguild predation and apparent competition. From a) to d) the complexity of the community is increased via increasing the number of species and the number of interactions among them. **B: Predicted trophic links** that have been observed using pairwise experimental settings**. C: Realized trophic links** occurring across community modules of increasing complexity, obtained from the experiments presented here, where interactions are measured in the presence of other components of the community. SC stands for secondary consumer, PC for primary consumer, PP for primary producer, and AF for alternative food. SC1 and SC2 are phytoseiid predatory mites, i.e. *Euseius stipulatus* and *Neoseiulus californicus*, respectively, PC is the tetranychid herbivore mite *Oligonychus perseae*, AF is pollen of *Carpobrotus edulis*, and PP is the avocado *Persea americana*. Solid arrows indicate negative direct interactions (who eats whom), whereas dotted and dashed arrows in Box 1A indicate negative indirect interactions (apparent competition and competition).

Figure 1. Mortality rates (average ± S.E.) of a) herbivores (*Oligonychus perseae* females) and b) IG-prey (*Neoseiulus californicus* juveniles), and c) oviposition rates (average ± S.E.) of IG-predators (*Euseius stipulatus* females), in 14 different treatments defined by presence or absence of either IG-predators, IG-prey, herbivores or alternative food (pollen), depicted in the lower part of the figure, that mimicked four different community configurations and their respective controls.

Figure 2. Mortality rates (average ± S.E.) of a) herbivores (*Oligonychus perseae* females) and b) IG-prey (*Euseius stipulatus* juveniles), and c) oviposition rates (average ± S.E.) of IG-predators (*Neoseiulus califonicus* females), in 14 different treatments defined by presence or absence of either IG-predators, IG-prey, herbivores or alternative food (pollen), depicted in the lower part of the figure, that mimicked four different community configurations and their respective controls.

715   Box 1.

A: FUNDAMENTAL COMMUNITY	B: PREDICTED TROPHIC LINKS	C: REALIZED TROPHIC LINKS
a) Trophic chain Secondary Consumer (SCi,j) ↓ Primary Consumer (PC) ↓ Primary Producer (PP)	a.1.1) *E. stipulatus* a.1.2) *N. californicus* (SC1) (SC2) ↓ ↓ (PC) (PC) ↓ ↓ (PP) (PP)	a.2.1) *E. stipulatus* a.2.2) *N. californicus* (SC1) (SC2) ↓ ↓ (PC) (PC) ↓ ↓ (PP) (PP)
b) Apparent competition Secondary Consumer (SCi,j) ↗ ↘ Primary Consumer (PC) Alternative Food (AF) ⋯▶ ↓ Primary Producer (PP)	b.1.1) *E. stipulatus* b.1.2) *N. californicus* (SC1) (SC2) ↗ ↘ ↓ (PC) (PC) (AF) ↓ (AF) ↓ (PP) (PP)	b.2.1) *E. stipulatus* b.2.2) *N. californicus* (SC1) (SC2) ↓ ↓ (AF) (PC) (AF) (PC) ↓ ↓ (PP) (PP)

Figure 1.

719   Figure 2.