The RHIM within the M45 protein from murine cytomegalovirus forms

heteromeric functional amyloid fibrils with RIPK1 and RIPK3

Chi L. L. Pham[1], Merryn Strange[1], Ailis O' Carroll[2], Nirukshan Shanmugam[1], Emma

Sierecki[2], Yann Gambin[2], Megan Steain[3] and Margaret Sunde[1*]

[1]Discipline of Pharmacology, School of Medical Sciences and Sydney Nano, University of

Sydney, NSW 2006, Australia

[2]EMBL Australia Node in Single Molecule Sciences, School of Medical Science, University

of New South Wales, NSW 2052, Australia

[3]Infectious Diseases and Immunology, Central Clinical School, Sydney Medical School,

University of Sydney, NSW 2006, Australia

*Corresponding author: margaret.sunde@sydney.edu.au

Abstract

The M45 protein from murine cytomegalovirus protects infected murine cells from necroptosis and can protect human cells from necroptosis induced by TNFR activation, when heterologously expressed. We show that this activity is associated with amyloid formation by the N-terminal 90 residues of the M45 protein, which contain a RIP Homotypic Interaction Motif (RHIM). The RHIM of M45 drives self-assembly into homomeric fibrils with the hallmark characteristics of amyloid and interacts with the RHIMs of host RIPK1 and RIPK3 to form heteromeric amyloid fibrils. These interactions require an intact core tetrad within the M45 RHIM. The heteromeric amyloid fibrils formed when the M45 RHIM is incorporated reduces the stability of RIPK1 and RIPK3 RHIM-based fibrillar assemblies. Our results suggest that M45 mimics the interactions made by RIPK1 with RIPK3. It represents a novel functional amyloid and anti-RHIM strategy. The entrapment of cellular RHIM-containing proteins in alternative heteromeric complexes that are unable to perform appropriate signaling functions may underlie the suppression of RIPK3-associated necroptosis by murine cytomegalovirus, and other viruses that express RHIM-containing proteins.

Introduction

The RIP homotypic interaction motif (RHIM) was first identified in receptor-interacting protein kinase 1 (RIPK1) and receptor-interacting kinase 3 (RIPK3)[1]. A RHIM contains 18–19 amino acids, with a tetrad (V/I)-Q-(V/I/L/C)-G core sequence (Fig. 1a). An intact RHIM is required for the interaction between RIPK1 and RIPK3 that occurs downstream of tumour necrosis factor receptor 1 (TNFR1) activation, during the programmed cell death response known as necroptosis[2,3]. In addition to the TNFR1 pathway that leads to RIPK1:RIPK3 association, two other pathways involving RIPK3 and specific RHIM-containing adapter proteins can result in necroptosis[4]. Binding of microbial pathogen-associated molecular patterns to Toll-like receptors 3 and 4 can lead to activation of RIPK3 through formation of a complex with the RHIM-containing TIR-domain-containing adapter-inducing interferon-β protein (TRIF). Additionally, the Z-DNA binding protein/DNA-dependent activator of interferon regulatory factors (ZBP-1/DAI) is a cytosolic nucleic acid sensor that binds foreign nucleic acid and acts as the RHIM adapter protein for RIPK3 in viral-induced necroptosis[5-8]. In all three pathways, oligomerisation and autophosphorylation of RIPK3 occurs after binding of RIPK3 to the adapter protein. The activation of RIPK3 leads to phosphorylation and oligomerisation of mixed lineage kinase domain-like protein (MLKL) and this commits the cell to membrane disruption, ion dyshomeostasis and lytic necrotic cell death[9-12].

Necroptosis has essential functions in adult tissue homeostasis and innate immune defence against intracellular pathogens, however undesirable necroptosis can be triggered by ischaemic tissue damage[13,14] and recent evidence suggests that necroptosis contributes to neurodegenerative conditions[15]. Small molecules that block necroptosis are under investigation as potential modulators of ischaemia-reperfusion injury associated with

myocardial infarction, stroke and solid organ transplantation[16]. Solid cancers present with

necrosis but additionally necroptosis can promote tumour cell growth through suppression of

the immune response against cancer[17]. A number of herpesviruses express RHIM-containing

proteins that are primarily implicated in inhibiting host cell necroptosis to ensure virus

survival[18,19]. Understanding the molecular basis for this viral RHIM-driven inhibition could

identify therapeutic targets and provide guidance for the development of anti-necroptosis

strategies with many clinical applications.

We have characterised the properties of the RHIM-containing protein M45 (Fig. 1a), also

known as the viral Inhibitor of RIP Activation (vIRA)[20] from murine cytomegalovirus

(MCMV) and the structural basis for its interactions with RIPK1 and RIPK3. MCMV inhibits

host apoptosis and necroptosis pathways to sustain infection[21,22] and specifically thwarts

necroptosis in infected endothelial cells and macrophages[23,24] through the actions of M45.

Thus far, M45 is the only viral RHIM-containing protein known to suppress necroptosis in

mouse and human cells[23,25,26] . The M45 protein is delivered within the infecting virions[27].

Inhibition of necroptosis by M45 has been shown to involve interactions between M45 and

RIPK1[23], RIPK3[26] and ZBP-1/DAI[7,28] and to require an intact RHIM[7,28].

In 2012 Wu and colleagues demonstrated that the RHIM-driven association between RIPK1

and RIPK3 results in the formation of an amyloid, fibrillar structure where the RHIM

residues form the β-sheet core of RIPK1:RIPK3 fibrils[29]. This necrosome complex was the

first functional amyloid structure described with an active signalling role[30] but with the report

of similar amyloid-based signalling platform in *Drosophila*, where the PGRP-LC and PGRP-

LE receptors and downstream Imd adapter protein interact through motifs bearing some

similarity to RHIMs[31], RHIM-based signalling may be widespread. The structure of the

necrosome RIPK1:RIPK3 core has now been determined by solid state nuclear magnetic

resonance spectroscopy (ssNMR), revealing a compact hydrophobic interface and

highlighting the central role of the RHIM tetrad[32].

Given the demonstrated ability of the M45 protein to protect against TNFR-initiated

necroptosis in human cells[25] and the interest in inhibition of necroptosis in clinical settings,

we have investigated the interactions between the RHIMs of M45 and human RIPK1 and

RIPK3, which are highly homologous to the murine proteins (Fig. 1a). We find that the

RHIM of M45 is itself amyloidogenic and the ability of M45 to self-assemble into fibrils

depends on the presence of the core sequence IQIG. A 90-residue fragment of M45

encompassing the RHIM confers protection against TNF-driven necroptosis in human cells.

The RHIM of M45 is readily incorporated into heteromeric amyloid fibrils with the minimum

portions of human RIPK1 or RIPK3 that are necessary for necrosome formation[29]. The viral

RHIM preferentially assembles with RIPK3 over RIPK1. We have used the recently

determined structure of the RIPK1:RIPK3 core to propose how M45 could form a hetero-

amyloid structure with RIPK3, in a manner analogous to RIPK1, which has the same IQIG

tetrad sequence. Incorporation of the viral M45 RHIM alters the properties of the human

RHIM-based complexes. These results suggest that the M45 protein sequesters host RHIM-

containing proteins in alternative amyloid structures that are incompatible with auto-

phosphorylation of RIPK3 and/or activation of the downstream necroptosis mediator MLKL.

The ability of the M45 protein revealed here, to form amyloid-based structures with host

RHIM-containing proteins, suggests a general mechanism for modulation of host RHIM-

based signalling by other viral RHIM-containing proteins.

Results

The M45 RHIM drives spontaneous amyloid fibril formation

We initially sought to determine whether the RHIM in M45 was itself able to assemble into

amyloid fibrils. A peptide of 19 residues, corresponding to the M45 RHIM sequence, readily

assembled into long fibrils with amyloid morphology (Fig. 1a, b). In order to investigate the nature

of the interactions between M45 and host proteins, we prepared fusion proteins containing portions

of M45, RIPK1 and RIPK3 (Fig. 1a) with different partner proteins (His$_6$-ubiquitin (His-Ub), His$_6$-

YPet, -ECFP or -mCHERRY fluorescent proteins (FPs) or maltose binding protein (MBP)). The

human RIPK protein constructs contained the minimum RHIM-containing regions shown to support

interactions between these two proteins, i.e. RIPK1-(497-583) and RIPK3-(387-518)[29]. Two RHIM-

encompassing fragments of M45 were tested, comprising 90 or 277 N-terminal residues of the viral

protein. The 1–90 region of M45 was identified as a likely autonomous fragment because analysis

of the predicted secondary structure of the protein highlighted a clear boundary between the β-sheet

elements flanking the RHIM and the strongly predicted helical elements from residue 93 onwards.

The 277 N-terminal residues of M45 have previously been shown to provide protection against

some forms of cell death[23] and to inhibit ZBP-1 signalling[7].

RHIM-encompassing fragments of M45 inhibit necroptosis in human HT-29 cells

We compared the ability of the 1–90 and 1–277 regions of M45 to protect against TNFR-induced

necroptosis in human HT-29 cells. The 90-residue and 277-residue portions were expressed as

fusion proteins with the B subunit of *E. coli* DNA gyrase[9]. Addition of the antibiotic coumermycin

A1 can induce dimerization of the gyrase subunit, to substitute for the potential dimerisation of the

inactive ribonucleotide reductase-like (RNR1-like) domain of M45, which may form a heterodimer

with RNR large and small subunits in the intact viral protein. Fig. 1c shows that the 90-residue and

277-residue portions of M45 protect human cells from TNFR-induced necroptosis through the

RIPK1:RIPK3 pathway to a similar extent, and this is independent of the addition of coumermycin

A1. Dimerisation of M45 is therefore unlikely to be necessary for its role in inhibition of host cell

death. For the majority of biophysical experiments we have therefore probed RHIM-driven

interactions using the minimum necrosome-forming regions of RIPK1 and RIPK3[29] and the 90-

residue N-terminal portion of M45.

The wild type core tetrad is critical for amyloid formation by M45 and RIPK1 but not RIPK3

The RHIM-containing regions were expressed as fusion proteins with His-Ub or His-FP to improve

expression yields and to provide a means to study the kinetics of amyloid assembly. These fusion

proteins were purified in the presence of denaturant, which prevented unwanted self-assembly. Prior

experience with other functional amyloid-forming proteins has demonstrated that upon dilution of

the denaturant, the ubiquitin domain refolds independently to its native state, allowing the

characteristics of the partner sequence to be studied[33]. In YPet, ECFP and mCHERRY-containing

RHIM fusion proteins, the fluorescent partner refolded correctly, as judged by recovery of

characteristic fluorescence profiles. Purified His-Ub-M45$_{1-90}$ spontaneously formed fibrils upon

removal of denaturant by dialysis and the fibrils displayed a cross-β X-ray fibre diffraction pattern,

with a strong, sharp meridional reflection at 4.7 Å and a weaker, diffuse equatorial reflection at

~10.5 Å, which are characteristic of amyloid fibrils[34] (Fig. 2a).

The kinetics of self-assembly, or homomeric fibril formation, of the RHIM-containing fusion

proteins were studied by monitoring the fluorescence of thioflavin T (ThT), a dye commonly used

as an indicator of amyloid structure. ThT exhibits increased fluorescence emission at 485 nm when

it binds to amyloid fibrils[35]. All wild type RHIM-containing His-Ub and fluorescent fusion proteins

formed homomeric amyloid (Fig. 2b). Tetra-alanine substitution of the RHIM core is a commonly

used strategy to probe the role of RHIM:RHIM interactions in cells and during viral infection[1,23]

(Fig. 1a). RHIM mutant constructs in which the four core amino acids were substituted by AAAA

were produced and the ability of these proteins to assemble into homomeric amyloid fibrils was

tested. The substitution of the IQIG tetrad by AAAA within the $M45_{1-90}$ and $RIPK1_{497-583}$ RHIM

constructs abolishes amyloid assembly (Fig. 2b). In contrast, the AAAA version of $RIPK3_{387-518}$

retains the ability to assemble into amyloid fibrils and has a similar kinetic profile to the WT form

but exhibits a reduced ThT fluorescence intensity, reflecting an alteration in the fibril structure. This

result indicates that in RIPK3, interactions outside of the core four residues are also important in the

formation of homomeric amyloid structures.

The M45 RHIM forms the core structure within the amyloid fibrils

Amyloid assembly by the RHIM-containing fragments was also characterised under native

conditions, when amyloid assembly occurred spontaneously during recombinant expression. When

$M45_{1-90}$, $RIPK1_{497-583}$ and $RIPK3_{387-518}$ were expressed as fusions with MBP, the fusion proteins

remained soluble and could be purified under native conditions. Size exclusion chromatography

after affinity purification separated the material into two main fractions (Fig. 2c). The first of these

eluted in the void volume, indicating the formation of structures with MW >100 000 Da. SDS

PAGE showed that the peak eluting in the void volume contained MBP-RHIM fusion protein while

the second peak contained MBP only. The high molecular weight material was examined by

negative stain transmission electron microscopy (TEM) and for MBP-$M45_{1-90}$, MBP-$RIPK1_{497-583}$

and MBP-$RIPK3_{387-518}$, fibrillar structures of up to ~200 nm in length were observed (Fig. 2d).

Congo red solution binding assays performed with this material showed increased absorbance at

540 nm for all three samples and a shift in the absorbance maxima, relative to a non-amyloid

control (soluble insulin), indicating that the fibrils had an amyloidogenic nature (Fig. 2e).

To probe the arrangement of the RHIM within the fibrils, we treated the fibrillar material with

thrombin to cleave specifically between MBP and the RHIM-containing regions. During incubation

with the protease, insoluble material became visible in all three samples. Separation of this material

from the soluble fraction by centrifugation, followed by SDS PAGE and SEC analysis of the

soluble material, revealed that M45$_{1-90}$, RIPK1$_{497-583}$ and RIPK3$_{387-518}$ had formed insoluble

precipitates while the MBP remained soluble. The insoluble material was resuspended and

examined by TEM and fibrils were observed that were longer and thinner than those formed by the

MBP fusion proteins (Fig. 2f). These results indicate that the RHIM-containing region of M45 is

amyloidogenic and forms the β-sheet core structure of the fibrils, in a similar fashion to that

demonstrated for RIPK1 and RIPK3 by Wu and coworkers[29]. The MBP fusion partner is arrayed on

the outside of the β-sheet core and limits the growth of the fibrils to ~200 nm; in its absence, the

RHIM-containing segments assemble into longer fibrils.

M45 interacts with RIPK1 and RIPK3 to form heteromeric assemblies

To address the interactions between the RHIM-containing portions of M45, RIPK1 and RIPK3, we

applied fluorescence polarisation and co-localisation experiments. When His-Ub-RIPK1$_{497-583}$ was

covalently labelled with Alexa Fluor488 and mixed with unlabelled His-Ub-RIPK1$_{497-583}$, His-Ub-

RIPK3$_{387-518}$ or His-Ub-M45$_{1-90}$ in a ratio of 1 labeled:1999 unlabelled, the Alexa488-labelled

RIPK1$_{497-583}$ was incorporated into growing His-Ub-RIPK1$_{497-583}$, His-Ub-RIPK3$_{387-518}$ or His-Ub-

M45$_{1-90}$ fibrils (Fig. 3a). The rates of incorporation of the RIPK1-RHIM construct into His-Ub-

RIPK1$_{497-583}$ and His-Ub-RIPK3$_{387-518}$ fibrils were similar and RIPK1-RHIM was incorporated into

His-Ub-M45$_{1-90}$ fibrils more slowly. The tetrad sequence in the RHIM of M45 is IQIG and identical

to that in RIPK1, whereas the RIPK3 tetrad sequence is VQVG. The observed difference in

incorporation rates indicates that residues outside the tetrad also influence the interactions between

RHIM-containing proteins (Fig. 1a). Taken together with the observed different effects of the

AAAA mutations on M45, RIPK1 and RIPK3 homomeric amyloid formation, we conclude that the

contributions of the tetrad sequences to homomeric and heteromeric structures are distinct and

differ between RHIM-containing proteins.

Fluorescence co-localisation experiments were applied to detect RHIM-based hetero-oligomers against a background of monomeric proteins and the potential competing formation of homo-oligomers[36,37]. In these experiments, multiple different RHIM-containing fusion proteins were mixed together and maintained in monomeric form in 8 M urea-containing buffer. Assembly or co-assembly were initiated by reduction of the urea concentration through dilution. Mixtures of YPet- and mCHERRY-tagged proteins were examined in solution, using a confocal microscope where the lasers create a very small confocal volume (~250x250x800 nm). Freely diffusing fluorophore-tagged proteins are only excited and detected when they are within the confocal volume (Fig. 3b). The number of molecules in the focal volume/millisecond over the range of concentrations used was ~10-500.

In initial fluorescence spectroscopy experiments, we analysed mixtures of YPet- and mCHERRY-tagged RIPK1$_{497-583}$ or YPET- and mCHERRY-tagged RIPK3$_{387-518}$. We observed coincidence of the signals from both fluorophores when oligomers formed, demonstrating that homomeric fibril assembly can be detected in this way (Fig. 3c,d). Mixtures of mCHERRY-RIPK1$_{497-583}$ and YPet-RIPK3$_{387-518}$ exhibited formation of heteromeric fibrils, indicated by coincidence of the signals from the two fluorophores attached to different RHIM-containing constructs (Fig. 3e). In addition, FRET could be observed between YPet and mCHERRY within the fibrils (Fig. 3f). These results provide the first direct single-molecule-level evidence for the formation of heteromeric fibrils containing RIPK1 and RIPK3 RHIM sequences, supporting the results from bulk solution previously reported[29]. The RIPK3$_{387-518}$ construct demonstrated a greater propensity to self-assemble than the RIPK1$_{497-583}$ construct, as evidenced by the formation of larger assemblies (Fig. 3c,d). In addition, all oligomeric RIPK1$_{497-583}$ appeared coincident with RIPK3$_{387-518}$ material, while some RIPK3$_{387-518}$-only fibrils were observed (Fig. 3e).

The self-assembly of $M45_{1-90WT}$ into fibrils could be detected by fluorescence co-localisation but

substitution of the IQIG tetrad with AAAA abolished the ability of this M45 fragment to self-

assemble (Fig. 4a,b). This was confirmed by electron microscopy, which detected the presence of

long fibrils formed by $M45_{1-90WT}$-mCHERRY and only amorphous clusters in the $M45_{1-90AAAA}$-

mCHERRY sample (Supplementary Fig. 1). When $M45_{1-90WT}$-mCHERRY and YPet-RIPK1$_{497-583}$

or YPet-RIPK3$_{387-518}$ were mixed under fibril assembly conditions, we observed coincidence of the

$M45_{1-90WT}$ with both RIPK fragments, demonstrating heteromeric complex formation between viral

and host RHIM-containing fragments (Fig. 4c,d). The complexes containing $M45_{1-90}$ and

RIPK3$_{387-518}$ appeared larger than those containing $M45_{1-90}$ and RIPK1$_{387-518}$, as judged by the

length of time taken by the heteromeric complexes to diffuse through the focal volume. The AAAA

mutation of the core tetrad of M45 abolished co-assembly with RIPK1 and RIPK3 (Fig. 4e,f).

We added non-fluorescent His-Ub-$M45_{1-90}$ into mixtures of YPet- and mCHERRY-tagged RIPK1

and RIPK3 fragments, to probe the effect of the amyloidogenic viral RHIM on heteromeric

complex formation by the two human RHIMs. The number of times a large complex generating

>5000 or >10000 photons above baseline intensity was detected in the focal volume for 1

millisecond during the last 200 seconds of a 10–minute incubation was quantified for a

RIPK1:RIPK3 mixture and a series of RIPK1:RIPK3:M45 mixtures. The addition of His-Ub-

$M45_{1-90}$ resulted in a decrease in the number of the large complexes detected, correlated with the

relative concentration of His-Ub-$M45_{1-90}$ (Fig. 4g). EM of samples formed with increasing amounts

of $M45_{1-90}$ relative to RIPK3$_{387-518}$ (1:1; 2:1 and 4:1) confirmed an increase in the size of the

fibrillar assemblies (Supplementary Fig. 2). The reduction in the number of large complexes

detected is therefore likely due to a reduction in the rate at which larger complexes diffuse in the

well, which results in fewer passes through the focal volume and hence detection. Collectively,

these results show that $M45_{1-90}$ is able to interact with RIPK1$_{497-583}$ and RIPK3$_{387-518}$ in a RHIM-

dependent manner but displays a preference for co-assembly with the RHIM of RIPK3.

The M45-RHIM alters the formation of fibrils by the RHIMs of RIPK1 and RIPK3

Self-assembly of single types of RHIM-containing proteins resulted in predominantly unbranched

single fibrils, as visualised by TEM (Fig. 5a). In contrast, when the $M45_{1-90}$ fragment was mixed

with either of the $RIPK1_{497-583}$ or $RIPK3_{387-518}$ fragments in a 1:1 molar ratio under conditions

allowing fibril assembly, extensive fibrillar networks were observed (Fig. 5a). These consisted of

large bundles of multiple fibrils that branched and reformed in different combinations, resulting in

an interweaved and dense assembly covering large areas of the TEM grids. By contrast, when

$RIPK1_{497-583}$ and His-Ub-$RIPK3_{387-518}$ were mixed together under conditions allowing fibril

assembly, isolated mostly single fibrils of <200 nm length were observed, similar to those reported

by Li et al (2012). When $M45_{1-90}$ or $RIPK1_{497-583}$ were allowed to assemble in the presence of an

unrelated protein with the same fusion partner, His_6-Ub-RodA, no thick multi-fibril bundles were

observed, suggesting that this is a feature associated with hetero-fibrils formed by two RHIM-

containing proteins (Supplementary Fig. 3).

The different RHIM-containing components cannot be distinguished in these TEM images so we

used confocal microscopy to visualise fibrils formed from mixtures of RHIM-containing fluorescent

fusion proteins (Fig. 5b–e). We observed the formation of long fibrils that contained $M45_{1-90}$ and

either one or both of human $RIPK1_{497-583}$ or $RIPK3_{387-518}$. These assemblies are thioflavin T

positive, confirming that these large heteromeric assemblies do have a cross-β substructure

(Supplementary Fig. 4). Fibrils prepared from a mixture containing mCHERRY-$M45_{1-90}$ and YPet-

$RIPK3_{387-518}$ were additionally imaged by super-resolution microscopy and transmission electron

microscopy to provide higher resolution images. We found that the viral and host RHIMs were

distributed throughout the fibrillar protein network formed by these proteins (Fig. 5f,g).

The incorporation of M45 RHIM alters fibril stability

We reasoned that the incorporation of the viral M45 RHIM into host heteromeric fibrils might alter

the properties of the fibrils, in addition to changing the morphology of the aggregates. Other studies

of functional amyloid fibrils have noted that they are less resistant to treatment with the detergent

sodium dodecyl sulphate (SDS) than some disease-associated amyloid fibrils[38,39]. We prepared

homomeric and heteromeric assemblies from $M45_{1-90WT}$ and $M45_{1-90AAAA}$ with $RIPK3_{387-518}$. These

assemblies were incubated with and without SDS, and then the preparations were analysed by

agarose gel electrophoresis (AGE) (Fig. 6). This separated the material into (1) very large SDS-

resistant fibrillar material that remained in the wells, (2) ThT-positive soluble amyloid oligomers

and (3) monomeric protein. The $M45_{1-90WT}$ and $RIPK3_{387-518}$ mixtures contained soluble amyloid

oligomers. The incorporation of the viral RHIM destabilised the complexes such that treatment with

2% SDS resulted in release of both constructs in monomeric form and, in the 1:1 mixtures, the

generation of RIPK3-only fibrils. RIPK3-only fibrils were predominantly SDS-resistant and

remained as large fibrils trapped in the wells (Fig. 6a). When the viral M45 RHIM was present in

four-fold molar excess, little RIPK3-only fibrillar material remained in the wells. These results were

supported by analysis of the soluble and insoluble material generated by dialysing mixtures of

RHIM-containing constructs together, which showed that the presence of the $M45_{1-90WT}$ fragment

reduced the amount of $RIPK3_{387-518}$ that was detected in the insoluble fraction (Supplementary Fig.

5). This activity of the M45 RHIM was dependent on the presence of the WT tetrad core sequence,

as in the presence of $M45_{1-90AAAA}$, a large amount of SDS-resistant RIPK3-only material remained

in the wells and very low levels of monomeric $RIPK3_{387-518}$ were detected, even with four times as

much $M45_{1-90AAAA}$ (Fig. 6a). The $M45_{1-277}$ construct displayed a similar activity (Fig. 6b). It self-

assembled into soluble homo-oligomers. It co-migrated with $RIPK1_{497-583}$ but interacted strongly

with $RIPK3_{387-518}$ as judged by the narrow, intense orange-coloured co-localised band

corresponding to soluble oligomers. In the absence of the intact RHIM core, $M45_{1-277AAAA}$ formed

large soluble oligomers with a wide mobility range but was unable to interact with $RIPK1_{497-583}$ or

$RIPK3_{387-518}$. Independently migrating oligomers of $RIPK1_{497-583}$ and large RIPK3-only homo-

oligomeric fibrils were observed (Fig. 6b). These results indicate that the M45 RHIM can compete

with, and prevent, homo-oligomerisation of RIPK3-containing oligomers, such as those found in

necrosomes.

Discussion

These results indicate that the RHIM within the MCMV M45 protein is amyloidogenic and the ability of M45 to self-assemble into homomeric fibrils or to form heteromeric amyloid-structured complexes with RIPK1 and RIPK3 depends on the presence of the core tetrad sequence IQIG within the RHIM. The N-terminal 90 residues of M45 are sufficient to support interactions with the RHIMs of human RIPK1 and RIPK3 and to protect against TNFR-mediated necroptosis in human cells *in vitro*. When probed with necroptosis-relevant RHIM-containing protein fragments, the viral RHIM preferentially interacts with the RIPK3 RHIM over the RIPK1 RHIM. In the presence of the M45 RHIM, a RHIM-encompassing region of RIPK3 is incorporated into metastable heteromeric fibrils rather than into SDS-resistant homo-oligomers. The RIPK1:RIPK3 interaction and formation of the necrosome is central to TNFR-mediated responses if caspase-8 is inhibited by viral or chemical inhibitors[3]. We propose that the presence of M45 in cells results in the trapping of RIPK1 and RIPK3 within alternative heteromeric "decoy" amyloid structures and prevents the native RIPK1:RIPK3 RHIM-based amyloid interactions that are required to trigger cell death.

An important difference between RHIM-based amyloids and other disease-associated or most functional amyloids is the formation of heteromeric assemblies[30]. Heteromeric RHIM-based amyloid is central to necroptosis induction[11]. RNA-binding proteins containing low complexity sequences have the capacity to co-assemble into phase-separated droplets and hetero-fibrils, driven by intermolecular interactions between the prion-like domains, however the *in vivo* significance of heteromeric fibril assembly by these proteins is currently less clear than is the case for RHIM-based amyloid[38,39]. The multi-protein RHIM amyloid structures are similar to other higher-order assemblies such as signalosomes, that form by polymerization and recruit multiple proteins with different activities[40].

The key features of the heteromeric RIPK1:RIPK3 necrosome core have now been revealed by

ssNMR[32]. The RHIMs of RIPK1 and RIPK3 adopt a serpentine fold and form parallel β-sheets

through intermolecular hydrogen binding. Two such sheets assemble to form a fibril through inter-

tetrad and other interactions, which generate an oblong-shaped hydrophobic core (Fig. 6c). The

structure is also stabilised by solvent-exposed side-chain-side-chain hydrogen bonds between

stacked Asn and Gln residues, and Tyr and Ser/Thr side chains that form stacked ladder-like

interactions on the periphery of the core.

We have used this structure to model the incorporation of the RHIM of M45. Since M45 and

RIPK1 have the same core IQIG tetrad sequence, and we have determined that M45 preferentially

forms heteromeric fibrils with RIPK3 over RIPK1, we have mutated the RIPK1 sequence to that of

M45 and residue substitutions were made without significant steric clashes (Fig. 6d). The core

tetrad interactions therefore are unchanged from those in the RIPK1:RIPK3 structure but with the

viral protein, two Tyr and one Asn ladder are not possible. Instead, a new Asp-Asn side-chain

interaction is likely and in the M45:RIPK3 structure, a Val-Cys ladder takes the place of the Ser-

Cys ladder observed with RIPK1:RIPK3. Our model suggests that the M45:RIPK3 RHIM core will

be similar to that formed by RIPK1:RIPK3 and that M45 mimics the host inter-RHIM interactions.

It is in line with our experimental observation that the AAAA mutation to M45 abolishes the ability

of this protein to form a hetero-amyloid with the RHIM of RIPK3. Our experimental data strongly

indicate that the interactions driving M45:RIPK3 heteromeric assembly are stronger than those

driving M45:RIPK1 or RIPK1:RIPK3 assembly. The ssNMR-derived structure does not reveal

potential heteromeric interactions between polypeptides that occupy the corresponding "rung" of

opposing sheets. It is possible that the viral protein could be involved in such interactions with host

partners. Interference by the viral protein with inter-sheet interactions would explain the branching

and network morphology we observe in our heteromeric assemblies (Fig. 5a,f; Supplementary Fig.

2). The concentration of viral protein in the cell may also influence the make-up of the RHIM-based

structures within cells, and hence control the final cellular outcome. The structure determined by

Mompean *et al.* is also consistent with confocal images we have collected of isolated RHIM fibrils,

which show fluorescent proteins along the two flanks of the fibril and a dark central core. This

organisation of fluorescent partner proteins is only consistent with an opposing arrangement of two

parallel β-sheets (Fig. 6e,f).

It is noteworthy that in our experiments the AAAA mutation of the core tetrad in $RIPK3_{387-518}$ does

not abolish self-assembly into an amyloid structure while this mutation in $RIPK1_{497-583}$ and $M45_{1-90}$

prevents self-assembly. This indicates that residues outside the tetrad of RIPK3 can support

oligomerisation but these remain to be identified by further mutagenesis and structural studies of the

entire RIPK3 RHIM. Although calculations based on the RIPK1:RIPK3 core structure suggest that

RIPK3:RIPK3 homo-amyloid formation is less favourable than RIPK1:RIPK3 or RIPK1:RIPK1[32],

the formation of the RIPK1:RIPK3 necrosome is known to nucleate further polymerisation of

RIPK3 that leads to activation of RIPK3 and interaction with MLKL[41]. It will be important to

determine whether the entrapment of RIPK3 by M45 in alternative assemblies prevents RIPK3

auto-phosphorylation and/or if it prevents the subsequent recruitment and activation of MLKL

which leads to MLKL oligomerisation and membrane lysis[42]. The results of other studies suggest

that the former mechanism may operate. M45 prevents ZBP1-induced RIPK3 phosphorylation[7] and

autophosphorylation of RIPK3 when co-expressed in 293Ts[43] (Huang 2015 CHM). HSV-1 and -2

encode RHIM-containing proteins, ICP-6 and ICP10, which have been shown to protect against

necroptosis in human cells [25,44] and ICP10 expression is known to prevent phosphorylation of

RIPK1 and RIPK3 compared to control[25]. Notably, in our results, the effect of the M45 RHIM is to

sequester RIPK1 and RIPK3 RHIM-containing fragments in amyloid-based assemblies that are

persistent but not as stable as RIPK1:RIPK3 fibrils. Any decoy amyloids involving M45 and host

RHIM-containing proteins should not trigger other host defense or cytotoxic mechanisms that

would be undesirable for continued viral infection of the cell.

An alternative to the viral RHIM sequestration studied here has been adopted by some bacterial

pathogens. Enteropathogenic *E. coli* (EPEC) is a gut pathogen that has been shown to encode a

protease, EspL, that is delivered through the type III secretion effector system and which rapidly

cleaves within the RHIM of RIPK1, RIPK3, ZBP-1/DAI and TRIF in non-fibrillar forms of these

proteins[45]. Amyloid fibril formation by endogenous RHIM-containing proteins protects them from

cleavage by EspL. Thus EspL appears to act by disabling the amyloid-based RHIM host defense

pathways that would otherwise curtail infection[45].

Several different RHIM:RHIM interactions also occur in the absence of infection. An intact RHIM

in RIPK1 suppresses undesirable ZBP-1:RIPK3 interactions during development that would lead to

necroptosis[46,47] and ZBP-1 mediates RIPK3-dependent necroptosis in HT29 cells[48]. The methods

we have developed to probe and compare competing RHIM interactions can be applied to determine

the full extent of the interface involved in heteromeric interactions and to test the relative strengths

of different RHIM:RHIM interactions. This will lead to elucidation of the basis for cellular control

of these potentially competing RHIM:RHIM interactions. Necroptosis is increasingly recognized in

pathophysiological processes including ischaemia-reperfusion injury, and inflammation of the skin

and intestinal epithelium[13,49]. Any development of therapies that target the necroptosis pathway will

benefit from a molecular-level understanding of the structure and regulation of the multiple RHIM-

based interactions that are important in human health and disease.

Acknowledgements

The work was supported by funds provided by Australian Research Council Discovery Project Grants DP150104227 and DP180101275 to M. Sunde. M. Steain was supported by National Health and Medical Research Council funding to Prof. Allison Abendroth and The University of Sydney Bridging Grant scheme. The authors acknowledge the facilities and the scientific and technical assistance of Sydney Microscopy & Microanalysis at the Australian Centre for Microscopy & Microanalysis at the University of Sydney. We thank Dr Louise Cole of the Bosch Institute Advanced Microscopy Facility for assistance with collection of confocal images, staff from the Bosch Institute Molecular Biology Facility for use of equipment, Charles Collyer for use of the X-ray generator, Sue McLennan for providing the HT-29 adenocarcinoma cells and James Murphy for providing the *E.coli* DNA gyrase subunit B. We thank Nicholas Della Marta, Monique Fischer, Laura Lau and Jake Campbell for preliminary experiments with RHIM peptides and cloning of fluorescent proteins. We thank Pascal Krotee and David Eisenberg for helpful discussions about the use of fluorescent proteins for studying RHIM interactions and models of heteromeric and homomeric amyloids, respectively.

Figure Legends

Figure 1. The M45 RHIM is amyloidogenic and 1-90 and 1-277 regions of M45 confer protection against TNF-induced necroptosis in human cells. (**a**) Schematic representation of M45, RIPK1 and RIPK3 proteins. Grey regions indicate RHIM-encompassing fragments used in this study. Black bar represents 19-residue RHIM. RNR-like represents inactive ribonucleotide reductase domain, DD represents the RIPK1 death domain. Amino acid sequences of the RHIMs of M45 and human and mouse RIPK1 and RIPK3, boxed region indicates the core tetrad that is substituted with AAAA in mutant protein constructs. (**b**) Negatively-stained transmission electron micrograph of fibrils formed by a synthetic peptide encompassing the M45 RHIM. Three samples were examined by microscopy. (**c**) HT29 cell survival 18h after treatment with TNF+BV-6+z-VAD-fmk to induce necroptosis. Cells expressed lentivirus constructs encoding 1-90 or 1-277 fragments of M45 and were cultured +/- the addition of coumermycin A1 to induce dimerisation. N=2 for M45-90 gyrase and 3 for M45-277 gyrase. Error bars represent SEM, **** = $p < 0.0001$.

Figure 2. The RHIM-containing fragment forms the amyloid fibril core. (**a**) X-ray fibre diffraction pattern collected from fibrils formed by His-Ub-M45$_{1-90}$. Two different samples were examined by XRFD. (**b**) Thioflavin T fluorescence as a function of time, following dilution of WT or AAAA His-Ub-M45$_{1-90}$, mCHERRY-RIPK1$_{497-583}$, and YPet-RIPK3$_{387-518}$ from 8 M-containing buffer to 100–300 mM urea. Buffer sample contains ThT but no protein. Samples were tested in triplicate and solid line is the average of the triplicates and the error bars (in grey) represent +/- SEM. (**c**) Size exclusion chromatography (SEC) of MBP-M45$_{1-90}$, MBP-RIPK1$_{497-583}$, and MBP-RIPK3$_{387-518}$ following amylose affinity purification. Grey oval indicates high MW material eluting in void volume. (**d**) Negatively-stained transmission electron micrograph of material eluted in the void volume. These are representative micrographs of the material observed. All fusion proteins expressed and purified at least twice. (**e**) Absorbance spectrum from solutions containing Congo red with soluble monomeric insulin, insulin amyloid fibrils or material eluted in the void volume containing MBP-M45$_{1-90}$ (green), MBP-RIPK1$_{497-583}$ (cyan), or MBP-RIPK3$_{387-518}$ (blue). Experiment conducted once. (**f**) Representative images of insoluble material generated after treatment with thrombin removed the MBP. Experiment repeated three times.

Figure 3. Assembly of RHIMs gives rise to homomeric and heteromeric structures. (**a**) Fluorescence polarization profiles indicating the rate of incorporation of Alexa488-tagged RIPK1$_{497-583}$ into unlabeled M45$_{1-90}$ (green), RIPK1$_{497-583}$ (cyan), or RIPK3$_{387-518}$ (blue), with labeled:unlabeled protein at 1:1999. The average polarization profile from triplicate readings is shown, with s.d. indicated. (**b**) Schematic representation of fluorescence spectroscopy experiments used to detect the formation of homo- and heteromeric amyloid fibrils. Fusion proteins containing RHIMs and different fluorescent partners are mixed together under assembly-permissive conditions and examined in the confocal volume. Co-assembly of two different fluorophores within one complex results in detection of coincident signals in the two channels. (**c**) Representative fluorescence intensity trace showing homomeric assembly over 40 s in a mixture of mCHERRY-RIPK1$_{497-583}$ and YPet-RIPK1$_{497-583}$. (**d**) Representative fluorescence intensity trace showing homomeric assembly over 40 s in a mixture of mCHERRY-RIPK3$_{387-518}$ and YPet-RIPK3$_{387-518}$. (**e**) Representative fluorescence intensity trace showing heteromeric assembly over 40 s in a mixture of mCHERRY-RIPK1$_{497-583}$ and YPet-RIPK3$_{387-518}$. Inset is expansion of region indicated by * on main trace. (**f**) Förster resonance energy transfer (FRET) signal between YPet- and mCHERRY fluorophores, detected with excitation by 488 nm laser only. All mixtures examined in at least three different experiments on three separate occasions. In panels c–f the signal from the mCHERRY-tagged protein is coloured magenta and the signal from the YPet-tagged protein is coloured green.

Figure 4. Heteromeric amyloid formation including M45 RHIM requires intact core tetrad.
Representative fluorescence intensity traces collected over 50 s for mixtures of (**a**) mCHERRY-M45$_{1-90WT}$ and YPet-only; (**b**) mCHERRY-M45$_{1-90AAAA}$ and YPet-only; (**c**) mCHERRY-M45$_{1-90WT}$ and YPet-RIPK1$_{497-583}$; (**d**) mCHERRY-M45$_{1-90WT}$ and YPet-RIPK3$_{387-518}$; (**e**) mCHERRY-M45$_{1-90AAAA}$ and YPet-RIPK1$_{497-583}$ and (**f**) mCHERRY-M45$_{1-90AAAA}$ and YPet-RIPK3$_{387-518.}$ (**g**) Number of bins where intensity (photons/ms) exceeded 5000 or 10000 in the last 200 s of a 10–min incubation, where samples contained no His-Ub-M45$_{1-90}$ or up to a 4-fold molar excess of His-Ub-M45$_{1-90}$. Coincidence experiments conducted three times or more on different days. Quantification performed once.

Figure 5. Co-assembly of M45$_{1-90WT}$ with RIPK1$_{497-583}$ and His-Ub-RIPK3$_{387-518}$ results in the formation of heteromeric fibrils and dense fibrillar networks. (**a**) Representative electron micrographs of fibrils formed by M45$_{1-90}$, RIPK1$_{497-583}$ or His-Ub-RIPK3$_{387-518}$ when dialysed individually or in mixtures, into phosphate buffer to remove urea. Confocal fluorescence images of heteromeric fibrils formed by co-assembly of (**b**) mCHERRY-M45$_{1-90}$ and YPet-RIPK1$_{497-583}$; (**c**) mCHERRY-M45$_{1-90}$ and YPet-RIPK3$_{387-518}$; (**d**) mCHERRY-RIPK1$_{497-583}$ and YPet-RIPK3$_{387-518}$; (**e**) ECFP-M45$_{1-90}$ with mCHERRY- RIPK1$_{497-583}$ and YPet-RIPK3$_{387-518}$. (**f**) STED super-resolution and (**g**) electron microscope images of heteromeric fibrils formed by co-assembly of mCHERRY-M45$_{1-90}$ and YPet-RIPK3$_{387-518}$.

Figure 6. SDS AGE analysis of RHIM-based assemblies and model of the heteromeric M45:RIPK3 core. (**a**) SDS AGE analysis of YPet-RIPK3$_{387-518}$ and mCHERRY-M45$_{1-90WT}$ or mCHERRY-M45$_{1-90AAAA}$ homomeric and heteromeric assemblies, formed by initial mixing in 8 M urea, followed by removal of urea by overnight dialysis. Dialysed samples were incubated with 0 or 2% SDS at RT for 10 min before electrophoresis. Monomeric forms of the protein constructs were maintained in 8 M urea prior to electrophoresis, as indicated. Protein components as indicated above each lane. (**b**) SDS AGE analysis of homomeric and heteromeric assemblies of YPet-RIPK1$_{497-583}$, YPet-RIPK3$_{387-518}$ and mCHERRY-M45$_{1-277WT}$ or mCHERRY-M45$_{1-277AAAA}$. BSA was included in M45-only samples to maintain the overall protein concentration constant. (**c**) Structure of the RIPK1:RIPK3 RHIM core. Figure prepared using PyMol Graphics System from PDB 5V7Z. The RIPK3 sequence is coloured in blue and RIPK1 is coloured in green. Tetrad residue side chains are shown as sticks; Tyr, Asn/Gln, Ser and Cys side chains are shown in wheat, red, grey and yellow, respectively. Core tetrad interactions across the opposing sheets are indicated by the dotted box. (**d**) Model of M45:RIPK3 RHIM core, prepared by appropriate mutation of RIPK1. The M45 sequence is coloured in cyan and RIPK3 in blue. Tetrad residue side chains are shown as sticks; Tyr, Asn/Gln, Val and Cys side chains are shown in wheat, red, grey and yellow, respectively. Core tetrad interactions across the opposing sheets are indicated by the dotted box. (**e**) Confocal microscopy image of a single fibril composed of YPet-M45. (**f**) Schematic representation showing associated globular domains (e.g. kinase or RNR, coloured green) flanking either side of the amyloid core formed by RHIMs (coloured grey).

Legends to Supplementary Figures

Supplementary Fig 1. The AAAA substitution of the core tetrad within the RHIM of M45$_{1-90}$ abolishes its ability to self-assemble into fibrils. Negatively stained EM of fibrils formed by (**a**) M45$_{1-90WT}$-mCHERRY and (**b**) M45$_{1-90AAAA}$-mCHERRY fibrils upon removal of 8 M urea by dialysis.

Supplementary Fig 2. An increase in the amount of M45$_{1-90}$ relative to RIPK3$_{387-518}$ results in the formation of increasingly large fibrillar assemblies. Negatively stained EMs of fibrillar assemblies formed from mixtures of M45$_{1-90WT}$-mCHERRY and YPetRIPK3$_{387-518}$ at (**a**) 1:1, (**b**) 2:1 and (**c**) 4:1 relative molar concentration. Monomeric forms of the proteins were mixed together and allowed to co-assemble upon removal of 8 M urea by dialysis.

Supplementary Fig 3. The presence of an unrelated His$_6$-ubiquitin fusion protein does not result in generation of fibrillar networks. Samples of either (a) His-Ub-M45$_{1-90}$ or (b) His-Ub-RIPK1$_{497-583}$ RHIM fusion proteins were mixed with the non-RHIM-containing protein His-Ub-RodA in urea-containing buffer and allowed to assemble upon removal of urea by dialysis.

Supplementary Fig 4. Heteromeric assemblies formed by RHIM-containing fragments are thioflavin T-positive, indicative of an amyloid structure. mCHERRY-M45$_{1-90}$ and YPet-RIPK3$_{387-518}$ were allowed to coassemble during dialysis, following which thioflavin T was added and the dialysate imaged using a Cytation 3 Imager (BioTek). CFP, YFP and Texas Red filter sets were used to detect Thioflavin T fluorescence, YPet and mCHERRY respectively.

Supplementary Fig 5. SDS-PAGE analysis of RHIM-containing assemblies resulting from overnight dialysis of individual constructs (a–c) or protein pairs (d–f). RHIM-containing fusion proteins dialysed overnight to allow fibril assembly, either individually or in pairs after initial mixing in urea in the monomeric form. All RHIM constructs carried mCHERRY as fusion protein partner. RHIM-containing individual construct or pairs as indicated in each panel. The stability of the resulting material towards SDS treatment was tested. Samples are total dialysate (total), supernatant from centrifugation at 16000g for 10 min (soluble), pellet from centrifugation resuspended in 20 mM NaH$_2$PO$_4$, 150 mM NaCl, 0.5 mM DTT, pH 7.4 (insoluble), pellet from centrifugation treated with 2% SDS for 10 min at 37 °C and then separated again into supernatant (SDS-soluble) and pellet (SDS-insoluble). SDS-insoluble material was incubated in 8 M urea to solubilise material for detection by PAGE. This SDS analysis was performed once.

Methods

Expression of RHIM fusion proteins

RHIM-containing portions of mCMV M45 (Q06A28; residues 1-90), human RIPK1 (Q13546;

residues 497-583), human RIPK3 (Q9Y572; residues 387-518), were produced as maltose binding

protein (MBP) fusion proteins using the pMAL vector, as His_6-Ubiquitin fusion proteins using the

pHUE vector system (Catanzariti et al) or as His-tagged N- or C-terminal ECFP, YPet or

mCHERRY fusion proteins using a vector prepared in-house. Successful cloning was confirmed by

sequencing at the Australian Genome Research Facility at the Westmead Millennium Institute.

Proteins were expressed in BL21(DE3) *E. coli* grown in LB media containing ampicillin at 37 °C

for 2–3 hours, with induction by 0.5–1.0 mM IPTG when the $OD_{600\,nm}$ of the culture reached

0.6–0.8. All pMAL plasmids were grown in LB media containing 0.2% glucose to prevent amylase

production. SDS PAGE analysis of cell lysates following induction of protein over-expression

indicated that the MBP-RHIM fusion proteins were present in the soluble cell fraction while the

His_6-Ub-RHIM fusion proteins and His_6-ECFP, His_6-YPet or His_6-mCHERRY fusion proteins were

located in the insoluble fraction.

Purification of MBP fusion proteins

Cells were lysed by incubation in 20 mM Tris.HCl, 200 mM NaCl, 1 mM EDTA and 0.5 mM DTT,

pH 8.0 with lysozyme at 100 µg/mL, followed by a freeze thaw cycle. DNaseI and $MgSO_4$ were

added to final concentrations of 100 µg/mL and 100 mM respectively, and samples were incubated

with stirring for 15 min at room temperature. Following the addition of AEBSF to 0.2 mM, samples

were centrifuged at 4 °C at 10,000*g* for 40 min, to separate soluble material from insoluble cell

debris. MBP fusion proteins were purified from the soluble fraction using amylose beads from New

England Biolabs (MA, USA), as per the manufacturer's instructions.

Purification of His_6-Ubiquitin fusion proteins

Cell pellets were lysed by suspension and incubation in 6 M GuHCl, 100 mM NaH_2PO_4, 20 mM

Tris.HCl, 5 mM β-mercaptoethanol, pH 8.0, and soluble material further purified on Ni-NTA

agarose (Life Technologies) under denaturing conditions, with exchange into 8 M urea, 100 mM

NaH_2PO_4, 20 mM Tris.HCl, 5 mM β-mercaptoethanol at pH 6.0 for washing and pH 4.0 for elution

from the Ni-NTA agarose.

Purification of His$_6$-ECFP, His$_6$-YPet and His$_6$-mCHERRY fusion proteins

Two methods for purification of polyhistidine-tagged fluorescent protein-RHIM fragment fusions

were used during the course of this study. The first method was batch purification where cell

pellets were lysed by suspension and incubation in 8 M urea, 100 mM NaH_2PO_4, 20 mM Tris.HCl,

5 mM β-mercaptoethanol, pH 8.0. Soluble material was purified using Ni-NTA agarose beads (Life

Technologies) under denaturing conditions, with 8 M urea, 100 mM NaH_2PO_4, 20 mM Tris.HCl,

5 mM β-mercaptoethanol at pH 8.0 and in the presence of 20mM imidazole for washing steps and

300mM imidazole for elution from the Ni-NTA agarose. The second method involved using a His-

trap (GE Healthcare) column. For this, cell pellets were lysed as described for the purification of

MBP fusion proteins. The insoluble pellet was resuspended in 8 M urea, 100 mM NaH_2PO_4,

20 mM Tris, pH 8.0 and allow to dissolve by stirring for 2 h. The sample was centrifuged at 4 °C at

31,000g for 45 min, filtered and then purified on a 5mL His-Trap column (GE Healthcare) under

denaturing conditions (8 M urea, 100 mM NaH_2PO_4, 20 mM Tris, pH 8.0 + 0.5mM DTT) using

AKTA FPLC. The elution of protein was achieved with a gradient of imidazole. Protein elution was

monitored by absorbance at 280 nm.

For both purification procedures, the eluted fractions were analysed by SDS-PAGE. Fractions

containing the fusion proteins were pooled and concentrated using Amicon Ultra-15 Centrifugal

Filter Units with MWCO 30kDa (Millipore). The concentration of the final concentrated protein

sample was determined by BCA assay (Pierce).

Removal of fusion tag (MBP and His$_6$-Ubiquitin)

MBP-RHIM fusion proteins at concentrations of 1–4 mg/mL in 20 mM Tris.HCl, 200 mM NaCl,

1 mM EDTA, 2.5 mM $CaCl_2$, 10 mM maltose pH 8.0 were incubated with 5 U Thrombin for 2

hours at 37 °C. Samples were centrifuged at 16 000g for 10 min to separate insoluble and soluble

material and both fractions were analysed by SDS PAGE. His$_6$-Ub-RHIM constructs in 8 M urea-

containing buffer were dialysed overnight at room temperature, against 20 mM NaH$_2$PO$_4$, 50 mM

NaCl pH 7.4 and then incubated with deubiquitylating enzyme UBP41 (produced in-house

according to Catanzariti et al) at 37°C for 3 hours. The RHIM-containing fragments self-assembled

to form insoluble material during dialysis and incubation with UBP41. Samples were centrifuged to

separate the supernatant, containing the cleaved His$_6$-Ubiquitin, from the pellet. The pellet

containing the RHIM fragment was washed twice with 20 mM NaH$_2$PO$_4$, 50 mM NaCl pH 7.4,

followed by addition of 6 M GuHCl, 100 mM NaH$_2$PO$_4$, 20 mM Tris.HCl pH 8.0 to solubilise the

pellet and maintain the RHIM fragments in a monomeric form for storage.

Size exclusion chromatography

MBP-RHIM fusion proteins, before and after treatment with enzyme to remove the fusion partner,

were analysed by size exclusion chromatography using a Superdex 75 10/30 GL column (GE

Healthcare Life Sciences), running in 20 mM NaH$_2$PO$_4$, 150 mM NaCl pH 7.4 with a flow rate of

0.4 mL/min. Protein elution was monitored by absorbance at 280 nm and peaks were analysed by

SDS-PAGE and negative stain transmission electron microscopy.

Congo red assays

Insulin (1 mg/mL) was incubated in 20 mM glycine pH 2.0, with shaking at 700 rpm for 4 hours at

60 °C to convert it into amyloid fibril form. Samples of monomeric insulin, insulin fibrils or MBP-

RHIM fusion proteins, at a concentration of 100 μM, were prepared with 2 μM Congo red in

20 mM NaH$_2$PO$_4$, 150 mM NaCl pH 7.4 in a 1 mL cuvette. The absorbance of each protein-

containing solution was measured over the wavelength range 400 to 650 nm and compared to the

absorbance profile of a solution containing only Congo red in buffer.

M45 RHIM peptide

A synthetic peptide encompassing the RHIM of M45 was purchased from Genscript with the

sequence VRIMNGVSGIQIGNHNAMS. The peptide was solubilised in DMSO and then diluted

with water to a final peptide concentration of 300 μM with addition of a trace amount of ammonium

hydroxide to aid solubilisation. Addition of Thioflavin T indicated the development of a ThT-

positive structure. Samples were analysed by transmission electron microscopy.

Transmission Electron Microscopy (TEM)

A droplet of protein or peptide-containing solution (20 µL) was placed onto Parafilm™ and a

carbon/formvar-coated copper grid (200 mesh, ProSciTech) was floated on the droplet surface for 1

minute. Excess solution was removed by touching the edge of the grid to filter paper and the grid

was washed with filtered water three times and then stained by floating on a 20–µL droplet of 2%

uranyl acetate for 2 minutes, followed by removal of excess solution by blotting. The grid was air-

dried overnight at room temperature and then imaged using a Philips CM12 microscope operating at

120 kV. Digital images were recorded using a Morada 11 MegaPixel CCD camera camera and

iTEM digital imaging system.

ThT Kinetic Assays

His-Ub or fluorescent RHIM-containing fusion proteins in 8 M urea-containing buffer were diluted

into phosphate buffer (25 mM NaH_2PO_4, 150 mM NaCl, 0.5 mM DTT, pH 7.4) containing 40 µM

Thioflavin T to a final volume of 200 µL, protein concentration 10 µM His-Ub-M45, 2.5 µM

mCHERRY-$RIPK3_{387-518}$ or 2.5 µM YPet-$RIPK1_{497-583}$ and residual urea concentration of 0.2 M.

Three replicates of each protein sample were analysed simultaneously in a 96-well black

fluorescence plate sealed with optically clear film (Corning®). Samples were incubated at 25 °C

and fluorescence was measured at 480 nm, with excitation at 440 nm, every 60 s in a POLARstar

Omega microplate reader (BMG Labtech).

Fibril co-assembly for TEM

His_6-Ub-$RIPK3_{387-518}$ and $RIPK1_{497-583}$ and $M45_{1-90}$ were mixed together in 1:1 molar ratios or with

an appropriate volume of urea-containing buffer and then dialysed against 25 mM NaH_2PO_4,

150 mM NaCl, 0.5 mM DTT pH 7.4 for a minimum of 4 hours. Samples were removed and used to

prepare negatively stained grids for TEM, as above.

Fluorescence anisotropy assays

Covalent labelling of RIPK1$_{497-583}$ with the Alexa 488 fluorescence tag was performed in the

amyloid state. Samples of 1–4 mL of 100 μM RHIM fragment in 6 M GuHCl, obtained from the

cleavage of His$_6$-Ub-RIPK1$_{497-583}$, were dialysed against phosphate buffer at pH 8.0 with 0.5 mM

DTT at room temperature for 3h, followed by further dialysis against 20 mM NaH$_2$PO$_4$, 150 mM

NaCl pH 7.4 at pH 8.0 with two buffer changes. After the dialysis process, RIPK1-RHIM protein

was in an amyloid state and the sample was centrifuged at 16 000g for 10 min. The supernatant was

discarded and the amyloid pellet was resuspended in 1mL of 20 mM NaH$_2$PO$_4$, 150 mM NaCl, pH

8.0. A 50–μL volume of 10 mg/mL Alexa 488 (10 mg/mL in DMSO) was added to the amyloid

sample. The sample was incubated on a rocker for 1 h at room temperature and then unbound dye

was removed by centrifugation at 16000g for 10 min. The pellet was washed 3 times with 1 mL of

20 mM NaH$_2$PO$_4$, 150 mM NaCl, pH 8.0 and then resuspended in 1 mL of 8 M urea, 20 mM

NaH$_2$PO$_4$, 150 mM NaCl, pH 7.4 for storage. The degree of labelling was calculated to be 5%.

For polarization experiments, samples containing a molar ratio of 1:1999 of Alexa 488-RIPK1$_{497-583}$

to unlabelled RIPK1$_{497-583}$, RIPK3$_{387-518}$ or M45$_{1-90}$ in 20 mM NaH$_2$PO$_4$, 150 mM NaCl, pH 7.4

were prepared to a final volume of 300 μL. Total final protein concentrations of unlabelled protein

were 0.85 μM for RIPK1, 0.69 μM RIPK3 and 1.05 μM for M45 and the residual urea

concentration was 0.052 M. Three replicates of each protein sample were analysed simultaneously

in a 96-well black fluorescence plate sealed with optically clear film (Corning®). Samples were

incubated at 25 °C in a POLARstar Omega microplate reader (BMG Labtech) and excited at

480 nm. The parallel and perpendicular fluorescence emission was recorded at 520 nm, every 60 s.

These fluorescence intensities were used to calculate the fluorescence polarization.

X-ray fibre diffraction

His$_6$-Ub-M45$_{1-90}$ protein was allowed to assemble into fibrils by dialysis out of 8 M urea-containing

buffer into water overnight. The insoluble fibrils were pelleted by centrifugation at 16000g for

10 min, resuspended in a small volume of water and allowed to air-dry from a droplet suspended

between the wax-filled ends of two glass capillaries. X-ray fibre diffraction images were obtained

using an in-house Cu Kα Rigaku rotating anode source (wavelength 1.5418 Å) and MARresearch

image plate detector.

SDS-stability assays

Protein samples (0.5–1 mL) of individual fluorescent RHIM-containing constructs as well as

mixtures of these proteins were diluted to final individual protein concentrations of 20 μM with 8 M

urea, 25 mM NaH_2PO_4, 150 mM NaCl, pH 7.4. Approximately 50 μL of each sample was removed

for analysis as the monomeric, non-assembled control. The remaining solutions were dialysed

overnight at room temperature against 25 mM NaH_2PO_4, 150 mM NaCl, 0.5 mM DTT, pH 7.4.

Samples were removed and prepared with 4% glycerol and 0.0008% bromophenol blue and 0 or 2%

SDS, incubated at room temperature for 10 min and then analysed by electrophoresis using a 1%

agarose gel running in TAE buffer containing 0.1% SDS. Gels were imaged on a Bio-Rad Chemi-

Doc imaging system with 605/50nm and 695/55nm emission filters. In order to study the stability of

the amyloid fibrils by SDS-PAGE, mCHERRY RHIM-containing constructs, individually or in

mixtures, were dialysed against phosphate buffer overnight, at room temperature. After dialysis,

two 50–μL samples were centrifuged at 16,000g for 10 min and the resulting supernatants removed

and pooled (soluble sample). One pellet was resuspended with 50 μL of 8 M urea, 25 mM

NaH_2PO_4, 150 mM NaCl, pH 7.4 to allow visualisation of the insoluble material produced during

dialysis (insoluble sample). The other pellet was resuspended in 50 μL of 2% SDS in 25 mM

NaH_2PO_4, 150 mM NaCl, pH 7.4 and incubated at 37 °C for 10 minutes. SDS-containing samples

were then centrifuged at 16,000g for 10 min. The supernatant was removed (SDS-soluble sample)

and the pellet (SDS-insoluble sample) resuspended in 50 μL urea-containing buffer. All samples

were boiled with NuPAGE LDS sample buffer and NuPAGE reducing agent (50 mM DTT, Thermo

Scientific) and then analysed by SDS PAGE.

Confocal microscopy

Protein samples (0.5 mL) in 8 M urea containing mixtures of monomeric, purified YPET, ECFP

and mCHERRY RHIM-containing constructs at final individual protein concentrations of either 2.5

or 5 µM were dialysed overnight at room temperature against 25 mM NaH_2PO_4, 150 mM NaCl,

0.5 mM DTT, pH 7.4. Samples (15 µL) were placed on glass slides, protected with cover slips and

imaged on a Zeiss LSM 510 Meta confocal microscope.

Single molecule fluorescence spectroscopy

Mixtures of fluorescent fusion proteins with YPet or mCHERRY fluorophores were prepared at

70 µM in 8 M urea, 25 mM NaH_2PO_4, 150 mM NaCl, 0.5 mM DTT, pH 7.4 and then diluted 100–

fold with 25 mM NaH_2PO_4, 150 mM NaCl, 0.5 mM DTT, pH 7.4 in a 192-well silicone plate for

single molecule measurement before monitoring. Plates were analyzed at room temperature on a

Zeiss Axio Observer microscope with a custom-built data acquisition setup. Fluorescence signal

was collected and separated using a 565 nm dichroic mirror. For detection of YPET-tagged

proteins, signal was passed through a 525/20 nm band pass filter; for detection of mCHERRY-

tagged proteins a 580 nm long pass filter was used. For experiments with all three RHIM-

containing proteins, protein mixtures containing YPet-$RIPK3_{387-518}$, mCHERRY-$RIPK1_{497-583}$ and

His_6-Ub-$M45_{1-90}$ were prepared in 8 M urea-containing buffer, to provide samples containing the

RIPK3:RIPK1:M45 RHIMs in the range ~4:4:1 to ~1:1:4. Samples were diluted 100-fold into

25 mM NaH_2PO_4, 150 mM NaCl, 0.5 mM DTT, pH 7.4 and 20 µL pipetted into a 192-well silicone

plate for single molecule measurement. For experiments with multiple fluorophores, the signal from

the two channels was recorded simultaneously in 1 ms time bins.

Generation of M45-expressing HT-29 cells and cell death assays

$M45_{1-90}$ and $M45_{1-277}$ fragments were amplified from MCMV (Smith) and cloned upstream of the

E.Coli DNA gyrase subunit B (residues 2-220) in the pCDH-MCS-EF1-neo lentivector. All

insertions were validated by sequencing (Garvan Institute, Sydney, Australia). Lentivirus for $M45_{1-}$

$_{90}$-gyrase, $M45_{1-277}$-gyrase or empty vector control were made in 293T cells by co-transfecting the

lentivectors with psPAX2 and pMD2G using Fugene HD (Promega). HT-29s were then transduced

with the lentivirus particles in the presence of polybrene (Santa Cruz), and after 3 days G418

(Roche) was added for the subsequent 10 days to select for successfully transduced cells. $M45_{1-90}$-

gyrase, $M45_{1-277}$-gyrase or empty-vector-control transduced HT-29s were seeded in 96 well plates

(10^4 cells/well) and allowed to adhere for 6 h before treating with 900 nM coumermycin A1 (Enzo)

or media (no treatment control), followed 1 hour later with combinations of TNF (30 ng/ml, R and

D), BV-6 (1 μM, Selleckchem) and z-VAD-fmk (25 μM, R and D), or DMSO only (Sigma-

Aldrich) as control. Cell viability was then assessed 17-18 hour post-treatment by measuring levels

of intracellular ATP, using the CellTitre-Glo2 Assay (Promega). Data was expressed as percentage

of cell survival relative to the DMSO only control. Luminescence was measured using an Infinite

M1000 Pro plate reader (TECAN).

References

757   **References**

759   1.   Sun, X., Yin, J., Starovasnik, M.A., Fairbrother, W.J. & Dixit, V.M. Identification of a novel homotypic
760   interaction motif required for the phosphorylation of receptor-interacting protein (RIP) by RIP3. *J*
761   *Biol Chem* **277**, 9505-11 (2002).
762   2.   Newton, K. RIPK1 and RIPK3: critical regulators of inflammation and cell death. *Trends Cell Biol* **25**,
763   347-53 (2015).
764   3.   Sun, L. & Wang, X. A new kind of cell suicide: mechanisms and functions of programmed necrosis.
765   *Trends Biochem Sci* **39**, 587-93 (2014).
766   4.   Vandenabeele, P., Galluzzi, L., Vanden Berghe, T. & Kroemer, G. Molecular mechanisms of
767   necroptosis: an ordered cellular explosion. *Nat Rev Mol Cell Biol* **11**, 700-14 (2010).
768   5.   Kaiser, W.J., Sridharan, H., Huang, C., Mandal, P., Upton, J.W., Gough, P.J., Sehon, C.A., Marquis,
769   R.W., Bertin, J. & Mocarski, E.S. Toll-like receptor 3-mediated necrosis via TRIF, RIP3, and MLKL. *J*
770   *Biol Chem* **288**, 31268-79 (2013).
771   6.   Kaiser, W.J., Upton, J.W. & Mocarski, E.S. Viral modulation of programmed necrosis. *Curr Opin Virol*
772   **3**, 296-306 (2013).
773   7.   Rebsamen, M., Heinz, L.X., Meylan, E., Michallet, M.C., Schroder, K., Hofmann, K., Vazquez, J.,
774   Benedict, C.A. & Tschopp, J. DAI/ZBP1 recruits RIP1 and RIP3 through RIP homotypic interaction
775   motifs to activate NF-kappaB. *EMBO Rep* **10**, 916-22 (2009).
776   8.   Thapa, R.J., Ingram, J.P., Ragan, K.B., Nogusa, S., Boyd, D.F., Benitez, A.A., Sridharan, H., Kosoff, R.,
777   Shubina, M., Landsteiner, V.J., Andrake, M., Vogel, P., Sigal, L.J., tenOever, B.R., Thomas, P.G.,
778   Upton, J.W. & Balachandran, S. DAI Senses Influenza A Virus Genomic RNA and Activates RIPK3-
779   Dependent Cell Death. *Cell Host Microbe* **20**, 674-681 (2016).
780   9.   Cook, W.D., Moujalled, D.M., Ralph, T.J., Lock, P., Young, S.N., Murphy, J.M. & Vaux, D.L. RIPK1- and
781   RIPK3-induced cell death mode is determined by target availability. *Cell Death Differ* **21**, 1600-12
782   (2014).
783   10.   Murphy, J.M., Czabotar, P.E., Hildebrand, J.M., Lucet, I.S., Zhang, J.G., Alvarez-Diaz, S., Lewis, R.,
784   Lalaoui, N., Metcalf, D., Webb, A.I., Young, S.N., Varghese, L.N., Tannahill, G.M., Hatchell, E.C.,
785   Majewski, I.J., Okamoto, T., Dobson, R.C., Hilton, D.J., Babon, J.J., Nicola, N.A., Strasser, A., Silke, J.
786   & Alexander, W.S. The pseudokinase MLKL mediates necroptosis via a molecular switch
787   mechanism. *Immunity* **39**, 443-53 (2013).
788   11.   Murphy, J.M. & Silke, J. Ars Moriendi; the art of dying well - new insights into the molecular
789   pathways of necroptotic cell death. *EMBO Rep* **15**, 155-64 (2014).
790   12.   Orozco, S., Yatim, N., Werner, M.R., Tran, H., Gunja, S.Y., Tait, S.W., Albert, M.L., Green, D.R. &
791   Oberst, A. RIPK1 both positively and negatively regulates RIPK3 oligomerization and necroptosis.
792   *Cell Death Differ* **21**, 1511-21 (2014).
793   13.   Linkermann, A. & Green, D.R. Necroptosis. *N Engl J Med* **370**, 455-65 (2014).
794   14.   Oberst, A. Death in the fast lane: what's next for necroptosis? *FEBS J* **EPub ahead of print.**,
795   2015/09/24 (2015).
796   15.   Caccamo, A., Branca, C., Piras, I.S., Ferreira, E., Huentelman, M.J., Liang, W.S., Readhead, B., Dudley,
797   J.T., Spangenberg, E.E., Green, K.N., Belfiore, R., Winslow, W. & Oddo, S. Necroptosis activation in
798   Alzheimer's disease. *Nat Neurosci* **20**, 1236-1246 (2017).
799   16.   Degterev, A. & Linkermann, A. Generation of small molecules to interfere with regulated necrosis.
800   *Cellular and Molecular Life Sciences* **73**, 2251-2267 (2016).
801   17.   Seifert, L., Werba, G., Tiwari, S., Giao Ly, N.N., Alothman, S., Alqunaibit, D., Avanzi, A., Barilla, R.,
802   Daley, D., Greco, S.H., Torres-Hernandez, A., Pergamo, M., Ochi, A., Zambirinis, C.P., Pansari, M.,
803   Rendon, M., Tippens, D., Hundeyin, M., Mani, V.R., Hajdu, C., Engle, D. & Miller, G. The Necrosome
804   Promotes Pancreas Oncogenesis via CXCL1 and Mincle Induced Immune Suppression. *Nature* **532**,
805   245-249 (2016).
806   18.   Mocarski, E.S., Guo, H. & Kaiser, W.J. Necroptosis: The Trojan horse in cell autonomous antiviral
807   host defense. *Virology* **479-480**, 160-6 (2015).

19. Baker, M., Shanmugam, N., Pham, C.L.L., Strange, M., Steain, M. & Sunde, M. RHIM-based
protein:protein interactions in anti-microbial defence against programmed cell death by
necroptosis. *Semin Cell Dev Biol* (2018).
20. Fliss, P.M. & Brune, W. Prevention of cellular suicide by cytomegaloviruses. *Viruses* **4**, 1928-49
(2012).
21. Daley-Bauer, L.P., Roback, L., Crosby, L.N., McCormick, A.L., Feng, Y., Kaiser, W.J. & Mocarski, E.S.
Mouse cytomegalovirus M36 and M45 death suppressors cooperate to prevent inflammation
resulting from antiviral programmed cell death pathways. *Proc Natl Acad Sci U S A* **114**, E2786-
e2795 (2017).
22. Mocarski, E.S., Kaiser, W.J., Livingston-Rosanoff, D., Upton, J.W. & Daley-Bauer, L.P. True grit:
programmed necrosis in antiviral host defense, inflammation, and immunogenicity. *J Immunol* **192**,
2019-26 (2014).
23. Upton, J.W., Kaiser, W.J. & Mocarski, E.S. Cytomegalovirus M45 cell death suppression requires
receptor-interacting protein (RIP) homotypic interaction motif (RHIM)-dependent interaction with
RIP1. *J Biol Chem* **283**, 16966-70 (2008).
24. Brune, W., Menard, C., Heesemann, J. & Koszinowski, U.H. A ribonucleotide reductase homolog of
cytomegalovirus and endothelial cell tropism. *Science* **291**, 303-5 (2001).
25. Guo, H., Omoto, S., Harris, P.A., Finger, J.N., Bertin, J., Gough, P.J., Kaiser, W.J. & Mocarski, E.S.
Herpes simplex virus suppresses necroptosis in human cells. *Cell Host Microbe* **17**, 243-51 (2015).
26. Upton, J.W., Kaiser, W.J. & Mocarski, E.S. Virus inhibition of RIP3-dependent necrosis. *Cell Host*
Microbe **7**, 302-13 (2010).
27. Lembo, D., Donalisio, M., Hofer, A., Cornaglia, M., Brune, W., Koszinowski, U., Thelander, L. &
Landolfo, S. The ribonucleotide reductase R1 homolog of murine cytomegalovirus is not a
functional enzyme subunit but is required for pathogenesis. *J Virol* **78**, 4278-88 (2004).
28. Upton, J.W., Kaiser, W.J. & Mocarski, E.S. DAI/ZBP1/DLM-1 complexes with RIP3 to mediate virus-
induced programmed necrosis that is targeted by murine cytomegalovirus vIRA. *Cell Host Microbe*
11, 290-7 (2012).
29. Li, J., McQuade, T., Siemer, A.B., Napetschnig, J., Moriwaki, K., Hsiao, Y.S., Damko, E., Moquin, D.,
Walz, T., McDermott, A., Chan, F.K. & Wu, H. The RIP1/RIP3 necrosome forms a functional amyloid
signaling complex required for programmed necrosis. *Cell* **150**, 339-50 (2012).
30. Pham, C.L., Kwan, A.H. & Sunde, M. Functional amyloid: widespread in Nature, diverse in purpose.
Essays Biochem **56**, 207-19 (2014).
31. Kleino, A., Ramia, N.F., Bozkurt, G., Shen, Y., Nailwal, H., Huang, J., Napetschnig, J., Gangloff, M.,
Chan, F.K., Wu, H., Li, J. & Silverman, N. Peptidoglycan-Sensing Receptors Trigger the Formation of
Functional Amyloids of the Adaptor Protein Imd to Initiate Drosophila NF-kappaB Signaling.
Immunity **47**, 635-647 e6 (2017).
32. Mompean, M., Li, W., Li, J., Laage, S., Siemer, A.B., Bozkurt, G., Wu, H. & McDermott, A.E. The
Structure of the Necrosome RIPK1-RIPK3 Core, a Human Hetero-Amyloid Signaling Complex. *Cell*
(2018).
33. Macindoe, I., Kwan, A.H., Ren, Q., Morris, V.K., Yang, W., Mackay, J.P. & Sunde, M. Self-assembly of
functional, amphipathic amyloid monolayers by the fungal hydrophobin EAS. *Proc Natl Acad Sci U S*
A **109**, E804-11 (2012).
34. Sunde, M. & Blake, C. The structure of amyloid fibrils by electron microscopy and X-ray diffraction.
Advances in Protein Chemistry, Vol 50 **50**, 123-159 (1997).
35. LeVine, H., 3rd. Quantification of beta-sheet amyloid fibril structures with thioflavin T. *Methods*
Enzymol **309**, 274-84 (1999).
36. Gambin, Y., Polinkovsky, M., Francois, B., Giles, N., Bhumkar, A. & Sierecki, E. Confocal
Spectroscopy to Study Dimerization, Oligomerization and Aggregation of Proteins: A Practical
Guide. *Int J Mol Sci* **17**(2016).
37. Sierecki, E., Giles, N., Bowden, Q., Polinkovsky, M.E., Steinbeck, J., Arrioti, N., Rahman, D., Bhumkar,
A., Nicovich, P.R., Ross, I., Parton, R.G., Bocking, T. & Gambin, Y. Nanomolar oligomerization and
selective co-aggregation of alpha-synuclein pathogenic mutants revealed by single-molecule
fluorescence. *Sci Rep* **6**, 37630 (2016).

38.   Hennig, S., Kong, G., Mannen, T., Sadowska, A., Kobelke, S., Blythe, A., Knott, G.J., Iyer, K.S., Ho, D.,
862   Newcombe, E.A., Hosoki, K., Goshima, N., Kawaguchi, T., Hatters, D., Trinkle-Mulcahy, L., Hirose, T.,
863   Bond, C.S. & Fox, A.H. Prion-like domains in RNA binding proteins are essential for building
864   subnuclear paraspeckles. *J Cell Biol* **210**, 529-39 (2015).
865   39.   Kato, M., Han, T.W., Xie, S., Shi, K., Du, X., Wu, L.C., Mirzaei, H., Goldsmith, E.J., Longgood, J., Pei, J.,
866   Grishin, N.V., Frantz, D.E., Schneider, J.W., Chen, S., Li, L., Sawaya, M.R., Eisenberg, D., Tycko, R. &
867   McKnight, S.L. Cell-free formation of RNA granules: low complexity sequence domains form
868   dynamic fibers within hydrogels. *Cell* **149**, 753-67 (2012).
869   40.   Wu, H. & Fuxreiter, M. The Structure and Dynamics of Higher-Order Assemblies: Amyloids,
870   Signalosomes, and Granules. *Cell* **165**, 1055-66 (2016).
871   41.   Wu, X.N., Yang, Z.H., Wang, X.K., Zhang, Y., Wan, H., Song, Y., Chen, X., Shao, J. & Han, J. Distinct
872   roles of RIP1-RIP3 hetero- and RIP3-RIP3 homo-interaction in mediating necroptosis. *Cell Death*
873   *Differ* **21**, 1709-20 (2014).
874   42.   Murphy, J.M. & Vince, J.E. Post-translational control of RIPK3 and MLKL mediated necroptotic cell
875   death. *F1000Res* **4**(2015).
876   43.   Huang, Z., Wu, S.Q., Liang, Y., Zhou, X., Chen, W., Li, L., Wu, J., Zhuang, Q., Chen, C., Li, J., Zhong,
877   C.Q., Xia, W., Zhou, R., Zheng, C. & Han, J. RIP1/RIP3 binding to HSV-1 ICP6 initiates necroptosis to
878   restrict virus propagation in mice. *Cell Host Microbe* **17**, 229-42 (2015).
879   44.   Guo, H., Kaiser, W.J. & Mocarski, E.S. Manipulation of apoptosis and necroptosis signaling by
880   herpesviruses. *Med Microbiol Immunol* **204**, 439-48 (2015).
881   45.   Pearson, J.S., Giogha, C., Muhlen, S., Nachbur, U., Pham, C.L., Zhang, Y., Hildebrand, J.M., Oates,
882   C.V., Lung, T.W., Ingle, D., Dagley, L.F., Bankovacki, A., Petrie, E.J., Schroeder, G.N., Crepin, V.F.,
883   Frankel, G., Masters, S.L., Vince, J., Murphy, J.M., Sunde, M., Webb, A.I., Silke, J. & Hartland, E.L.
884   EspL is a bacterial cysteine protease effector that cleaves RHIM proteins to block necroptosis and
885   inflammation. *Nat Microbiol* **2**, 16258 (2017).
886   46.   Lin, J., Kumari, S., Kim, C., Van, T.M., Wachsmuth, L., Polykratis, A. & Pasparakis, M. RIPK1
887   counteracts ZBP1-mediated necroptosis to inhibit inflammation. *Nature* **540**, 124-128 (2016).
888   47.   Newton, K., Wickliffe, K.E., Maltzman, A., Dugger, D.L., Strasser, A., Pham, V.C., Lill, J.R., Roose-
889   Girma, M., Warming, S., Solon, M., Ngu, H., Webster, J.D. & Dixit, V.M. RIPK1 inhibits ZBP1-driven
890   necroptosis during development. *Nature* **540**, 129-133 (2016).
891   48.   Sridharan, H., Ragan, K.B., Guo, H., Gilley, R.P., Landsteiner, V.J., Kaiser, W.J. & Upton, J.W. Murine
892   cytomegalovirus IE3-dependent transcription is required for DAI/ZBP1-mediated necroptosis.
893   *EMBO Rep* **18**, 1429-1441 (2017).
894   49.   Zhou, W. & Yuan, J. Necroptosis in health and diseases. *Semin Cell Dev Biol* **35**, 14-23 (2014).

Figure 1.

Figure 2.

Figure 3.

Figure 4.

Figure 5.

a.

Figure 6.

a.

b.

c.

d.

e.

f.