# When parasites are selected to kill the young

Ryosuke Iritani[*,1,2], Elisa Visher[2] & Mike Boots[2]
May 21, 2018

Author for correspondance: RI, E-mail: Lambtani@gmail.com

Contact information:

1. Biosciences, College of Life and Environmental Science, University of Exeter, Exeter, United Kingdom.
2. Department of Integrative Biology, University of California, Berkeley, CA 94720, USA.

**Abstract**

The impact of infectious disease is often very different in juveniles and adults, but theory has focused on the drivers of stage-dependent defense in hosts rather than stage-dependent virulence evolution. Stage-structure has the potential to be important to the evolution of pathogens because it exposes parasites to heterogeneous environments in terms of both host characteristics and transmission routes. We develop a stage-structured (juvenile-adult) epidemiological model and examine the evolutionary outcomes of stage-specific virulence under the classic assumption of a transmission-virulence trade-off. We show that selection on virulence against adults remains consistent with the classic theory. However, the evolution of juvenile virulence is sensitive to both demography and contact structure with higher virulence against juveniles being favored either when the contact structure is assortative (juveniles preferentially interact together) and the juvenile stage is short, or in contrast when the contact structure is disassortative and the juvenile stage is long. These results can be understood in the context of Fisher's reproductive value theory and highlight the potentially profound effects of host stage-structure on determining parasite virulence in nature. This new perspective may have broad implications for both understanding and managing disease severity.

## Introduction

Understanding how parasites are selected to exploit their hosts remains a central research question in the evolutionary ecology of host-parasite interactions [1–11], with important implications for host persistence [12, 13], disease management [14], and host-parasite coevolution [15]. Theory on parasite evolution is typically based on trade-offs imposed between transmission rate and virulence [defined in this literature as the increased death rate due to infection; 3, 16]. Specifically, the transmission-virulence trade-off hypothesis posits that high host exploitation by the parasite leads to high transmission but also results in higher virulence (reviewed in [8, 16]). Theoretically, evolutionarily stable exploitation occurs when the marginal increase in transmission due to exploitation equals the marginal increase in host mortality due to exploitation. This consequently optimizes parasite fitness [e.g., 3, 17–19] (but see [20]). Another, less studied trade-off is between virulence and rate of recovery, with more rapidly growing parasites being harder to clear but causing more damage [3]. Such trade-offs are fundamental to understanding the evolutionary drivers of the virulence of infectious diseases and offer a number of important insights for disease management [21]. However, despite the considerable variation in virulence that is found at different life-stages in nature [22], the implications of host stage structure to parasite virulence has not been examined and the differences in virulence between adults and juveniles are typically explained in terms of differences in host defense [22, 23].

A number of recent ecological studies have examined the impact of a host populations' stage-related heterogeneities in disease response on disease epidemiology [e.g., 24–26]. In these studies, the differences in the impact of infection between life stages have been assumed to be properties of the host and driven by processes like maternal and acquired immunity and age-related variation in tolerance, resistance, exposure, immunocompetence, and susceptibility [22, 23]. In principle, however, this variation in disease outcomes across different host life-stages (e.g., egg, larvae, juvenile, adult, etc) has the potential to significantly impact the evolutionary dynamics of virulence. Additionally, stage structure in host populations can also lead to variation in transmission routes between and across stages [reviewed in 27–29]. For instance, juvenile-juvenile and adult-adult contacts might be more likely than juvenile-adult contacts if juvenile and adult hosts are segregated in spatio-temporal niches, typical of humans [30], amphibians [31], and insects [32]. Such variation in assortativity or disassortivity of transmission between life stages can create differential selection pressures where a parasite may be selected to bias its virulence towards different stages ("stage-specific virulence"). We propose that the differences in virulence at different ages may not necessarily be properties of the host, but instead could be selected for

as adaptive properties of the parasites (Fig 1).

Here, we show that hosts' stage-related contact structure and the period of reproductive (and thus adult) stage can in concert drive the evolution of stage-specific virulence. We develop a mathematical model and explore the evolutionary outcomes of stage-specific virulence under different patterns of contacts across stages. We explicitly model the stage-structured host population dynamics for juveniles and adults including explicit epidemiology, and analyze the evolutionary dynamics using the adaptive dynamics toolbox [33, 34]. We first show that the evolutionary outcomes of virulence against adults ("adult-virulence") follow classic results such that background mortality rate in adults favors higher virulence. Second, we show that the evolutionary outcomes of virulence against juveniles ("juvenile-virulence") are critically impacted by the interplay between assortativity and maturation. We explain the results in terms of Fisher's reproductive value to account for life-history evolution of parasites [19, 35–43] and in terms of transmission pathways (and thus the pathways to reproductive success of parasites). We find a number of robust examples in the empirical literature that match our predictions and therefore highlight the importance of age-structure to the evolution of infectious disease.

Figure 1: Empirical data on stage-specific virulence. In the following, H indicates "host" while P indicates "parasites" : (a) H: Pink Salmon (*Oncorhynchus gorbuscha*); P: Salmon Louse [44, 45]. (b) H: Gerbil (*Gerbillus andersoni*); P: Ectoparasites [46, 47]. (c) H: European Rabbit (*Oryctolagus cuniculus*); P: Nematode [48, 49]. (d) H: Rabbits (*Leporidae*); P: RHD Virus [50, 51]. (e) H: Common Guillemot (*Uria aalge*); P: Great Island Virus [52–54]. (f) H: Asian Elephant (*Elephas maximus*); P: Parasites [55, 56]. (g) H: Pigeon (*Columba livia*) P: Blood parasites [57, 58]

Figure 2: A schematic illustration of the model on the force of infection and assortativity. (A) Force of infection from Y to X, $\phi_{XY}$, is a compound factor of susceptibility (likeliness of X-hosts becoming infected), contact intensity (likeliness of contacts between X and Y), and infectiousness (propagule production of the parasite carried by Y). (B-D) Degrees of assortativity. Negative assortativity indicates that contacts occur more frequently between stages than within stages (panel B). The contact structure is unbiased (random) when $\rho = 0$ (panel C). Positive assortativity indicates that contacts occur more frequently within stages than between stages (panel D).

## Method

We consider a host population subdivided into juvenile (J) and adult (A) stages, in which juveniles are obviously incapable of reproduction. The density of susceptible or infected juveniles is denoted $S_J$ or $I_J$ respectively, and that of susceptible or infected adults is denoted $S_A$ or $I_A$ respectively. Combining an epidemiological SI-model with a stage-structured model [59] yields the following ordinary differential equations (ODEs):

$$
\begin{aligned}
\frac{dS_J}{dt} &= \left(r - \kappa(S_A + I_A)\right) \cdot (S_A + I_A) - \left(u + \phi_{JA} + \phi_{JJ} + m_J\right) S_J, \\
\frac{dS_A}{dt} &= uS_J - \left(m_A + \phi_{AJ} + \phi_{AA}\right) S_A \\
\frac{dI_J}{dt} &= \left(\phi_{JJ} + \phi_{JA}\right) S_J - \left(u + m_J + v_J\right) I_J \\
\frac{dI_A}{dt} &= \left(\phi_{AJ} + \phi_{AA}\right) S_A + uI_J - \left(m_A + v_A\right) I_A,
\end{aligned}
\tag{1}
$$

where: $r$ represents intrinsic growth rate of the adult hosts per capita (and we assume that susceptible and infected hosts have the same fecundity); there is no reproduction from juveniles. Reproduction is

reduced by s density-dependent factor $\kappa$; $u$ represents the maturation rate; $\phi_{XY}$ represents the rate at which a susceptible stage-X host gets transmitted from an infected stage-Y host (i.e., force of infection from infectious Y to susceptible X per capita; Fig 2); $m_X$ represents the background mortality for a stage-X host; $v_X$ represents the virulence against a stage-X host, as an evolving trait. Note that we choose to consider discrete stages instead of continuous-stages to quantify the relative strength of exploitation against juveniles and adults. Other approach including physiologically structured population model (synthesized in [60]) or infection-age model [61, 62] is also applicable, but we chose to use a discrete stage model for the sake of simplicity.

The relative, average time a host individual spends as an adult becomes longer with faster maturation for fixed natural mortality rates. For the sake of relative quantification of this, we first calculate the hosts' average time that is spent in the juvenile (or adult) stage in the absence of disease, denoted $T_J$ (or $T_A$). Using the basic property of the exponential distributions, we get:

$$
\begin{aligned}
T_J &= \frac{1}{m_J + u}, \\
T_A &= \frac{1}{m_A},
\end{aligned}
\tag{2}
$$

where the latter is conditional on successful maturation to adulthood. Second, the probability of successful maturation of a susceptible juvenile to its adulthood, denoted $P_{AJ}$, is given by:

$$
P_{AJ} = \frac{u}{u + m_J}.
\tag{3}
$$

Therefore, the fraction of the average time that is spent in the adult stage ("post-maturation span") is given by:

$$
\theta_A := \frac{P_{AJ} T_A}{T_J + P_{AJ} T_A} = \frac{u}{m_A + u}
\tag{4}
$$

from which we can see that faster maturation results in the longer post-maturation span. We use $\theta_A$ as a characteristic parameter of the stage-structured host populations.

The force of infection for a stage-X host from a stage-Y host (where X and Y run across J and A) involves with three processes: susceptibility $\alpha_X$ (the rate at which a stage-X host gets infected given exposure to infectious propagules), contact structure $\sigma_{XY}$ (which represents the fraction of time of a stage-Y host spends with a stage-X host), and infectiousness $\beta_Y$ (the rate of propagule production from

 a stage-Y host; reviewed in [28]):

$$\phi_{JJ} = \frac{\alpha_J \sigma_{JJ} \beta_J I_J}{S_J + S_A + I_J + I_A},$$
$$\phi_{JA} = \frac{\alpha_J \sigma_{JA} \beta_A I_A}{S_J + S_A + I_J + I_A},$$
$$\phi_{AJ} = \frac{\alpha_A \sigma_{AJ} \beta_J I_J}{S_J + S_A + I_J + I_A}, \tag{5}$$
$$\phi_{AA} = \frac{\alpha_A \sigma_{AA} \beta_A I_A}{S_J + S_A + I_J + I_A}$$

(see Fig 2). Here, Eqn (5) assumes that the transmission follows a frequency-dependent mass-action model [63]. To quantify the contact structure, we use a single parameter of contct structure $\sigma = 1 - \sigma_{AJ} = \sigma_{JJ} = 1 - \sigma_{JA} = \sigma_{AA}$, accounting for the assumption that within-class contacts decrease with between-class contacts [and vice versa; 27, 30, 64]. With this symmetry, "assortativity" is given by:

$$\rho = \sigma_{JJ} + \sigma_{AA} - 1 = 2\sigma - 1, \tag{6}$$

where $\rho$ varies from $-1$ to $1$ [65–68]. If $-1 \leq \rho < 0$, then within-stage contact is less frequent compared to between-stage contact (such a contact is said to be "disassortative"). Instead, if $0 < \rho \leq 1$, then within-stage contact is more likely than between-stage contact ("assortative" contact). $\rho = 0$ indicates that contact is unbiased ("random" contact). In the extreme case, $\rho = 1$ (or $-1$) indicates that contact (and consequently transmission) occurs exclusively within stages.

A final ingredient is the transmission-virulence trade-off, formulated by:

$$\beta_J = b_J \frac{k_J v_J}{1 + k_J v_J},$$
$$\beta_A = b_A \frac{k_A v_A}{1 + k_A v_A}, \tag{7}$$

where $k_X$ tunes the degree of steepness or the efficiency of virulence for infectiousness from stage-X hosts; $b_X$ represent the upper bounds of infectiousness from stage-X hosts. Here, we have assumed that infectiousness increases with virulence; hence, the trade-off is explicitly imposed between infectiousness and virulence.

We use the adaptive dynamics toolbox [33, 34] to study the long-term evolutionary dynamics. First, suppose that the demographic and epidemiological dynamics have quickly reached a steady state:

$(S_J, S_A, I_J, I_A) = (S_J^*, S_A^*, I_J^*, I_A^*)$, which is uniquely determined by the ODE. We then introduce a rare mutant of small phenotypic changes in host stage-specific virulence, $\mathbf{v}' := (v_J', v_A')$ attempting to invade a monomorphic, resident type ("wild type") virulence $\mathbf{v} := (v_J, v_A)$. We assume that the differences in virulence between mutant and wild types are very small (phenotypically weak selection).

To assess the possibility of mutant invasion, we define the invasion fitness, denoted $w$, by using the Next-Generation Theorem [19, 69, 70]. The "next-generation matrix" (that governs the population dynamics of the rare mutant and thus its long term growth) can be written as the product of five matrices, given by:

$$
\mathbf{G}' = \underbrace{\begin{pmatrix} S_J^* & 0 \\ 0 & S_A^* \end{pmatrix}}_{\text{availability}} \underbrace{\begin{pmatrix} \alpha_J & 0 \\ 0 & \alpha_A \end{pmatrix}}_{\text{susceptibility}} \underbrace{\begin{pmatrix} \sigma_{JJ} & \sigma_{JA} \\ \sigma_{AJ} & \sigma_{AA} \end{pmatrix}}_{\text{contact}} \underbrace{\begin{pmatrix} \frac{\beta_J'}{H^*} & 0 \\ 0 & \frac{\beta_A'}{H^*} \end{pmatrix}}_{\text{infectiousness}} \underbrace{\begin{pmatrix} \frac{1}{\mu_J'} & 0 \\ \frac{u}{\mu_J'\mu_A'} & \frac{1}{\mu_A'} \end{pmatrix}}_{\text{infectious period}} , \tag{8}
$$

where $H^* = S_J^* + S_A^* + I_J^* + I_A^*$ (the total density of the hosts), $\mu_J' = u + m_J + v_J'$ (the loss rate of infected juveniles with maturation being included), and $\mu_A' = m_A + v_A'$ (the mortality rate of infected adults). The decomposition of $\mathbf{G}'$ into the product of matrices allows for a natural interpretation by partitioning the epidemiological process and is consistent with the proposed approach to transmission dynamics in heterogeneous host populations (reviewed in [27–29]). The first matrix restores the availability of susceptible hosts, each with a specific susceptibility (the second matrix); the third matrix represents the contact pattern with infected hosts; the fourth matrix represents the infectiousness of infected hosts per capita, and parasites impact the infectious period among hosts with the effect of maturation from juveniles to adults being included (the fifth matrix).

The invasion fitness is given by the dominant eigenvalue of $\mathbf{G}'$ (denoted $\Lambda[\mathbf{G}']$), which turns out to exhibit a complicated expression; therefore, we choose to use a simpler but equivalent measure for invasion fitness using graph-theoretical method developed by Camino Beck & Lewis [71, 72] and Camino Beck, Lewis, & Driessche [73], given by:

$$
w(\mathbf{v}', \mathbf{v}) = \alpha_J \frac{S_J^*}{H^*} \sigma_{JJ} \frac{\beta_J'}{\mu_J'} + \frac{u}{\mu_J'} \cdot \alpha_J \frac{S_J^*}{H^*} \sigma_{JA} \frac{\beta_A'}{\mu_A'} + \alpha_A \frac{S_A^*}{H^*} \sigma_{AA} \frac{\beta_A'}{\mu_A'} - \alpha_J \alpha_A \frac{\rho S_J^* S_A^*}{(H^*)^2} \cdot \frac{\beta_J' \beta_A'}{\mu_J' \mu_A'} \tag{9}
$$

(also see [39, 68, 70, 74]). The condition for the mutant type to outcompete the wild type (i.e., invadability condition), $w(\mathbf{v}', \mathbf{v}) > 1$, holds if and only if $\Lambda[\mathbf{G}'] > 1$ (for more details, see Appendix).

Virulence evolves in the direction of selection gradient, given by:

$$g_J(\mathbf{v}) = \frac{\partial w(\mathbf{v}', \mathbf{v})}{\partial v_J'},$$

$$g_A(\mathbf{v}) = \frac{\partial w(\mathbf{v}', \mathbf{v})}{\partial v_A'},$$

(10)

where the partial derivatives are evaluated at $\mathbf{v}' = \mathbf{v}$ ("neutrality"). Evolution ceases at which both gradients are nullified ("Singular Strategy", SS).

We assess two stability criteria of the singular strategy. The first criterion, attainability [75, 76], concerns whether recurrent substitutions of genes from wild to mutant can lead to the convergence of the strategy to SS. The second is referred to as evolutionary stability [77], which assures that SS can resist against any invasion of alternative, mutant strategies. If SS meets both of these criteria, it is then called as Continuously Stable Strategy (CSS [78]). Analytical investigation revealed that the SS is always a CSS, and thus we do not detail the stability analyses below (Appendix). We will hereafter superscriptize an asterisk ($^*$) on CSS.

We use the following default parameter-values: $r = 6$, $\kappa = 0.06$, $h = 0$, $m_J = 1$, $\alpha_J = \alpha_A = 1$, $k_J = k_A = 1$, $b_J = b_A = 10$, while varying $u$ and $\rho$. That is, the parameter values are symmetric for juveniles and adults. We subsequently check the effects of the difference in $\alpha$ (susceptibility), $k$ (efficiency of exploitation for transmission), and $b$ (upper bound in infectiousness). Finally, we check whether recovery or tolerance in the host can affect the results.

## Results

We first derive the selection gradient along $v_A$:

$$g_A(\mathbf{v}) = \left\{ \frac{\alpha_A S_A^* \sigma_{AA}}{H^*} \cdot \left(1 - \frac{\alpha_J S_J^* \sigma_{JJ}}{H^*} \cdot \frac{\beta_J}{\mu_J}\right) + \frac{\alpha_J S_J^* \sigma_{JA}}{H^*} \cdot \left(\frac{u}{\mu_J} + \frac{\alpha_A S_A^* \sigma_{AJ}}{H^*} \cdot \frac{\beta_J}{\mu_J}\right) \right\}$$

$$\times \frac{\beta_A}{\mu_A} \cdot \left(\frac{1}{\beta_A} \cdot \frac{d\beta_A}{dv_A} - \frac{1}{\mu_A}\right),$$

(11)

which is consistent with a number of previous studies: under the transmission-virulence trade-off, higher exploitation is expected to increase the infectiousness (i.e., a marginal benefit) at the immediate (marginal) costs owing to reduced infectious period [8, 11, 39, 42, 79–81]. Therefore, the direction of selection on adult virulence is completely determined by the balance between such benefits and costs.

In terms of juvenile-virulence, however, an additional term emerges in the present model because of the host maturation rate $u$. To make the biological meaning of this term clearer, we use the reproductive-value based form of the selection gradient [19, 35, 36, 38–43], which reads:

$$g_J(\mathbf{v}) \propt \left( \ell_J^* \cdot \underbrace{\frac{\alpha_J S_J^* \sigma_{JJ}}{H^*}}_{\text{from J to J}} + \ell_A^* \cdot \underbrace{\frac{\alpha_A S_A^* \sigma_{AJ}}{H^*}}_{\text{from J to A}} \right) \cdot \frac{\beta_J}{\mu_J} \times \left( \frac{1}{\beta_J} \cdot \frac{d\beta_J}{dv_J} - \frac{1}{\mu_J} \right) - \ell_A^* \frac{u}{\mu_J} \cdot \frac{1}{\mu_J}, \qquad (12)$$

where $(\ell_J^*, \ell_A^*)$ represents the pair of individual reproductive values of the parasites carried by juvenile and adult hosts (or the left eigenvector of $\mathbf{G}$ at neutrality; Appendix). The first term represents the sum of reproductive success owing to transmission from an infected juvenile hosts to a susceptible juvenile and to a susceptible adult; once transmitted to a juvenile (or adult) host, the parasites can gain the relative reproductive success $\ell_J^*$ (or $\ell_A^*$, respectively). In total, the first term obeys classic marginal value theorem [8, 11, 17, 18, 39, 42, 79–81] such that the marginal increase in transmission due to exploitation confers a benefit but the marginal decrease in infectious period imposes a cost on exploitation. The second term represents the reduction in successful maturation due to killing the juveniles and this term involves $\ell_A^*$ (the individual reproductive value of the parasites infecting adults) times the probability of maturation of infected juveniles $u/\mu_J$. This is because killing juveniles can lead to the loss of expected reproductive success via adult hosts that the parasites could otherwise gain through the maturation of the juvenile host [42]. In other words, killing the juvenile hosts can result in the reduction of prospective fitness. The final multiplicative factor $1/\mu_J$ represents the marginal increase in juvenile mortality due to increasing virulence.

We looked at the effects of (i) post-maturation span $\theta_A$ and (ii) stage-assortativity $\rho$, on SS. Strikingly, the SS for adult virulence is necessarily $v_A^* = \sqrt{m_A/k_A}$, which is independent of any demographic and disease characteristics of juveniles. This is because the parasites infecting adults can utilize a single transmission pathway from adults (to any susceptible hosts in the population). Hence, we used $v_A^*$ as a benchmark result and compared it with $v_J^*$.

In contrast, SS for juvenile-virulence is dramatically affected by the densities of susceptible hosts (ecological feedback), the adult virulence (epidemiological feedback), and stage-assortativity (demographic feedback). This is because the parasites infecting the juveniles can utilize two pathways of transmission: either from the juvenile (to any susceptible hosts), or from the adult who has successfully

matured from the juvenile stage. The analytical expression for $v_J^*$ is intractable, and thus we numerically

evaluated $v_J^*$.

We can immediately see that $v_A^*$ increases with $m_A$, in agreement with the previous studies [reviewed

in 8, 11]. Although we can isolate the pure effect of each parameter (e.g., $m_J$, $k_J$) on $v_J^*$, we chose to plot the

relative virulence $v_J^*/v_A^*$ against the assortatvity ($\rho$, abscissa) and post-maturation span ($\theta_A$, ordinate).

We found that either disassortative hosts with a long post-maturation span or assortative hosts with

a short post-maturation span select for higher virulence against juveniles. This result slightly changes

given stage-specific mortality rates ($m_J \neq m_A$), but the general trend is robust (Fig 3A-C).

By relaxing the assumptions of the symmetry in disease-related parameters $k_J, k_A$ (efficiency

of exploitation for transmission), $b_J, b_A$ (maximum transmissibility), and $\alpha_J, \alpha_A$ (susceptibility) for

juveniles and adults, or by incorporating recovery or tolerance, we showed that the results are robust

and qualitatively unchanged. Therefore, we conclude that the combined effects of maturation and

assortativity are critical to the evolution of virulence.

Figure 3: Left panels: Evolutionary outcomes of relative virulence ($v_J^*/v_A^*$). Red indicates $v_J^* > v_A^*$ and blue indicates $v_J^* < v_A^*$. Color scales used are the same in the three panels. Middle panels: Densities of infected juveniles at equilibrium, $I_J^*$. Right panels: Densities of susceptible juveniles at equilibrium, $S_J^*$. In each panel, abscissa: assortativity; ordinate: post-maturation span $\theta_A$; from (A) to (C): $m_A = 0.8, 1.0, 1.25$ as indicated; HE: host-extinction; PE: parasite-extinction (or evolutionary suicide); dotted curve: $v_J^* = v_A^*$; parameters: default values.

## Discussion

We have shown theoretically how parasites are subject to different selective pressures when they infect adults or juveniles. In particular, the key prediction is that the combination between maturation and contact-structure – fast maturation with disassortativity, or slow maturation with assortativity – has a dramatic impact on optimal juvenile-virulence. Higher virulence against juveniles is favored either if: (i) adult-stage is relatively long and the contact-structure is disassortative (between age class interactions are high; Fig 3, left-top zone), (ii) juvenile-stage is relatively long and the contact structure is assortative (interactions occur preferentially within classes; Fig 3, right-bottom zone). This result can be understood as follows: given that post-maturation span is long, and the contact structure is disassortative, adult hosts are abundant in the population and the transmission from juveniles to adults is more likely than from adults to juveniles; in this case, the availability of adult hosts is higher, which selects for higher exploitation against juveniles to obtain more abundant resource (adults hosts). Spatial and/or temporal segregation in the niches of juveniles and adults therefore has the potential to be an important evolutionary driver of virulence. Previous theory has overlooked the phenomena that virulence is highly sensitive to stage-structured life-history characteristics of hosts such as ontogeny and associated, spatio-temporal niche-shifts.

The incorporation of the maturation of the hosts in our model shows that higher parasite exploitation against juveniles incurs an additional cost associated with increased maturation failure [42]. In addition, non-random assortativity generates additional selective pressures [39, 82]. In particular, while marginal value theorem does correctly predict the evolutionary outcomes of adult virulence it does not predict that of juvenile-virulence. Therefore, sources of heterogeneity in hosts can clearly lead to different predictions than classic virulence evolution theory based on the marginal value theorem. Gandon [39] and Osnas & Dobson [82] introduced multiple hosts' types or species and studied conditional virulence against them while Williams and colleagues [41, 42, 83, 84] have proposed to use reproductive value theory to study parasite evolution in heterogeneous host populations. However, none of these studies are devoted to stage structure with associated stage-specific virulence. Our novel results arise because we explicitly assumed stage structure with maturation from juveniles to adults and reproduction by adults rather than more generic heterogeneity between different types of hosts.

Finding examples of stage-specific virulence in empirical systems can be difficult due to the intricacies of specific host-pathogen systems. Stage-related trends in virulence can be complicated by age-related trends in maternal immunity, adaptive immunity, and exposure rate, and specific host-parasite system

characteristics including maladaptation and immuno-pathogenicity [22, 23]. Additionally, studies looking at age-related virulence or case mortality do not exclusively look at differences between adult and juvenile stages and may focus on old age-mediated declines in immuno-competence. However, despite these issues, we found data on several empirical systems that lend support to our predictions and may offer opportunities for testing our hypotheses Fig 1. In particular, Wanelik *et al.* [54] showed that Great Island Virus (GIV) transmission in Guillemots (*Uria aalge*) is assortative across age classes because of the spatial structure of breeding grounds. GIV is transmitted by poorly motile ticks and pre-breeder stages of Guillemots do not enter breeding areas of the colony. As a consequence, the virus does not readily transmit between guillemot age-stages [54]. Previous work on guillemot life history shows that the birds spend more than three quarters of their life-span as mature breeders [52], so the combination of assortative transmission and fast maturation predicts that GIV should be more virulent in breeders. In line with the predictions of our model, infection associated mortality risk is 1.45 times higher for adults than for juveniles [53]. In contrast, [45] showed that salmon louse caused morality in juvenile pink salmon (*Oncorhynchus gorbuscha*), but had no effect on mortality risk for adults. Salmon louse is also assortatively transmitted between age classes, because pink salmon have strict two-year lifespans where they are only ever associated with individuals of their same age class [44]. The salmon only reproduce once at the very end of their lives (semelparity), and therefore have a short adult period by our model. This short post-maturation stage and assortative transmission predicts the higher salmon louse virulence in juveniles.

Better data on mixing matrices for more disease systems could provide interesting insights into the maintenance of either high juvenile or high adult virulence. One system where these insights could prove especially important is in Bd (*Batrachochytrium dendrobatidis*, or chytrid fungus) infection in frogs, which has been causing catastrophic worldwide declines in frog populations [31]. Bd infection has been shown to have different virulence effects in the different frog life-stages [26, 85] and these effects also vary by frog species [86, 87]. Recent work has shown that adult virulence in several frog populations has not decreased even after 20 years of Bd presence [88]. Already, frog demography has been implicated as an important factor for population persistence in the face of Bd with frog species where adults move away from breeding waters being more resistant to population declines [89, 90], but habitats with multi-year larvae have more severe epidemics because the older stages maintain high levels of infection that then spill over to infect other stages and species [26, 85]. Changes in the assortiveness of mixing clearly has important implications for disease transmission across stages, and our model suggests that it could also have implications for the maintenance of high virulence in different age stages.

While data on age-related contact patterns are difficult to access for wildlife populations, a wealth of mixing data exists for humans [30, 91]. These suggest that contacts relevant for the transmission of directly transmitted pathogens are highly assortative by age. While the evolutionary drivers of human pathogens is sometimes complicated, we posit that chickenpox (varicella virus) virulence in humans proves an intriguing case study. Given that humans have a long juvenile period in the context of our model, even when we only consider pre-reproductive and reproductive periods [92], the higher virulence in adults of Chickenpox (23-29 times higher mortality risk in adults [93]) fits the predictions of our model. This higher mortality risk corresponds to increased viral titers with age [94] and, perhaps most interestingly, while varicella virus infects many cell-types, T cell infection is thought to be important for transport and pathogenesis [95]. Therefore, age-related trends in T-cell abundance could be implicated in chickenpox pathogenesis, although this relationship is complicated by the fact that VSV-specific T cell responses are also correlated with decreased viral titer and diminish with age [94, 96, 97]. Still, this example points towards one mechanism that may underlie the mediation of age-specific virulence in pathogens.

Our models have implications for disease management especially in farmed and other managed animal populations. For instance, if the post-maturation span is short (i.e. if $u$ is small), then artificial restriction of the contacts between stages is predicted to select for higher virulence and population crashes in juveniles. However, if the post-maturation span is long, restricting the contacts into juvenile-juvenile and adult-adult (by e.g., separating the cohorts) can lead to the parasite extinction as a result of overexploitation against the juveniles. These contrasting outcomes can occur for any given host species, depending on how management modulates host stage-structure. Our models thus predict that, to prevent evolutionary changes towards higher virulence, one needs to carefully take into account the host stage and contact structure.

For simplicity and tractability we chose to use simple two-stage models rather than a more continuous "infection-age" models (which would entail the formalism based on partial differential equations and dynamic programming approach). Future studies that capture more continuous age structure are an important next step. Also, although we assumed that parasites can express conditional virulence depending on the stage of the hosts they infect with, more data are needed to test this idea. In addition, coevolutionary models are likely to give further important insights to the determinants of age-dependent disease interactions in nature. Our approach offers the basis for modeling these coevolutionary dynamics between hosts and parasites when there is stage structure.

## Declaration

*Acknowledgement*

We thank Troy Day and Jessica Hite for the helpful comments.

*Funding*

R.I. and M.B. are supported by Natural Environment Research Council (NERC; NE/K014617/1); E.V. is supported by the National Science Foundation Graduate Research Fellowship Program (Grant Number DGE 1752814).

*Author contribution*

R.I and M.B. designed research; R.I. and E.V. performed research; R.I., E.V., and M.B. wrote the paper.

## References

[1]  Smith, T. "Some problems in the life history of pathogenic microorganisms". *Science* **20**.520 (1904), pp. 817–832.

[2]  Ball, G. H. "Parasitism and evolution". *The American Naturalist* **77**.771 (1943), pp. 345–364.

[3]  Anderson, R. M. & May, R. "Coevolution of hosts and parasites". *Parasitology* **85**.2 (1982), pp. 411–426.

[4]  Read, A. F. "The evolution of virulence". *Trends in Microbiology* **2**.3 (1994), pp. 73–76.

[5]  Ebert, D. & Herre, E. A. "The evolution of parasitic diseases". *Parasitology Today* **12**.3 (1996), pp. 96–101.

[6]  Frank, S. A. "Models of parasite virulence". *Quarterly Review of Biology* (1996), pp. 37–78.

[7]  Mideo, N., Alizon, S., & Day, T. "Linking within-and between-host dynamics in the evolutionary epidemiology of infectious diseases". *Trends in Ecology & Evolution* **23**.9 (2008), pp. 511–517.

[8]  Alizon, S., Hurford, A., Mideo, N., & Van Baalen, M. "Virulence evolution and the trade-off hypothesis: history, current state of affairs and the future". *Journal of Evolutionary Biology* **22**.2 (2009), pp. 245–259.

[9]  Schmid-Hempel, P. *Evolutionary parasitology: the integrated study of infections, immunology, ecology, and genetics*. Oxford University Press, New York, 2011.

[10]  Bull, J. J. & Lauring, A. S. "Theory and empiricism in virulence evolution". *PLoS Pathogens* **10**.10 (2014), e1004387.

[11]  Cressler, C. E., McLeod, D. V., Rozins, C., Hoogen, J. van den, & Day, T. "The adaptive evolution of virulence: a review of theoretical predictions and empirical tests". *Parasitology* **143**.7 (2016), pp. 915–30.

[12] Boots, M. & Sasaki, A. "Parasite evolution and extinctions". *Ecology Letters* **6**.3 (2003), pp. 176–182.

[13] De Castro, F. & Bolker, B. "Mechanisms of disease-induced extinction". *Ecology Letters* **8**.1 (2005), pp. 117–126.

[14] Dieckmann, U. *Adaptive dynamics of infectious diseases: in pursuit of virulence management*. **2**. Cambridge University Press, 2005.

[15] Boots, M., Best, A., Miller, M. R., & White, A. "The role of ecological feedbacks in the evolution of host defence: what does theory tell us?" *Philosophical Transactions of the Royal Society B: Biological Sciences* **364**.1513 (Jan. 2009), p. 27.

[16] Ewald, P. W. "Host-parasite relations, vectors, and the evolution of disease severity". *Annual Review of Ecology and Systematics* **14**.1 (1983), pp. 465–485.

[17] Charnov, E. L. "Optimal foraging, the marginal value theorem". *Theoretical Population Biology* **9**.2 (1976), pp. 129–136.

[18] Bulmer, M. G. *Evolutionary theoretical ecology*. Sinauer Associates, Sunderland, MA, 1994.

[19] Otto, S. P. & Day, T. *A biologist's guide to mathematical modeling in ecology and evolution*. **13**. Princeton University Press, 2007.

[20] Lion, S. & Metz, J. A. J. "Beyond $R_0$ maximisation: on pathogen evolution and environmental dimensions". *Trends in Ecology & Evolution* **33** (2018), pp. 75–90.

[21] Van Baalen, M. & Sabelis, M. W. "The scope for virulence management: a comment on Ewald's view on the evolution of virulence". *Trends in microbiology* **3**.11 (1995), pp. 414–416.

[22] Hudson, P. J. & Dobson, A. P. "Macroparasites: observed patterns". *Ecology of Infectious Diseases in Natural Populations*. Ed. by Grenfell, B. T. & Dobson, A. P. Cambridge University Press, 1995, pp. 144–176.

[23] Wilson, K., Bjørnstad, O., Dobson, A., Merler, S., Poglayen, G, Randolph, S., Read, A., & Skorping, A. "Heterogeneities in macroparasite infections: patterns and processes". *The Ecology of Wildlife Diseases* **44** (2002), pp. 6–44.

[24] Dwyer, G. "The roles of density, stage, and patchiness in the transmission of an insect virus". *Ecology* **72**.2 (1991), pp. 559–574.

[25] Fleming-Davies, A. E., Dukic, V., Andreasen, V., & Dwyer, G. "Effects of host heterogeneity on pathogen diversity and evolution". *Ecology Letters* **18**.11 (2015), pp. 1252–1261.

[26] Hite, J. L., Penczykowski, R. M., Shocket, M. S., Strauss, A. T., Orlando, P. A., Duffy, M. A., Cáceres, C. E., & Hall, S. R. "Parasites destabilize host populations by shifting stage-structured interactions". *Ecology* **97**.2 (2016), pp. 439–449.

[27] Craft, M. E. "Infectious disease transmission and contact networks in wildlife and livestock". *Philosophical Transactions of the Royal Society of London B: Biological Sciences* **370**.1669 (2015), p. 20140107.

[28] VanderWaal, K. L. & Ezenwa, V. O. "Heterogeneity in pathogen transmission: mechanisms and methodology". *Functional Ecology* **30**.10 (2016). Ed. by Hawley, D., pp. 1606–1622.

[29] White, L. A., Forester, J. D., & Craft, M. E. "Using contact networks to explore mechanisms of parasite transmission in wildlife". *Biological Reviews* **92**.1 (2017), pp. 389–409.

[30] Rohani, P., Zhong, X., & King, A. A. "Contact network structure explains the changing epidemiology of pertussis". *Science* **330**.6006 (2010), pp. 982–985.

[31] Kilpatrick, A. M., Briggs, C. J., & Daszak, P. "The ecology and impact of chytridiomycosis: an emerging disease of amphibians". *Trends in Ecology & Evolution* **25**.2 (2010), pp. 109–118.

[32] Briggs, C. & Godfray, H. "The dynamics of insect-pathogen interactions in stage-structured populations". *The American Naturalist* **145**.6 (1995), pp. 855–887.

[33] Hofbauer, J. & Sigmund, K. "Adaptive dynamics and evolutionary stability". *Applied Mathematics Letters* **3**.4 (1990), pp. 75–79.

[34] Dieckmann, U. & Law, R. "The dynamical theory of coevolution: a derivation from stochastic ecological processes". *Journal of Mathematical Biology* **34**.5-6 (1996), pp. 579–612.

[35] Fisher, R. A. *The genetical theory of natural selection.* Tech. rep. Dover Publications, 1958.

[36] Taylor, P. D. "Allele-frequency change in a class-structured population". *The American Naturalist* **135** (1990), pp. 95–106.

[37] Frank, S. A. *Foundations of social evolution*. Princeton University Press, 1998.

[38] Caswell, H. *Matrix population models*. Wiley Online Library, 2001.

[39] Gandon, S. "Evolution of multihost parasites". *Evolution* **58**.3 (2004), pp. 455–469.

[40] Grafen, A. "A theory of Fisher's reproductive value". *Journal of Mathematical Biology* **53**.1 (2006), pp. 15–60.

[41] Williams, P. D. "New insights into virulence evolution in multigroup hosts". *The American Naturalist* **179**.2 (2011), pp. 228–239.

[42] Williams, P. D. & Kamel, S. J. "The evolution of pathogen virulence: Effects of transitions between host types". *Journal of Theoretical Biology* **438** (2018), pp. 1–8.

[43] Lion, S. "Class structure, demography, and selection: reproductive-value weighting in nonequilibrium, polymorphic populations". *The American Naturalist* (2018), pp. 000–000.

[44] Heard, W. R. "Life history of pink salmon (Oncorhynchus gorbuscha)". *Pacific salmon life histories* (1991), pp. 119–230.

[45] Jones, S, Kim, E, & Bennett, W. "Early development of resistance to the salmon louse, Lepeophtheirus salmonis (Krøyer), in juvenile pink salmon, Oncorhynchus gorbuscha (Walbaum)". *Journal of fish diseases* **31**.8 (2008), pp. 591–600.

[46] Wassif, K. & Soliman, S. "Population studies on gerbils of the western desert of Egypt, with special reference to Gerbillus andersoni de Winton". *Proceedings of the 9th Vertebrate Pest Conference (1980)*. 1980, p. 41.

[47] Hawlena, H., Abramsky, Z., & Krasnov, B. R. "Ectoparasites and age-dependent survival in a desert rodent". *Oecologia* **148**.1 (2006), pp. 30–39.

[48] Holst, D. von, Hutzelmeyer, H., Kaetzke, P., Khaschei, M., Rödel, H. G., & Schrutka, H. "Social rank, fecundity and lifetime reproductive success in wild European rabbits (*Oryctolagus cuniculus*)". *Behavioral Ecology and Sociobiology* **51**.3 (2002), pp. 245–254.

[49] Cornell, S. J., Bjornstad, O. N., Cattadori, I. M., Boag, B., & Hudson, P. J. "Seasonality, cohort-dependence and the development of immunity in a natural host–nematode system". *Proceedings of the Royal Society of London B: Biological Sciences* **275**.1634 (2008), pp. 511–518.

[50] Morisse, J., Le, G. G., & Boilletot, E. "Hepatitis of viral origin in Leporidae: introduction and aetiological hypotheses." *Revue scientifique et technique (International Office of Epizootics)* **10**.2 (1991), pp. 269–310.

[51] Reluga, T. C., Medlock, J., Poolman, E., & Galvani, A. P. "Optimal timing of disease transmission in an age-structured population". *Bulletin of Mathematical Biology* **69**.8 (2007), pp. 2711–2722.

[52] Harris, M. & Wanless, S. "Survival and non-breeding of adult Common Guillemots Uria aalge". *Ibis* **137**.2 (1995), pp. 192–197.

[53] Nunn, M., Barton, T., Wanless, S, Hails, R., Harris, M., & Nuttall, P. "Tick-borne Great Island Virus:(II) Impact of age-related acquired immunity on transmission in a natural seabird host". *Parasitology* **132**.2 (2006), pp. 241–253.

[54] Wanelik, K. M., Burthe, S. J., Harris, M. P., Nunn, M. A., Godfray, H. C. J., Sheldon, B. C., McLean, A. R., & Wanless, S. "Investigating the effects of age-related spatial structuring on the transmission of a tick-borne virus in a colonially breeding host". *Ecology and Evolution* **7**.24 (2017), pp. 10930–10940.

[55] Sukumar, R, Krishnamurthy, V, Wemmer, C., & Rodden, M. "Demography of captive Asian elephants (*Elephas maximus*) in southern India". *Zoo Biology: Published in affiliation with the American Zoo and Aquarium Association* **16**.3 (1997), pp. 263–272.

[56] Lynsdale, C. L., Mumby, H. S., Hayward, A. D., Mar, K. U., & Lummaa, V. "Parasite-associated mortality in a long-lived mammal: Variation with host age, sex, and reproduction". *Ecology and Evolution* **7**.24 (2017), pp. 10904–10915.

[57] Lack, D. L. "Ecological adaptations for breeding in birds" (1968).

[58] Sol, D., Jovani, R., & Torres, J. "Parasite mediated mortality and host immune response explain age-related differences in blood parasitism in birds". *Oecologia* **135**.4 (2003), pp. 542–547.

[59] Schreiber, S. & Rudolf, V. H. "Crossing habitat boundaries: coupling dynamics of ecosystems through complex life cycles". *Ecology Letters* **11**.6 (2008), pp. 576–587.

[60] Roos, A. M. de & Persson, L. *Population and community ecology of ontogenetic development*. Princeton University Press, 2013.

[61] Day, T., Alizon, S., & Mideo, N. "Bridging scales in the evolution of infectious disease life histories: theory". *Evolution* **65**.12 (2011), pp. 3448–3461.

[62] Mideo, N., Nelson, W. A., Reece, S. E., Bell, A. S., Read, A. F., & Day, T. "Bridging scales in the evolution of infectious disease life histories: application". *Evolution* **65**.11 (2011), pp. 3298–3310.

[63] McCallum, H., Barlow, N., & Hone, J. "How should pathogen transmission be modelled?" *Trends in Ecology & Evolution* **16**.6 (2001), pp. 295–300.

[64] Glasser, J., Feng, Z., Moylan, A., Del Valle, S., & Castillo-Chavez, C. "Mixing in age-structured population models of infectious diseases". *Mathematical Biosciences* **235**.1 (2012), pp. 1–7.

[65] Massol, F. & Cheptou, P.-O. "Evolutionary syndromes linking dispersal and mating system: the effect of autocorrelation in pollination conditions". *Evolution* **65**.2 (2011), pp. 591–598.

[66] Rodrigues, A. M. M. & Gardner, A. "Evolution of helping and harming in heterogeneous populations". *Evolution* **66**.7 (2012), pp. 2065–2079.

[67] Massol, F. & Débarre, F. "Evolution of dispersal in spatially and temporally variable environments: The importance of life cycles". *Evolution* **69**.7 (2015), pp. 1925–1937.

[68] Iritani, R. & Cheptou, P.-O. "Joint evolution of differential seed dispersal and self-fertilization". *Journal of Evolutionary Biology* **30**.8 (2017), pp. 1526–1543.

[69] Driessche, P. Van den & Watmough, J. "Reproduction numbers and sub-threshold endemic equilibria for compartmental models of disease transmission". *Mathematical Biosciences* **180**.1 (2002), pp. 29–48.

[70] Hurford, A., Cownden, D., & Day, T. "Next-generation tools for evolutionary invasion analyses". *Journal of The Royal Society Interface* **7**.45 (2010), pp. 561–571.

[71] Camino Beck, T. de & Lewis, M. A. "A new method for calculating net reproductive rate from graph reduction with applications to the control of invasive species". *Bulletin of Mathematical Biology* **69**.4 (2007), pp. 1341–1354.

[72] Camino Beck, T. de & Lewis, M. A. "On Net Reproductive Rate and the Timing of Reproductive Output". *The American Naturalist* **172**.1 (2008), pp. 128–139.

[73] Camino Beck, T. de, Lewis, M. A., & Driessche, P. van den. "A graph-theoretic method for the basic reproduction number in continuous time epidemiological models". *Journal of Mathematical Biology* **59**.4 (2008), pp. 503–516.

[74] Best, A., White, A., & Boots, M. "The coevolutionary implications of host tolerance". *Evolution* **68**.5 (2014), pp. 1426–1435.

[75] Takada, T. & Kigami, J. "The dynamical attainability of ESS in evolutionary games". *Journal of Mathematical Biology* **29**.6 (1991), pp. 513–529.

[76] Christiansen, F. B. "On conditions for evolutionary stability for a continuously varying character". *The American Naturalist* (1991), pp. 37–50.

[77] Maynard Smith, J. & Price, G. R. "The Logic of Animal Conflict". *Nature* **246** (1973), p. 15.

[78] Eshel, I. "Evolutionary and continuous stability". *Journal of Theoretical Biology* **103**.1 (1983), pp. 99–111.

[79] Day, T. "Parasite transmission modes and the evolution of virulence". *Evolution* **55**.12 (2001), pp. 2389–2400.

[80] Gandon, S., Jansen, V. A., & Van Baalen, M. "Host life history and the evolution of parasite virulence". *Evolution* **55**.5 (2001), pp. 1056–1062.

[81] Day, T. & Proulx, S. R. "A general theory for the evolutionary dynamics of virulence". *The American Naturalist* **163**.4 (2004), E40–E63.

[82] Osnas, E. E. & Dobson, A. P. "Evolution of virulence in heterogeneous host communities under multiple trade-offs". *Evolution* **66**.2 (2011), pp. 391–401.

[83] Williams, P. D., Day, T., Fletcher, Q., & Rowe, L. "The shaping of senescence in the wild". *Trends in Ecology & Evolution* **21**.8 (2006), pp. 458–463.

[84] Williams, P. D., Dobson, A. P., Dhondt, K. V., Hawley, D. M., & Dhondt, A. A. "Evidence of trade-offs shaping virulence evolution in an emerging wildlife pathogen". *Journal of Evolutionary Biology* **27**.6 (2014), pp. 1271–1278.

[85] Medina, D., Garner, T. W., Carrascal, L. M., & Bosch, J. "Delayed metamorphosis of amphibian larvae facilitates Batrachochytrium dendrobatidis transmission and persistence". *Diseases of aquatic organisms* **117**.2 (2015), pp. 85–92.

[86] Berger, L., Speare, R., Daszak, P., Green, D. E., Cunningham, A. A., Goggin, C. L., Slocombe, R., Ragan, M. A., Hyatt, A. D., McDonald, K. R., *et al.* "Chytridiomycosis causes amphibian mortality associated with population declines in the rain forests of Australia and Central America". *Proceedings of the National Academy of Sciences* **95**.15 (1998), pp. 9031–9036.

[87] Blaustein, A. R., Romansic, J. M., Scheessele, E. A., Han, B. A., Pessier, A. P., & Longcore, J. E. "Interspecific variation in susceptibility of frog tadpoles to the pathogenic fungus Batrachochytrium dendrobatidis". *Conservation Biology* **19**.5 (2005), pp. 1460–1468.

[88] Voyles, J., Woodhams, D. C., Saenz, V., Byrne, A. Q., Perez, R., Rios-Sotelo, G., Ryan, M. J., Bletz, M. C., Sobell, F. A., McLetchie, S., *et al.* "Shifts in disease dynamics in a tropical amphibian assemblage are not due to pathogen attenuation". *Science* **359**.6383 (2018), pp. 1517–1519.

[89] Lips, K. R., Brem, F., Brenes, R., Reeve, J. D., Alford, R. A., Voyles, J., Carey, C., Livo, L., Pessier, A. P., & Collins, J. P. "Emerging infectious disease and the loss of biodiversity in a Neotropical amphibian community". *Proceedings of the national academy of sciences of the United States of America* **103**.9 (2006), pp. 3165–3170.

[90] McCaffery, R., Richards-Zawacki, C. L., & Lips, K. R. "The demography of Atelopus decline: Harlequin frog survival and abundance in central Panama prior to and during a disease outbreak". *Global Ecology and Conservation* **4** (2015), pp. 232–242.

[91] Mossong, J., Hens, N., Jit, M., Beutels, P., Auranen, K., Mikolajczyk, R., Massari, M., Salmaso, S., Tomba, G. S., Wallinga, J., *et al.* "Social contacts and mixing patterns relevant to the spread of infectious diseases". *PLoS Medicine* **5**.3 (2008), e74.

[92] Bogin, B. & Smith, B. H. "Evolution of the human life cycle". *American Journal of Human Biology* **8**.6 (1996), pp. 703–716.

[93] Heininger, U. & Seward, J. F. "Varicella". *The Lancet* **368**.9544 (2006), pp. 1365–1376.

[94] Malavige, G. N., Jones, L., Kamaladasa, S., Wijewickrama, A, Seneviratne, S., Black, A. P., & Ogg, G. S. "Viral load, clinical disease severity and cellular immune responses in primary varicella zoster virus infection in Sri Lanka". *PloS one* **3**.11 (2008), e3789.

[95] Zerboni, L. & Arvin, A. M. "The pathogenesis of varicella-zoster virus neurotropism and infection". *Neurotropic viral infections*. Springer, 2016, pp. 135–173.

[96] Erkeller-Yuksel, F. M., Deneys, V, Yuksel, B, Hannet, I, Hulstaert, F, Hamilton, C, Mackinnon, H, Stokes, L. T., Munhyeshuli, V. a., Vanlangendonck, F, *et al.* "Age-related changes in human blood lymphocyte subpopulations". *The Journal of Pediatrics* **120**.2 (1992), pp. 216–222.

[97] Nader, S., Bergen, R., Sharp, M., & Arvin, A. M. "Age-related differences in cell-mediated immunity to varicella-zoster virus among children and adults immunized with live attenuated varicella vaccine". *Journal of Infectious Diseases* **171**.1 (1995), pp. 13–17.