Title

RNA-antimicrobial peptide LL-37 complexes fuel a TLR- and NET-

mediated inflammatory cycle of neutrophil activation in psoriasis

Running title

RNA and LL37 activate neutrophils in psoriasis

Authors

Franziska Herster[1], Zsofia Bittner[1], Sabine Dickhöfer[1], David Eisel[1,$], Tatjana Eigenbrod[2], Thomas

Knorpp[3], Nicole Schneiderhan-Marra[3], Markus W. Löffler[1,4], Hubert Kalbacher[5], Dominik Hartl[6], Lukas

Freund[7], Knut Schäkel[7], Martin Heister[8], Kamran Ghoreschi[8,9], Alexander N.R. Weber[1,]*

Affiliations

[1] Department of Immunology, University of Tübingen, Auf der Morgenstelle 15, 72076 Tübingen,

Germany

[2] Department of Infectious Diseases, Medical Microbiology and Hygiene, University Hospital

Heidelberg, Im Neuenheimer Feld 324, 69120 Heidelberg, Germany

[3] NMI Natural and Medical Sciences Institute at the University of Tübingen, Markwiesenstr. 55, 72770

Reutlingen, Germany

[4] Department of General, Visceral and Transplant Surgery, University Hospital Tübingen, Hoppe-

Seyler-Str. 3, 72076 Tübingen, Germany

[5] Interfaculty Institute of Biochemistry, University of Tübingen, Hoppe-Seyler-Str. 4, 72076 Tübingen,

Germany

[6] University Children's Hospital and Interdisciplinary Center for Infectious Diseases, University of

Tübingen, Hoppe-Seyler-Str. 1, 72076 Tübingen, Germany

[7] Department of Dermatology, University Hospital Heidelberg, Im Neuenheimer Feld 440, 69120

Heidelberg, Germany

[8] Department of Dermatology, University Hospital Tübingen, Liebermeisterstr. 25, 72076 Tübingen,

Germany

[9] Department of Dermatology, Charité – Universitätsmedizin Berlin, Charitéplatz 1, 10117 Berlin,

Germany

* Contact information (Corresponding Author)

Alexander N. R. Weber, Interfaculty Institute for Cell Biology, Department of Immunology, University

of Tübingen, Auf der Morgenstelle 15, 72076 Tübingen, Germany. Tel.: +49 7071 29 87623. Fax: +49

7071 29 4579. Email: alexander.weber@uni-tuebingen.de

[$] Change of address/additional addresses

David Eisel, GMP & T Cell Therapy Unit (G183), German Cancer Research Center, Im Neuenheimer

Feld 280, 69120 Heidelberg, Germany

Dominik Hartl, Galapagos Pharma, Basel, Switzerland

Keywords

Antimicrobial peptide LL-37, RNA, psoriasis, neutrophil, Toll-like receptor (TLR), neutrophil extracellular trap (NET)

Summary

Human and bacterial RNA in complex with LL37 activates neutrophils via TLR8 to release cytokines, chemokines and neutrophil extracellular traps (NETs). NETs and neutrophil-rich areas in psoriatic skin contain RNA and LL37, suggesting RNA-LL37 may fuel a PMN-mediated and self-sustaining inflammatory cycle in psoriasis.

Abstract

Psoriasis is an inflammatory autoimmune disease characterized by skin lesions showing strong neutrophil (PMN) infiltration and high levels of the antimicrobial peptide, LL37, but the role of PMNs in this context remains unclear. We here show that primary human PMNs, especially PMNs from psoriasis patients, not only respond via TLR8 to human and bacterial RNA in complexed with LL37 by cytokine-, chemokine- and neutrophil extracellular trap (NET)-release; they also actively release additional RNA and LL37 in response to stimulation by the same complex and both RNA and LL37 were found to be highly abundant in psoriatic skin. Moreover, RNA-LL37-induced NETs propagated PMN activation and could thus fuel a PMN-mediated and self-sustaining inflammatory loop that may represent an unexpected early initiator or amplifying event in psoriasis. Given that TLR inhibitory oligodeoxynucleotides prevented the cytokine production and NETosis of PMNs by RNA-LL37 complexes *in vitro*, our study also highlights TLR blockade as a potential therapeutic intervention strategy in psoriasis.

Introduction

Psoriasis is an autoimmune disease of the skin with high incidence in Western countries (1.5-3%),
causing high socioeconomic and disease burden with limited but increasing treatment options
(Eberle et al., 2016; Griffiths and Barker, 2007). The most common form of psoriasis, plaque psoriasis,
is characterized by epidermal hyperplasia due to keratinocyte (KC) hyper-proliferation, increased
endothelial proliferation and an infiltrate of leukocytes, such as dendritic cells, T cells and,
prominently, polymorphonuclear neutrophils (PMNs)(Griffiths and Barker, 2007). The high and early
accumulation of PMNs in psoriatic plaques and micro-abscesses is well documented, as well as an
increase of PMNs in the circulation of psoriasis patients (Sen et al., 2013) (Schon et al., 2017).
However, it is unclear whether PMN infiltration represents a bystander phenomenon or has a causal
role in the development of the disease.

PMNs are known as a source of LL37, an amphipathic, positively-charged 37 amino acid peptide
generated from a precursor protein, the cathelicidin hCAP18, and highly abundant in psoriatic lesions
(Griffiths and Barker, 2007; Morizane and Gallo, 2012). LL37 is stored in the secondary granules of
PMNs, from which it can be released upon activation (Sorensen et al., 1997). Apart from acting as a
self-antigen for CD4 helper T cells in psoriasis patients (Lande et al., 2014), LL37 was previously
shown to form complexes with DNA or RNA that resisted nuclease degradation, were readily taken
up by plasmacytoid dendritic cells (pDCs), and triggered high Interferon (IFN) α and low Tumor
necrosis factor (TNF) or Interleukin (IL) 6 secretion from these cells (Ganguly et al., 2009; Lande et al.,
2007). Although type I IFNs contribute to psoriasis (Afshar et al., 2013; Yao et al., 2008), chemokines
and pro-inflammatory cytokines such as TNF, IL-1β or IL-23 are additionally required for immune cell
infiltration and T cell polarization, respectively (Ghoreschi et al., 2007). Since pDCs are comparatively
low producers of these psoriasis-initiating inflammatory cytokines and do not accumulate to high
numbers in psoriatic skin, this raises the question whether nucleic acid-LL37 sensing by other
immune cells, particularly PMNs, may contribute more prominently to initial inflammatory
cytokine/chemokine milieu observed in psoriasis.

In contrast to pDCs, PMNs are known producers of considerable amounts of various cytokines and
chemokines (Kruger et al., 2015; Tecchio et al., 2014; Terui et al., 2000). Additionally, PMNs are able
to undergo neutrophil extracellular trap (NET) formation (NETosis), an activation-induced process
leading to the extrusion of nuclear DNA (Brinkmann et al., 2004), as well as cellular proteins,
including LL37 (Radic and Marion, 2013). Like pDCs, PMNs express multiple pattern recognition
receptors (PRRs) of the Toll-like receptor (TLR) family, namely TLR2, 4, 8 (Berger et al., 2012) and 9
(Lindau et al., 2013), but not TLR3 or TLR7 (Berger et al., 2012). TLRs directly bind activating ligands,
typically so-called microbe-associated molecular patterns (MAMPs) absent from the host (Kawasaki

and Kawai, 2014); however, when they can gain access to endosomal compartments - for example by
complex formation with LL37 - host nucleic acids are able to trigger the endosomal TLRs 3, 7, 8
(sensing double-stranded or single-stranded RNA, respectively) or TLR9 (sensing DNA). TLR7
expression in humans is restricted to pDC and (only inducibly) to B cells. TLR8 thus appears to be the
primary receptor for foreign (and potentially host-derived) single-stranded RNA in humans
(Eigenbrod et al., 2015).

PMNs theoretically combine the abilities (i) to release LL37 and nucleic acids as components of
immuno-stimulatory ligands via NETosis; (ii) to potentially sense such ligands via TLRs; and (iii) to
secrete chemo-/cytokines. These mechanisms were never shown to act in concert but we speculated
whether LL37-mediated DNA or RNA sensing via TLRs in PMNs might initiate and fuel inflammatory
chemo-/cytokine production and thus inflammation and immune cell infiltration in psoriatic skin. We
here present experimental evidence that human primary PMNs readily respond to RNA-LL37
complexes, but not RNA alone, leading to the release of a broad array of chemokines and cytokines à
and can be therapeutically blocked by inhibitory oligodeoxynucleotides (iODNs). Unexpectedly, RNA-
LL37 also triggered the release of NETs, which contained additional LL37, RNA and DNA and could
activate further PMN, thereby providing the basis for a self-propagating inflammatory cycle.

Results

LL37 promotes RNA uptake and induces TLR8-mediated PMN activation

Previous results indicated that human primary PMNs can respond to RNA and DNA when stimulated for >12 h, albeit at much lower levels than when stimulated with the nucleoside analog TLR7/8 agonist, R848 (Janke et al., 2009; Lindau et al., 2013). We sought to re-evaluate these findings using highly purified primary PMNs assayed within a short time period (4 h) that excludes secondary release effects, e.g. by apoptosis (Fig. S1A). The TLR7/8 agonist R848, like the TLR4 agonist, LPS, elicited robust IL-8 release but only LPS triggered CD62L shedding; phospho-thioate (PTO) synthetic CpG ODN, a typical TLR9 agonist, also strongly activated IL-8 release and CD62L shedding (Fig. 1A, Table S1). However, unmodified, natural phospho-diester DNA ODN or human genomic DNA elicited neither IL-8 release nor CD62L shedding, irrespective of whether they were complexed with LL37 (Fig. 1B). In contrast, in pDC LL37 binding of natural DNA triggered potent TLR9 responses (Lande et al., 2007). In the absence of LL37, single-stranded synthetic RNA40 (henceforward referred to as 'RNA') barely caused IL-8 release when applied at equimolar concentrations with R848 (Fig. 1C). This suggests that on their own neither RNA nor DNA are able to trigger primary PMN responses. This may be due to the endosomal localization of TLR8 and 9, which R848 and CpG can apparently access, whereas RNA and normal DNA cannot (Kuznik et al., 2011). However, robust IL-8 release and moderate CD62L shedding were observed when RNA was complexed with LL37 (Fig. 1C and *cf.* Fig. 2E). To check whether PMNs could engage RNA-LL37 complexes, live PMNs were seeded and RNA-LL37 complexes added for 20 min before fixing the cells for electron microscopy analysis. As shown in Fig. 1D, PMNs were found in proximity to fiber-like structures corresponding to RNA-LL37 complexes (Schmidt et al., 2015). Flow cytometry using AlexaFluor (AF) 647-labeled RNA showed that LL37 also promoted the uptake of complexed RNA: >20% of PMNs incubated with labeled RNA in the presence of LL37 were AF647-positive within 60 min, compared to 5% in the absence of LL37 (Fig. 1E). ImageStream bright-field cytometry confirmed that this was not due to external/surface binding of labeled RNA but rather internalization (Fig. 1F, G). Bright-field fluorescence microscopy with Atto488-labeled LL37 showed co-localization of AF647-RNA and Atto488-LL37 in intracellular compartments (Fig. 1H) where endosomal TLRs are expressed (Berger et al., 2012). To pharmacologically explore whether an endosomal TLR might be involved, we investigated the effect of chloroquine, a well-known inhibitor of endosomal TLRs that was non-toxic for PMNs (Fig. S1B). Indeed IL-8 release from primary PMNs in response to RNA-LL37 and CpG ODN (positive control) was significantly reduced by chloroquine addition (Fig. 1I and S1C, respectively). As chloroquine does not affect cytoplasmic RNA sensors (e.g. RIG-I) (Matsukura et al., 2007), this implicates an endosomal TLR in RNA-sensing. We next sought to check whether the observed effect on cytokine release extended to normal RNA from

mammalian cells which may be more physiologically relevant in the context of psoriasis. As shown in
Fig. 1J, human mRNA significantly induced IL-8 release, but only in combination with LL37. Since
microbiome analyses have recently revealed alterations in staphylo- and streptococcal skin
colonization in psoriasis patients (Alekseyenko et al., 2013) and bacterial RNA is an emerging
immuno-stimulatory pattern (Eigenbrod and Dalpke, 2015), we also tested whether LL37 promoted *S.*
aureus, i.e. bacterial RNA (bRNA) activation of PMNs. Fig. 1K shows that bRNA significantly
stimulated IL-8 release similarly to RNA40 when complexed with LL37. Taken together this suggests
that uptake of RNA is promoted by LL37 and directs RNA to intracellular compartments from which
TLR sensing and activation leading to cytokine release can occur – both in response to mammalian
and bacterial/foreign RNA.

RNA-LL37 complexes prompt release of multiple pro-inflammatory cytokines and chemokines,
especially in psoriasis PMNs.
To test whether RNA-LL37 also promoted the release of cytokines other than IL-8, we screened
supernatants of stimulated PMNs by Luminex multiplex analysis and detected TNF, IL-1β, IL-6, and –
unexpectedly – IL-16 and MIP-1β primarily or exclusively in RNA-LL37-stimulated cell supernatants,
respectively (Fig. S2A,B). Despite variation between donors, this was confirmed by cytometric bead
arrays analyses (Fig. 2A-C) and ELISA (Fig. 2D, E) in samples from more donors. To our knowledge,
active receptor-mediated release of the chemoattractants, IL-16 (also known as Lymphocyte
Chemoattractant Factor, LCF) and MIP-1β (also known as Chemokine (C-C motif) Ligand, CCL4) from
primary human PMNs has so far not been reported. In fact, release of IL-16 was also boosted by LL37
alone (Fig. 2D) and MIP-1β was strongly released from PMNs in response to RNA-LL37 complexes but
not RNA alone (Fig. 2E).

Since IL-16 and MIP-1β are known chemoattractants for CD4 T cells and a variety of other immune
cells (Cruikshank and Little, 2008; Menten et al., 2002; Roth et al., 2016), we checked whether IL-16
or MIP-1β, at the concentrations observed in this study, could influence the migration and infiltration
of additional immune cells, which is a hallmark of psoriatic plaques (Schon et al., 2017). Transwell
experiments with peripheral blood mononuclear cells (PBMCs) from healthy human donors (upper
well) and IL-16 (300 and 1500 pg/ml), MIP-1β (30 and 150 pg/ml) or SDF-1α as a control (lower well,
Fig. S2C-E) showed that the lowest concentration of both cytokines induced a donor-dependent
increase in the number of CD3+CD4+ helper, CD3+CD8+ cytotoxic T cells as well as CD14+HLA-DR+
monocytes (Figs. 2F-H). Possibly due to a non-linear dose response relationship observed for some
cytokines (Atanasova and Whitty, 2012), higher concentrations had a weaker effect in some donors.
Owing to the donor-to-donor variation generally observed throughout in the human system, only a
non-significant, moderate effect was observed. We separately tested whether RNA or RNA-LL37 in
the lower chamber had any direct influence on the migration of these cell populations and noted an

unexpected and significant chemoattractive effect on CD4[+] T cells, in response to RNA-LL37 (Fig. 2I).

Interestingly, the presence of PMNs, regardless of stimulus, was sufficient to attract a variety of

other cells (results not shown) due to a so far unknown mechanism.

When repeating the above described stimulation experiments with psoriasis PMNs compared to

PMNs from healthy donor, both release of both IL-8 and MIP-1β was significantly increased two-fold

in response to RNA-LL37 (Fig. 2J, K), but not to LPS (Fig. S2F, G). Whereas IL-8 release in response to

'RNA alone' (i.e. without LL37) was indistinguishable from 'unstimulated' samples in healthy donor

PMNs (*cf.* also Fig. 1C), psoriasis PMNs treated with 'RNA alone' (Fig. 2J, arrow) produced significantly

more IL-8 than both 'unstimulated' psoriasis PMNs and healthy donor PMNs stimulated with 'RNA

alone') Since in healthy donors, RNA seemed to strongly depend on LL37 for efficient PMN uptake

and stimulation (*cf.* Fig. 1E, F) and PMNs are the major contributors to elevated LL37 in psoriasis, we

speculated whether psoriasis PMNs might constitutively secrete LL37 and this endogenous LL37

might promote an increased responsiveness to exogenously added RNA alone in psoriasis PMNs.

Psoriasis PMNs constitutively showed an increased baseline release of LL37 compared to healthy

donors (Fig. 2L). We conclude that the combination of RNA with LL37 triggers the release of an

extended array of pro-inflammatory cytokines and effective chemoattractants not triggered by RNA

alone. This was more prominent in psoriasis PMNs, potentially due to a higher abundance of PMN-

derived LL37.

RNA-LL37 complexes trigger the release of RNA- and LL37-containing NETs

Our data so far indicate that RNA and LL37 might contribute to cytokine-mediated inflammation and

immune infiltration in neutrophil-containing skin lesions in psoriasis patients. Such neutrophil

responses would probably be temporary, unless RNA-LL37 triggered the release of additional RNA

and LL37. PMN NETs are known to contain extracellular DNA that acts as an immune stimulant of

pDCs when complexed with LL37 (Lande et al., 2011). But DNA-LL37 cannot activate PMNs (*cf.* Fig.

1B) and RNA release by NETosis has not been shown. NETs would thus only propagate PMN

activation in case they also contained RNA. We therefore next tested (i) whether RNA-LL37 induces

NETosis and (ii) whether NETs contain RNA. Interestingly, electron microscopy (Fig. 3A) and a

neutrophil elastase-based NET assay (see Methods) confirmed significant neutrophil elastase release,

a hallmark of NETosis, in response to RNA-LL37 and PMA (positive control), but also with LL37 alone

(Fig. 3B). However, fluorescence microscopy analysis of fixed PMN samples established that only

RNA-LL37 complexes, and not LL37 alone, prompted the formation of LL37-positive NET-like

structures (Fig. 3C). To check whether the NETs also contained RNA, an RNA-selective fluorescent

dye, SYTO RNAselect, was used to stain NETs. This staining showed released endogenous RNA (green)

in both PMA and RNA-LL37-mediated NETs (Fig. 3C). Although the dye's specificity for RNA has been

established already (Li et al., 2006) we confirmed that RNA staining, but not DNA staining, was

sensitive to RNase A and thus RNA-specific (Fig. S3A, B). The presence of cellular RNA in NET
structures was also confirmed by the use of specific Abs against pseudo-uridine (ΨU), a nucleotide
absent from both DNA and also the synthetic RNA40 used for LL37 complex formation and
stimulation (Figs. 3D). Furthermore, using AF647-labeled or AF488-labeled synthetic RNA for in vitro
stimulation both the extracellular SYTO RNAselect and anti-ΨU signals could be unequivocally
attributed to de-novo released 'cellular RNA' and distinguished from exogenously added 'stimulant
(synthetic) RNA' by (Fig. S3C, D). The same applied to LL37, using Atto488-LL37 (Fig. S3E). To further
exclude artefacts that might arise from staining fixed samples, RNA-LL37 mediated NETosis was also
analyzed and confirmed using live-cell time-lapse analysis of PMNs (Fig. 3E and Movies S1-S3,
quantified in Fig. 3F). Interestingly, in PMNs captured on the verge of NETosis, RNA staining in a
granula-like fashion could be clearly observed (Fig. S3F). Thus, RNA-LL37 activates primary PMNs to
release NETs. In addition to the well established NET components DNA and LL37, these NETs contain
further RNA as a so far unappreciated component.

Increased levels of NETs containing DNA have already been reported in the blood and skin of
psoriasis patients (Hu et al., 2016). To check whether this was also the case for RNA, we stained skin
sections with SYTO RNAselect, anti-LL37 and anti-NE Abs. As expected, psoriatic skin was highly
positive for LL37; interestingly, the RNA signal was also generally stronger when compared to healthy
skin, using identical staining and microscopy settings (Fig. 3G). RNA and LL37 were frequently co-
localized in samples with high PMN infiltration (anti-NE staining), (Fig. 3H and Movie S4). This was
SYTO RNAselect specificity was confirmed using anti-Ψ staining on skin sections (Fig. S3G). This
indicates that RNA-LL37 complexes may act as physiologically relevant activators in psoriatic skin.

However, if RNA-LL37 complexes truly triggered a NET-mediated self-propagating inflammatory loop
of repetitive PMN activation, we would expect that isolated NET content would act in a manner
similar to RNA-LL37 and activate naïve PMNs to NETose. Indeed, when we harvested NETs from
PMNs stimulated either with PMA, RNA-LL37 complexes, LL37 or left untreated (mock),and
transferred them to unstimulated naïve PMNs, these responded with de-novo NET release in turn,
monitored by fixed microscopy of stimulated PMNs (Fig. 3J, quantified in I).. Collectively, RNA-LL37
complexes were found in psoriatic skin and in vitro prompted the release of NETosis. These NETs in
turn had the capacity to activate naive PMNs, thus providing the requirements for a self-propagating
feed forward loop of immune activation that might apply to skin PMNs.

RNA-LL37 complexes and NETs activate PMNs via TLR8/TLR13

In order to gain an insight which receptor mediates the observed effects of RNA-LL37 in primary
PMNs, we first compared BM-PMNs from WT mice and mice deficient for *Unc93b1*, a critical
chaperone of endosomal TLRs (Eigenbrod et al., 2012). Evidently, not only the cytokine response of

BM-PMNs to bRNA+LL37 (and the control stimulus R848) was strictly dependent on *Unc93b1* and
thus endosomal TLRs (Fig. 4A); NETosis triggered by bRNA-LL37 complexes, but not the TLR-
independent control stimulus PMA, was also *Unc93b1* dependent (Fig. 4B). Given that RNA is a
known dual TLR7 and TLR8 agonist (Colak et al., 2014) and primary human PMNs do not express *TLR7*
(Berger et al., 2012), we suspected TLR8 to mediate RNA-LL37-mediated PMN activation.
Unfortunately, human primary PMNs are not amenable to RNAi or genetic editing. Therefore, we first
tested the responsiveness of BM-PMNs from mice deficient for TLR13, the functional counterpart of
human TLR8 and primary bRNA sensor in mice (Eigenbrod and Dalpke, 2015). In murine PMNs both
cytokine and NET responses to bRNA+LL37 were strictly dependent on TLR13 (Fig. 4A, C), whereas
the negative controls LPS, *E. co*li or PMA were not, respectively (*cf.* Fig. 4A or B, respectively).
Complementarily, the failure of *TLR8* CRISPR-deleted BlaER1 monocytes (Vierbuchen et al., 2017) to
respond to RNA-LL37 complexes and a novel TLR8-specific small molecule, TL8, (Fig. 4D) indicated a
dependence of RNA-LL37 immunostimulatory activity on TLR8 in the human cells, including PMNs.
Thus the stimulatory potential of RNA-LL37 complexes can be considered to be dependent on
TLR8/TLR13 as a NET-triggering sensing system.

PMN cytokine release and NET formation by RNA-LL37 can be blocked by iODNs

Having identified TLR8 as the responsible receptor in humans, we next wondered whether PMN
activation by RNA-LL37 could be blocked at the level of TLR activation by so-called inhibitory
oligonucleotides (iODNs). IRS869, IRS661 and IRS954 represent such TLR7 and 9-inhibitory iODNs that
were recently proposed for the treatment of another autoimmune disease, namely systemic lupus
erythematosus (SLE), whereas IRS546 was used as a non- 'control' ODN inhibitory (Barrat and
Coffman, 2008; Barrat et al., 2005; Duramad et al., 2005; Guiducci et al., 2010). Given the similarities
between endosomal TLRs (Colak et al., 2014), we speculated that some of these iODNs could
potentially also block TLR8, but the effects of iODNs on this receptor have not been studied
systematically. HEK293T cells do not express TLR7-9 but can be rendered responsive to their agonists
by exogenously expressing TLR7, TLR8 or TLR9 (control) upon transfection (Colak et al., 2014). In
HEK293T cells transfected and stimulated in this way, IRS869 and IRS954 blocked NF-κB activation in
response to R848 for TLR7 and 8 (Figs. S4A and B) and CpG for TLR9 (Fig. S4C). IRS661 effectively
blocked TLR7- but not TLR9-mediated activation. Paradoxically, IRS661 strongly increased R848-
mediated NF-κB activation via TLR8 (Fig. S4B), similar to observations made for T-rich ODNs
combined with R848 (Jurk et al., 2006) and highlighting the differences between R848 and RNA
reported previously (Colak et al., 2014). However, for RNA complexed with DOTAP (a synthetic
complexing agent promoting endosomal uptake (Yasuda et al., 2005), similarly to LL37), all three
iODNs blocked TLR8-mediated NF-κB activation (Fig. 4E). As the 'control' ODN IRS546 blocked NF-κB
activation in response to TLR7-9 stimulation, in our hands it did not represent a proper control unlike

previously published (Barrat et al., 2005). Nevertheless, the effects seemed specific for endosomal
TLR signaling as the effect of all iODNs on NF-κB activity induced by TNF (a non-TLR receptor pathway
activating NF-κB, Fig. S4D) or MyD88 overexpression (TLR receptor-independent NF-κB activation,
Fig. S4E) were minor. Having excluded a toxic effect on primary PMNs (Fig. S4F), we investigated
whether the observed effects could be transferred to endogenous TLR8 activation in primary PMNs,
where LL37 as a physiologically relevant uptake reagent was again used. Here, we found that iODNs
IRS661 and IRS954 were able to inhibit IL-8 release by primary PMNs from healthy donors in
response to RNA-LL37 at nanomolar iODN concentrations (Fig. 4F), whereas LPS-mediated IL-8
release was unaffected (Fig. S4G and (Saitoh et al., 2012)). The blocking effect extended to MIP-1β
(Fig. 4G), similar to the effect observed for chloroquine (Fig. S4H). Importantly, RNA-LL37-elicited
NETosis by primary human PMNs could be effectively blocked by nanomolar concentrations of
IRS661 (Fig. 4H, quantified in Fig. 4I). These results are in agreement with the observed TLR8-
dependence and iODN-mediated inhibition of NETosis in HIV infections (Saitoh et al., 2012) and
indicate an applicability to RNA-LL37 complexes. In conclusion, known pre-clinical iODNs are able to
block RNA-LL37-mediated cytokine and NET release.

Discussion

Although systemic therapies with biologicals have brought benefit for many psoriasis patients in
industrialized countries (Eberle et al., 2016), psoriasis remains a significant health problem
worldwide. One reason is that triggers and mechanisms leading to disease chronification,
exacerbation and flare-ups are not well understood. This uncertainty also applies to the role of PMNs
in these processes (Schon et al., 2017). Since PMNs can release nucleic acids, constitute the main
sources of LL37 in humans and congregate in psoriatic skin, we chose to study PMN RNA-LL37-
mediated responses after identifying TLR8 as an RNA receptor on PMNs (Berger et al., 2012). We
now provide novel evidence that RNA-LL37 can directly and TLR8-dependently activate PMNs to
release chemo-/cytokines and NETs, and that RNA originated from NETs could serve as a novel
immunostimulatory component within NETs, which has not been appreciated so far. Interestingly,
insect NETs contain primarily RNA rather than DNA (Altincicek et al., 2008), and eosinophils pre-store
RNA within their granules (Behzad et al., 2010). RNA may thus be a common and functionally
important NET component that has not been described for mammalian immune cells and ought to be
explored further. Given that RNA can also amplify responses to other PRR ligands (Noll et al., 2017),
the role of RNA will thus be interesting to study in other NET-related disease, e.g. SLE (Gestermann et
al., 2018), atherosclerosis (Warnatsch et al., 2015) and even cancer (Gregoire et al., 2015), where the
focus so far has been largely on DNA.

Recently, the effects of DNA-/RNA-LL37 complexes on pDCs were reported as a novel mechanism in
psoriasis inflammation (Ganguly et al., 2009; Lande et al., 2007). However, since pDC are only a
minor cell population even in psoriatic skin and they cannot release nucleic acids, their contribution
to cell-mediated inflammation and the provision of self-ligands is restricted to amplifying an already
ongoing pathology, first requiring pDC-attracting chemokines and an existing presence of self-ligands
in the skin. We found that the combination of RNA and LL37 produces an inflammatory ligand that is
not only directly sensed by PMNs; it also prompted the release of further components for additional
RNA-LL37 and DNA-LL37 complexes from PMNs. Whereas for pDCs both "inflammatory ingredients",
RNA or DNA and LL37, will need to be provided by another cellular source first, a small number of
activated and NETing PMNs – by releasing both, LL37 and RNA/DNA – may spark potent, self-
propagating inflammation *inter pares* and *in situ.* Following an initial activation of PMNs by
endogenous RNA from damaged skin cells (so-called Köbner phenomenon) or by bacterial
infection/RNA (Eigenbrod and Dalpke, 2015 and manuscript in press[1];*cf.* Fig. 1K) immune activation
may get out of control due to chemokine release and NET formation (and thus nucleic acid and LL37

---

[1] Anna Hafner, Ulrike Kolbe, Isabel Freund, Virginia Castiglia, Pavel Kovarik, Tanja Poth, Franziska Herster, Markus A. Weigand, Alexander N. Weber, Alexander H. Dalpke and Tatjana Eigenbrod: Crucial role of nucleic acid sensing via endosomal Toll-like receptors for the defense of *Streptococcus pyogenes in vitro* and *in vivo*; Front. Immunol. | doi: 10.3389/fimmu.2019.00198.

extrusion (Neumann et al., 2014)) which has been observed in the skin of psoriasis patients (Hu et al.,

2016). Although the concentration of the cytokines released after 4 h stimulation was moderate,

release over longer time periods and the sheer number of PMNs found in psoriatic skin may

nevertheless contribute substantially to local inflammation and attraction of additional leukocytes

over time. The PMN-mediated ongoing release of NET DNA, RNA and LL37 would eventually enable

pDCs and other immune cells to join the vicious cycle of activation fueled by endogenous nucleic

acids (Fig. S5) (Chamilos et al., 2012; Ganguly et al., 2009; Lande et al., 2007). Additionally, cytokine-

mediated inflammation is known to favor KC hyper-proliferation (Hanel et al., 2013), LL37 production

(Schauber et al., 2007), TLR9 responsiveness (Morizane et al., 2012) and IFN production (Zhang et al.,

2016), as well as T cell and monocyte attraction and polarization (Ghoreschi et al., 2007). In

conclusion, our study provides a plausible explanation how PMNs may contribute to early disease

development and, based on the observed effects of iODNs on PMN activation, warrants the further

exploration of TLR pathways and PMNs as targets for restricting innate immune activation in psoriatic

skin.

Materials and Methods

Reagents

All chemicals were from Sigma unless otherwise stated. PRR agonists and LL37 were from Invivogen

except RNA40 (iba Lifescience, normal and AF647- or AF488-labeled) and CpG PTO 2006 (TIB

Molbiol), see Table S2 and 3. iODNs used in this study with their respective sequences are listed in

Table S3 and were from TIB Molbiol. Total human mRNA was isolated from HEK293T cells using the

RNeasy kit on a QIAcube, both from QIAGEN. *S. aureus* RNA was isolated as described (Eigenbrod et

al., 2015). Genomic human DNA was isolated from whole blood using QIAamp DNA Blood Mini Kit

from Qiagen (51106) and phosphodiester DNA from TIB Molbiol. LL37 was from InvivoGen (see Table

S2) and DOTAP from Roth, L787.2. LL37 was Atto488 (from Atto-Tec as a carboxy-reactive reagent)-

labeled using standard procedures. For complex formation 5.8 µM RNA40 (approximately 34.4 µg/ml

and equimolar to R848 used in this setting), 1 µM ssDNA (approximately 20 µg/ml and equimolar to

CpG used in this setting, sequence see Table S3), genomic DNA (20 µg/ml) or bacterial RNA (10

µg/ml) was mixed together with 10 µg LL37 (see Table S2, Atto-488 where indicated) and left for one

hour at RT. For experiments with BM-PMNs, 1.25 µg bacterial RNA was complexed with 2.5 µg LL37.

For the RNA-only or LL37-only conditions, the same amounts and volumes were used replacing one

of the constituents by sterile, endotoxin-free H_2O. Ficoll was from Millipore. Antibodies used for flow

cytometry, ImageStream analysis or fluorescence microscopy are listed in Table S4 as well as the

recombinant cytokines used in this study. Constructs used for HEK293T transfection are listed in

Table S5.

Study participants and sample acquisition

All patients and healthy blood donors included in this study provided their written informed consent

before study participation. Approval for use of their biomaterials was obtained by the local ethics

committees at the University Hospitals of Tübingen and Heidelberg, in accordance with the principles

laid down in the Declaration of Helsinki as well as applicable laws and regulations. All blood or skin

samples obtained from psoriasis patients (median age 41.8 years, PASI >10, no systemic treatments

at the time of blood/skin sampling) were obtained at the University Hospitals Tübingen or

Heidelberg, Departments of Dermatology, respectively, and were processed simultaneously with

samples from at least one healthy donor matched for age and sex (recruited at the University of

Tübingen, Department of Immunology). Skin sections were from 12 patients with Plaque Psoriasis

and 1 patient with Psoriasis guttata.

Mice and isolation of bone-marrow derived PMNs (BM-PMNs)

Unc93b1$^{3d/3d}$-(Tabeta et al., 2006), *Tlr13*-deficient (Li and Chen, 2012) mice (both C57BL/6

background) and WT C57BL/6 mice between 8 and 20 weeks of age were used in accordance with

local institutional guidelines on animal experiments and under specific locally approved protocols for

sacrificing and *in vivo* work. All mouse colonies were maintained in line with local regulatory

guidelines and hygiene monitoring. Bone-marrow (BM)-PMNs were isolated from the bone marrow

using magnetic separation (mouse Neutrophil isolation kit, Miltenyi Biotec, 130-097-658) and

following the manufacturer's instructions. PMN stimulation was carried out for 5 hours at 37 °C and

5% CO_2. Thereafter supernatants were harvested and used for ELISA. For microscopy the cells were

stimulated for 16 h and subsequently stained.

Neutrophil isolation and stimulation

Whole blood (EDTA-anticoagulated) was diluted in PBS (Thermo Fisher, 14190-169), loaded on Ficoll

(1.077 g/ml, Biocoll, ab211650) and centrifuged for 25 min at 366 x g at 21 °C without brake. All

layers were discarded after density gradient separation except for the erythrocyte-granulocyte

pellet. Thereafter, erythrocyte lysis (using 1x ammonium chloride erythrocyte lysis buffer, see Table

S6) was performed twice (for 20 and 10 min) at 4 °C on a roller shaker. The remaining cell pellet was

carefully resuspended in culture medium (RPMI culture medium (Sigma Aldrich, R8758) + 10% FBS

(heat inactivated, sterile filtered, TH Geyer, 11682258)) and 1.6 x 10^6 cells/ml were seeded (24 well

plate, 96 well plate). After resting for 30 min at 37°C, 5% CO_2, the cells were pre-treated with

inhibitors (where indicated) for 30 min and subsequently stimulated with the indicated agonists for 4

hours (for ELISA) or for 30 min to 3 h (for FACS analysis or microscopy).

PBMC isolation

Whole blood (EDTA anticoagulant) was diluted in PBS. After density gradient separation using Ficoll

(described above), the PBMC layer was then carefully transferred into a new reaction tube and

diluted in PBS (1:1). The cell suspension was spun down at 1800 rpm for 8 min. The cells were then

washed twice more in PBS, resuspended in culture medium (RPMI + 10% FBS (heat inactivated) + 1%

L-glutamine), before counting and seeding.

Generation and harvest of NETs

Neutrophils were isolated as described above. $5x10^6$cells/ml were seeded per 10 cm dish, rested for

30 min and subsequently stimulated with RNA-LL37 complex, LL37 alone, PMA or left untreated

('Mock' NETs) for 4 hours. Then the medium was removed and the cells were washed with PBS for

three times. Thereafter the adherent cell debris and generated NETs were scratched from the

surface, resuspended in culture medium and stored at -80°C before use.

BlaER1 cells culture, transdifferentiation and stimulation

BlaER1 cells (WT and TLR8$^{-/-}$, kindly provided and validated by Tim Vierbuchen and Holger Heine

from Borstel, Germany {Vierbuchen, 2017 #6405}) were cultured in VLE-RPMI (Merck Biochrom,

FG1415) + 10 % FBS, 1 % Pen/Strep (Gibco, 15140122), 1 % sodium pyruvate (100 mM, Gibco,

11360070), 1 % HEPES soluation (Sigma, H0887)). After reaching a cell concentration not higher than

2×10^6/ml the cells were seeded in a 6 well plates (1×10^6 cells per well) and transdifferentiated in

culture medium adding 150 nM β-estradiol, 10 ng/ml M-CSF and 10 ng/ml hIL-3 for 6 days including 2

medium changes (on day 2 and 5). On day 7 the adherent cells were counted again, seeded (96 well

plate, 5×10^4 cells per well) and left to rest for 1 h. Subsequently, they were stimulated for 18 h and

the supernatants were harvested and collected for ELISA measurements. The transdifferentiation

efficiency was verified by FACS analysis, using CD19, CD14 and CD11b as cell surface markers as

described {Rapino, 2013 #6445}.

Flow cytometry

After PMN isolation and stimulation, the purity and activation status of neutrophils was determined

by flow cytometry. 200 μl of the cell suspension was transferred into a 96 well plate (U-shape) and

spun down for 5 min at 322 x g, 4 °C. FcR block was performed using pooled human serum diluted

1:10 in FACS buffer (PBS, 1 mM EDTA, 2% FBS heat inactivated) for 15 min at 4 °C. After washing, the

samples were stained for approximately 20-30 min at 4°C in the dark. Thereafter, fixation buffer (4%

PFA in PBS) was added to the cell pellets for 10 min at RT in the dark. After an additional washing

step, the cell pellets were resuspended in 150 μl FACS buffer. Measurements were performed on a

FACS Canto II from BD Bioscience, Diva software. Analysis was performed using FlowJo V10 analysis

software.

433 ELISA

434 Cytokines were determined in half-area plates (Greiner, Bio-one) using duplicates or triplicates and

435 measuring with a standard plate reader. The assays were performed according to the manufacturer's

436 instructions (Biolegend, R&D systems), using appropriate dilutions of the supernatants. For LL37

437 determination a kit from HycultBiotech (HK321-02) was used following the manufacturer's

438 instructions.

439 ImageStream analysis

ImageStream analysis was used to analyze internalization of RNA-LL37 complexes using spot-counts

and tracking single cells. The cells were first seeded in a 96 well plate, 8 x 10^6 cells/ml, 125 µl per

well. Subsequently, they were stimulated for 1 hour with RNA-AF647 (IBA technologies) and/or LL37-

Atto488 (kindly provided by Hubert Kalbacher, University of Tübingen). FcR block and surface

staining (here CD15 PE) was performed as described above. After fixation, the cells were

permeabilized with 0.05 % Saponin (Applichem, A4518.0100) for 15 min at RT in the dark. After

washing, nuclei were stained with Hoechst 33342 (Sigma, B2261, 1 µg/ml) for 5 min at RT in the dark,

washed and resuspended in 50 µl FACS buffer and transferred into a 1.5 ml Eppendorf tube. At least

10.000 cells were acquired for each sample with 40x magnification using an ImageStream X MKII with

the INSPIRE instrument controller software (Merck-Millipore/Amnis). Data were analyzed with IDEAS

Image analysis software. All samples were gated on single cells in focus.

451 Fluorescence Microscopy of fixed neutrophils

The cells were seeded in a 96 well plate at 1.6 x 10^6 cells/ml, 125 µl per well. Subsequently they were

stimulated with the complexes for 30 min and 1 h using RNA-AF647/AF488 and/or LL37-Atto488. FcR

block, staining, fixation and permeabilization were performed as for Flow cytometry. The cell pellets

were resuspended in 50-100 µl FACS buffer. 40 µl of the cell suspension was pipetted on a Poly-L-

Lysine coated coverslip (Corning, 734-1005) and the cells were left to attach for one hour in the dark.

ProLong Diamond Antifade (Thermo Fisher, P36965) was used to mount the coverslips on uncoated

microscopy slides. For NET analysis PMNs were seeded in 24 well plates, containing Poly-L-Lysine

coated coverslips and stimulated with RNA-LL37 complexes or PMA (600 nM) for 3 hours. NETs were

fixed and stained using the protocol from Brinkmann et al.(Brinkmann et al., 2010). Where indicated,

100 µg/ml RNase A (DNase, protease-free, ThermoFisher, EN0531) was added after fixation and

incubated overnight at 37°C. RNA was stained using SYTO RNAselect Green fluorescent dye (Thermo

Fisher, 50 µM) or anti-ΨU antibody (see Table S4) and nuclear DNA was stained with Hoechst33342

(Thermo Fisher, 1 µg/ml). LL37 and PMNs were visualized using an unconjugated rabbit anti-LL37

antibody or unconjugated mouse anti-NE with subsequent staining with an AF647-conjugated anti-

rabbit or an AF594-conjugated anti-mouse antibody, respectively (see Supplementary Table S4), after

blocking with pooled human serum (1:10 in PBS). Secondary antibodies alone did not yield any

significant staining. The slides were left to dry overnight at RT in the dark and were then stored at 4

°C before microscopy. The measurements were conducted with a Nikon Ti2 eclipse (100x

magnification) and the analysis was performed using Fiji analysis software.

Fluorescence Microscopy of tissue samples

Skin samples from psoriasis patients with a PASI ≥ 10 and without systemic treatment, and healthy

skin samples paraffin-embedded according to standard procedures were deparaffinized and

rehydrated using Roti Histol (Roth, 6640.1) and decreasing concentrations of ethanol (100%, 95%,

80% and 70%). After rinsing in ddH$_2$O, antigen retrieval was performed by boiling for 10-20 min in

citrate buffer (0.1 M, pH=6). The skin tissue was then washed 3 times for 5 min with PBS. Blocking

was performed using pooled human serum (1:10 in PBS) for 30 min at RT. The primary antibody was

added either overnight at 4°C or for 1 hour at RT. After 3 washes, the secondary antibody was added

for 30 min at RT in the dark. After another 3 washes, SYTO RNAselect Green fluorescent dye (Thermo

Fisher, 50 µM) was added for 20 min at RT in the dark. Thereafter, the samples were washed again

and Hoechst 33342 (ThermoFisher, 1 µg/ml) was added for 5 min. Then 3 last washes were

performed before using ProLong Diamond Antifade (Thermo Fisher, P36965) for mounting. The

samples were left to dry overnight at RT in the dark before being used for microscopy or stored at

4°C. The specimens were analyzed on a Nikon Ti2 eclipse bright-field fluorescence microscope (10x-

60x magnification) and the analysis was performed using Fiji analysis software. Autofluorescence in

multiple channels typical for the stratum corneum was labeled "AF".

Live cell imaging of primary neutrophils

Human neutrophils were isolated by magnetic separation using MACSxpress whole blood neutrophil

isolation kit (Miltenyi Biotec, 130-104-434) and were seeded into a micro-insert 4 well dish (Ibidi

80406). Hoechst 33342 (1 µg/ml) and SYTO RNAselect Green fluorescent dye (50 µM) were added to

the cells and incubated for 20 min at 37°C, 5 % CO$_2$. Live cell imaging was performed by using Nikon

Ti2 eclipse bright-field fluorescence microscope (40x magnification) including a CO$_2$-O$_2$ controller

from Okolab. Measurements were started immediately after adding of stimuli. Time-lapse analysis

was performed by taking pictures every 3 min for at least 2 hours. Image analysis was performed

using NIS Elements from Nikon and Fiji analysis software.

Luminex cytokine multiplex analysis

All samples were stored at -70 °C until testing. The samples were thawn at room temperature,

vortexed, spun at 18,000 x g for 1 min to remove debris and the required sample volumes were

removed for multiplex analysis according to the manufacturer's recommendations. The samples

were successively incubated with the capture microspheres, a multiplexed cocktail of biotinylated,

reporter antibodies, and a streptavidin-phycoerythrin (PE) solution. Analysis was performed on a

Luminex 100/200 instrument and the resulting data were interpreted using proprietary data analysis

software (Myriad RBM). Analyte concentrations were determined using 4 and 5 parameter, weighted

and non-weighted curve fitting algorithms included in the data analysis package.

Cytometric bead array

A cytometric bead array was performed using the "Human inflammatory cytokine kit" from BD

Bioscience (551811) and following the manufacturer's instruction. 25 µl of samples and standards

were added to 25 µl of the capturing bead mixture. Additionally, 25 µl of PE detection reagent was

added to all tubes and incubated for 3 h at RT in the dark. Thereafter, 1 ml of wash buffer was added

to each tube and centrifuged at 200 x g for 5 minutes. The supernatant was carefully removed and

the pellet was resuspended in 300 µl wash buffer. Measurements were performed with the FACS

Canto II from BD Bioscience operated using Diva software. Analysis was performed with Soft Flow

FCAP Array v3 analysis software from BD Bioscience.

Transwell experiments

PBMCs and neutrophils were always used from the same donor in each experiment. Transwell inserts

were loaded with 100 µl of PBMC suspension (0.8×10^6 cells/insert). In the lower chamber, either

PMNs were seeded as described above (same plate size, same volume, same cell concentration)

using polycarbonate, 24 well plates, 3 µm pores, Corning, 734-1570) and stimulated for 4 h with

stimuli as indicated. Alternatively, media containing the stimuli only (i.e. no PMNs) or media

containing only MIP-1β (30 and 150 pg /ml), IL-16 (300 and 1500 pg/ml) or SDF-1α (control, 100

ng/ml) and no PMNs were added. After 4 h, the lower compartment was harvested and FACS staining

was performed as described above. The total number of migrated cells was acquired using counting

beads (Biolegend, 424902) on a FACS Canto II (BD Bioscience) with Diva software. Analysis was

performed using FlowJo V10 analysis software.

Neutrophil elastase NETosis assay

Neutrophil extracellular trap formation was determined using the colorimetric NETosis Assay Kit from

Cayman Chemicals based on the enzymatic activity of NET-associated neutrophil elastase. PMNs from

various healthy donors were isolated as described above and stimulated with RNA-LL37 complex, or

PMA and a calcium ionophore (A-23187) as positive controls for 1 or 3 hours. The assay was

performed following the manufacturer's instructions. The absorbance was then measured at 405 nm

using a standard plate reader.

Transient transfection of HEK293T cells

HEK293T cells were transiently transfected using the CaPO$_4$ method as described (Colak et al., 2014).

Cells were seeded in 24-well plates at a density of 14 x 10^5 cells/ml 2-3 h prior to transfection. For the

transfection of one well, 310 ng of plasmid DNA (100 ng TLR plasmid, 100 ng firefly luciferase NF-κB

reporter, 10 ng *Renilla* luciferase control reporter, and 100 ng EGFP plasmid) was mixed with 1.2 μl of

a 2 M CaCl$_2$ solution and filled up with sterile endotoxin-free H$_2$0 to obtain a total reaction volume of

10 μl. After the addition of 10 μl of 2X HBS solution (50 mM HEPES (pH 7.05), 10 mM KCl, 12 mM

Glucose, 1.5 mM Na$_2$HPO$_4$), the mixture was then added to the cell suspension. As negative controls,

TLR coding plasmids were replaced by empty vectors carrying the appropriate backbone of the TLR

plasmids. After the addition of the transfection complexes, the cells were incubated either for 24 h

followed by stimulation, or kept for 48 h without stimulation (MyD88 expression). For stimulation,

the media was aspirated and replaced by fresh growth medium in which TNFα or the different TLR

ligands (R848, CpG, RNA40) with or without IRS were diluted to appropriate concentrations. TLR8

activation with RNA40 was facilitated by complexation of the RNA with DOTAP (L787.1, Roth). RNA40

and DOTAP were first diluted in 1X HBS separately. Next, RNA40/HBS was diluted 1:3 in DOTAP/HBS.

The solution was carefully mixed by pipetting up and down. After 15 min of incubation at RT, the

mixture was 1:6.7 diluted in growth medium (with or without IRS) and finally dispensed (500 μl/well)

into the wells containing transfected HEK293T cells. Each tested condition was measured in

triplicates. The cells were stimulated and inhibited for 24 h at 37°C.

Dual Luciferase Reporter Assay

After checking transfection efficiency via EGFP fluorescence microscopy, HEK293T supernatants were

aspirated and 60 μl of 1X passive lysis buffer (E194A, Promega) added per well. The plate was then

incubated for 15 min at RT on the plate shaker and subsequently stored at -80 °C for at least 15

minutes to facilitate complete cell lysis. After thawing, 60 μl were transferred into a V-bottom 96-

well plate and centrifuged for 10 min at 2500 rpm and 4 °C to pellet cell debris. 10 μl supernatant

were then transferred into a white microplate and each condition was measured in triplicates using

the FLUOstar OPTIMA device (BMG Labtech). Firefly and Renilla luciferase activity were determined

using the Promega Dual luciferase kit. Both enzyme activities were measured for 12.5 s with 24

intervals of 0.5 s, respectively. The data was analyzed by calculating the ratio of the two measured

signals, thereby normalizing each firefly luciferase signal to its corresponding Renilla luciferase signal.

The ratios were represented as the relative light units (RLU) of NF-κB activation.

Statistics

Experimental data was analyzed using Excel 2010 (Microsoft) and/or GraphPad Prism 6 or 7,

microscopy data with ImageJ/Fiji, flow cytometry data with FlowJo 10. In case of extreme values,

outliers were statistically identified using the ROUT method at high (0.5%) stringency. Normal

distribution in each group was always tested using the Shapiro-Wilk test first for the subsequent

choice of a parametric (ANOVA, Student's t-test) or non-parametric (e.g. Friedman, Mann-Whitney U

or Wilcoxon) test. p-values (α=0.05) were then calculated and multiple testing was corrected for in

Prism, as indicated in the figure legends. Values < 0.05 were generally considered statistically as

significant and denoted by * throughout. Comparisons made to unstimulated control, unless

indicated otherwise, denoted by brackets.

Materials & Correspondence

Please address requests to: Alexander N. R. Weber, Interfaculty Institute for Cell Biology, Department

of Immunology, University of Tübingen, Auf der Morgenstelle 15, 72076 Tübingen, Germany. Tel.:

+49 7071 29 87623. Fax: +49 7071 29 4579. Email: alexander.weber@uni-tuebingen.de

Supplementary Material

Is uploaded separately and includes 5 Supplementary figures and 6 supplementary tables.

Authorship contributions

F.H., Z.B., S.D., D.E., T.K. and N.S.M. performed experiments; F.H., Z.B., S.D., D.E., N.S.M. and A.N.R.W. analyzed data; M.H., H. K., M.W.L., L.F., K.S., K.G. and T.E. were involved in sample and reagent acquisition; F.H., D.H., K.G. and T.E. contributed to the conceptual development of the study; F.H. and A.N.R.W. wrote the manuscript and all authors commented on an revised the manuscript; A.N.R.W. coordinated and supervised the entire study. None of the authors declare competing interests.

Acknowledgements

We thank S. Pöschel, J. Berger, S. Haen, K. Preissner, O. Sorensen and A. Dalpke for provision of reagents, samples, technical support and/or helpful discussions and all healthy donors and patients for participation in our study. This study was supported by the Deutsche Forschungsgemeinschaft (German Research Foundation, DFG) CRC TR156 "The skin as an immune sensor and effector organ – Orchestrating local and systemic immunity", the University of Tübingen and the University Hospital Tübingen.

References

Afshar, M., A.D. Martinez, R.L. Gallo, and T.R. Hata. 2013. Induction and exacerbation of psoriasis with Interferon-alpha therapy for hepatitis C: a review and analysis of 36 cases. J Eur Acad Dermatol Venereol 27:771-778.

Alekseyenko, A.V., G.I. Perez-Perez, A. De Souza, B. Strober, Z. Gao, M. Bihan, K. Li, B.A. Methe, and M.J. Blaser. 2013. Community differentiation of the cutaneous microbiota in psoriasis. Microbiome 1:31.

Altincicek, B., S. Stotzel, M. Wygrecka, K.T. Preissner, and A. Vilcinskas. 2008. Host-derived extracellular nucleic acids enhance innate immune responses, induce coagulation, and prolong survival upon infection in insects. J Immunol 181:2705-2712.

Atanasova, M., and A. Whitty. 2012. Understanding cytokine and growth factor receptor activation mechanisms. Crit Rev Biochem Mol Biol 47:502-530.

Barrat, F.J., and R.L. Coffman. 2008. Development of TLR inhibitors for the treatment of autoimmune diseases. Immunol Rev 223:271-283.

Barrat, F.J., T. Meeker, J. Gregorio, J.H. Chan, S. Uematsu, S. Akira, B. Chang, O. Duramad, and R.L.
Coffman. 2005. Nucleic acids of mammalian origin can act as endogenous ligands for Toll-like
receptors and may promote systemic lupus erythematosus. The Journal of experimental medicine
202:1131-1139.

Behzad, A.R., D.C. Walker, T. Abraham, J. McDonough, S. Mahmudi-Azer, F. Chu, F. Shaheen, J.C.
Hogg, and P.D. Pare. 2010. Localization of DNA and RNA in eosinophil secretory granules. Int Arch
Allergy Immunol 152:12-27.

Berger, M., C.Y. Hsieh, M. Bakele, V. Marcos, N. Rieber, M. Kormann, L. Mays, L. Hofer, O. Neth, L.
Vitkov, W.D. Krautgartner, D. von Schweinitz, R. Kappler, A. Hector, A. Weber, and D. Hartl. 2012.
Neutrophils express distinct RNA receptors in a non-canonical way. The Journal of biological
chemistry 287:19409-19417.

Brinkmann, V., B. Laube, U. Abu Abed, C. Goosmann, and A. Zychlinsky. 2010. Neutrophil
extracellular traps: how to generate and visualize them. J Vis Exp

Brinkmann, V., U. Reichard, C. Goosmann, B. Fauler, Y. Uhlemann, D.S. Weiss, Y. Weinrauch, and A.
Zychlinsky. 2004. Neutrophil extracellular traps kill bacteria. Science 303:1532-1535.

Chamilos, G., J. Gregorio, S. Meller, R. Lande, D.P. Kontoyiannis, R.L. Modlin, and M. Gilliet. 2012.
Cytosolic sensing of extracellular self-DNA transported into monocytes by the antimicrobial peptide
LL37. Blood 120:3699-3707.

Colak, E., A. Leslie, K. Zausmer, E. Khatamzas, A.V. Kubarenko, T. Pichulik, S.N. Klimosch, A. Mayer, O.
Siggs, A. Hector, R. Fischer, B. Klesser, A. Rautanen, M. Frank, A.V. Hill, B. Manoury, B. Beutler, D.
Hartl, A. Simmons, and A.N. Weber. 2014. RNA and imidazoquinolines are sensed by distinct TLR7/8
ectodomain sites resulting in functionally disparate signaling events. J Immunol 192:5963-5973.

Cruikshank, W., and F. Little. 2008. Interleukin-16: The Ins and Outs of Regulating T-Cell Activation.
Critical Reviews in Immunology 28:467-483.

Duramad, O., K.L. Fearon, B. Chang, J.H. Chan, J. Gregorio, R.L. Coffman, and F.J. Barrat. 2005.
Inhibitors of TLR-9 act on multiple cell subsets in mouse and man in vitro and prevent death in vivo
from systemic inflammation. J Immunol 174:5193-5200.

Eberle, F.C., J. Bruck, J. Holstein, K. Hirahara, and K. Ghoreschi. 2016. Recent advances in
understanding psoriasis. F1000Res 5:

Eigenbrod, T., and A.H. Dalpke. 2015. Bacterial RNA: An Underestimated Stimulus for Innate Immune
Responses. J Immunol 195:411-418.

Eigenbrod, T., L. Franchi, R. Munoz-Planillo, C.J. Kirschning, M.A. Freudenberg, G. Nunez, and A.
Dalpke. 2012. Bacterial RNA mediates activation of caspase-1 and IL-1beta release independently of
TLRs 3, 7, 9 and TRIF but is dependent on UNC93B. J Immunol 189:328-336.

Eigenbrod, T., K. Pelka, E. Latz, B. Kreikemeyer, and A.H. Dalpke. 2015. TLR8 Senses Bacterial RNA in
Human Monocytes and Plays a Nonredundant Role for Recognition of Streptococcus pyogenes. J
Immunol 195:1092-1099.

Ganguly, D., G. Chamilos, R. Lande, J. Gregorio, S. Meller, V. Facchinetti, B. Homey, F.J. Barrat, T. Zal,
and M. Gilliet. 2009. Self-RNA-antimicrobial peptide complexes activate human dendritic cells
through TLR7 and TLR8. The Journal of experimental medicine 206:1983-1994.

Gestermann, N., J. Di Domizio, R. Lande, O. Demaria, L. Frasca, L. Feldmeyer, J. Di Lucca, and M.
Gilliet. 2018. Netting Neutrophils Activate Autoreactive B Cells in Lupus. J Immunol 200:3364-3371.

Ghoreschi, K., C. Weigert, and M. Rocken. 2007. Immunopathogenesis and role of T cells in psoriasis.
Clin Dermatol 25:574-580.

Gregoire, M., F. Guilloton, C. Pangault, F. Mourcin, P. Sok, M. Latour, P. Ame-Thomas, E. Flecher, T.
Fest, and K. Tarte. 2015. Neutrophils trigger a NF-kappaB dependent polarization of tumor-
supportive stromal cells in germinal center B-cell lymphomas. Oncotarget 6:16471-16487.

Griffiths, C.E., and J.N. Barker. 2007. Pathogenesis and clinical features of psoriasis. Lancet 370:263-
271.

Guiducci, C., M. Gong, Z. Xu, M. Gill, D. Chaussabel, T. Meeker, J.H. Chan, T. Wright, M. Punaro, S.
Bolland, V. Soumelis, J. Banchereau, R.L. Coffman, V. Pascual, and F.J. Barrat. 2010. TLR recognition of
self nucleic acids hampers glucocorticoid activity in lupus. Nature 465:937-941.

Hanel, K.H., C. Cornelissen, B. Luscher, and J.M. Baron. 2013. Cytokines and the skin barrier. Int J Mol
Sci 14:6720-6745.

Hu, S.C., H.S. Yu, F.L. Yen, C.L. Lin, G.S. Chen, and C.C. Lan. 2016. Neutrophil extracellular trap
formation is increased in psoriasis and induces human beta-defensin-2 production in epidermal
keratinocytes. Sci Rep 6:31119.

Janke, M., J. Poth, V. Wimmenauer, T. Giese, C. Coch, W. Barchet, M. Schlee, and G. Hartmann. 2009.
Selective and direct activation of human neutrophils but not eosinophils by Toll-like receptor 8. The
Journal of allergy and clinical immunology 123:1026-1033.

Jurk, M., A. Kritzler, B. Schulte, S. Tluk, C. Schetter, A.M. Krieg, and J. Vollmer. 2006. Modulating
responsiveness of human TLR7 and 8 to small molecule ligands with T-rich phosphorothiate
oligodeoxynucleotides. European Journal of Immunology 36:1815-1826.

Kawasaki, T., and T. Kawai. 2014. Toll-like receptor signaling pathways. Front Immunol 5:461.

Kruger, P., M. Saffarzadeh, A.N. Weber, N. Rieber, M. Radsak, H. von Bernuth, C. Benarafa, D. Roos, J.
Skokowa, and D. Hartl. 2015. Neutrophils: Between host defence, immune modulation, and tissue
injury. PLoS Pathog 11:e1004651.

Kuznik, A., M. Bencina, U. Svajger, M. Jeras, B. Rozman, and R. Jerala. 2011. Mechanism of
endosomal TLR inhibition by antimalarial drugs and imidazoquinolines. Journal of Immunology
186:4794-4804.

Lande, R., E. Botti, C. Jandus, D. Dojcinovic, G. Fanelli, C. Conrad, G. Chamilos, L. Feldmeyer, B.
Marinari, S. Chon, L. Vence, V. Riccieri, P. Guillaume, A.A. Navarini, P. Romero, A. Costanzo, E.
Piccolella, M. Gilliet, and L. Frasca. 2014. The antimicrobial peptide LL37 is a T-cell autoantigen in
psoriasis. Nat Commun 5:5621.

Lande, R., D. Ganguly, V. Facchinetti, L. Frasca, C. Conrad, J. Gregorio, S. Meller, G. Chamilos, R.
Sebasigari, V. Riccieri, R. Bassett, H. Amuro, S. Fukuhara, T. Ito, Y.J. Liu, and M. Gilliet. 2011.
Neutrophils activate plasmacytoid dendritic cells by releasing self-DNA-peptide complexes in
systemic lupus erythematosus. Sci Transl Med 3:73ra19.

Lande, R., J. Gregorio, V. Facchinetti, B. Chatterjee, Y.H. Wang, B. Homey, W. Cao, B. Su, F.O. Nestle,
T. Zal, I. Mellman, J.M. Schroder, Y.J. Liu, and M. Gilliet. 2007. Plasmacytoid dendritic cells sense self-
DNA coupled with antimicrobial peptide. Nature 449:564-569.

Li, Q., Y. Kim, J. Namm, A. Kulkarni, G.R. Rosania, Y.H. Ahn, and Y.T. Chang. 2006. RNA-selective, live
cell imaging probes for studying nuclear structure and function. Chem Biol 13:615-623.

Li, X.D., and Z.J. Chen. 2012. Sequence specific detection of bacterial 23S ribosomal RNA by TLR13.
Elife 1:e00102.

Lindau, D., J. Mussard, B.J. Wagner, M. Ribon, V.M. Ronnefarth, M. Quettier, I. Jelcic, M.C. Boissier,
H.G. Rammensee, and P. Decker. 2013. Primary blood neutrophils express a functional cell surface
Toll-like receptor 9. European Journal of Immunology 43:2101-2113.

Matsukura, S., F. Kokubu, M. Kurokawa, M. Kawaguchi, K. Ieki, H. Kuga, M. Odaka, S. Suzuki, S.
Watanabe, T. Homma, H. Takeuchi, K. Nohtomi, and M. Adachi. 2007. Role of RIG-I, MDA-5, and PKR
on the expression of inflammatory chemokines induced by synthetic dsRNA in airway epithelial cells.
Int Arch Allergy Immunol 143 Suppl 1:80-83.

Menten, P., A. Wuyts, and J. Van Damme. 2002. Macrophage inflammatory protein-1. Cytokine
Growth Factor Rev 13:455-481.

Morizane, S., and R.L. Gallo. 2012. Antimicrobial peptides in the pathogenesis of psoriasis. J Dermatol
39:225-230.

Morizane, S., K. Yamasaki, B. Muhleisen, P.F. Kotol, M. Murakami, Y. Aoyama, K. Iwatsuki, T. Hata,
and R.L. Gallo. 2012. Cathelicidin antimicrobial peptide LL-37 in psoriasis enables keratinocyte
reactivity against TLR9 ligands. The Journal of investigative dermatology 132:135-143.

Neumann, A., E.T. Berends, A. Nerlich, E.M. Molhoek, R.L. Gallo, T. Meerloo, V. Nizet, H.Y. Naim, and
M. von Kockritz-Blickwede. 2014. The antimicrobial peptide LL-37 facilitates the formation of
neutrophil extracellular traps. Biochem J 464:3-11.

Noll, F., J. Behnke, S. Leiting, K. Troidl, G.T. Alves, H. Muller-Redetzky, K.T. Preissner, and S. Fischer.
2017. Self-extracellular RNA acts in synergy with exogenous danger signals to promote inflammation.
PLoS One 12:e0190002.

Radic, M., and T.N. Marion. 2013. Neutrophil extracellular chromatin traps connect innate immune
response to autoimmunity. Seminars in immunopathology 35:465-480.

Roth, S., W. Solbach, and T. Laskay. 2016. IL-16 and MIF: messengers beyond neutrophil cell death.
Cell Death Dis 7:e2049.

Saitoh, T., J. Komano, Y. Saitoh, T. Misawa, M. Takahama, T. Kozaki, T. Uehata, H. Iwasaki, H. Omori,
S. Yamaoka, N. Yamamoto, and S. Akira. 2012. Neutrophil extracellular traps mediate a host defense
response to human immunodeficiency virus-1. Cell Host Microbe 12:109-116.

Schauber, J., R.A. Dorschner, A.B. Coda, A.S. Buchau, P.T. Liu, D. Kiken, Y.R. Helfrich, S. Kang, H.Z.
Elalieh, A. Steinmeyer, U. Zugel, D.D. Bikle, R.L. Modlin, and R.L. Gallo. 2007. Injury enhances TLR2
function and antimicrobial peptide expression through a vitamin D-dependent mechanism. The
Journal of clinical investigation 117:803-811.

Schmidt, N.W., F. Jin, R. Lande, T. Curk, W. Xian, C. Lee, L. Frasca, D. Frenkel, J. Dobnikar, M. Gilliet,
and G.C. Wong. 2015. Liquid-crystalline ordering of antimicrobial peptide-DNA complexes controls
TLR9 activation. Nat Mater 14:696-700.

Schon, M.P., S.M. Broekaert, and L. Erpenbeck. 2017. Sexy again: the renaissance of neutrophils in
psoriasis. Exp Dermatol 26:305-311.

Sen, B.B., E.N. Rifaioglu, O. Ekiz, M.U. Inan, T. Sen, and N. Sen. 2013. Neutrophil to lymphocyte ratio
as a measure of systemic inflammation in psoriasis. Cutan Ocul Toxicol

Sorensen, O., K. Arnljots, J.B. Cowland, D.F. Bainton, and N. Borregaard. 1997. The human
antibacterial cathelicidin, hCAP-18, is synthesized in myelocytes and metamyelocytes and localized to
specific granules in neutrophils. Blood 90:2796-2803.

Tabeta, K., K. Hoebe, E.M. Janssen, X. Du, P. Georgel, K. Crozat, S. Mudd, N. Mann, S. Sovath, J.
Goode, L. Shamel, A.A. Herskovits, D.A. Portnoy, M. Cooke, L.M. Tarantino, T. Wiltshire, B.E.
Steinberg, S. Grinstein, and B. Beutler. 2006. The Unc93b1 mutation 3d disrupts exogenous antigen
presentation and signaling via Toll-like receptors 3, 7 and 9. Nature Immunology 7:156-164.

Tecchio, C., A. Micheletti, and M.A. Cassatella. 2014. Neutrophil-derived cytokines: facts beyond
expression. Front Immunol 5:508.

Terui, T., M. Ozawa, and H. Tagami. 2000. Role of neutrophils in induction of acute inflammation in T-
cell-mediated immune dermatosis, psoriasis: a neutrophil-associated inflammation-boosting loop.
Experimental dermatology 9:1-10.

Vierbuchen, T., C. Bang, H. Rosigkeit, R.A. Schmitz, and H. Heine. 2017. The Human-Associated
Archaeon Methanosphaera stadtmanae Is Recognized through Its RNA and Induces TLR8-Dependent
NLRP3 Inflammasome Activation. Front Immunol 8:1535.

Warnatsch, A., M. Ioannou, Q. Wang, and V. Papayannopoulos. 2015. Inflammation. Neutrophil
extracellular traps license macrophages for cytokine production in atherosclerosis. Science 349:316-
320.

Yao, Y., L. Richman, C. Morehouse, M. de los Reyes, B.W. Higgs, A. Boutrin, B. White, A. Coyle, J.
Krueger, P.A. Kiener, and B. Jallal. 2008. Type I interferon: potential therapeutic target for psoriasis?
PLoS One 3:e2737.

Yasuda, K., P. Yu, C.J. Kirschning, B. Schlatter, F. Schmitz, A. Heit, S. Bauer, H. Hochrein, and H.
Wagner. 2005. Endosomal translocation of vertebrate DNA activates dendritic cells via TLR9-
dependent and -independent pathways. J Immunol 174:6129-6136.

Zhang, L.J., G.L. Sen, N.L. Ward, A. Johnston, K. Chun, Y. Chen, C. Adase, J.A. Sanford, N. Gao, M.
Chensee, E. Sato, Y. Fritz, J. Baliwag, M.R. Williams, T. Hata, and R.L. Gallo. 2016. Antimicrobial
Peptide LL37 and MAVS Signaling Drive Interferon-beta Production by Epidermal Keratinocytes
during Skin Injury. Immunity 45:119-130.

Abbreviations

APC – antigen-presenting cell; bRNA – bacterial ribonucleic acid; Enhanced green fluorescence
protein – EGFP; Human embryonic kidney – HEK; HLA – Human leucocyte antigen; iODN – inhibitory
oligonucleotides; Microbe-associated molecular pattern – MAMPs; MHC – Major histocompatibility
complex; Inhibitory oligodeoxynucleotides – iODNs; Interferon – IFN; Interleukin – IL; Keratinocytes –
KC; PASI – psoriasis area and severity; Plasmacytoid dendritic cells – pDCs; PMA – phorbol myristate

acetate; Pattern recognition receptor – PRR; Polymorphonuclear leukocytes – PMNs; NE - Neutrophil

elastase; Neutrophil extracellular trap - NET; Scanning electron microscopy – SEM; Systemic lupus

erythematosus – SLE; TLR – Toll-like receptor; Tumor necrosis factor (TNF).

Figure legends

Figure 1: Human neutrophils take up and respond to RNA when complexed to LL37

(A-C) IL-8 release (ELISA, 4 h stimulation) and CD62L shedding (flow cytometry, 1 or 2 h) by PMNs from healthy donors stimulated with LPS, R848 or CpG ODN (A, n=6-7), ssDNA and genomic DNA (B, n=5), or RNA40 (C, n=8-15), with or without LL37. (D) Electron microscopy of PMNs incubated with RNA-LL37 for 20 min (n=1). (E-H) FACS (E, n=6), ImageStream cytometry (F, G, scale bar = 10 μm) or conventional bright-field microscopy (H, n=2, scale bar = 10 μm) of PMNs incubated for 60 min with RNA-AF647 complexed with LL37 (n=6). In F the % of cells with 'internalized' features are shown (n=2, see Methods), in G selected cells from one representative donor are shown. In H unmodified LL37 was replaced with Atto488-labeled LL37, one representative donor shown. (I) as in C but including chloroquine (CQ) pre-incubation for 30 min (n=6-7). PMN activation by human mRNA (J, hRNA, n=4-6) and bacterial RNA (K, bRNA from *S. aureus,* n=4) with and without LL37. A-C, E, F, I-K represent combined data (mean+SD) from 'n' biological replicates (each dot represents one donor). In D, G and H one representative of 'n' biological replicates (donors) is shown (mean+SD). * p<0.05 according to one-way ANOVA with Dunnett's correction (A, B, C, I, J) or Friedman test with Dunn's correction (E, K).

Figure 2: RNA-LL37 complexes promote release of multiple pro-inflammatory cytokines and chemokines, especially in psoriasis PMN

Cytometric bead array (A-C) or ELISA (D, E) for TNF (A), IL-6 (B), IL-1β (C), IL-16 (D) or MIP-1β (E) secreted from PMNs stimulated for 4 h as indicated (n=6). Flow cytometric cell counts of migrated CD4 T cells (F, I), CD8 T cells (G) and CD14$^+$HLA-DR$^+$ monocytes (H) quantified in transwell assays with total PBMCs in the upper and either MIP-1β (30 and 150 pg/ml) or IL-16 (300 and 1500 pg/ml) (F-H, n=6-7, p>0.05 for treatments vs. media) or R848, RNA with and without LL37 (I, n=3-7) in the lower chamber. (J, K) ELISA of IL-8 and MIP-1β secreted from psoriasis PMNs (n=3 patients, , squares, chequered bars) or PMNs from sex-and age-matched healthy donors (n=10, circles, filled bars). (L) as in J or K but ELISA for LL37 (n=4 patients, 7 healthy donors). A-K represent combined data (mean+SD) from 'n' biological replicates (each dot represents one donor). * p<0.05 according to Friedmann test with Dunn's correction (A, B, D-H), or one-way ANOVA with Dunnett's correction for multiple testing (C, I, J-L).

Figure 3: RNA-LL37 trigger the release of NETs containing further RNA, DNA and LL37

(A) EM pictures from PMNs stimulated with RNA-LL37 (n=2). (B) Neutrophil elastase (NE) release from PMNs stimulated for 3 h (n=8, each dot representing one donor). Fluorescence microscopy of fixed and Hoechst, anti-LL37, SYTO RNAselect- (C) and/or anti-ΨU-stained (D, with or without RNase

A treatment) or live (E) PMNs stimulated as indicated (n=6, scale bar = 10 µm or n=4, scale bar = 20

µm). (F) Quantification of live cell microscopy), see also Movies S1-S3. (G, H) Skin sections from

healthy (G) or psoriasis- (G, H) affected skin (n=12 patients and 3 healthy controls, scale bar = 20 µm).

(I, J) as in C (left panels) and stimulated with transferred NETs (right panels). AF = autofluorescence.

B, F and I represent combined data (mean+SD) from 'n' biological replicates. In A, C-E, H and J

representative samples of 'n' replicates or donors are shown. Arrowheads indicate released

RNA/NETs (C-E) or RNA-LL37 co-localization (H). * p<0.05 according to one-way ANOVA (B), two-way

ANOVA (F) with Dunnett's correction or Kruskal-Wallis test with Dunn's correction (I).

Figure 4: RNA-LL37 effects in primary PMNs are mediated by TLR8/TLR13 and can be inhibited by

Inhibitory oligodeoxynucleotides

(A) Tnf release (ELISA, 5 h stimulation as indicated) by BM-PMNs from different mouse strains (n=4-8

each). (B, C) Fluorescence microscopy of fixed, Hoechst and SYTO RNAselect-stained stimulated BM-

PMNs (n=4 $Tlr13^{-/-}$, n=5 $Unc93b1^{3d/3d}$, n=8 WT, scale bar = 10 or 20 µm, respectively). (D) Relative IL-

8release (ELISA, 18 h stimulation as indicated) by WT and $TLR8$ CRISPR-edited BlaER1 cells,

normalized to $E.\ coli$ control (n=4). (E) NF-κB dual luciferase reporter assay in HEK293T cells,

transfected with NF-κB firefly luciferase reporter, $Renilla$ control reporter, TLR8 plasmid or empty

vector (EV), and subsequently stimulated with RNA without (arrow) or with IRS661, IRS954, IRS869

and IRS546 (0.15-3.5 µM, n=2 each). IL-8 (F, n=6) or MIP-1β (G, n=4) release from stimulated PMNs

with or without IRS661 (1 nM) or IRS954 (50 nM) pre-incubation (30 min) quantified by ELISA. (H)

Fluorescence microscopy of fixed Hoechst and anti-ΨU-stained RNA-LL37-stimulated PMNs (n=3,

scale bar = 10 µm). (I) Quantification of H. A, D- G and I represent combined data (mean+SD) from 'n'

biological replicates (each dot represents one mouse, donor or measurement). In B, C, E and H one

representative of 'n' replicates is shown (mean+SD of technical triplicates). * p<0.05 according to

two-way ANOVA with Dunnett correction for multiple testing (E, comparison against 'no inhibitor'

condition, arrow), one-way ANOVA with Sidak correction (A, D, F), Friedmann test with Dunn

correction (G) or Mann-Whitney test (I).

Figure 1

Figure 2

Figure 3

Figure 4