Profiles of cooperative brains: A discriminant analysis of cleaner and client fish monoaminergic responses to different social contexts

Caio Maximino[1*], Ana Cristina R. Gomes[2*], Murilo S. Abreu[3], Sónia C Cardoso[2], Monica Lima-Maximino[4], Svante Winberg[5], Marta C. Soares[2]

*shared first authorship

[1] IESB, Instituto de Estudos em Saúde e Biológicas, Universidade Federal do Sul e Sudeste do Pará, Unidade III, Marabá, Brazil

[2] CIBIO, Centro de Investigação em Biodiversidade e Recursos Genéticos, Universidade do Porto, Campus Agrário de Vairão, 4485-661 Vairão, Portugal.

[3] Bioscience Institute, University of Passo Fundo (UPF), Passo Fundo, RS, Brazil.

[4] DMCF, Departamento de Morfologia e Ciências Fisiológicas, Universidade do Estado do Pará, Marabá, Brazil

[5] Department of Neuroscience, Uppsala University, Box 593, Husargatan 3, 75124, Uppsala, Sweden

Abstract

Vertebrate cognitive function requires a dynamic coordination of multiple specialized areas of the brain. The challenge here is to understand how these brain areas respond in dependence to the neurophysiological mechanisms in place, as to enable the successful processing of information. For instance, social and cooperative behaviour has been linked to the activation of some specific brain areas, mostly associated with reward processing, however mechanisms could in principle change, in accordance to species social demands and taxon involved. Here we chose a classic model system of cooperation between species of fish and compared the brain monoaminergic response of cleaners *Labroides dimidiatus* exposed to several social-related treatments to the response of one client species (*Naso elegans)* introduced to similar contexts. We demonstrate that the variable appraisal of each social challenge contributes to brain dopaminergic and serotonergic changes, in cleaners and clients, with both showing the diencephalon and optic tectum as main areas of metabolite response. However, the role of the serotoninergic system activation was mostly demonstrated at the diencephalon and cerebellum of cleaners, a response that was driven by the exposure to clients. Our current results are the first to jointly demonstrate the level of selective similarity in brain monoaminergic mechanisms that underlie fish mutualistic and social engagement, for both sides of these partnerships.

Key-words:

1. **Introduction**

The neural substrate of cooperative behaviour is gaining increasing attention in the latter years, due to the appearance of novel techniques to study neurochemistry and neurophysiology as well as to the recognition that cooperation is widespread in nature [1]. In recent years, the focus on fish species as model organisms to study cooperative behaviour increased [2].

Cleaner wrasses (*Labroides dimidiatus*) are mutualistic fish that engage in cooperative behaviour, inspecting the surface, gills and sometimes the mouth of 'client' reef fish, eating ectoparasites, mucus, scales and dead or infected tissue [3]. These cleaning mutualisms involve repeated iterations with multiple partners (referred to as clients) that have asymmetrical options of behaving cooperatively or to defect [2]. Cleaners display flexibility in adjusting their cleaning services in accordance with clients, varying between "cooperating" (removing ectoparasites) or "defecting" (eating fish mucus) [3]. Cleaners adjust their behaviour towards clients by changing cheating frequency and/or providing a form of physical contact with clients ("tactile stimulation"), which is thought to "appease" the client after a cheating event, to prolong interactions with clients about to leave the cleaning station, or to act as a pre-conflict management strategy with predators [4–6].

There is limited evidence for a role of monoamines on cooperative behaviour. In healthy humans, tryptophan depletion reduced the level of cooperation shown by participants in the first day (but not the second day) of an iterared Prisoner's dillema game, and a diminished probability of cooperative responding given previous mutually cooperative behaviour [7]. In cleaner wrasses, fluoxetine or the serotonin (5-HT) 1A receptor agonist 8-OH-DPAT increased motivation to engage in cleaning behaviour and the frequency of tactile stimulation, while blocking serotonin synthesis with pCPA or

antagonizing the 5-HT$_{1A}$ receptor with WAY 100,635 decreased cleaner's cheating levels [8]□. Blocking dopamine (DA) D1 receptors increases tactile stimulation and induces cleaners to initiate more interactions with clients [9]□; these effects appear to be mediated by the degree of familiarity with the client, with D1 antagonists increasing tactile stimulation only when interacting with novel clients [10]□. It has been suggested [2] that dopamine is tracking predicted outcomes, with reductions in DA signalling worse outcomes than predicted (e.g., lower probability of getting food during an interaction, or higher likelihood of being punished). Interestingly, blocking D1 receptors in cleaners which were trained to wait for a larger reward improves learning [11], while blocking 5-HT$_{1A}$ receptors impair learning of the task [12]□, suggesting that 5-HT and DA have opponent roles in tracking outcomes.

A relevant question regarding the role of monoamines in this cooperative behaviour regards the importance of physical contact. Monoaminergic signalling has been proposed to mediate social behaviour in fish, with increases in dopamine after visual contact with conspecifics in zebrafish [13]. For a cleaning mutualism to take place in fish, however, physical contact is needed. Monoamines could act at various points of the cleaning sequence: client evaluation, motivation to approach the client, maintenance of cleaning behaviour after initial contact, and during inspection, choosing whether to clean or to cheat. There is some evidence that both DA and 5-HT participate in client evaluation [12]. Dopamine, for example, is the main neurotransmitter in the mesolimbic reward system [14], which interacts with the social behaviour network as to reinforce responses to salient social stimuli [15], which could include the motivation to cooperate. Here we test the hypothesis that monoamines are also associated with the motivation to approach the client by presenting cleaner wrasse *Labroides dimidiatus* with either a novel, non-living stimulus, a conspecific with only visual access, a

conspecific with both visual and tactile access, a client with visual (but not tactile) access, and a client with both visual and tactile access, and measuring levels of DA and 5-HT, as well as their metabolites, in the forebrain, diencephalon, optic tectum, cerebellum, and brain stem. We predict that, if monoamines are related to the motivation to approach the client, their increase should be related to presentation of heterospecifics vs. conspecifics, but not visual vs. tactile access.

A secondary aim refers to neurochemical changes in the client brain. Clients are not passive receivers in a cleaning mutualism, and indeed may react to cheating by cheating in return (if they are piscivores), switch between cleaner stations, or by punishing the cheating cleaner [16–19]. Indeed, cleaners can establish mutualisms with herbivore fishes which can reciprocate by chasing cheaters or swimming away, or with piscivores which can, in addition to those strategies, "cheat back" by eating the cleaner in a subsequent encounter [3]. In an encounter with herbivore clients, then, the cooperation is asymmetric [17]. If monoamines participate in this relationship, it is to be expected that these neurotransmitters also modulate client behaviour. In that case, monoamines could act at the approach motivation or as a "reward signal" as well, or they could be produced in response to the reception of tactile stimulation (which decreases cortisol in clients [6]). Our second hypothesis is that monoamines are also associated with the motivation to approach the cleaner by presenting Naso tang *Naso elegans* with either a novel, non-living stimulus, a conspecific with both visual and tactile access, and a client with both visual and tactile access, and measuring monoamines in the same brain regions as in cleaners. If monoamines are thus related to the motivation to approach the client, their increase should be related to presentation of heterospecifics vs. conspecifics, but not to a novel stimulus.

Material and Methods

2.1.   Experimental conditions

Experiments were conducted at the fish housing facilities of the Oceanário de Lisboa (Lisbon, Portugal). In total, 53 Indo-Pacific bluestreak cleaner wrasses, *Labroides dimidiatus* and 18 adult blond naso tang *Naso elegans* (family Acanthuridae; from now on referred to as "clients") were used. For similar methodologies please see [20–22].

A) Experiment 1: cleaners

Cleaners were exposed to five different contextual treatments. Indeed, on each experimental day, one of the five following treatments was randomly allocated to each subject cleaner wrasse aquarium: a) conspecific (*L. dimidiatus*, n = 10), b) client (*N. elegans*, n = 11), c) conspecific inside another smaller aquarium (n = 12), d) client inside another smaller aquarium (n = 10) and e) ball (white ball, about 5 cm in diameter, which stayed at the bottom, completely sessile, n = 10), see Fig 1. Cleaners were exposed to treatments for the next 60 minutes. At the end of experiments, each subject cleaner wrasse was captured and immediately sacrificed with an overdose of tricaine solution (MS222, Pharmaq; 500–1000 mg/L) and the spinal cord was sectioned, the period between anaesthesia and the spinal cord section does not exceed 2 min, no change of monoamines levels. The brain was immediately dissected under a stereoscope (Zeiss; Stemi 2000) into five macro-areas: forebrain (FB - which included the olfactory bulbs and the telencephalon), diencephalon (DL), optic tectum (OT), cerebellum (CB) and brain stem (BS) (for a drawing of the major brain areas, please see [20–22]. Major brain areas were frozen with dry ice and the stored at - 80°C.

B) Experiment 2: Clients

Three different contextual treatments were allocated to our subject clients (*N. elegans*): a) conspecific (*N. elegans*, n =7), b) a cleaner (*L. dimidiatus*, n = 7) and c) ball (same as above, n = 4) see Fig 1. Clients were distributed across experimental treatments as follows: 7 clients for the sympatric cleaner group, 7 clients for the conspecific group and 4 for the ball group. Each client was introduced inside the experimental tank and left for at least 2-5 min until normal behaviour was restored. Cleaners were exposed to treatments for the next 60 minutes. At the end of experiments, each subject client was captured and immediately sacrificed and brain micro-dissected as explained above (experiment 1).

2.2. Quantification of monoamines by high performance liquid chromatography with electrochemical detection (HPLC-EC)

High performance liquid chromatography with electrochemical detection (HPLC-EC) was used to analyze the monoamines dopamine (DA) and serotonin (5-HT, 5-hydroxytryptamine), the DA metabolite DOPAC (3,4-dihydroxyphenylacetic acid), the 5-HT metabolite 5-HIAA (5-hydroxy indole acetic acid), as described by [23] (for the complete protocol used in both experiments please see [21,22]). The ratios of 5-HIAA/5-HT and DOPAC/DA were calculated and used as an index of serotonergic and dopaminergic activity, respectively. For normalization of brain monoamine levels, brain protein weights were determined with Bicinchoninic acid protein determination (Sigma–Aldrich, Sweden) according to the manufacturer's instructions. The assay was read on Labsystems multiskan 352 plate reader (Labsystems, Thermo Fisher Scientific) wavelength of 570 nm.

2.3. Statistical analysis

We used multivariate discriminant analysis (MDA) to test for differences in the pattern of metabolite expression among the 5 social contexts for cleaners (with

conspecific, with client, conspecific inside another aquarium, client inside another

aquarium, and control), or the 3 social contexts for clients (with conspecific, with

cleaner, and control). We focused our measures to the dopaminergic system and

measure levels of dopamine and those of DOPAC (3,4- Dihydroxyphenylacetic acid), a

dopamine's metabolite and also levels of serotonin (5-hydroxytryptamine, 5-HT), and

serotonin's metabolite 5HIAA (5-Hydroxyindoleacetic acid). One set of MDA uses as

data the concentrations of each monoamine (DA and 5-HT) and metabolite (DOPAC

and 5-HIAA) at each brain area. This allows us to find the combination of metabolites

and brain areas that discriminate the most the different social contexts. Another set of

MDA uses as data the ratios of DOPAC/DA and 5-HT/5-HIAA as these will enable us to

interpret the relation between monoaminergic production and its subsequent metabolic

use. We interpret the two principal discriminant functions (DF1 and DF2) of each MDA

based on trait loadings higher than 0.40 [24]. We assessed the significance of the

discrimination within the entire set of social contexts using Wilk's Lambda, and we

used ANOVAs with post-hoc Tukey HSD tests to find which pairs of social context

differ significantly in DF1 or DF2. All values were standardized using script available

[25] and normality was checked through q-q plots. All analyses were run with R

(version 3.3.3; R Core Team, 2017) package car (version 2.1-4; [26] and MASS (version

7.3-47; [27]).

2.4. Ethical note

The protocols were carried out in accordance to the approved guidelines by the

Oceanário de Lisboa (fish housing facilities), where the experiments were then

developed. Animal procedures used in this study were also approved by the Portuguese

Veterinary Office (Direccao Geral de Veterinaria, license # 0420/000/000/2009).

2.5. Open science practices

The analyses performed were not formally preregistered. Data for this manuscript can be found at our GitHub repository (https://github.com/lanec-unifesspa/cleaners).

3. **Results**

3.1) Experiment 1: cleaners

Overall, social contexts differed in the levels of metabolite expression (Wilk's lambda = 0.03, p = 0.001; Fig.2A) and in metabolite ratios (Wilk's lambda = 0.23, p = 0.02; Fig.2 B) across the brain tissues of cleaners.

For metabolite levels, DF1 and DF2 explained 83.9% of the variation between all the contexts. DF1 differed significantly between most pairs of contexts, except for the comparison between client vs. conspecific inside another aquarium, which instead were found to differ in DF2 (Fig. 2A, Table 1). DF1 was characterized mostly by strong negative loadings of DOPAC at the brain stem, and of DA, 5-HIAA and 5-HT at the diencephalon of cleaners (loadings >-0.40; all other loadings <|0.29|; Fig. 2C and Table 2). DF2 was characterized mostly by a strong positive loading of DA at the optic tectum (loading = 0.58; all other loadings <|0.36|; Fig. 2C and Table 2).

When using instead ratios of metabolite expression in the brain tissues of cleaners, DF1 and DF2 together explained 75% of the variation among contexts. DF1 differed significantly between cleaners that were prevented to interact with clients (inside a smaller aquarium) from those that could engage freely with clients (Fig 1B; Table 1). DF1 also differed between fully interactive contexts (i.e. interaction with clients) and either controls or interaction with conspecifics, and between clients in

separate aquaria and conspecifics in separate aquaria (Table 1). DF2 differed only

between interaction with clients and either conspecifics in separate aquaria or

conspecifics (Table 1). DF1 was characterized mostly by loadings of 5-HIAA/5-HT

ratios (loadings = -0.51 at the diencephalon and 0.45 at the cerebellum; all other

loadings <|0.33|; Fig. 2D and Table 3). DF2 was characterized mostly by a strong

negative loading of 5-HIAA/5-HT ratio at the diencephalon and cerebelum (loading = -

0.50 and -0.41, respectively; all other loadings <|0.27|; Fig. 2D and Table 3).

3.2.)   Experiment 2: clients

Regarding clients, the exposure to different social contextual treatments was

solely found to differ significantly in the levels of metabolite expression (Wilk's lambda

= 0.01, p = 0.02), showing a clear separation between the individuals exposed to

different contexts (Fig. 3 A) but not for metabolite ratios (Wilk's lambda = 0.42 , p =

0.51 , Fig 3B; Table 4).

For monoaminergic metabolite production, DF1 and DF2 explained 99% of the

variation and differ significantly among all pairs of contexts, except for the context of a

cleaner with a client, which was not significant for DF2. DF1 seems to be explained by

two areas of client fish *N. elegans*: the diencephalon (varying in terms of DOPAC and

5-HIAA, in opposite directions) and the optic tectum (solely for DOPAC; Fig. 3, Table

2), while DF2 seems to be explained by the DOPAC on the TC (Fig. 3, Table 2).

Surprisingly, no differences were found in the overall ratio patterns (monoaminergic

ratios DOPAC/DA and 5-HIAA/5-HT; Wilk's lambda = 0.42, p = 0.51).

4. Discussion

Discriminant analysis is a powerful statistical tool that enabled us to determine

the contribution of social and mutualistic behaviour to the modulation of brain

monoaminergic levels of cleaners and clients, and specifically those linked to the

activation of some localized brain areas. We found that brain monoaminergic response

to social challenges shared interesting similarities for cleaners and clients,

distinguishing broadly between stimuli that fish could physically interact with, from all

those that were either physically inaccessible (stimuli inside aquaria) or not able to

respond (sessile, non-living stimulus), with most relevant alterations occurring at the

diencephalon and optic tectum, in reference to DAergic and 5-HTergic monoaminergic

change. And while client fish clearly responded to differences posed by conspecifics

compared to heterospecifics, we discovered that it was cleaners' 5-HT system that

mostly contributed to separate mutualistic activities from all others, changing crucially

whenever these individuals were given the possibility to physically engage in cleaning

(occurring at the cerebellum) or were prevented to interact (occurring at the

diencephalon) (but see also [22]).

Broad visual analysis of cleaners' monoaminergic response in respect to all different

treatments suggests the distinction of 3 groups: a first with inaccessible clients and

control, a second composed by inaccessible conspecifics and a third that included

conspecifics and heterospecifics that were fully accessible. Specifically, the DF1 axis

represented most of the variation observed, being negatively associated with DA, 5-HT

and 5-HIAA at the diencephalon and DOPAC at the brain stem; while the DF2 mostly

distinguished those stimuli that cleaners could physically access from those they could

not and was positively associated with DA levels at the optic tectum. It is usually

considered that monoaminergic increase (in this case, DA or 5-HT) means an increment

in production while the elevation of metabolite levels (such as DOPAC or 5-HIAA)

signals for the actual system activation and monoamine usage (see [13]). One potential explanation for the positive response of the DAergic system at the optic tectum is cleaners' visual stimulation, while the DOPAC at the brain stem should be related to an increase of motor activity. The overall decrease of monoamine levels, particularly those concerning the 5-HT system, may mostly occur in response to the inability to contact clients, as the opposite occurs following the exogenous treatment with serotonin activity antagonists (PCPA and WAY) which produced a general decrease in interspecific activities (but not the conspecific ones, see [8]). Interestingly, the exogenous infusion of DAergic antagonists is also known to produce a decrease to the actual foraging (aka cleaning) of cleaners [9], with these behaving as if cleaning was not secured, which would be similar to the inability to interact produced in two of our treatment groups, being that correlated to a decrease of DA and DOPAC. And while there was a clear distinction between the cleaners that could freely contact with stimuli and those that were unable (increment of DA levels at the optic tectum), it was solely by looking at the metabolite/neurotransmitter ratios that the differences between being able to clean or not, were finally disclosed: the DF2 axis distinguishing between mutualistic and non-mutualistic activities and the DF1 discriminating between the actual physical contact, of those that were able to fully inspect a client (positive association with 5-HIAA/5-HT at the cerebellum) and those that could not engage in cleaning (negative association with 5-HIAA/5-HT at the diencephalon). These points for an extra expenditure of 5-HTergic metabolites at the diencephalon whenever cleaners fail to achieve their needs and an increase of availability at the cerebellum during mutualistic activities.

Clients brain monoaminergic levels was also tested in respect to 3 main social challenges, revealing interesting similarities but also some relevant dissimilarities with cleaners' overall response; for instance, all 3 groups clearly discriminated, with the DF1

axis again representing the majority of the variation observed, being negatively

associated with 5-HIAA at the diencephalon and DOPAC at the optic tectum but

positively with DOPAC at the diencephalon; while the DF2 mostly distinguishing living

from non-living stimuli and positively associated with DA levels at the diencephalon.

Thus, clients seem to experience a general stimulation of the diencephalic DAergic

system at the diencephalon contrasting with the overall decrease function at the optic

tectum or brain stem in cleaners. One possible explanation is that clients' increase in DA

and DOPAC levels at the diencephalon is associated with positive social engagement

(when introduced with cleaners and other conspecifics), contrary to negative values

reported for diencephalons of zebrafish and matrinxã (*Brycon amazonicus)* when

subject to agonistic contexts [28,29]. The diencephalon is also associated with the

modulation of the posterior tuberculum/hypothalamic 5-HT neuronal populations, but

however, usually give emphasis to rises of 5-HT activity, normally correlated with overt

aggression, for instance, in fish introduced to a real opponent situation [28] or following

the loss of a fight [28], while the event of winning a fight seems to contribute to a

reduction of 5-HIAA / 5-HT [29,30].

Teleost fish (and other vertebrate groups) cognitive function requires a dynamic

coordination of multiple specialized areas of the brain. Here we aimed to understand

how these brain areas responded in dependence to the monoaminergic mechanisms in

place, as to enable the successful processing of social and mutualistic information. We

demonstrate that the variable appraisal of each social challenge contributes to brain

dopaminergic and serotonergic changes, in cleaners and clients, with both showing the

diencephalon and optic tectum as main areas of metabolite response. However, the role

of the serotoninergic system activation was mostly demonstrated at the diencephalon

and cerebellum of cleaners, a response that was driven by the exposure to mutualistic

stimuli. Our current results are the first to jointly demonstrate the level of selective

similarity in brain monoaminergic mechanisms that underlie fish mutualistic and social

engagement, for both sides of these partnerships.

Author contributions

M.C.S. conceived the study. A.C.R.G. performed statistical analysis. All authors

participated in writing and provided comments and criticism and gave final approval for

publication. The authors declare no competing interests.

Acknowledgements

We thank Gonçalo Cardoso for helpful discussions during the statistical analysis

and comments on earlier drafts of this manuscript.

Funding

Data collection was supported by the Portuguese Foundation for Science and

Technology-FCT (grant PTDC/MAR/105276/2008 given to M.C.S.). A.C.R.G. is

currently supported by SFRH/BD/129002/2017 and M.C.S. by FRH/BPD/109433/2015.

S.W. lab is supported by the Swedish research council (VR) and the Swedish research

council FORMAS.

References

1. Dugatkin LA. 1997 The evolution of cooperation. *Bioscience* **47**, 355–362.

2. Soares MC, Cardoso SC, Carvalho T dos S, Maximino C. 2018 Using model fish to study the biological mechanisms of cooperative behaviour: A future for translational research concerning social anxiety disorders? *Progress in Neuro-Psychopharmacology and Biological Psychiatry* **82**, 205–215. (doi:https://doi.org/10.1016/j.pnpbp.2017.11.014)

3. Bshary R, Côté I. 2008 New perspectives on marine cleaning mutualism. In *Fish behaviour*, Enfield: Science Publishers.

4. Bshary R, Würth M. 2001 Cleaner fish Labroides dimidiatus manipulate client reef fish by providing tactile stimulation. *Proceedings of the Royal Society of London. Series B: Biological Sciences* **268**, 1495.

5. Grutter AS. 2004 Cleaner Fish Use Tactile Dancing Behavior as a Preconflict Management Strategy. *Current Biology* **14**, 1080–1083. (doi:10.1016/j.cub.2004.05.048)

6. Soares MC, Oliveira RF, Ros AFH, Grutter AS, Bshary R. 2011 Tactile stimulation lowers stress in fish. *Nature Communications* **2**, 534. (doi:10.1038/ncomms1547 https://www.nature.com/articles/ncomms1547#supplementary-information)

7. Wood RM, Rilling JK, Sanfey AG, Bhagwagar Z, Rogers RD. 2006 Effects of Tryptophan Depletion on the Performance of an Iterated Prisoner's Dilemma Game in Healthy Adults. *Neuropsychopharmacology* **31**, 1075. (doi:10.1038/sj.npp.1300932)

8. Paula JR, Messias JP, Grutter AS, Bshary R, Soares MC. 2015 The role of serotonin in the modulation of cooperative behavior. *Behavioral Ecology* **26**, 1005–1012. (doi:10.1093/beheco/arv039)

9. Messias JPM, Paula JR, Grutter AS, Bshary R, Soares MC. 2016 Dopamine disruption increases negotiation for cooperative interactions in a fish. *Scientific Reports* **6**, 20817. (doi:10.1038/srep20817)

10. Soares MC, Santos TP, Messias JPM. 2017 Dopamine disruption increases cleanerfish cooperative investment in novel client partners. *Royal Society Open Science* **4**.

11. Messias JPM, Santos TP, Pinto M, Soares MC. 2016 Stimulation of dopamine D(1) receptor improves learning capacity in cooperating cleaner fish. *Proceedings of the Royal Society B: Biological Sciences* **283**, 20152272. (doi:10.1098/rspb.2015.2272)

12. Soares MC, Paula JR, Bshary R. 2016 Serotonin blockade delays learning performance in a cooperative fish. *Animal Cognition* **19**, 1027–1030. (doi:10.1007/s10071-016-0988-z)

13. Saif M, Chatterjee D, Buske C, Gerlai R. 2013 Sight of conspecific images induces changes in neurochemistry in zebrafish. *Behavioural Brain Research* **243**, 294–299. (doi:https://doi.org/10.1016/j.bbr.2013.01.020)

14. Alcaro A, Huber R, Panksepp J. 2007 Behavioral functions of the mesolimbic dopaminergic system: an affective neuroethological perspective. *Brain research reviews* **56**, 283–321. (doi:10.1016/j.brainresrev.2007.07.014)

15. O'Connell LA, Hofmann HA. 2011 The Vertebrate mesolimbic reward system and social behavior network: A comparative synthesis. *The Journal of Comparative Neurology* **519**, 3599–3639. (doi:10.1002/cne.22735)

16. Bshary R, Grutter AS. 2002 Asymmetric cheating opportunities and partner

control in a cleaner fish mutualism. *Animal Behaviour* **63**, 547–555. (doi:https://doi.org/10.1006/anbe.2001.1937)

17. Bshary R. 2002 Building up relationships in asymmetric co-operation games between the cleaner wrasse Labroides dimidiatus and client reef fish. *Behavioral Ecology and Sociobiology* **52**, 365–371. (doi:10.1007/s00265-002-0527-6)

18. Grutter AS, Bshary R. 2003 Cleaner wrasse prefer client mucus: support for partner control mechanisms in cleaning interactions. *Proceedings of the Royal Society of London. Series B: Biological Sciences* **270**, S242.

19. Bshary R, Grutter AS. 2005 Punishment and partner switching cause cooperative behaviour in a cleaning mutualism. *Biology Letters* **1**, 396–399. (doi:10.1098/rsbl.2005.0344)

20. Soares MC, Cardoso SC, Mazzei R, André GI, Morais M, Gozdowska M, Kalamarz-Kubiak H, Kulczykowska E. 2017 Region specific changes in nonapeptide levels during client fish interactions with allopatric and sympatric cleaner fish. *PLoS One* **12**, e0180290. (doi:10.1371/journal.pone.0180290)

21. Abreu MS, Messias JPM, Thörnqvist PO, Winberg S, Soares MC. 2018 Monoaminergic levels at the forebrain and diencephalon signal for the occurrence of mutualistic and conspecific engagement in client reef fish. *Sci Rep* **8**, 7346. (doi:10.1038/s41598-018-25513-6)

22. Abreu MS, Messias JPM, Thörnqvist PO, Winberg S, Soares MC. 2018 The variable monoaminergic outcomes of cleaner fish' brains when facing different social and mutualistic contexts. *PeerJ*

23. Øverli Ø, Harris CA, Winberg S. 1999 Short-Term Effects of Fights for Social Dominance and the Establishment of Dominant-Subordinate Relationships on Brain Monoamines and Cortisol in Rainbow Trout. *Brain, Behavior and Evolution* **54**, 263–275.

24. Hair JF. 2010 Black, WC, Babin, BJ, & Anderson, RE (2010). *Multivariate data analysis* **7**.

25. Coghlan A. 2013 A little book of R for multivariate analysis. *Wellcome Trust Sanger Institute, Cambridge, UK* , 1–47.

26. Fox J, Weisberg S. 2011 *An R Companion to Applied Regression*. SAGE Publications. See https://books.google.com.br/books?id=YH6NotdvzF0C.

27. Venables WN, Ripley BD. 2010 *Modern Applied Statistics with S*. Springer New York. See https://books.google.com.br/books?id=uj5oewAACAAJ.

28. Teles MC, Dahlbom SJ, Winberg S, Oliveira RF. 2013 Social modulation of brain monoamine levels in zebrafish. *Behav Brain Res* **253**, 17–24. (doi:10.1016/j.bbr.2013.07.012)

29. Wolkers CP, Serra M, Urbinati EC. 2015 Social challenge increases cortisol and hypothalamic monoamine levels in matrinxã (Brycon amazonicus). *Fish Physiol Biochem* **41**, 1501–1508. (doi:10.1007/s10695-015-0102-5)

30. Winberg S, Thörnqvist P-O. 2016 Role of brain serotonin in modulating fish behavior. *Current Zoology* **62**, 317–323. (doi:10.1093/cz/zow037)

433   Figures

434   Figure 1. Experimental Schematics

435
436   Figure 2. Cleaner's discriminant analysis plot of DF1 versus DF2 to visualize how the

437   discriminant functions discriminate between contextual treatments, for both

438   monoaminergic metabolites and monoaminergic ratios, synthesized at each brain area.

439   Bigger symbols with dotted drop line in the linear discriminant plot indicate centroid of

440   each group. DF1 and DF2 explained variation is, respectively, for monoaminergic

441   metabolites analysis (A) 63.8% and 20.1% and for monoaminergic ratios (B), 50.8%

442   and 25.1%. (C) and (D), Trait loadings of each variable on DF1 and DF2; the most

443   relevant metabolites to interpret DF1 or DF2 (trait loadings > 0.40) are shown in green.

444   See the methods section for acronyms.

446   Figure 3. Clients' discriminant analysis plot of DF1 versus DF2 to visualize how the

447   discriminant functions discriminate between contextual treatments, for both

448   monoaminergic metabolites and monoaminergic ratios, synthesized at each brain area.

449   Bigger symbols with dotted drop line in the linear discriminant plot indicate centroid of

450   each group. DF1 and DF2 explained variation is, respectively, for monoaminergic

451   metabolites analysis (A), 87.4% and 12.6% and for monoaminergic ratios (B), 70.2%

452   and 29.8%. In (C) and (D), relevant metabolites (correlational coefficient > 0.40) are

453   shown in green. See the methods section for acronyms.

Tables

Table 1. Discriminant model significance and pairwise combinations of contextual groups' differences that are significantly explained by each of the discriminant functions, for cleaners.

	Monoamine		Monoamine ratio	
	DF1	DF2	DF1	DF2
Conspecific vs. Control	0.02*	6.85E-06*	1.00	0.75
Conspecific inside aquarium vs. Control	5.07E-11*	0.98	0.63	0.34
Heterospecific vs. Control	2.92E-07*	<0.01*	<0.01*	0.42
Heterospecific inside aquarium vs. Control	0.02*	0.37	0.11	0.51
Conspecific inside aquarium vs. Conspecific	2.47E-05*	5.55E-07*	0.58	0.99
Heterospecific vs. Conspecific	0.03*	0.65	<0.01*	0.05*
Heterospecific inside aquarium vs. Conspecific	1.48E-06*	<0.01*	0.21	0.07
Heterospecific vs. Conspecific inside aquarium	0.16	2.23E-05*	0.08	<0.01*
Heterospecific inside aquarium vs. Conspecific inside aquarium	2.49E-13*	0.12	<0.01*	0.01
Heterospecific inside	7.95E-12*	0.05	8.49E-07*	1.00

aquarium vs. **Heterospecific**			

Table 2. Coefficients of each monoaminergic metabolites, synthesized at each brain

area, for both clients and cleaners. DF1 and DF2 explained variation is, respectively,

87.4% and 12.6% for clients, and 63.8% and 20.1% for cleaners. * important

metabolites (correlational coefficient > 0.40).

	Cleaners		Clients	
	DF1	**DF2**	**DF1**	**DF2**
BS_DOPAC	*-0.419	0.137	-0.321	0.220
BS_5-HIAA	0.185	-0.110	-0.147	0.011
BS_DA	0.196	0.364	-0.219	0.116
BS_5-HT	0.233	0.102	0.053	0.180
DL_DOPAC	-0.087	-0.257	*0.427	0.062
DL_5-HIAA	*-0.536	0.116	*-0.401	0.085
DL_DA	*-0.503	0.022	-0.214	*0.503
DL_5-HT	*-0.588	0.167	-0.154	0.396
FB_DOPAC	-0.153	-0.305	-0.363	-0.107
FB_5-HIAA	0.167	0.212	-0.228	-0.329
FB_DA	-0.068	-0.272	-0.267	-0.081
FB_5-HT	-0.276	0.190	0.166	-0.302
OT_DOPAC	0.045	-0.260	*-0.444	0.117
OT_5-HIAA	0.187	-0.012	-0.041	0.007
OT_DA	-0.165	*0.582	-0.214	0.051

OT_5-HT	0.075	0.233	0.011	-0.211
CB_DOPAC	0.067	-0.316	-	-
CB_5-HIAA	-0.201	-0.235	-	-
CB_DA	0.050	-0.151	-	-
CB_5-HT	-0.292	-0.183	-	-

464
465
466   Table 3. Coefficients of each monoaminergic ratios, synthesized at each brain area, for

467   both clients and cleaners. DF1 and DF2 explained variation is, respectively, 70.2% and

468   29.8% for clients, and 50.8% and 25.1% for cleaners. * important metabolites

469   (correlational coefficient > 0.40).

	Cleaners		Clients	
	DF1	**DF2**	**DF1**	**DF2**
BS_DOPAC/DA	-0.332	-0.276	-0.081	-0.033
BS_5-HIAA/5-HT	-0.182	-0.252	-0.341	0.310
DL_DOPAC/DA	0.324	-0.166	0.710	-0.269
DL_5-HIAA/5-HT	*-0.514	*-0.495	-0.138	-0.514
FB_DOPAC/DA	-0.165	0.265	0.136	-0.357
FB_5-HIAA/5-HT	0.227	0.233	-0.551	-0.012
OT_DOPAC/DA	0.077	0.220	-0.278	0.380
OT_5-HIAA/5-HT	0.252	-0.156	0.309	0.526
CB_DOPAC/DA	0.309	-0.210	-	-
CB_5-HIAA/5-HT	*0.446	*-0.413	-	-

470
471
472

473   Table 4. Discriminant model significance and pairwise combinations of contextual

474   groups' differences that are significantly explained by each of the discriminant

475   functions, for clients.

| | Monoamine | | Monoamine ratio |
	DF1	DF2	DF1 and DF2
Heterospecific vs. Control	4.67E-11*	6.26E-06*	-
Conspecific vs. Control	2.96E-05*	3.38E-07*	-
Conspecific vs. Heterospecific	4.72E-14*	0.29	-