Title: Alternative activation of macrophages is accompanied by chromatin remodeling

and short-term dampening of macrophage secondary response

Authors: Mei San Tang[1,6§], Emily R. Miraldi[2,3], Natasha M. Girgis[1], Richard A. Bonneau[4,5],

P'ng Loke[1§]

Affiliations:

[1]Department of Microbiology, New York University School of Medicine, New York, NY

10016, USA.

[2] Divisions of Immunobiology and Biomedical Informatics, Cincinnati Children's Hospital

Medical Center, Cincinnati, OH 45229, USA.

[3] Department of Pediatrics, University of Cincinnati College of Medicine, Cincinnati, OH,

45267, USA.

[4] Department of Biology, Center for Genomics and Systems Biology, New York University,

New York, NY 10003, USA.

[5] Simons Center for Data Analysis, Simons Foundation, New York, NY 10011, USA.

[6] Current address: Department of Pathology and Immunology, Washington University in St

Louis, MO 63110, USA.

[§]Correspondence to:
meisan@wustl.edu
Png.Loke@nyulangone.org
22

Abstract

Interleukin-4 (IL-4) activates macrophages to adopt a distinct phenotype associated with

clearance of helminth infections and tissue repair. Here, we describe changes in the accessible

chromatin landscape following IL-4 stimulation of terminally differentiated mouse peritoneal

macrophages. This chromatin remodeling process occurs in both tissue resident and

monocyte-derived macrophages, but the regions gaining accessibility post-stimulation are

macrophage-specific. PU.1 motif is similarly associated with tissue resident and monocyte-

derived IL-4 induced regions, but has macrophage-specific DNA shape and predicted co-

factors. In addition, IL-4 stimulation leads to short-term dampening of macrophage secondary

response. However, the degree of dampening differs between macrophages derived from

different genetic backgrounds. Together, these results lead us to propose that DNA sequence

variations can alter parts of the accessible chromatin landscape and differences in secondary

responses due to host genetics can contribute to phenotypic variations in immune responses.

Introduction

Macrophage activation is a process by which macrophages transition from a resting state to adopt different phenotypes, in response to specific external stimuli that can either be danger signals or homeostatic and metabolic signals (1). The macrophage activation process is accompanied by changes in transcriptional activities and histone modifications genome-wide, orchestrated by combinatorial actions of different transcription factors (TFs) that include lineage-determining TFs (such as PU.1) and stimulus-dependent TFs (such as the STAT and IRF proteins) (1-6). However, such molecular events have almost exclusively been described for bone marrow derived macrophages (BMDMs) in response to toll-like receptor (TLR) signaling, which is often used as a reductionist model to mimic type 1 immune response to acute infections that gives rise to classically activated macrophages.

In contrast, alternatively activated macrophages (AAMs) induced by type 2 cytokines, such as interleukin-4 (IL-4) and IL-13, adopt a distinct phenotype that can promote helminth expulsion and limit tissue damage during helminth infection (7-10). We have previously demonstrated that macrophages of tissue resident and monocytic origins are phenotypically different following IL-4 stimulation (11, 12). Here, we expand on these macrophage-specific differences by characterizing changes in accessible chromatin landscape following IL-4 stimulation of these different types of macrophages. The effects of IL-4 on BMDMs have also been documented, particularly highlighting the reduced response to interferon gamma after IL-4 stimulation that is mediated by the action of TFs such as the PPARγ:RXR heterodimer and STAT6 (13-15). However, the effects of IL-4 on the chromatin of macrophages from different cellular lineages *in vivo* have yet to be carefully investigated.

In vivo, chromatin accessibility changes have mostly been associated with the cellular differentiation process, but we find that IL-4 stimulation alone can give rise to new accessible regions in terminally differentiated peritoneal macrophages. These IL-4 induced regions are macrophage-specific, with differences in TF motifs. In addition, IL-4 stimulation *in vivo* leads to a short-term dampening of macrophage response to a repeat IL-4 stimulation. While there have been considerable number of studies that described the opposing effects of different

cytokines (16, 17), none has so far demonstrated that a single cytokine can dampen its own

response upon repeated stimulation. Notably, this dampening in response to the second IL-4

stimulation occurs to different degrees in C57BL/6 and BALB/c mouse strains. These

differences may be important for the outcome of helminth infections, since during

Litomosoides sigmodontis infection the more susceptible BALB/c mice undergo less tissue

resident macrophage expansion and more monocyte infiltration than the resistant C57BL/6

strain (10). Finally, because our knowledge of macrophage biology is largely based on the

C57BL/6 mouse strain, our study highlights the need to account for genetic diversity in

mouse immune models.

Results

IL-4 stimulation leads to remodeling of open chromatin landscape in peritoneal

macrophages.

To examine chromatin remodeling on different types of tissue macrophages, we

injected recombinant IL-4–antibody complex (IL-4c) into the peritoneal cavity of C57BL/6

mice to induce accumulation of alternatively activated $F4/80^{hi}CD206^-$ macrophages of

embryonic origin (AAM^{res}) and compared these with $F4/80^{int}CD206^+$ macrophages derived

from $Ly6C^{hi}$ inflammatory blood monocytes (AAM^{mono}) in mice injected with IL-4c and

thioglycollate (12). We then used ATAC-seq (18) to profile the open chromatin landscape of

these macrophages, in comparison to non-stimulated $F4/80^{hi}CD206^-$ macrophages of naïve

mice and $F4/80^{int}CD206^+$ macrophages from thioglycollate-treated mice (12).

The overall differences in accessible chromatin landscape (a total of 61,713 open

chromatin regions) could be attributed mainly to the type of macrophage (27% of total

variance), but alternative activation by IL-4 also altered the accessible chromatin profiles

(Figures 1A, B). *Arg1* and *Ucp1*, which are known to be IL-4 inducible (12), had

constitutively accessible chromatin regions, whereas *Retnla*, another IL-4 inducible gene, had

chromatin regions that gained accessibility in response to IL-4 (Figure 1B). This IL-4 induced

chromatin remodeling process can be cell-type-specific at certain regions (e.g. regions

adjacent to the loci of *Tgfb2*, *Ccl2*) (Figure 1B). Of the 61,713 total accessible regions, we

identified 1572 regions induced by IL-4 for AAM^{res} and 1462 regions for AAM^{mono} (Figure

S1A). IL-4-dependent regions also had the largest contribution to the differences in open

chromatin profiles between non-stimulated and IL-4 stimulated macrophages (Figure 1C).

The IL-4 induced regions almost all (99% in AAM^{res} and 97% in AAM^{mono}) gained

accessibility from undetectable levels at baseline (Figure 1D). We made comparisons of

several sequence characteristics between constitutively accessible and IL-4 induced regions

(Figure S1B). IL-4 induced regions were more likely to reside in non-coding intronic regions

– in tissue-resident macrophages, 781 of 1572 IL-4 induced peaks were intronic, as compared

to 2199 of 8061 constitutively accessible regions (binomial enrichment, p = 2.5×10^{-5}), while

in monocyte-derived macrophages, 761 of 1462 IL-4 induced peaks were intronic, as

compared to 5282 of 14045 constitutively accessible regions (binomial enrichment, p = $8.5 \times$

10^{-4}) (Figure 1E). In the monocyte-derived macrophages, IL-4 induced regions were overall

closer to IL-4 induced genes (two-sided Mann-Whitney test, p = 3.4×10^{-25} in AAM^{mono})

(Figure 1F). In tissue-resident macrophages, IL-4 induced regions contained lower GC

content (two-sided Mann-Whitney test, p = 8.6×10^{-46} in AAM^{res}) (Figure 1G, left) and were

also less likely to overlap with a CpG island (Figure 1G, right). Hence, IL-4 stimulation can

lead to reorganization of the chromatin landscape in terminally differentiated peritoneal

macrophages, giving rise to newly accessible regions that have distinct sequence properties

when compared to constitutively accessible regions.

IL-4 induced regions are associated with PU.1, KLF and AP-1 motifs.

Even though both AAM^{mono} and AAM^{res} received the same stimulation within the peritoneal

tissue environment, the regions that were remodeled by IL-4 were largely macrophage-

specific (Figure 2A). Of all the 2855 IL-4 induced regions, only 179 regions (6% of total IL-4

induced regions) were shared between both AAM^{mono} and AAM^{res} (Figure S1C). While the

IL-4 induced regions from AAM^{mono} and AAM^{res} were largely distinct, the DNA motifs

discovered from these distinct regions were grouped into similar families of TFs, which

included PU.1, KLF and the AP-1 family of motifs (Figure 2B). However, AAM^{mono} had

significantly higher number of accessible regions with the AP-1 motif (two-sided Fisher's

exact test, p = 9.5 × 10^{-10}), while AAM^{res} had significantly higher number of accessible

regions with the KLF motif (two-sided Fisher's exact test, p = 9.2 × 10^{-8}), suggesting the use

of different TFs by the two macrophage types during chromatin remodeling upon IL-4

induced alternative activation (Figure 2C). Such differences were not observed with the PU.1

motifs. We next examined the expression levels of these TF families (KLF vs. AP-1) to

determine if specific members within each family could be differentially expressed between

AAM^{res} and AAM^{mono}. 20 TFs of KLF and AP-1 families were highly expressed in peritoneal

macrophages (Figure S2A). These TFs almost all demonstrated lineage-specific expression,

both at baseline (Clusters 1 and 2) and with IL-4 stimulation (Clusters 3 and 4) (Figure 2D).

These results indicate that while KLF and AP-1 family of TFs may have lineage-specific

functions, PU-1 is likely important for both macrophage lineages.

We next used an over-representation approach to identify additional motifs enriched

in the IL-4 induced regions of AAM^{res} or AAM^{mono} (Data S1). Since there were some overlaps

between the motifs discovered by the *de novo* discovery method and over-representation

method, we combined the two sets of motifs for a clustering analysis and merged motifs that

were redundant. Using this approach, we identified macrophage-specific motifs beyond those

from *de novo* motif discovery (Figure 2E). The GATA motifs and basic helix-loop-helix

(bHLH) motifs were specific to IL-4 induced regions of AAM^{res}. TFs with these binding

motifs have been implicated to be important in proliferation of tissue resident macrophages

(19, 20). In contrast, the NFY and STAT motifs were specific to IL-4 induced regions of

AAM^{mono}. These macrophage-specific motifs were only detected in 15-21% of IL-4 induced

regions (236 of 1572 IL-4 induced peaks in AAM^{res} and 317 of 1462 IL-4 induced peaks in

AAM^{mono}), while the PU.1, KLF and AP-1 motifs discovered by the *de novo* method were

present in approximately 75% of IL-4 induced peaks in AAM^{res} and AAM^{mono}. Therefore,

while there are specific TF motifs enriched in IL-4 induced regions of AAM^{res} (GATA and

bHLH) and AAM^mono (NFY and STAT), the majority of IL-4 induced regions are enriched for

a common set of TF motifs.

PU.1 motifs in AAM^res and AAM^mono are associated with macrophage-specific sequence

features.

PU.1 motif was the most frequently found motif from the IL-4 induced regions across

both AAM^res (639 of 1572 IL-4 induced regions) and AAM^mono (552 of 1462 IL-4 induced

regions) (Figure 2C). We focused on these predicted PU.1 binding sites and further

characterized their local sequence features in both types of macrophages. We first quantified

the accuracy of PU.1 motif prediction with actual PU.1 binding, by comparing our predicted

PU.1 motifs from thioglycollate-elicited macrophages with PU.1 ChIP-seq in the same cell

type (5, 21). 78% of the PU.1 motif sites predicted from thioglycollate-induced macrophages

in our study (4,282 of total 5,492 predicted PU.1 motif sites) overlapped with a PU.1 binding

site defined by ChIP-seq (Figure S2B).

We next characterized the PU.1 motifs discovered from the IL-4 induced regions of

AAM^res and AAM^mono. Motif score is commonly used as a proxy of TF-DNA binding affinity.

When compared to PU.1 motifs from AAM^res, PU.1 motifs from AAM^mono had significantly

lower motif scores and also demonstrated greater variability in their values (Figure 3A). To

identify potential co-factors that could bind in collaboration with PU.1 and contribute to

macrophage-specific PU.1 accessibility, we performed motif scanning using sequences from

PU.1 motifs ±25bp flanking sequences. We identified TF motifs that were specific for

AAM^res vs. AAM^mono in these PU.1 regions (Figure 3B, Data S2). These predicted

macrophage-specific co-factors are largely from different families. We observed a greater

diversity in TF families enriched around the IL-4 induced PU.1 motifs of AAM^mono. These

results indicate that PU.1 may function cooperatively with different co-factors to bind

different genomic regions depending on the macrophage lineage.

Since DNA shape has been shown to be a predictor of TF binding pattern (22, 23), we

computationally predicted four DNA shape configurations (minor groove width, propeller

twist, helical twist and roll) at the IL-4 induced PU.1 regions of AAM^{res} and AAM^{mono} (24).

The PU.1 motifs of AAM^{res} have a more conserved DNA configuration, while PU.1 motifs of

AAM^{mono} demonstrated greater variability in DNA configuration. These differences were

most pronounced in the propeller and helical twist configurations of the 8^{th} base pair in the

PU.1 motif (Figure 3C, Data S3). In summary, while the PU.1 motif is the most enriched

motif in the IL-4 induced regions for both AAM^{res} and AAM^{mono}, sequence characteristics in

the PU.1 motifs differ between the two types of peritoneal macrophages, suggesting PU.1

might cooperate with distinct co-regulators in these different lineages.

AAMs from C57BL/6 and BALB/c mice are functionally distinct.

We next characterized the lineage-specific response to IL-4 stimulation on a different genetic

background, by comparing the transcriptional profiles of C57BL/6 and BALB/c AAMs. Most

of the differences in transcriptional profiles were driven by macrophage types (45% of total

variance), although strain differences also contributed to the considerable variation in

transcriptional profiles (18% of total variance) (Figure 4A). Consistent with this finding, most

of the macrophage-specific functions were conserved across mouse strains and not affected

by genetic differences (Figure 4B, left panel). In contrast, functional differences secondary to

genetics were largely specific to the different types of macrophages (Figure 4B, right panel).

We next examined the strain-specific functional differences in AAM^{res} and AAM^{mono},

respectively (Figure 4C, Data S4). BALB/c AAM^{res} expressed lower levels of cell-cycle-

related genes, in line with the previously reported observation that peritoneal AAM^{res} have

lower proliferation capacity during *Litomosoides sigmodontis* infection in BALB/c mice (10).

Furthermore, the expressions of PD-L2 (*Pdcd1lg2*) (Figure 4D) and MHCII molecules

(Figure 4E), which are cellular markers typically used to characterize alternative activation

(12) in AAM^{mono} of C57BL/6 background, were significantly reduced in BALB/c AAM^{mono}.

Notably, while 3 of the 5 MHCII genes (*H2-Aa*, *H2-Ab1*, *H2-Eb1*) had significantly higher

expression in C57BL/6 AAM^{mono}, *H2-Ea-ps* expression was specific to BALB/c AAM^{mono}.

Hence, our studies, together with published findings (10), indicate that macrophages from

BALB/c and C57BL/6 mice are functionally distinct in how they respond to IL-4 activation.

AAMs from C57BL/6 and BALB/c mice respond differently to secondary stimulation by

IL-4.

Primary stimulation of macrophages could lead to the formation of *de novo* enhancer and

transcriptional memory, which consequently hastens the kinetics of activation upon a second

repeated stimulation (4, 13). We hypothesized that the chromatin remodeling that occurred

with IL-4 stimulation in AAMs may translate into enhanced secondary responses with

repeated IL-4 stimulation. To test this hypothesis, we compared the responsiveness of *in vivo*

derived F4/80^{int}CD206$^+$ AAM^{mono} and M^{mono} to a secondary *ex vivo* stimulus of 24 hours

(Figure 5A). The secondary stimulation was performed after the harvested macrophages were

rested overnight in the absence of IL-4. These differences in secondary responses were also

compared between C57BL/6 and BALB/c mice.

AAM^{mono} from C57BL/6 mice were transcriptionally more distinct from M^{mono} after

24 hours of *ex vivo* culture without IL-4 stimulation, when compared to AAM^{mono} from

BALB/c mice (Figure 5B). In general, across both mouse strains, *in vivo* IL-4 stimulation led

to greater transcriptional changes than *ex vivo* IL-4 stimulation (Figure 5B-5C). AAM^{mono}

were also less responsive to the *ex vivo* IL-4 stimulation, as compared to M^{mono}. This

indicated that pre-treatment with *in vivo* IL-4 led to reduced response to a secondary *ex vivo*

IL-4 stimulation and was contrary to what we had hypothesized. However, the degree of

unresponsiveness to the secondary *ex vivo* IL-4 stimulation was more pronounced in

AAM^{mono} from C57BL/6 mice than that in BALB/c AAM^{mono} (Figure 5D).

By taking a supervised approach, we determined that 203 of the 252 genes that were

inducible by *ex vivo* IL-4 in C57BL/6 macrophages would no longer be responsive if the

macrophages were pre-stimulated with *in vivo* IL-4. This was significantly greater (two-sided

Fisher's exact test, p = 1 × 10^{-9}) than that observed for BALB/c macrophages, where 111 of

the 206 genes that were inducible by *ex vivo* IL-4 would not be responsive after pre-treatment

with *in vivo* IL-4 (Figure 5E). Some of the genes that were unresponsive to the secondary IL-

4 stimulation were genes that remained persistently upregulated overnight in the absence of

IL-4 stimulation (92 of the 203 unresponsive genes in C57BL/6 AAM^{mono} and 39 of the 111

unresponsive genes in BALB/c AAM^{mono}). On the other hand, BALB/c macrophages had

significantly higher number of genes that were enhanced in upregulation upon IL-4

restimulation (49 of 206 genes in BALB/c vs. 15 of 252 genes in C57BL/6, two-sided

Fisher's exact test, $p < 2.2 \times 10^{-16}$) (Figure 5E). When we examined the subset of genes that

were only upregulated in BALB/c macrophages with repeated IL-4 stimulation (Figure 5F,

Cluster C6), this included *Arg1* (Figure 5F). In AAM^{mono} of C57BL/6 mice, *Arg1* was

downregulated overnight in the absence of IL-4 and was equally responsive to *ex vivo* IL-4

stimulation in both M^{mono} and AAM^{mono}. In contrast, in macrophages of BALB/c mice, ex

vivo *Arg1* expression was only induced with repeated IL-4 stimulations (Figure 5G). Overall,

these results support the conclusion that macrophages from BALB/c and C57BL/6 mice are

functionally distinct and indicates that this may be particularly important when responding to

a secondary stimulation.

Discussion

248   In this study, we define and characterize IL-4 induced chromatin accessibility with *in*

vivo alternative activation of tissue resident and monocyte-derived peritoneal macrophages.

We also show that *in vivo* alternative activation led to a dampening of macrophage response

to a repeat stimulation. Hence, *in vivo* IL-4 activation did not only lead to remodeling of the

accessible chromatin landscape, but also led to persistent differences in cellular response.

However, this dampening of macrophage secondary response occurs to different degrees in

mice with different genetic backgrounds.

255   We propose that PU.1 is one of the key regulators of IL-4 induced chromatin

accessibility and PU.1 binding can be mediated through DNA shape readout. DNA shape

features, particularly the DNA minor groove width and roll configuration, have been used to

distinguish between functional PU.1 binding sites and randomly occurring PU.1 motif (25).

However, since DNA shape is a consequence of DNA sequence, both modes of DNA

recognition are confounded and difficult to dissociate from one another. For example, it is

unclear if the differences of DNA shape in the PU.1 regions of AAM^{res} and AAM^{mono} are

simply due to differences in co-factor binding, or if the PU.1 protein in these different

lineages of macrophages have different post-translational modifications and recognize

binding sites with different DNA shape. Therefore, it is important for future studies to

identify which amino acid residue(s) in the PU.1 protein could be involved with DNA shape

readout, as mutating these residues could potentially be a strategy to identify PU.1 binding

sites that are solely dependent on DNA shape readout, without being confounded by DNA

sequences (26).

Several studies have recently described the combinatorial effects of cytokines on

macrophage activation, although most of these studies have focused on the opposing effects

of type 1 and type 2 cytokines (16, 17). When repeated stimulation with the same cytokine

was performed and demonstrated "transcriptional memory" (4, 13), these studies were

conducted using short-term stimulation in BMDMs. As such, no comparable studies have so

far demonstrated the dampening of response to repeated cytokine stimulation after *in vivo*

alternative activation. Whereas an enhanced secondary response in macrophage function is

now considered to be a key component of "trained immunity" (27), our results here suggest

that a secondary response may also include a dampening of responses under certain

circumstances that may be dependent on the genetic background of the host. In future work, it

would be important to determine if this dampening of response was specific to *in vivo* IL-4

stimulation, particularly in the setting of a physiological helminth infection model, which

should have continuous production of type 2 cytokines at high abundance and also over a

chronic infection course. Furthermore, while we have only studied this phenomenon of

dampened immune response in AAM^{mono}, it should also be investigated in macrophages of

other tissue origins, as well as other phagocytes, such as neutrophils and dendritic cells.

Finally, given that most human individuals would be exposed to different environmental

stimulations throughout their lives, it would be interesting to determine if such repeated

stimulations would cause dampening in human macrophage response.

Materials and Methods

Experimental methods

Mice Wild type (WT) C57BL/6 mice were purchased from Jackson laboratory and bred onsite for the first set of experiments that compared the effect of IL-4 stimulation on the accessible chromatin profiles from M^{res}, AAM^{res}, M^{mono} and AAM^{mono}. For experiments directly comparing AAMs of C57BL/6 and BALB/c backgrounds, mice of both strains were purchased from Jackson laboratory and immediately used for experiments. Mice were age (7-8 weeks of age) and gender matched. IL-4/anti-IL-4 monoclonal antibody (mAb) complexes (IL-4c) were prepared as described previously (28). To induce AAM^{res}, mice were injected intraperitoneally (i.p.) with IL-4c on days 0 and 2. Mice were also treated with 4% thioglycollate alone (to induce M^{mono}) or in combination with IL-4c to induce AAM^{mono} (12). All animal procedures were approved by the New York University Institutional Animal Care and Use Committee (IACUC) under protocol numbers 131004 and 130504.

Peritoneal cell isolation and cell sorting Peritoneal cells were isolated by washing the peritoneal cavity twice with cold PBS 1x. Peritoneal exudate were then treated with ACK Lysis buffer to lyse red blood cells and washed once with PBS. Cells were then re-suspended to single-cell suspensions for staining with fluorescently conjugated antibodies at 1:100 dilutions, unless otherwise noted. Antibodies were diluted using 2% fetal bovine serum (FBS). Cells were stained with one of either LIVE/DEAD^{TM} Blue (Invitrogen) or LIVE/DEAD^{TM} Near-IR (Invitrogen), blocked with 4μg/ml anti-CD16/32 (2.4G2; Bioxcell) and stained with anti-CD11b Pacific Blue (M1/70; Biolegend), F4/80 PECy7 (BM8; Biolegend), CD206 APC (C068C2; Biolegend), Siglec-F PE (E50-2440; BD Biosciences), CD3 PE (145-2C11; Biolegend), CD19 PE (6D5; Biolegend), CD49b PE (DX5; Biolegend), Ly6G (1A8; Biolegend), PD-L2 (PerCP-Cy55; Miltenyi; diluted at 1:20), MHCII (APC-Cy7; Biolegend). Cells were gated on singlet, live, Dump-negative (CD3−, CD19−, DX5−, Siglec-F−, Ly6G−), CD11b+, then subsequently gated on their M^{res} and AAM^{res} (F4/80hi, CD206−)

or M^{mono} and AAM^{mono} (F4/80int, CD206+) phenotype. Cell surface expression of PD-L2 and

MHCII were acquired for analysis. Cells were sorted using 100μm nozzle into FBS, on either

BD FACSAriaII or SONY HAPS1, depending on instrument availability.

Assay for Transposase-Accessible Chromatin with Sequencing (ATAC-seq) ATAC-seq

was performed as described by Buenrostro et al (18). 50,000 FACS-purified cells per sample

were spun down at 400g for 5 min at 4°C and washed once with 50μl cold PBS. Cells were

lysed with 50μl lysis buffer (10 mM Tris-HCl, pH7.4, 10 mM NaCl, 3 mM MgCl2, 0.1%

IGEPAL CA-630) and immediately spun down at 1500rpm for 10 min at 4°C. The isolated

cell nuclei were then incubated for 30 min at 37°C with 50μl of transposase reaction, which

contained 25μl Tagment DNA buffer (Illumina), 2.5μl Tagment DNA enzyme (Illumina) and

22.5μl nuclease-free water. The transposed DNA was immediately purified using the Qiagen

MinElute PCR Purification Kit (Qiagen) following manufacturer's guide and eluted at 10ul

volume. PCR amplification of the transposed DNA was done using a low-cycle number

protocol and with primers published by Buenrostro et al (18). Each PCR mix contained of

25μl of NEB 2x PCR Mix (New England Biolabs), 2.5μl of 25μM forward primer (Primer

Ad1_noMX), 2.5μl of 25μM reverse barcoded primer, 0.3μl of 100x SYBR Green

(Invitrogen) and 10μl of transposed DNA. PCR was carried out with the cycling protocol:

72°C for 5 min, 98°C for 30s, followed by 5 cycles of 98°C for 10s, 63°C for 30s, 72°C for 1

min. The reaction was held at 4°C after the 5th cycle. A side qPCR was set up using the PCR

product from these 5 cycles of amplification. Each qPCR mix contained 5μl NEB 2x PCR

Mix, 0.25μl 25μM forward primer, 0.25μl 25μM reverse barcoded primer, 0.06μl 100x SYBR

Green, 4.44μl nuclease-free water and 5μl of the PCR-amplified product. qPCR was carried

out using the cycling protocol: 98°C for 30s, followed by 25 cycles of 98°C for 10s, 63°C for

30s, 72°C for 1 min and plate read. The qPCR amplification plot was then used to calculate

the additional number of cycles needed for the PCR to achieve maximum amount of product

without going into saturation. Each sample was amplified for a total of 14-16 cycles. The

amplified libraries were then purified using Qiagen MinElute PCR Purification kit following

manufacturer's guide and eluted at 20μl volume. Libraries were sequenced on the HiSeq 2000

with 2 x 50 cycles and for an average of 50 million paired-end reads per sample. We

performed the IL-4c stimulation experiment twice and generated two independent sets of

libraries to obtain an optimal number of biological replicates for each macrophage population.

The two independent sets of libraries are referred here after as "Run 1" and "Run 2",

respectively. ATAC-seq libraries for C57BL/6 and BALB/c AAMs were generated using the

same protocol and sequenced in a single run.

Transcriptional profiling of BALB/c and C57BL/6 AAMs 100,000 cells were sorted per

sample as described above. FACS-purified cells were spun down and washed once with PBS

before lysis with 350μl of Buffer RLT from the RNeasy Mini Kit (QIAgen). RLT lysates

were homogenized by 1 minute of vortexing and were immediately stored at -80°C until RNA

isolation. RNA was isolated using the RNeasy Mini Kit (QIAgen) based on manufacturer's

protocol, with an additional DNase digestion step using the RNase-free DNase set (QIAgen).

Transcriptional profiling was done using the CEL-seq2 protocol (29) and library preparation

was performed at the NYU School of Medicine Genome Technology Center core facility.

CEL-seq2 libraries were sequenced on the HiSeq 4000 with 2 x 50 cycles. While CEL-seq2

was originally developed as a single-cell assay, we used this protocol in this study as a bulk

transcriptional profiling assay and use the more commonly-used terminology "RNA-seq" to

describe data generated from this assay.

Re-stimulation experiment 300,000 cells (M^{mono} or AAM^{mono} from C57BL/6 and BALB/c

mice, induced using the injection protocols described above) were sorted per sample as

described above. FACS-purified cells were washed twice, then re-suspended to 1ml PBS for

cell counting on the hemocytometer. Cells were spun down again and re-suspended to a

single-cell suspension of 100,000 cells per 200μl, using DMEM with 10% FBS and 1%

Penicillin/Streptomycin. Two aliquots of 100,000 cells were obtained from each sample and

plated separately onto wells of a 48-well tissue-culture plate in 200μl. Cells were incubated

overnight (approximately 18 hours) at 37°C, 5% CO_2. After the overnight incubation, the

initial culture media was discarded and replaced with fresh media. An *ex vivo* dose of IL-4

(20ng/ml) was added for designated wells. Control wells received fresh media only. After 24

hours of stimulation, media was aspirated completely from each well and 350μl of Buffer

RLT was immediately added for cell lysis. The lysed cells of each sample was transferred to a

1.5ml Eppendorf tube, vortexed for 1 minute and immediately stored at -80°C until RNA

isolation. RNA isolation and transcriptional profiling by CEL-seq2 were performed as

described above.

 Bioinformatics and computational methods

ATAC-seq sequence processing Raw ATAC-Seq reads were aligned to the reference mouse

genome mm9 using bowtie2 (v2.2.9) (30), with the parameters --maxin 2000 and --local,

while keeping all other parameters at default settings. To keep only highly unique alignments,

reads with MAPQ score less than 30 were removed. We further removed all duplicate reads,

as well as reads mapping to mitochondrial DNA and chromosome Y. Read filtering steps

were done using the suite of tools from samtools (v1.2 and v1.3.1) (31), ngsutils (v0.5.9) (32)

and picard-tools (http://broadinstitute.github.io/picard/, v1.1.1 and v2.8.2). After all filtering

steps, reads were merged across all replicates from the same macrophage population. This

resulted in a median depth of 15,235,324 reads per macrophage population in Run 1 and

9,865,310 reads per macrophage population in Run 2. For visualization of accessibility reads

on the Integrative Genomics Viewer (IGV), we merged reads from the same macrophage

population across samples from both runs, generated tiled data format (TDF) files using

IGVtools and finally normalized the merged reads to reads per million (RPM) (33).

Identification of accessible chromatin regions We used the merged reads for each

macrophage population to identify accessible chromatin regions, using the PeaKDEck (v1.1)

peak calling algorithm, which measures signal density from randomly sampled bins genome-

wide before generating a data set-specific probability distribution to identify regions with

significant signal enrichment (34). We ran PeaKDEck using sampling bins that consist of a

75bp central bin (-bin) and a 10000bp background bin (-back). Sampling along the genome

was done in steps (-STEP) of 25bp and the background probability distribution was generated

using 100000 randomly selected sites (-npBack). Significance was defined using a p-value of

less than 0.0001 and regions with significant p-values were defined as a "peak" (i.e. an

accessible chromatin region). Peak calling was done independently on libraries generated

from Run 1 and Run 2.

Generation of a union set of accessible chromatin regions We next counted the number of

reads present at each accessible region in order to analyze the ATAC-Seq data using

quantitative approaches downstream. To do this, we first generated a set of consensus peaks

across the data set by taking the union of peaks called from each macrophage population.

Peaks were merged if overlapping by 1bp or more. The number of reads at each peak within

the union peak sets were then counted for each sample. Finally, each peak was re-centered

±100bp on its summit, defined as the position with maximum pile up of reads. Re-centering

on peak summits was performed as this should coincide with the binding event of a

transcription factor within an accessible chromatin region. We implemented the read counting

and peak summit re-centering steps directly using the dba.count function from the

Bioconductor package DiffBind (version 1.14.2) (35). The final count matrix, which

consisted of 61,713 peaks, was used for downstream analyses.

Quantitative ATAC-seq analysis ATAC-seq read counts were transformed using the

regularized logarithmic (rlog) transformation as implemented in the Bioconductor package

DESeq2 (36). To manage batch effect from the two separate libraries, we first modeled the

rlog accessibility read counts to the batch variable using a linear model and subtracted out the

coefficient contributed by the batch variable – this was implemented directly using the

removeBatchEffect function in limma (37). We next chose a set of 30,856 regions with high

variance, using the varFilter function in the genefilter package with default parameters, which

keeps only features with variance inter-quartile range > 0.5 (38). We performed principal

component analysis (PCA) using the batch-subtracted rlog read counts of these regions with

high variance using the prcomp function in R.

430   To identify IL-4 dependent accessible regions, we directly compared the ATAC-seq

profiles of IL-4 stimulated macrophages to their reference non-stimulated macrophages, using

a differential analyses workflow directly implemented through DESeq2. We fit the negative

binomial model in DESeq2 using the raw accessibility reads from all 61,713 regions, with the

model ~ Batch + Population, where Batch is a variable describing if a sample belonged in

Run 1 or Run 2, while Population is a variable describing if the sample is M^{res}, AAM^{res}, M^{mono}

or AAM^{mono}. IL-4 dependency was defined using a significance threshold of False Discovery

Rate (FDR) of 10%. To visualize IL-4 dependent regions, we scaled the batch-subtracted rlog

read counts of these IL-4 dependent regions by z-score transformation and next performed k-

means clustering on these scaled, rlog-transformed reads (K = 4). The clustered matrix was

visualized as a heatmap.

Comparison of sequence properties between constitutively accessible and IL-4 induced

regions *Identification of constitutively accessible and IL-4 induced regions*: To define a set of

constitutively accessible regions, we used only peaks from M^{res} and M^{mono}, respectively, that

were identified in both Run 1 and Run 2. This resulted in 8061 constitutively accessible

regions in M^{res} and 14,045 constitutively accessible regions in M^{mono}. IL-4 induced peaks

were defined using the differential analysis outlined above. All region overlap analyses

throughout this study were performed using the intersect function from the BEDTools suite

(39) and overlaps were defined as any regions overlapping by at least 1bp, unless otherwise

noted.

Genomic elements enrichment analysis: We downloaded genome-wide annotations of five

different genomic elements (promoter, start exon, coding exon, end exon, intron) from the

UCSC Known Gene database for mm9 (40). We defined promoter elements as the 200bp-

region upstream of a transcriptional start site (TSS). We next assigned each of the 61,713

accessible regions in our data set to a unique genomic element label. Where an accessible

region overlapped two different types of genomic elements, we assigned it to the element

with higher number of overlapping base pairs. Finally, any chromatin regions not assigned to

one of these five genomic elements were labeled as intergenic. To determine the enrichment

levels of a particular type of genomic element G within a given set of accessible regions A

(either constitutively accessible or IL-4 induced regions in AAM^{res} or AAM^{mono}), we used the

binomial cumulative probability distribution, b(x; n,p), where x = number of success, n =

number of trials and p = background probability of success. We used the pbinom function in

R. We defined x to be the number of accessible regions in A that were labeled as the genomic

element G that was being tested, n to be the total number of genomic elements G detected in

our combined data set and p to be the proportion of the accessible region A to the total 61,713

accessible regions. This then gave the enrichment levels of G in A, relative to all the

accessible regions identified across the different macrophage populations.

G/C content analysis: To calculate percentage GC, we first used the hgGcPercent function

from   the   kentTools   suite   (v20170111,   UCSC   Genome   Bioinformatics   Group,

https://github.com/ucscGenomeBrowser/kent) to quantitate the number of G or C bases in

each accessible region. This value was next normalized using the length of the accessible

region. CpG island track was downloaded from the UCSC Genome Annotation Database for

mm9 (http://hgdownload.soe.ucsc.edu/goldenPath/mm9/database/). Enrichment levels of CpG

island in a given set of accessible regions A was based on the binomial cumulative probability

as described above, where x = number of accessible regions in A that overlapped a CpG

island, n = number of CpG island in the total data set of 61,713 regions and p = proportion of

A to the total 61,713 regions.

Calculation of distance to IL-4 induced genes: IL-4 induced genes were first identified for

AAM^{res} and AAM^{mono} from the microarrays generated by Gundra and Girgis *et al* (12), using

the linear model and empirical Bayes statistics as implemented in limma, with genes

significantly induced by IL-4 defined using the thresholds FDR 10% and $\log_2$ fold change >

1. The distance between each accessible region and its closest IL-4 induced gene body was

calculated using the closest function in BEDTools.

Transcription factor (TF) motif analysis Sequences of IL-4 induced regions were fetched

using the BEDTools getfasta function for TF motif analysis with the MEME Suite tools (41).

Whole genome fasta file for mm9 was downloaded from the Illumina igenome database

(https://support.illumina.com/sequencing/sequencing_software/igenome.html). Background

file was generated using the function fasta-get-markov in MEME, based on the total 61,713

accessible regions at a Markov model order of 3. TF motif databases (which included mouse

and human TF motifs) were curated as described in (42). We performed *de novo* motif

discovery by running MEME (as part of MEMEChIP, which randomly sampled 600

sequences) with the parameters: -mod zoops -nmotifs 3 -minw 6 -maxw 30. Over-

representation analysis to identify macrophage-specific TF motifs was performed by running

HOMER (43) using sequences from the opposing macrophages as background sequences (i.e.

to identify TF motifs specific to AAM^{res}, sequences of IL-4 induced regions from AAM^{mono}

were used as background sequences) and the parameters: -mask, -size 8,10,12,16, -mset

vertebrates, -nlen 3. For motifs from each macrophage lineage, we combined all three *de novo*

motifs discovered by MEME and motifs with enrichment $\log_2$ p-values < -15 by HOMER for

clustering analysis (we used the known motifs output from HOMER). This resulted in 14

motifs for AAM^{res} and AAM^{mono}, respectively. Clustering was done using STAMP (44, 45),

with the frequency matrices of motifs and the default parameters of: column comparison

metric – Pearson correlation coefficient, alignment method – ungapped Smith-Waterman,

tree-building algorithm – UPGMA, multiple alignment strategy – iterative refinement.

Comparisons between predicted PU.1 motif and ChIP-seq defined PU.1 binding sites

PU.1 ChIP-seq regions identified in thioglycollate-induced macrophages, generated from two

different experiments, were downloaded as BED files that had been directly deposited on

Gene Expression Omnibus (GEO) (GSM1131238 and GSM1183968) (5, 21). A set of 55,386

reproducible PU.1 binding sites were defined by intersecting these two sets of PU.1 ChIP-seq

regions. We used the PU.1 motif discovered *de novo* from all the constitutively accessible

regions in M^{mono} and ran FIMO to identify all PU.1 motif sites from M^{mono}, using a p-value

threshold of 0.0001 and the background file generated as described above. Since the

published PU.1 ChIP-seq regions were of 200bp length, we extended the predicted PU.1

motifs from M^{mono} by ±100bp to match the comparison. Overlapping rate was calculated as

total number of predicted PU.1 motifs from M^{mono} overlapping a reproducible PU.1 ChIP-seq

region / total number of predicted PU.1 motifs from M^{mono} × 100%.

PU.1 motif analysis We identified IL-4 induced PU.1 motif sites by performing motif

scanning with FIMO (46), using the PU.1 motif discovered *de novo* from the IL-4 induced

peaks of AAM^{res} and AAM^{mono}, respectively. FIMO was run with a p-value threshold of

0.0001 (as part of MEMEChIP). Motif scores were also calculated as part of FIMO.

The detected IL-4 induced PU.1 motif sites were then extended ±25bp using the

BEDTools slop function. These PU.1 motif sites ± 25bp are hereafter referred to as "PU.1

regions". To identify potential co-factors for PU.1, these PU.1 regions were specifically

subjected to motif scanning by FIMO, with a background model that was based on the 55,386

PU.1 ChIP-seq peaks described above and at a Markov model order of 3. To determine which

of these detected motifs were macrophage-specific, the odds ratio of a motif being detected in

the PU.1 regions of AAM^{res} vs. AAM^{mono} were calculated. P-value was determined using two-

sided Fisher's test, with the null hypothesis of a motif being equally likely to be detected in

the IL-4 induced PU.1 regions of AAM^{res} and AAM^{mono} (i.e. $\log_2$ odds ratio of zero). When

multiple motifs of the same TF were present in the database, we used the motif that was most

frequently detected in PU.1 regions for odds ratio calculations. TFs from mouse and human

were kept as separate analyses, but included in the same visualization. To account for the

multiple hypotheses testing performed over 633 mouse TFs and 835 human TFs, the

Benjamini-Hochberg procedure was used to perform p-value adjustment by calculating the

FDR and significance threshold was set at FDR 10%. Hence, statistically significant TFs were

macrophage-specific, with $\log_2$ odds ratio > 0 indicating AAM^{res}-specificity and $\log_2$ odds

ratio < 0 indicating AAM^{mono}-specificity. For visualization of significant results, TFs were

summarized at the family level as defined in (47) (Figure 3B). The maximum absolute $\log_2$

odds ratio of the family was visualized and the specific TF with the maximum absolute $\log_2$

odds ratio value was stated in parenthesis. Where TF family annotation was not available, the

$\log_2$ odds ratio of the specific TF itself was used.

DNA shape features of PU.1 motifs were analyzed using the DNAshape algorithm

(24, 48) for 4 different DNA shape configurations at single nucleotide resolution. Sequences

on the anti-sense strand were reverse complemented prior to DNA shape prediction.

Processing of CEL-seq reads CEL-seq reads were first demultiplexed using the

bc_demultiplex script from https://github.com/yanailab/CEL-Seq-pipeline (29).

Demultiplexed reads were aligned to the mm10 mouse reference genome using bowtie2

(version 2.2.9). Aligned reads were counted for each gene using a modified htseq-count script

(from https://github.com/yanailab/CEL-Seq-pipeline) adapted for CEL-seq reads with unique

molecular identified (UMI). We included only reads with MAPQ score > 30 and removed

singleton genes. This resulted in a final median read depth of 737,848 reads per sample,

covering a median of 11,096 genes per sample. PCA was performed using 7431 genes with

high variance, defined using the varFilter function in the genefilter package with default

parameters, which keeps only features with variance inter-quartile range > 0.5. Differential

analysis was done using DESeq2 by fitting the negative binomial model using ~ Strain +

CellType + Strain:CellType. Significantly differential genes were extracted using a threshold

of FDR 10% for the four different comparisons of: (1) AAM^{res} vs. AAM^{mono} in C57BL/6, (2)

AAM^{res} vs. AAM^{mono} in BALB/c, (3) BALB/c vs. C57BL/6 in AAM^{res} and (4) BALB/c vs.

C57BL/6 in AAM^{mono}. Overlapping genes were defined as genes identified as differential in

two different comparisons. Pathway enrichment analysis was done using strain-specific genes

in AAM^{res} and AAM^{mono}, respectively, through Ingenuity Pathway Analysis (IPA) with the

parameter Organism = Mouse and keeping all other parameters at default settings.

Defining gene classes from re-stimulation experiment CEL-seq reads were processed as

described above. This resulted in a final median read depth of 979,547 reads per sample,

covering a median of 10,820 genes per sample. PCA was separately performed on the rlog

read counts from 6528 high variance genes from C57BL/6 and BALB/c, respectively. The

differences in transcriptional profiles induced by *ex vivo* IL-4 stimulation were represented as

Euclidean distances, which were calculated between transcriptional profiles (rlog read counts

of all non-singleton genes) of cell aliquots from the same cell type that either received a

secondary *ex vivo* IL-4 stimulation or cultured in control media. Differential analysis was

done using DESeq2 by fitting the negative binomial model using ~ Group, where Group is a

factor consisting of 8 different levels (M^{mono} from C57BL/6, AAM^{mono} from C57BL/6, M^{mono}

with *ex vivo* IL-4 stimulation from C57BL/6, AAM^{mono} with *ex vivo* IL-4 stimulation from

C57BL/6, M^{mono} from BALB/c, AAM^{mono} from BALB/c, M^{mono} with *ex vivo* IL-4 stimulation

from BALB/c, AAM^{mono} with *ex vivo* IL-4 stimulation from BALB/c). Significantly

differential genes were extracted using a threshold of FDR 10% and the comparisons are

described on Figure 7B.

To define "Unresponsive", "Enhanced" and "Equal" genes, we identified genes that

were either uniquely or commonly differential in the comparisons "AAM^{mono} with *ex vivo* IL-

4 vs. AAM^{mono}" and "M^{mono} with *ex vivo* IL-4 vs. M^{mono}" (49). "Unresponsive" genes were

defined as genes that were uniquely differential in "M^{mono} with *ex vivo* IL-4 vs. M^{mono}", while

"Enhanced" genes were uniquely differential in "AAM^{mono} with *ex vivo* IL-4 vs. AAM^{mono}".

For genes that were differential in both comparisons, we further examined the magnitude of

fold change – if the fold change (defined as a differences in $\log_2$ fold change of >1) is greater

in the "M^{mono} with *ex vivo* IL-4 vs. M^{mono}" comparison, it is defined as an "Unresponsive"

genes and if the fold change is greater in the "AAM^{mono} with *ex vivo* IL-4 vs. AAM^{mono}"

comparison, it is defined as an "Enhanced" gene. Finally, if the magnitude of fold change is

similar in both comparisons (defined as a differences in $\log_2$ fold change of <1), the gene is

defined as "Equal". This categorization process of different gene classes was performed for

C57BL/6 and BALB/c macrophages, respectively. For visualization of these different gene

groups, the rlog read counts from all of these genes were extracted, scaled to a mean of 0 and

standard deviation of 1 across each sample, and subjected to k-means clustering, with K=6.

Heatmap was visualized using the R package pheatmap.

Acknowledgements

We thank the NYU School of Medicine Genome Technology Center and Cytometry and Cell Sorting Laboratory core facilities. These shared resources are partially supported by the Laura and Isaac Perlmutter Cancer Center support grant P30CA016087. We also thank the NYU IT High Performance Computing resources, services, and staff expertise. This work was supported through the NIH, NIAID grants AI093811 and AI094166 (P.L.) and NIDDK grant DK103788 (P.L.), American Association of Immunologists (M.S.T.), Vilcek Foundation (M.S.T.).

Author contributions

Conceptualization, M.S.T. and P.L.; Methodology, M.S.T, E.R.M., N.M.G., R.A.B. and P.L.; Formal Analysis, M.S.T.; Investigation, M.S.T. and N.M.G.; Writing – Original Draft, M.S.T. and P.L.; Writing – Review and Editing, E.R.M. and P.L.; Visualization, M.S.T.; Data Curation, M.S.T.; Supervision, R.A.B. and P.L.; Funding acquisition, P.L. All authors read and approved of the final draft.

Data availability

ATAC-Seq and RNA-Seq data have been deposited on NCBI database Gene Expression Omnibus (GEO) under the SuperSeries GSE116108 (ATAC-Seq subseries GSE116107 and RNA-Seq subseries GSE116105).

References

623
624

1.   Glass, C. K., and G. Natoli. 2016. Molecular control of activation and priming in
macrophages. *Nature immunology* 17: 26.
2.   Ghisletti, S., I. Barozzi, F. Mietton, S. Polletti, F. De Santa, E. Venturini, L. Gregory,
L. Lonie, A. Chew, C.-L. Wei, J. Ragoussis, and G. Natoli. 2010. Identification and
Characterization of Enhancers Controlling the Inflammatory Gene Expression
Program in Macrophages. *Immunity* 32: 317-328.
3.   Natoli, G., S. Ghisletti, and I. Barozzi. 2011. The genomic landscapes of
inflammation. *Genes & development* 25: 101-106.
4.   Ostuni, R., V. Piccolo, I. Barozzi, S. Polletti, A. Termanini, S. Bonifacio, A. Curina,
E. Prosperini, S. Ghisletti, and G. Natoli. 2013. Latent enhancers activated by
stimulation in differentiated cells. *Cell* 152: 157-171.
5.   Kaikkonen, M. U., N. J. Spann, S. Heinz, C. E. Romanoski, K. A. Allison, J. D.
Stender, H. B. Chun, D. F. Tough, R. K. Prinjha, and C. Benner. 2013. Remodeling
of the enhancer landscape during macrophage activation is coupled to enhancer
transcription. *Molecular cell* 51: 310-325.
6.   Comoglio, F., M. Simonatto, S. Polletti, X. Liu, S. T. Smale, I. Barozzi, and G.
Natoli. 2019. Dissection of acute stimulus-inducible nucleosome remodeling in
mammalian cells. *Genes & development* 33: 1159-1174.
7.   Herbert, D. R., C. Holscher, M. Mohrs, B. Arendse, A. Schwegmann, M. Radwanska,
M. Leeto, R. Kirsch, P. Hall, H. Mossmann, B. Claussen, I. Forster, and F.
Brombacher. 2004. Alternative macrophage activation is essential for survival during
schistosomiasis and downmodulates T helper 1 responses and immunopathology.
Immunity 20: 623-635.
8.   Chen, F., W. Wu, A. Millman, J. F. Craft, E. Chen, N. Patel, J. L. Boucher, J. F.
Urban, Jr., C. C. Kim, and W. C. Gause. 2014. Neutrophils prime a long-lived
effector macrophage phenotype that mediates accelerated helminth expulsion. *Nature
immunology* 15: 938-946.
9.   Chen, F., Z. Liu, W. Wu, C. Rozo, S. Bowdridge, A. Millman, N. Van Rooijen, J. F.
Urban, Jr., T. A. Wynn, and W. C. Gause. 2012. An essential role for TH2-type
responses in limiting acute tissue damage during experimental helminth infection.
Nat Med 18: 260-266.
10.   Campbell, S. M., J. A. Knipper, D. Ruckerl, C. M. Finlay, N. Logan, C. M. Minutti,
M. Mack, S. J. Jenkins, M. D. Taylor, and J. E. Allen. 2018. Myeloid cell recruitment
versus local proliferation differentiates susceptibility from resistance to filarial
infection. *eLife* 7.
11.   Gundra, U. M., N. M. Girgis, M. A. Gonzalez, M. San Tang, H. J. P. Van Der Zande,
J.-D. Lin, M. Ouimet, L. J. Ma, J. Poles, N. Vozhilla, E. A. Fisher, K. J. Moore, and
P. n. Loke. 2017. Vitamin A mediates conversion of monocyte-derived macrophages
into tissue-resident macrophages during alternative activation. *Nature immunology*
advance online publication.
12.   Gundra, U. M., N. M. Girgis, D. Ruckerl, S. Jenkins, L. N. Ward, Z. D. Kurtz, K. E.
Wiens, M. S. Tang, U. Basu-Roy, A. Mansukhani, J. E. Allen, and P. n. Loke. 2014.
Alternatively activated macrophages derived from monocytes and tissue macrophages
are phenotypically and functionally distinct. *Blood* 123: 22.
13.   Daniel, B., G. Nagy, A. Horvath, Z. Czimmerer, I. Cuaranta-Monroy, S. Poliska, T.
T. Hays, S. Sauer, J. Francois-Deleuze, and L. Nagy. 2018. The IL-4/STAT6/PPAR
γ signaling axis is driving the expansion of the RXR heterodimer cistrome,
providing complex ligand responsiveness in macrophages. *Nucleic Acids Research*:
gky157-gky157.
14.   Daniel, B., G. Nagy, Z. Czimmerer, A. Horvath, D. W. Hammers, I. Cuaranta-
Monroy, S. Poliska, P. Tzerpos, Z. Kolostyak, T. T. Hays, A. Patsalos, R. Houtman,
S. Sauer, J. Francois-Deleuze, F. Rastinejad, B. L. Balint, H. L. Sweeney, and L.
Nagy. 2018. The Nuclear Receptor PPARgamma Controls Progressive Macrophage

Polarization as a Ligand-Insensitive Epigenomic Ratchet of Transcriptional Memory. *Immunity* 49: 615-626.e616.

15. Czimmerer, Z., B. Daniel, A. Horvath, D. Rückerl, G. Nagy, M. Kiss, M. Peloquin, M. M. Budai, I. Cuaranta-Monroy, Z. Simandi, L. Steiner, B. Nagy, Jr., S. Poliska, C. Banko, Z. Bacso, I. G. Schulman, S. Sauer, J.-F. Deleuze, J. E. Allen, S. Benko, and L. Nagy. 2018. The Transcription Factor STAT6 Mediates Direct Repression of Inflammatory Enhancers and Limits Activation of Alternatively Polarized Macrophages. *Immunity* 48: 75-90.e76.

16. Piccolo, V., A. Curina, M. Genua, S. Ghisletti, M. Simonatto, A. Sabo, B. Amati, R. Ostuni, and G. Natoli. 2017. Opposing macrophage polarization programs show extensive epigenomic and transcriptional cross-talk. *Nature immunology* 18: 530-540.

17. Kang, K., S. H. Park, J. Chen, Y. Qiao, E. Giannopoulou, K. Berg, A. Hanidu, J. Li, G. Nabozny, K. Kang, K. H. Park-Min, and L. B. Ivashkiv. 2017. Interferon-gamma Represses M2 Gene Expression in Human Macrophages by Disassembling Enhancers Bound by the Transcription Factor MAF. *Immunity* 47: 235-250.e234.

18. Buenrostro, J. D., P. G. Giresi, L. C. Zaba, H. Y. Chang, and W. J. Greenleaf. 2013. Transposition of native chromatin for fast and sensitive epigenomic profiling of open chromatin, DNA-binding proteins and nucleosome position. *Nature methods* 10: 1213-1218.

19. Rosas, M., L. C. Davies, P. J. Giles, C. T. Liao, B. Kharfan, T. C. Stone, V. B. O'Donnell, D. J. Fraser, S. A. Jones, and P. R. Taylor. 2014. The transcription factor Gata6 links tissue macrophage phenotype and proliferative renewal. *Science* 344: 645-648.

20. Jarjour, N. N., E. A. Schwarzkopf, T. R. Bradstreet, I. Shchukina, C. C. Lin, S. C. Huang, C. W. Lai, M. E. Cook, R. Taneja, T. S. Stappenbeck, G. J. Randolph, M. N. Artyomov, J. F. Urban, Jr., and B. T. Edelson. 2019. Bhlhe40 mediates tissue-specific control of macrophage proliferation in homeostasis and type 2 immunity. *Nature immunology* 20: 687-700.

21. Heinz, S., C. E. Romanoski, C. Benner, K. A. Allison, M. U. Kaikkonen, L. D. Orozco, and C. K. Glass. 2013. Effect of natural genetic variation on enhancer selection and function. *Nature* 503: 487-492.

22. Dror, I., T. Golan, C. Levy, R. Rohs, and Y. Mandel-Gutfreund. 2015. A widespread role of the motif environment in transcription factor binding across diverse protein families. *Genome Res* 25: 1268-1280.

23. Gordân, R., N. Shen, I. Dror, T. Zhou, J. Horton, R. Rohs, and M. L. Bulyk. 2013. Genomic regions flanking E-box binding sites influence DNA binding specificity of bHLH transcription factors through DNA shape. *Cell reports* 3: 1093-1104.

24. Zhou, T., L. Yang, Y. Lu, I. Dror, A. C. Dantas Machado, T. Ghane, R. Di Felice, and R. Rohs. 2013. DNAshape: a method for the high-throughput prediction of DNA structural features on a genomic scale. *Nucleic acids research* 41: W56-W62.

25. Barozzi, I., M. Simonatto, S. Bonifacio, L. Yang, R. Rohs, S. Ghisletti, and G. Natoli. 2014. Coregulation of transcription factor binding and nucleosome occupancy through DNA features of mammalian enhancers. *Molecular cell* 54: 844-857.

26. Abe, N., I. Dror, L. Yang, M. Slattery, T. Zhou, H. J. Bussemaker, R. Rohs, and R. S. Mann. 2015. Deconvolving the recognition of DNA shape from sequence. *Cell* 161: 307-318.

27. Netea, M. G., L. A. B. Joosten, E. Latz, K. H. G. Mills, G. Natoli, H. G. Stunnenberg, L. A. J. O'Neill, and R. J. Xavier. 2016. Trained immunity: A program of innate immune memory in health and disease. *Science (New York, N.Y.)* 352: aaf1098-aaf1098.

28. Jenkins, S. J., D. Ruckerl, P. C. Cook, L. H. Jones, F. D. Finkelman, N. van Rooijen, A. S. MacDonald, and J. E. Allen. 2011. Local macrophage proliferation, rather than recruitment from the blood, is a signature of TH2 inflammation. *Science* 332: 1284-1288.

29. Hashimshony, T., N. Senderovich, G. Avital, A. Klochendler, Y. de Leeuw, L.
Anavy, D. Gennert, S. Li, K. J. Livak, O. Rozenblatt-Rosen, Y. Dor, A. Regev, and I.
Yanai. 2016. CEL-Seq2: sensitive highly-multiplexed single-cell RNA-Seq. *Genome*
biology 17: 77.
30. Langmead, B., and S. L. Salzberg. 2012. Fast gapped-read alignment with Bowtie 2.
Nat Meth 9: 357-359.
31. Li, H., B. Handsaker, A. Wysoker, T. Fennell, J. Ruan, N. Homer, G. Marth, G.
Abecasis, and R. Durbin. 2009. The Sequence Alignment/Map format and SAMtools.
Bioinformatics (Oxford, England) 25: 2078-2079.
32. Breese, M. R., and Y. Liu. 2013. NGSUtils: a software suite for analyzing and
manipulating next-generation sequencing datasets. *Bioinformatics (Oxford, England)*
29: 494-496.
33. Thorvaldsdóttir, H., J. T. Robinson, and J. P. Mesirov. 2013. Integrative Genomics
Viewer (IGV): high-performance genomics data visualization and exploration.
Briefings in Bioinformatics 14: 178-192.
34. McCarthy, M. T., and C. A. O'Callaghan. 2014. PeaKDEck: a kernel density
estimator-based peak calling program for DNaseI-seq data. *Bioinformatics (Oxford,*
England) 30: 1302-1304.
35. Stark, R., and G. Brown. 2011. DiffBind: differential binding analysis of ChIP-Seq
peak data.
36. Love, M. I., W. Huber, and S. Anders. 2014. Moderated estimation of fold change
and dispersion for RNA-seq data with DESeq2. *Genome biology* 15: 550.
37. Ritchie, M. E., B. Phipson, D. Wu, Y. Hu, C. W. Law, W. Shi, and G. K. Smyth.
2015. limma powers differential expression analyses for RNA-sequencing and
microarray studies. *Nucleic Acids Res* 43: e47.
38. Gentleman, R., V. Carey, W. Huber, and F. Hahne. 2016. genefilter: methods for
filtering genes from high-throughput experiments. R package version 1.56.0 ed.
39. Quinlan, A. R., and I. M. Hall. 2010. BEDTools: a flexible suite of utilities for
comparing genomic features. *Bioinformatics (Oxford, England)* 26: 841-842.
40. Hsu, F., W. J. Kent, H. Clawson, R. M. Kuhn, M. Diekhans, and D. Haussler. 2006.
The UCSC Known Genes. *Bioinformatics (Oxford, England)* 22: 1036-1046.
41. Bailey, T. L., M. Boden, F. A. Buske, M. Frith, C. E. Grant, L. Clementi, J. Ren, W.
W. Li, and W. S. Noble. 2009. MEME SUITE: tools for motif discovery and
searching. *Nucleic Acids Res* 37: W202-208.
42. Karwacz, K., E. R. Miraldi, M. Pokrovskii, A. Madi, N. Yosef, I. Wortman, X. Chen,
A. Watters, N. Carriero, A. Awasthi, A. Regev, R. Bonneau, D. Littman, and V. K.
Kuchroo. 2017. Critical role of IRF1 and BATF in forming chromatin landscape
during type 1 regulatory cell differentiation. *Nature immunology* 18: 412.
43. Heinz, S., C. Benner, N. Spann, E. Bertolino, Y. C. Lin, P. Laslo, J. X. Cheng, C.
Murre, H. Singh, and C. K. Glass. 2010. Simple combinations of lineage-determining
transcription factors prime cis-regulatory elements required for macrophage and B
cell identities. *Molecular cell* 38: 576-589.
44. Mahony, S., P. E. Auron, and P. V. Benos. 2007. DNA Familial Binding Profiles
Made Easy: Comparison of Various Motif Alignment and Clustering Strategies. *PLoS*
computational biology 3: e61.
45. Benos, P. V., and S. Mahony. 2007. STAMP: a web tool for exploring DNA-binding
motif similarities. *Nucleic Acids Research* 35: W253-W258.
46. Grant, C. E., T. L. Bailey, and W. S. Noble. 2011. FIMO: scanning for occurrences of
a given motif. *Bioinformatics (Oxford, England)* 27: 1017-1018.
47. Wingender, E., T. Schoeps, M. Haubrock, M. Krull, and J. Dönitz. 2018. TFClass:
expanding the classification of human transcription factors to their mammalian
orthologs. *Nucleic Acids Research* 46: D343-D347.
48. Chiu, T.-P., F. Comoglio, T. Zhou, L. Yang, R. Paro, and R. Rohs. 2016.
DNAshapeR: an R/Bioconductor package for DNA shape prediction and feature
encoding. *Bioinformatics (Oxford, England)* 32: 1211-1213.

49.   Kaufmann, E., J. Sanz, J. L. Dunn, N. Khan, L. E. Mendonca, A. Pacis, F. Tzelepis,
788   E. Pernet, A. Dumaine, J. C. Grenier, F. Mailhot-Leonard, E. Ahmed, J. Belle, R.
789   Besla, B. Mazer, I. L. King, A. Nijnik, C. S. Robbins, L. B. Barreiro, and M.
790   Divangahi. 2018. BCG Educates Hematopoietic Stem Cells to Generate Protective
791   Innate Immunity against Tuberculosis. *Cell* 172: 176-190.e119.
792
793

Figures legends

Figure 1: IL-4 stimulation leads to remodeling of open chromatin landscape in

peritoneal macrophages.

(A) PCA scores of individual ATAC-seq samples. PCA was performed using rlog-

transformed ATAC-seq read counts of 30,856 regions with high variance (only regions with

variance inter-quartile range > 0.5 were retained). N = 4-6 mice per macrophage population.

Data points represent independent biological replicates. (B) Genome browser views of

representative (boxed) constitutively accessible and IL-4 induced regions. (C) The

contributions of individual accessible regions to PCs 1 and 2 are represented in the PCA

loadings plot. Each data point is color-coded based on the direction of its IL-4 dependency.

Hence, IL-4 induced regions (red) are highly associated with IL-4 stimulated macrophages,

while IL-4 repressed regions (yellow) are highly associated with non-stimulated

macrophages. We compared (D) enrichment levels for different types of genomic elements,

(E) distance from a closest IL-4-induced gene and (F) G/C content between constitutively

accessible and IL-4-induced regions in AAM^{mono} and AAM^{res}, respectively. G/C content

information is represented in two different ways – percentage of G/C bases in an accessible

region (F, left panel) and CpG island enrichment for a given group of accessible regions (F,

right panel). Number of IL-4 induced regions = 1572 in AAM^{res} and 1462 in AAM^{mono};

number of constitutively accessible regions = 8061 in AAM^{res} and 14045 in AAM^{mono}.

Enrichment p-values are from binomial test while two-class comparison p-values are from

two-sided Mann-Whitney test.

Figure 2: IL-4 induced regions are associated with PU.1, KLF and AP-1 motifs.

(A) Heatmap visualizing the macrophage-specific IL-4 dependent regions. Each row

represents one of the 2855 IL-4 dependent regions and each column a unique sample. Values

are rlog-transformed, batch-subtracted read counts, scaled using a z-score transformation for

each region. (B) Motifs discovered *de novo* from IL-4 induced regions in AAM^{res} and

AAM^{mono}. (C) Frequency of IL-4 induced peaks delineated by the presence of *de novo* PU.1,

KLF and AP-1 motifs. (D) 20 highly expressed TF genes related to the *de novo* motifs

discovered from the IL-4 induced regions. Values are log$_2$ intensity values of microarrays

(12). (E) Clustering analysis of *de novo* motifs and macrophage-specific motifs identified

using an over-representation approach, from the IL-4 induced regions of AAM^{res} (left) and

AAM^{mono} (right). Asterisks indicate macrophage-specific motifs uniquely identified via the

over-representation approach. Only macrophage-specific motifs with log$_2$ p-value < -15 are

included in this visualization and the complete list of motifs identified by over-representation

are included as Data S1.

Figure 3: PU.1 motifs in AAM^{res} and AAM^{mono} are associated with macrophage-specific

sequence features.

(A) Comparison of PU.1 motif scores derived by FIMO in IL-4 induced regions of AAM^{res}

vs. AAM^{mono}, with horizontal lines in the violin plots representing values at 25th, 50th and 75th

percentiles. P-value is from a two-sided Mann-Whitney test. Number of IL-4 induced regions

= 1572 in AAM^{res} and 1462 in AAM^{mono}. (B) Macrophage-specific TF motifs found within

IL-4 induced PU.1 motifs ± 25bp regions, represented using log2 odds ratio values (two-sided

Fisher's test, adjusted p-values < 0.1). Motifs are summarized as TF families and the specific

TF with the maximum absolute log$_2$ odds ratio value is stated in parenthesis. Where TF

family annotation was not available, the log$_2$ odds ratio of the specific TF itself is used. (C)

Predicted DNA shape at the 8th base pair of PU.1 motif of AAM^{res} and AAM^{mono}. Frequency

distributions are represented by smoothed kernel density estimates. Complete comparisons at

all nucleotides are presented in Data S3. ProT = propeller twist, HelT = helical twist, MGW =

minor groove width.

Figure 4: AAMs from C57BL/6 and BALB/c are functionally distinct.

(A) PCA of 7431 genes with high variance (only genes with variance inter-quartile range >

0.5 were retained). Data points represent independent biological replicates. (B) Venn

diagrams indicating the number of genes that were commonly and uniquely identified as

significantly differential (FDR < 0.1) in different comparisons – (left) macrophage-specific

genes in C57BL/6 and BALB/c AAMs; (right) strain-specific genes in AAM^{res} and AAM^{mono}.

(C) Enrichment values from Ingenuity Pathway Analysis visualized as $-\log_{10}$ P-value for the

four different groups of genes – (1) BALB/c specific in AAM^{res}, (2) C57BL/6 specific in

AAM^{res}, (3) BALB/c specific in AAM^{mono} and (4) C57BL/6 in AAM^{mono}. Only the top 10

pathways (as defined by enrichment p-values) are included in this visualization. Specific

pathways are highlighted for clarity. The complete lists of enriched pathways are provided as

Supplemental Materials Data S2. (D) Representative flow cytometric analysis of F4/80 and

PD-L2 surface expressions in AAM^{mono} of C57BL/6 and BALB/c mice. Boxplots show

frequency of CD11b+ F4/80+ PD-L2+ singlet, live cells. P-value is based on a two-sided

unpaired t-test. (E) Expression of the *Pdcd1lg2* gene in AAM^{mono} of C57BL/6 vs. BALB/c

mice, represented by size-factor normalized read counts. P-value is from DESeq2 and

adjusted by Benjamini-Hochberg procedure. (F) Representative flow cytometric analysis of

F4/80 and MHCII surface expressions in AAM^{mono} of C57BL/6 and BALB/c mice. Boxplots

show frequency of CD11b+ F4/80+ MHCII+ singlet, live cells. P-value is based on a two-

sided unpaired t-test. (G) Expression values of all MHCII genes in AAM^{mono} of C57BL/6 vs.

BALB/c mice, represented by size-factor normalized read counts. P-values are from DESeq2

and adjusted by Benjamini-Hochberg procedure. Hinges of all boxplots correspond to values

of the 25th, 50th and 75th percentiles, while boxplot whiskers extend to no more than 1.5 ×

inter-quartile range, beyond which the outlier data points will be plotted individually.

Transcriptional profiling analysis: N = 8 AAM^{res} (C57BL/6), 4 AAM^{res} (BALB/c), 7

AAM^{mono} (C57BL/6), 6 AAM^{mono} (BALB/c). Flow cytometric analysis: N = 6 AAM^{mono}

(C57BL/6), 6 AAM^{mono} (BALB/c).

Figure 5: AAMs from C57BL/6 and BALB/c mice respond differently to secondary stimulation by IL-4.

(A) Schematic illustrating the study designed to assess differences in response to a secondary IL-4 stimulation between AAMmono and Mmono of C57BL/6 and BALB/c backgrounds. (B) Number of significantly differential genes (FDR < 10%) identified for each comparison. (C) PCA performed separately on 6528 genes with high variance (only genes with variance inter-quartile range > 0.5 were retained) in C57BL/6 and BALB/c macrophages. Data points represent independent biological replicates. (D) Differences in global transcriptional profiles induced by *ex vivo* IL-4, quantitated using Euclidean distance that is calculated between transcriptional profiles (rlog read counts of all non-singleton genes) of cell aliquots from the same cell type that either received a secondary *ex vivo* IL-4 stimulation or cultured in control media. (E) Frequency of genes with enhanced, equal or dampened upregulation after pre-treatment with in vivo IL-4. (F) A union set of 334 genes upregulated by *ex vivo* IL-4 in C57BL/6 and BALB/c macrophages, separated into 6 different clusters by k-means clustering. (G) Expression values of *Arg1* across the different conditions, represented by size-factor normalized read counts. N = 3 mice for each group.

Supplemental Figure 1: IL-4 stimulation leads to remodeling of open chromatin

landscape in peritoneal macrophages.

(A) Comparisons between the accessible chromatin profiles of IL-4 stimulated macrophages

and non-stimulated macrophages, presented as MA plots (left – AAM^{res}; right – AAM^{mono}).

Differential chromatin regions (FDR 10%, |LFC| > 0) are highlighted in red. (B) Venn

diagram illustrating the minimal overlap between IL-4 induced regions in AAM^{res} and

AAM^{mono}. Values in Venn diagrams represent the number of unique accessible regions in

each corresponding set. (C) Schematic outlining the workflow to identify IL-4 induced

regions and constitutively regions for comparisons of the sequence properties associated with

these two classes of genomic elements.

Supplemental Figure 2: Motif analyses of IL-4 induced regions in AAM^{res} and AAM^{mono}.

(A) Expression values of all TF genes related to the *de novo* motifs discovered from the IL-4

induced regions. Values are log$_2$ intensity values of microarrays (12) and were clustered by

hierarchical clustering. TF genes falling into the 2nd cluster were excluded from the analysis

presented in Figure 2D due to their low expression. (B) Schematic outlining the workflow

used to verify the accuracy of predicted PU.1 motif sites by comparing to published ChIP-Seq

data sets. (C) Schematic outlining the workflow used to analyze features of IL-4 induced

PU.1 motif sites.

911
912

Data S1. Complete lists of TF motifs and associated statistics identified by the over-

representation approach. Related to Figure 2E.

Data S2. Complete list of macrophage-specific motifs detected within ±25bp of PU.1 motifs

in AAM^{res} and AAM^{mono}. Related to Figure 3B.

Data S3. Composite plots of predicted DNA shape values at single nucleotide resolution,

between base pair 3-13 of the PU.1 motifs in AAM^{res} and AAM^{mono}. Related to Figure 3D.

Data S4. List of pathways enriched among strain-specific genes from AAM^{res} and AAM^{mono}.

Related to Figure 4C.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

A

B

C

IL-4 induced (AAM^res)   IL-4 induced (AAM^mono)

1393   **179**   **1283**

Supplemental Figure 1

A

Klf2
Klf4
Klf10
Klf3
Batf3
Jund
Jun
Klf6
Fosb
Fosl1
Klf5
Klf14
Klf15
Klf1
Klf11
Klf12
Klf9
Fos
Klf7
Batf
Jdp2
Batf2
Klf13
Fosl2
Junb
Klf17
Klf16
Klf8

AAM^res   M^res   AAM^mono   M^mono

log$_2$ expression

8   10   12   14   16

B

M^mono constitutively accessible regions

↓

De novo motif discovery

↓

Identify sites of *de novo* PU.1 motifs

↓

Extend +/- 100bp

5,492 regions in M^mono

Published PU.1 binding regions defined by ChIP-Seq BED files from Heinz et al (Nature 2013), Kaikkonen et al (Mol Cell 2013)

↓

55,386 reproducible PU.1 binding regions

Overlap → 4,282 predicted PU.1 motifs from M^mono overlap a PU.1 ChIP-Seq region

C

IL-4 induced regions

n = 1572 AAM^res
n = 1462 AAM^mono

→ *De novo* motif discovery

→ Identify sites of *de novo* PU.1 motifs

865 in AAM^res
812 in AAM^mono

→ Extend +/- 25bp and identify centrally enriched motifs

PU.1 motif
Collaborating TFs?   Collaborating TFs?

→ DNA shape prediction

Minor groove width (MGW)
Propeller twist (ProT)
Helical twist (HelT)
Roll

Supplemental Figure 2