The community and ecosystem consequences of intraspecific

diversity: a meta-analysis

Allan Raffard[1,2*], Frédéric Santoul[2], Julien Cucherousset[3] and Simon

Blanchet[1,3]

[1]*CNRS, Université Toulouse III Paul Sabatier, Station d'Écologie Théorique et*

Expérimentale du CNRS à Moulis, UMR-5321, 2 route du CNRS, F-09200 Moulis,

France

[2]*UPS, CNRS, INP, EcoLab (Laboratoire d'écologie fonctionnelle et environnement),*

Université de Toulouse, 118 route de Narbonne, UMR-5245, FR-31062 Toulouse Cedex

9, France

[3]*Laboratoire Évolution and Diversité Biologique (EDB UMR 5174), Université de*

Toulouse, CNRS, IRD, UPS, 118 route de Narbonne, Toulouse 31062, France

[*]Author for correspondence at address 1 (Tel.: (+33) 561556747; E-mail:

allan.raffard@sete.cnrs.fr).

ABSTRACT

Understanding the relationships between biodiversity and ecosystem functioning has

major implications. Biodiversity–ecosystem functioning relationships are generally

investigated at the interspecific level, although intraspecific diversity (i.e. within-

species diversity) is increasingly perceived as an important ecological facet of

biodiversity. Here, we provide a quantitative and integrative synthesis testing, across

diverse plant and animal species, whether intraspecific diversity is a major driver of

community dynamics and ecosystem functioning. We specifically tested (*i*) whether the

number of genotypes/phenotypes (i.e. intraspecific *richness*) or the specific identity of

genotypes/phenotypes (i.e. intraspecific *variation*) in populations modulate the structure

of communities and the functioning of ecosystems, (*ii*) whether the ecological effects of

intraspecific richness and variation are strong in magnitude, and (*iii*) whether these

effects vary among taxonomic groups and ecological responses. We found a non-linear

relationship between intraspecific richness and community and ecosystem dynamics

that follows a saturating curve shape, as observed for biodiversity–function

relationships measured at the interspecific level. Importantly, intraspecific richness

modulated ecological dynamics with a magnitude that was equal to that previously

reported for interspecific richness. Our results further confirm, based on a database

containing more than 50 species, that intraspecific variation also has substantial effects

on ecological dynamics. We demonstrated that the effects of intraspecific variation are

twice as high as expected by chance, and that they might have been underestimated

previously. Finally, we found that the ecological effects of intraspecific variation are not

homogeneous and are actually stronger when intraspecific variation is manipulated in

primary producers than in consumer species, and when they are measured at the

ecosystem rather than at the community level. Overall, we demonstrated that the two

facets of intraspecific diversity (richness and variation) can both strongly affect

community and ecosystem dynamics, which reveals the pivotal role of within-species

biodiversity for understanding ecological dynamics.

Key words: ecosystem functioning, community, genotype, phenotype, meta-analysis,

biodiversity, eco-evolutionary dynamics.

CONTENTS

I. INTRODUCTION

Understanding the relationships between biodiversity and ecosystem functioning is an

intensely active field of research informing on the services provided by biodiversity

(Chapin *et al.*, 2000; Loreau, 2000; Hooper *et al.*, 2005; Cardinale *et al.*, 2012).

Biodiversity is generally quantified as the taxonomic, functional and/or phylogenetic

diversity of a species assemblage, and most studies on biodiversity–ecosystem

functioning relationships have to date focused on the interspecific facet of biodiversity

(Naeem *et al.*, 1994; Downing & Leibold, 2002; Hillebrand & Matthiessen, 2009).

However, biodiversity also includes an intraspecific facet that is defined as the

phenotypic, functional and genetic diversity measured within a single species (Odling-

Smee, Laland & Feldman, 2003; Bolnick *et al.*, 2003). During the last two decades,

intraspecific diversity has been demonstrated to account for a non-negligible part of the

total biodiversity measured in plants and animals (Fridley & Grime, 2010; de Bello *et*

al., 2011), representing in some cases up to a quarter of the total variability measured in

communities (Fridley & Grime, 2010; de Bello *et al.*, 2011; Siefert *et al.*, 2015).

In parallel, the hypothesis that intraspecific diversity may affect ecological

dynamics at levels higher than the population level (for instance the composition and

the dynamics of communities and/or the dynamics of ecosystem functions) has been

conceptualized (Bolnick *et al.*, 2003, 2011; Hughes *et al.*, 2008; Bailey *et al.*, 2009;

Violle *et al.*, 2012). These conceptual insights have been validated by several key

experiments both in plants and animals (Whitham *et al.*, 2003; Madritch, Greene &

Lindroth, 2009; Matthews *et al.*, 2016; Rudman & Schluter, 2016). For instance, the

experimental manipulation of fish phenotypes from several evolutionarily independent

fish populations has been shown to generate significant changes in both the community

structure of invertebrate prey and the primary productivity of the ecosystem (Harmon *et*

al., 2009; Matthews *et al.*, 2016).

Intraspecific diversity can be characterized based on the *richness* of populations,

which corresponds to the differences in the number of genotypes and/or phenotypes

composing populations. For instance, populations are often characterized according to their 'allelic, genotypic or phenotypic richness', which is a population parallel of species richness, a common metric measured at the interspecific level and classically used to investigate biodiversity–ecosystem function (BEF) relationships (Crutsinger *et al.*, 2006). Intraspecific richness can also affect ecological dynamics hence generating 'intraspecific BEF' (Whitham *et al.*, 2006; Crutsinger *et al.*, 2006). The basic hypothesis for intraspecific BEF is that increasing the number of genotypes/phenotypes in a population should alter (either negatively or positively) key ecological functions such as the decomposition rate of organic matter or the structure of communities. For instance, experiments manipulating the number of genotypes (from one to 12 genotypes) in plant (*Solidago altissima*) populations have shown that richer populations contained a higher diversity of invertebrates (Crutsinger *et al.*, 2006). Actually, the ecological consequences of intraspecific richness should follow a saturating curve (i.e. a rapid increase followed by a plateau) as often described for BEF observed at the interspecific level (Hughes *et al.*, 2008). Although rarely tested empirically, this saturating shape could be due to the combined effects of several mechanisms. Populations with different richness could have different ecological consequences because of ecological complementarity among genotypes/phenotypes (i.e. niche partitioning, facilitation occurring when genotypes use different resources), inhibition among genotypes/phenotypes (when multiple genotypes are in competition for resources), or functional redundancy among genotypes/phenotypes that can make populations ecologically equivalent (Johnson, Lajeunesse & Agrawal, 2006; Hughes *et al.*, 2008). Yet, the shape of the relationship between intraspecific richness and ecological dynamics has rarely been investigated empirically and to our knowledge has never been quantified across species.

The ecological consequences of intraspecific diversity can also be investigated

through the lens of *variation* in genotypic or phenotypic attributes. Adaptive and non-

adaptive evolutionary processes such as natural selection, plasticity or genetic drift can

generate unique phenotypic differences among populations. These differences can be

associated to key functional processes such as food acquisition or nutrient cycling (e.g.

Grant & Grant, 2006; Rudgers & Whitney, 2006; Lowe, Kovach & Allendorf, 2017),

resulting in both trophic and non-trophic effects of intraspecific variation on ecological

dynamics (Odling-Smee *et al.*, 2003; Whitham *et al.*, 2003; Matthews *et al.*, 2011). For

instance, it has been shown experimentally that plant genotypes differing in their

susceptibility to herbivores harbour different communities of herbivores (Crutsinger,

Cadotte & Sanders, 2009*a*; Barbour *et al.*, 2009*b*). Similarly, mesocosm experiments

have shown that differences in diet within predator populations can modify prey

community structure (Post *et al.*, 2008; Harmon *et al.*, 2009; Howeth *et al.*, 2013). Non-

trophic interactions can also have an important role. For instance, differences in the

chemical composition of individuals can result in differences in excretion rate or in leaf

chemistry that can then affect ecosystem functions such as primary production or

nutrient recycling (Lecerf & Chauvet, 2008; El-Sabaawi *et al.*, 2015). Recently, Des

Roches *et al.* (2018) demonstrated that intraspecific variation can affect ecological

dynamics to the same extent as the removal or replacement of a species in the

environment. Although based on a relatively limited number of studies (25 studies on

15 species), their study confirmed the hypothesis that intraspecific variation might be a

non-negligible driver of ecological dynamics.

Here, we investigated – across various species and ecosystems – the extent to

which both intraspecific richness and intraspecific variation affect the structure of

communities and the functioning of ecosystems, and whether intraspecific diversity is a

major driver of ecological dynamics. We reviewed published studies testing the causal

effects of intraspecific diversity on ecological dynamics in two meta-analyses

synthesizing published data across taxa and ecosystems for intraspecific richness and

variation, respectively, and to fulfil three specific objectives. First, we tested the

significance and the shape of the relationship between intraspecific richness and

ecological dynamics. We expected to find a significant saturating relationship between

intraspecific richness and ecological dynamics, because of potential facilitation and

functional redundancy among genotypes and phenotypes (Hughes *et al.*, 2008). Second,

we tested whether manipulating intraspecific richness has similar effects on ecological

dynamics to manipulating interspecific richness, by comparing the ecological effects of

intraspecific richness with those of interspecific richness obtained from experimental

studies manipulating species richness (Duffy, Godwin & Cardinale, 2017). Finally, we

provided a novel and extensive quantitative synthesis testing for the effects of

intraspecific variation on ecological dynamics. Des Roches *et al.* (2018) previously

focused on studies removing or replacing the target species (by which intraspecific

variation was manipulated) to investigate the ecological consequences of intraspecific

variation. This strongly restricted the number of available studies for which effects sizes

could be calculated, and potentially upwardly biased the resulting estimates (Des

Roches *et al.*, 2018). We here relax this restriction by considering all studies

manipulating intraspecific variation, and use a null-model approach to provide a more

accurate relative effect size of intraspecific variation on ecological dynamics. We also

built on this extended data set to partition variance in the ecological consequences of

intraspecific variation according to the type of organism manipulated and the type of

response variable measured. We tested whether the magnitude of the effects of

intraspecific variation on ecological dynamics vary among organism types (primary

producers *versus* consumers) and levels of biological organization (community *versus*

ecosystem levels). Because primary producers form the basis of trophic chains, we

expect stronger ecological effects of intraspecific variation in producers than in

consumers. We also expect stronger effects of intraspecific variation on ecosystem

functions than on metrics describing community structure because ecosystem functions

are affected by both trophic and non-trophic effects of biodiversity (Matthews *et al.*,

2014).

II. MATERIALS AND METHODS

(1) Data collection

We compiled data from published articles quantifying the effects of intraspecific

diversity in a single species on community structure and/or ecosystem functioning. We

focused only on intraspecific diversity that represented the integrative phenotypic effect

of multiple evolutionary processes including selection, drift and/or plasticity. As a

result, we did not consider articles focusing on experimentally induced intraspecific

diversity through induced plastic responses to particular predatory or environmental

cues [for example see Werner & Peacor (2003) for a review]. We reviewed several

experimental studies manipulating intraspecific variation and/or richness within a single

species to test their respective ecological effects. We also reviewed some observational

studies with strong biological hypotheses and adequate design allowing inferring causal

links from intraspecific diversity to ecological dynamics (e.g. Post *et al.*, 2008). Studies

varying intraspecific diversity within a set of multiple species (e.g. Booth & Grime,

2003) were not included in this meta-analysis. The literature search was carried out

using the ISI *Web of Knowledge* and *Scopus* platforms (last accessed 25th July 2018).

We also scrutinized the reference list of each article to obtain additional articles. The

following key words were used in various combinations: *community genetics* AND

intraspecific variation, *eco-evolutionary dynamics* AND *ecosystem function*, *community*

genetics AND *ecosystem function*, and *intraspecific genetic variation* AND *ecosystem*

function. We selected articles describing the effects of genotypic and/or phenotypic

richness (intraspecific *richness*) and/or different genotypes/phenotypes (intraspecific

variation) in a single target species on community and/or ecosystem dynamics. A total

of 90 studies with available statistics were selected (see online Supporting information,

Appendix S1 and Fig. S1). Among these, 23 studies (100% experimental studies)

focused on intraspecific richness and 75 studies (90% experimental studies, 10%

empirical studies) focused on intraspecific variation.

For each study, we recorded the Latin name of the target species and classified them as

primary producers or consumers (including primary and secondary consumers as well as

predators) and according to the major taxonomic categories represented in our data sets:

arthropods (9 species), fishes (6 species), herbaceous plants (14 species), trees (31

species), and fungi (5 species). Overall, this led to 52 species for studies focusing on

intraspecific variation, and 17 species for studies focusing on intraspecific richness. We

recorded seven main response variables related to community structure and ecosystem

functioning. A community is here defined as a group of at least two species, and we

focused on three types of response variables describing the structure of communities:

(1) species abundance: total number of individuals of all species; (2) biomass: total

mass of individuals of all species; (3) community structure: number of species (e.g.

Simpson or Shannon indices), species evenness and/or species richness.

Regarding response variables at the ecosystem level, we considered four main

ecosystem functions: (1) decomposition rate: rate at which organic matter is recycled;

(2) elemental cycling: quantity of organic or inorganic materials; (3) primary

productivity: measured as several proxies of primary producers: biomass of primary

producers excluding the productivity of the target species, chlorophyll *a* concentration,

daily oxygen production, and photosynthetically active radiation; (4) ecosystem

respiration: rate of oxygen consumption.

(2) Meta-analysis

(*a*) *The ecological consequences of intraspecific richness*

To test for the consequences of intraspecific richness on ecological dynamics, we

focused only on studies investigating the consequences of genotypic richness since this

was the intraspecific diversity facet most commonly manipulated to test for the effects

of intraspecific richness on ecological dynamics. Here, we used the log-transformed

response ratio (ln*RR*) as an effect size. ln*RR* was computed as: $\ln\left(\frac{XG}{1G}\right)$, where *1G* is the

average of the response variable measured for the treatment with a single genotype (i.e.

monoculture), and *XG* is the average of the response variable measured for each

treatment independently including more than one genotype. For each response variable,

ln*RR* increases as the difference in the mean response variable measured in the

treatment with a single genotype and treatments including more than one genotype

increases. We also recorded the difference in the number of genotypes between the

single genotype treatment (monoculture) and all other treatments separately (i.e.

treatments including 2–12 genotypes) as the 'difference in intraspecific richness'. In our

data set, difference in intraspecific richness therefore varied between one and 11

genotypes. This approach allowed quantifying the ecological consequences of

increasing the number of genotypes for each target species. Since each study generally

assessed the effects of intraspecific richness on more than one response variable, our

data set included a total of 135 assays.

We wanted to test the shape and the significance of the relationship between

$\ln RR$ and the difference in intraspecific richness across all case studies. The general

expectation is that ecological differences between treatments increase as differences in

intraspecific richness increase, although this increase may be non-linear (Hughes *et al.*,

2008). We therefore used non-linear mixed-effect models to test the significance and

shape of the relationship between absolute values of $\ln RR$ ($|\ln RR|$) and differences in

intraspecific richness. More precisely, we modelled this relationship using four different

models to determine the most likely shape of the relationship between $|\ln RR|$ and

difference in intraspecific richness: (1) a null model (one parameter) was computed for

the null-effect hypothesis (i.e. no significant relationship between $|\ln RR|$ and difference

in intraspecific richness); (2) a linear model (two parameters) suggesting a positive and

linear relationship between $|\ln RR|$ and difference in intraspecific richness; (3) a

Michaelis–Menten model (two parameters) in which $|\ln RR|$ increases with intraspecific

richness, until a plateau is reached (saturating shape); (4) an asymptotic exponential

model (two parameters) with a shape similar to the Michaelis–Menten model, except

that the plateau is reached sooner.

All models cited above included article ID and the monoculture ID (i.e. the monoculture

treatment to which each other treatment of richness was compared for a given response

variable within each study) as random terms to account for non-independence of effect

sizes (Noble *et al.*, 2017), and the inverse of the sample size as a weighting parameter

giving greater weight to articles including more replicates. Models were compared using

the Akaike information criterion (AIC) and we retained (as "best models") all models

that fell within a ΔAIC < 4 (Burnham & Anderson, 2002). We also calculated for each

model the Akaike weight that provides a conditional probability for each model to be

best supported by the data (Burnham & Anderson, 2002).

We then compared the magnitude (absolute effect size) of ecological effects of

intraspecific and interspecific richness. We extracted from each study and for each

response variable the lnRR corresponding to the most extreme levels of genotypic

richness (x_{max}) manipulated in each study ($N = 63$ lnRR). These values were

subsequently compared to published lnRR values calculated following a similar method

for experiments ($N = 35$) manipulating interspecific richness (Duffy et $al.$, 2017).

Because absolute effect sizes follow a folded-normal distribution, we used an 'analyse

and transform' approach ($sensu$ Morrissey, 2016a,b) to estimate the absolute means of

effect sizes. This approach consists first of estimating the mean and variance of lnRR

(using non-absolute values), and then deriving the mean absolute value from these

estimates. To do so, we estimated the mean of lnRR for interspecific and intraspecific

richness, respectively, using two independent intercepts models with lnRR as the

response variable, article ID as the random effect and the inverse of the sample size as

the weighting parameter. These intercepts models were implemented using the

MCMCglmm package in R (Hadfield, 2010). Markov chain Monte Carlo (MCMC)

chains were run on 15×10^5 iterations, with a burn-in interval of 3×10^4, a thinning

interval of 1×10^3, and an inverse-Wishart prior ($V = 1$ and $\eta = 0.002$). Finally, the

estimated means' lnRR values were converted into absolute-magnitude |lnRR| values

(following Morrissey, 2016b), and we compared the magnitudes of the ecological

effects of interspecific and intraspecific richness based on visual inspection of 95%

percentile intervals (PIs).

Finally, to compare the ecological consequences of intraspecific richness

between levels of biological organization, we performed the same 'analyse and

transform' approach described above. We used a linear mixed-effect model

(implemented in the MCMCglmm package in R, and parameterized similarly than

above) with the lnRR as the dependent variable, the inverse of the sample size as the

weighting parameter, article ID and monoculture ID as random factors, and with the

level of biological organization (community *versus* ecosystem response variables)

treated as a fixed effect. The type of organism was not included in this analysis given

that studies on intraspecific richness focused almost exclusively on primary producers

(with two exceptions on fungi).

(*b*) *The ecological consequences of intraspecific variation*

Given that most studies (86%) did not include a control (i.e. a treatment without the

target species), we compared the strength of effects among all unique genotypes and/or

phenotypes that were considered in each study. Studies generally compared the

consequences of 2–10 different unique genotypes and/or phenotypes (i.e. 2–10

treatments, with each treatment corresponding to a unique genotype/phenotype) on

community and/or ecosystem dynamics; we gathered from each study the statistic (*t*, *F*,

Chi-squared, Pearson's *r*, Spearman's *r*, R^2 or Hedges' *g*) associated with the between-

treatments comparison (i.e. the variation of the phenotypes/genotypes). The higher the

absolute value of the statistic, the higher the community and ecosystem consequences

due to the variation of genotypes and/or phenotypes. The value of each statistic was

converted into a correlation coefficient (*r*) ranging from 0 to 1 (see Table S1 for the

formulae used). We did not use the direction of the statistic (i.e. positive or negative)

because this depended upon the ecological response variable that was considered, which

complicates comparisons on the direction of effects. The *Z*-Fisher transformation then

was used to obtain a standardized effect size using the formula: $Zr = 0.5\ln \frac{(1+r)}{(1-r)}$. For

each *Zr* value, we calculated the corresponding standard error as $se_{Zr} = \frac{1}{\sqrt{n-3}}$

(Nakagawa & Cuthill, 2007). Since each study generally focused on more than one

response variable, we obtained a total of 502 observed Zr values, each corresponding to

the effect size of intraspecific variation observed within one species on a single

response variable. The mean global Zr or mean effect size observed (MES_{obs}) and its

95% confidence interval (CI) were calculated using an intercept-only model. This

intercept-only model was run as a mixed model with no fixed effect, article ID as the

random effect and se_{Zr} included as a weighting parameter to give more weight to

studies with a larger sample size (Koricheva, Gurevitch & Mengersen, 2013).

Because Zr ranged between 0 and $+ \infty$, the CIs of the MES_{obs} do not theoretically

overlap 0, which makes it difficult to assess the significance of the strength of the

MES_{obs}. We therefore used a null-model approach to test if MES_{obs} was significantly

different from that expected under the null hypothesis, i.e. the true effect of intraspecific

variation in all studies was zero. We resampled each statistic (e.g. t, F) from each

empirical study in their respective null distribution with the adequate degree of freedom.

This resampled set of statistics ($N = 502$) was transformed into Zr as described above to

create a set of resampled Zr values. We used this set of resampled Zr values to fit an

intercept-only model with no fixed effect, the corresponding article ID as a random term

and se_{Zr} included as a weighting parameter (as for MES_{obs}). The mean global

resampled Zr (MES_{res}) was extracted from the model, and we repeated this resampling

procedure 1000 times to obtain 1000 values of MES_{res}. This resampled distribution of

1000 MES_{res} approximates the range of possible MES values expected if the null

hypothesis was true. Finally, we calculated the probability of MES_{obs} to be larger than

expected under this null hypothesis using a one-tailed test (Manly, 1997).

We then compared the median of effect sizes (MES_{common}) of studies that were in

common between our extended data set and that used by Des Roches *et al.* (2018) (i.e.

15 studies that were used both in our meta-analysis and that of Des Roches *et al.*) to a

selection of 15 studies randomly sampled from our extended data set (i.e. 75 studies in

our extended data set). We calculated the median effect size for the subset of random

studies (MES$_{ran}$) and repeated this resampling procedure 1000 times to obtain 1000

values of MES$_{ran.}$ We then compared MES$_{common}$ to each MES$_{ran}$ value to calculate the

probability that MES$_{common}$ was higher than a random subset of 15 studies taken from

the whole data set (Manly, 1997).

We then investigated the variability in effect sizes (Zr) and the potential

moderators. We analysed the heterogeneity in effect sizes across articles using the I^2

statistic, which was calculated using an intercept model with the article ID as the

random effect and se_{Zr} as the weighting parameter (Higgins & Thompson, 2002; Senior

et al., 2016). Finally, we tested whether effect sizes (Zr) differed among organism types

with intraspecific variation manipulation, and among the ecological response variables

considered. We hence computed meta-regressions based on linear mixed-effect models

with Zr values (for all 75 studies and 502 measures) as the dependent variable, and

organism type or ecological response variable as fixed effects. The article ID was

included as a random effect, and se_{Zr} was included as a weighting parameter. Four

models were run to assess the differences of effect sizes (*i*) between organism types

classified as consumers or primary producers, and (*ii*) between detailed taxonomic

categories (arthropods, fishes, herbaceous plants and trees). We then tested whether the

effect sizes of intraspecific variation differed among ecological response variables (*iii*)

classified as community or ecosystem variables, and (*iv*) classified according to more

detailed categories (abundance, biomass, community structure, decomposition, nutrient

cycling, primary productivity and respiration of the ecosystem).

(3) Publication bias

For both intraspecific variation and intraspecific richness*,* we assessed potential publication bias by combining Egger's regressions and funnel plots (Egger *et al.*, 1997). Egger's regressions and funnel plots were computed using the residuals of meta-regressions related effect sizes to the main modifiers (i.e. the explanatory variables) and a measure of study size (the inverse of se_{Zr} and sample size for intraspecific variation and intraspecific richness*,* respectively; Horvathova, Nakagawa & Uller, 2012; Nakagawa & Santos, 2012). Typically, for intraspecific variation we ran an Egger's regression model including the residuals of the meta-regression linking intraspecific variation to the modifiers as a response variable and the inverse of se_{Zr} as the explanatory variable. A similar approach was used for intraspecific richness. The intercept α and the slope β of the Egger's regressions are expected not to differ significantly from zero if the data sets are not biased towards significant results. Finally, funnel plots were produced as a scatterplot linking the residuals described above to the respective measure of the study size. An unbiased data set is expected to generate a funnel plot in which articles with larger sample sizes will be close to the mean effect size, whereas articles with small sample sizes will show more variance around the mean effect size (Horvathova *et al.*, 2012; Nakagawa & Santos, 2012).

Overall, and after accounting for important modifiers we found that there was no strong visual sign of publication bias, neither for intraspecific variation nor for intraspecific richness (Fig. S2). This visual inspection of funnel plots was confirmed by the Egger's regressions since parameter values were not significant for intraspecific variation (α = 0.015, *P* = 0.404; β = –0.001, *P* = 0.501) or for intraspecific richness (α= –0.001, *P* = 0.914; β < 0.001, *P* = 0.961).

397   All statistical analyses were performed using the R environment (R core team,

398 2013, Appendix S2). The nlme package (Pinheiro *et al.*, 2014) was used to compute

399 linear and non-linear mixed-effect models, unless specified otherwise.

401 **III. RESULTS**

402 All articles (N = 90) selected for investigating the effects of intraspecific richness and

403 intraspecific variation were published between 2000 and 2018, and 74% used primary

404 producers as target species (Fig. 1). The first studies focusing on consumers were

405 published in 2008, using fish (60%), arthropods (32%), and fungi (8%) as model

406 species.

408 **(1) The intraspecific richness–ecological dynamics relationship**

409 As expected, we found a significant, positive and non-linear relationship between

410 intraspecific richness and ecological dynamics that approximated a saturating curve

411 (Fig. 2A). The AIC selection procedure revealed that one out of the four tested models

412 was highly likely to be supported by the data (i.e. 99% chance of being the best fitting

413 model according to the Akaike weight and ΔAIC > 15.721 for the other models, Table

414 1). The model that best supported the data was the exponential asymptotic model,

415 suggesting that the relationship between intraspecific richness and changes in

416 community structure and ecosystem functioning (i.e. effect size: ln*RR*) likely followed a

417 saturating shape (Fig. 2A).

418   We further found that the ecological effects of intraspecific richness were similar

419 to the ecological effects induced by interspecific richness (Fig. 2B). Indeed, the two

420 distributions largely overlapped and the estimated means were similar (intraspecific

421 richness |ln*RR*| = 0.132, PI = 0.048–0.216; interspecific richness |ln*RR*| = 0.134 PI =

0.012–0.462). The ecological effects of intraspecific richness tended to be higher,

although the difference was not significant, for community metrics ($|\ln RR| = 0.156$, PI =

0.070–0.242) than for ecosystem metrics ($|\ln RR| = 0.045$, PI = 0.004–0.137) (see Fig. S3

for details of ecological metrics).

(2) The ecological consequences of intraspecific variation

We extended the meta-analysis performed by Des Roches *et al.* (2018) to 52 species (15

species were used in Des Roches *et al.*, 2018). We found that that the observed effect

size of intraspecific variation on community structure and ecosystem dynamics was

significant, and was twice as large as the resampled effect size expected under the null

expectation ($MES_{obs} = 0.521$, 95% confidence interval (CI) = 0.444–0.598; $MES_{null} =$

0.259, CI = 0.258–0.259; resampled test, $P < 0.001$; see Figs 3 and S4).

We tested the extent to which the more restricted data set of Des Roches *et al.*

(2018) was representative of our extended data set, or whether it was upwardly biased

as expected by Des Roches *et al.* (2018). We found that that effect sizes for studies in

common with the Des Roches *et al.* (2018) data set ($MES_{common} = 0.299$, 95% percentile

interval (PI) = 0.033–1.092) were not significantly different from the distribution of

effect sizes measured in our extended data set ($MES_{ran} = 0.418$, PI = 0.255–0.616;

resampling test, $P = 0.118$; Fig. S5), and in fact showed a tendency to be downwardly

biased.

Finally, a relatively low heterogeneity in effect size (*Zr*) was detected across

articles ($I^2 = 0.151$). The ecological effects induced by intraspecific variation were

stronger when primary producers rather than consumers were manipulated ($F = 3.968$

d.f. = 1, 425, $P = 0.047$; Fig. 4A). Nonetheless, the strongest ecological effects of

intraspecific variation tended to be observed in arthropods and herbaceous species,

whereas the smallest effects were observed in fish and tree species ($F = 2.475$ d.f. = 3,

417, $P = 0.061$; Fig. 4A). Irrespective of organism type, the effects of intraspecific

variation were significantly stronger when the response variables were measured at the

ecosystem level rather than at the community level ($F = 7.295$, d.f = 1, 425, $P = 0.007$;

Fig. 4B). The strongest effects were detected when response variables concerned

nutrient cycling and the assembly of community, whereas the lowest effects were found

for general measures of abundance and density of species ($F = 2.725$, d.f = 6, 417, $P =$

0.013; Fig. 4B).

IV. DISCUSSION

Intraspecific diversity is increasingly recognized as an important facet of biodiversity

that can affect all biological levels (Bailey *et al.*, 2009). Several studies have

experimentally tested the ecological effects of intraspecific diversity, and we here

provide the first global and quantitative estimates of the consequences of intraspecific

richness and variation on community structure and ecosystem functioning. We

demonstrated for the first time that the intraspecific BEF followed – as theoretically

expected – a non-linear saturating curve with a plateau at 4–6 genotypes per population.

Importantly, we demonstrated also for the first time that intraspecific richness affects

community and ecosystem dynamics with a magnitude comparable to that of

biodiversity measured at the species level. We further confirmed and extended the result

that genotypic and/or phenotypic variation observed between populations can have non-

negligible effects on community structure and ecosystem functions, and we

demonstrated that previous estimates (Des Roches *et al.*, 2018) of these ecological

effects of intraspecific variation actually tended to be underestimated. Finally, our

exhaustive quantitative survey identified that the ecological consequences of

intraspecific variation differ among biological level of organization, and among

organism types. These findings provide novel and integrative insights, as well as

multiple research perspectives, into the ecological role of intraspecific diversity.

(1) Intraspecific diversity and the dynamics of communities and ecosystems

Although the form of the relationship between intraspecific richness and

ecological consequences has already been discussed conceptually (Hughes *et al.*, 2008),

our meta-analysis provides for the first time a qualitative and quantitative assessment of

intraspecific BEF measured experimentally. Specifically, although considering mostly

primary producers, our results demonstrated that an increase in intraspecific richness

resulted in a non-linear (saturating) increase in the magnitude of its effects on ecological

dynamics. This finding supports the idea that ecological divergence between an

environment hosting populations composed of a single genotype and an environment

hosting populations composed of multiple genotypes increases until a plateau is reached

as the number of genotypes increases. This result echoes the BEF as defined at the

interspecific level (Reiss *et al.*, 2009; Cadotte, Carscadden & Mirotchnick, 2011) and

suggests that the saturating shape might arise from similar mechanisms occurring at the

intraspecific and interspecific levels (Johnson *et al.*, 2006; Hughes *et al.*, 2008). More

specifically, the initial linear increase is assumed to be due to complementarity and

facilitation among genotypes, whereas the plateau likely occurs due to functional

redundancy among genotypes (Johnson *et al.*, 2006). Redundant genotypes probably

display functionally similar traits since two genotypes do not necessarily generate two

functionally different traits (e.g. through synonymous mutations or trait convergences).

Thus manipulating trait richness rather than genotypic richness, or more precisely

manipulating functional effect traits [i.e. traits with ecological effects (Violle *et al.*,

2007)], in future experiments should allow us to explore the mechanisms underlying the

intraspecific biodiversity–ecological dynamics relationship.

We found that effect sizes for intraspecific richness were very similar to values

reported recently reported for experimental interspecific BEF, indicating that the

ecological effects of varying phenotypic/genotypic richness within populations are close

to those induced by varying species richness within communities. This finding raises

several questions regarding the general relationships among intraspecific diversity,

community structure, ecosystem functioning and common abiotic constraints. A large

body of literature has demonstrated that intraspecific genetic diversity and species

diversity (a measure of community structure) might co-vary because of common

environmental drivers and/or reciprocal causal relationships between intraspecific

genetic diversity and species diversity [i.e. the species–genetic diversity correlation

(SGDC) framework (Vellend & Geber, 2005; Vellend, 2005)]. Because most studies

considered in our meta-analysis are experimental, our findings confirm that intraspecific

diversity can directly influence the structure of communities irrespective of the abiotic

environments, hence adding weight to the SGDC framework. Additionally, we suggest

expanding the SGDC framework since intraspecific diversity can also affect ecosystem

functioning. This suggests that intraspecific diversity, community structure and

ecosystem functioning may actually be tightly linked in a tripartite relationship. A major

future challenge will be to tease apart the causal relationships linking these three

components within a common abiotic environment. These relationships might be direct

(e.g. intraspecific diversity directly affects community structure), indirect (e.g.

intraspecific diversity indirectly affects ecosystem functions through its direct effect on

community structure such as the trophic cascade), and/or due to the parallel effects of

common abiotic drivers (e.g. temperature directly affects intraspecific diversity,

community structure and ecosystem functions). As has been done recently for the BEF

(Grace *et al.*, 2016; Duffy *et al.*, 2016) and the SGDC (Fourtune *et al.*, 2016; Lamy *et*

al., 2017) frameworks, we argue that a future important step will be to combine

powerful statistical methods (e.g. path analysis; Shipley, 2000; Grace, 2006) with

appropriate experimental designs to disentangle causal relationships between

intraspecific diversity, community structure, ecosystem functions and their common

environment.

We further demonstrated that intraspecific variation has significant ecological

effects across a large set of species (52 species and 75 articles), hence confirming and

refining a previous estimate based on a more restricted species set (Des Roches *et al.*,

2018). By more than tripling the number of species being investigated in this meta-

analysis, we extend the conclusion to a greater taxonomic set that intraspecific variation

is involved in shaping ecological dynamics, and that the ecological effects of

intraspecific variation might be more common than expected. Moreover, we

demonstrated that previous estimates (Des Roches *et al.*, 2018) were not upwardly

biased (as expected from their focus on a non-random species pool), but were well

within the range of estimates we report here and actually tended to be slight

underestimates. Our finding hence strongly supports the idea that adaptive and non-

adaptive processes can lead to unique populations differentially and significantly

affecting ecological systems.

Although our conclusions held true for many species, the ecological effects of

intraspecific variation were not homogeneous across species, and this was partly

explained by their trophic level. Indeed, and according to expectations, the ecological

effects of intraspecific variation were stronger when the target species was a primary

producer than when it was a consumer. Several non-exclusive mechanisms might

explain this result. For instance, many primary producers considered here provide a

habitat for many invertebrate species (Southwood, Brown & Reader, 1979) (this is not

the case for the consumer species), and this habitat can be modulated by changes in

plant structure. The relative biomass of primary producers is higher than that of

consumers, thus primary producers could generate stronger effects on communities and

ecosystems than consumers simply because of this biomass effect. However, a more

detailed analysis showed that the effects of intraspecific variation tended to be stronger

for arthropod and herbaceous species than for fish (and to a lesser extent tree) species.

This suggests that the trophic level of a species may not be the only predictor of the

ecological effects of intraspecific diversity, and we argue that future work should aim to

test specifically why intraspecific variation matters more for some species than others.

Finally, the effects of intraspecific variation were globally higher for ecosystems

than for communities, hence generalizing across organism and ecosystem types a

previous conclusion for freshwater consumer species (Palkovacs *et al.*, 2015). We can

speculate that this difference arises because intraspecific variation acts on community

dynamics through trophic mechanisms, whereas ecosystem functions can be modulated

through both trophic and non-trophic interactions [e.g. excretion rate or leaf chemistry

(Vanni, 2002; Schmitz *et al.*, 2014)]. For instance, a consumer species that shows

intraspecific variation in resource selectivity and/or consumption rate could affect both

the community structure and productivity of its resource (Harmon *et al.*, 2009). Non-

trophic mechanisms such as variability in organismal stoichiometry could reinforce the

effect of the consumer species on several ecosystem functions [e.g. primary production

or soil mineralization (Schmitz, Hawlena & Trussell, 2010; Hawlena *et al.*, 2012)].

Alternatively, changes in ecosystem functions might be due to both direct effects of

intraspecific diversity and indirect effects of intraspecific diversity mediated through

changes in community structure, which may overall strengthen the effects of

intraspecific diversity at the ecosystem level. However, a more detailed analysis

revealed that much variation exists between sub-categories of response variables (Fig.

4B), and that the dichotomy between variables measured at the community and

ecosystem levels is not straightforward. Although providing the first attempts to

separate variance in the ecological effects of intraspecific variation into its component

parts, our findings call for further studies on various taxa and in different ecosystems in

order to understand fully the ecological effects of intraspecific diversity.

(2) Research perspectives on the relationships between intraspecific diversity and
the dynamics of communities and ecosystems

We highlight several research avenues that may greatly enhance our understanding of

the relationships between intraspecific diversity and the dynamics of communities and

ecosystems.

Our review demonstrates that the ecological effect size of intraspecific diversity varies

among species and that this interspecific variance in effect size can be partly explained

by the type of organism (i.e. primary producer or consumer). However, species

composing a community also vary according to abundance, role in the ecosystem (e.g.

keystone species; Paine, 1969), body size (or height for plants), life-history strategy

(e.g. r–K strategy), recent history (e.g. whether the species is non-native), functional

characteristics (e.g. stoichiometry, metabolism), etc. The next challenge will be to

partition this interspecific variance in effect size better by determining the species

characteristics that best predict the strength of effect sizes; this is a pre-requisite to

design coherent conservation plans at the intraspecific level (Mimura *et al.*, 2016).

Intraspecific diversity is often manipulated with respect to a single target

species. However, in nature, species are interacting and we argue that future studies

should manipulate intraspecific diversity within multiple interacting species to reflect

natural conditions, and to allow partitioning the relative importance of intraspecific

diversity between interacting (and potentially co-evolving) species.

Ecosystems are interconnected through cross-ecosystem fluxes of matter

(Loreau, Mouquet & Holt, 2003). For instance, freshwater ecosystems such as rivers

receive a large amount of material from surrounding terrestrial ecosystems [e.g. dead

leaves falling from trees (Bartels *et al.*, 2012)]. A major future challenge would be to

assess the relative effects of intraspecific diversity on allochtonous ecosystems *versus*

autochtonous ecosystems; for example testing whether the consequences of intraspecific

diversity manipulation in a fish species are greater in associated aquatic ecosystems than

on nearby terrestrial ecosystems [see Jackrel & Wootton (2014), Crutsinger *et al.*

(2014)*b* and Rudman *et al.* (2015) for attempts to quantify across-ecosystem effects of

intraspecific diversity]. This is an important next step to quantify in a more integrative

way the importance of intraspecific diversity in natural systems.

Previous studies have mainly assessed the consequences of intraspecific

diversity by considering the genetic or the phenotypic characteristics of organisms as a

whole. However, some traits have been shown to be more important for ecological

dynamics than others; this is the case for functional effect traits that are defined as traits

with an ecological impact [e.g. excretion rate or leaf nutrient content (Violle *et al.*,

2007; Díaz *et al.*, 2013)]. We suggest that direct manipulation of the functional trait

diversity of individuals within a species, rather than their genotypic or phenotypic

diversity as a whole would be a powerful means to understanding the mechanisms by

which intraspecific diversity acts on ecological dynamics.

Although we emphasize that intraspecific diversity is an essential component of

the community and ecosystem facets of ecological dynamics, comparisons with the

effects of other major ecological constraints (e.g. temperature, interspecific biodiversity,

human disturbance) have rarely been conducted (but see Burkle *et al.*, 2013; El-Sabaawi

et al., 2015). To evaluate and quantify the importance of intraspecific diversity in

natural systems better, it is important to assess the relative contributions of intraspecific

diversity and other determinants of ecosystem functioning. This will be an essential step

in confirming that intraspecific variation is a key determinant, and not just a random

signal in complex systems.

V. CONCLUSIONS

(1) This study provides empirical evidence that the ecological effects of intraspecific

richness increase asymptotically, paralleling well-known patterns observed at the

interspecific level (Loreau, 2000; Hooper *et al.*, 2005) and confirming previous

hypotheses (Hughes *et al.*, 2008).

(2) We found that experimental manipulations of intraspecific richness caused

community and ecosystem differentiations as large as those generated by interspecific

richness. This suggests that variation in richness within populations can have similar

ecological consequences to variation in richness among species.

(3) We also demonstrated that variation in phenotypes or genotypes within species is an

important driver of community and ecosystem dynamics. These major ecological effects

of intraspecific diversity held true for a range of organisms including plants and

animals, although much remains to be tested.

(4) Overall these findings demonstrate that intraspecific diversity - beyond its

importance for species to adapt to environmental changes - is an important facet of

biodiversity for understanding and predicting the ecological dynamics of communities

and ecosystems, reinforcing the need for a better appraisal of the causes and

consequences of intraspecific diversity in natural populations and for improved

conservation plans.

VI. ACKNOWLEDGEMENTS

We warmly thank José M. Montoya for valuable comments on a previous version of this

review. We also thank three anonymous reviewers for their constructive and useful

comments. A.R. was supported financially by a Doctoral scholarship from the

Université Fédérale de Toulouse. This work was undertaken at SETE, which is part of

the "Laboratoire d'Excellence" (LABEX) entitled TULIP (ANR-10-LABX-41).

VII. REFERENCES
659

ABBOTT, J.M., GROSBERG, R.K., WILLIAMS, S.L. & STACHOWICZ, J.J. (2017). Multiple
dimensions of intraspecific diversity affect biomass of eelgrass and its
associated community. *Ecology* **98**, 3152–3164.

BAILEY, J.K., SCHWEITZER, J.A., UBEDA, F., KORICHEVA, J., LEROY, C.J., MADRITCH,
M.D., REHILL, B.J., BANGERT, R.K., FISCHER, D.G., ALLAN, G.J. & WHITHAM,
T.G. (2009). From genes to ecosystems: a synthesis of the effects of plant
genetic factors across levels of organization. *Philosophical Transactions of the*
Royal Society B: Biological Sciences **364**, 1607–1616.

BARBOUR, M.A., FORTUNA, M.A., BASCOMPTE, J., NICHOLSON, J.R., JULKUNEN-TIITTO,
R., JULES, E.S. & CRUTSINGER, G.M. (2016). Genetic specificity of a plant–
insect food web: Implications for linking genetic variation to network
complexity. *Proceedings of the National Academy of Sciences* **113**, 2128–2133.

BARBOUR, R.C., FORSTER, L.G., BAKER, S.C., STEANE, D.A. & POTTS, B.M. (2009*a*).
Biodiversity consequences of genetic variation in bark characteristics within a
foundation tree species. *Conservation Biology* **23**, 1146–1155.

BARBOUR, R.C., O'REILLY-WAPSTRA, J.M., DE LITTLE, D.W., JORDAN, G.J., STEANE,
D.A., HUMPHREYS, J.R., BAILEY, J.K., WHITHAM, T.G. & POTTS, B.M. (2009*b*).
A geographic mosaic of genetic variation within a foundation tree species and its
community-level consequences. *Ecology* **90**, 1762–1772.

BARBOUR, R.C., STORER, M.J. & POTTS, B.M. (2009*c*). Relative importance of tree
genetics and microhabitat on macrofungal biodiversity on coarse woody debris.
Oecologia **160**, 335–342.

BARRIOS-GARCIA, M.N., RODRIGUEZ-CABAL, M.A., RUDGERS, J.A. & CRUTSINGER,
G.M. (2016). Soil fertilization does not alter plant architectural effects on
arthropod communities. *Journal of Plant Ecology*, rtw087.

BARTELS, P., CUCHEROUSSET, J., STEGER, K., EKLÖV, P., TRANVIK, L.J. & HILLEBRAND,
H. (2012). Reciprocal subsidies between freshwater and terrestrial ecosystems
structure consumer resource dynamics. *Ecology* **93**, 1173–1182.

BASSAR, R.D., MARSHALL, M.C., LOPEZ-SEPULCRE, A., ZANDONA, E., AUER, S.K.,
TRAVIS, J., PRINGLE, C.M., FLECKER, A.S., THOMAS, S.A., FRASER, D.F. &
REZNICK, D.N. (2010). Local adaptation in Trinidadian guppies alters ecosystem
processes. *Proceedings of the National Academy of Sciences* **107**, 3616–3621.

BOLNICK, D.I., AMARASEKARE, P., ARAÚJO, M.S., BÜRGER, R., LEVINE, J.M., NOVAK,
M., RUDOLF, V.H., SCHREIBER, S.J. & URBAN, M.C. (2011). Why intraspecific
trait variation matters in community ecology. *Trends in ecology & evolution* **26**,
183–192.

BOLNICK, D.I., SVANBÄCK, R., FORDYCE, J.A., YANG, L.H., DAVIS, J.M., HULSEY, C.D.
& FORISTER, M.L. (2003). The ecology of individuals: incidence and
implications of individual specialization. *The American Naturalist* **161**, 1–28.

BOOTH, R.E. & GRIME, J.P. (2003). Effects of genetic impoverishment on plant
community diversity. *Journal of Ecology* **91**, 721–730.

BURKLE, L.A., SOUZA, L., GENUNG, M.A. & CRUTSINGER, G.M. (2013). Plant genotype,
nutrients, and G X E interactions structure floral visitor communities. *Ecosphere*
4.

BURNHAM, K.P. & ANDERSON, D.R. (2002). *Model selection and multimodel inference:
a practical information-theoretic approach.* Springer, New York.

BUSTOS-SEGURA, C., POELMAN, E.H., REICHELT, M., GERSHENZON, J. & GOLS, R.
(2017). Intraspecific chemical diversity among neighbouring plants correlates
positively with plant size and herbivore load but negatively with herbivore
damage. *Ecology Letters* **20**, 87–97.

CADOTTE, M.W., CARSCADDEN, K. & MIROTCHNICK, N. (2011). Beyond species:
functional diversity and the maintenance of ecological processes and services.
Journal of Applied Ecology **48**, 1079–1087.

CAMPOS-NAVARRETE, M.J., MUNGUÍA-ROSAS, M.A., ABDALA-ROBERTS, L., QUINTO, J.
& PARRA-TABLA, V. (2015). Effects of tree genotypic diversity and species
diversity on the arthropod community associated with big-leaf mahogany.
Biotropica **47**, 579–587.

CARDINALE, B.J., DUFFY, J.E., GONZALEZ, A., HOOPER, D.U., PERRINGS, C., VENAIL, P.,
NARWANI, A., MACE, G.M., TILMAN, D., WARDLE, D.A., KINZIG, A.P., DAILY,

G.C., LOREAU, M., GRACE, J.B., LARIGAUDERIE, A., ET AL. (2012). Biodiversity
loss and its impact on humanity. *Nature* **486**, 59–67.

CHAPIN, S.F., ZAVALETA, E.S., EVINER, V.T., NAYLOR, R.L., VITOUSEK, P.M.,
REYNOLDS, H.L., HOOPER, D.U., LAVOREL, S., SALA, O.E., HOBBIE, S.E., MACK,
M.C. & DÍAZ, S. (2000). Consequences of changing biodiversity. *Nature* **405**,
234–242.

CHISLOCK, M.F., SARNELLE, O., OLSEN, B.K., DOSTER, E. & WILSON, A.E. (2013).
Large effects of consumer offense on ecosystem structure and function. *Ecology*
94, 2375–2380.

CLASSEN, A.T., CHAPMAN, S.K., WHITHAM, T.G., HART, S.C. & KOCH, G.W. (2007).
Genetic-based plant resistance and susceptibility traits to herbivory influence
needle and root litter nutrient dynamics. *Journal of Ecology* **95**, 1181–1194.

COOK-PATTON, S.C., MCART, S.H., PARACHNOWITSCH, A.L., THALER, J.S. &
AGRAWAL, A.A. (2011). A direct comparison of the consequences of plant
genotypic and species diversity on communities and ecosystem function.
Ecology **92**, 915–923.

CORNELISSEN, J.H.C., PEREZ-HARGUINDEGUY, N., GWYNN-JONES, D., DIAZ, S.,
CALLAGHAN, T.V. & AERTS, R. (2000). Autumn leaf colours as indicators of
decomposition rate in sycamore (*Acer pseudoplatanus L.*). *Plant and Soil* **225**,
33–38.

CRAWFORD, K.M. & RUDGERS, J.A. (2013). Genetic diversity within a dominant plant
outweighs plant species diversity in structuring an arthropod community.
Ecology **94**, 1025–1035.

CRUTSINGER, G.M., CADOTTE, M.W. & SANDERS, N.J. (2009*a*). Plant genetics shapes
inquiline community structure across spatial scales. *Ecology Letters* **12**, 285–
292.

CRUTSINGER, G.M., CARTER, B.E. & RUDGERS, J.A. (2013). Soil nutrients trump
intraspecific effects on understory plant communities. *Oecologia* **173**, 1531–
1538.

CRUTSINGER, G.M., COLLINS, M.D., FORDYCE, J.A., GOMPERT, Z., NICE, C.C. &
SANDERS, N.J. (2006). Plant genotypic diversity predicts community structure
and governs an ecosystem process. *Science* **313**, 966–968.

CRUTSINGER, G.M., COLLINS, M.D., FORDYCE, J.A. & SANDERS, N.J. (2008*a*).
Temporal dynamics in non-additive responses of arthropods to host-plant
genotypic diversity. *Oikos* **117**, 255–264.

CRUTSINGER, G.M., REYNOLDS, W.N., CLASSEN, A.T. & SANDERS, N.J. (2008*b*).
Disparate effects of plant genotypic diversity on foliage and litter arthropod
communities. *Oecologia* **158**, 65–75.

CRUTSINGER, G.M., RODRIGUEZ-CABAL, M.A., RODDY, A.B., PEAY, K.G., BASTOW,
J.L., KIDDER, A.G., DAWSON, T.E., FINE, P.V.A. & RUDGERS, J.A. (2014*a*).

Genetic variation within a dominant shrub structures green and brown community assemblages. *Ecology* **95**, 387–398.

CRUTSINGER, G.M., RUDMAN, S.M., RODRIGUEZ-CABAL, M.A., MCKOWN, A.D., SATO, T., MACDONALD, A.M., HEAVYSIDE, J., GERALDES, A., HART, E.M., LEROY, C.J. & EL-SABAAWI, R.W. (2014*b*). Testing a 'genes-to-ecosystems' approach to understanding aquatic-terrestrial linkages. *Molecular Ecology* **23**, 5888–5903.

CRUTSINGER, G.M., SANDERS, N.J. & CLASSEN, A.T. (2009*b*). Comparing intra- and inter-specific effects on litter decomposition in an old-field ecosystem. *Basic and Applied Ecology* **10**, 535–543.

DE BELLO, F., LAVOREL, S., ALBERT, C.H., THUILLER, W., GRIGULIS, K., DOLEZAL, J., JANECEK, S. & LEPS, J. (2011). Quantifying the relevance of intraspecific trait variability for functional diversity. *Methods in Ecology and Evolution* **2**, 163–174.

DE GRAAFF, M.-A., SIX, J., JASTROW, J.D., SCHADT, C.W. & WULLSCHLEGER, S.D. (2013). Variation in root architecture among switchgrass cultivars impacts root decomposition rates. *Soil Biology & Biochemistry* **58**, 198–206.

DES ROCHES, S., POST, D.M., TURLEY, N.E., BAILEY, J.K., HENDRY, A.P., KINNISON, M.T., SCHWEITZER, J.A. & PALKOVACS, E.P. (2018). The ecological importance of intraspecific variation. *Nature Ecology & Evolution* **2**, 57–64.

DES ROCHES, S., SHURIN, J.B., SCHLUTER, D. & HARMON, L.J. (2013). Ecological and evolutionary effects of stickleback on community structure. *PLoS ONE* **8**, e59644.

DÍAZ, S., PURVIS, A., CORNELISSEN, J.H.C., MACE, G.M., DONOGHUE, M.J., EWERS, R.M., JORDANO, P. & PEARSE, W.D. (2013). Functional traits, the phylogeny of function, and ecosystem service vulnerability. *Ecology and Evolution* **3**, 2958–2975.

DOWNING, A.L. & LEIBOLD, M.A. (2002). Ecosystem consequences of species richness and composition in pond food web. *Nature* **416**, 837–841.

DUFFY, J.E., GODWIN, C.M. & CARDINALE, B.J. (2017). Biodiversity effects in the wild are common and as strong as key drivers of productivity. *Nature* **549**, 261–264.

DUFFY, J.E., LEFCHECK, J.S., STUART-SMITH, R.D., NAVARRETE, S.A. & EDGAR, G.J. (2016). Biodiversity enhances reef fish biomass and resistance to climate change. *Proceedings of the National Academy of Sciences* **113**, 6230–6235.

EGGER, M., SMITH, G.D., SCHNEIDER, M. & MINDER, C. (1997). Bias in meta-analysis detected by a simple graphical test. *Bmj* **315**, 629–634.

EL-SABAAWI, R.W., BASSAR, R.D., RAKOWSKI, C., MARSHALL, M.C., BRYAN, B.L., THOMAS, S.N., PRINGLE, C., REZNICK, D.N. & FLECKER, A.S. (2015). Intraspecific phenotypic differences in fish affect ecosystem processes as much as bottom-up factors. *Oikos* **124**, 1181–1191.

FARKAS, T.E., MONONEN, T., COMEAULT, A.A., HANSKI, I. & NOSIL, P. (2013).
Evolution of camouflage drives rapid ecological change in an insect community.
Current Biology **23**, 1835–1843.

FOURTUNE, L., PAZ-VINAS, I., LOOT, G., PRUNIER, J.G. & BLANCHET, S. (2016).
Lessons from the fish: a multi-species analysis reveals common processes
underlying similar species-genetic diversity correlations. *Freshwater Biology* **61**,
1830–1845.

FRIDLEY, J.D. & GRIME, J.P. (2010). Community and ecosystem effects of intraspecific
genetic diversity in grassland microcosms of varying species diversity. *Ecology*
91, 2272–2283.

FRYXELL, D.C., ARNETT, H.A., APGAR, T.M., KINNISON, M.T. & PALKOVACS, E.P.
(2015). Sex ratio variation shapes the ecological effects of a globally introduced
freshwater fish. *Proceedings of the Royal Society B: Biological Sciences* **282**,
20151970.

GENUNG, M.A., LESSARD, J.-P., BROWN, C.B., BUNN, W.A., CREGGER, M.A.,
REYNOLDS, W.N., FELKER-QUINN, E., STEVENSON, M.L., HARTLEY, A.S.,
CRUTSINGER, G.M., SCHWEITZER, J.A. & BAILEY, J.K. (2010). Non-additive
effects of genotypic diversity increase floral abundance and abundance of floral
visitors. *Plos One* **5**, e8711.

GOITOM, E., KILSDONK, L.J., BRANS, K., JANSEN, M., LEMMENS, P. & DE MEESTER, L.
(2018). Rapid evolution leads to differential population dynamics and top-down
control in resurrected *Daphnia* populations. *Evolutionary Applications* **11**, 96–
111.

GRACE, J.B. (2006). *Structural equation modeling and natural systems.* Cambridge
University Press.

GRACE, J.B., ANDERSON, T.M., SEABLOOM, E.W., BORER, E.T., ADLER, P.B., HARPOLE,
W.S., HAUTIER, Y., HILLEBRAND, H., LIND, E.M., PÄRTEL, M., BAKKER, J.D.,
BUCKLEY, Y.M., CRAWLEY, M.J., DAMSCHEN, E.I., DAVIES, K.F., ET AL. (2016).
Integrative modelling reveals mechanisms linking productivity and plant species
richness. *Nature* **529**, 390–393.

GRANT, P.R. & GRANT, B.R. (2006). Evolution of character displacement in darwin's
finches. *Science* **313**, 224–226.

HADFIELD, J.D. (2010). MCMC methods for multi-response generalized linear mixed
models: the MCMCglmm R package. *Journal of Statistical Software* **33**, 1–22.

HARMON, L.J., MATTHEWS, B., DES ROCHES, S., CHASE, J.M., SHURIN, J.B. &
SCHLUTER, D. (2009). Evolutionary diversification in stickleback affects
ecosystem functioning. *Nature* **458**, 1167–1170.

HARRISON, J.G., PHILBIN, C.S., GOMPERT, Z., FORISTER, G.W., HERNANDEZ-ESPINOZA,
L., SULLIVAN, B.W., WALLACE, I.S., BELTRAN, L., DODSON, C.D., FRANCIS, J.S.,
SCHLAGETER, A., SHELEF, O., YOON, S.A. & FORISTER, M.L. (2018).
Deconstruction of a plant-arthropod community reveals influential plant traits

839 with nonlinear effects on arthropod assemblages. *Functional Ecology* **32**, 1317–
840 1328.

841 HAWLENA, D., STRICKLAND, M.S., BRADFORD, M.A. & SCHMITZ, O.J. (2012). Fear of
842 predation slows plant-litter decomposition. *Science* **336**, 1434–1438.

843 HIGGINS, J.P.T. & THOMPSON, S.G. (2002). Quantifying heterogeneity in a meta-
844 analysis. *Statistics in Medicine* **21**, 1539–1558.

845 HILLEBRAND, H. & MATTHIESSEN, B. (2009). Biodiversity in a complex world:
846 consolidation and progress in functional biodiversity research. *Ecology Letters*
847 **12**, 1405–1419.

848 HINES, J., REYES, M., MOZDER, T.J. & GESSNER, M.O. (2014). Genotypic trait variation
849 modifies effects of climate warming and nitrogen deposition on litter mass loss
850 and microbial respiration. *Global Change Biology* **20**, 3780–3789.

851 HOOPER, D.U., CHAPIN, F.S., EWEL, J.J., HECTOR, A., INCHAUSTI, P., LAVOREL, S.,
852 LAWTON, J.H., LODGE, D.M., LOREAU, M., NAEEM, S., SCHMID, B., SETÄLÄ, H.,
853 SYMSTAD, A.J., VANDERMEER, J. & WARDLE, D.A. (2005). Effects of
854 biodiversity on ecosystem functioning: a consensus of current knowledge.
855 *Ecological Monographs* **75**, 3–35.

856 HORVATHOVA, T., NAKAGAWA, S. & ULLER, T. (2012). Strategic female reproductive
857 investment in response to male attractiveness in birds. *Proceedings of the Royal*
858 *Society B: Biological Sciences* **279**, 163–170.

859 HOWETH, J.G., WEIS, J.J., BRODERSEN, J., HATTON, E.C. & POST, D.M. (2013).
860 Intraspecific phenotypic variation in a fish predator affects multitrophic lake
861 metacommunity structure. *Ecology and Evolution* **3**, 5031–5044.

862 HUANG, J., LIU, M., CHEN, X., CHEN, J., LI, H. & HU, F. (2015). Effects of intraspecific
863 variation in rice resistance to aboveground herbivore, brown planthopper, and
864 rice root nematodes on plant yield, labile pools of plant, and rhizosphere soil.
865 *Biology and Fertility of Soils* **51**, 417–425.

866 HUGHES, A.R. (2014). Genotypic diversity and trait variance interact to affect marsh
867 plant performance. *Journal of Ecology* **102**, 651–658.

868 HUGHES, A.R., INOUYE, B.D., JOHNSON, M.T.J., UNDERWOOD, N. & VELLEND, M.
869 (2008). Ecological consequences of genetic diversity. *Ecology Letters* **11**, 609–
870 623.

871 INGRAM, T., SVANBÄCK, R., KRAFT, N.J.B., KRATINA, P., SOUTHCOTT, L. & SCHLUTER,
872 D. (2012). Intraguild predation drives evolutionary niche shift in threespine
873 stickleback. *Evolution* **66**, 1819–1832.

874 ITO, M. & OZAKI, K. (2005). Response of a gall wasp community to genetic variation in
875 the host plant *Quercus crispula*: a test using half-sib families. *Acta Oecologica*
876 **27**, 17–24.

JACKREL, S.L. & WOOTTON, J.T. (2014). Local adaptation of stream communities to
intraspecific variation in a terrestrial ecosystem subsidy. *Ecology* **95**, 37–43.

JOHNSON, M.T. & AGRAWAL, A.A. (2005). Plant genotype and environment interact to
shape a diverse arthropod community on evening primrose (*Oenothera biennis*).
Ecology **86**, 874–885.

JOHNSON, M.T.J., LAJEUNESSE, M.J. & AGRAWAL, A.A. (2006). Additive and interactive
effects of plant genotypic diversity on arthropod communities and plant fitness.
Ecology Letters **9**, 24–34.

JOHNSON, M.T.J., VELLEND, M. & STINCHCOMBE, J.R. (2009). Evolution in plant
populations as a driver of ecological changes in arthropod communities.
Philosophical Transactions of the Royal Society B: Biological Sciences **364**,
1593–1605.

KAGIYA, S., YASUGI, M., KUDOH, H., NAGANO, A.J. & UTSUMI, S. (2018). Does
genomic variation in a foundation species predict arthropod community structure
in a riparian forest? *Molecular Ecology* **27**, 1284–1295.

KANAGA, M.K., LATTA, L.C., MOCK, K.E., RYEL, R.J., LINDROTH, R.L. & PFRENDER,
M.E. (2009). Plant genotypic diversity and environmental stress interact to
negatively affect arthropod community diversity. *Arthropod-Plant Interactions*
3, 249–258.

KATANO, O. (2011). Effects of individual differences in foraging of pale chub on algal
biomass through trophic cascades. *Environmental Biology of Fishes* **92**, 101–
112.

KORICHEVA, J., GUREVITCH, J. & MENGERSEN, K. (2013). *Handbook of meta-Analysis*
in ecology and evolution. Princeton University press, Princeton, New Jersey.

KOTOWSKA, A.M., CAHILL JR, J.F. & KEDDIE, B.A. (2010). Plant genetic diversity
yields increased plant productivity and herbivore performance. *Journal of*
Ecology **98**, 237–245.

LAGERSTROM, A., NILSSON, M.-C. & WARDLE, D.A. (2013). Decoupled responses of
tree and shrub leaf and litter trait values to ecosystem retrogression across an
island area gradient. *Plant and Soil* **367**, 183–197.

LAMY, T., LAROCHE, F., DAVID, P., MASSOL, F. & JARNE, P. (2017). The contribution of
species-genetic diversity correlations to the understanding of community
assembly rules. *Oikos* **126**, 759–771.

LECERF, A. & CHAUVET, E. (2008). Intraspecific variability in leaf traits strongly affects
alder leaf decomposition in a stream. *Basic and Applied Ecology* **9**, 598–605.

LEROY, C.J., WHITHAM, T.G., WOOLEY, S.C. & MARKS, J.C. (2007). Within-species
variation in foliar chemistry influences leaf-litter decomposition in a Utah river.
Journal of the North American Benthological Society **26**, 426–438.

LOREAU, M. (2000). Biodiversity and ecosystem functioning: recent theoretical
916   advances. *Oikos* **91**, 3–17.

917 LOREAU, M., MOUQUET, N. & HOLT, R.D. (2003). Meta-ecosystems: a theoretical
918   framework for a spatial ecosystem ecology. *Ecology Letters* **6**, 673–679.

919 LOWE, W.H., KOVACH, R.P. & ALLENDORF, F.W. (2017). Population genetics and
920   demography unite ecology and evolution. *Trends in Ecology & Evolution* **32**,
921   141–152.

922 MADRITCH, M., DONALDSON, J.R. & LINDROTH, R.L. (2006). Genetic identity of
923   *Populus tremuloides* litter influences decomposition and nutrient release in a
924   mixed forest stand. *Ecosystems* **9**, 528–537.

925 MADRITCH, M.D., GREENE, S.L. & LINDROTH, R.L. (2009). Genetic mosaics of
926   ecosystem functioning across aspen-dominated landscapes. *Oecologia* **160**, 119–
927   127.

928 MADRITCH, M.D. & HUNTER, M.D. (2003). Intraspecific litter diversity and nitrogen
929   deposition affect nutrient dynamics and soil respiration. *Oecologia* **136**, 124–
930   128.

931 MADRITCH, M.D. & HUNTER, M.D. (2004). Phenotypic diversity and litter chemistry
932   affect nutrient dynamics during litter decomposition in a two species mix. *Oikos*
933   **105**, 125–131.

934 MADRITCH, M.D. & HUNTER, M.D. (2005). Phenotypic variation in oak litter influences
935   short- and long-term nutrient cycling through litter chemistry. *Soil Biology &*
936   *Biochemistry* **37**, 319–327.

937 MADRITCH, M.D. & LINDROTH, R.L. (2011). Soil microbial communities adapt to
938   genetic variation in leaf litter inputs. *Oikos* **120**, 1696–1704.

939 MANLY, B.F. (1997). *Randomization, bootstrap and Monte Carlo methods in biology.*
940   Chapman and Hall, London.

941 MATTHEWS, B., AEBISCHER, T., SULLAM, K.E., LUNDSGAARD-HANSEN, B. &
942   SEEHAUSEN, O. (2016). Experimental evidence of an eco-evolutionary feedback
943   during adaptive divergence. *Current Biology* **26**, 483–489.

944 MATTHEWS, B., DE MEESTER, L., JONES, C.G., IBELINGS, B.W., BOUMA, T.J.,
945   NUUTINEN, V., DE KOPPEL, J. VAN & ODLING-SMEE, J. (2014). Under niche
946   construction: an operational bridge between ecology, evolution, and ecosystem
947   science. *Ecological Monographs* **84**, 245–263.

948 MATTHEWS, B., NARWANI, A., HAUSCH, S., NONAKA, E., PETER, H., YAMAMICHI, M.,
949   SULLAM, K.E., BIRD, K.C., THOMAS, M.K., HANLEY, T.C. & TURNER, C.B.
950   (2011). Toward an integration of evolutionary biology and ecosystem science.
951   *Ecology Letters* **14**, 690–701.

MCART, S.H., COOK-PATTON, S.C. & THALER, J.S. (2012). Relationships between
arthropod richness, evenness, and diversity are altered by complementarity
among plant genotypes. *Oecologia* **168**, 1013–1021.

MIMURA, M., YAHARA, T., FAITH, D.P., VÁZQUEZ-DOMÍNGUEZ, E., COLAUTTI, R.I.,
ARAKI, H., JAVADI, F., NÚÑEZ-FARFÁN, J., MORI, A.S., ZHOU, S.,
HOLLINGSWORTH, P.M., NEAVES, L.E., FUKANO, Y., SMITH, G.F., SATO, Y.-I., ET
AL. (2016). Understanding and monitoring the consequences of human impacts
on intraspecific variation. *Evolutionary Applications* **10**, 121–139.

MORRISSEY, M.B. (2016*a*). Meta-analysis of magnitudes, differences and variation in
evolutionary parameters. *Journal of Evolutionary Biology* **29**, 1882–1904.

MORRISSEY, M.B. (2016*b*). Rejoinder: Further considerations for meta-analysis of
transformed quantities such as absolute values. *Journal of Evolutionary Biology*
29, 1922–1931.

NAEEM, S., THOMPSON, LINDSEY J, LAWLER, SHARON P, LAWTON, JOHN H & WOODFIN,
RICHARD M (1994). Declining biodiversity can alter the performance of
ecosystems. *Nature* **368**, 734–737.

NAKAGAWA, S. & CUTHILL, I.C. (2007). Effect size, confidence interval and statistical
significance: a practical guide for biologists. *Biological Reviews* **82**, 591–605.

NAKAGAWA, S. & SANTOS, E.S.A. (2012). Methodological issues and advances in
biological meta-analysis. *Evolutionary Ecology* **26**, 1253–1274.

NELL, C.S., MEZA-LOPEZ, M.M., CROY, J.R., NELSON, A.S., MOREIRA, X., PRATT, J.D.
& MOONEY, K.A. (2018). Relative effects of genetic variation sensu lato and
sexual dimorphism on plant traits and associated arthropod communities.
Oecologia **187**, 389–400.

NOBLE, D.W.A., LAGISZ, M., O'DEA, R.E. & NAKAGAWA, S. (2017). Nonindependence
and sensitivity analyses in ecological and evolutionary meta-analyses. *Molecular*
Ecology **26**, 2410–2425.

ODLING-SMEE, J., LALAND, K.N. & FELDMAN, M.W. (2003). *Niche construction the*
neglected process in evolution. Princeton University Press.

PAINE, R.T. (1969). A note on trophic complexity and community stability. *The*
American Naturalist **103**, 91–93.

PALKOVACS, E.P., FRYXELL, D.C., TURLEY, N.E. & POST, D.M. (2015). Ecological
effects of intraspecific consumer biodiversity for aquatic communities and
ecosystems. In *Aquatic Functional Biodiversity* pp. 37–51. Elsevier.

PALKOVACS, E.P., MARSHALL, M.C., LAMPHERE, B.A., LYNCH, B.R., WEESE, D.J.,
FRASER, D.F., REZNICK, D.N., PRINGLE, C.M. & KINNISON, M.T. (2009).
Experimental evaluation of evolution and coevolution as agents of ecosystem
change in Trinidadian streams. *Philosophical Transactions of the Royal Society*
B-Biological Sciences **364**, 1617–1628.

PALKOVACS, E.P. & POST, D.M. (2009). Experimental evidence that phenotypic
divergence in predators drives community divergence in prey. *Ecology* **90**, 300–
305.

PINHEIRO, J., BATES, D., DEBROY, S., SARKAR, D. & R CORE TEAM (2014). nlme: linear
and nonlinear mixed effects models. R package version 3.1-117.

POST, D.M., PALKOVACS, E.P., SCHIELKE, E.G. & DODSON, S.I. (2008). Intraspecific
variation in a predator affects community structure and cascading trophic
interactions. *Ecology* **89**, 2019–2032.

PRATT, J.D., DATU, A., TRAN, T., SHENG, D.C. & MOONEY, K.A. (2017). Genetically
based latitudinal clines in *Artemisia californica* drive parallel clines in arthropod
communities. *Ecology* **98**, 79–91.

PREGITZER, C.C., BAILEY, J.K. & SCHWEITZER, J.A. (2013). Genetic by environment
interactions affect plant-soil linkages. *Ecology and Evolution* **3**, 2322–2333.

REISS, J., BRIDLE, J.R., MONTOYA, J.M. & WOODWARD, G. (2009). Emerging horizons
in biodiversity and ecosystem functioning research. *Trends in Ecology &*
Evolution **24**, 505–514.

RENNEVILLE, C., ROUZIC, A.L., BAYLAC, M., MILLOT, A., LOISEL, S. & EDELINE, E.
(2016). Morphological drivers of trophic cascades. *Oikos* **125**, 1193–1202.

ROBINSON, K.M., INGVARSSON, P.K., JANSSON, S. & ALBRECTSEN, B.R. (2012). Genetic
variation in functional traits influences arthropod community composition in
aspen (*Populus tremula L.*). *PLoS ONE* **7**, e37679.

RODRIGUEZ-CABAL, M.A., BARRIOS-GARCIA, M.N., RUDMAN, S.M., MCKOWN, A.D.,
SATO, T. & CRUTSINGER, G.M. (2017). It is about time: genetic variation in the
timing of leaf-litter inputs influences aquatic ecosystems. *Freshwater Biology*
62, 356–365.

ROYAUTÉ, R. & PRUITT, J.N. (2015). Varying predator personalities generates
contrasting prey communities in an agroecosystem. *Ecology* **96**, 2902–2911.

RUDGERS, J.A. & WHITNEY, K.D. (2006). Interactions between insect herbivores and a
plant architectural dimorphism. *Journal of Ecology* **94**, 1249–1260.

RUDMAN, S.M., RODRIGUEZ-CABAL, M.A., STIER, A., SATO, T., HEAVYSIDE, J., EL-
SABAAWI, R.W. & CRUTSINGER, G.M. (2015). Adaptive genetic variation
mediates bottom-up and top-down control in an aquatic ecosystem. *Proceedings*
of the Royal Society B: Biological Sciences **282**, 125–132.

RUDMAN, S.M. & SCHLUTER, D. (2016). Ecological impacts of reverse speciation in
threespine stickleback. *Current Biology* **26**, 490–495.

RUDOLF, V.H.W. & RASMUSSEN, N.L. (2013*a*). Ontogenetic functional diversity: Size
structure of a keystone predator drives functioning of a complex ecosystem.
Ecology **94**, 1046–1056.

RUDOLF, V.H.W. & RASMUSSEN, N.L. (2013*b*). Population structure determines
functional differences among species and ecosystem processes. *Nature*
Communications **4**.

SCHMITZ, O.J., HAWLENA, D. & TRUSSELL, G.C. (2010). Predator control of ecosystem
nutrient dynamics. *Ecology Letters* **13**, 1199–1209.

SCHMITZ, O.J., RAYMOND, P.A., ESTES, J.A., KURZ, W.A., HOLTGRIEVE, G.W., RITCHIE,
M.E., SCHINDLER, D.E., SPIVAK, A.C., WILSON, R.W., BRADFORD, M.A.,
CHRISTENSEN, V., DEEGAN, L., SMETACEK, V., VANNI, M.J. & WILMERS, C.C.
(2014). Animating the carbon cycle. *Ecosystems* **17**, 344–359.

SCHWEITZER, J.A., BAILEY, J.K., FISCHER, D.G., LEROY, C.J., LONSDORF, E.V.,
WHITHAM, T.G. & HART, S.C. (2008). Plant–soil–microorganism interactions:
heritable relationship between plant genotype and associated soil
microorganisms. *Ecology* **89**, 773–781.

SEMCHENKO, M., SAAR, S. & LEPIK, A. (2017). Intraspecific genetic diversity modulates
plant-soil feedback and nutrient cycling. *New Phytologist* **216**, 90–98.

SENIOR, A.M., GRUEBER, C.E., KAMIYA, T., LAGISZ, M., O'DWYER, K., SANTOS, E.S.A.
& NAKAGAWA, S. (2016). Heterogeneity in ecological and evolutionary meta-
analyses: its magnitude and implications. *Ecology* **97**, 3293–3299.

SHIPLEY, B. (2000). *Cause and correlation in biology: a user's guide to path analysis,*
structural equation and causal inference. Cambridge University Press.

SIEFERT, A., VIOLLE, C., CHALMANDRIER, L., ALBERT, C.H., TAUDIERE, A., FAJARDO,
A., AARSSEN, L.W., BARALOTO, C., CARLUCCI, M.B., CIANCIARUSO, M.V., DE L.
DANTAS, V., DE BELLO, F., DUARTE, L.D.S., FONSECA, C.R., FRESCHET, G.T., ET
AL. (2015). A global meta-analysis of the relative extent of intraspecific trait
variation in plant communities. *Ecology Letters* **18**, 1406–1419.

SILFVER, T., KONTRO, M., PAASO, U., KARVINEN, H., KESKI-SAARI, S., KEINÄNEN, M.,
ROUSI, M. & MIKOLA, J. (2018). Intrapopulation genotypic variation in leaf litter
chemistry does not control microbial abundance and litter mass loss in silver
birch, *Betula pendula. Plant and Soil* **426**, 253–266.

SILFVER, T., MIKOLA, J., ROUSI, M., ROININEN, H. & OKSANEN, E. (2007). Leaf litter
decomposition differs among genotypes in a local *Betula pendula* population.
Oecologia **152**, 707–714.

SILFVER, T., PAASO, U., RASEHORN, M., ROUSI, M. & MIKOLA, J. (2015). Genotype x
herbivore effect on leaf litter decomposition in *Betula Pendula* saplings:
ecological and evolutionary consequences and the role of secondary metabolites.
PLoS One **10**.

SILFVER, T., ROUSI, M., OKSANEN, E. & ROININEN, H. (2014). Genetic and
environmental determinants of insect herbivore community structure in a *Betula*
pendula population. *F1000Research* **3**, 34–34.

SOUTHWOOD, T.R., BROWN, V.K. & READER, P.M. (1979). The relationships of plant
and insect diversities in succession. *Biological Journal of the Linnean Society*
12, 327–348.

STHULTZ, C.M., WHITHAM, T.G., KENNEDY, K., DECKERT, R. & GEHRING, C.A. (2009).
Genetically based susceptibility to herbivory influences the ectomycorrhizal
fungal communities of a foundation tree species. *New Phytologist* **184**, 657–667.

TACK, A.J.M., OVASKAINEN, O., PULKKINEN, P. & ROSLIN, T. (2010). Spatial location
dominates over host plant genotype in structuring an herbivore community.
Ecology **91**, 2660–2672.

TACK, A.J.M. & ROSLIN, T. (2011). The relative importance of host-plant genetic
diversity in structuring the associated herbivore community. *Ecology* **92**, 1594–
1604.

TOVAR-SANCHEZ, E., VALENCIA-CUEVAS, L., CASTILLO-MENDOZA, E., MUSSALI-
GALANTE, P., PEREZ-RUIZ, R.V. & MENDOZA, A. (2013). Association between
individual genetic diversity of two oak host species and canopy arthropod
community structure. *European Journal of Forest Research* **132**, 165–179.

TRAP, J., HÄTTENSCHWILER, S., GATTIN, I. & AUBERT, M. (2013). Forest ageing: An
unexpected driver of beech leaf litter quality variability in European forests with
strong consequences on soil processes. *Forest Ecology and Management* **302**,
338–345.

UTSUMI, S. (2015). Feeding evolution of a herbivore influences an arthropod
community through plants: implications for plant-mediated eco-evolutionary
feedback loop. *Journal of Ecology* **103**, 829–839.

VANNI, M.J. (2002). Nutrient cycling by animals in freshwater ecosystems. *Annual
Review of Ecology and Systematics* **33**, 341–370.

VELLEND, M. (2005). Species diversity and genetic diversity: parallel processes and
correlated patterns. *The American Naturalist* **166**, 199–215.

VELLEND, M. & GEBER, M.A. (2005). Connections between species diversity and
genetic diversity. *Ecology Letters* **8**, 767–781.

VIOLLE, C., ENQUIST, B.J., MCGILL, B.J., JIANG, L., ALBERT, C.H., HULSHOF, C., JUNG,
V. & MESSIER, J. (2012). The return of the variance: intraspecific variability in
community ecology. *Trends in Ecology & Evolution* **27**, 244–252.

VIOLLE, C., NAVAS, M.-L., VILE, D., KAZAKOU, E., FORTUNEL, C., HUMMEL, I. &
GARNIER, E. (2007). Let the concept of trait be functional! *Oikos* **116**, 882–892.

WAGG, C., BOLLER, B., SCHNEIDER, S., WIDMER, F. & VAN DER HEIJDEN, M.G.A.
(2015). Intraspecific and intergenerational differences in plant-soil feedbacks.
Oikos **124**, 994–1004.

WALSH, M.R., DELONG, J.P., HANLEY, T.C. & POST, D.M. (2012). A cascade of
evolutionary change alters consumer-resource dynamics and ecosystem

function. *Proceedings of the Royal Society B-Biological Sciences* **279**, 3184–
3192.

WANG, X.-Y., MIAO, Y., YU, S., CHEN, X.-Y. & SCHMID, B. (2014). Genotypic diversity
of an invasive plant species promotes litter decomposition and associated
processes. *Oecologia* **174**, 993–1005.

WEIS, J.J. & POST, D.M. (2013). Intraspecific variation in a predator drives cascading
variation in primary producer community composition. *Oikos* **122**, 1343–1349.

WERNER, E.E. & PEACOR, S.D. (2003). A review of trait-mediated indirect interactions
in ecological communities. *Ecology* **84**, 1083–1100.

WHITHAM, T.G., BAILEY, J.K., SCHWEITZER, J.A., SHUSTER, S.M., BANGERT, R.K.,
LEROY, C.J., LONSDORF, E.V., ALLAN, G.J., DIFAZIO, S.P., POTTS, B.M.,
FISCHER, D.G., GEHRING, C.A., LINDROTH, R.L., MARKS, J.C., HART, S.C., ET AL.
(2006). A framework for community and ecosystem genetics: from genes to
ecosystems. *Nature Reviews Genetics* **7**, 510–523.

WHITHAM, T.G., YOUNG, W.P., MARTINSEN, G.D., GEHRING, C.A., SCHWEITZER, J.A.,
SHUSTER, S.M., WIMP, G.M., FISCHER, D.G., BAILEY, J.K., LINDROTH, R.L.,
WOOLBRIGHT, S. & KUSKE, C.R. (2003). Community and ecosystem genetics: A
consequence of the extended phenotype. *Ecology* **84**, 559–573.

WILKINSON, A., ALEXANDER, I. & JOHNSON, D. (2012). Genotype identity determines
productivity and CO2 efflux across a genotype-species gradient of
ectomycorrhizal fungi. *Fungal Ecology* **5**, 571–580.

WILKINSON, A., SOLAN, M., TAYLOR, A.F.S., ALEXANDER, I.J. & JOHNSON, D. (2010).
Intraspecific diversity regulates fungal productivity and respiration. *Plos One* **5**,
e12604.

ZYTYNSKA, S.E., FAY, M.F., PENNEY, D. & PREZIOSI, R.F. (2011). Genetic variation in a
tropical tree species influences the associated epiphytic plant and invertebrate
communities in a complex forest ecosystem. *Philosophical Transactions of the
Royal Society B: Biological Sciences* **366**, 1329–1336.

ZYTYNSKA, S.E., KHUDR, M.S., HARRIS, E. & PREZIOSI, R.F. (2012). Genetic effects of
tank-forming bromeliads on the associated invertebrate community in a tropical
forest ecosystem. *Oecologia* **170**, 467–475.

1138

1139   **VIII. SUPPORTING INFORMATION**

1140

Additional supporting information may be found in the online version of this article.

Appendix S1. Papers used in the meta-analysis.

Fig. S1. Flow diagram of manuscript screening and eligibility for this meta-analysis.

Table S1. Formulae used to convert different statistical values into an *r* correlation

coefficient.

Fig. S2. Funnel plots describing the residuals effect size distribution against an estimate

of the study size for (A) intraspecific variation and (B) intraspecific richness.

Appendix S2. R script of the principal functions used in analyses.

Fig, S3. Distribution of effect sizes (ln*RR*) across ecological metrics.

Fig. S4. Distribution of resampled mean effect size (MES) expected under the null

hypothesis, i.e. the effect of intraspecific identity in all studies is zero.

Fig. S5. Distribution of median effect sizes calculated for each of the 1000 sets of 15

studies randomly selected within our extended data set of 75 studies.

Figure legends

Fig. 1. Publication year of the 90 selected articles used in the meta-analysis.

1156

Fig. 2. (A) Relationship between intraspecific richness and effect size (lnRR) on

community (points) and ecosystem (crosses) dynamics. The line represents the shape of

the relationship as predicted using an exponential asymptotic non-linear mixed effect

model. The blue shadow represents 95% CI. Symbol colours denote the target species:

herbaceous plant (black), tree (blue) or fungus (grey). (B) Density of absolute effect size

(lnRR) for intraspecific and interspecific richness on ecological dynamics. Posterior

means and 95% percentile intervals (points and horizontal lines, respectively) were

estimated using a model including article ID as the random effect and the inverse of

sample size as a weighting parameter.

1166

1167 **Fig. 3.** Density probability of effect size Zr. The vertical broken line represents the

1168 resampled Zr mean under the null hypothesis (confidence intervals not shown because

1169 they are too narrow); the black curve shows the distribution of observed Zr and its mean

1170 (vertical black straight line) and 95% CIs (grey shading).

1171

Fig. 4. (A) Mean effect size Zr for different species groups. The sample sizes (N)

represent the number of species and the number of effect sizes, respectively. The

horizontal broken line represents the mean effect size; error bars represent ± 1 SE. (B)

Mean Zr for the ecological response variables. The sample sizes (N) of the number of

effect sizes are given. The horizontal broken line represents the mean effect size; error

bars represent ± 1 SE.

1178

1179   Fig. 1

1180

1181

1182   Fig. 2

1183

1184

1185   Fig. 3

1186

1187

1188   Fig. 4

1189

1190

Table 1. Summary table of model selection by Akaike information criterion (AIC)
comparison to explain the shape of the relationship between the ecological
consequences and the intraspecific richness. Models were run as non-linear mixed-
effect models with the article ID as a random factor; equations and parameters estimates
are also shown. IR, intraspecific richness; ln*RR*, effect size of intraspecific richness on
ecological dynamics.
1197

Model	AIC	ΔAIC	AIC weight	Equation	Parameter estimates
Asymptotic exponential model	−176.695	0	0.999	$lnRR = a * (1 - e^{-b*IR})$	$a = 0.221$ $b = 0.617$
Michaelis–Menten model	−11.873	164.821	< 0.001	$lnRR = V * \dfrac{IR}{k + IR}$	$V = 0.054$ $k = -2.712$
Linear model	−160.974	15.721	< 0.001	$lnRR = b0 + b1 * IR$	$b1 = 0.012$ $b0 = 0.122$
Null model	−144.191	32.503	< 0.001	$lnRR = b0$	$b0 = 0.183$

1198