The community and ecosystem consequences of intraspecific

diversity: a meta-analysis

Allan. Raffard[1, 2*], Frédéric. Santoul[2,], Julien. Cucherousset[3] and Simon.

Blanchet[1, 3*]

[1] CNRS, Université Toulouse III Paul Sabatier, Station d'Écologie Théorique et

Expérimentale du CNRS à Moulis, UMR-5321, F-09200 Moulis, France.

[2] UPS, CNRS, INP, EcoLab (Laboratoire d'écologie fonctionnelle et environnement),

Université de Toulouse, 118 route de Narbonne, UMR-5245, FR-31062 Toulouse

Cedex 9, France;

[3] Laboratoire Évolution and Diversité Biologique (EDB UMR 5174), Université de

Toulouse, CNRS, IRD, UPS, 118 route de Narbonne, Toulouse 31062, France.

Corresponding authors:

Allan RAFFARD and Simon BLANCHET
Station d'Ecologie Théorique et Expérimentale du CNRS à Moulis.
UMR-5321(CNRS-UPS)
2 route du CNRS, F-09200 Moulis, France
Phone: (+33) 5.61.55.67.47
E-mail: allan.raffard@sete.cnrs.fr; simon.blanchet@sete.cnrs.fr
24

25 **Co-author's e-mails:**

26 Frédéric SANTOUL: frederic.santoul@univ-tlse3.fr

27 Julien CUCHEROUSSET: julien.cucherousset@univ-tlse3.fr

Brief statement of Authorship: *A.R and S.B designed the study. A.R performed*

statistical analyses. A.R, S.B, F.S and J.C interpreted and discussed the results. A.R and

S.B wrote the first draft of the paper, A.R, S.B, F.S and J.C corrected and improved the

paper. All authors have read and approved this version of the manuscript.

Keywords: ecosystem functioning, community, genotype, phenotype, meta-analysis,

biodiversity, eco-evolutionary dynamics

Abstract

Understanding the relationships between biodiversity and ecosystem functioning has major implications. Biodiversity-function relationships are generally investigated at the interspecific level, although intraspecific diversity is increasingly perceived as an important ecological facet of biodiversity. Here, we provide a quantitative synthesis testing –across diverse plant and animal species- whether intraspecific diversity (measured either as the *richness* or *variation* of genotypes/phenotypes in a population) is a major driver of community dynamics and ecosystem functioning. We found a significant relationship between intraspecific *richness* and community and ecosystem dynamics that followed a saturating shape, as observed for biodiversity-function relationships measured at the interspecific level. The strength of the ecological effects of intraspecific *richness* were similar to those reported for interspecific richness. We further found that intraspecific *variation* has substantial effects on ecological dynamics that are twice higher than expected by chance, and that these effects might have been previously underestimated. We finally found that the ecological effects of intraspecific *variation* were not homogeneous and were actually stronger in primary producer than in consumer species, and when measured at the ecosystem than at the community level. Overall, we demonstrated that intraspecific diversity can have strong consequences on community and ecosystem dynamics, which unfolds the pivotal role of this biodiversity facet for understanding ecological dynamics.

Table of Contents

I. Introduction

Understanding the relationships between biodiversity and ecosystem functioning is a field of intense investigations informing on the services provided by biodiversity (Chapin *et al.*, 2000; Loreau, 2000; Hooper *et al.*, 2005; Cardinale *et al.*, 2012). Biodiversity is generally quantified as the taxonomic, functional and/or phylogenetic diversity of a species assemblage, and most studies on biodiversity-ecosystem functioning relationships have to date focused on the *interspecific* facet of biodiversity (Naeem *et al.*, 1994; Downing & Leibold, 2002; Hillebrand & Matthiessen, 2009). However, biodiversity also includes an *intraspecific* facet that is defined as the phenotypic, functional and genetic diversity measured within a single species (Odling-Smee, Laland, & Feldman, 2003; Bolnick *et al.*, 2003). During the last two decades, intraspecific diversity has been demonstrated to account for a non-negligible part of the total biodiversity measured in plants and animals (Fridley & Grime, 2010; de Bello *et al.*, 2011), representing in some cases up to a quarter of the total variability measured in communities (Fridley & Grime, 2010; de Bello *et al.*, 2011; Siefert *et al.*, 2015). Furthermore, there is now mounting evidence that intraspecific diversity can strongly affect ecological dynamics at levels higher than the population, i.e. at the community and ecosystem levels (Bolnick *et al.*, 2003, 2011; Hughes *et al.*, 2008; Bailey *et al.*, 2009; Violle *et al.*, 2012).

Investigations on the relationships between intraspecific diversity and community and ecosystem dynamics (hereafter "ecological dynamics") have attracted many scientists, since this facet of biodiversity is at the thin frontier between evolution and ecology (Schoener, 2011; Hendry, 2016). Consequently, many conceptual insights on the relationships between intraspecific diversity and ecological dynamics arose from studies on eco-evolutionary dynamics. These dynamics aimed at understanding how

evolutionary processes can affect ecological dynamics, and how, in turn, these

ecological changes can lead to rapid evolutionary responses (Thompson, 1998; Odling-

Smee *et al.*, 2003; Whitham *et al.*, 2003; Schoener, 2011; Matthews *et al.*, 2011;

Hendry, 2016). These key concepts have been substantiated by major experimental

studies demonstrating how stress-induced phenotypic changes (e.g. through predation)

(Werner & Peacor, 2003; Schmitz *et al.*, 2008; Hawlena *et al.*, 2012), individual

genotypes (Whitham *et al.*, 2003; Madritch, Greene, & Lindroth, 2009), and rapid

adaptation (Harmon *et al.*, 2009; Matthews *et al.*, 2016; Rudman & Schluter, 2016) can

alter community and ecosystem dynamics.

Intraspecific diversity can be quantified based on the *richness* of populations,

which corresponds to the differences in the number of genotypes/phenotypes composing

populations. For instance, populations are often characterized according to their "allelic

richness", which is a population parallel of species richness, a common metric measured

at the community level and used to investigate biodiversity-functions relationships

(BEF, Crutsinger, 2006). Intraspecific richness can also affect ecological dynamics

(hence generating "intraspecific BEF", Whitham *et al.*, 2006; Crutsinger, 2006), and the

ecological consequences of intraspecific richness should follow a saturating curve as

often described for BEF observed at the interspecific level (Hughes *et al.*, 2008).

Although rarely tested empirically, the saturating shape could be due to the combined

effects of several mechanisms. Populations with different richness could have different

ecological   consequences   because   of   ecological   complementarity   among

genotypes/phenotypes (i.e. niche partitioning, facilitation occurring when genotypes use

different resources), inhibition among genotypes/phenotypes (when multiple genotypes

are   in   competition   for   resources),   or   functional   redundancy   among

genotypes/phenotypes that can make populations ecologically equivalent (Johnson,

Lajeunesse, & Agrawal, 2006; Hughes *et al.*, 2008). Yet, the shape of the relationship

between intraspecific richness and ecological dynamics has rarely been investigated

empirically and –up to our knowledge- never been quantified across species.

The ecological consequences of intraspecific diversity can also been investigated

through the lens of *variation* in genotypic or phenotypic attributes. Adaptive and non-

adaptive evolutionary processes such as natural selection, plasticity or genetic drift can

generate unique phenotypic differences among populations. These differences can be

associated to key functional processes such as food acquisition or nutrient cycling (e.g.

Grant & Grant, 2006; Rudgers & Whitney, 2006; Lowe, Kovach, & Allendorf, 2017),

resulting in both trophic and non-trophic effects of intraspecific variation on ecological

dynamics (Odling-Smee *et al.*, 2003; Whitham *et al.*, 2003; Matthews *et al.*, 2011). For

instance, plant genotypes differing in their susceptibility to herbivores have diverging

influences on herbivore communities (Barbour *et al.*, 2009). Similarly, differences in

diet within predator populations can modify prey community structure (Post *et al.*,

2008; Howeth *et al.*, 2013). Non-trophic interactions can also have an important role.

For instance, differences in the chemical composition of individuals can result in

differences in excretion rate or in leaf chemistry that can then affect ecosystem

functions such as primary production or nutrient recycling (Lecerf & Chauvet, 2008; El-

Sabaawi *et al.*, 2015). Recently Des Roches *et al.* (2018) demonstrated that intraspecific

variation can affect ecological dynamics as much as the removal or replacement of a

species in the environment. Although based on a restricted number of pioneer studies

(25 studies and 15 species), this study confirms the hypothesis that intraspecific

variation might be a non-negligible driver of ecological dynamics.

Here, we tested –across various species and ecosystems- to which extent

intraspecific *richness* and intraspecific *variation* affect the structure of communities and

the functioning of ecosystems, and hence whether intraspecific diversity is a major

driver of ecological dynamics. We ran two meta-analyses synthesizing published data

across taxa and ecosystems for intraspecific richness and variation respectively to fulfil

three specific objectives. First, we tested for the significance and the shape of the

relationship between intraspecific richness and ecological dynamics. We expect to find

a significant saturating relationship between intraspecific richness and ecological

dynamics, mainly because of potential facilitation and functional redundancy among

genotypes and phenotypes (Hughes *et al.*, 2008). Second, we compared the ecological

effects of intraspecific richness to those of interspecific richness obtained from

experimental studies manipulating species richness (Duffy, Godwin, & Cardinale,

2017). Finally, we provided a novel and extensive quantitative synthesis testing for the

effects of intraspecific variation on ecological dynamics. Des Roches *et al.* (2018)

previously focused on studies removing or replacing the target species (this for which

intraspecific variation is manipulated) to investigate the ecological consequences of

intraspecific variation. This strongly restricted the number of available studies on which

effects sizes were calculated, and potentially upwardly biased these estimates (Des

Roches *et al.*, 2018). We here relaxed this restriction by considering all studies

manipulating intraspecific variation, and we used a null model approach to provide a

more accurate effect size of intraspecific variation on ecological dynamics. We also

built on this extended dataset to thoroughly decompose variance in the ecological

consequences of intraspecific variation according to the type of organism that was

manipulated and the type of response variable being measured. Amongst other, we

tested whether the magnitude of the effects of intraspecific variation on ecological

dynamics vary among organism types (primary producers *vs*. consumers) and levels of

biological organisation (community *vs*. ecosystem levels). Because primary producers

form the basis of trophic chains, we expect stronger ecological effects of intraspecific

variation in producers than in consumers. We also expect stronger effects of

intraspecific variation on ecosystem functions than on metrics describing community

structure because ecosystem functions are affected by both trophic and non-trophic

effects of biodiversity (Matthews *et al.*, 2014).

II.   Material & Methods
(1)   Data collection

We compiled data from published articles quantifying the effects of intraspecific

diversity in a single species on community structure and/or ecosystem functioning. We

focused only on intraspecific diversity that was naturally observed in populations and

that represented the integrative phenotypic effect of multiple evolutionary processes

including selection, drift and/or plasticity. As a result, we did not consider articles

focusing on experimentally-induced intraspecific diversity through induced plastic

responses to particular predatory or environmental cues for instance (see Werner &

Peacor, 2003 for a review on this topic). Studies varying intraspecific diversity within a

set of multiple species (e.g. Booth & Grime, 2003) were not included in this meta-

analysis. The literature search was done using the ISI Web of Knowledge® platform

(last accessed January 2016). We also scrutinized the reference list of each article to

obtain additional articles. When conducted the literature search, the following keywords

were used under various combinations: "community genetics", "intraspecific variation",

"ecosystem functions", "eco-evolutionary dynamics", and "intraspecific genetic

variation" (see Appendix S1 for details). We selected articles describing the effects of

genotypic and/or phenotypic richness (*intraspecific richness*) and/or different genotypes

phenotypes (*intraspecific variation*) in a single species (i.e. the target species) on

community and/or ecosystem dynamics. A total of 77 articles with available statistics

were selected (see Appendix S2). Among them, 18 focused on intraspecific richness and

65 focused on intraspecific variation.

For each study, we recorded the Latin name of the target species and we

classified them as primary producers or consumers (including here primary and

secondary consumers as well as predators) and according to the major taxonomic

categories that were represented in our dataset, namely arthropods (7 species), fishes (6

species), herbaceous (8 species) and trees (26 species). We also recorded the duration

(in days, mean duration of survey ± SD: 662 ± 1310, range: from 1 to 6205 days) of the

study as a co-variable because we can expect that the ecological divergence among

treatments could increase with time. We finally recorded seven main response variables

related to community structure and ecosystem functioning. A community was here

defined as a group of at least two species, and we focused on three types of response

variables describing the structure of communities:

- *Species abundance*: total number of individuals of all species;

- *Biomass*: total mass of individuals of all species;

- *Community structure*: number of species such as Simpson or Shannon indices,

species evenness and/or species richness.

Regarding response variables at the ecosystem level, we considered four main

ecosystem functions:

- *Decomposition rate*: rate at which organic matter is recycled;

- *Elemental cycling*: quantity of organic or inorganic materials;

- *Primary productivity*: measured as several proxies of primary producers: biomass

of primary producers excluding the productivity of the target species, chlorophyll *a*

concentration, daily oxygen production, and photosynthetically active radiation;

- *Ecosystem respiration*: rate of oxygen consumption.

(2)   Meta-analysis
(a) The ecological consequences of intraspecific richness

To test for the consequences of intraspecific richness on ecological dynamics, we

focused only on studies investigating the consequences of *genotypic* richness since it

was the intraspecific diversity facet most commonly manipulated to test for the effects

of intraspecific richness on ecological dynamics. Since measures of variance associated

to the effects of intraspecific richness on ecological dynamics were not available for all

response variables, we used the log-transformed response ratio (lnRR) as an effect size.

lnRR was computed using the following formula: $\ln\left(\frac{XG}{1G}\right)$, in which $1G$ is the average of

the response variable measured for the treatment with a single genotype (i.e.

monoculture), and XG is the average of the response variable measured for each

treatment independently including more than one genotype. For each response variable,

lnRR increases as the difference in the mean response variable measured in the

treatment with a single genotype and treatments including more than one genotype

increases. We also recorded the difference in the number of genotypes between the

single genotype treatment (monoculture) and all other treatments separately (i.e.

treatment including more than one genotype, from two to twelve genotypes) that was

termed "difference in intraspecific richness". In our dataset, difference in intraspecific

richness varied between one and eleven genotypes. This approach allowed quantifying

the ecological consequences of increasing the number of genotypes for each target

species. Since each study generally assessed the effects of intraspecific richness on

more than one response variable, our dataset included a total of 101 assays.

We aimed at testing the shape and the significance of the relationship between

lnRR and the difference in intraspecific richness across all case studies. The general

expectation is that ecological differences between treatments increase as the differences

in intraspecific richness increase, although this increase could be non-linear (Hughes *et*

al., 2008). We therefore used non-linear mixed effect models to test the significance and

shape of the relationship between absolute values of ln*RR* and differences in

intraspecific richness. More precisely, we modelled this relationship according to four

different models to determine the most likely shape of the relationship between ln*RR*

and difference in intraspecific richness:

- A null model (1 parameter) was computed for the null effect hypothesis (i.e. no

significant relationship between ln*RR* and difference in intraspecific richness);

- A linear model (2 parameters) suggesting a positive and linear relationship between

ln*RR* and difference in intraspecific richness;

- A Michaelis-Menten model (2 parameters) suggesting that ln*RR* increases with the

differences in intraspecific richness, until a plateau is reached (saturating shape);

- An asymptotic exponential model (2 parameters) suggesting a shape similar to the

Michaelis-Menten model, except that the plateau is reached sooner.

All models cited above included article ID as a random term, and the reverse of the

sample size as a weighting parameter so as to give more weight to articles including

more replicates. Models were compared according to AIC and we retained (as "best

models") all models that fell within a ΔAIC < 4 (Burnham & Anderson, 2002). We also

calculated for each model the Akaike weight that provides a conditional probability for

each model to be best supported by the data (Burnham & Anderson, 2002).

We then compared the ecological effects of intraspecific and interspecific

richness by extracting from each study and for each response variable the |ln*RR*|

corresponding to the most extreme levels of genotypic richness (x_{max}) manipulated in

each study (N = 48). These values were subsequently compared to published ln*RR*

calculated following a similar method for experiments manipulating interspecific

richness (Duffy *et al.*, 2017) (N = 35) using a Kruskal-Wallis rank sum test.

Finally, to compare the ecological consequences of intraspecific richness

between levels of biological organisation, we performed meta-regression as a linear

mixed effect model. The absolute value of ln*RR* was used as the dependent variable, the

articles ID was used as a random factor, whereas the level of biological organisation

(community *vs.* ecosystem response variables) was fixed effects. The type of organisms

was not included in the analysis given that studies on intraspecific richness focused

almost exclusively on primary producers (excepted one on fungus).

(b) The ecological consequences of intraspecific variation

Given that most studies (81 %) did not include a control (i.e. a treatment without the

target species), we compared the strength of effects among all unique genotypes and/or

phenotypes that were considered in each study. Studies generally compared the

consequences of 2 to 10 different unique genotypes and/or phenotypes (i.e. 2 to 10

treatments, with each treatment corresponding to a unique genotype/phenotype) on

community and/or ecosystem dynamics; we gathered from each study the statistic (*t*, *F*,

Chi-squared, Pearson's *r*, Spearman's *r*, R^2 or Hedge's *g*) associated with the between-

treatments comparison (i.e. the variation of the phenotypes/genotypes). The higher the

absolute value of the statistic, the higher the community and ecosystem consequences

due to the variation of genotypes and/or phenotypes. The absolute value of each statistic

was converted into a correlation coefficient (*r*) ranging from 0 to 1 (see Table S1 for the

formulas used to convert each statistic into a *r*). We did not use the direction of the

statistic (i.e. positive or negative) because it depended upon the ecological response

variable that was considered, which makes inconvenient comparisons on the direction

of effects. Then, the *Z*-Fisher transformation was used to obtain a standardized effect

size using the following formulae: $Zr = 0.5\ln\frac{(1+r)}{(1-r)}$. For each Zr value, we calculated

the corresponding standard error using the following formulae: $se_{Zr} = \frac{1}{\sqrt{n-3}}$ (Nakagawa

& Cuthill, 2007). Since each study generally focused on more than one response

variable, we obtained a total of 430 observed Zr values, each corresponding to the effect

size of intraspecific variation observed within one species on a single response variable.

The mean global Zr (MES$_{obs}$ for mean effect size observed) and its 95% confidence

interval (CI) were calculated using an intercept-only model. This intercept-only model

was run as a mixed model with no fixed effect, the article ID as random effect and se_{Zr}

was included as a weighting parameter to give more weight to studies with larger

sample size (Koricheva, Gurevitch, & Mengersen, 2013).

Because Zr ranged between 0 and $+\infty$, the CIs of the MES$_{obs}$ do not theoretically

overlap 0, which makes difficult the assessment of the significance of the strength of the

MES$_{obs}$. We therefore used a null model approach to test if MES$_{obs}$ was significantly

different from what was expected under the null hypothesis, i.e. the true effect of

intraspecific variation in all studies was zero. We resampled each statistic (e.g. t, F)

from each empirical study in their respective null distribution (Student, Fisher…) with

the adequate degree of freedom. This resampled set of statistics (N = 430) was

transformed into Zr as explained above to build a set of resampled Zr values. We used

this set of resampled Zr values to fit an intercept-only model with no fixed effect, the

corresponding article ID as a random term and se_{Zr} included as a weighting parameter

(as for MES$_{obs}$). The mean global resampled Zr (MES$_{res}$) was extracted from the model,

and we repeated this resampling procedure 1000 times to obtain 1000 values of MES$_{res}$.

This resampled distribution of 1000 MES$_{res}$ corresponds to the range of possible MES

values expected if the null hypothesis was true. Finally, we calculated the probability of

MES_{obs} to be larger than expected under this null hypothesis using a one-tailed test

(Manly, 1997).

We then compared the median of effect sizes (MES_{common}) of studies that were

common between our extended dataset and the dataset used by Des Roches *and*

colleagues (Des Roches *et al.*, 2018) (i.e. 15 studies) to a selection of 15 studies

randomly sampled from our extended dataset. We calculated the median effect size for

the subset of random studies (MES_{ran}) and repeated this resampling procedure 1000

times to obtain 1000 values of MES_{ran}. We then compared MES_{common} to each MES_{ran}

value to calculate the probability that MES_{common} was higher than a random subset of 15

studies taken from the whole dataset (Manly, 1997).

We finally tested whether effect sizes (Zr) differed among organism types, for

which intraspecific variation was manipulated, and among the ecological response

variables considered. We hence computed meta-regressions based on linear mixed

effect models with Zr values (for all 65 articles and 430 measures) as the dependent

variable, and the organism type or the ecological response variable as fixed effects. The

article ID was included as a random effect, and se_{Zr} was included as a weighting

parameter. Four models were run to assess the differences of effect sizes (i) between

organism types classified as consumers or primary producers, and (ii) between detailed

taxonomic categories (arthropods, fishes, herbaceous and trees). We then tested whether

the effect sizes of intraspecific variation differed among ecological response variables

(iii) classified as community or ecosystem variables, and (iv) classified according to

more detailed categories (abundance, biomass, community structure, decomposition,

nutrient cycling, primary productivity and respiration of the ecosystem).

(3)   Publications biases

For both intraspecific variation and intraspecific richness, we assessed potential

publication biases by combining Egger's regressions and funnel plots (Egger *et al.*,

1997). Egger's regressions and funnel plots were computed using the residuals of meta-

regressions related effect sizes to the main modifiers (i.e. the explicative variables) and

a measure of the study size (the inverse of se_{zr} and the sample size for intraspecific

variation and intraspecific richness, respectively, (Horvathova, Nakagawa, & Uller,

2012; Nakagawa & Santos, 2012). Typically, for intraspecific variation we ran an

Egger's regression model including the residuals of the meta-regression linking

intraspecific variation to the modifiers as a response variable and the inverse of se_{zr} as

the explicative variable. A similar approach was used for intraspecific richness. The

intercept α and the slope β of the Egger's regressions are expected not to be

significantly different from zero if the data sets are unbiased towards significant results.

Finally, funnel plots were produced as a scatterplot linking the residuals described

above to the respective measure of the study size. An unbiased data set is expected to

generate a funnel plot in which articles with larger sample sizes will be close to the

mean effect size, whereas articles with small sample sizes will show more variance

around the mean effect size (Horvathova *et al.*, 2012; Nakagawa & Santos, 2012).

Overall -and after accounting for important modifiers- we found that there was

no strong visual signs of publication biases, neither for intraspecific variation nor for

intraspecific richness (Fig S1). This visual inspection of funnel plots was confirmed by

the Egger's regressions since parameter values were not significant, neither for

intraspecific variation ($\alpha = 0.014$, *p* $= 0.485$; $\beta = -0.0014$, *p* $= 0.562$) nor for

intraspecific richness ($\alpha = -0.007$, *p* $= 0.654$; $\beta = 0.0001$, *p* $= 0.736$).

All statistical analyses were performed using the R environment (R core team,

2013). The nlme package (Pinheiro *et al.*, 2014) was used to compute linear and non-

linear mixed effect models.

III.   Results

All articles (n = 77) selected for investigating the effects of intraspecific richness and

intraspecific variation were published between 2000 and 2016, and 70 % used primary

producers as target species, which were also the first model organisms used (Fig. 1).

The first studies focusing on consumers were published only recently (in 2008), using

fish (70 %) and arthropods (30 %) as model species.

(1)   The intraspecific richness-ecological dynamics relationship

As expected, we found a significant, positive and non-linear relationship between

intraspecific richness and ecological dynamics that likely followed a saturating shape

(Fig. 2A). Indeed, the AIC selection procedure revealed that two out of the four tested

models were likely to be supported by the data (ΔAICs < 4, Table 1). Interestingly these

two models were both non-linear models varying slightly in the specification of the non-

linear term, and both the linear and null models were strongly rejected (ΔAICs > 12,

Table 1). Among these two supported models, the model that best supported the data

was the exponential asymptotic model (86% of chances of being the best fitting model

according to the Akaike weight and ΔAIC = 3.694 for the other model, Table 1),

suggesting that the relationship between intraspecific richness and changes in

community structure and ecosystem functioning (effect size: ln*RR*) likely followed a

saturating shape (Fig. 2A).

We further found that the ecological effects of intraspecific richness represented

66% of the ecological effects induced by interspecific richness (Fig. 2B). Although

effect sizes measured for interspecific and intraspecific richness respectively were

significantly different ($\chi^2 = 5.704$, d.f = 1, $p = 0.016$), they largely overlapped (Fig. 2B).

Interestingly, this difference was mainly driven by few studies on interspecific richness

with strong ecological effects (Fig. 2B). The ecological effects of intraspecific richness

did not significantly vary between the community ($\ln RR = 0.18$, SE = 0.02) and

ecosystem ($\ln RR = 0.12$, SE = 0.02) response variables ($F = 0.504$, d.f = 1, 16, $p = $

0.488).

(2)   The ecological consequences of intraspecific variation

We extended the meta-analysis performed by Des Roches *and colleagues* to 44 species

(*vs*. 15 species in Des Roches *et al.*, 2018). We found that that the observed effect size

of intraspecific variation on community structure and ecosystem dynamics was

significant, and actually twice higher than the resampled effect size expected under a

null expectation ($MES_{obs} = 0.52$, 95% confident interval (CI) = 0.44-0.59, and $MES_{null} = $

0.26, CI = 0.26-0.27, resampled test, $p < 0.001$, Fig. 3, see also Appendix S3 and Fig

S2).

We tested to which extent the restricted dataset analysed by Des Roches *et al.*

(2018) was representative of our extended dataset, or on the contrary biased upward as

expected by Des Roches *et al.* (2018). We found that that effect sizes measured from

Des Roches et al's dataset ($MES_{common} = 0.29$, 95% percentile interval (PI) = 0.03-1.05)

was not significantly different from the distribution of effect sizes measured in our

extended dataset ($MES_{ran} = 0.41$, PI = 0.02-1.43, resampling test, $p = 0.091$, Fig S3),

and actually tended to be downwardly biased.

Finally, the ecological effects induced by intraspecific variation were stronger

when primary producers rather than consumers were manipulated ($F = 4.082$ d.f. = 1,

364, $p = 0.044$, Fig. 4A). Nonetheless, the strongest ecological effects of intraspecific

variation were observed in arthropods and herbaceous species, whereas the smallest

effects were observed in fish and tree species ($F = 2.445$ d.f. = 3, 362, $p = 0.063$, Fig.

4A). Irrespectively of organism type, the effects of intraspecific variation were

significantly stronger when the response variables were measured at the ecosystem level

rather than at the community level ($F = 5.226$, d.f = 1, 364, $p = 0.023$, Fig. 4B). The

strongest effects were detected when response variables concerned nutrient cycling and

the assembly of community, whereas the lowest effects were found for general

measures of abundance and density of species ($F = 3.068$, d.f = 6, 358, $p = 0.006$, Fig.

4B).

IV.   Discussion

Intraspecific diversity is increasingly recognized as an important facet of biodiversity

that can affect all biological levels (Bailey *et al.*, 2009). We here provide quantitative

estimates of the consequences of intraspecific richness and variation on community

structure and ecosystem functioning. We demonstrated for the first time that the

intraspecific BEF followed –as theoretically expected- a non-linear saturating shape.

We confirmed that genotypic and/or phenotypic variation observed between populations

can have non-negligible effects on community structure and ecosystem functions, and

we demonstrated that previous estimates (Des Roches *et al.*, 2018) of these ecological

effects of intraspecific variation actually tended to be underestimated. Finally, our

exhaustive quantitative survey pointed out that the ecological consequences of

intraspecific variation differed among biological level of organisation, and among

organism types. These findings provide novel and integrative insights into the

ecological role of intraspecific diversity.

(1) Intraspecific diversity and the dynamics of communities and ecosystems

Although the form of the relationship between intraspecific richness and

ecological consequences has already been discussed conceptually (Hughes *et al.*, 2008),

our meta-analysis provides for the first time a qualitative and quantitative assessment of

intraspecific BEF. Specifically -and although considering almost only primary

producers-, our results demonstrated that an increase in intraspecific richness resulted in

a non-linear (saturating) increase in the magnitude of its effects on ecological dynamics.

Our finding supports the idea that ecological divergences between an environment

hosting populations composed of a single genotype and an environment hosting

populations composed of multiple genotypes increase until a plateau is reached as the

number of genotypes composing the later increases. This result echoes BEF defined at

the interspecific level (Reiss *et al.*, 2009; Cadotte, Carscadden, & Mirotchnick, 2011)

and suggests that the saturating shape might arise from similar mechanisms occurring at

the intraspecific and interspecific levels (Johnson *et al.*, 2006; Hughes *et al.*, 2008).

More specifically, the initial linear increase is assumed to be due to complementarity

and facilitation among genotypes, whereas the plateau is likely due to functional

redundancy among genotypes (Johnson *et al.*, 2006). Redundant genotypes probably

display functionally similar traits since two genotypes do not necessarily generate two

functionally different traits (e.g. through synonymous mutations or trait convergences).

Thus manipulating traits richness -rather than genotypic richness only-, and more

precisely functional effect traits (i.e. traits having ecological effects, Violle *et al.*, 2007),

in future experiments should confirm mechanisms underlying intraspecific biodiversity-

ecological dynamics relationship.

We found that effect sizes found for intraspecific richness were within the range

of the values recently reported for experimental interspecific BEF, indicating that the

ecological effects of varying phenotypic/genotypic richness within populations are in

some cases similar to those induced by varying species richness within communities.

This later finding posits several questions regarding the general relationships between

intraspecific diversity, community structure, ecosystem functioning and common

abiotic constraints. A large body of literature has demonstrated that intraspecific genetic

diversity and species diversity (a measure of community structure) might co-vary over

space because of common environmental drivers and/or reciprocal causal relationships

between intraspecific genetic diversity and species diversity (i.e. the species-genetic

diversity correlation framework, SGDC, Vellend & Geber, 2005; Vellend, 2005).

Because most studies considered in our meta-analysis are experimental, our findings

confirm that intraspecific diversity can directly influence the structure of communities -

irrespective of the abiotic environments-, hence adding weight to the SGDC framework.

Additionally, we suggest expanding the SGDC framework since intraspecific diversity

can also affect ecosystem functioning. This suggests that intraspecific diversity,

community structure and ecosystem functioning may actually be tightly linked into a

tripartite relationship. A major future challenge will be to tease apart the causal

relationships linking these three components within a common abiotic environment.

These relationships might indeed be direct (e.g. intraspecific diversity directly affects

community structure), indirect (e.g. intraspecific diversity indirectly affects ecosystem

functions through its direct effect on community structure such as the trophic cascade),

and/or due to the parallel effects of common abiotic drivers (e.g. temperature affects

directly intraspecific diversity, community structure and ecosystem functions). As it has

been done recently for the BEF (Grace *et al.*, 2016; Duffy *et al.*, 2016) and the SGDC

(Fourtune *et al.*, 2016; Lamy *et al.*, 2017) frameworks, we argue that a future important

step will be to combine appropriate statistical methods (e.g. path analysis, Shipley,

2000; Grace, 2006) and powerful experimental designs to disentangle causal

relationships between intraspecific diversity, community structure, ecosystem functions

and their common environment.

We further demonstrated that intraspecific variation has significant ecological

effects across a large set of species, hence confirming and refining a previous estimate

based on a restricted set of species (Des Roches *et al.*, 2018). By tripling the number of

species being investigated in this meta-analysis, we taxonomically broaden the

conclusion that intraspecific variation matters for shaping ecological dynamics, and that

the ecological effects of intraspecific variation in nature might be more common than

expected. Moreover, we demonstrated that previous estimates provided by (Des Roches

et al., 2018) were not upwardly biased (which was expected due to the focus on a non-

random pool of species), but were well within the range of estimates we report here and

actually tended to be slightly underestimated. Our finding hence strongly supports the

idea that adaptive and non-adaptive processes can lead to unique populations

differentially and significantly affecting ecological systems.

Although our conclusions held true for many species, the ecological effects of

intraspecific variation were not homogeneous across species, and this is partly

explained by the trophic level of species. Indeed –and according to our expectations-,

the ecological effects of intraspecific variation were stronger when the target species

was a primary producer than when it was a consumer. Several non-exclusive

mechanisms might explain this result. For instance, many primary producers considered

here provide a habitat for many invertebrates species (Southwood, Brown, & Reader,

1979) (this is not the case for considered consumer species), and this habitat can be

modulated by changes in plant structures. Non-exclusively, the relative biomass of

primary producers is higher than that of consumers; as such primary producers could

generate stronger effects on communities and ecosystems than consumers because of a

pure biomass effect. However, when decomposing variance in the ecological effects of

intraspecific variation into a greater detail, we found that these effects were stronger for

arthropod and herbaceous species than for fish (and to a lesser extent tree) species. This

suggests that the trophic level of species cannot be seen as the only predictor of the

ecological effects of intraspecific diversity, and we argue that future works should aim

at testing specifically why intraspecific variation ecologically matters for some species

and not for others.

Finally, the effects of intraspecific variation were globally higher on ecosystems

than on communities, hence generalizing across organism and ecosystem types a similar

observation on freshwater consumer species (Palkovacs *et al.*, 2015). We can speculate

that this difference of effects holds because intraspecific variation acts on community

dynamics through trophic mechanisms, whereas ecosystem functions can be modulated

through both trophic *and* non-trophic interactions (e.g. excretion rate or leaf chemistry,

Vanni, 2002; Schmitz *et al.*, 2014). For instance, a consumer species varying

intraspecifically in resource selectivity and/or consumption rate can affect both the

community structure and productivity of its resource (Harmon *et al.*, 2009). Non-trophic

mechanisms such as variability in organisms stoichiometry can reinforce the effect of

the consumer species on several ecosystem functions (e.g. primary production or soil

mineralization, (Schmitz, Hawlena, & Trussell, 2010; Hawlena *et al.*, 2012).

Alternatively -yet non-exclusively-, changes in ecosystem functions might be due to

both direct effects of intraspecific diversity and indirect effects of intraspecific diversity

mediated through changes in community structure, which may overall strengthened the

effects of intraspecific diversity at the ecosystem level. However, as observed for the

taxonomic predictors, an analysis accounting for a greater detail revealed that a lot of

variation exists between sub-categories of response variables (Fig. 4B), and that the

dichotomy between variables measured at the community and ecosystem levels is not a

straightforward. Although providing the first attempts to decompose variance in the

ecological effects of intraspecific variation, our findings call for further studies on

various taxa and in different ecosystems to fully appreciate the intensity of ecological

effects of intraspecific diversity (Box 1).

V.   Conclusions

(1)   We provided empirical evidence that the ecological effects of intraspecific

richness increase asymptotically, paralleling well-known patterns observed at the

interspecific level (Loreau, 2000; Hooper *et al.*, 2005) and confirming conceptual ideas

(Hughes *et al.*, 2008).

(2)   We also demonstrated that the variation of phenotypes or genotypes within

species is an important driver of community and ecosystem dynamics. These major

ecological effects of intraspecific diversity held true for organisms including several

species of plants and animals, although much remains to be tested (Box 1).

(3)   Overall these findings demonstrated that intraspecific diversity -beyond its

importance for species to adapt to environmental changes- is an important facet of

biodiversity for understanding and predicting the ecological dynamics of communities

and ecosystems, which reinforces the needs for better appraising the causes and

consequences of intraspecific diversity in natural populations and for improving

associated conservation plans.

VI.   Acknowledgements

We warmly thank José M. Montoya for valuable comments on a previous version of the manuscript. We also thank two anonymous reviewers for their constructive and useful comments on a previous version of the manuscript. AR is financially supported by a Doctoral scholarship from the Université Fédérale de Toulouse. This work was undertaken at SETE, which is part of the "Laboratoire d'Excellence" (LABEX) entitled TULIP (ANR-10-LABX-41).

VII.   References

579
Bailey, J.K., Schweitzer, J.A., Ubeda, F., Koricheva, J., LeRoy, C.J., Madritch, M.D., Rehill, B.J., Bangert, R.K., Fischer, D.G., Allan, G.J. & Whitham, T.G. (2009) From genes to ecosystems: a synthesis of the effects of plant genetic factors across levels of organization. *Philosophical Transactions of the Royal Society B: Biological Sciences* **364**, 1607–1616.

Barbour, R.C., O'Reilly-Wapstra, J.M., De Little, D.W., Jordan, G.J., Steane, D.A., Humphreys, J.R., Bailey, J.K., Whitham, T.G. & Potts, B.M. (2009) A geographic mosaic of genetic variation within a foundation tree species and its community-level consequences. *Ecology* **90**, 1762–1772.

Bartels, P., Cucherousset, J., Steger, K., Eklöv, P., Tranvik, L.J. & Hillebrand, H. (2012) Reciprocal subsidies between freshwater and terrestrial ecosystems structure consumer resource dynamics. *Ecology* **93**, 1173–1182.

de Bello, F., Lavorel, S., Albert, C.H., Thuiller, W., Grigulis, K., Dolezal, J., Janecek, S. & Leps, J. (2011) Quantifying the relevance of intraspecific trait variability for functional diversity. *Methods in Ecology and Evolution* **2**, 163–174.

Bolnick, D.I., Amarasekare, P., Araújo, M.S., Bürger, R., Levine, J.M., Novak, M., Rudolf, V.H., Schreiber, S.J. & Urban, M.C. (2011) Why intraspecific trait variation matters in community ecology. *Trends in ecology & evolution* **26**, 183–192.

Bolnick, D.I., Svanbäck, R., Fordyce, J.A., Yang, L.H., Davis, J.M., Hulsey, C.D. & Forister, M.L. (2003) The Ecology of Individuals: Incidence and Implications of Individual Specialization. *The American Naturalist* **161**, 1–28.

Booth, R.E. & Grime, J.P. (2003) Effects of genetic impoverishment on plant
community diversity. *Journal of Ecology* **91**, 721–730.

Burkle, L.A., Souza, L., Genung, M.A. & Crutsinger, G.M. (2013) Plant genotype,
nutrients, and G X E interactions structure floral visitor communities. *Ecosphere*
4, UNSP 113.

Burnham, K.P. & Anderson, D.R. (2002) *Model Selection and Multimodel Inference: A*
*Practical Information-Theoretic Approach*Springer, New York.

Cadotte, M.W., Carscadden, K. & Mirotchnick, N. (2011) Beyond species: functional
diversity and the maintenance of ecological processes and services: Functional
diversity in ecology and conservation. *Journal of Applied Ecology* **48**, 1079–
1087.

Cardinale, B.J., Duffy, J.E., Gonzalez, A., Hooper, D.U., Perrings, C., Venail, P.,
Narwani, A., Mace, G.M., Tilman, D., Wardle, D.A., Kinzig, A.P., Daily, G.C.,
Loreau, M., Grace, J.B., Larigauderie, A., et al. (2012) Biodiversity loss and its
impact on humanity. *Nature* **486**, 59–67.

Chapin, S.F., Zavaleta, E.S., Eviner, V.T., Naylor, R.L., Vitousek, P.M., Reynolds,
H.L., Hooper, D.U., Lavorel, S., Sala, O.E., Hobbie, S.E., Mack, M.C. & Díaz,
S. (2000) Consequences of changing biodiversity. *Nature* **405**, 234–242.

Crutsinger, G.M. (2006) Plant Genotypic Diversity Predicts Community Structure and
Governs an Ecosystem Process. *Science* **313**, 966–968.

Crutsinger, G.M., Rudman, S.M., Rodriguez-Cabal, M.A., McKown, A.D., Sato, T.,
MacDonald, A.M., Heavyside, J., Geraldes, A., Hart, E.M., LeRoy, C.J. & El-
Sabaawi, R.W. (2014) Testing a 'genes-to-ecosystems' approach to
understanding aquatic-terrestrial linkages. *Molecular Ecology* **23**, 5888–5903.

Des Roches, S., Post, D.M., Turley, N.E., Bailey, J.K., Hendry, A.P., Kinnison, M.T.,
Schweitzer, J.A. & Palkovacs, E.P. (2018) The ecological importance of
intraspecific variation. *Nature Ecology & Evolution* **2**, 57–64.

Díaz, S., Purvis, A., Cornelissen, J.H.C., Mace, G.M., Donoghue, M.J., Ewers, R.M.,
Jordano, P. & Pearse, W.D. (2013) Functional traits, the phylogeny of function,
and ecosystem service vulnerability. *Ecology and Evolution* **3**, 2958–2975.

Downing, A.L. & Leibold, M.A. (2002) Ecosystem consequences of species richness
and composition in pond food web. *Nature* **416**, 837–-841.

Duffy, J.E., Godwin, C.M. & Cardinale, B.J. (2017) Biodiversity effects in the wild are
common and as strong as key drivers of productivity. *Nature* **549**, 261–264.

Duffy, J.E., Lefcheck, J.S., Stuart-Smith, R.D., Navarrete, S.A. & Edgar, G.J. (2016)
Biodiversity enhances reef fish biomass and resistance to climate change.
Proceedings of the National Academy of Sciences **113**, 6230–6235.

Egger, M., Smith, G.D., Schneider, M. & Minder, C. (1997) Bias in meta-analysis
detected by a simple, graphical test. *Bmj* **315**, 629–634.

El-Sabaawi, R.W., Bassar, R.D., Rakowski, C., Marshall, M.C., Bryan, B.L., Thomas,
S.N., Pringle, C., Reznick, D.N. & Flecker, A.S. (2015) Intraspecific phenotypic
differences in fish affect ecosystem processes as much as bottom-up factors.
Oikos **124**, 1181–1191.

Fourtune, L., Paz-Vinas, I., Loot, G., Prunier, J.G. & Blanchet, S. (2016) Lessons from
the fish: a multi-species analysis reveals common processes underlying similar
species-genetic diversity correlations. *Freshwater Biology* **61**, 1830–1845.

Fridley, J.D. & Grime, J.P. (2010) Community and ecosystem effects of intraspecific
genetic diversity in grassland microcosms of varying species diversity. *Ecology*
91, 2272–2283.

Grace, J.B. (2006) *Structural equation modeling and natural systems*Cambridge
University Press.

Grace, J.B., Anderson, T.M., Seabloom, E.W., Borer, E.T., Adler, P.B., Harpole, W.S.,
Hautier, Y., Hillebrand, H., Lind, E.M., Pärtel, M., Bakker, J.D., Buckley, Y.M.,
Crawley, M.J., Damschen, E.I., Davies, K.F., et al. (2016) Integrative modelling
reveals mechanisms linking productivity and plant species richness. *Nature* **529**,
390–393.

Grant, P.R. & Grant, B.R. (2006) Evolution of Character Displacement in Darwin's
Finches. *Science* **313**, 224–226.

Harmon, L.J., Matthews, B., Des Roches, S., Chase, J.M., Shurin, J.B. & Schluter, D.
(2009) Evolutionary diversification in stickleback affects ecosystem functioning.
Nature **458**, 1167–1170.

Hawlena, D., Strickland, M.S., Bradford, M.A. & Schmitz, O.J. (2012) Fear of
Predation Slows Plant-Litter Decomposition. *science* **336**, 1434–1438.

Hendry, A.P. (2016) *Eco-Evolutionary Dynamics*Princeton University Press.

Hillebrand, H. & Matthiessen, B. (2009) Biodiversity in a complex world: consolidation
and progress in functional biodiversity research: Consolidation and progress in
BDEF research. *Ecology Letters* **12**, 1405–1419.

Hooper, D.U., Chapin, F.S., Ewel, J.J., Hector, A., Inchausti, P., Lavorel, S., Lawton,
J.H., Lodge, D.M., Loreau, M., Naeem, S., Schmid, B., Set?l?, H., Symstad,
A.J., Vandermeer, J. & Wardle, D.A. (2005) Effects of biodiversity on
ecosystem functioning: a consensus of current knowledge. *Ecological*
Monographs **75**, 3–35.

Horvathova, T., Nakagawa, S. & Uller, T. (2012) Strategic female reproductive
investment in response to male attractiveness in birds. *Proceedings of the Royal*
Society B: Biological Sciences **279**, 163–170.

Howeth, J.G., Weis, J.J., Brodersen, J., Hatton, E.C. & Post, D.M. (2013) Intraspecific
phenotypic variation in a fish predator affects multitrophic lake metacommunity
structure. *Ecology and Evolution* **3**, 5031–5044.

Hughes, A.R., Inouye, B.D., Johnson, M.T.J., Underwood, N. & Vellend, M. (2008)
Ecological consequences of genetic diversity. *Ecology Letters* **11**, 609–623.

Jackrel, S.L. & Wootton, J.T. (2014) Local adaptation of stream communities to
intraspecific variation in a terrestrial ecosystem subsidy. *Ecology* **95**, 37–43.

Johnson, M.T.J., Lajeunesse, M.J. & Agrawal, A.A. (2006) Additive and interactive
effects of plant genotypic diversity on arthropod communities and plant fitness.
Ecology Letters **9**, 24–34.

Koricheva, J., Gurevitch, J. & Mengersen, K. (2013) *Handbook of Meta-Analysis in*
*Ecology and Evolution*Princeton University press, Princeton, New Jersey.

Lamy, T., Laroche, F., David, P., Massol, F. & Jarne, P. (2017) The contribution of
species-genetic diversity correlations to the understanding of community
assembly rules. *Oikos* **126**, 759–771.

Lecerf, A. & Chauvet, E. (2008) Intraspecific variability in leaf traits strongly affects
alder leaf decomposition in a stream. *Basic and Applied Ecology* **9**, 598–605.

Loreau, M. (2000) Biodiversity and ecosystem functioning: recent theoretical advances.
Oikos **91**, 3–17.

Loreau, M., Mouquet, N. & Holt, R.D. (2003) Meta-ecosystems: a theoretical
framework for a spatial ecosystem ecology. *Ecology Letters* **6**, 673–679.

Lowe, W.H., Kovach, R.P. & Allendorf, F.W. (2017) Population Genetics and
Demography Unite Ecology and Evolution. *Trends in Ecology & Evolution* **32**,
141–152.

Madritch, M.D., Greene, S.L. & Lindroth, R.L. (2009) Genetic mosaics of ecosystem
functioning across aspen-dominated landscapes. *Oecologia* **160**, 119–127.

Manly, B.F.. (1997) *Randomization, Bootstrap and Monte Carlo Methods in*
*Biology*Chapman and Hall, London.

Matthews, B., Aebischer, T., Sullam, K.E., Lundsgaard-Hansen, B. & Seehausen, O.
(2016) Experimental Evidence of an Eco-evolutionary Feedback during
Adaptive Divergence. *Current Biology* **26**, 483–489.

Matthews, B., De Meester, L., Jones, C.G., Ibelings, B.W., Bouma, T.J., Nuutinen, V.,
de Koppel, J. van & Odling-Smee, J. (2014) Under niche construction: an
operational bridge between ecology, evolution, and ecosystem science.
Ecological Monographs **84**, 245–263.

Matthews, B., Narwani, A., Hausch, S., Nonaka, E., Peter, H., Yamamichi, M., Sullam,
K.E., Bird, K.C., Thomas, M.K., Hanley, T.C. & Turner, C.B. (2011) Toward an
integration of evolutionary biology and ecosystem science: Integration of
evolutionary biology and ecosystem science. *Ecology Letters* **14**, 690–701.

Mimura, M., Yahara, T., Faith, D.P., Vázquez-Domínguez, E., Colautti, R.I., Araki, H.,
Javadi, F., Núñez-Farfán, J., Mori, A.S., Zhou, S., Hollingsworth, P.M., Neaves,

L.E., Fukano, Y., Smith, G.F., Sato, Y.-I., et al. (2016) Understanding and
719   monitoring the consequences of human impacts on intraspecific variation.
720   *Evolutionary Applications* **10**, 121–139.

721   Naeem, S., Thompson, Lindsey J, Lawler, Sharon P, Lawton, John H & Woodfin,
722   Richard M (1994) Declining biodiversity can alter the performance of
723   ecosystems. *Nature* **368**, 734–737.

724   Nakagawa, S. & Cuthill, I.C. (2007) Effect size, confidence interval and statistical
725   significance: a practical guide for biologists. *Biological Reviews* **82**, 591–605.

726   Nakagawa, S. & Santos, E.S.A. (2012) Methodological issues and advances in
727   biological meta-analysis. *Evolutionary Ecology* **26**, 1253–1274.

728   Odling-Smee, J., Laland, K.N. & Feldman, M.W. (2003) *Niche Construction The*
729   *neglected process in evolution*Princeton University Press.

730   Paine, R.T. (1969) A note on trophic complexity and community stability. *The*
731   *American Naturalist* **103**, 91–93.

732   Palkovacs, E.P., Fryxell, D.C., Turley, N.E. & Post, D.M. (2015) Ecological Effects of
733   Intraspecific Consumer Biodiversity for Aquatic Communities and Ecosystems.
734   In *Aquatic Functional Biodiversity* pp. 37–51. Elsevier.

735   Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D. & R Core Team (2014) nlme: Linear and
736   Nonlinear Mixed Effects Models_. R package version 3.1-117.

737   Post, D.M., Palkovacs, E.P., Schielke, E.G. & Dodson, S.I. (2008) Intraspecific
738   variation in a predator affects community structure and cascading trophic
739   interactions. *Ecology* **89**, 2019–2032.

740   Reiss, J., Bridle, J.R., Montoya, J.M. & Woodward, G. (2009) Emerging horizons in
741   biodiversity and ecosystem functioning research. *Trends in Ecology & Evolution*
742   **24**, 505–514.

743   Rudgers, J.A. & Whitney, K.D. (2006) Interactions between insect herbivores and a
744   plant architectural dimorphism. *Journal of Ecology* **94**, 1249–1260.

745   Rudman, S.M., Rodriguez-Cabal, M.A., Stier, A., Sato, T., Heavyside, J., El-Sabaawi,
746   R.W. & Crutsinger, G.M. (2015) Adaptive genetic variation mediates bottom-up
747   and top-down control in an aquatic ecosystem. *Proceedings of the Royal Society*
748   *B: Biological Sciences* **282**, 125–132.

749   Rudman, S.M. & Schluter, D. (2016) Ecological Impacts of Reverse Speciation in
750   Threespine Stickleback. *Current Biology* **26**, 490–495.

751   Schmitz, O.J., Grabowski, J.H., Peckarsky, B.L., Preisser, E.L., Trussell, G.C. &
752   Vonesh, J.R. (2008) From individuals to ecosystem function: toward an
753   integration of evolutionary and ecosystem ecology. *Ecology* **89**, 2436–2445.

754   Schmitz, O.J., Hawlena, D. & Trussell, G.C. (2010) Predator control of ecosystem
755   nutrient dynamics. *Ecology Letters* **13**, 1199–1209.

Schmitz, O.J., Raymond, P.A., Estes, J.A., Kurz, W.A., Holtgrieve, G.W., Ritchie,
M.E., Schindler, D.E., Spivak, A.C., Wilson, R.W., Bradford, M.A.,
Christensen, V., Deegan, L., Smetacek, V., Vanni, M.J. & Wilmers, C.C. (2014)
Animating the Carbon Cycle. *Ecosystems* **17**, 344–359.

Schoener, T.W. (2011) The Newest Synthesis: Understanding the Interplay of
Evolutionary and Ecological Dynamics. *Science* **331**, 426–429.

Shipley, B. (2000) *Cause and correlation in biology: a user's guide to path analysis,*
*structural equation and causal inference*Cambridge University Press.

Siefert, A., Violle, C., Chalmandrier, L., Albert, C.H., Taudiere, A., Fajardo, A.,
Aarssen, L.W., Baraloto, C., Carlucci, M.B., Cianciaruso, M.V., de L. Dantas,
V., de Bello, F., Duarte, L.D.S., Fonseca, C.R., Freschet, G.T., et al. (2015) A
global meta-analysis of the relative extent of intraspecific trait variation in plant
communities. *Ecology Letters* **18**, 1406–1419.

Southwood, T.R., Brown, V.K. & Reader, P.M. (1979) The relationships of plant and
insect diversities in succession. *Biological Journal of the Linnean Society* **12**,
327–348.

Thompson, J.N. (1998) Rapid evolution as an ecological process. *Trends in Ecology &*
Evolution **13**, 329–332.

Vanni, M.J. (2002) Nutrient Cycling by Animals in Freshwater Ecosystems. *Annual*
Review of Ecology and Systematics **33**, 341–370.

Vellend, M. (2005) Species Diversity and Genetic Diversity: Parallel Processes and
Correlated Patterns. *The American Naturalist* **166**, 199–215.

Vellend, M. & Geber, M.A. (2005) Connections between species diversity and genetic
diversity: Species diversity and genetic diversity. *Ecology Letters* **8**, 767–781.

Violle, C., Enquist, B.J., McGill, B.J., Jiang, L., Albert, C.H., Hulshof, C., Jung, V. &
Messier, J. (2012) The return of the variance: intraspecific variability in
community ecology. *Trends in Ecology & Evolution* **27**, 244–252.

Violle, C., Navas, M.-L., Vile, D., Kazakou, E., Fortunel, C., Hummel, I. & Garnier, E.
(2007) Let the concept of trait be functional! *Oikos* **116**, 882–892.

Werner, E.E. & Peacor, S.D. (2003) A review of trait-mediated indirect interactions in
ecological communities. *Ecology* **84**, 1083–1100.

Whitham, T.G., Bailey, J.K., Schweitzer, J.A., Shuster, S.M., Bangert, R.K., LeRoy,
C.J., Lonsdorf, E.V., Allan, G.J., DiFazio, S.P., Potts, B.M., Fischer, D.G.,
Gehring, C.A., Lindroth, R.L., Marks, J.C., Hart, S.C., et al. (2006) A
framework for community and ecosystem genetics: from genes to ecosystems.
Nature Reviews Genetics **7**, 510–523.

Whitham, T.G., Young, W.P., Martinsen, G.D., Gehring, C.A., Schweitzer, J.A.,
Shuster, S.M., Wimp, G.M., Fischer, D.G., Bailey, J.K., Lindroth, R.L.,

Woolbright, S. & Kuske, C.R. (2003) Community and ecosystem genetics: A
consequence of the extended phenotype. *Ecology* **84**, 559–573.

Figure captions

Fig. 1. Publication year of the 77 selected articles used in the present study.

Fig. 2. (A) Relationship between intraspecific richness and effect size (ln*RR*) on

ecological dynamics. The line represents the shape of the relationship as predicted by

the exponential asymptotic non-linear mixed effect model. Blue shadow represents 95%

CI. **(B)** Medians (vertical dotted lines) and density of absolute effect size (ln*RR*) for

intraspecific and interspecific richness on ecological dynamics.

Fig. 3. Density probability of *Zr* effect size. The vertical grey straight line represents the

resampled *Zr* mean under the null hypothesis (confidence intervals not shown because

they were to narrow); the black curve shows the distribution of observed *Zr* and its

mean (vertical black straight line), grey shades represent 95% CIs.

Fig. 4. (A) Mean of *Zr* effect size in function of the model species. The sample sizes

(N) are given as indication, and represent the number of species and the number of

effect sizes respectively. The horizontal dotted line represents the mean effect size; error

bars represent ± 1 SE. **(B)** Mean of *Zr* effect size in function of the ecological response

variables. The sample sizes (N) of the number of effect sizes are given as indication.

The horizontal dotted line represents the mean effect size; error bars represent ± 1 SE.

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Table 1. Summary table of the model selection by AIC comparison to explain the shape
of the relationship between the ecological consequences and the intraspecific richness.
Models were run as non-linear mixed-effect models with the articles ID as random
factor, equations and parameters estimates are given as indication. IR = intraspecific
richness, lnRR = effect size of intraspecific richness of ecological dynamics.
834

Model	AIC	ΔAIC	AIC Weight	Equation	Parameter estimates
Asymptotic exponential model	-141.726	0	0.862	$lnRR = a * (1 - e^{-b*IR})$	$a = 0.201$ $b = 0.572$
Michaelis-Mentens model	-138.032	3.694	0.136	$lnRR = V * \dfrac{IR}{k + IR}$	$V = 0.225$ $k = 1.001$
Linear model	-129.038	12.689	0.001	$lnRR = b0 + b1 * IR$	$b1 = 0.011$ $b0 = 0.118$
Null model	-120.455	-21.2714	< 0.001	$lnRR = b0$	$b0 = 0.172$

Box 1. **Research perspectives on the relationships between intraspecific diversity and the dynamics of communities and ecosystems.**

We hereafter highlight several research avenues that may greatly favour the understanding of the relationships between intraspecific diversity and the dynamics of communities and ecosystems.

(i) Our review demonstrated that the ecological effect size of intraspecific diversity strongly varies from one species to another. We demonstrated that this inter-species variance in effect size is partly explained by the type of organism (i.e. primary producer or consumer). However, species composing a community also vary according to its commonness, its role in the ecosystem (e.g. keystone species or not, (Paine, 1969)), its body length (or height for plants), its life-history strategy (e.g. r-K strategy), its recent history (e.g. whether or not the species is non-native), functional characteristics (e.g. stoichiometry, metabolism), etc... A next challenge will be to better partition this inter-species variance in effect size by determining the species characteristics that best predict the strength of effect sizes, which is a pre-requisite to set coherent conservation plans at the intraspecific level (Mimura *et al.*, 2016);

(ii) Intraspecific diversity has often been manipulated for a single target species at a time. However, in nature, species are interacting and we argue that next studies should manipulate intraspecific diversity within multiple interacting species to better reflect reality, and to partition the relative importance of intraspecific diversity between interacting (potentially co-evolving) species;

(iii) Ecosystems are connected one each others through cross-ecosystem fluxes of matter (Loreau, Mouquet, & Holt, 2003). For instance, freshwater ecosystems such as rivers receive a large amount of material from the surrounding terrestrial ecosystem (e.g. dead leaves falling from trees, (Bartels *et al.*, 2012). A major -yet fascinating- challenge would be to assess the relative consequence of intraspecific diversity on allochtonous ecosystems *vs* autochtonous ecosystems; for instance testing whether or not the consequences of intraspecific diversity manipulated in a fish species is higher on aquatic ecosystems than on terrestrial ecosystems (Jackrel & Wootton, 2014; Crutsinger *et al.*, 2014; Rudman *et al.*, 2015), for the first attempts to quantify across-ecosystem effects of intraspecific diversity). This is an important next step to quantify in a more integrative way the importance of intraspecific diversity in natural systems;

(iv) Previous studies mainly assessed the consequences on intraspecific diversity by considering the genetic or the phenotypic characteristics of organisms as a whole. However, some traits have been shown to be more important for ecological dynamics than others; this is the case of *functional effect traits* that are defined as the traits having an ecological impact (e.g. excretion rate or leaf nutrient content, (Violle *et al.*, 2007; Díaz *et al.*, 2013). We suggest that directly manipulating functional traits diversity of individuals within species, rather than genotypic or phenotypic diversity as a whole would be a powerful mean to understand the mechanisms by which intraspecific diversity act on ecological dynamics;

(v) Although we contributed to generalise the idea that intraspecific diversity is essential to the community and ecosystem facets of ecological dynamics, comparisons to other major and indisputable ecological constraints (e.g. temperature, interspecific

biodiversity, human disturbances) have rarely been conducted (but see (Burkle *et al.*, 2013; El-Sabaawi *et al.*, 2015). To better evaluate and quantify the importance of intraspecific diversity in natural system, it is urgent to assess the relative contribution of intraspecific diversity compare to indisputable determinants of ecosystem functioning. This is a key step to confirm that intraspecific is actually part of these indisputable determinants, and not a random signal in complex systems.