1

An amplitude code increases the efficiency of information transmission across a visual synapse

Ben James[1], Léa Darnet[1], José Moya-Díaz[1], Sofie-Helene Seibel[1] and Leon Lagnado[1]*

School of Life Sciences, University of Sussex, Brighton BN1 9QG, UK

[1]These authors contributed equally to this work.
*Corresponding author: l.lagnado@sussex.ac.uk

Most neurons in the brain transmit information digitally using sequences of spikes that trigger release of synaptic vesicles of fixed size. The first stages of vision and hearing are distinct in operating with analogue signals, but it is unclear how these are recoded for synaptic transmission. By imaging the release of glutamate in live zebrafish, we demonstrate how ribbon synapses of retinal bipolar cells transmit analogue visual signals by changes in both the rate and amplitude of synaptic events. Higher contrasts released glutamate packets composed of more vesicles and coding by amplitude often continued after rate coding had saturated. Glutamate packets equivalent to five vesicles transmitted four times as many bits of information per vesicle compared to independent release events. By discretizing analogue signals into sequences of numbers ranging up to eleven, ribbon synapses increase the dynamic range, temporal precision and efficiency with which visual information is transmitted.

The spike code of neurons has been studied in detail[1,2] but much less is known about the vesicle code transmitting information across the synapse[3]. Since the work of Katz[4-6], the general view has been that changes in presynaptic potential are communicated by modulating the mean rate of a Poisson process in which vesicles at the active zone are triggered to fuse independently[7-9]. At most synapses, the voltage signal controlling this process is digital, arriving in the form of an all-or-nothing spike, and the output is also digital, in the form of a vesicle releasing a fixed packet of neurotransmitter[2]. But in the first stages of vision and hearing the neural signal is in analogue form - continuous changes in membrane

potential that are graded with the strength of the stimulus [10]. How are these sensory signals

recoded for transmission across a synapse?

Synapses driven by analogue signals are distinguished from those driven digitally by

the presence of a specialized structure, the ribbon, that holds many tens of vesicles just

behind the active zone[11,12]. It has generally been assumed that these ribbon synapses carry

out the key step of analogue-to-digital conversion when the graded voltage signal is

represented as changes in the rate of events releasing just one quantum of

neurotransmitter[7,8,13-15].   A number of *in vitro* studies have, however, shown that ribbon

synapses are also capable of modulating the size of the packet of neurotransmitter through

a process termed coordinated multivesicular release (CMVR)[15-20].   This observation is

intriguing because it suggests that the output of a ribbon synapse may not be a single

"symbol", but a number of symbols varying in amplitude[21]. The functional role of CMVR has,

however, been unclear because it has not been observed in response to light or sound.

To understand whether the coordinated release of vesicles contributes to the

transmission of visual information we used the fluorescent glutamate reporter iGluSnFR[22] to

image the output of ribbon synapses in the retina of zebrafish. Optimizing the signal-to-

noise ratio allowed us to count vesicles released from individual active zones *in vivo* and

investigate how they are used to transmit the visual signal while it is still in analogue form.

We show that ribbon synapses of bipolar cells employ a coding strategy that is a hybrid of

the known rate code and of changes in the amplitude of synaptic events, which we term the

amplitude code. We describe how amplitude coding confers several advantages, including

the ability to signal contrast beyond the range where the rate code has saturated, improving

the temporal precision of transmission and increasing the efficiency of communication by

transmitting more bits of information per vesicle.

Results

63
64
65   **Optical detection of co-ordinated multivesicular release *in vivo***
66
All visual signals flow through bipolar cells because these are the only neurons that connect

the photoreceptors to ganglion cells the inner retina (Fig. 1a). To investigate how these

signals are represented as they cross the synapse, expression of iGluSnFR was driven in

bipolar cells using the *ribeye* promoter[14]. A periodic stimulus (5 Hz) triggered a train of

glutamate transients which were sampled by performing line scans through the terminal at 1

kHz (Fig 1b and c; Video S2). These transients were generated by the electrical signal

arriving from the cell body rather than glutamate spillover from neighbouring synapses

because they were destroyed by ablating the soma of the cell (Fig. S8). Indeed, closer

analysis of the intensity profile along a line demonstrated that iGluSnFR signals were

constrained to spatial scales of ~1-2 mm making it possible to distinguish fusion events at

neighbouring active zones by spatial demixing (Fig. 1c and Supplementary Figure 2).

iGluSnFR signals at adjacent active zones did not always coincide in time, reflecting the

stochastic nature of vesicle release (Fig. 1c) and, most notably, the size of glutamate

transients varied widely (Fig. 1c and d).

82   **Figure 1 near here**

84   To improve estimates of the timing and amplitude of glutamatergic events, a Wiener

filter was applied[9] in which the raw iGluSnFR trace (Fig. 1d, top) was deconvolved (middle)

using a temporal kernel estimated by averaging a number of isolated events (Supplementary

Figures 3 and 4; described in detail in the Supplementary Methods). Histograms of event

amplitudes measured from deconvolved traces displayed several peaks at equal intervals

indicating that transients of different sizes were composed of multiples of an underlying

quantal event (Fig. 1e). These quanta are very likely to correspond to synaptic vesicles

given the electrophysiological evidence that ribbon synapses in auditory hair cells and the

retina can sometimes coordinate the fusion of two or more vesicles rather than releasing

them all independently[16,17,19]. Based on amplitude histograms allowing estimates of the

quantal signal, we partitioned iGluSnFR signals of different sizes into numbers of quanta

using a maximum likelihood estimate, as shown in the lower panel of Fig. 1d and described

in Supplementary Methods.

97   Two further observations indicated that larger glutamate events involved the

synchronized release of vesicles rather than coincidental and independent fusion occurring

within a narrow time window[16-19]. First, the time-courses of iGluSnFR signals containing

multiple quanta were indistinguishable from those containing just one (Supplementary Figure

4). Second, the distribution of time intervals between events consistently deviated from the

Poisson statistics expected for vesicles released independently with constant probability

(Supplementary Figure 9). Coordinated release of two or more vesicles was observed in all

of 186 active zones that we investigated in both the ON and OFF channels of the retina (74

ON and 112 OFF synapses).

Reverse-correlation at an active zone: the transmitter-triggered average

The existence of a process that synchronizes the release of two or more vesicles has been

recognized at ribbon synapses for some years, but it's functional role has not been clear[16-19].

Particularly puzzling has been the observation that, although the frequency of CMVR in hair

cells and bipolar cells is calcium-dependent, the amplitude of events is not[17,18].   At face

value, this would suggest that CMVR does not encode the loudness of sound or contrast of

a visual stimulus.   These experiments did not, however, measure responses to a

physiological stimulus but instead stimulated the synapse through a patch pipette[17,18].

Counting vesicles with iGluSnFR allowed us to make an *in vivo* investigation of how CMVR

contributes to the encoding of visual stimuli.

We began investigating this question by adapting an approach that has been widely

used to explore the information represented by spikes – calculation of the spike-triggered

average (STA). The STA is calculated by reverse-correlating the spike train generated by a

random ("white noise") stimulus to the stimulus itself[2,23] and provides an estimate of the

neurons tuning in the form of its linear filter in time or in space and time.   By reverse-

correlating events measured using iGluSnFR in response to full-field noise, we instead

constructed the "transmitter triggered average" (TTA) to describe the output of an active

zone.   This approach to understanding the function of a synapse was first proposed in

2004[3], but has not been realized until now.   A fundamental distinction between the STA and

TTA was that the while the spike output of a neuron contains just one symbol, here we are

dealing with a vesicle code involving a number of symbols (1, 2, 3 quanta etc). We therefore

calculated the TTA separately for synaptic events composed of different numbers of vesicles

(Fig. 2a).

The TTA revealed that the more quanta within an event the higher, on average, the

temporal contrast driving it (Fig. 2b).   The relationship between the number of quanta

released per event (Q_e) and the contrast in the temporal filter (C) could be described by a

first-order saturation of the form $C = C_{max} \times Q_e /(Q_e + Q_{1/2})$, where C_{max} = 19% and $Q_{1/2}$ = 2.4

vesicles (Fig. 2c).   Thus, CMVR encodes one of the most fundamental properties of a visual

stimulus – temporal contrast – in a simple and direct way.

137   **Figure 2 near here**

The TTA revealed a second distinction between events of different amplitude:

uniquantal events were characterized by temporal filters that were monophasic through both

ON and OFF channels, corresponding to a low-pass filter, but the TTA from ON synapses

were biphasic corresponding to band-pass characteristics with peak transmission at ~4Hz

(Fig. 2b). In other words, a period of hyperpolarization before depolarization favoured larger

synaptic events through the ON channel.   Such biphasic antagonism in the time-domain is

commonly observed in the temporal receptive fields of neurons early in the visual system [24],

where it has been proposed to underly the suppression of redundant signals [25].

Synchronizing the release of multiple vesicles is expected to amplify the output of a

bandpass filter to enhance the signaling of change in the time-domain. Together, the results

in Figs. 1 and 2 demonstrate that CMVR plays a fundamental role in transmitting the visual

signal[16,17,19].

A hybrid rate and amplitude code

To investigate further how changes in the amplitude of synaptic events transmitted

information about a visual stimulus we constructed contrast-response functions for individual

active zones. Fig. 3a shows a protocol used to quickly assess this function, in which the

contrast of a 5 Hz stimulus was increased in 10% steps. The first measure of response that

we used was the total number of quanta per cycle of the stimulus (Q_c), from which we

estimated the contrast generating the half-maximal response, $C_{1/2}$, as shown by the example

in Fig. 3b. This value occurs at the steepest part of the contrast-response function, and so

also defines the range in which the synapse signaled a change in contrast with the highest

sensitivity[2]. A first inspection of these traces shows that an increase in contrast increases

both the frequency and amplitude of glutamatergic events, as can also be seen in the

records in Fig, 4a, Fig. 5a and Fig. 6a.

Figure 3 near here

What are the relative contributions of the rate and amplitude codes? To assess how

these two modes of signaling operated in parallel, we focused on the range where contrast

sensitivity was highest, $C_{1/2} \pm 10\%$, using a second stimulus protocol in which different

contrasts were applied in 2 s steps separated by 2 s intervals, as shown in Fig. 6a. Q_c was

then factorized into two quantities: the number of events per cycle (E_c, representing the

contribution of the rate code) and the number of quanta per event (Q_e, the amplitude code).

The relative change in E_c and Q_e varied across synapses: in 38 of 55 active zones, an

increase in contrast caused E_c to rise more steeply than Q_e (Fig. 3c) but in the remaining 17

the amplitude code was dominant and Q_e rose more rapidly than E_c (Fig. 3d). Synapses in

which the amplitude of synaptic events was modulated more strongly displayed a second

striking property: when the rate code saturated, further increases in contrast were

represented *wholly* as increases in the size of the glutamate packets released (arrowed in

Fig. 3d). CMVR therefore extends the range of contrasts that can be signaled beyond those

allowed by the rate code alone.

One factor determining the shift towards release events composed of more quanta at

higher contrasts was the ON/OFF identity of the bipolar cell. The change in the distribution of

event amplitudes when the contrast was increased from 20% to 100% are shown in Fig. 4.

At 20% contrast, Q_e was not significantly different in the ON and OFF channels (0.48 ± 0.19

and 0.58 ± 0.18; n= 18 and 33, respectively). But at 100% contrast, the average event

amplitude in the OFF channel (5.07± 0.45 quanta) was significantly higher than the ON (2.89

± 0.5; p<0.0001 students t-test). The distribution of event amplitudes at 100% contrast were

also significantly different through ON and OFF channels (Chi-squared;p<0.0001). For

instance, there was a cut-off above which larger events uniquely coded higher contrasts

which was about six quanta and for the OFF channel and four quanta for the ON (Fig. 4b

and c). The largest synaptic events in both ON and OFF bipolar cells were composed of

about eleven vesicles (Fig. 4b). Thus, while most synapses in the brain signal with zeros

and ones, the ribbon synapses of bipolar cells go up to eleven.

Figure 4 near here

Ribbon synapses of bipolar cells therefore encode a visual stimulus through a hybrid

strategy involving changes in both the rate and amplitude of synaptic events. These

observations present a fundamentally distinct picture to that obtained by making

electrophysiological measurements of CMVR from hair cell ribbon synapses *in vitro*, where

the rate of events depended on the amplitude of modulation of the sinusoidal command

voltage but the average event size did not[20]. It may therefore be that CMVR at the ribbon

synapse of hair cells is not used to encode loudness, although further experiments in a more

intact preparation would be required to confirm this (see Discussion).

The amplitude code improves the temporal resolution of visual signals

Events of larger amplitude improved the temporal precision with which the visual signal was

transmitted. The timing of uniquantal events relative to the phase of a 5 Hz stimulus is

shown in Fig. 5a for contrasts of 20% and 100%. Uniquantal events displayed a standard

deviation ("temporal jitter") of 24 to 28 ms over a range of contrasts, while events composed

of 7 or more quanta jittered by as little as 2.5 ms (Fig. 5b). The spike trains of retinal

ganglion cells (RGCs) post-synaptic to bipolar cells can also encode visual signals of high

contrast with millisecond precision[26] and the stronger excitatory input from large

glutamatergic events may be one mechanism by which this is achieved.

Figure 5 near here

Stronger synchronization of larger synaptic events has also been observed by
making electrophysiological measurements of the output from hair cell ribbon synapses
driven by a sinusoidal command voltage[20]. This property can be quantified by a metric called
the vector strength, which varies between zero (random occurrence of events) and one
(perfect synchronization with the stimulus) [27]. The vector strength (s) is related to the
temporal jitter (r) by the relation $s = (2(1-r))^{1/2}/2Pf$ where f is the frequency. In the
experiments shown in Fig. 5, the vector strength achieved a near perfect value of 0.98 at
100% contrast. In comparison, *in vitro* experiments in hair cells have measured
multivesicular events to have vector strengths up to 0.6 at frequencies of 400 Hz[20]. These
different degrees of synchronization might reflect the higher operating frequencies of
auditory synapses compared to visual in combination with an inherent temporal limit in the
processes triggering CMVR.

The degree to which synaptic events were consistent in time depended not only on
event amplitude, but also the contrast of the stimulus eliciting the event (Fig. 5b). For
instance, the largest packets of glutamate consistently observed at 20% contrast contained
8 quanta with a temporal jitter of 12.1 ± 1.4 ms, while 8 quantal events at 100% contrast
jittered by 4.6 ± 0.6 ms. These observations run counter to electrophysiological
measurements of CMVR made from hair cell ribbon synapses, where the amplitude of the
changes in presynaptic potential had no effect on the size of the evoked events[20]. The
properties of the amplitude code shown in Fig. 5 are, however, similar to the properties of
the spike train in post-synaptic ganglion cells, which also become more temporally precise
as the contrast is increased[26]. A shift to larger glutamatergic inputs of higher temporal
precision may be one of the mechanisms that cause the spike trains of ganglion cells to
become less variable as contrast is increased.

Amplitude coding improves the efficiency of information transmission
The ionic mechanisms that generate neural signals and the synaptic processes that transmit
them are a major energetic cost to the brain[28,29]. The need to transmit information in an
energy-efficient manner has provided a unifying principle by which to understand the design
of sensory circuits[15,30,31]. This framework provides, for instance, a teleological explanation
for the use of analogue signaling to transmit early visual and auditory information: graded
changes in membrane potential transmit information more efficiently than spikes[15]. The
reason that analogue signals do not find more widespread use in the vertebrate nervous
system is that they dissipate over relatively short distances, requiring a switch to digital
signaling with regenerative spikes to relay information beyond the distance of one or two
small neurons. The finding that ribbon synapses employ a hybrid rate and amplitude code to

transmit analogue signals therefore led us to compare the amount of information contained
within uniquantal and multiquantal events.

Using information theory[32], we quantified the mutual information between a set of
stimuli of varying contrasts (Fig. 6a) and release events containing different numbers of
quanta (Fig. 6b).   Vesicles released individually carried an average of 0.125 bits of
information which is, as expected, significantly less than the 1-3.6 bits transmitted per spike
in post-synaptic ganglion cells[26,33].   The amount of information contained within events of
different size is shown in Fig. 6b. Larger events transmitted progressively more information
because they were rarer[34] (Fig. 4b and c) and driven preferentially by higher contrasts rather
than occurring randomly (Fig. 2). The relation between the specific information (i, bits) and
Q_e, the number of vesicles comprising the event, could be described as a power function of
the form $i = i_o + A.Q_e^x$, with i_o = 0.12 bits, A= 0.008, x = 2.8.   This supralinear relation
indicates that larger synaptic events transmit more information per vesicle.

269   **Figure 6 near here**

To explore this idea more directly we divided the amount of information in an event
by the number of vesicles it contained to provide a measurement of "vesicle efficiency"
before averaging across synapses to quantify the trend.   The change in this quantity is
plotted in Fig. 6c after normalizing to the value measured in the same synapse for one
vesicle.   This analysis confirmed that multiquantal events transmit more information *per*
vesicle. For instance, events composed of five quanta carried, on average, four times as
much information per vesicle as uniquantal events (Fig. 6c). The amplitude code at ribbon
synapses counteracts the reduction in the efficiency of information transmission that
accompanies the conversion from analogue to digital signaling[15].

Discussion
Achieving single-vesicle resolution with iGluSnFR has provided the opportunity to investigate
the synaptic code by counting vesicles released from individual active zones, much as
electrophysiology has been used to investigate the spike code of neurons[2]. This *in vivo*
approach demonstrates that the coordinated release of two or more vesicles is a
fundamental aspect of the strategy by which ribbon synapses recode an analogue signal for
transmission across the synapse. CMVR discretizes the analogue signal into about eleven
output values, with larger modulations of intensity increasing the average quantal content of
synaptic events (Fig. 4b and c). As a result, vesicles are used to generate a hybrid code
that consists of changes in both the rate and amplitude of events (Figs. 3-5). The amplitude

code complements the rate code by increasing the temporal accuracy (Fig. 5), efficiency

(Fig. 6) and operating range (Fig. 3d) of synapses transmitting the visual signal to RGCs.

An amplitude code at ribbon synapses

In essence, CMVR provides a mechanism by which the excitatory effects of glutamatergic

vesicles can be summed.   Normally, this process occurs on the dendrites of the post-

synaptic neuron, with signals from different vesicles adding up on the time scales of ~5-20

ms over which synaptic potentials decay[35]. An increase in stimulus strength increases the

rate of vesicle release and, therefore, the chances of summation. The key distinctions of

CMVR are vesicles are summed presynaptically and effectively instantaneous. It will

therefore be important to understand how much more effectively single multiquantal events

trigger spikes in ganglion cells compared to transient increases in the rate of release of

single vesicles. Reliable experimental answers to this question will be difficult to obtain

using electrophysiology because a patch pipette records all the current from the

compartment it is contacting, making it difficult to distinguish true CMVR from the

coincidental arrival of vesicles released from different synapses. A very direct and reliable

assessment of the postsynaptic effects of CNVR has, however, been possible in the auditory

system of bullfrog where afferents contacting only one hair cell ribbon can be patched.

Measuring the currents injected into the afferent by CMVR relative to the currents needed to

depolarize the fiber beyond threshold demonstrates that larger glutamatergic events will

reliably trigger spikes[20].

One factor determining how efficiently presynaptic summation of glutamate converts

into a postsynaptic current is whether or not post-synaptic receptors glutamate receptors

saturate. Saturation does not appear to be an issue at AII amacrine cells postsynaptic to

rod-driven bipolar cells[17] or fibers postsynaptic to hair cells[20], but it may nonetheless be that

there are variations in the efficiency of connections between different cell types. Given the

observation that different bipolar cells utilize CMVR to different degrees (Figs. 3 and 4), it

may even be that the distribution of sizes of glutamate packets is matched to the properties

of receptors on the postsynaptic neuron.

This study demonstrates that the average amplitude of synaptic events is determined

by the amplitude of the modulations in light intensity, while previous measurements of

CMVR have failed to detect any dependence on the strength of an electrophysiological

stimulus[17,20]. One possibility is that the normal calcium-dependence of CMVR was

compromised during whole-cell recordings because of wash-out of diffusible intracellular

components.   For instance, ribbon synapses contain high concentrations of diffusible

calcium buffers which act to localize calcium signals around calcium channels and inhibit the

release of vesicles docked further away[36]. Wash-out of these buffers through a pipette

allows more vesicles to be released by very brief calcium transients and may also disrupt the

normal relationship between the calcium current and CMVR. Regenerative calcium signals

("calcium spikes") within the bipolar cell terminal are also subject to washout[27] and may be

one of the mechanisms which coordinates the release of multiple vesicles. These problems

might be overcome by using the perforated patch technique to maintain intracellular

molecules while stimulating a bipolar cell or hair cell[36]. Certainly, the relationship between

the pre-synaptic calcium signal and the triggering of CMVR requires further investigation.

Potential mechanisms of CMVR

The cellular mechanisms synchronizing the release of two or more vesicles remain to be

discovered. One suggestion is that vesicles attached to the ribbon can fuse to each other to

create a multiquantal packet of glutamate which can subsequently fuse with the surface

membrane to generate a larger post-synaptic event[16]. This idea is supported by two lines of

evidence. Electron microscopy of bipolar cell terminals demonstrates the appearance of

large tubular structures on or around the ribbon after a period of stimulation[37,38] and

fluorophore-assisted light inactivation of the ribbon reduces both the rate and amplitude of

excitatory post-synaptic potentials[19].   It has been suggested that variable amounts of

glutamate might also be released from a single vesicle because of the dynamics of a fusion

pore[39], but this runs counter to evidence that bipolar cells transmit by full collapse of vesicles

into the membrane surface[40,41]. Electrophysiological evidence provides stronger support for

the idea that ribbon synapses can coordinate the simultaneous fusion of two or more

vesicles rather than releasing them all independently[16,17,19] and this idea is corroborated by

the quantization of iGluSnFR events shown in Fig. 1.

The presence of ribbon structures at synapses driven by analogue signals correlates

with a second functional specialization - a continuous mode of operation that allows the

transmission of sensory information to be maintained over prolonged periods[12,42](Figs. 4 and

5). The ribbon is thought to support continuous release by capturing vesicles from a mobile

pool in the cytoplasm[37] and then transporting them to the active zone[19]. The maximum rates

of continuous release that we observed using iGluSnFR were in the range of 50-100

vesicles s^{-1} per active zone (Fig. 3b), which is in agreement with measurements made using

the membrane dye FM1-43[42,43] and the fluorescent reporter protein sypHy[14].

Amplitude and rate coding in relation to the output of the retina

For vision to be useful, information about important stimuli must be transmitted over an

appropriately short time window. Multiquantal events were rare and strongly dependent on

temporal contrast, so their arrival provided more information about a preceding stimulus than

vesicles released individually (Fig. 6). Crucially, CMVR also encoded the timing of a stimulus

more precisely: the largest glutamatergic events jittered by just a few milliseconds relative to
a stimulus (Fig. 5), which is similar to the spike responses observed in postsynaptic ganglion
cells. These spike trains often consist of brief increases in firing rate from longer
background periods of silence, making it difficult to describe activity as a time-varying rate.
Rather, it has been suggested that "Firing events containing single spikes or bursts of spikes
are elicited precisely enough to convey distinct packets of visual information, and hence may
constitute the fundamental symbols in the neural code of the retina"[26]. It seems likely that
these symbols originate in the amplitude code of bipolar cell synapses.

To understand why a coding strategy based on amplitude might have arisen, it is
useful to think about the temporal requirements of a simple rate code. If a Poisson synapse
releasing all vesicles independently encodes an event by changing the rate of vesicle
release from R to kR, the signal-to-noise ratio achieved over an observation time Dt will be

$$SNR = \frac{(kR\Delta t - R\Delta t)}{\sqrt{(kR\Delta t + R\Delta t)}}$$

(Online Methods, Equation 4). If one considers the range of contrasts around $C_{1/2} \pm 10\%$,
the rate of vesicle release was modulated by a factor k of 1.8 (Q_e x E_c; Fig. 3c), from a basal
rate R of no more than 20 vesicles s^{-1}. To detect such a change with a SNR of 4 (i.e good
reliability) would require an observation time of ~420 ms. In comparison, within an OFF
synapse, any release event with amplitude greater than four quanta would *immediately*
signal an increase in contrast beyond 20% (Fig. 4c). This simple comparison illustrates one
of the potential advantages of recoding an analogue signal using symbols varying in
amplitude rather than as rates of digital events: an unexpected symbol immediately imparts
new information, while a stochastic rate code composed of a single event must be observed
over a time window sufficiently long to establish a significant change relative to the noise.
Further analysis of the statistics of CMVR will likely shed light on the properties of the vesicle
code transmitting visual information.

Acknowledgments

Many thanks to Jamie Johnston for discussions and to Hazel Smulders and Natasha
Bashford for looking after our zebrafish. Funding: Wellcome Trust and BBSRC. Author
contributions: This was a team project. S-HS performed molecular biology, fish transgenesis
and initial functional analysis; JM-D carried out experiments and analysis and helped
prepare the manuscript and supplemental information; LD performed molecular biology, and
carried out experiments and analysis; BJ wrote software, conceived and designed
information theory experiments and estimates of the transmitter-triggered average, helped
prepare the manuscript and supplemental information; LL conceived the project, designed
experiments, wrote software, analyzed results and prepared the manuscript.
407

408 **Competing interests**
409 The authors declare no competing financial interests.
410
411
412

Figure 1. Quantized release of glutamate imaged at individual active zones

a) Multiphoton section through the eye of a zebrafish larva expessing iGluSnFr in bipolar cells at 7dpf. **b)** Linescan through a single terminal. Arrowheads shows the position of two active zones. **c)**. The kymograph (top) shows the intensity profile along the line in B as a function of time. The red and black traces (middle) show the time-course of the iGluSnFR signal over the two active zones marked to the right of the kymograph and the stimulus is shown immediately below. After 2 s there was a switch from constant illumination to full field modulation at 20% contrast, 5 Hz. A section of the record (green box) is expanded Note variations in the amplitude of glutamate transients. Sometimes both active zones release glutamate (green dashed line) while on other occasions only AZ 1 (black dash) or AZ 2 (red dash). **d)** Raw (top) and deconvolved (middle) traces from a synapse stimulated at 50% contrast. Transients in the deconvolved trace were considered for analysis if they crossed a threshold (red line). The bottom panel shows the estimation of the number of quanta per event, based on the quantal size estimated in e. **e)** Amplitude histogram extracted from the deconvolved trace in **d**. The red line is a fit of six Gaussians, constrained to have equal distance between each peak but with all other parameters free. The peaks are separated by an inter-peak distance of ~0.22, which is the amplitude of the event equivalent to a single vesicle.

Figure 2. The transmitter-triggered average depends on the number of quanta in a release event

a) Example of iGluSnFR signals (top) elicited by a "white noise" stimulus (bottom). **b)** Upper traces: Linear filters extracted by reverse-correlation of responses composed of one quantum (left) and four to six quanta (right) from ON synapses (green, n = 9) and OFF synapses (red, n = 8). Multiquantal events encoded larger modulations in intensity. The plots below show the power spectra of the linear filters. Through the OFF channel, both uniquantal and multiquantal events were driven through low-pass filters, but in the ON channel multiquantal events were driven through a band-pass filter peaking at ~4 Hz. **c)** Relation between the Michelson contrast represented in the TTA and the number of quanta released within the events used for reverse correlation in ON and OFF terminals. The relation could be described by a first-order saturation of the form $C = C_{max}.N/(N + N_{1/2})$, where $C_{max} = 19\%$ and $N_{1/2} = 2.4$. Bars show standard errors.

449 **Figure 3. The relative contributions of coding by rate and amplitude**
450

a) Example of the iGluSnFR signal (top) elicited by a full-field stimulus of increasing contrast
delivered at 5 Hz (bottom). This protocol was used to quickly assess the half-point of the
contrast-response function ($C_{1/2}$). **b)** Contrast-response function extracted from the synapse
in **a**, with response (R) quantified as the total number of quanta per cycle of the stimulus.
The curve is a Hill equation of the form $C = R_{max}.(C^h/C^h + C^h_{1/2})$, where R_{max} = 9.8 quanta per
cycle (49 quanta s^{-1}), h = 3.5 and $C_{1/2}$ = 39% (dashed arrow). **c)** The relative change in
synaptic activity around $C_{1/2}$. The average number of events per cycle (E_c, black) is
compared with the average number of quanta per event (Q_e, red). Stimuli were delivered in 2
s episodes with 2 s rest as shown in Fig. 6a. In these 38 synapses an increase in contrast
caused E_c to rise more steeply than Q_e. **d)** In the remaining 17 synapses, Q_e rose more
steeply than E_c, which then saturated (dashed line) such that further increases in contrast
were signaled only by increases in the number of quanta per event (vertical arrow).
463
464
465 **Figure 4. Changes in the distribution of event amplitudes**
466

a) Representative traces of synaptic responses in ON and OFF terminals stimulated at 20%
and 100% contrast (5 Hz). Note the increase in the frequency and average amplitude of
events at higher contrast. **b)** Probability density of events composed of different numbers of
quanta in OFF synapses (n = 33). Events composed of more than 6 quanta were only
observed at contrasts greater than 20%. **c)** Distributions for ON synapses (n = 18). Events
composed of more than 4 quanta were only observed at contrasts greater than 20%. The
distribution at 100% contrast was significantly different to that observed in OFF synapses
(Chi-squared; p<0.0001).
475

476
477

Figure 5.   Multivesicular events increased the temporal precision of synaptic
transmission

a) Top: iGluSnFR signals from a terminal stimulated at 100% and 20% contrast. Note the
large variation in event amplitudes at lower contrast. Bottom: expanded time-scale showing
responses at 20% and 100% contrast superimposed on the phase of the stimulus (grey).
Events composed of fewer quanta are less synchronized than multiquantal events. Two
examples of events occurring at phases different to the majority are shown by the blue
arrows**.   b)**   Temporal jitter (s) as a function of the number of quanta per event (Qe) at
contrasts of 20% (blue, n= 57 synapses), 50% (red, n= 61) and 100% (black, n= 63). Events
composed of more quanta exhibit lower jitter at a given contrast and events of a given
amplitude exhibit lower jitter at higher contrasts. The relationship between s and Q_e could
be described as $s = s_{min} + (s_{max} - s_{min}) \cdot (e^{-Qe/k})$. At 100% contrast, s_{min} = 2.7 ms (dashed black
line), s_{max} = 30 ms and k = 2.6 quanta. At 20% contrast, s_{min} = 9.5 ms (dashed blue line),
s_{max} = 28 ms and k = 2.4 quanta.

492
493

Figure 6. Multivesicular events increased the efficiency of the vesicle code
a) Stimulus set for estimating mutual information. Stimuli of varying contrasts were selected
to span ± 10% of the range around which the contrast sensitivity was highest in steps of 2%.
Note variations in the frequency and amplitude of events at different contrast levels. Stimuli
were applied in two different sequences which were repeated alternately.   **b)** Specific
information per event (i, bits) as a function of Q_e, the number of vesicles comprising the
event. The curve describing the points is a power function of the form $i = i_o + A \cdot Q_e^x$, with i_o =
0.12 bits, A= 0.008, x = 2.8. Results pooled from 17 synapses. Bars show standard errors.
c) Specific information per vesicle normalized to the value measured for a uniquantal event
(i'). The curve describing the points is a power function of exponent 2.1. Results from the
same 17 synapses in **b**.

Online Methods

Zebrafish husbandry

Fish were raised and maintained under standard conditions on a 14 h light/10 h dark cycle[14]. To aid imaging, fish were heterozygous or homozygous for the casper mutation which results in hypopigmentation and they were additionally treated with1-phenyl-2-thiourea (200 μM final concentration; Sigma) from 10 hours post fertilization (hpf) to reduce pigmentation further. All animal procedures were performed in accordance with the Animal Act 1986 and the UK Home Office guidelines and with the approval of the University of Sussex local ethical committee.

Molecular Biology

The zebrafish ribeye a (ctbp2) promoter was used to drive expression of iGluSnFR in all neurons with ribbon synapses (17). Tg(−1.8ctbp2:Gal4VP16_BH) fish that drive the expression of the transcriptional activator protein Gal4VP16 were generated by co-injection of I-SceI meganuclease and endofree purified plasmid into wild-type zebrafish with a mixed genetic background. A myocardium-specific promoter that drives the expression of mCherry protein was additionally cloned into the plasmid to allow for phenotypical screening of founder fish. Tg(10xUAS:iGluSnFR_MH) fish driving the expression of the glutamate sensor iGluSnFR[22] under the regulatory control of the 10 x UAS enhancer elements were generated by co-injection of purified plasmid and tol2 transposase RNA into offspring of AB wildtype fish outcrossed to casper wildtype fish. The sequences for the myocardium-specific promoter driving the expression of enhanced green fluorescent protein (mossy heart) were added to the plasmid to facilitate the screening process. Crossing the stable lines of Tg(−1.8ctbp2:Gal4VP16_BH) with Tg(10xUAS:iGluSnFR_MH) results in the targeted expression of the transcriptional activator protein Gal4VP16 that subsequently binds to the UAS elements and switches on the expression of the sensor.

Plasmids for the transgenesis protocol were generated via the Gibson Assembly Cloning method. The table shows the plasmid and primer information, the underlined nucleotide sequences show the part of the primer that anneals to the template during the polymerase chain reaction.

Plasmid	Amplicon	5'-Forward Primer-3'	5'-Reverse Primer-3'
Tol2 pDest 10 x UASiGluSnFR, kind gift from Michael Orger Lab	Whole plasmid	TCCTGTGGTGTCTGAAACACCTGTGCTGCTCGCAGCTGCTGA	GTTAGGGATAACAGGGTAATTCCAAAATCAGCCACAGGATCAAGAGCA
14 x UASiGluSnFR _Mossy Heart	Cmcl2-eGFP-SV40	ATCCTGTGGCTGATTTTGGAATTACCCTGTTATCCCTAACGCC	AGCAGCTGCGAGCAGCACAGGTGTTTCAGACACCACAGGAA
I-SceI pBKS Ribeye SyGCaMP6.10.500 _Bleeding heart	Whole plasmid without SyGCaMP6 cds	GGCGCTCTGGATATGTAGCGGCGGCCGCGACTCTAGATCA	CTTTCAGGAGGCTTGCTTCAGGTGGCTCGAGATCTGAGTCC
Brn3cGal4, kind gift from Martin Meyer Lab	Gal4 VP16	GACTCAGATCTCGAGCCACCTGAAGCAAGCCTCCTGAAAGATGAAG	TGATCTAGAGTCGCGGCCGCCGCTACATATCCAGAGCGCC

Multiphoton Imaging *In Vivo*

Zebrafish larvae (7–9 days post-fertilization) were immobilized in 3 % low melting point
agarose (Biogene) in E2 medium on a glass coverslip (0 thickness) and mounted in a
chamber where they were superfused with E2, as described previously[14]. Imaging of bipolar
cell terminals was carried out using a two-photon microscope (Scientifica) equipped with a
mode-locked titanium-sapphire laser (Chameleon, Coherent) tuned to 915 nm and an
Olympus XLUMPlanFl 20x water immersion objective (NA 0.95). To prevent eye
movements, the ocular muscles were paralyzed by injection of 1 nL of α-bungarotoxin
(2 mg/mL) behind the eye. The signal-to-noise ratio for imaging was optimized by collecting
photons through both the objective and a sub-stage oil condenser (Olympus, NA 1.4).
Emission was filtered through GFP filters (HQ 535/50, Chroma Technology) before detection
with GaAsP photomultipliers (H7422P-40, Hamamatsu). The signal from each detector
passed through a current-to-voltage converter and then the two signals were added by a
summing amplifier before digitization. Scanning and image acquisition were controlled under

ScanImage v.3.6 software[44]. Images were typically acquired at 10 Hz (128 × 100 pixels per
frame, 1 ms per line) while linescans were acquired at 1 kHz.

Full-field light stimuli were generated by an amber LED (lmax = 590 nm, Thorlabs),
filtered through a 590/10 nm BP filter (Thorlabs), and delivered through a light guide placed
close to the eye of the fish. Stimuli were normally delivered as modulations around a mean
intensity of ~165 nW/mm^2 and intensity was modulated through Igor Pro software
(Wavemetrics) running mafPC (courtesy of M. A. Xu-Friedman) through an ITC-18 DAC
interface (HEKA). The multiphoton microscope was synchronized to visual stimulation.

Extraction of glutamatergic events

An analysis package was created within Igor Pro (Wavemetrics) to detect and quantize
glutamatergic events from two-photon line-scans across the terminals of bipolar cells
expressing iGluSnFR. The analysis comprised of six major steps which are described in
detail in the Supplementary Information:
1.   Separation of ROIs by spatial decomposition
2.   Time series extraction by weighted averaging
3.   Baseline correction and calculation of ΔF/F
4.   Identification of events by Wiener deconvolution
5.   Extraction of events
6.   Amplitude clustering to create a time series of quantized events

Calculation of temporal jitter
In order to compute the temporal jitter of the glutamatergic events, we first calculated the
vector strength:

$$VS_q = \frac{1}{N_q}\sqrt{\left(\sum_{i=1}^{N_q} cos\left(\frac{2\pi t_{q_i}}{T}\right)\right)^2 + \left(\sum_{i=1}^{N_q} sin\left(\frac{2\pi t_{q_i}}{T}\right)\right)^2} \quad \#(1)$$

592   where t_{qi} is the time of the i^{th} q-quantal event, T is the stimulus period, and N_q is the total
593   number of events of composed of q-quanta. The temporal jitter can then be computed by:

$$TJ_q = \frac{\sqrt{2(1 - VS_q)}}{2\pi f} \#(2)$$

595   where f is the stimulus frequency.

The signal-to-noise ratio associated with a change in the rate of a Poisson process

598   Imagine the mean rate of a Poisson process, R, changes by a factor k. The signal, S,

599   generated by comparing two observation times Dt will be the change in the number of

600   events counted in each period (kR Dt - R Dt), and the variance in that signal will be the sum

601   of the number of events counted in each period (kR Dt + R Dt). Defining the SNR as

$$SNR = \frac{S}{\sqrt{variance}} \qquad (3)$$

603   yields

$$SNR = \frac{(kR\Delta t - R\Delta t)}{\sqrt{(kR\Delta t + R\Delta t)}} \qquad (4)$$

604   which can be rearranged to obtain the time period Dt required to obtain a given SNR, as

$$\Delta t = \frac{SNR}{R} \frac{\sqrt{k+1}}{k-1} \qquad (5)$$

Calculation of the Transmitter Triggered Average (TTA)

609   The TTA for an event containing a specific number of quanta (s_q) was calculated by

610   averaging the stimuli that preceded each release event of that type:

$$\vec{s_q} = \frac{1}{n_q} \sum_{i=1}^{n_q} \vec{s}(t_i) \#(6)$$

where q is the quantal event type, n_q is the number of n-quantal events within the recording,

and s(t) is the stimulus window ending at time t. Thus, for instance, a synapse that releases

up to 11 vesicles would in principle produce 11 filters, one for each quantal event type. Two

different stimuli were used to drive the cell for the TTA analysis. We used Gaussian White

Noise (GWN) in which intensities were drawn from a Gaussian distribution spanning the LED

drivers input range and updated at a specific stimulus frame rate (varied between 10 and 30

Hz). The GWN was then discretized into eight equally spaced bins to produce a stimulus

that approximates a Gaussian distribution while driving synaptic responses effectively.

Calculations based on information theory

To quantify the amount of information about a visual stimulus that is contained within the sequence of release events from an active zone we first needed to convert bipolar cell outputs into a probabilistic framework from which we could evaluate the specific information (I_2), a metric that quantifies how much information about one random variable is conveyed by the observation a specific symbol of another random variable[21]. The time series of quantal events was converted into a probability distribution by dividing into time bins of 20 ms, such that each time bin contained either zero events or one events of an integer amplitude. We then counted the number of bins containing events of amplitude 1, or 2, or 3 etc. By dividing the number of bins of each type by the total number of bins for each different stimulus, we get the conditional distribution of Q given S, $p(Q|S)$, where Q is the random variable representing the *quanta/bin* and S is the random variable representing the *stimulus contrasts* presented throughout the course of the experiment. We then compute the joint probability distribution by the chain rule for probability (given the experimentally defined uniform distribution of stimuli S):

$$p(S,Q) = p(Q|S)p(S) \#(7)$$

In order to convert this distribution into the conditional distribution of S given Q, we use the definition of the conditional distribution:

$$p(S|Q) = \frac{p(S,Q)}{p(Q)} \#(8)$$

From these distributions we can now compute the specific information as the difference between the entropy of the stimulus S minus the conditional entropy of the stimulus given the observed symbol in the response q:

$$I_2(S,q) = H(S) - H(S|q) \#(9)$$

$$I_2(S,q) = -\sum_{s \in S} p(s) \log p(s) + \sum_{s \in S} p(s|q) \log p(s|q) \qquad (10)$$

representing the amount of information observing each quantal event type q ϵ Q carries about the stimulus distribution S.

1   Rieke, F., Bialek, W., Warland, D. & Ruyter van Steveninck, R. *Spikes: Exploring the*
Neural Code. (The MIT Press, 1999).
2   Dayan, P. & Abbott, L. R. *Theoretical Neuroscience: Computational and*
Mathematical Modeling of Neural Systems. (The MIT Press 2005).
3   Abbott, L. F. & Regehr, W. G. Synaptic computation. *Nature* **431**, 796-803, (2004).
4   Del Castillo, J. & Katz, B. Quantal components of the end-plate potential. *The*
Journal of Physiology **124**, 560-573 (1954).
5   Del Castillo, J. & Katz, B. Statistical factors involved in neuromuscular facilitation and
depression. *The Journal of Physiology* **124**, 574-585 (1954).
6   Katz, B. *Nerve, Muscle and Synapse.* (McGraw Hill, New York, 1966).
7   Choi, S. Y. *et al.* Encoding light intensity by the cone photoreceptor synapse. *Neuron*
48, 555-562, (2005).
8   Jackman, S. L. *et al.* Role of the synaptic ribbon in transmitting the cone light
response. *Nature neuroscience* **12**, 303-310, (2009).
9   Malagon, G., Miki, T., Llano, I., Neher, E. & Marty, A. Counting Vesicular Release
Events Reveals Binomial Release Statistics at Single Glutamatergic Synapses. *The*
Journal of Neuroscience **36**, 4010-4025, (2016).
10   Masland, Richard H. The Neuronal Organization of the Retina. *Neuron* **76**, 266-280,
(2012).
11   Matthews, G. & Fuchs, P. The diverse roles of ribbon synapses in sensory
neurotransmission. *Nat Rev Neurosci* **11**, 812-822, (2010).
12   Lagnado, L. & Schmitz, F. Ribbon Synapses and Visual Processing in the Retina.
Annual Review of Vision Science **1**, 235-262, (2015).
13   Freed, M. A. Quantal encoding of information in a retinal ganglion cell. *Journal of*
Neurophysiology **94**, 1048-1056, (2005).
14   Odermatt, B., Nikolaev, A. & Lagnado, L. Encoding of luminance and contrast by
linear and nonlinear synapses in the retina. *Neuron* **73**, 758-773, (2012).
15   Sterling, P. & Laughlin, S. B. *Principles of Neural Design.* (MIT Press, 2015).
16   Glowatzki, E. & Fuchs, P. A. Transmitter release at the hair cell ribbon synapse.
Nature neuroscience **5**, 147-154, (2002).
17   Singer, J. H., Lassova, L., Vardi, N. & Diamond, J. S. Coordinated multivesicular
release at a mammalian ribbon synapse. *Nature Neuroscience* **7**, 826-833, (2004).
18   Li, G. L., Keen, E., Andor-Ardo, D., Hudspeth, A. J. & von Gersdorff, H. The unitary
event underlying multiquantal EPSCs at a hair cell's ribbon synapse. *The Journal of*
Neuroscience **29**, 7558-7568, (2009).

19   Mehta, B., Snellman, J., Chen, S., Li, W. & Zenisek, D. Synaptic ribbons influence
the size and frequency of miniature-like evoked postsynaptic currents. *Neuron* **77**,
516-527, doi:10.1016/j.neuron.2012.11.024 (2013).

20   Li, G. L., Cho, S. & von Gersdorff, H. Phase-locking precision is enhanced by
multiquantal release at an auditory hair cell ribbon synapse. *Neuron* **83**, 1404-1417,
(2014).

21   DeWeese, M. R. & Meister, M. How to measure the information gained from one
symbol. *Network: Computation in Neural Systems* **10**, 325-340 (1999).

22   Marvin, J. S. *et al.* An optimized fluorescent probe for visualizing glutamate
neurotransmission. *Nature Methods* **10**, 162-170, doi:10.1038/nmeth.2333 (2013).

23   Meister, M., Pine, J. & Baylor, D. A. Multi-neuronal signals from the retina: acquisition
and analysis. *Journal of Neuroscience Methods* **51**, 95-106 (1994).

24   Meister, M. & Berry, M. J., 2nd. The neural code of the retina. *Neuron* **22**, 435-450
(1999).

25   Srinivasan, M. V., Laughlin, S. B. & Dubs, A. Predictive coding: a fresh view of
inhibition in the retina. *Proceedings of the Royal Society of London. Series B,*
Biological sciences **216**, 427-459 (1982).

26   Berry, M. J., Warland, D. K. & Meister, M. The structure and precision of retinal spike
trains. *Proceedings of the National Academy of Sciences of the United States of*
America **94**, 5411-5416 (1997).

27   Baden, T., Esposti, F., Nikolaev, A. & Lagnado, L. Spikes in retinal bipolar cells
phase-lock to visual stimuli with millisecond precision. *Current Biology* **21**, 1859-1869
(2011).

28   Laughlin, S. B., de Ruyter van Steveninck, R. R. & Anderson, J. C. The metabolic
cost of neural information. *Nature Neuroscience* **1**, 36-41, doi:10.1038/236 (1998).

29   Attwell, D. & Laughlin, S. B. An energy budget for signaling in the grey matter of the
brain. *J Cereb Blood Flow Metab* **21**, 1133-1145, (2001).

30   Barlow, H. B. Possible principles underlying the transformations of sensory
messages. In "Sensory communication", pp. 217-234. Rosenblith W.A. (Ed.). MIT
Press, Cambridge, MA (1961).

31   Niven, J. E. & Laughlin, S. B. Energy limitation as a selective pressure on the
evolution of sensory systems. *J Exp Biol* **211**, 1792-1804, (2008).

32   Strong, S. P., Koberle, R., de Ruyter van Steveninck, R. R. & Bialek, W. Entropy and
Information in Neural Spike Trains. *Physical Review Letters* **80**, 197-200 (1998).

33   Koch, K. *et al.* How Much the Eye Tells the Brain. *Current Biology* **16**, 1428-1434,
(2006).

34   Zador, A. Impact of Synaptic Unreliability on the Information Transmitted by Spiking
Neurons. *Journal of Neurophysiology* **79**, 1219-1229, (1998).

35   Magee, J. C. Dendritic integration of excitatory synaptic input. *Nat Rev Neurosci* **1**,
181-190, (2000).

36   Burrone, J., Neves, G., Gomis, A., Cooke, A. & Lagnado, L. Endogenous calcium
buffers regulate fast exocytosis in the synaptic terminal of retinal bipolar cells.
Neuron **33**, 101-112 (2002).

37   Holt, M., Cooke, A., Neef, A. & Lagnado, L. High mobility of vesicles supports
continuous exocytosis at a ribbon synapse. *Current Biology,* **14**, 173-183, (2004).

38   Matthews, G. & Sterling, P. Evidence that vesicles undergo compound fusion on the
synaptic ribbon. *The Journal of Neuroscience* **28**, 5403-5411, (2008).

39   Chapochnikov, N. M. *et al.* Uniquantal release through a dynamic fusion pore is a
candidate mechanism of hair cell exocytosis. *Neuron* **83**, 1389-1403, (2014).

40   Llobet, A., Beaumont, V. & Lagnado, L. Real-time measurement of exocytosis and
endocytosis using interference of light. *Neuron* **40**, 1075-1086 (2003).

41   Zenisek, D., Davila, V., Wan, L. & Almers, W. Imaging calcium entry sites and ribbon
structures in two presynaptic cells. *The Journal of neuroscience : the official journal*
of the Society for Neuroscience **23**, 2538-2548 (2003).

42   Lagnado, L., Gomis, A. & Job, C. Continuous vesicle cycling in the synaptic terminal
of retinal bipolar cells. *Neuron* **17**, 957-967 (1996).

43   Neves, G. & Lagnado, L. The kinetics of exocytosis and endocytosis in the synaptic
terminal of goldfish retinal bipolar cells. *The Journal of Physiology* **515 (Pt 1)**, 181-
202 (1999).

44   Pologruto, T. A., Sabatini, B. L. & Svoboda, K. ScanImage: flexible software for
operating laser scanning microscopes. *Biomed Eng Online* **2**, 13 (2003).

745
746
747

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

a

b

Figure 6

a

ΔF/F=2

20% contrast

Sequence 1

Sequence 2

5 s

b

Specific information per event (bits)

Q_e, Number of quanta in event

c

Specific information per vesicle (normalized)

Q_e, Number of quanta in event