Bacteria associated with jellyfish during bloom and post-bloom periods

Maja Kos Kramar[1], Tinkara Tinta[1], Davor Lučić[2], Alenka Malej[1] and Valentina Turk[1*]

[1]Marine Biology Station Piran, National Institute of Biology, Piran, Slovenia

[2]Institute for Marine and Coastal Research, University of Dubrovnik, Dubrovnik, Croatia

 * Corresponding author

E-mail: Valentina.Turk@nib.si

Abstract

This study is the first to investigate bacterial community associated with live medusa *Aurelia sp.* in the Gulf of Trieste (northern Adriatic Sea) using both culture independent and culture-based methods. We have analysed bacterial community composition of different body parts of medusa: exumbrella surface, oral arms ('outer' body parts) and of gastric cavity ('inner' body part) and investigated possible differences in medusa associated bacterial community structure at the time of jellyfish population peak and during senescent phase at the end of bloom, when jellyfish start to decay. Based on 16S rRNA clone libraries and denaturing gradient gel electrophoresis (DGGE) analysis, we demonstrated significant difference between bacterial community associated with *Aurelia* and the ambient seawater bacterial assemblage. Comparing bacterial community composition between different *Aurelia* medusa body parts, communities differed significantly, especially the one within the gastral cavity. The pronounced difference is dominance of *Betaproteobacteria* (*Burkholderia*, *Cupriavidus* and *Achromobacter*) in gastral cavity of medusa and *Alpha-* (*Phaeobacter, Ruegeria*) and *Gamma- proteobacteria* (*Stenotrophomonas*, *Alteromonas*, *Pseudoalteromonas* and *Vibrio*) on 'outer' body parts. This suggests that body-part specific bacterial association might have an important functional roles for the host. The results of bacterial isolates showed the dominance of *Gammaproeteobacteria*, especially *Vibrio* and *Pseudoalteromonas* in all body parts. Finally, comparison of medusa associated bacterial community structure, at the time of jellyfish population peak and during senescent phase at the end of bloom showed increased abundance of *Gammaproteobacteria*, especially *Vibrio*. Our results suggest members of *Vibrio* group are possible commensal opportunistic visitors, later becoming consumer of moribund jellyfish biomass and that the structure of jellyfish bacterial community might be affected by anthropogenic pollution in the marine environment.

Introduction

Surfaces of marine animals were found to be a unique habitat for colonization by

microorganisms, and the microbial communities associated with living surfaces showed a

pronounced variety [1]. Till recently, studies focused on the colonization of benthic organisms

such as sponges [2–6], bryozoans [7], and cnidarians, within which are included mainly corals

[8–13]. Recent studies of bacteria colonizing crustacean surfaces in the marine pelagic

environment, showed considerable dissimilarities with bacterial communities in the

surrounding seawater [14–16]. Recently, associated bacteria were reported for gelatinous

plankton such as appendicularians [17], ctenophores [18–21], and also cnidarian jellyfish

[18,22–28]. Moreover, several studies investigated the role of microbes during jellyfish

blooms, and demonstrated high bacterial growth, changes in bacterial community structure in

the surroundings of live or decaying jellyfish, and subsequently consequences in altering

trophic interactions with higher trophic levels and implications for the carbon, nitrogen, and

phosphorus cycles [29–37].

However, very few studies focused on microbial associations with scyphozoan jellyfish

during their life span. Studies show a presence of endobiotic bacteria in jellyfish tentacles [28]

and suggest that jellyfish could be vectors of bacterial pathogens and implicated in infections

of farmed salmons [24,25]. Cleary *et al.* [22] presented data on the bacterial community

composition associated with scyphozoan *Mastigias* cf. *papua etpisoni* and box jellyfish

Tripedalia cf. cystophora, while Weiland-Bräuer *et al.* [27] and Daley *et al.* [18] focused on

Aurelia aurita s.l. bacterial associates. These studies showed a diverse and specific associated

bacterial community, which in composition differs among different marine

ecosystems/different jellyfish populations, and has little similarity to the surrounding

seawater.

Furthermore, Weiland-Bräuer *et al.* [27] showed that *Aurelia aurita* harbours a different

bacterial community on its outer, mucus-covered surface of the exumbrella and gastral cavity,

and that microbial community composition differs at different life stages, especially between

benthic (polyps and strobila) and sequential planktonic life stages (ephira and juvenile and

adult medusa). In addition, an intracellular *Mycoplasma* strain, a possible endosymbiont, has

been detected. Studying microbiomes in the gastral cavity of *Cotylorhiza tuberculata,*

Mycoplasma-like bacteria was one of four bacterial taxa composing a community of reduced

diversity [24]. Most of the bacteria were suggested to have an intracellular lifestyle and

established a cooperative relationship with their host. In addition, a new candidate bacterial

taxa were proposed [23,26].

Bacterial colonization of given surface is determined by the availability of nutrients, host

immune responses, and competition between bacteria from the surrounding environment for

attachment space [38]. The epidermis and gastrodermis of jellyfish, including *A. aurita*,

contain numerous types of unicellular mucus producing gland cells, leading to the formation

of thin, constantly renewing mucus layers over external medusa [39,40]. Under certain

conditions like stress, during reproduction and digestion, and also when dying, the amount of

released mucus is even more pronounced ([40] and the references within). Mucus on jellyfish

surfaces was also found to have a role in surface cleaning and defense against predators.

Shanks and Graham [41] characterized mucus secretion as an important chemical defense

mechanism, since it contained toxins and discharged and undischarged nematocysts. The

contribution to jellyfish chemical defense is, besides mucus, the production of toxins or

antimicrobial compounds, such as isolated antibacterial peptide aurelin from mesoglea of

Aurelia aurita [42].

Further, secreted mucus, whether still covering the surface of jellyfish or already dissociated

in forms of blobs, is an attractive niche for bacteria. Since jellyfish mucus is composed mainly

of proteins, lipids, and lower percentage of carbohydrates [43], it is a high quality energy

source which is readily utilized by bacteria, especially those with a competitive advantage and

specialized for settling from surrounding seawater. This indicates that jellyfish as a host can

actively or passively affect/select bacterial associates. In addition, bacterial community

structure can be also influenced by bacterium-bacterium antagonism, as seen on particles [44],

and by environmental conditions determining the presence of metabolically active bacteria

and physiological responses of the host [45]. Whether bacteria directly adhere to external cell

layers of jellyfish, or are only associated in the thin mucus layer is not clear, however all of

the above indicates that the association of bacteria with jellyfish is highly dynamic and

complex.

Aurelia are among the most widespread scyphozoan medusae that form large aggregations in

coastal areas, fjords, and estuaries, and other semi-enclosed or enclosed systems. Moon jelly,

Aurelia sp. 8 [46] recently designated as *A. solida* [47], is also a very common jellyfish in the

northern Adriatic Sea, where 200 years of data show the stabilization of its massive

reoccurrence after 2002 [48]. Medusae are generally present from February until late June

[49], with peak abundance in the spring [48,50]. The jellyfish outbreaks worldwide seem to

become more frequent and last longer in recent years [51]. Whether this is just a rising phase

of a natural pattern of decadal oscillations or a true increase of gelatinous zooplankton blooms

is still unclear [52]. Still, some data show more frequent and abundant jellyfish aggregations

in some coastal areas around the world [48,53], causing numerous socio-economic and

ecological problems [54,55]. It has been hypothesized that jellyfish benefit from human-

caused changes in environment such as climate change, overfishing, eutrophication, habitat

modification, and species introductions [56–59].

This study is the first to investigate the associations of bacteria with live moon jellyfish in the

Gulf of Trieste (northern Adriatic Sea) using both culture-independent and culture-based

methods. Our hypotheses were the following: (i) the bacterial community associated with

medusa is specific and significantly different from the ambient bacterial population in the

environment; (ii) the bacterial community composition of different body parts of medusa, i.e.

exumbrella surface, oral arms, and of gastral cavity vary; and (iii) medusa-associated bacterial

community structure at the time of jellyfish population peak and during senescent phase at the

end of bloom, when jellyfish start to decay, differ.

Materials and Methods

Sampling site and sampling

The Gulf of Trieste is the northernmost part of the Adriatic Sea. It is characterized by a

shallow water column, with salinity and temperature variations, and strong seasonal

stratification in late summer [60]. In such an environment, *Aurelia* populations show clear

seasonality with late-autumn-early winter recruitment of ephyrae from attached polyps, spring

medusa growth, and their decay at high early summer temperatures [59].

Sampling of *Aurelia* medusae was performed during the warmer part of the year in the

beginning of May and late June 2011, at the time of adult medusa biomass accumulation.

While in May, at the time of population peak, jellyfish were viable and swimming actively, in

June, at the end of blooming period, sampled jellyfish were already in the senescent phase and

started to decay. Jellyfish were sampled individually by divers or from a boat with a sample

bucket. Each individual was placed in plastic bag with some seawater and was transported to

the laboratory. Before further analysis, each jellyfish was measured and rinsed twice with

sterile seawater (0.2 µm pre-filtered and autoclaved). For determination of the total bacterial

community associated with *Aurelia*, samples of exumbrella and oral arms of about 8 cm^2 in

size, were cut out with a sterile razor blade and stored at -80 °C. At the same time, mucus

from gastral cavity was sampled with a sterile syringe and stored under the same conditions.

At the same time of medusa sampling, ambient seawater samples were collected with a Niskin

sampler (V= 5L) at 5 m depth at the oceanographic buoy Vida (45° 32' 55. 68" N, 13°33'

1.89" E), where most of jellyfish were restrained at the time of sampling. Each time before

sampling standard physical properties, including seawater temperature, salinity and oxygen

concentration were measured with a CTD fine-scale probe (Microstructure Profiler MSS90,

Sea & Sun Technology GmbH).

Bacterial isolates from jellyfish and seawater samples

Viable bacterial cells from the surfaces of jellyfish and seawater samples were determined

with the spread plate method on modified ZoBell marine agar media [61]. For jellyfish

exumbrella, the whole exumbrella surface was inoculated on the plate to create jellyfish

imprints of exumbrella-associated bacteria, while the gastro vascular cavity was scraped with

a sterile cotton swab and spread evenly over the surface of agar plates. For seawater samples,

100 µL was evenly spread on an agar plate. Inoculated plates were incubated in the dark at an

in situ temperature for 21 days. For each plate, the number of colony forming units (CFU)

was determined and distinctive morphological types of colonies were described. Since direct

prints of jellyfish exumbrella were created on agar plates, we estimated the abundance of

bacteria as CFU/cm^2. For DNA extraction, individual colonies were aseptically picked and

streaked onto fresh agar plate until single colonies were obtained. A single colony of each

bacterial isolate was inoculated in modified liquid ZoBell media and incubated in the dark at

the *in situ* temperature until growth was observed (increased turbidity). Part of the liquid

cultures was stored in 30% glycerol (final concentration) at – 80 °C for bacterial culture

collection. Data on the seawater cultivable bacterial community from the Gulf of Trieste were

taken from the dataset gathered during two-year sampling campaign of cultivable bacterial

community, of which the sampling time and location coincided with the sampling time of

jellyfish (May and June 2011) (Acc. No. KC307273- KC307520). Also, for this dataset

sampling was performed at 5 m depth and the bacterial culture collection was obtained as

described above.

DNA extraction and PCR

Bacterial DNA was extracted from a liquid culture with a modified Chelex-based procedure

[62], or with a commercial kit (NucleoSpin Tissue, Macherey - Nagel) according to

manufacturer's protocol. Bacterial cells were harvested by centrifugation and washed twice

with 1x PBS buffer (stock 10x; 1.4 M NaCl, 27 mM KCl, 101 mM Na_2HPO_4, 18 mM

KH_2PO). When using Chelex (BioRad), the washed cells were re-suspended in 200 µl of 5%

Chelex solution. The suspension was incubated for 15 min at 99 °C and transferred at 4 °C for

10 min., and was centrifuged at 4000 rpm for 10 min. The water phase with dissolved DNA

was transferred to a new tube and stored at -20 °C until further downstream applications.

Bacterial 16S rRNA genes were amplified using universal bacterial primers 27F and 1492R.

The PCR reaction mix (50 µl) contained 2 µl (50-100 ng) of extracted genomic DNA, 1x

reaction buffer (TrisKCl-MgCl$_2$), 2 mM MgCl$_2$, 0.2 mM dNTP, 1 µM of each primer, and Taq

polymerase (5U/µl, Fermentas). The PCR temperature cycling conditions were as follows:

initial denaturation at 94°C for 2 min.; 30 cycles of denaturation at 94°C for 1 min., annealing

at 55°C for 2 min., and elongation at 72°C for 2 min. The final cycle was followed by

extension at 72°C for 5 min. The quality and size of PCR products was confirmed by agarose

gel electrophoresis (1% agarose (Sigma) in 1x TAE buffer) with etidium bromide (10 mg/ml)

and visualized using an UV transilluminator (BioDocAnalyze Gel documentation system,

Biometra). The bacterial 16S rRNA genes were partially sequenced with 27F primer at

Macrogen Inc.

Total bacterial community composition

Jellyfish-associated bacterial community DNA extraction

Exumbrella and oral arms samples were thawed down. Bacterial DNA was extracted with

CTAB (cetyl- trimethyl-ammonium bromide) as described before [21] with slight

modification. Samples were placed into a tube containing 2% CTAB solution (1.4 M NaCl,

100 mM Tris_Cl pH= 8, 2% CTAB, 20 mM EDTA pH=8, 0.2% β- mercaptoethanol), and

were incubated at 65°C for 1 h. Afterwards, a STE buffer (6.7% sucrose, 50 mM Tris-Cl pH=

8, 1 mM EDTA pH= 8) and proteinase K (100 µg/ml final concentration, Sigma) were added

and the mixture was incubated at 55°C overnight. Following chloroform-isoamylacohol (24:1,

v/v, Sigma) and phenol- chloroform- isoamylacohol (25:24:1, v/v, Sigma) purification steps,

DNA was precipitated at - 20°C overnight with isopropanol. The pellet was washed with 70%

ice-cold ethanol and dried in a speed vac. The precipitated DNA was re-suspended in 0.22 µm

pre-filtered, autoclaved 1X TE buffer, and kept at – 20°C.

Seawater's total bacterial community DNA extraction

Seawater samples were filtered onto 0.2 µm polyethersulfone membrane filters (47 mm

diameter, PALL Inc.), which were stored at - 80°C. DNA was extracted from the filters (one

quarter per sample) as described in Böstrom *et al.* [63], with slight modifications. DNA was

precipitated at - 20°C for 1 h, with 0.1 volume of sodium acetate (3 M NaAc, pH= 5.2) and

0.6 volume of isopropanol. The pellet was washed with 70% ice-cold ethanol and dried in a

speed vac. Precipitated DNA was re-suspended in 0.22 µm, pre-filtered, autoclaved TE

buffer, and kept at - 20°C.

Denaturing Gradient Gel Electrophoresis

For DGGE analysis of jellyfish-associated and seawater's total bacterial community, the

bacterial 16S rRNA genes were amplified using a universal primer set, 341F with a 40 bp GC-

clamp and 907R as described before [64,65]. The PCR reaction mix with a final volume 50 µl

contained 2 µl of extracted DNA (50–100 ng), 1x reaction buffer (Tris KCl-MgCl$_2$,

Fermentas), 1.5 mM MgCl$_2$ (Fermentas), 0.2 mM dNTP (Fermentas), 0.5 µM of each primer

(Sigma), 0.38 µg/ml BSA (Fermentas), and Taq polymerase (5 U/µl, Fermentas). The PCR

touchdown protocol according to Don *et al.* [66] was used: with initial denaturation at 94°C

for 5 min., followed by 10 touchdown cycles and 20 standard cycles: denaturation for 1 min.

at 94°C, primer annealing for 1 min. at 55°C, and primer extension for 3 min. at 72°C. The

last cycle was followed by 2 min. incubation at primer extension temperature of 72°C.

When we were unable to obtain a sufficient quantity of PCR products from jellyfish samples,

we used a two-step nested PCR- DGGE strategy [67], modified to analyze the marine

bacterial community. Bacterial 16S rRNA genes were first amplified with universal primer

set, 27F and 1492R. The PCR reaction mix, with a final volume 50 µl, contained 2 µl of

extracted DNA (50-100 ng), and was prepared the same as described above in this section.

The PCR temperature cycling conditions were as follows: initial denaturation for 2 min. at

94°C, followed by 25 standard cycles: denaturation at 94°C for 1 min., primer annealing for 1

min. at 50°C, and primer extension at 72°C for 1 min. The last cycle was followed by 5 min.

incubation at the primer extension temperature of 72°C. Second, nested amplification was

performed using a DGGE primer set, PCR mixture, and a touchdown annealing protocol, as

described above in this section. The quality and size of PCR products were tested by agarose

gel electrophoresis. PCR products were analyzed by DGGE electrophoresis, as previously

described in [33].

Distinct bands were excised from the gel and placed in 100 µl of sterile Sigma water

overnight to elute DNA. The eluted DNA was re-amplified using primer set 341F and 907R

and the same reaction mix (a final volume 50 µl) with 2 µl of eluted DNA, as described above

in the first paragraph in this section. The cycling protocol used was the same as to amplify the

DNA of bacterial isolates (see section on Bacterial isolates DNA extraction and PCR). The

bacterial 16S rRNA genes were partially sequenced with 341F primer at Macrogen Inc.

Bacterial 16S rRNA gene clone libraries

For jellyfish and seawater samples clone libraries construction, bacterial 16S rRNA gens were

amplified using the same DNA as for DGGE and universal primer set, 27F and 1492R, as

described before [33]. The PCR reaction mix with a final volume 50 µl contained 2 µl of

extracted DNA (50-100 ng), and was prepared as described above (see section on DGGE, first

paragraph). For samples with low DNA concentration (extracted from jellyfish samples), a

nested PCR-libraries approach was used [68], and modified to analyze marine bacterial

community. Again, bacterial 16S rRNA gene was first amplified with a universal primer set,

27F and 1492R using same protocol and reaction mix as in the first amplification step of the

nested PCR-DGGE strategy (see the section on DGGE, second paragraph). Second, nested

amplification was performed using primers 341F and 907R. The PCR reaction mixture and

cycling protocol were the same as used for clone library construction (described above in this

section [33]). The PCR products were immediately ligated into a commercially available pCR

2.1 vector and transformed into competent *E. coli* TOP 10 cells using a commercially

available TA Cloning Kit (Invitrogen), and according to the manufacturer's protocol. The

plasmid inserts from of each clone library were partially sequenced using M13F primer or

27F primer at Macrogen Inc.

Sequence analyses

Raw sequence data recovered from bacterial isolates and clone libraries were passed through

the DNA Baser program (www.DNAbaser.com) to remove traces of sequencing primer, and

to trim away ambiguous bases at the end of a sequence. The clone libraries sequences were

also checked for vector contamination and were analyzed with the program Bellerophon

(https://greengenes.lbl.gov/) to detect chimeric sequences, which were removed. Additionally,

Mothur software [69] was used to further reduce poor quality sequence data. Sequence

taxonomic identities (with ≥ 97%) of bacterial isolates and sequences recovered from clone

libraries and DGGE bands were assigned using the BLAST (Basic Local Alignment Tool)

algorithm available at NCBI (National Centre for Biotechnology Information). Around 50%

of the sequences recovered from clone libraries and DGGE bands exhibited <97% similarity

to previously published GenBank entries (omitted from further analysis). Classification of

bacterial isolates was done down to the genus level, and of clones and DGGE bands down to

the family level. The number (N) of high quality sequences, with $\geq$ 97% similarity to the

nearest GenBank entry, and the number of distinct bacterial genus and families (S_{obs}) is

presented in Supporting Information (S1 Table, S2 Table). The contribution of distinct

bacterial genus or families was expressed as a percentage of the total number of sequences in

each sample or library (relative abundance) (S1 Table, S2 Table). Chloroplast sequences were

omitted from further analysis.

Nucleotide sequence accession numbers

The 16S rRNA gene sequences, for all bacterial isolates, clone libraries, and DGGE bands,

obtained in this study have been deposited in the GenBank (NCBI) under following accession

numbers: from KF816449 to KF816471, and KF816480 to KF816592 for bacterial isolates

(Supporting information, Table S5), from KF816761 to KF816832, from KF817469 to

KF817519, from MF952738 to MF952748, and from MF952764 to MF952865 for sequences

obtained from clone libraries, and from MF952749 to MF952763 for sequences obtained from

DGGE bands.

Diversity indices and statistical analyses

To compare the diversity of viable bacterial isolates from jellyfish and surrounding seawater,

ecological diversity indices were calculated for each sample: the number of different bacterial

genus (species richness (Sobs)), Shannon diversity index (H'), Margalef's index (d), Pielou's

evenness index (J') and Chao-1 index. The same parameters were calculated for 16S rRNA

bacterial clone libraries at the family level. Additionally, in order to estimate how well the

actual species composition was captured, for each clone library a coverage value was

calculated as $C = 1 - n_1/N$, where n1 is the number of phylotypes appearing only once in the

library, and N is the library size [70].

Non-metric multi-dimensional scaling (nMDS) plots were used to determine the similarities

between DGGE banding patterns. For this purpose, a similarity matrix was calculated (using

Jaccard resemblance measure) based on the presence/absence matrix of align bands. Analysis

of similarity (ANOSIM) was used to verify the significance of similarity among bacterial

communities, as indicated by nMDS, by testing the hypothesis that bacterial communities

from the same cluster are more similar to each other than to communities in different clusters.

Cluster analysis was used to determine scaled similarities between 16S rRNA gene clone

libraries (total bacterial communities) and between bacterial isolates (culturable bacterial

communities). For cluster analysis of 16S rRNA gene clone libraries, a Bray-Curtis similarity

matrix was constructed from arcsine-transformed relative abundances of distinct bacterial

families in each clone library. For bacterial isolates, a Bray-Curtis similarity matrix was

constructed from untransformed relative abundances of distinct bacterial genus in each

culturable bacterial community. Based on the similarity matrix, dendrogram was produced

with group-average linkage algorithm. The similarity profile test (SIMPROF) was used to

define statistically significant clusters in samples.

To examine the difference between communities associated to different jellyfish body parts

and seawater, one-way ANOSIM statistic with 999 permutations, based on Bray-Curtis

similarity matrix, was made. Samples were grouped according to isolation source

(communities of jellyfish exumbrella (AK), jellyfish oral arms (AR) and jellyfish gastral

cavity (AG) and communities of seawater (W)). Similarly, one- way ANOSIM statistic with

999 permutations was made to examine the difference between communities associated with

jellyfish at the time of population peak and at the end of blooming period. Samples were

grouped according to time (communities associated to jellyfish sampled at time of population

peak (May) and those sampled at the end of the bloom, when jellyfish were in senescent phase

(June)). Additionally, similarities percentage (SIMPER) analysis was used to determine which

bacterial group contribute the most to the differences between communities of different body

parts of jellyfish and water communities (for culturable and total bacterial community).

Diversity indices and statistical analysis were performed using Primer v6 [71] and PAST,

version 3.9 [72].

Results

The composition of the bacterial community associated with scyphomedusae *Aurelia,* which

frequently blooms in the Northern Adriatic, was studied and compared with the community

composition from the surrounding seawater, in order to understand if the jellyfish-associated

community is specific and significantly distinct from the ambient seawater bacterial

assemblage. The composition of bacterial community associated with different jellyfish

compartments (exumbrella surface, oral arms, and in the gastral cavity) was analysed to

examine the compartment-specificity of associated bacterial consortia. In addition, we

compared the composition of the bacterial community associated with jellyfish collected

during two different time points of bloom development/progression: (i) at the peak of

population, and (ii) at the end of the blooming period/at the decay of the bloom. The bacterial

community composition/structure was determined using both culture-independent and

culture-dependent techniques.

Comparison of jellyfish-associated and ambient seawater bacterial community

composition

The phylogenetic analyses of the bacterial 16S rRNA gene clone libraries revealed significant

difference between jellyfish-associated and ambient seawater bacterial communities

(ANOSIM, global R= 0.777, p< 0.01) (Fig 1A). The bacterial communities associated with

Aurelia showed the dominance of bacterial phyla *Proteobacteria*, which consisted of

Alphaproteobacteria (up to 75%), *Gammaproteobacteria* (up to 45.5%), and

Betaproteobacteria (up to 53.5%), with different relative contributions in the individual

jellyfish sample (Fig 1B). At the family level, *Alphaproteobacteria* were dominated by

Rhodobacteraceae (mostly *Phaeobacter*, *Ruegeria*)) and *Betaproteobacteria* by

Burkholderiaceae (*Burkholderia*) (Fig 1B, S1 Table). Within *Gammaproteobacteria,* mostly

Vibrionaceae (*Vibrio*), *Pseudoalteromonadaceae* (*Pseudoalteromonas*), *Xanthomonadaceae*

(*Stenotrophomonas*), and *Pseudomonadaceae* (*Pseudomonas*) (Fig 1B, S1 Table) were

detected.

Fig 1. Bacterial 16S rRNA gene clone libraries constructed from samples of *Aurelia*

jellyfish and ambient sweater.

(**A**) Cluster analysis based on bacterial 16S rRNA gene clone libraries. AK-jellyfish

exumbrella surface, AR-jellyfish oral arms, AG-mucus from gastral cavity and W-ambient

seawater. Samples were collected in May (grey squares) and June (inverted black

triangles).The dendrogram was inferred with the group average algorithm, based on the Bray–

Curtis similarity matrix of arcsine transformed averaged abundances. Grey branches do not

differ significantly (SIMPROF test, p> 0.05). (**B**) Composition of bacterial 16S rRNA gene

clone libraries (expressed as percentage of clones) constructed from samples of jellyfish

exumbrella surface (AK1, AK2), oral arms (AR1), mucus from gastral cavity (AG1) and the

ambient seawater (W_May) sampled in May and jellyfish exumbrella surface (AK6, AK7),

oral arms (AR6) and the ambient seawater (W_Jun) sampled in June. Cumulative column

charts represent relative abundances of bacterial family and area chart in the background

represent relative abundances of major bacterial groups and *Proteobacteria* class. Taxa with

relative abundance of< 3% across all samples are subsumed under Other Bacteria.

In comparison to the jellyfish-associated bacterial community, the ambient seawater bacterial

community was more diverse and dominated by three bacterial phyla: *Proteobacteria*,

Flavobacteria, and Cyanobacteria (Fig 1B, S5 Table). *Alphaproteobacteria* (up to 38.6%)

were dominated by *Rhodobacteraceae* and SAR11, *Gammaproteobacteria* (up to 21.4%) by

Litoricolaceae and SAR86; *Flavobacteria* (up to 17.4%) by *Flavobacteriaceae* and

Cryomorphaceae, and *Cyanobacteria* (up to 15.4%) by *Synechococcus*. We also detected

Actinobacteria (10.3% in May) with the representative from the *Microbacteriaceae* family.

(Fig 1B, S1 Table). According to SIMPER analysis *Synechococcus*, SAR11 and

Flavobacteriaceae contributed the most to difference between jellyfish-associated and water

column bacterial community (S3 Table).

The bacterial community composition associated with different body parts of jellyfish

The results of 16S rRNA gene clone libraries analysis pointed on the statistically significant

differences between bacterial communities associated with different body parts of jellyfish

(exumbrella surface, oral arms, and gastral cavity) (ANOSIM, global R= 0.571, p< 0.05)(Fig

1A). The bacterial communities' composition associated with different body parts of jellyfish

sampled at the peak of population, were as follows. The bacterial communities associated

with jellyfish exumbrella were dominated by *Alphaproteobacteria* (up to 75%), followed by

Gammaproteobacteria (up to 22.2%) and *Betaproteobacteria* (up to 12.5%) (Fig 1B). The

population of *Alphaproteobacteria* was dominated by *Rhodobacteraceae*, mostly

Phaeobacter, *Ruegeria*, but also *Rhizobiaceae*, *Hyphomicrobiaceae*, and *Sphingomonadaceae*

were detected. Within *Gammaproteobacteria*, mostly *Xanthomonadaceae*

(*Stenotrophomonas*), but also *Alteromonadaceae* (*Alteromonas*), and within

Betaproteobacteria exclusively *Comamonadaceae* were detected. (Fig 1B, S1 Table). The

bacterial community of jellyfish oral arms was more diverse than the bacterial community

associated with exumbrella and gastral cavity (S5 Table). The bacterial community associated

with oral arms consisted of *Alphaproteobacteria* (50%, exclusively *Rhodobacteriaceae*) and a

higher percentage of *Gammaproteobacteria* (31.3%) composed mainly of *Vibrionaceae*

(*Vibrio*), but also *Pseudoalteromonadaceae*, *Moraxellaceae*, and *Pseudomonadaceae*.

Betaproteobacteria were detected (12.5%, only *Burkholderiaceae*), and also a small

percentage of *Actinobacteria* (6.3%) (Fig 1B, S1 Table). In contrast, the bacterial community

in the gastral cavity, was dominated by *Betaproteobacteria* (53.5%), followed by

Gammaproteobacteria (27.9%) and *Actinobacteria* (11.6%, dominated by *Micrococcaceae*).

At the family level, *Burkholderiaceae* (*Burkholderia*) and *Alcaligenaceae* dominated the

Betaproteobacteria class. The gammaproteobacterial population was almost exclusively

Pseudomonadaceae (*Pseudomonas*) (Fig 1B, S1 Table).

The bacterial community structure of different jellyfish body parts was also studied using

denaturing gradient gel electrophoresis (DGGE). Bacterial community fingerprints varied,

both within and between sample types (exumbrella, oral arms, and gastral cavity). Despite the

observed heterogeneity (S1 Fig), the DGGE-based non-metric multidimensional scaling

(nMDS) analysis showed that bacterial communities clustered according to jellyfish body part

(S1 Fig) (ANOSIM, global R= 0.633, p< 0.05). Jellyfish-associated bacterial community

composition was more similar between replicates of the samples collected from the same

body parts (40%) than different jellyfish body parts (Fig 2). Phylogenetic information

obtained from excised DGGE bands showed that bacterial taxa across all samples mostly

belonged to *Alphaproteobacteria* (*Roseobacter*, *Phaeobacter*, *Ruegeria* all

Rhodobacteraceae), but also *Gammaproteobacteria* (*Vibrio*, *Pseudoalteromonas*,

Stenotrophomonas) and *Betaproteobacteria* (*Burkholderia*) (S1 Fig, S7 Table).

Fig 2. **Non-metric multidimensional (nMDS) analysis based on bacterial community**

DGGE banding patterns of *Aurelia* jellyfish samples.

AK- jellyfish exumbrella surface, AR-jellyfish oral arms, AG- mucus from gastral cavity and

W- ambient seawater. Samples were collected in May (grey squares) and June (inverted black

triangles). Resemblance circles: grey line - 40% similarity; black line - 50% similarity.

Bacterial community structure shifts due to jellyfish population senescence

Our results show the difference between the bacterial communities associated with jellyfish

collected during the peak of the jellyfish bloom and one month later, at the jellyfish

population senescence (Fig 1). Changes in bacterial communities, due to jellyfish population

senescence were evident as the shift towards *Gammaproteobacteria*, mostly at the expense of

Betaproteobacteria, and to a lesser extent at the expense of *Alphaproteobacteria*, whose

dominance became less pronounced. In addition, diversity was lower in the bacterial

community associated with senescent jellyfish S5 Table). Bacterial communities associated

with exumbrella surface were composed of *Gammaproteobacteria* (up to 42.9%) and

Alphaproteobacteria (up to 66.7%; almost exclusively *Rhodobacteriaceae* of which mostly

Roseovarius, *Ruegeria*). Within *Gammaproteobacteria*, previously dominant

Xanthomonadaceae were replaced with *Alteromonadaceae* (*Marinobacter*) and *Vibrionaceae*

(*Vibrio*). In addition, *Pseudoalteromonadaceae* (*Pseudoalteromonas*) and *Moraxellaceae*

were detected (Fig 1B, S1 Table). A similar change/shift was evident in the bacterial

community structure determined in the sample of oral arms. *Gammaproteobacteria* were

dominant (45.5%; taxa composition similar to that associated with exumbrella surface, with

even more pronounced *Vibrio* dominance), followed by *Alphaproteobacteria* (36.4%,

exclusively *Rhodobacteraceae*) and *Betaproteobacteria* (18.2%, exclusively

Burkholderiaceae). The shift in bacterial community structure was supported by SIMPER

analysis showing that *Rhodobacteriaceae* and *Comamonadaceae* were more frequent in the

bacterial community associated with jellyfish at the peak of the bloom, and

Rhodobacteriaceae, *Vibrionaceae*, and *Alteromonadaceae* in the bacterial community

associated with senescent jellyfish (S8 Table).

Culturable bacterial community composition

Altogether, 135 bacteria were isolated from the exumbrella surface and gastral cavity of

jellyfish, sampled during the peak of jellyfish population (AK1, AK3, AK6, AG1, AG6), and

at the end of the blooming period (AK8, AK10, AK11, AG8, AG11). Regarding the

morphology of bacterial colonies, we detected nine morphotypes in May, and only three

morphotypes in June. The estimated abundance of cultural bacteria associated to jellyfish

exumbrella was 1.9 CFU/cm^2 in May, and 2.0 CFU/cm^2 in June. Identification of 16S rRNA

gene sequence revealed that across all samples, bacteria predominantly belonged to

Proteobacteria (*Gammaproteobacteria* followed by *Alphaproteobacteria and*

Betaproteobacteria), *Actinobacteria,* and *Firmicutes* (*Bacilli)* (exhibiting ≥ 97% identity to

previously described bacterial species). Analysis of bacterial isolates again showed significant

differences between seawater and jellyfish-associated communities (ANOSIM, global

R=0.393, p< 0.05) (Fig 3), with seawater communities being more diverse (Fig 4, S6 Table).

Gammaproteobacteria (mostly *Vibrio* and *Pseudoalteromonas*) dominated the exumbrella

surface-associated community (up to 100%) and the community of gastral cavity (> 60%),

while bacterial isolates obtained from ambient seawater were mainly affiliated with

Alphaproteobcateria (88% in May and 42% in June, mostly *Erythrobacter*

(*Erythrobacteraceae*) and *Brevundimonas* (*Caulobacteraceae*)), followed by

Gammaproteobacteria (32%, mostly *Halomonas* (*Halomonadaceae*), *Idiomarina*

(*Idiomarinaceae*)) and *Actinobacteria* (25%, *Brevibacterium* (*Brevibacteriaceae*)) which

were more abundant in June (Fig 4, S2 Table). According to SIMPER analysis,

Erythrobacter, *Brevibacterium*, and *Brevundimonas* contributed the most to the difference

between jellyfish-associated and water communities (S4 Table).

Fig 3. Cluster analysis based on culturable fraction of bacterial community associated

with *Aurelia* jellyfish.

AK- jellyfish exumbrella surface, AG- mucus from gastral cavity and W- ambient seawater.

Samples were collected in May (grey squares) and June (inverted black triangles). The

dendrogram was inferred with the group average algorithm, based on the Bray–Curtis

similarity matrix of arcsine transformed averaged abundances. Grey branches do not differ

significantly (SIMPROF test, p> 0.05).

Fig 4. Bacterial isolates obtained from *Aurelia* jellyfish and ambient seawater.

(**A**) Bacterial isolates obtained from jellyfish exumbrella surface (AK1, AK3, AK6), gastral

cavity (AG1, AG6) and seawater (W_May) in May in the Gulf of Trieste. (**B**) Bacterial

isolates obtained from exumbrella surface (AK8, AK10, AK11) and gastral cavity (AG8,

AG11) of jellyfish and seawater (W_Jun) in June in the Gulf of Trieste. Cumulative column

charts represent relative abundances of bacterial genus and area chart in the background

represent relative abundances of major bacterial groups and *Proteobacteria* class. Taxa with

relative abundance of< 3% across all samples are subsumed under Other Bacteria..

Differences between culturable bacterial communities of different body parts were small.

Bacterial communities associated with jellyfish exumbrella at the time of population peak

were dominated by *Gammaproteobacteria* (from 44.4% up to 100%), while

Alphaproteobacteria and *Actinobacteria* represented up to 22.2% of the community (Fig 4A).

Small percentages of isolates belonged to *Betaproteobacteria* (11%, exclusively *Delftia*

Comamonadaceae) *Bacilli* (9%; mostly *Exiguobacterium*) and *Bacteriodetes* (12.5%).

Considering the main representatives within *Gammaproteobacteria, Vibrio* (*Vibrionaceae*),

Pseudoalteromonas (*Pseudoalteromonadaceae*), and *Stenotrophomonas* (Xanthomonadaceae)

dominated , but also *Pseudomonas* (*Pseudomonadaceae*), *Alteromonas* (*Alteromonadaceae*),

and *Psyhrobacter* (*Moraxellaceae*) were detected. Representatives of *Alphaproteobacteria*

were mostly *Labrenzia* and *Phaeobacter* (*Rhodobacteraceace*), and representatives of

Actinobacteia mostly *Kocuria* (*Micrococcaceae*) and *Microbacterium* (*Microbacteriaceae*)

(Fig 4A, S2 Table). The composition of bacterial communities in jellyfish gastral cavity was

similarly dominated by *Gammaproteobacteria* (> 60%), followed by *Alphaproteobacteria* (up

to 40%) and *Betaproteobacteria* and *Bacilli* (both 10%). The population of

Gammaproteobacteria was dominated by *Pseudoalteromonas* followed by *Pseudomonas,*

Vibrio, and *Stenotrophomonas* (Fig 4A, S2 Table).

Changes in culturable bacterial community composition, presumably related to jellyfish

population decay, at the end of blooming period were significant (ANOSIM, global R= 0.362,

p< 0.05). Changes were evident as there was even more pronounced dominance of

Gammaproteobacteria (from 66.7% up to 100%) in communities associated to jellyfish

exumbralla and of gastral cavity (Fig 4B). Within the Gammaproteobacterial population,

predominant community members *Pseudoalteromonas*, *Stenotrophomonas*, and *Pseudomonas*

almost or completely 'disappeared', and *Vibrio* became highly dominant or even formed a

monoculture representative (Fig 4B, S2 Table). The change in community structure towards

Vibrio was also confirmed by SIMPER analysis (S9 Table). Consequently, the prevalence of

Vibrio also resulted in lower diversity (S6 Table).

Discussion

***Aurelia* associated bacterial community**

Comparison of jellyfish-associated and ambient seawater bacterial community composition

Our results on bacterial community composition, assessed by culture-dependent and cultural-

independent approaches, demonstrated significant differences between bacterial community

associated with *Aurelia* and the ambient seawater bacterial assemblage. Phylogenetic analysis

showed a wide diversity of bacterial community associated with jellyfish, including members

of Proteobacteria (*Alpha-*, *Gamma-*, and *Beta-proteobacteria),* which dominated the

community, and members of *Actinobacteria* and *Cyanobacteria*. Ambient seawater bacterial

communities were more diverse, and dominated by three bacterial phyla: *Proteobacteria*,

Flavobacteria, and *Cyanobacteria*. Also, within Proteobacterial groups associated with

jellyfish and the one detected within ambient seawater assemblage, had different taxonomic

affiliations, and the dominance or exclusivity of only one taxon was hardly present.

Similar observations of the jellyfish-specific bacterial community, distinct from the

community in ambient seawater, were reported previously for *A. aurita* [18,27] and also other

marine animals [45]. Since associated bacterial assemblages differed from the ambient

seawater bacterial community, and from bacteria associated with other types of

substrates/surfaces found in the water column, it was suggested that associations with animals

might be specific to some degree [45]. According to Taylor *et al*. [73] sponge bacterial

associates could be separated/split into three groups: (i) bacterial specialists - found on only

one host species; (ii) host associates - found on multiple hosts; and (iii) generalists - found on

multiple hosts and within the seawater community. In our study most bacteria associated with

Aurelia were not detected in the ambient seawater, however, they were closely related to

bacteria previously found in association with other host animals (exhibiting at least 97%

similarity), indicating that this relationship is not host-specific. Previous studies on *A. aurita*

bacterial associates also did not reveal the presence of any *Aurelia*-bacterial specialists, with

the exception of *Mycoplasma* sp. (class *Mollicutes*), a possible/hypothetical endosymbiont

[18,27]. However, in our study, we were not able to detect any *Mycoplasma* members. In

addition, bacterial community composition was very different to the community associated

with *A. aurita* from the North West Atlantic and the Baltic Sea [18,27]. This might suggest

the possible effect of host genetics background (different populations of *Aurelia* species in

geographically distant locations), and also the important's of environmental and

anthropogenic conditions, determining the presence, activity, and composition of bacterial

community in jellyfish's environment.

We found another interesting result, which is the consistently unsuccessful amplification of

bacterial 16S rRNA genes from jellyfish samples, unless an additional nested PCR reaction

was performed. Problems with DNA amplification were reported before in the analyses of the

tissue of healthy corals, and were attributed to the low abundance of bacterial associates [74],

confirming previous observation of rare isolated bacterial cells within coral tissue by in situ

hybridization [75]. This indicates that unsuccessful DNA amplification in our study also could

be the consequence of the overall low bacterial number on jellyfish exumbrella, the oral arm

surface, and in the gastral cavity. The speculations on the low number of *Aurelia* jellyfish-

associated bacteria, was also confirmed by scanning electron microscopy of the adult medusa

umbrella surface in our parallel study (data not shown). Our results show that the surface was

covered with mucus, and we observed mucus secretions in the form of flocs on all external

surfaces, but with no bacteria observed. Unlike the epidermal umbrella surface, the

examination of mucus secreted from exumbrella surface, revealed the presence of

considerable amounts of bacteria (data not shown). Using the same microscopic method,

Johnston and Rohwer [76] similarly found that external cell layers of coral are invariably

clean of adhering microbes. They did, however, suggest the possibility of a dynamic

community hovering in the boundary layers above the coral epidermis. So, the possibility that

the majority of bacteria dwell in mucus produced by medusa rather than being present within

the mesoglea or attached to epidermis is also in agreement with observations by Weiland-

Bräuer *et al*. [27], detecting the majority of bacteria located on the outer surface of coating

mucus, covering *A. aurita* polips. This could also support our results of bacterial colonies

grown after imprints of jellyfish surfaces on agar plates, where it was estimated CFU less than

two bacterial colonies/cm^2 of jellyfish surface in May and June. The presence of rare bacterial

cells could be due to the fact that adult medusa has evolved mechanisms of defense against

epibiotic organisms. One type of mechanism could be the production of antibacterial peptide

aurelin, extracted from mesoglea of *A. aurita* [42]. Based on that, we can speculate that *A.*

aurita is a 'hostile' environment for bacteria. It is also known that jellyfish surfaces, including

A. aurita, are covered by a constantly renewing mucus layer, which was found to have

implications in surface cleaning and defense against predators [39–41]. Similarly as Garren

and Azam [77] demonstrated for corals, surface cleaning by mucus production could be used

by jellyfish to regulate an abundance of bacterial associates. Even more so, during certain

conditions, including stress, the mucus production is more pronounced [41]. In our study,

extensive mucus release (from surface and gastral cavity) was detected during the processing

of jellyfish, possibly leading to bacterial loss and low bacterial numbers of jellyfish associates

in our samples.

Despite being rare, the total and the culturable part of the bacterial community associated with

Aurelia jellyfish from the Gulf of Trieste was found to be diverse. It was composed mostly of:

(i) bacteria belonging to genus *Ruegeria*, *Phaeobacter*, and *Pseudoalteromonas*, known for

their extraordinary ability to successfully compete and colonize surfaces, and also to enhance

survival chances of the host organism [78–81]; (ii) bacteria belonging to genus *Alteromonas*

and *Vibrio*, known as particle and surface colonizers with the ability to degrade and utilize a

broad spectrum of organic substrates [81]; and (iii) bacteria belonging to genus

Stenotrophomonas, *Burkholderia*, *and Achromobacter*, mostly known as medically important

strains, but also with high bioremediation potential due to ability of PAHs and xenobiotic

degradation and with broad antibiotic resistance [82–87]. With the exception of *Ruegeria*,

Burkholderia, *and Achromobacter*, other bacteria were also recovered by culturing. The

presence of *Ruegeria* and *Phaeobacter* (*Alphaproteobacteria - Roseobacter* clade) and

Pseudoalteromonas, *Alteromonas*, and *Vibrio* (*Gammaproteobacteria*) is not surprising, since

they are known as successful and dominant particle/surface colonizers [80,88,89], and

regularly associated with marine sponges [4], corals (tissue and mucus) [8,9,13,90–93], and

ctenophores [18,20,94]. A more interesting feature of *Aurelia* jellyfish associated bacterial

community was the high relative abundance of *Burkholderia* and *Achromobacter*

(*Betaproteobacteria*). This group is not characteristic of a marine environment but was found

in environments characterized with lower salinity and higher nutrient concentrations, such as

estuaries [95]. *Betaproteobacteria* were also found in association with sponges [3,5,96],

corals [9,11,12], ctenophores [19,94], and cnidarian Hydra [97], but, with the exception of the

last, their relative abundances were lower than in our study. Of the above bacterial taxa, only

Phaeobacter, *Vibrio,* and *Pseudoalteromonas* were previously found in association with *A.*

aurita [18,27,33].

The bacterial community composition associated with different body parts of jellyfish

Bacterial community composition differed significantly between different *Aurelia* medusa

body parts, especially the one within the gastral cavity. The communities of exumbrella and

oral arms shared dominant bacterial groups, *Alphaproteobacteria* followed by

Gammaproteobacteria, while the community in the gastral cavity was dominated by

Betaproteobacteria, followed by *Gammaproteobactera* and *Actinobacteria*. Within

Alphaproteobacteria, bacterial communities of the exumbrella surface and oral arms were

affiliated with *Phaeobacter, Ruegeria*, and within *Gammaproteobacteria* with

Stenotrophomonas, *Alteromonas*, *Pseudoalteromonas*, and *Vibrio.* In the gastral cavity were

members of *Betaproteobacteria* affiliated with *Burkholderia*, *Cupriavidus*, and

Achromobacter. Members of *Gammaproteobacteria* affiliated mostly with *Pseudomonas*, and

members of *Actinobacteria* with *Kocuria.*

In contrast to the total bacterial community, bacteria isolated from jellyfish (from both the

exumbrella surface and gastral cavity) were mostly affiliated with *Gammaproteobacteria,*

within the most relevant members affiliated with *Vibrio*, *Stenotrophomonas*, and

Pseudoalteromonas. The observed dominance of different bacterial classes within the total

bacterial community and cultured bacterial community in our study is not that surprising,

since both culturing and molecular-based methods are biased towards certain microbial

groups. Similar observations, were reported previously by Rohwer [11] studying coral-

associated bacterial communities. In addition, the culturing approach revealed the presence of

bacteria affiliated with *Microbacterium* (*Microbacteriaceae*), *Sphingobacterium*

(*Sphingobacteriaceae*), *Brevundimonas* (*Caulobacteraceae*), and *Delftia* (*Comamonadaceae*).

However, considering the main representatives within bacterial groups, molecular-based

studies and the culture-dependent method more or less pointed to the presence/dominance of

the same bacterial taxa.

The pronounced difference in composition between gastral cavity bacterial community and

communities of exumbrella and oral arms surface could be the consequence of different

surface/epithelial structures and their function. Exumbrella and oral arms are densely covered

with cilia. The epidermis contains numerous mucus cells, especially in densely ciliated area

[39]. Mucus cells were thought to contribute to the constantly renewing mucus layer involved

in surface cleaning [40], and potentially controlling the density of associated bacteria. The

exumbrella and oral arms surfaces are in constant contact with bacteria in surrounding

ambient seawater, attracted by secreted mucus, which is potentially a high quality energy

source and settling niche. The bacteria of genus *Phaeobacter* and genus *Ruegeria,* which

belongs to *Silicibacter-Ruegeria* subgroup, are members of *Roseobacter* clade, known as the

successful surface colonizers, and as the fastest utilizers of nutrients in the marine

environment [80]. They produce acylated homoserine lactons (AHLs), the quorum-sensing

signals involved in biofilm formation and function [80]. Besides the production of broad

range biologically active metabolites, bacteria of the *Pseudoalteromanas* genus produce

extracellular enzymes and exopolysaccharides, which all together enable them to successfully

compete for nutrients and colonization of surfaces [79]. Bacteria of *Alteromonas* and *Vibrio*

genus are widespread in the marine environment and are common surface and particle

colonizers [81]. *Alteromonas* produces and secretes a variety of extracellular enzymes that

contribute to the hydrolysis of biopolymers, including polysaccharides, proteins, nucleic

acids, and lipids, which are the major components of marine organic particles [81]. According

to Allers *et al.* [98], their versatile metabolism helps them exploit a complex substrate source,

such as coral mucus, which in composition resembles to mucus produced by jellyfish *A.*

aurita [43]. *Vibrio* species are major chitin utilizers, largely contributing to global carbon and

nitrogen cycling. Although association with insoluble chitinous surface of detritus and life

zooplankton is a preferable lifestyle for vibrios [81], they were found in association with other

marine animals, including jellyfish, and there are indices that could be highly enriched in the

seawater at the end of the jellyfish blooms [33]. Ritchie [90] characterized the *Vibrio* species

more as 'visitors' than true residents, and as commensal microorganisms that can potentially

become opportunistic under certain conditions. *Vibrio coralliilyticus* was found in high

abundances in coral tissue slurry [9] and proven to infect and cause tissue damage in corals at

higher temperatures [99]. The *Stenotrphomonas* genus was usually represented in low

abundances in communities associated with marine animals [3,5,9], but found to be producing

antimicrobial compounds [100]. Otherwise, *Stenotrophomonas* species are found in many

environments, but mostly associated with terrestrial plants that provide plant protection and

growth promotion. They were also found to be resistant to heavy metals and antibiotics, and

to degrade pollutants like polycyclic aromatic hydrocarbons (PAHs) and xenobiotics [85].

In contrast, the gastral cavity is somehow isolated from the surrounding environment. The

gastral cavity surface is covered by finger-like villi, with numerous cilia at their apical region,

and with vesiculous receptacles at the basal region. Mucus cells in the gastrodermis are

present mostly at the apical region of the villi, while gastrodermis at the basal region is

composed mainly by serous cells, producing digestive enzymes [39]. As a niche, the jellyfish

'gut' could somehow impose strict requirements of bacteria to survive. Studies on *Cotylorhiza*

tuberculata showed possible intracellular symbiotic bacteria within organs of the gastral

cavity (gastric filaments), with possible involvement in the digestion process [26]. The

dominance of *Betaproteobacteria* in medusa gastral cavity detected within our study is

somehow surprising, since they are more characteristic for organic aggregates in limnetic

ecosystems [89]. However, bacteria of the *Burkholderia* and *Achromobacter* genus were also

isolated from the marine environment, including animals [83,87]. Both were found to be able

to degrade PAHs and to be resistant to multiple antibiotics [82–84,86,87]. Similarly,

Achromobacter species were found to be n- alkane degrader and to remove also anthracene,

phenanthrene, and pyrence from the environment [82]. In addition, *Achromobacter* sp. HZ01

possesses genes related to the metabolism of secondary metabolites [83]. The *Cupriavidus*

species were not detected in the marine environment, to our knowledge, however they were

attributed with the ability to degrade aliphatic hydrocarbons [101]. Similarly, were the marine

Pseudomonas species found to be able to degrade hydrocarbons like naphthalene, present

within petroleum [86,102]. However, they were also found in association with sponges,

producing antimicrobial compounds [2,100]. *Kocuria* isolated from marine sponges were

found to produce the antibiotic kocurin, active against *Staphylococcus aureus* (MRSA)

[103,104]. Even more so, *Kocuria* isolated from the marine environment was found to be able

to utilize polyethylene as a sole carbon source [105]. The Gulf of Trieste, an ecosystem where

Aurelia used in our study was collected, is known to be impacted by different anthropogenic

pressures. Consequently, the sediment and the water column is polluted with PAHs and other

chemical compounds [106] as well as by faecal bacteria, originating from coastal run off and

municipal wastewater discharges [106]. This suggests that the bacterial community associated

with jellyfish from this environment could be adapted to such environmental conditions.

Furthermore, supporting our hypothesis, polyps generating *Aurelia* medusa were previously

found attached to artificial structures such as port pillars [107]. This indicates that pollution

adapted bacterial community could evolved and prosper at polyp and medusa stages. This

altogether suggests possible impact of anthropogenic pollution on the structure of bacterial

community associated with jellyfish and possible adaptation mechanism of jellyfish

associated bacterial population.

Hosts recruit bacteria, which are beneficial for their development or contribute to their well-

being. Selection of certain bacteria in different medusa body parts could be the strategy of

Aurelia, to harbor bacteria with specific functions needed in different body parts. Bacterial

associates found on *Aurelia* exumbrella and oral arms surface were previously supposed to

assist in host defense against pathogens and fouling organisms from surrounding seawater

[13,93,108,109]. This is not surprising, since *Vibrio* and pigmented strains of *Phaeobacter*,

Ruegeria, and *Pseudoalteromonas* produce antimicrobial compounds when attached to live or

inert surfaces [44,109–112]. In addition, some of extracellular compounds produced by

Pseudoalteromonas bacteria were found to enhance the chances of host organisms to survive

in specific marine habitats [79]. Apprill *et al.* [78] even proposed possible role of

Pseudoalteromonas in coral planula settlement and adhesion process, while were the

Roseobacter clade bacteria supposed to be important in coral development. The role in host

defense was also proposed for bacteria of *Stenotrophomonas,* detected within all body part

communities and for *Pseudomonas* [2,100], and *Actinobacteria* [103,113], present within the

gastral cavity. However, bacteria in 'digestive system' should be more involved in food

digestion and nutrition, which is reasonable, since jellyfish are supposed to 'lack' some

digestive enzymes to utilize prey [114]. An especially intriguing ability of the gastral cavity

bacteria is degradation of PAH's, xenobiotics, and plastic. Jellyfish mucus was found to have

structural properties to effectively accumulate nanoparticles [40] and PAHs [115]. This

property could be also applied to entrap micro- plastic particles, which was demonstrated for

corals [116]. PAHs were found to be highly toxic for zooplankton organisms, however, adult

medusa *A. aurita* and *M. leidyi* showed a higher tolerance to exposure [117]. *A. aurita* under

stress conditions, release blobs of mucus [41], detected also under exposure to crude oil

(containing PAHs) [115]. This suggests that sloughing as a possible way to reduce the toxic

effect of crude oil. In addition, when PAHs were entrapped within jellyfish mucus,

hydrocarbon-degrading bacteria cell densities doubled, which resulted in a significant increase

in oil degradation [115]. Some of toxic particles, entrapped within mucus and covered with

degrading bacteria, could be also transferred by ciliary currents and boundary layer flow to a

marginal umbrella groove, and then to gastral cavity, since this is one type of prey capture

recognized for *A. aurita* [114], which could explain high abundances of hydrocarbon and

plastic degrading bacteria found in the gut of *Aurelia* jellyfish within our study. Even more so,

since toxic compounds are utilized by associated bacteria, less are accumulated within

jellyfish tissue and are not transferred to higher trophic levels.

Bacterial community structure shifts due to jellyfish *Aurelia* population collapse

The shift in bacterial community composition within a one-month period (from May to June)

was observed. It resulted in a higher abundance of *Gammaproteobacteria*, especially *Vibrio*,

which became a dominant member of community. This shift towards *Vibrio* was even more

evident in cultural bacterial community.

The major difference between both studied months was a rise in the temperature and the

viability state of *Aurelia* jellyfish in the Gulf of Trieste. In June was the end of blooming

period and jellyfish were in the phase of dying, which was evident as typical signs of

moribund jellyfish: degenerated tentacles, oral structures, and gonads, reduced swimming

ability and necrosis of the epithelial bell tissue [118]. The process is normally triggered by

environmental stress like change in temperature or salinity, food availability, parasitism, and

spawning or even more likely, interacting stressors [118]. Nevertheless, in summer, the

greater part of the *Aurelia* jellyfish population was found to be parasitized, along with altered

morphology, growth, and swimming pattern in the Big Lake (Mljet Island, Croatia) [119].

This could indicate that jellyfish defense mechanism was probably disturbed due to

environmental stress (higher temperature), which resulted in parasitism and mortality.

The senescing process in jellyfish indirectly affects interaction between jellyfish (host) and

bacterial associates, which leads to a shift in the associated microbial community. Moribund

jellyfish, without their own defense mechanisms, represent organic rich particles, where the

structure of associated bacterial community is influenced solely by bacterium-bacterium

antagonism and environmental conditions, determining the presence of metabolically active

bacteria. Jellyfish tissue was found to be high quality labile organic substrate for bacteria [33-

35]. Previous bacterial degradation experiments performed on *Aurelia* jellyfish in the Gulf of

Trieste, resulted in the increase in *Vibrio* abundance [33].

Vibrio was recognized before as 'visitor,' exploiting the nutrient-rich niche [90]. As such, this

commensal microorganism probably dwells in the 'cloud' of jellyfish mucus and under the

right conditions becomes opportunistic. Vibrios rapidly grow in organic-rich environments

[120], and together with tolerance to higher temperature [121] (documented in June in the

Gulf of Trieste), up-regulating virulence determinants such as motility, resistance to

antimicrobial compounds, hemolysis, and cytotoxicity detected in coral pathogens [81] and

the references within), it can outcompete other bacterial residents and become highly

dominant.

Conclusion

Both culture-dependent and independent methods have been extensively used to study and to

understand the role of microbial communities associated with marine animals, especially

crustacean zooplankton, and benthic sponges and corals. Data available on *A. aurita* are still

limited. With the exception of *Mycoplasma* bacteria, a possible endosymbiont detected within

A. aurita tissue [18,27], the nature of the relationship between *Aurelia* jellyfish and bacterial

associates is not straight forward. In addition, it is hard to say whether or not these bacteria

are true residents of jellyfish, forming a species-specific association with the host or are just

opportunistic microbes residing in niche of an organically- rich environment. So far, we only

speculate on the role of bacterial associates, although Weiland-Bräuer *et al.* [27] suggested

associated bacteria may play important functional roles during the life cycle of *A. aurita*.

Bacteria associated with *Aurelia* jellyfish in the Gulf of Trieste were found to be mostly

generalists, composing for the host beneficial assemblage possibly involved in food digestion

and protection from toxic compounds, pathogens, and other fouling organisms. With

speculation on the active and passive role of *Aurelia* jellyfish in selection of bacterial

associates, demonstrated for other animals [45], jellyfish may form a relationship with diverse

metabolically active microorganisms, providing more effective adaptation of host to changing

environmental and anthropogenic conditions. From this perspective, the relationship somehow

resembles suggested coral probiotic theory [122].

Further investigation of such a relationship is necessary to understand the relevance of the

associated bacteria for the host during its life spam and during/after the bloom period,

especially in areas experiencing seasonal blooms, influencing food webs, and biogeochemical

cycles in those regions. Even more so, despite the small number of experimental data, our

results suggest that the jellyfish - bacteria link could be applied as an effective pollution-

control method in marine environments affected by crude oil and micro plastic. Finally, we

would also like to emphasize the importance of culturing organisms. Although the method is

biased towards certain bacterial groups, it remains important to obtain complete genome

sequences, to identify properties of organisms, and to help understand the biology and

ecology of microbial species.

Acknowledgments

We would like to thank R/V Sagita crew and Dr. Mateja Grego for her help with Primer v6
analyses. We are grateful to anonymous reviewers for their critical and valuable comments on
the manuscript.

References

1.   Apprill A. Marine Animal Microbiomes: Toward understanding host–microbiome
interactions in a changing ocean. Front Mar Sci. 2017;4(July):1–9. Available from:
http://journal.frontiersin.org/article/10.3389/fmars.2017.00222/full
2.   Thakur NL, Anil AC. Antibacterial activity of the sponge *Ircinia ramosa*: Importance
of its surface- associated bacteria. J Chem Ecol. 2000;26(1):57–71.
3.   Radwan M, Hanora A, Zan J, Mohamed NM, Abo- Elmatty DM, Abou-El-Ela SH, et
al. Bacterial community analyses of two Red Sea sponges. Mar Biotechnol.
2010;12:350–60.
4.   Webster NS, Negri AP, Munro MMHG, Battershill CN. Diverse microbial
communities inhabit Antarctic sponges. Environ Microbiol. 2004;6(3):288–300.
5.   Li ZY, He LM, Wu J, Jiang Q. Bacterial community diversity associated with four
marine sponges from the South China Sea based on 16S rDNA-DGGE fingerprinting. J
Exp Mar Bio Ecol. 2006;329(1):75–85.
6.   Hentschel U, Hopke J, Horn M, Friedrich AB, Wagner M, Hacker J, et al. Molecular
Evidence for a uniform microbial community in sponges from different oceans. Appl
Environ Microbiol. 2002;68(9):4431–40.
7.   Kittelmann S, Harder T. Species- and site-specific bacterial communities associated
with four encrusting bryozoans from the North Sea, Germany. J Exp Mar Bio Ecol.
2005;327(2):201–9.
8.   Koren O, Rosenberg E. Bacteria Associated with Mucus and Tissues of the Coral
Oculina patagonica in Summer and Winter. Appl Environ Microbiol.
2006;72(8):5254–9. Available from: http://aem.asm.org/cgi/doi/10.1128/AEM.00554-
06
9.   Bourne DG, Munn CB. Diversity of bacteria associated with the coral *Pocillopora
damicornis* from the Great Barrier Reef. Environ Microbiol. 2005;7(8):1162–74.
10.   Rohwer F, Seguritan V, Azam F, Knowlton N. Diversity and distribution of coral-
associated bacteria. Mar Ecol Prog Ser. 2002;243:1–10.
11.   Rohwer F, Breitbart M, Jara J, Azam F, Knowlton N. Diversity of bacteria associated
with the Caribbean coral *Montastraea franksi*. Coral Reefs. 2001;20(1):85–91.
12.   Webster NS, Bourne D. Bacterial community structure associated with the Antarctic
soft coral, *Alcyonium antarcticum*. FEMS Microbiol Ecol. 2007;59:81–94.
13.   Harder T, Lau SCK, Dobretsov S, Fang TK, Qian P-Y. A distinctive epibiotic bacterial
community on the soft coral *Dendronephthya* sp. and antibacterial activity of coral
tissue extracts suggest a chemical mechanism against bacterial epibiosis. FEMS
Microbiol Ecol. 2003;43:337–47.
14.   Tang KW, Turk V, Grossart H. Linkage between crustacean zooplankton and aquatic
bacteria. Aquat Microb Ecol. 2010;61(3):261–77.
15.   Gerdts G, Brandt P, Kreisel K, Boersma M, Schoo KL, Wichels A. The microbiome of
North Sea copepods. Helgol Mar Res. 2013;67(4):757–73.
16.   Grossart HP, Riemann L, Tang KW. Molecular and functional ecology of aquatic
microbial symbionts. Front Microbiol. 2013;4(MAR):2012–3.
17.   Flood PR. Architecture of, and water circulation and flow rate in, the house of the
planktonic tunicate *Oikopleura labradorensis*. Mar Biol. 1991;111:95–111.
18.   Daley MC, Urban-Rich J, Moisander PH. Bacterial associations with the hydromedusa
Nemopsis bachei and scyphomedusa *Aurelia aurita* from the North Atlantic Ocean.
Mar Biol Res. 2016;12(10):1088–100.
19.   Daniels C, Breitbart M. Bacterial communities associated with the ctenophores
Mnemiopsis leidyi and *Beroe ovata*. FEMS Microbiol Ecol. 2012;82(1):90–101.
20.   Dinasquet J, Granhag L, Riemann L. Stimulated bacterioplankton growth and selection

for certain bacterial taxa in the vicinity of the ctenophore *Mnemiopsis leidyi*. Front
Microbiol. 2012;3(AUG):1–8.

21. Hao W. Bacterial communities associated with jellyfish. Bremen University; 2014.

22. Cleary DFR, Becking LE, Polónia ARM, Freitas RM, Gomes NCM. Jellyfish-
associated bacterial communities and bacterioplankton in Indonesian Marine lakes.
FEMS Microbiol Ecol. 2016;92(5):1–14.

23. Cortés-Lara S, Urdiain M, Mora-Ruiz M, Prieto L, Rosselló-Móra R. Prokaryotic
microbiota in the digestive cavity of the jellyfish *Cotylorhiza tuberculata*. Syst Appl
Microbiol. 2015;38(7):494–500. Available from:
http://dx.doi.org/10.1016/j.syapm.2015.07.001

24. Delannoy CMJ, Houghton JDR, Fleming NEC, Ferguson HW. Mauve Stingers (
Pelagia noctiluca) as carriers of the bacterial fi sh pathogen Tenacibaculum
maritimum. Aquaculture. 2011;311(1–4):255–7. Available from:
http://dx.doi.org/10.1016/j.aquaculture.2010.11.033

25. Ferguson HW, Delannoy CMJ, Hay S, Nicolson J, Sutherland D, Crumlish M. Jellyfish
as vectors of bacterial disease for farmed salmon (*Salmo salar*). J Vet Diagn Invest.
2010;22:376–82.

26. Viver T, Orellana LH, Hatt JK, Urdiain M, Díaz S, Richter M, et al. The low diverse
gastric microbiome of the jellyfish *Cotylorhiza tuberculata* is dominated by four novel
taxa. Environ Microbiol. 2017;19(8):3039–58.

27. Weiland-Bräuer N, Neulinger SC, Pinnow N, Künzel S, Baines JF, Schmitz RA.
Composition of bacterial communities associated with *Aurelia aurita* changes with
compartment, life stage, and population. Appl Environ Microbiol. 2015;81(17).

28. Schuett C, Doepke H. Endobiotic bacteria and their pathogenic potential in cnidarian
tentacles. Helgol Mar Res. 2010;64:205–12.

29. Riemann L, Titelman J, Båmstedt U. Links between jellyfish and microbes in a
jellyfish dominated fjord. Mar Ecol Prog Ser. 2006;325(2003):29–42.

30. Manzari C, Fosso B, Marzano M, Annese A, Caprioli R, D'Erchia AM, et al. The
influence of invasive jellyfish blooms on the aquatic microbiome in a coastal lagoon
(Varano, SE Italy) detected by an Illumina-based deep sequencing strategy. Biol
Invasions. 2015;17(3):923–40.

31. Condon RH, Steinberg DK, del Giorgio PA, Bouvier TC, Bronk DA, Graham WM, et
al. Jellyfish blooms result in a major microbial respiratory sink of carbon in marine
systems. Proc Natl Acad Sci. 2011;108(25):10225–30. Available from:
http://www.pnas.org/cgi/doi/10.1073/pnas.1015782108

32. Blanchet M, Pringault O, Bouvy M, Catala P, Oriol L, Caparros J, et al. Changes in
bacterial community metabolism and composition during the degradation of dissolved
organic matter from the jellyfish *Aurelia aurita* in a Mediterranean coastal lagoon.
Environ Sci Pollut Res. 2015; 22, 18: 13638-13653.

33. Tinta T, Kogovšek T, Malej A, Turk V. Jellyfish modulate bacterial dynamic and
community structure. PLoS One. 2012;7(6):1–11.

34. Tinta T, Kogovšek T, Turk V, Shiganova TA, Mikaelyan AS, Malej A. Microbial
transformation of jellyfish organic matter affects the nitrogen cycle in the marine water
column - A Black Sea case study. J Exp Mar Bio Ecol. 2016;475:19–30.

35. Tinta T, Malej A, Kos M, Turk V. Degradation of the Adriatic medusa *Aurelia* sp. by
ambient bacteria. Hydrobiologia. 2010;645:179–91.

36. Titelman J, Riemann L, Sørnes TA, Nilsen T, Griekspoor P, Båmstedt U. Turnover of
dead jellyfish□: stimulation and retardation of microbial activity. Mar Ecol Prog Ser.
2006;325:43–58.

37. Pitt KA, Welsh DT, Condon RH. Influence of jellyfish blooms on carbon , nitrogen and
phosphorus cycling and plankton production. Hydrobiologia. 2009;616:133–49.

38. Bosch TCG. Cnidarian-microbe interactions and the origin of innate immunity in
Metazoans. Annu Rev Microbiol. 2013;67(1):499–518. Available from:
http://www.annualreviews.org/doi/10.1146/annurev-micro-092412-155626

39. Heeger T, Möller H. Ultrastructural observations on prey capture and digestion in the
scyphomedusa *Aurelia aurita*. Mar Biol. 1987;96(3):391–400.

40. Patwa A, Thiéry A, Lombard F, Lilley MKS, Boisset C, Bramard J, et al.
Accumulation of nanoparticles in " jellyfish " mucus: a bio-inspired route to

952 decontamination of nano- waste. Nat Sci Reports. 2015;1–8. Available from:
953 http://dx.doi.org/10.1038/srep11387

954 41. Shanks A, Graham W. Chemical defense in a scyphomedusa. Mar Ecol Prog Ser.
955 1988;45:81–6.

956 42. Ovchinnikova T V., Balandin S V., Aleshina GM, Tagaev A a., Leonova YF,
957 Krasnodembsky ED, et al. Aurelin, a novel antimicrobial peptide from jellyfish *Aurelia*
958 *aurita* with structural features of defensins and channel-blocking toxins. Biochem
959 Biophys Res Commun. 2006;348(2):514–23.

960 43. Ducklow HW, Mitchell R. Composition of mucus released by coral reef coelenterates.
961 Limnol Oceanogr. 1979;24(4):706–14.

962 44. Long RA, Azam F. Antagonistic interactions among marine pelagic bacteria. Appl
963 Environ Microbiol. 2001;67(11):4975–83.

964 45. Wahl M, Goecke F, Labes A, Dobretsov S, Weinberger F. The second skin: ecological
965 role of epibiotic biofilms on marine organisms. Front Microbiol. 2012;3:1–21.

966 46. Ramšak A, Stopar K, Malej A. Comparative phylogeography of meroplanktonic
967 species, *Aurelia* spp. and Rhizostoma pulmo (Cnidaria: Scyphozoa) in European Seas.
968 Hydrobiologia. 2012;690(1):69–80.

969 47. Scorrano S, Aglieri G, Boero F, Dawson MN, Piraino S. Unmasking *Aurelia* species in
970 the Mediterranean Sea: An integrative morphometric and molecular approach. Zool J
971 Linn Soc. 2017;180(2):243–67.

972 48. Kogovšek T, Bogunović B, Malej A. Recurrence of bloom-forming scyphomedusae:
973 Wavelet analysis of a 200-year time series. Hydrobiologia. 2010;645(1):81–96.

974 49. Malej A, Kogovšek T, Ramšak A, Catenacci L. Blooms and population dynamics of
975 moon jellyfish in the northern Adriatic. Cah Biol Mar. 2012;53(3):337–42.

976 50. Kogovšek T, Klun K, Ikeda H, Tinta T, Uye S. Starvation - an important factor
977 controlling scyphozoan population?. In: Fifth International Jellyfish Bloom
978 Symposium￼: Abstract book. Barcelona: Barcelona University. 2016; 2016. p. 33.

979 51. Brotz L, Cheung WWL, Kleisner K, Pakhomov E, Pauly D. Increasing jellyfish
980 populations: Trends in Large Marine Ecosystems. Hydrobiologia. 2012;690(1):3–20.

981 52. Condon RH, Graham WM, Duarte CM, Pitt KA, Lucas CH, Haddock SHD, et al.
982 Questioning the rise of gelatinous zooplankton in the World's Oceans. Bioscience.
983 2012;62(2):160–9. Available from:
984 http://bioscience.oxfordjournals.org/content/62/2/160.abstract

985 53. Condon RH, Duarte CM, Pitt KA, Robinson KL, Lucas CH, Sutherland KR, et al.
986 Recurrent jellyfish blooms are a consequence of global oscillations. PNAS.
987 2013;110(3):1000–5.

988 54. Purcell JE, Uye S, Lo W. Anthropogenic causes of jellyfish blooms and their direct
989 consequences for humans: a review. Mar Ecol Prog Ser. 2007;350:153–74.

990 55. Graham WM, Gelcich S, Robinson KL, Duarte CM, Brotz L, Purcell JE, et al. Linking
991 human well-being and jellyfish: Ecosystem services, impacts, and societal responses.
992 Front Ecol Environ. 2014;12(9):515–23.

993 56. Richardson AJ, Bakun A, Hays GC, Gibbons MJ. The jellyfish joyride: causes,
994 consequences and management responses to a more gelatinous future. Trends Ecol
995 Evol. 2009;24(6):312–22. Available from:
996 http://linkinghub.elsevier.com/retrieve/pii/S0169534709000883

997 57. Purcell JE. Jellyfish and ctenophore blooms coincide with human proliferations and
998 environmental perturbations. Ann Rev Mar Sci. 2012;4(1):209–35. Available from:
999 http://www.annualreviews.org/doi/abs/10.1146/annurev-marine-120709-142751

1000 58. Vodopivec M, Peliz AJ, Malej A. Offshore marine constructions as propagators of
1001 moon jellyfish dispersal. Environ Res Lett. 2017;12(8).

1002 59. Kogovšek T, Vodopivec M, Raicich F, Uye S, Malej A. Comparative analysis of the
1003 ecosystems in the northern Adriatic Sea and the Inland Sea of Japan: Can
1004 anthropogenic pressures disclose jellyfish outbreaks? Sci Total Environ [Internet].
1005 2018;626:982–94. Available from:
1006 http://www.sciencedirect.com/science/article/pii/S0048969718300111

1007 60. Malačič V, Petelin B. Gulf of Trieste. In: Cushman-Roisin B, Gacic M, Poulain P-M,
1008 Artegiani A, editors. Physical oceanography of the Adriatic Sea: past, present and
1009 future. Dordrecht: Kluwer Academic Press; 2001. p. 167–77.

1010   61.   Zobell CE. Marine microbiology. Marine microbiology. A monograph on
1011   hydrobacteriology. Chronica Botanica.; 1946. 240 p.
1012   62.   Giraffa G, Rossetti L, Neviani E. An evaluation of chelex-based DNA purification
1013   protocols for the typing of lactic acid bacteria. J Microbiol Methods. 2000;42(2):175–
1014   84.
1015   63.   Boström KH, Simu K, Hagström Å, Riemann L. Optimization of DNA extraction for
1016   quantitative marine bacterioplankton community analysis. Limnol Oceanogr Methods.
1017   2004;2(1988):365–73.
1018   64.   Muyzer G, De Waal EC, Uitterlinden AG. Profiling of complex microbial populations
1019   by denaturing gradient gel electrophoresis analysis of polymerase chain reaction-
1020   amplified genes coding for 16S rRNA. Appl Environ Microbiol. 1993;59(3):695–700.
1021   65.   Muyzer G, Smalla K. Application of denaturing gradient gel electrophoresis (DGGE)
1022   and temperature gradient gel electrophoresis (TGGE) in microbial ecology. Antonie
1023   Van Leeuwenhoek. 1998;73:127–41.
1024   66.   Don RH, Cox PT, Wainwright BJ, Baker K, Mattick JS. "Touchdown" PCR to
1025   circumvent spurious priming during gene amplification. Nucleic Acids Res.
1026   1991;19(14):4008.
1027   67.   Dar SA, Kuenen JG, Muyzer G. Nested PCR-Denaturing Gradient Gel Electrophoresis
1028   approach to determine the diversity of sulfate-reducing bacteria in complex microbial
1029   communities. Appl Environ Microbiol. 2005;71(5):2325–30.
1030   68.   Giloteaux L, Go~ni-Urriza M, Duran R. Nested PCR and new primers for analysis of
1031   sulfate-reducing bacteria in low-cell-biomass environments. Appl Environ Microbiol.
1032   2010;76(9):2856–65.
1033   69.   Schloss PD, Westcott SL, Ryabin T, Hall JR, Hartmann M, Hollister EB, et al.
1034   Introducing mothur: Open-source, platform-independent, community-supported
1035   software for describing and comparing microbial communities. Appl Environ
1036   Microbiol. 2009;75(23):7537–41.
1037   70.   Good IJ. No Title. Biometrika. 1953;40(3/4):237–64.
1038   71.   Clarke KR, Gorley RN. PRIMER v6: Manual/Tutorial. Prim Plymouth. 2006;
1039   72.   Hammer Ø, Harper DAT, Ryan PD. PAST-PAlaeontological STatistics, ver. 1.89.
1040   Palaeontol Electron. 2001;4(1):1–9.
1041   73.   Taylor MW, Schupp PJ, Dahllöf I, Kjelleberg S, Steinberg PD. Host specifity in marine
1042   sponge- associated bacteria. Environ Microbiol. 2004;6(2):121–30.
1043   74.   Cooney RP, Pantos O, Tissier MDA Le, Barer MR, Donnell AGO, Bythell JC.
1044   Characterization of the bacterial consortium associated with black band disease in coral
1045   using molecular microbiological techniques. Environ Microbiol. 2002;4(7):401–13.
1046   75.   Bythell JC, Barer MR, Cooney RP, Guest JR, O'Donnell AG, Pantos O, et al.
1047   Histopathological methods for the investigation of microbial communities associated
1048   with disease lesions in reef corals. Lett Appl Microbiol. 2002;34:359–64.
1049   76.   Johnston IS, Rohwer F. Microbial landscapes on the outer tissue surfaces of the reef-
1050   building coral *Porites compressa*. Coral Reefs. 2007;26(2):375–83.
1051   77.   Garren M, Azam F. Corals shed bacteria as a potential mechanism of resilience to
1052   organic matter enrichment. ISME J [Internet]. 2012;6:1159–65. Available from:
1053   http://dx.doi.org/10.1038/ismej.2011.180
1054   78.   Apprill A, Marlow HQ, Martindale MQ, Rappé MS. Specificity of associations
1055   between bacteria and the coral *Pocillopora meandrina* during early development. Appl
1056   Environ Microbiol. 2012;78(20):7467–75.
1057   79.   Holmström C, Kjelleberg S. Marine P*seudoalteromonas* species are associated with
1058   higher organisms and produce biologically active extracellular agents. FEMS
1059   Microbiol Ecol. 1999;30:285–93.
1060   80.   Dang H, Li T, Chen M, Huang G. Cross-Ocean Distribution of *Rhodobacterales*
1061   bacteria as primary surface colonizers in temperate coastal marine waters. Appl
1062   Environ Microbiol. 2008;74(1):52–60.
1063   81.   Dang H, Lovell CR. Microbial surface colonization and biofilm development in marine
1064   environments. Microbiol Mol Biol Rev. 2016;80(1):91–138.
1065   82.   Deng M-C, Li J, Liang F-R, Yi M, Xu X-M, Yuan J-P, et al. Isolation and
1066   characterization of a novel hydrocarbon-degrading bacterium *Achromobacter* sp. HZ01
1067   from the crude oil-contaminated seawater at the Daya Bay, southern China. Mar Pollut

1068   Bull. 2014;83(1):79–86.
1069   http://www.sciencedirect.com/science/article/pii/S0025326X14002306
1070 83. Hong YH, Ye CC, Zhou QZ, Wu XY, Yuan JP, Peng J, et al. Genome sequencing
1071   reveals the potential of *Achromobacter* sp. HZ01 for bioremediation. Front Microbiol.
1072   2017;8:1–14.
1073 84. Juhasz AL, Britz ML, Stanley GA. Degradation of fluoranthene, pyrene, benz [a]
1074   anthracene and dibenz [a,h] anthracene by *Burkholderia cepacia*. J Appl Microbiol.
1075   1997;83(2):189–98.
1076 85. Ryan RP, Monchy S, Cardinale M, Taghavi S, Crossman L, Avison MB, et al. The
1077   versatility and adaptation of bacteria from the genus *Stenotrophomonas*. Nat Rev
1078   Microbiol. 2009;7:514–25. http://dx.doi.org/10.1038/nrmicro2163
1079 86. Harayama S, Kishira H, Kasai Y, Shutsubo K. Petroleum biodegradation in marine
1080   environments. J Mol Microbiol Biotechnol. 1999;1(1):63–70.
1081 87. Maravić A, Skočibušić M, Šprung M, Šamanić I, Puizina J, Pavela-Vrančić M.
1082   Occurrence and antibiotic susceptibility profiles of *Burkholderia cepacia* complex in
1083   coastal marine environment. Int J Environ Health Res. 2012;22(6):531–42.
1084   http://www.tandfonline.com/doi/abs/10.1080/09603123.2012.667797
1085 88. DeLong EF, Franks DG, Alldredge AL. Phylogenetic diversity of aggregate-attached
1086   marine bacterial assemblages. Limnol Oceanogr. 1993;38(5):924–34.
1087 89. Simon M, Grossart H, Schweitzer B, Ploug H. Microbial ecology of organic aggregaes
1088   in aquatic ecosystems. Aquat Microb Ecol. 2002;28:175–211.
1089 90. Ritchie KB. Regulation of microbial populations by coral surface mucus and mucus-
1090   associated bacteria. Mar Ecol Prog Ser. 2006;322:1–14.
1091 91. Daniels CA, Zeifman A, Heym K, Ritchie KB, Watson CA, Berzins I, et al. Spatial
1092   heterogeneity of bacterial communities in the mucus of *Montastraea annularis*. Mar
1093   Ecol Prog Ser. 2011;426:29–40.
1094 92. Lampert Y, Kelman D, Dubinsky Z, Nitzan Y, Hill RT. Diversity of culturable bacteria
1095   in the mucus of the Red Sea coral *Fungia scutaria*. FEMS Microbiol Ecol.
1096   2006;58(1):99–108.
1097 93. Shnit-Orland M, Kushmaro A. Coral mucus-associated bacteria: a possible first line of
1098   defense. FEMS Microbiol Ecol. 2009;67:371–80.
1099 94. Hao W, Gerdts G, Peplies J, Wichels A. Bacterial communities associated with four
1100   ctenophore genera from the German Bight (North Sea). FEMS Microbiol Ecol.
1101   2015;(October 2014):1–11.
1102   http://femsec.oxfordjournals.org/cgi/doi/10.1093/femsec/fiu006
1103 95. Kirchman DL, Dittel AI, Malmstrom RR, Cottrell MT. Biogeography of major
1104   bacterial groups in the Delaware Estuary. Limnol Oceanogr. 2005;50(5):1697–706.
1105 96. Webster NS, Wilson KJ, Blackall LL, Hill RT. Phylogenetic diversity of bacteria
1106   associated with the marine sponge *Rhopaloeides odorabile*. Appl Environ Microbiol.
1107   2001;67(1):434–44.
1108 97. Franzenburg S, Walter J, Künzel S, Wang J, Baines JF, Bosch TCG. Distinct
1109   antimicrobial peptide expression determines host species-specific bacterial
1110   associations. Proc Natl Acad Sci U S A. 2013;110(39):E3730–8.
1111 98. Allers E, Niesner C, Wild C, Pernthaler J. Microbes enriched in seawater after addition
1112   of coral mucus. Appl Environ Microbiol. 2008;74(10):3274–8.
1113 99. Ben- Haim Y, Thompson FL, Thompson CC, Cnockaert MC, Hoste B, Swings J, et al.
1114   *Vibrio coralliilyticus* sp . nov., a temperature- dependent pathogen of the coral
1115   *Pocillopora damicornis*. Int J Syst Evol Microbiol. 2003;53:309–15.
1116 100. Santos OCS, Pontes PVML, Santos JFM, Muricy G, Giambiagi-deMarval M, Laport
1117   MS. Isolation, characterization and phylogeny of sponge- associated bacteria with
1118   antimicrobial activities from Brazil. Res Microbiol. 2010;161:604–12
1119 101. Bacosa HP, Suto K, Inoue C. Bacterial community dynamics during the preferential
1120   degradation of aromatic hydrocarbons by a microbial consortium. Int Biodeterior
1121   Biodegrad. 2012;74:109–15. Available from:
1122   http://dx.doi.org/10.1016/j.ibiod.2012.04.022
1123 102. Prince RC. Petroleum spill bioremediation in marine environments. Crit Rev
1124   Microbiol. 1993;19(4):217–40.
1125 103. Abdelmohsen UR, Bayer K, Hentschel U. Diversity, abundance and natural products of

marine sponge-associated actinomycetes. Nat Prod Rep. 2014;31(3):381–99. Available from: http://xlink.rsc.org/?DOI=C3NP70111E

104. Palomo S, González I, De La Cruz M, Martín J, Tormo JR, Anderson M, et al. Sponge-derived *Kocuria* and *Micrococcus* spp. as sources of the new thiazolyl peptide antibiotic kocurin. Mar Drugs. 2013;11(4):1071–86.

105. Harshvardhan K, Jha B. Biodegradation of low-density polyethylene by marine bacteria from pelagic water, Arabian Sea, India. Mar Pollut Bull. 2013;77:100–6.

106. Turk V, Mozetič P, Malej A. Overview of eutrophication-related events and other irregular episodes in Slovenian Sea (Gulf of Trieste, Adriatic Sea). Annales Ser hist nat 2007;17:197-216.

107. Hočevar S, Malej A, Boldin B, Purcell JE. Seasonal fluctuations in ppulaton dynamics of aurelia aurita polyp in situ with a modelling perspective. Mar Ecol Prog Ser. 2018; 591:155-166.

108. Dobretsov S, Qian P-Y. The role of epibotic bacteria from the surface of the soft coral *Dendronephthya* sp. in the inhibition of larval settlement. J Expr Mar Biol Ecol. 2004;299:35–50.

109. Holmström C, Egan S, Franks A, McCloy S, Kjelleberg S. Antifouling activities expressed by marine surface associated *Pseudoalteromonas* species. FEMS Microbiol Ecol. 2002;41:47–58.

110. Bruhn JB, Nielsen KF, Hjelm M, Hansen M, Schulz S, Gram L, et al. Ecology, inhibitory activity, and morphogenesis of a marine antagonistic bacterium belonging to the *Roseobacter* clade. Appl Environ Microbiol. 2005;71(11):7263–70.

111. Gram L, Melchiorsen J, Bruhn JB. Antibacterial activity of marine culturable bacteria collected from a global sampling of Ocean surface waters and surface swabs of marine organisms. Mar Biotechnol. 2010;12:439–51.

112. Porsby CH, Nielsen KF, Gram L. *Phaeobacter* and *Ruegeria* Species of the *Roseobacter* clade colonize separate niches in a Danish Turbot (*Scophthalmus maximus*)- rearing farm and antagonize *Vibrio anguillarum* under different growth conditions. Appl Environ Microbiol. 2008;74(23):7356–64.

113. Selvin J, Joseph S, Asha KRT, Manjusha W a., Sangeetha VS, Jayaseema DM, et al. Antibacterial potential of antagonistic *Streptomyces* sp. isolated from marine sponge *Dendrilla nigra*. FEMS Microbiol Ecol. 2004;50(2):117–22.

114. Arai MN. A Functional Biology of Scyphozoa. First edit. London: Chapman & Hall; 1997. 316 p.

115. Gemmell BJ, Bacosa HP, Liu Z, Buskey EJ. Can gelatinous zooplankton influence the fate of crude oil in marine environments? Mar Pollut Bull [Internet]. 2016;113(1–2):483–7. Available from: http://dx.doi.org/10.1016/j.marpolbul.2016.08.065

116. Hall NM, Berry KLE, Rintoul L, Hoogenboom MO. Microplastic ingestion by scleractinian corals. Mar Biol. 2015;162(3):725–32.

117. Almeda R, Wambaugh Z, Chai C, Wang Z, Liu Z, Buskey EJ. Effects of Crude Oil Exposure on bioaccumulation of polycyclic aromatic hydrocarbons and survival of adult and larval stages of gelatinous zooplankton. PLoS One. 2013;8(10):20–1.

118. Pitt KA, Chelsky Budarf A, Browne JG, Condon RH. Bloom and Bust: Why do blooms of jellyfish collapse? In: Pitt KA, Lucas CH, editors. Jellyfish Blooms. Netherlands: Springer; 2014. p. 79–103.

119. Graham WM, Chiaverano L, D'ambra I, Mianzan H, Colombo GA, Acha EM, et al. Fish and jellyfish: using the isolated marine " lakes" of Mljet Island, Croatia, to explore larger marine ecosystem complexities and ecosystem-based management approaches. In: Annales series historia naturalis. 2009. p. 39–48.

120. Eilers H, Pernthaler J, Amann R. Succession of pelagic marine bacteria during enrichment: A close look at cultivation-induced shifts. Appl Environ Microbiol. 2000;66(11):4634–40.

121. Thompson FL, Iida T, Swings J. Biodiversity of *Vibrios*. Microbiol Mol Biol Rev. 2004;68(3):403–31.

122. Reshef L, Koren O, Loya Y, Zilber-Rosenberg I, Rosenberg E. The coral probiotic hypothesis. Environ Microbiol. 2006;8(12):2068–73.

Supporting Information

S1 Table. Composition of 16S rRNA gene clone libraries (% of clones) from samples of jellyfish exumbrella (AK), oral arms (AR) and mucus from gastral cavity (AG) and seawater samples (W) at 5m depth collected on May and June 2011 in the Gulf of Trieste. Classification of bacterial clones was done down to the family level. The contribution of distinct bacterial families is expressed as a percentage of the total number of sequences in each sample. N is the total number of bacterial clones in the library; Sobs is the number of distinct bacterial families. In brackets beside bacterial family are the only representatives detected within the family. Numbers (N) in light grey and with asterisk (*) are total number of sequences recovered from clone library, including sequences affiliated with Chloroplast (%; presented at the bottom of table) that were omitted from further analysis.

S2 Table. Bacterial isolates obtained from samples of jellyfish exumbrella (AK) and mucus from gastral cavity (AG) and seawater samples (W) at 5 m depth collected on May and June 2011 in the Gulf of Trieste. Classification of bacterial isolates was done down to the genus level. The contribution of distinct bacterial genus or families is expressed as a percentage of the total number of sequences in each sample. N is the total number of isolated bacteria; Sobs is the number of distinct bacterial genus.

S3 Table. Similarities percentage (SIMPER) analysis of 16S rRNA gene clone libraries from samples of jellyfish exumbrella (AK), oral arms (AR) and mucus from gastral cavity (AG) and seawater samples (W) collected on May and June 2011 in the Gulf of Trieste. Seawater group (W) includes water samples collected at 5m depth on May and June.

S4 Table. Similarities percentage (SIMPER) analysis of culturable fraction of bacterial community associated with jellyfish exumbrella (AK), mucus from gastral cavity (AG) and seawater (W) collected in May and June 2011 in the Gulf of Trieste. Seawater group (W) includes the results of the water samples collected at 5m depth on May and June.

S5 Table. The diversity indices S, H', d, J', Chao- 1 and library coverage's (C) describing composition of total bacterial community associated with jellyfish exumbrella (AK), oral arms (AR) and mucus from gastral cavity (AG) and seawater (W) collected in May and June 2011 in the Gulf of Trieste. S represents the number of distinct bacterial families detected in each bacterial 16S rRNA gene clone library. C is a coverage value (C = (1–n1/N), where n1 is number of phylotypes appearing only once in the library and N is the library size.

S6 Table. The diversity indices S, H', d, J', Chao- 1 describing composition of culturable fraction of bacterial community associated with jellyfish exumbrella (AK) and mucus from gastral cavity (AG) and seawater (W) collected in May and June 2011 in the Gulf of Trieste. S represents the number of distinct bacterial genus detected in each sample.

S7 Table. Bacterial 16S rRNA sequences obtained from DGGE bands from jellyfish samples with their accession numbers. In the table is the name and an accession number of their closest relative in GeneBank (NCBI) with % of similarity, family, taxon and isolation source.

S8 Table. Similarities percentage (SIMPER) analysis of 16S rRNA gene clone libraries
from jellyfish samples collected at the time of population peak and at the end of the
bloom in the Gulf of Trieste. Group May includes samples of jellyfish exumbrella surface
(AK1, AK2) and oral arms (AR1) sampled in May. Group June includes samples of jellyfish
exumbrella surface (AK6, AK7) and oral arms (AR6) collected in June.

S9 Table. Similarities percentage (SIMPER) analysis of culturable fraction of bacterial
community associated with jellyfish at the time of population peak and at the end of the
bloom in the Gulf of Trieste. Group May includes samples of jellyfish exumbrella surface
(AK1, AK3, AK6) and gastral cavity (AG1, AG6) collected in May. Group June includes
samples of jellyfish exumbrella surface (AK8, AK10, AK11) and gastral cavity (AG8, AG11)
collected in June.

S1 Fig. DGGE profile of bacterial 16S rRNA gene fragments of samples from *Aurelia*
jellyfish exumbrella surface, oral arms and mucus from gastral cavity. AK1, AK2:
exumbrella surface of jellyfish collected in May; AK6, AK7: exumbrella surface of jellyfish
collected in June; AR1: sample of oral arms of jellyfish collected in May; AR6: oral arms of
jellyfish collected in June; AG1: gastral cavity mucus sample; S: standard. Numbers on the
figure represent bands that were cut from the gel and successfully sequenced; color dots place
sequence in one of bacterial groups.