1 *Plasmodium falciparum* **schizont stage transcriptome variation among clinical isolates and**
2 **laboratory-adapted clones**
3

Sarah J Tarr[1,*], Ofelia Díaz-Ingelmo[1], Lindsay B Stewart[1], Suzanne E Hocking[1], Lee Murray[1],

Craig W Duffy[1], Thomas D Otto[2], Lia Chappell[2], Julian C Rayner[2], Gordon A Awandare[3] and

David J Conway[1,*]

[1] Pathogen Molecular Biology Department, London School of Hygiene and Tropical

Medicine, London, UK

[2] Wellcome Sanger Institute, Hinxton, Cambridgeshire, UK

[3] West African Centre for Cell Biology of Infectious Pathogens, Department of Biochemistry,

Cell and Molecular Biology, University of Ghana, Legon, Ghana

[*] Corresponding authors

sarah.tarr@lshtm.ac.uk

david.conway@lshtm.ac.uk

Abstract

Malaria parasite genes exhibit variation in both sequence and expression level. There is much information on sequence polymorphism, but less resolution on natural variation in transcriptomes of parasites at specific developmental stages. This is largely because it is challenging to obtain highly replicated sampling of transcriptomes to overcome potentially confounding technical and biological variation. We address the issue in the major human parasite *Plasmodium falciparum* by obtaining RNA-seq profiles of multiple independent replicate preparations of mature schizont-stage parasites from a panel of clinical isolates recently established in culture and from long-term laboratory-adapted clones. With a goal of robustly identifying variably expressed genes, we show that increasing the numbers of biological sample replicates greatly improves the discovery rate. Generally, six independent replicates of each parasite culture is recommendable as being significantly to lower numbers, although for highly expressed genes variable expression can be detected when fewer replicates are available. A broad comparison identifies genes differing in relative expression between cultured clinical isolates and laboratory-adapted clones. Genes more highly expressed in the laboratory-adapted clones include an AP2 transcription factor gene Pf3D7_0420300 and putative methyl transferase genes. The variable expression of several known merozoite invasion ligands is confirmed, and previously uncharacterised genes are shown to be differentially expressed among clinical isolates. New RT-qPCR assays validate the variation in transcript levels of these genes, and allow quantitation of expression to be extended to a wider panel of clinical isolate samples. These variably expressed genes are new candidates for investigation as potential determinants of alternative parasite developmental pathways or targets of immunity.

Author summary

Understanding parasite diversity and adaptation may require characterisation of gene expression variation, and is vital if chemotherapeutic or vaccine development is to consider new candidate targets, but it is technically challenging to generate precise data on clinical isolates. Here, we analyse the transcriptomes of mature *Plasmodium falciparum* schizonts using RNA-sequencing, using large numbers of biological replicate samples to minimise the impact of inter-replicate variation on observed patterns of differential expression. This identifies genes that are differentially expressed in long term laboratory-adapted parasites

and recently cultured clinical isolates, as well as among different clinical isolates. In

additional samples of schizonts grown in the first cycle *ex vivo* prior to any erythrocyte

invasion, expression levels of a selected panel of these genes vary among isolates, but mean

levels are similar to those in the continuously cultured clinical isolates, indicating that the

latter are useful for experimental studies requiring biological replication.

Introduction

Variation in gene expression is key to survival and reproduction of microbes, affecting

diverse phenotypes such as sexual differentiation [1], adaptation to different growth

conditions [2] and evasion of host immunity [3]. Malaria occurs as parasites undergo cycles

of invasion and asexual replication inside erythrocytes. Towards the end of each cycle

schizonts develop which contain multiple merozoites that need to burst from the host cell

and invade a new erythrocyte. The transcriptional program is highly synchronised

throughout the asexual parasite replication cycle in erythrocytes [4], but some variation

exists between parasite clones [5,6]. Analysis of naturally occurring sequence polymorphism

in *Plasmodium falciparum* has shown that selection maintains multiple alleles of many

merozoite-stage genes [7,8], some of which encode targets of naturally acquired immunity

[9-11]. Genes expressed at this stage can also exhibit pronounced variation in transcription

among parasites [7,12-14], although the extent of this variation among laboratory-adapted

parasite lines and clinical isolates has not been deeply surveyed.

Accurate quantitation of differential gene expression between organisms or groups of

organisms in whole-transcriptome analyses requires biological replicate sampling [15,16].

The importance of minimising sampling error in order to detect real differential gene

expression is well recognised [17], and some bioinformatic tools can guide the

determination of replicate numbers appropriate to experimental designs [18,19]. In

transcriptomic studies of malaria parasites, achieving large numbers of replicate sample

preparations is difficult, particularly from clinical isolates [5], and most current

understanding is derived from a small number of parasite lines that have been culture-

adapted for many years. Examination of genome sequences has identified premature stop

codon mutants affecting some transcriptional factor genes and cell signalling protein genes

[20,21], as well as particular gene deletions and amplifications only observed in parasites

adapted to *in vitro* culture [22,23]. Although culture-adapted parasites are phenotypically

and transcriptionally diverse [6,12,24], it is not clear to what extent they reflect the diversity

of parasites causing clinical malaria. Parasites may be studied in the first cycle of *ex vivo*

growth following isolation from patients, and maturation to schizont stages shows variable

expression of invasion-related genes [7,13,14], but a lack of biological replicate

measurements limits the precision of whole transcriptome analysis of such samples.

Here we present gene expression profiles of schizont-stage malaria parasites from multiple

cultured clinical isolates and laboratory-adapted lines. We conduct RNA-seq analysis with

large numbers of replicates per sample, and show that high numbers of biological replicates

improves the true-positive discovery rate for identifying differentially expressed genes. This

identifies schizont-stage genes that are differentially expressed between laboratory-adapted

and cultured clinical isolates, as well as those that are variably expressed among cultured

clinical isolates. The results confirm variable transcription in particular genes encoding

ligands involved in erythrocyte invasion, and variation in expression of genes that have been

less characterised was quantitatively validated by targeted RT-qPCR assays, allowing analysis

to be extended to additional isolates from which material is more limited. The expression

levels of these genes in a panel of clinical isolates sampled at schizont stage in the first *ex*

vivo cycle were similar on average to those observed in cultured clinical isolates. This

indicates that cultured parasite lines are useful for studying expression of most genes,

although a minority of genes are affected by culture adaptation.

Results

Replicate samples of schizont-stage transcriptomes from *P. falciparum* laboratory-adapted

clones and clinical isolates

Nineteen biological replicate preparations of schizont-stage parasites were made from

cultured *P. falciparum* clone 3D7, and between five and ten biological replicate preparations

were made from each of three other laboratory-adapted clones of diverse origin and six

clinical isolates from Ghana. Transcriptomes were obtained by sequencing of oligo-dT-

primed cDNA (RNA-seq), and mapping reads to the 3D7 reference genome sequence. To

minimize potential effects of sequence differences between the different parasites, regions

of genes with high levels of allelic polymorphism were masked, so that for polymorphic

genes only reads mapping to conserved portions were analysed (S1 file). Genes that have

been described as deleted in some parasites were not excluded, as most individual deletions

are very rare and exclusion would result in unnecessary loss of data for genes unaffected in

the isolates sampled here, but we chose to rather filter out cases where deletion was

apparent in analysis. The subtelomeric large gene families *var*, *rifin* and *stevor* were

excluded as short read sequence mapping is not generally locus-specific for these, so 5,134

genes were analysed in total. Mapped RNA-seq reads for each gene were converted to

fragments per kilobase of transcript per million mapped reads (FPKM). Analysis of four

paired replicates of schizont preparations from E64-treated and untreated cultures of clone

3D7 showed that E64 treatment had no major effect, with only a single gene showing $\log_2$

fold difference > 2 (S2 File).

As multiple biological replicates could not have exactly the same distribution of parasite

developmental stages, despite being schizont-enriched, to assess the extent of earlier stage

transcripts the distributions of FPKMs for each replicate were correlated with FPKM values

in reference RNA-seq data for seven cultured time points over a ~48 hour *P. falciparum*

cycle [25]. This identified only a small number of replicates that did not have a maximum

Spearman's correlation with parasites at either 40 or 48 hours post-invasion, and these

were excluded from further analyses (Fig 1). Within each isolate, pairwise correlations of

FPKMs among replicates correlated highly, with average Spearman's ρ > 0.95 (Fig 2).

Numbers of biological replicates improve detection of differentially expressed genes

We assessed the impact of increasing numbers of biological replicates on detection of genes

with differing transcript levels between parasite lines. Data were first compared for the 3D7

and D10 laboratory clones, which each had material from schizonts purified by density

gradient centrifugation from ten independent cultures. Comparing all ten replicates of 3D7

and D10 identified 123 genes with $\log_2$ differences of > 2 in relative transcript levels

between the two clones (equivalent to at least 4-fold differences). Achieving this high

number of biological replicate samples is difficult, so we assessed what proportion of these

123 genes were captured as being differentially expressed in 100 random samples of two,

four, six and eight replicates within each group. As expected, the true-positive proportion of

differentially expressed genes detected (using the same statistical criteria of absolute $\log_2$

fold difference > 2) increased with the number of replicates within each group, median true-

positive rates being 0.28, 0.52, 0.67 and 0.79, for two, four, six and eight replicates

respectively (Fig 3). While the true-positive rate of genes detected increased with the

number of replicates, the median false-positive rates were very low in all cases (among the

genes that did not show differences with 10 replicates, the proportions showing apparent

differences with two, four, six and eight replicates were 0.001, 0.004, 0.005 and 0.004

respectively).

We also examined the ability to detect differential expression when focusing specifically on

genes that are highly expressed in schizont-stage samples, as these are the genes most likely

to be functionally important for the merozoite invasive stage. The FPKM values for each

sample in each of the 100 comparisons were averaged and the genes with non-zero FPKM

values were ranked by their maximum expression level in either strain. Then, the

differentially expressed genes that fell within the top quartile of most highly expressed

genes in the subsampled data were compared to those identified as differentially expressed

among the top quartile of genes in the full analysis. This yielded median true-positive rates

of 0.45, 0.63, 0.78 and 0.86 for comparisons respectively containing two, four, six and eight

replicates, higher proportions than for the analysis of all genes. Based on these data, it is

evident that the use of six independent biological replicates for parasite RNA-seq will detect

most genes that are differentially expressed, and that slightly fewer replicates are adequate

if analyses only need to focus on highly expressed genes.

Comparison of gene expression in cultured clinical and laboratory-adapted isolates.

Comparing the transcriptomes of long-term laboratory-adapted and cultured clinical

isolates, 134 genes (2.6 % of those analysed) had different transcript levels between the two

groups (S1 Table). Among the genes within the top quartile of expression values overall, 20

genes (1.6 %) were differentially expressed between the laboratory-adapted and clinical

isolate groups, a lower proportion than among the rest of the genes (Odds Ratio 0.52,

Fisher's exact P = 0.006). It is possible that there is a higher background of false positive

differences for the less expressed genes, due to lower read coverage giving greater noise in

the relative ratios. Examining the 20 genes with differing transcript levels among the top

quartile of expressed genes, several were seen to contain a deletion in one of the parasite

lines or to be members of multigene families. For example, PF3D7_1371600 (*ebl-1*) and

PF3D7_0935700 are deleted in the HB3 strain and D10 strain respectively (Fig 4), as

previously documented [26,27]. After excluding these genes and others for which one or

more of the samples were presumed to contain genetic deletions, and excluding members

of sub-telomeric multi-gene families, ten single-locus genes were identified as showing

differential expression between laboratory and clinical isolates (Table 1).

Eight of the ten highly expressed genes showing a difference between the groups had higher

transcript levels in the long-term laboratory-adapted clones than in the cultured clinical

isolates (Table 1). These include an AP2 transcription factor gene PF3D7_0420300, as well as

two predicted methyltransferase genes (PF3D7_0422900 and PF3D7_0522300), and gene

PF3D7_0104300 encoding ubiquitin-binding protein 1 which is involved in protein turnover

[28] (Fig 4). The only two genes that had higher transcript levels in cultured clinical isolates

were PF3D7_1036300 which encodes the merozoite surface protein MSPDBL2, and

PF3D7_1302100 which encodes the gamete antigen 27/25 (Fig 4). Previous data has shown

MSPDBL2 to be variably expressed among *P. falciparum* isolates [7], and experimental

analysis has shown transcript levels to be increased in parasite cultures when the

heterochromatin protein 1 is suppressed by gametocyte development protein 1 (GDV1)

leading to gametocytogenesis [29,30], whereas gamete antigen 27/25 is a known marker of

early gametocytogenesis [30,31]. Gametocyte gene transcripts were not generally more

highly expressed in the clinical isolates, and gene PF3D7_1327300 that has been previously

described to be transcribed in gametocytes [32] showed lower expression.

We investigated whether there was an overall association between the genes differentially

expressed and targets of HP1 regulation [29,33]. Of the HP1-regulated genes previously

identified [29], 118 genes are among those analysed here (*var, rifin* and *stevor* gene families

were excluded). Of the 134 genes differentially expressed between the two groups, 24 (17.9

%) were targets of HP1 regulation, a higher proportion than among the rest of the genes

(Odds Ratio 11.3, Fisher's exact P = 1.3 x 10^{-5}). Among 20 differentially expressed genes in

the top quartile of expression values, two (10 %) were targets of HP1 regulation, a higher

proportion than among the remaining genes in the top quartile (Odds Ratio 7.6, Fisher's

exact P = 0.037).

Differential expression of schizont-stage genes among cultured clinical isolates

Overall, 214 genes (4.2 %) showed a $\log_2$ fold difference > 2 for at least one of the fifteen

pairwise comparisons among cultured clinical isolates. Of those genes in the top quartile of

expression levels, 39 genes (3.0 %) showed a difference for at least one pairwise

comparison, a lower proportion than among the less expressed genes (Odds Ratio 0.66,

Fisher's Exact P = 0.02) (S2 Table). Some of the variably expressed genes encode merozoite

proteins previously characterised as transcriptionally variable among *ex vivo* clinical isolates

[14], including *dblmsp2* and *msp6*, as well as erythrocyte binding antigen genes *eba-175*,

eba-181, and *eba-140* (which is also deleted in one of the laboratory adapted clones) (S1

Fig).

There was an over-representation of GDV1-regulated genes [30] among the 214 genes

differentially expressed in cultured clinical isolate comparisons (Fisher's exact test odds ratio

12.5, Fisher's Exact P = 1.9×10^{-9}). Of the genes differentially expressed among cultured

clinical isolates, 64 % were also differentially expressed in comparisons among the long-

term laboratory-adapted clones (Odds Ratio 17.4, Fisher's exact P = 2.2×10^{-16}). There was

also strong overlap when only considering genes in the top quartile of expression overall, 44

% of genes differentially expressed among cultured clinical isolates also being differentially

expressed among laboratory-adapted lines (Odds Ratio 7.9, Fisher's exact P = 2.4×10^{-8}).

To validate the data obtained through RNA-seq, a subset of genes was selected for

quantitation by reverse-transcription quantitative PCR (RT-qPCR). We chose differentially

expressed genes encoding proteins containing predicted transmembrane domains or signal

peptides, thereby likely to enter the parasite secretory pathway, excluding genes previously

characterised as variably expressed or members of multi-gene families (S2 Table). Eight

genes were selected for assay (S2 Fig), two of which encode known antigens (merozoite-

associated tryptophan-rich antigen [34] liver-stage antigen-3 [35]). These gene transcripts

were quantified by RT-qPCR in 58 RNA preparations from the laboratory clones and clinical

isolates that had been analysed by RNA-seq. Transcript levels were normalised to those of a

house-keeping gene (serine-tRNA ligase PF3D7_0717700) and correlated with similarly

normalised FPKM expression values for the RNA-seq data. Each gene showed strong positive

and highly significant correlation between RNA-seq and RT-qPCR-derived expression

measures, all except one having correlation coefficients > 0.8 (Fig 5).

Variable expression in first *ex vivo* cycle schizonts

We next determined the expression levels in additional clinical samples from Ghana

matured during the first cycle *ex vivo* until a large proportion of parasites were schizonts.

For nine of these isolates there was sufficient material for RNA-seq to be performed, and

results were analysed for seven of these (two isolates were excluded from analysis as

whole-transcriptome staging of FPKM values showed maximal correlation with stages earlier

than schizonts in a reference time-course). For six isolates grown *ex vivo* to schizont stage

there was sufficient material for quantification of the selected eight gene panel by RT-qPCR.

Characterisation of the *ex vivo* samples using either method showed considerable variation

in relative expression levels of each gene, but the mean levels of each gene across the

isolates were similar to those determined in the cultured parasites that had extensive

sample replication (Fig 6). Without sample replication of the first cycle *ex vivo* clinical

samples, the contribution of random sampling variation and uncontrolled effects on the

expression levels cannot be accounted for, but the observation that for each gene the

normalised expression values have a similar mean to that for cultured clinical lines indicates

no systematic differences in expression between the initial *ex vivo* cycle and under

continuous culture conditions. These results support the use of cultured clinical isolates as a

means of studying parasite gene expression to reflect that occurring in original populations,

while enabling the necessary experimental replication for many analyses.

Discussion

Transcriptomic analyses perform optimally with increased levels of sample replication to

minimise the impact of stochastic or technical variation in individual samples [17,36]. This is

particularly important for studies that focus on a defined developmental stage, in which

error can be generated due to subtle differences between samples. However, the nature of

clinical samples of malaria parasites is such that replication is technically very difficult. In

order to assess the degree of replication useful to robustly identify differentially expressed

genes, we first generated RNA-seq profiles from multiple replicates of laboratory-adapted

clones, and used the data to calculate the true- and false-discovery rates obtained using

fewer replicates. Consistent with studies on other cultured eukaryotes [17], we show that

for malaria parasites increasing the sample replication improves the true-positive discovery

rate in identifying variably expressed genes, and indicate that six independent replicates are

useful to balance accuracy and experimental feasibility. Where these numbers of replicates

are not obtainable, it is recommended to focus on genes that are highly expressed in order

to achieve as much accuracy as possible per gene.

It has been proposed that spontaneous transcriptional variation within parasite populations

is a strategy that ensures fitness of parasites facing a range of potential and changing

selective pressures [6], and it is likely that the rates and mechanisms of regulation differ

among most genes. In comparison between long-term laboratory-adapted clones and

cultured clinical isolates, we found that among the genes most differentially expressed were

an AP2-transcription factor [37] and two methyltransferases, the functional roles of which

remain to be investigated. Most of the differentially expressed genes had lower

transcription in the clinical isolates, suggesting that these may be normally repressed. HP1 is

responsible for extensive gene silencing in malaria parasites [30], and we showed an over

representation of HP1-regulated genes among those differentially expressed between

laboratory-adapted clones and cultured clinical isolates. It would be interesting to

undertake further studies directly on HP1 and other chromatin modifications in recent

clinical isolates, as experimental studies have so far focused on long-term laboratory-

adapted clones. Interestingly, among the top quartile of genes expressed, the only two

genes that were more highly expressed in clinical isolates were an early marker of

gametocytogenesis (*Pfs25/27*) and a merozoite surface protein gene (*mspdbl2*) encoding a

product expressed all merozoites within only a minority of schizonts [7] which may be a

marker of parasites committed to sexual development [29]. Studies of gametocytogenesis in

clinical isolates will be important to understand the process of commitment to sexual

differentiation and transmission, as also indicated by transcriptome comparisons that

sampled different stages without having multiple replicates for each stage [5].

Through pairwise comparisons of individual cultured clinical isolates, we identified many

genes that were differentially expressed among clinical isolates. Some of the genes

identified have previously been shown to be differentially expressed through targeted qRT-
PCR assays of schizont-stage *ex vivo* cultured clinical isolates [7,13,14]. Focusing on genes
for which variable expression has not been previously studied, we selected a panel of eight
highly differentially expressed genes that encode secreted proteins. Of these, the
merozoite-associated tryptophan-rich antigen and liver stage antigen 3 have been
previously identified as merozoite proteins [34,35], and another gene (PF3D7_0423900) is
adjacent to loci encoding the cysteine-rich protective antigen (CyRPA) and reticulocyte
binding homologue 5 (Rh5) which are targets of antibodies that inhibit merozoite invasion.
We extensively validated the RNA-seq data by RT-qPCR for these genes, and extended
quantitation to additional *ex vivo* clinical isolate samples. Despite the lack of sample
replicates for the latter, results showed that average expression levels for each gene were
similar to those for the cultured samples that had biological replicates.
324
Together, these data highlight new candidates for investigation as potential determinants of
alternative developmental pathways or targets of immunity. Characterisation of gene
transcript levels is only one level of analysis, as proteomic characterisation [38] and analysis
of phosphorylation and other protein modifications [39,40] may also identify functional
determinants, and large-scale analysis of parasite protein variation has not yet been
attempted on clinical isolates. A more complete description of transcriptome variation may
also be derived in future by analysis of individual infected cells, which has not been
undertaken on parasites from clinical isolates, as initial single-cell sequencing studies of
laboratory model malaria parasites have only recently begun to probe the expression
pathways of different life-cycle stages [41-43]. Scaling up transcriptomic analyses to clinical
studies is challenging, and biological replicate preparations of parasite samples may only be
feasible in a limited number of cases. In an epidemiological context, reduction of error also
requires large sample sizes and tight definition of clinical phenotypes, and more statistical
analysis and modelling of covariates may be appropriate, ideally including data on host
transcriptome variation within the same samples [44].
340
Methods
Ethical approval. The study of clinical parasite transcriptomes was approved by the Ethics
committees of the Ghana Health Service, the Noguchi Memorial Institute for Medical

Research, University of Ghana, the Kintampo Health Research Centre, the Navrongo Health

Research Centre and the London School of Hygiene and Tropical Medicine. Written

informed consent was obtained from parents or legal guardians of all participating children,

and additional assent was received from children over 10 years old.

P. falciparum isolates sampled from clinical malaria cases. Blood samples were collected

from clinical malaria cases attending Ghana government health facilities between 2012 and

2013, in Kintampo (Brong-Ahafo Region of central Ghana), and Navrongo (Kassena-Nankana

East Municipality, in the Upper East Region of northern Ghana). Patients were eligible to

participate in the study if they had uncomplicated clinical malaria, were aged 2–14 years,

tested positive for _P. falciparum_ malaria by lateral flow rapid diagnostic test and slide

microscopy, and had not taken antimalarial drugs during the 72 hours preceding sample

collection. Venous blood samples (up to 5 ml) were collected into heparinized Vacutainer

tubes (BD Biosciences). Blood samples were centrifuged, plasma separated and leukocyte

buffy coat cell layer removed, and erythrocytes were cryopreserved in glycerolyte and

stored frozen at −80°C or in liquid nitrogen until shipment on dry ice to the London School

of Hygiene and Tropical Medicine.

Parasite culture and enrichment for schizont stages. Parasites from _ex vivo_ clinical samples

were matured within the original patient erythrocytes at 2–5 % haematoctit for up to 48 h in

RPMI 1640 medium containing 2 % human AB serum (GE Healthcare) and 0.3 % Albumax II

(Thermo Fisher Scientific) under an atmosphere of 5 % O_2, 5 % CO_2, and 90 % N_2 at 37 °C,

until most parasites were at the schizont stage of parasite development, at which point RNA

was extracted directly from the bulk cultures. Laboratory-adapted clones and cultured

clinical isolates were grown in erythrocytes at 2–5 % haematocrit in RPMI 1640 medium

containing 0.5 % Albumax II, at 37 °C. Laboratory-adapted clones were maintained under

atmospheric air with 5 % CO_2, and cultured clinical isolates were maintained in 5 % O_2, 5 %

CO_2, 90 % N_2. Schizonts were isolated by either MACS or discontinuous density

centrifugation methods.

For MACS purification of cultured clinical isolates, parasites were prepared from 100 ml

cultures with at least 0.7 % of erythrocytes containing schizonts. Parasites were isolated by

magnetic purification using magnetic LD Separation columns (Miltenyi Biotech). One column

was used per 25 ml culture, and washed twice in 3 ml of culture medium at room

temperature. Parasite culture was pelleted at 500 g for 5 minutes, then re-suspended in 3

ml culture medium per 1 ml of packed cell volume. The re-suspended material was bound to

the MACS column, which was then washed three times with 3 ml culture medium, and

schizonts were eluted twice by removing the magnet from the column and forcing 2 ml

culture medium through the column into a 15 ml collection tube. Finally, the schizonts were

pelleted by centrifugation at 500 g for 5 minutes and the pellet volume was estimated, with

0.5 µl used for Giemsa-stained microscopical examination to assess staging, 1 µl added back

to 250 µl culture at 0.8 % hematocrit to follow the progression. Remaining parasites were

re-suspended in 1.5 ml of culture medium with 10 µM E64 in a 12 well plate, and parasites

were incubated for 5.5 hours in 5 % CO_2 at 37 °C, before centrifugation in a 1.5 ml tube and

processing for RNA extraction.

For discontinuous density centrifugation purification, parasites were maintained as 25 ml

cultures at 2.5 % hematocrit. Cultures were used when > 1 % erythrocytes contained

parasites with multiple nuclei. Schizonts were purified using 70 % Percoll (GE

Healthcare)/2.93 % sorbitol/PBS overlayed with 35 % Percoll/1.47 % sorbitol/PBS, which was

overlayed with parasitized erythrocytes, allowing the schizonts to be separated by

centrifugation at 2500 g for 10 minutes at 24 °C, with a light brake at the end of

centrifugation. Purified schizonts were washed once in complete medium and the pellet

volume was estimated. Six pellet volumes of 50 % haematocrit erythrocytes were added to

the pellet and resuspended for slide examination. The sample was smeared, and

resuspended in 6 ml of complete culture medium. Of this, 1 ml was used as a control

untreated sample to track parasite egress. E64 was added to the remaining 5 ml at a final

conentraction of 10 µM. Control and E64-treated samples were placed at 37 °C, 5 % CO_2

static incubator for 5.5 hours. Schizonts from the E64-treated culture were overlaid on 70 %

Percoll/2.93 % sorbitol/PBS and separated by centrifugation at 2500 g for 10 minutes at 24

°C, with a light brake at the end of centrifugation. The schizont layer was washed once in

complete culture medium and final cell pellets of approximately 10 - 20 µl were used for

RNA extraction.

RNA extraction. Pellets were resuspended 500 µl in TRIzol® reagent (Thermo Fisher

Scientific) and stored at -80 °C until RNA extraction following manufacturer's instructions.

Final RNA pellets were resuspended in 100 µl RNase-free H_2O. A second RNA clean-up and

on-column DNase treatment was carried out using RNeasy mini columns (Qiagen), with RNA

eluted in 30-50 µl RNase-free H_2O and concentration quantified by Qubit High Sensitivity

RNA Assay (Thermo Fisher Scientific). Samples containing at least 500 ng RNA were

considered for analysis after the RNA integrity was checked on an Agilent Bioanalyzer using

RNA 6000 Nano reagents and chips (Agilent Genomics).

RNA-seq library preparation and sequencing. Laboratory isolate and cultured clinical

isolate RNA-seq libraries were prepared using TruSeq Stranded mRNA Library Prep Kit

(Illumina) using 500 ng − 1 µg RNA as per the Illumina TruSeq Stranded mRNA protocol for

MiSeq sequencers. Libraries were validated on an Agilent Bioanalyzer using DNA 1000

reagents and chips (Agilent Genomics) to quantify library sizes and confirm the absence of

primer dimers. Libraries were quantified using a KAPA Universal Library Quantification kit

(Roche Diagnostics Limited) on a 7500 Fast Real-Time PCR System (Thermo Fisher Scientific)

and library concentrations were adjusted for library size. 12 − 15 pM pooled libraries were

sequenced on a MiSeq System (Illumina) using a MiSeq Reagent Kit v3 (Illumina) with 2 x 75

cycles.

Ex vivo schizont-enriched *P. falciparum* isolate RNA-seq libraries were prepared using a

modified protocol. PolyA+ RNA (mRNA) was selected using magnetic oligo-d(T) beads.

mRNA was reverse transcribed using Superscript III® (Thermo Fisher Scientific), primed using

oligo-d(T) primers. dUTP was included during second-strand cDNA synthesis. The resulting

double stranded cDNA was fragmented using a Covaris AFA sonicator. Sheared double

stranded cDNA was dA-tailed, end repaired, and "PCR-free" barcoded sequencing adaptors

(Bioo Scientific) [45] were ligated (NEB). Libraries were cleaned up twice, using solid phase

reversible immobilisation beads, and eluted in EB buffer (Qiagen). Second strand cDNA was

removed using uracil-specific excision reagent enzyme mix (NEB). Libraries were quantified

by quantitative PCR prior to sequencing on an Illumina MiSeq sequencer.

Data curation and analysis. Raw Illumina sequence reads were aligned to the *P. falciparum*

3D7 v3 genome and converted to 'bam' format using samtools [46]. Reads with MAPQ

scores < 60 were removed. Reads were counted using the "summarizeOverlaps" feature of

the GenomicAlignments package [47] in R, against a previously published *P. falciparum*

genome annotation file that had been masked for regions of polymorphism (detailed in S2

File) to remove known regions of genomic polymorphism, highly polymorphic gene families

and duplicated genes.

Fragments Per Kilobase of transcript per Million mapped reads (FPKMs) for the data here

and those of previously published data on a developmental time course of the clone 3D7

[25] were calculated using the 'fpkm' function of DESeq2 [48] in R. FPKMs for all genes in

each of our parasite preparations were correlated using a Spearman's Rank correlation with

FPKMs for each of the seven time points (0, 8, 16, 24, 32, 40 and 48 hours post-invasion; S1

Table). Replicates with peak Spearman correlation values of $\rho > 0.7$ at the latest time points

(40 or 48 hours) were included for further analysis. Differential expression analysis was

conducted using DESeq2 in R.

RT-PCR assays. 150 – 500 ng total RNA from each preparation of parasite schizonts was

reverse transcribed using Superscript II® (Thermo Fisher Scientific) with 250 ng random

hexamer oligonucleotide primers per 20 μl reaction. Quantitative PCR (qPCR) was carried

out using SYBR® Select Master Mix (Thermo Fisher Scientific) with 500 nM forward and

reverse primers, in a Prism 7500 Fast qPCR machine. For each gene, threshold-cycle values

were quantified against a serially diluted genomic DNA (Dd2 strain) standard curve, run on

the same plate. Cycling parameters were: 50°C for 2 minutes, 95°C for 2 minutes followed

by 40 cycles of 95 °C for 15 seconds and 60 °C for 1 minute. All wells were run as 10 μl

volumes in technical duplicate. The qPCR copy numbers were normalised against copies of a

house-keeping gene, PF3D7_0717700 [49]. PCR primer pair sequences are as follows:

PF3D7_0102700 5'-CAACCAGACAAACAACAAGAAA-3' and 5'-TCCATTCTGATGCTTTCCAC-3',

PF3D7_0220000 5'-GTAAATGGTGAATTAGCTAGTGAAG-3' and 5'-

CCTTTATATCTTCGGCTTCTTCT-3', PF3D7_0423900 5'-GAGAAGAAGCCAAAGTAAATGAAC-3'

and 5'-GAATGTGTCAAGTGCATCATAA-3', PF3D7_0605900 5'-CGCAATAACAAGAAGTCACAAA-

3' and 5'-AAGATTGTAGGAGGGTGTTAAT-3', PF3D7_0935300 5'-GGGCTGCTGTTATACCTTG-3'

and 5'-AGAATGAGGAGGTTCAGTTTG-3', PF3D7_1252900 5'-

CCTTAGTAGAACTTCAAAGTGACA-3' and 5'-TGTAACCAACTACGAAATTCCC-3',

PF3D7_1461700 5'-TGCTGACTTTCAAGAGATAAGG-3' and 5'-

TTTCATTTGTTCATTTTGTACAACC-3', PF3D7_1476500 5'-CTTCGATTCACAAATGCAGAAA-3' and

5'-CGTATTCCACACCTTGTTCTAT-3', PF3D7_0717700 5'-AAGTAGCAGGTCATCGTGGTT-3' and

5'-GTTCGGCACATTCTTCCATAA-3'.

Data open access

475   All *P. falciparum* RNA-seq data from the laboratory-adapted and cultured clinical isolates

476   have been submitted for open access at the Gene Expression Omnibus

477   (https://www.ncbi.nlm.nih.gov/geo/), entry: GSE113718. The *P. falciparum* RNA-seq data

478   from the first cycle e*x vivo* isolates have been submitted to the European Nucleotide

479   Accession Archive (https://www.ebi.ac.uk/ena), study: ERP103955.

Acknowledgements

482   We are grateful to malaria patients who contributed blood samples for analysis of parasite

483   transcription, and to colleagues who supported the clinical sample collection and initial

484   laboratory processing. We thank Alistair Miles and Antoine Claessens for help with

485   identification of genomic coordinates containing highly polymorphic sequences within

486   genes.

References

1. Mata J, Wilbrey A, Bahler J (2007) Transcriptional regulatory network for sexual differentiation in
fission yeast. Genome Biol 8: R217.
2. Cases I, de Lorenzo V, Ouzounis CA (2003) Transcription regulation and environmental adaptation
in bacteria. Trends Microbiol 11: 248-253.
3. Cestari I, Stuart K (2018) Transcriptional Regulation of Telomeric Expression Sites and Antigenic
Variation in Trypanosomes. Curr Genomics 19: 119-132.
4. Bozdech Z, Llinás M, Pulliam BL, Wong ED, Zhu J, et al. (2003) The transcriptome of the
intraerythrocytic developmental cycle of *P. falciparum*. PLoS Biology 1: e5.
5. Mackinnon MJ, Li J, Mok S, Kortok MM, Marsh K, et al. (2009) Comparative transcriptional and
genomic analysis of *Plasmodium falciparum* field isolates. PLoS Pathog 5: e1000644.
6. Rovira-Graells N, Gupta AP, Planet E, Crowley VM, Mok S, et al. (2012) Transcriptional variation in
the malaria parasite *Plasmodium falciparum*. Genome Res 22: 925-938.
7. Amambua-Ngwa A, Tetteh KK, Manske M, Gomez-Escobar N, Stewart LB, et al. (2012) Population
genomic scan for candidate signatures of balancing selection to guide antigen
characterization in malaria parasites. PLoS Genet 8: e1002992.
8. Ochola LI, Tetteh KK, Stewart LB, Riitho V, Marsh K, et al. (2010) Allele frequency-based and
polymorphism-versus-divergence indices of balancing selection in a new filtered set of
polymorphic genes in *Plasmodium falciparum*. Mol Biol Evol 27: 2344-2351.
9. Osier FH, Mackinnon MJ, Crosnier C, Fegan G, Kamuyu G, et al. (2014) New antigens for a
multicomponent blood-stage malaria vaccine. Science Translational Medicine 6: 247ra102-
247ra102.

10. Richards JS, Arumugam TU, Reiling L, Healer J, Hodder AN, et al. (2013) Identification and
prioritization of merozoite antigens as targets of protective human immunity to *Plasmodium*
falciparum malaria for vaccine and biomarker development. J Immunol 191: 795-809.
11. Tetteh KK, Osier FH, Salanti A, Kamuyu G, Drought L, et al. (2013) Analysis of antibodies to newly
described *Plasmodium falciparum* merozoite antigens supports MSPDBL2 as a predicted
target of naturally acquired immunity. Infect Immun 81: 3835-3842.
12. Stubbs J, Simpson KM, Triglia T, Plouffe D, Tonkin CJ, et al. (2005) Molecular mechanism for
switching of *P. falciparum* invasion pathways into human erythrocytes. Science 309: 1384-
1387.
13. Gomez-Escobar N, Amambua-Ngwa A, Walther M, Okebe J, Ebonyi A, et al. (2010) Erythrocyte
invasion and merozoite ligand gene expression in severe and mild *Plasmodium falciparum*
malaria. J Infect Dis 201: 444-452.
14. Bowyer PW, Stewart LB, Aspeling-Jones H, Mensah-Brown HE, Ahouidi AD, et al. (2015) Variation
in *Plasmodium falciparum* erythrocyte invasion phenotypes and merozoite ligand gene
expression across different populations in areas of malaria endemicity. Infect Immun 83:
2575-2582.
15. Hansen KD, Wu Z, Irizarry RA, Leek JT (2011) Sequencing technology does not eliminate biological
variability. Nat Biotechnol 29: 572-573.
16. Elowitz MB, Levine AJ, Siggia ED, Swain PS (2002) Stochastic gene expression in a single cell.
Science 297: 1183-1186.
17. Schurch NJ, Schofield P, Gierlinski M, Cole C, Sherstnev A, et al. (2016) How many biological
replicates are needed in an RNA-seq experiment and which differential expression tool
should you use? RNA 22: 839-851.
18. Seo J, Gordish-Dressman H, Hoffman EP (2006) An interactive power analysis tool for microarray
hypothesis testing and generation. Bioinformatics 22: 808-814.
19. Busby MA, Stewart C, Miller CA, Grzeda KR, Marth GT (2013) Scotty: a web tool for designing
RNA-Seq experiments to measure differential gene expression. Bioinformatics 29: 656-657.
20. Claessens A, Affara M, Assefa SA, Kwiatkowski DP, Conway DJ (2017) Culture adaptation of
malaria parasites selects for convergent loss-of-function mutants. Sci Rep 7: 41303.
21. Kafsack BF, Rovira-Graells N, Clark TG, Bancells C, Crowley VM, et al. (2014) A transcriptional
switch underlies commitment to sexual development in malaria parasites. Nature 507: 248-
252.
22. Cheeseman IH, Gomez-Escobar N, Carret CK, Ivens A, Stewart LB, et al. (2009) Gene copy number
variation throughout the *Plasmodium falciparum* genome. BMC Genomics 10: 353.
23. Jeffares DC, Pain A, Berry A, Cox AV, Stalker J, et al. (2007) Genome variation and evolution of
the malaria parasite *Plasmodium falciparum*. Nat Genet 39: 120-125.
24. Llinas M, Bozdech Z, Wong ED, Adai AT, DeRisi JL (2006) Comparative whole genome
transcriptome analysis of three *Plasmodium falciparum* strains. Nucleic Acids Res 34: 1166-
1173.
25. Otto TD, Wilinski D, Assefa S, Keane TM, Sarry LR, et al. (2010) New insights into the blood-stage
transcriptome of *Plasmodium falciparum* using RNA-Seq. Mol Microbiol 76: 12-24.
26. Peterson DS, Wellems TE (2000) EBL-1, a putative erythrocyte binding protein of *Plasmodium*
falciparum, maps within a favored linkage group in two genetic crosses. Mol Biochem
Parasitol 105: 105-113.
27. Nacer A, Roux E, Pomel S, Scheidig-Benatar C, Sakamoto H, et al. (2011) Clag9 is not essential for
PfEMP1 surface expression in non-cytoadherent *Plasmodium falciparum* parasites with a
chromosome 9 deletion. PLoS One 6: e29039.
28. Henriques G, Hallett RL, Beshir KB, Gadalla NB, Johnson RE, et al. (2014) Directional selection at
the pfmdr1, pfcrt, pfubp1, and pfap2mu loci of *Plasmodium falciparum* in Kenyan children
treated with ACT. J Infect Dis 210: 2001-2008.

29. Filarsky M, Fraschka SA, Niederwieser I, Brancucci NMB, Carrington E, et al. (2018) GDV1 induces
sexual commitment of malaria parasites by antagonizing HP1-dependent gene silencing.
Science 359: 1259-1263.
30. Brancucci NMB, Bertschi NL, Zhu L, Niederwieser I, Chin WH, et al. (2014) Heterochromatin
protein 1 secures survival and transmission of malaria parasites. Cell Host Microbe 16: 165-
176.
31. Carter R, Graves PM, Creasey A, Byrne K, Read D, et al. (1989) *Plasmodium falciparum*: an
abundant stage-specific protein expressed during early gametocyte development. Exp
Parasitol 69: 140-149.
32. Lu XM, Batugedara G, Lee M, Prudhomme J, Bunnik EM, et al. (2017) Nascent RNA sequencing
reveals mechanisms of gene regulation in the human malaria parasite *Plasmodium*
falciparum. Nucleic Acids Res 45: 7825-7840.
33. Fraschka SA, Filarsky M, Hoo R, Niederwieser I, Yam XY, et al. (2018) Comparative
heterochromatin profiling reveals conserved and unique epigenome signatures linked to
adaptation and development of malaria parasites. Cell Host Microbe 23: 407-420 e408.
34. Ntumngia FB, Bouyou-Akotet MK, Uhlemann AC, Mordmuller B, Kremsner PG, et al. (2004)
Characterisation of a tryptophan-rich *Plasmodium falciparum* antigen associated with
merozoites. Mol Biochem Parasitol 137: 349-353.
35. Morita M, Takashima E, Ito D, Miura K, Thongkukiatkul A, et al. (2017) Immunoscreening of
Plasmodium falciparum proteins expressed in a wheat germ cell-free system reveals a novel
malaria vaccine candidate. Sci Rep 7: 46086.
36. Robinson MD, Smyth GK (2007) Moderated statistical tests for assessing differences in tag
abundance. Bioinformatics 23: 2881-2887.
37. Balaji S, Babu MM, Iyer LM, Aravind L (2005) Discovery of the principal specific transcription
factors of Apicomplexa and their implication for the evolution of the AP2-integrase DNA
binding domains. Nucleic Acids Research 33: 3994-4006.
38. Kumar K, Srinivasan P, Nold MJ, Moch JK, Reiter K, et al. (2017) Profiling invasive *Plasmodium*
falciparum merozoites using an integrated omics approach. Sci Rep 7: 17146.
39. Doerig C, Rayner JC, Scherf A, Tobin AB (2015) Post-translational protein modifications in malaria
parasites. Nat Rev Microbiol 13: 160-172.
40. Jones ML, Collins MO, Goulding D, Choudhary JS, Rayner JC (2012) Analysis of protein
palmitoylation reveals a pervasive role in *Plasmodium* development and pathogenesis. Cell
Host Microbe 12: 246-258.
41. Poran A, Notzel C, Aly O, Mencia-Trinchant N, Harris CT, et al. (2017) Single-cell RNA sequencing
reveals a signature of sexual commitment in malaria parasites. Nature 551: 95-99.
42. Reid AJ, Talman AM, Bennett HM, Gomes AR, Sanders MJ, et al. (2018) Single-cell RNA-seq
reveals hidden transcriptional variation in malaria parasites. Elife 7.
43. Walzer KA, Kubicki DM, Tang X, Chi JT (2018) Single-cell analysis reveals distinct gene expression
and heterogeneity in male and female *Plasmodium falciparum* gametocytes. mSphere 3.
44. Lee HJ, Georgiadou A, Otto TD, Levin M, Coin LJ, et al. (2018) Transcriptomic studies of malaria: a
paradigm for investigation of systemic host-pathogen interactions. Microbiol Mol Biol Rev
82.
45. Kozarewa I, Ning Z, Quail MA, Sanders MJ, Berriman M, et al. (2009) Amplification-free Illumina
sequencing-library preparation facilitates improved mapping and assembly of (G+C)-biased
genomes. Nat Methods 6: 291-295.
46. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, et al. (2009) The Sequence Alignment/Map
format and SAMtools. Bioinformatics 25: 2078-2079.
47. Lawrence M, Huber W, Pages H, Aboyoun P, Carlson M, et al. (2013) Software for computing and
annotating genomic ranges. PLoS Comput Biol 9: e1003118.
48. Love MI, Huber W, Anders S (2014) Moderated estimation of fold change and dispersion for RNA-
seq data with DESeq2. Genome Biol 15: 550.

49. Salanti A, Staalsoe T, Lavstsen T, Jensen AT, Sowa MP, et al. (2003) Selective upregulation of a
single distinctly structured var gene in chondroitin sulphate A-adhering *Plasmodium*
falciparum involved in pregnancy-associated malaria. Mol Microbiol 49: 179-191.

Table 1. *P. falciparum* genes in the top quartile of expression showing a difference in

schizont stage transcript levels between long-term laboratory-adapted clones and recent

clinical isolates in culture

Gene ID	Product annotation	Log$_2$ fold difference	Wald test
Higher transcription in long-term laboratory-adapted clones			
PF3D7_0422900	methyltransferase, putative	-2.50	-6.48
PF3D7_0501300	skeleton-binding protein 1	-2.50	-4.98
PF3D7_0410000	erythrocyte vesicle protein 1	-2.38	-10.62
PF3D7_0104300	ubiquitin carboxyl-terminal hydrolase 1, putative	-2.18	-6.13
PF3D7_0522300	18S rRNA (guanine-N(7))-methyltransferase, putative	-2.15	-8.67
PF3D7_0420300	AP2 domain transcription factor, putative	-2.02	-7.19
PF3D7_1030200	claudin-like apicomplexan microneme protein, putative	-2.00	-9.21
PF3D7_1327300	conserved *Plasmodium* protein, unknown function	-2.00	-4.97
Higher transcription in clinical isolates			
PF3D7_1036300	duffy binding-like merozoite surface protein 2	3.02	5.32
PF3D7_1302100	gamete antigen 27/25	2.02	3.99

Figure 1. Multiple replicate biological preparations of *P. falciparum* transcriptomes from

four long-term laboratory adapted clones and six Ghanaian clinical isolates, showing FPKM

values across all genes correlated with previous data from seven time-points across the *P.*

falciparum asexual erythrocytic cycle [25]. Each individual parasite culture was enriched for

schizont-stage parasites, with egress blocked using E64 treatment for 5.5 hours, and

parasites were purified using MACS purification for all of the lines, except for the D10 and

HB3 clones and ten of the 3D7 replicates which were purified using Percoll density gradient

centrifugation. Red lines plot data for samples with peak correlation at either 40 or 48 hours

post-invasion, and grey lines plot replicates that did not maximally correlate with either of

these timepoints and which were subsequently excluded from the analysis.

Figure 2. Distribution of correlations of FPKM expression values among biological sample

replicate transcriptomes for each isolate or clone. The final numbers of replicates, and

median Spearman's correlation coefficients among replicates were: 3D7 n=19, ρ=0.89; D10

n=10, ρ=0.93; Dd2 n=6, ρ=0.95; HB3 n=7, ρ=0.89; 271 n=3, ρ=0.94; 278 n=5, ρ=0.91; 280

n=6, ρ=0.93; 286 n=6, ρ=0.97; 293 n=6, ρ=0.93; 296 n=3, ρ=0.90.

Figure 3. Increased sample replication improves discovery of differentially expressed

genes. Assessment of the proportion of genes captured as being differentially expressed

between two different parasite clones (3D7 and D10), by taking 100 random samples of two,

four, six and eight replicates of each (out of ten initially analysed replicates that identified

123 genes with $\log_2$ differences of > 2 in relative transcript levels between the two clones).

Figure 4 Differential gene transcript levels among preparations of schizont-enriched P.

falciparum cultures of laboratory-adapted clones and clinical isolates. Plots of normalised

RNA-seq read counts for eight of the genes showing differences between four laboratory-

adapted clones and six clinical isolates. Each point shows the value for an independent

biological replicate sample. The two top left panels show examples of genes for which one

of the laboratory-adapted clones had a deletion (HB3 and D10 for the respective genes).

These, and other genes with suspected or known deletions in the sampled parasites, are

excluded from the list of differentially expressed genes in Table 1.

Figure 5. High correlations between RNA-seq and RT-qPCR expression measures. Eight

genes identified as differentially expressed among clinical isolates by RNA-seq were

validated by RT-qPCR for 49 of the independent RNA preparations from the four laboratory-

adapted clones and six cultured clinical isolates under study. Scatter plots for each gene

show the gene of interest (GOI) copies ($\log_2$-transformed) normalised to house-keeping

gene copies (HKG; PF3D7_0717700), correlated with GOI FPKM values ($\log_2$-transformed)

normalised to HKG FPKM values. Spearman's correlation coefficients are shown for each

plot. For PF3D7_1461700, one sample had an FPKM value of 0 which was not plotted but is

included in the calculation of correlation.

Figure 6. Transcript levels of variably expressed genes for first cycle *ex vivo* clinical

samples compared with continuously cultured isolates. For the laboratory and cultured

clinical isolates, each point represents the mean $\log_2$ normalised expression value for

multiple replicates of each sample under study. For the *ex vivo* clinical isolates, each point

represents $\log_2$ normalised expression values for the individual, unreplicated samples under

study. Horizontal bars indicate the mean $\log_2$ normalised expression value for all samples

within the group.

Supplementary Material

S1 Figure. Differential expression of merozoite invasion-related genes among schizonts

from different parasite cultures. Distributions of read counts (normalised to library size) for

eight genes, for replicated laboratory-adapted and clinical isolate samples, showing data

from each replicate culture preparation of each strain.

S2 Figure. Gene expression levels for eight genes differentially expressed among clinical

isolates. Distributions of read counts (normalised to library size) for eight genes, showing

data from each replicate culture preparation of each strain.

S1 Table. Genes differentially expressed between clinical and laboratory lines by absolute

log2 fold > 2 (at least 4 fold difference on average). For genes among the top quartile of

expression values genome-wide (top 18 genes in the table), members of multigene families

and genes in which strain-specific deletions may be responsible for differences are

annotated with asterisks (*).

S2 Table. Log_2 fold changes of genes differentially expressed among pairwise comparisons

of six cultured clinical isolates with multiple schizont preparations of each, among genes

within the top quartile of expression overall.

S1 File. Masking of *P. falciparum* genome annotation file for polymorphic regions of genes.

S2 File. Analysis of paired E64-treated and untreated *P. falciparum* 3D7 schizonts

S3 File. Sequence annotation file Pf3D7.May2015.NoSplice.LSHTM.gtf (see S1 Table for

details)

S4 File. Sequence annotation file GTF_VarRifStev_filtered out.gtf (see S1 Table for details)