1

Native tau structure is disrupted by disease-associated mutations that promote aggregation
4

Kenneth W. Drombosky[1], Dailu Chen[1], DaNae Woodard[1], Levent Sari[3,4], Milo Lin[3,4], Marc I. Diamond[1],

Lukasz A. Joachimiak[1,2,*]

[1] Center for Alzheimer's and Neurodegenerative Diseases, University of Texas Southwestern Medical
Center, Dallas, TX 75235, United States
10

[2] Department of Biochemistry, University of Texas Southwestern Medical Center, Dallas, TX 75235,
United States
13

[3] Green Center for Molecular, Computational, and Systems Biology, University of Texas Southwestern
Medical Center, Dallas, TX 75235, United States
16

[4] Department of Biophysics, University of Texas Southwestern Medical Center, Dallas, TX 75235,
United States
19
20

* To whom correspondence should be addressed: Lukasz.Joachimiak@utsouthwestern.edu

ABSTRACT

Tauopathies are neurodegenerative diseases characterized by intracellular amyloid deposits of tau protein. Missense mutations in *MAPT* cause dominantly inherited tauopathies, but a biophysical mechanism driving its amyloid formation is poorly understood. Many disease-associated mutations localize within tau's repeat domain at inter-repeat interfaces proximal to amyloidogenic sequences, such as ^{306}VQIVYK^{311}. Using crosslink mass spectrometry, intramolecular FRET, *in vitro* peptide systems, simulations, and cell models, we provide evidence that the aggregation prone ^{306}VQIVYK^{311} motif forms metastable compact structures with upstream sequences, which modulates aggregation propensity. Disease-associated mutations, isomerization of critical prolines, or alternative splicing are all sufficient to destabilize local structure and trigger spontaneous aggregation. These local structural rearrangements provide a biophysical framework supporting a model in which perturbations initiate early events in sporadic and genetic tau pathogenesis.

INTRODUCTION

Tauopathies belong to a class of over 20 neurodegenerative diseases in which the protein tau

aggregates in neurons and glia. Tau aggregation correlates strongly with the degree of dementia and

neurodegeneration, especially in Alzheimer's Disease. The structural mechanisms by which disease-

associated mutations, alternative splicing, or other events promote aggregation and pathology is

unknown. Understanding the molecular mechanisms of tau aggregation is critical for identifying

therapeutic targets and diagnostic strategies.

The N-terminal ~200 and C-terminal ~80 residues of tau are largely disordered, rendering this

system refractory to high-resolution structural biology methods[1]. In contrast, tau repeat domain (tau

RD), which spans residues 243 to 365, is predicted to be more structured[2], forms the core of amyloid

fibrils[3], and is the minimal region to propagate strains[4]. Tau RD contains an amyloid motif

(^{306}VQIVYK^{311}) (**Figure 1 A**), which is the basis of conversion between the soluble and insoluble states;

^{306}VQIVYK^{311}-mediated self-assembly is sufficient and necessary to form amyloid *in vitro*[5] and generate

pathology *in vivo*[6], but its detailed mechanism of action is poorly understood.

Recent data from our group has indicated a switch occurs in soluble monomeric tau, which

exists in at least two conformational ensembles: inert monomer (M_i) which does not spontaneously self-

assemble, and seed-competent monomer (M_s) which spontaneously self-assembles into amyloid[7]. M_s

has been purified from multiple sources: recombinant fibrils, Alzheimer's disease brain lysate, or brief

exposure to heparin[7]. Recent solution nuclear magnetic resonance (NMR) methods mapped the

heparin binding site to repeat 2, but how this interaction modulates tau conformation remains unclear[8].

Cryo-EM structures suggest an extended conformation of tau when bound to tubulin[9]. Building on this

observation, double electron-electron resonance experiments indicated an expansion of this region

upon heparin binding[10]. Other work mapping the recruitment of molecular chaperones with tau indicates

many chaperones, including Hsp70, Hsp40, and Hsp90, localize around ^{306}VQIVYK^{311} [11]. Furthermore,

expansion of tau RD appears to promote chaperone binding to the amyloid motif, suggesting that local

conformational changes help recruit factors to suppress aggregation[12]. In contrast, intramolecular

disulfide bridges in tau RD are predicted to form a local loop-like structure spanning repeats 2 and 3

that are incompatible with the formation of amyloid[13].

Further solution NMR data is available for a fragment of tau RD in complex microtubules[14]

where portions of the protein appear to form local contacts with upstream flanking sequences.

Structural models guided by experimental restraints from crosslinking mass spectrometry are consistent

with independent NMR data[14]: ^{306}VQIVYK^{311} has potential to form local structural elements.

We hypothesized that disease-associated mutations may contribute to tau's molecular

rearrangement from inert to an early seed-competent form. Of the missense mutations genetically

linked to tau pathology in humans, many disproportionally occur within tau RD and specifically cluster

near ^{306}VQIVYK^{311} [15] (**Figure 1 A, B**). Disease-associated mutations found near tau's amyloid motif,

such as P301L or P301S, have no definitive biophysical mechanism of action, but are nevertheless

widely used in cell and animal models[16,17]. These observations suggest that the local environment of

tau's amyloid motif influences aggregation propensity. A mechanistic understanding of how mutations in

tau drive pathology bears directly on how we might better develop conformation-specific diagnostics

and therapies.

As with disease-associated mutations, alternative splicing also impacts the primary sequence N-

terminal to ^{306}VQIVYK^{311}. Tau is expressed in the adult brain primarily as two major splice isoforms: 3-

repeat and 4-repeat[18]. In the truncated 3-repeat isoform, the second of four imperfectly repeated

segments in tau RD is absent. Expression of the 4-repeat isoform correlates with the deposition of

aggregated tau tangles in many tauopathies[19]; non-coding mutations that increase preferential splicing

or expression of the 4-repeat isoform are genetically linked to some tauopathies[19–21]. It is not obvious

why the incorporation or absence of the second repeat correlates with disease, as the primary

sequences, although imperfectly repeated, are relatively conserved.

Finally, previous reports have focused on studying intra-repeat interactions with the assumption

that each repeat functions independently within tau RD[22]. These reports have shown a relationship

between the length of an intra-repeat fragment, its propensity to spontaneously aggregate, and its

seeding capacity in cells[22]. However, inter-repeat interactions may also influence aggregation given that

both alternative splicing and many disease-associated mutations cluster around the repeat interfaces.

We hypothesize that wild-type tau aggregates less efficiently because flanking sequences shield

^{306}VQIVYK^{311}, while disease-causing mutations, alternative splicing, or other factors destabilize local

structures around ^{306}VQIVYK^{311} allowing the transition of tau monomer to an aggregation-prone

conformation.

RESULTS

***In Silico* modeling indicates tau RD local structure.** To initiate our studies, we first used

molecular modeling to gain insight into how putative hairpin structures in tau RD might behave. We built

predictive models using CS-ROSETTA, which implied metastable local β-hairpin structures across

repeat interfaces. We illustrate one of these interface fragments using a model built with CS-ROSETTA

for the R2-R3 [306]VQIVYK[311]-containing interface (**Figure 1 C**). We focused on the [306]VQIVYK[311]-

containing interface over others in tau RD for several reasons. The [306]VQIVYK[311]-containing interface

has the highest frequency of proximal disease-associated mutations, particularly P301L and P301S,

which are commonly used as the genetic basis of disease modelling (**Figure 1 A**). Secondly, other

potential amyloid regions, such as [275]VQIINK[280], are nevertheless capable of aggregation

(**Supplemental Figure 1**), but are absent in recent cryo-EM structures of tau aggregates[3,23].

A standard fragment selection protocol in *ab initio* ROSETTA yielded similar local β-hairpin

structures of the same sequence, further suggesting that the amino acid composition near the amyloid

motif is compatible with beta-turns (**Figure 1 D**). The P-G-G-G sequences preceding [306]VQIVYK[311]

found in tau RD are compatible with the formation of β-hairpins and are typically found in β-turn

secondary structures. Simulating 5,000 structures using each protocol for each of the four potential tau

hairpins led to structures consistent with published Rg measurements[24]. Ensemble-wide analysis of

each dataset showed an approximate ~25-59% propensity to form hairpin-like structures across repeat

interfaces centered on the P-G-G-G sequences (**Supplemental Figure 2**). Mapping known missense

mutations onto the *ab initio* β-hairpin structure (**Figure 1 E**), we hypothesized that this cluster of

disease-associated mutations could act through a single mechanism of action. Mutations that

destabilize this β-hairpin secondary structure may expose the amyloid motif [306]VQIVYK[311], allowing for

aggregation. This model is compatible with recent cryo-EM findings that indicate a disengagement of

[306]VQIVYK[311]'s N-terminal flanking sequence in a fibril structure[3].

P301L promotes extended forms of tau. *In silico* modelling corroborated recent biochemical

findings[7] and suggested a minimal sequence necessary to form a collapsed structure around

[306]VQIVYK[311]. To understand how these structures might self-assemble, we employed molecular

dynamics (MD) simulations of two tau fragments comprising the minimally structured fragment centered

around the R2 to R3 interface: R2R3-WT and R2R3-P301L (**Table 1**). To enable sufficient sampling of

oligomer structures, we employed an unbiased algorithm based on a recently-developed symmetry-

constraint approach[25]. The trimer conformations obtained in simulations are depicted on a root-mean-

square deviation (RMSD) matrix for both R2R3-WT (**Figure 2 A**) and for the R2R3-P301L mutant

(**Figure 2 B**). For R2R3-WT, we observed a dominant population of trimeric conformations composed of

primarily hairpins, while the P301L disease-associated mutation stabilized an extended form. The

energy basin for a R2R3-WT peptide hairpin was predicted to be 5 to 6 kJ/mol lower than an

extended/fibril state, whereas the R2R3-P301L peptide is 3 kJ/mol lower in a fibril state than a hairpin

(**Figure 2 C**). Additionally, the free energy surface suggested an energy barrier of approximately 5

kJ/mol to convert R2R3-WT from hairpin to fibril. That same barrier was less than 1 kJ/mole for R2R3-

P301L, predicting a faster rate of kinetic conversion between the two conformational states. Thus, MD

predicted that a P301L mutation would promote amyloid assembly by destabilizing monomeric hairpin

structures.

Tau amyloid motif has metastable compact structure. Cross-linking mass spectrometry (XL-

MS) defines contact points in proteins and protein complexes and can guide the determination of

structure for large protein complexes or transient protein-protein interactions[26,27]. Tau RD cross-linked

samples are confirmed by SDS-PAGE (**Supplemental Figure 3**) and monomers are purified by Ultra

Performance Liquid Chromatography (UPLC). Cross-linked samples are trypsin digested and analyzed

by mass spectrometry to yield intramolecular protein contact information (**Methods**). In each dataset,

the cross-links reported represent consensus data across five independent samples with a low false

discovery rate (**Methods**). XL-MS of recombinant tau RD revealed higher order contacts at 37 °C

(**Figure 3 D**) that decrease in frequency at 75 °C (**Figure 3 E**), consistent with heat denaturation. We

found a bimodal distribution of short and long-range contacts within multiple repeat domains, which are

consistent with compact local metastable structures.

In contrast, XL-MS of recombinant tau RD with a P301L missense mutation (tau RD-P301L)

revealed an increased susceptibility to heat denaturation. At 37 °C, the crosslinks found in tau RD-

P301L (**Figure 3 G**) are similar than wild type at the same temperature (**Supplemental Figure 4**).

However, at 75 °C there were fewer crosslinks within tau RD-P301L compared to wildtype (**Figure 3 H**),

particularly within the N-terminal sector, which harbors the P301L perturbation.

To further examine the stability of the amyloid motif in tau, we labeled recombinant tau RD

monomer with complementary Fluorescence Resonance Energy Transfer (FRET) fluorophores

(**Methods**, **Table 1**) at the two endogenous cysteines in tau located at positions C291 and C322, which

serendipitously flank the amyloid motif [306]VQIVYK[311] around the predicted β-hairpin structure (**Figure 3**

A-C). In a closed conformation, the two fluorophores should have relatively high FRET, while a

disordered, extended, or denatured chain should yield relatively low FRET signal. To test our system,

we first examined two previously identified monomer populations of FL WT tau: inert monomer (M_i) and

seed-competent monomer (M_s). M_i requires cofactors, such as heparin, to spontaneously aggregate *in*

vitro, while the M_s monomer, derived from tau fibrils or Alzheimer patient brain material, readily self-

assembles to form amyloid[7]. M_i and M_s were labeled at C291 and C322 and their steady-state FRET

intensities measured. M_i, which is predicted by XL-MS to be in a more closed conformation, showed

higher FRET intensities (E = 0.82 ± 0.08, 52.9 Å ± 4.5) compared to M_s (E = 0.46 ± 0.11, 67.7 Å ± 4.4)

(**Supplemental Figure 5**). Denaturing M_i and M_s at 95 °C overnight yielded similar FRET levels (68.9 Å

± 1.8, 65.1 Å ± 1.2 respectively). Surprisingly, this region in M_s is denatured at room temperature,

suggesting an extended conformation of the R2R3 interface for M_s, versus a relatively compact in M_i.

Using wild type recombinant tau RD, a time course heat denaturation of tau monomer (M_i) between 65

°C to 95 °C gradually decreased FRET efficiency, consistent with denaturation of relatively stable local

structure (**Figure 3 F**). The predicted end-to-end distance of a denatured random chain 33-mer is

approximately 66 Å[28]. Consistent with the previous experiment, we observe M_i to be collapsed at room

temperature, and expand to a theoretical fully denatured structure with high temperature (**Figure 3 F**).

We next examined tau RD monomer, M_i, in the context of a P301L mutation, which sits between

the two FRET probes immediately upstream [306]VQIVYK[311] (**Figure 3 B**). A P301L missense mutation is

associated with neurodegeneration in model systems[17,29] and is genetically linked to dementias in

human patients[30], though its biophysical mechanism of action is poorly understood. We found no

detectable difference between monomeric wildtype tau RD and tau RD-P301L FRET energies at room

temperature. Consistent with XL-MS data, tau RD-P301L rapidly loses FRET signal and local structure

with heat denaturation (**Figure 3 I**), suggesting that tau RD-P301L lacks the thermostability associated

with a wildtype tau RD.

Tau amyloid formation is governed by flanking residues. In tau RD, [306]VQIVYK[311] is

sufficient and necessary to form an amyloid[5,6]. In solution, [306]VQIVYK[311] hexapeptides aggregate

spontaneously and rapidly as measured by Thioflavin T (ThT) fluorescence intensity ($t_{1/2}$ < 1 hr,

Supplemental Figure 1) whereas the N-terminal sequence [295]DNIKHV[300] does not aggregate.

Combining these sequences yields the minimal structural element spanning repeat 2 into repeat 3

([295]DNIKHVPGGGSVQIVYK[311], R2R3) predicted by *in silico* modelling (**Table 1, Figure 4 A**). This

sequence is also where numerous disease-associated mutations cluster. Indeed, WT peptide

fragments representing this motif did not aggregate readily, with no ThT detected up to 96 hrs (**Figure 4**

C). By contrast, disease-associated mutations (**Figure 4 B**) substituted into the R2R3 peptide fragment

were sufficient to generate spontaneous amyloid formation: R2R3-P301S ($t_{1/2}$ = 4.1 ± 1.3 hours), R2R3-

P301L ($t_{1/2}$ = 7.2 ± 0.2 hours), R2R3-N296Δ ($t_{1/2}$ = 31.9 ± 0.2 hours), R2R3-G303V ($t_{1/2}$ = 32.1 ± 0.7

hours), R2R3-S305N ($t_{1/2}$ = 41.2 ± 0.2 hours), and R2R3-V300I ($t_{1/2}$ = 77.8 ± 1.3 hours, **Figure 4 C**).

Each of these peptides was confirmed to form amyloid-like fibril morphologies by transmission electron

microscopy, except for the wildtype R2R3 peptide where no large structures were found (**Figure 5 B-**

H).

We next assessed whether these mutations perturb local structure. We performed circular

dichroism on both the R2R3-WT and R2R3-P301L peptides (**Figure 4 C**). Compared to R2R3-WT,

R2R3-P301L monomer had diminished 198nm signal consistent with more random coil and less β-

sheet. Deconvolution of CD spectra using BeStSel (beta structure selection) revealed predominantly

anti-parallel signal (**Table 3, Supplemental Figure 6**)[31]. Both peptides contained significant random

coil structure, which is expected for metastable structures, intrinsically disordered proteins, and short

peptides that cannot gain stability from tertiary structure. The compact structures identified are

consistent with intrinsically disordered proteins having residual structural preferences.

To test the structural compatibility of aggregates formed by *in vitro* tau models, we employed tau

biosensor HEK293 cells that stably express tau RD (P301S) fused to cyan and yellow fluorescent

proteins. These cells sensitively report a FRET signal (tau RD-CFP/tau RD-YFP) only when exposed to

tau amyloid seeds, and are unresponsive to aggregates formed by other proteins, such as huntingtin or

α-synuclein[32]. The tau biosensor cells responded to all disease-associated peptides that aggregated

spontaneously *in vitro*, but not to the wild-type R2R3 peptide (which did not aggregate *in vitro*) (**Figure**

5 A). Qualitatively, biosensor cells retained their diffuse tau localization when untreated or exposed to a

wild-type R2R3 peptide, but formed puncta when cultured with aggregated mutant peptides (**Figure 5 I-**

P). Interestingly, the biosensor cells responded to disease-associated mutant peptides with varying

degrees of sensitivity and created dissimilar aggregate morphologies. This is consistent with amyloid

structures that act as distinct templates, as we have previously observed to form the basis of tau prion-

like strains[4,33].

Tau splice variants reveal differential aggregation propensity. Tau is expressed in the adult

brain as 6 major splice isoform types that include either 3 or 4 repeated segments within RD (**Figure 6**

A). 3R lacks the second of four repeats. 4R tau correlates strongly with aggregation in most

tauopathies[19] and mutations that increase splicing of the 4R isoform correlate with disease[19,21]. We

examined whether this splice isoform affects the propensity of ^{306}VQIVYK^{311}-mediated aggregation due

to the R1 to R3 junction having a different sequence composition from R2 to R3. We constructed a

series of peptides to encompass the R1 to R3 repeat interface (R1R3, **Table 1**). This wild type

sequence peptide mimicking a 3R splice isoform (R1R3) did not spontaneously aggregate. Surprisingly,

an R1R3 peptide with a corresponding P301L mutation also did not aggregate (**Figure 6**). We

hypothesized that differences in amino acid composition between the two splice isoforms led to the

sensitivity to aggregate in the presence of disease-associated mutations.

The R1 leading sequence ^{264}**EN**L**KHQ**PGGG**K**^{273} differs from R2 ^{295}**DN**I**KHV**PGGG**S**^{304} at four

amino acid positions. To test the stabilizing effect of the R1 leading sequence, we constructed 16

peptides with a P301L disease-associated mutation to represent every combinatorial sequence

between the two leading strands (**Figure 6 B**). In this way, we could identify which amino acid(s)

governed R1's stronger inhibitory effects. We identified a general trend where R2R3-P301L aggregates

in hours with zero or one R1 substitutions. With 2 R1 substitutions, R2R3-P301L aggregation was

delayed roughly an order of magnitude to tens of hours. With 3 R1 substitutions, R2R3-P301L

aggregation was further delayed to hundreds of hours. With all four R1 substitutions (R1R3-P301L), no

ThT signal was observed within a week (**Figure 6 B**). Thus, all four amino acids contributed to the ability of the R1 leading sequence to delay [306]VQIVYK[311]-mediated spontaneous aggregation in a 3R splice isoform. This may explain the differential aggregation propensities of tau isoforms in human pathology.

Stabilizing β-hairpin structure blocks P301L-mediated aggregation. Our model predicted sequestration of the [306]VQIVYK[311] motif in tau *via* local β-structure. To further test this, we hypothesized that artificially enhancing the local β-hairpin structure by stabilizing the termini would promote a more inert, closed conformation. Consequently, we modified the R2R3-P301L peptide with a tryptophan zipper (Trp-R2R3-P301L-Trp, **Table 1**), which stabilizes a β-hairpin structure approximately -2.5 to -7 kJ/mol[34]. We confirmed a structural change using circular dichroism of Trp-R2R3-P301L-Trp, where we found an increase in β-sheet signal at 198 nm compared to R2R3-P301L (**Supplemental Figure 6, Table 3**). Subtracting the R2R3-P301L signal from the Trp-R2R3-P301L-Trp revealed a curve with significantly increased signal at 198 nm, indicative of β-sheet formation. Consistent with our model, Trp-R2R3-P301L-Trp does not spontaneously aggregate *in vitro* (**Figure 7 B**).

To ensure that this effect wasn't a result of adding bulky tryptophan residues, we constructed control peptides that contain only the N-term (Trp-R2R3-P301L) or the C-term (R2R3-P301L-Trp) portion of the tryptophan zipper sequence (**Figure 7 A**). Both half-sequence controls spontaneously aggregated, implying that a tryptophan in either position is insufficient to block aggregation (**Figure 7 B**). Only a fully intact tryptophan zipper that stabilizes a β-hairpin conformation ameliorates aggregation propensity. Alternative methods to stabilize a β-hairpin architecture, such as introducing isoelectric interactions, also delayed aggregation: peptides containing additional aspartic acids on the N-terminus and lysines on the C-terminus (R2R3-IEZip, **Table 1**) retarded R2R3-P301L aggregation over an order of magnitude ($t_{1/2}$ = 8 hours to $t_{1/2}$ = 83 hours, **Supplemental Figure 7**).

To test this effect in cells, we generated biosensor HEK293 cells expressing tau RD with a comparable tryptophan zipper sequence. These biosensors had a significantly diminished capacity to be seeded; R2R3-P301S peptide aggregates triggered aggregation in 14.5 ± 1.6% of tau biosensor cells, but only 0.3 ± 0.12% of the tryptophan zipper stabilized biosensor cells (**Supplemental Figure 8**).

Proline 301 *cis-trans* isomerization modulates aggregation. Many proteins in the cell utilize

proline isomerization as a molecular switch, such as heat shock protein activation[35] or cell cycle

regulation[36]. In some proteins, proline isomerization directly induces or mitigates aggregation into

amyloid[37–39]. Proline isomerization events in tau have been proposed to play a role in aggregation and

disease[38], but P301 isomerization has not been linked to tau aggregation and pathology. Serine or

leucine substitutions at P301 proximal to [306]VQIVYK[311] drastically alter aggregation propensity. We

hypothesized that P301 plays a crucial role inducing a β-turn in a P-G-G-G motif, which mediates a

collapsed structure. We tested whether isomerization of P301 could influence spontaneous amyloid

formation. We constructed a series of R2R3 peptides with proline analogs that preferentially populate

either: (1) a *cis* rotamer (2S,4S)-fluoroproline; (2) a *trans* rotamer (2S,4R)-fluoroproline; or (3) a proline

analog that easily interconverts between *cis* and *trans* (4,4)-difluoroproline (**Table 1, Figure 7 C**). Only

R2R3-Trans spontaneously aggregated (**Figure 7 D**), indicating the potential for proline isomerization

events in tau pathogenesis.

DISCUSSION

Here we establish the molecular and functional basis for how a series of prominent tau mutations drive spontaneous aggregation. Using intramolecular FRET and XL-MS, we identified local structure within the inter-repeat junctions of tau RD. Local compact structure in the R2-R3 junction, which encompasses the amyloidogenic ^{306}VQIVYK^{311} motif, is perturbed by disease-associated mutations, such a P301L. Using peptide model systems, we demonstrate that splice isoforms of tau are functionally dissimilar in their capacity to spontaneously aggregate with missense mutations. Furthermore, these peptide models indicate a potential role of proline isomerization in tau aggregation at the critical P301 position. Conversely, stabilizing local structure with tryptophan zippers or other means drastically limits the aggregation *in vitro* and *in vivo*. Taken together, our work provides a biophysical framework for how pathogenic mutations, splice isoforms, and other perturbations can drive early events in tau aggregation.

Our study has suggested that local structure encompassing the amyloid motif ^{306}VQIVYK^{311} regulates aggregation of the protein. Modeling of tau RD by *ab initio* or CS-ROSETTA indicated that repeat interfaces encode local structure that is compatible with a β-hairpin. Importantly, a cluster of disease-associated mutations localized to the repeat2-repeat3 interface suggested a possible mechanism. To explore the energetic landscape of local compact structure proximal to ^{306}VQIVYK^{311}, we employed molecular dynamics simulations. A WT peptide encoding the ^{306}VQIVYK^{311}-containing repeat 2/3 interface preferentially populated a hairpin conformation with a relatively low frequency of extended fibril-like structures. Conversely, a disease-associated mutation, P301L, dramatically shifted the equilibrium away from monomeric hairpins to fibril-like ensembles. To test the extent of structure formed in tau RD, we utilized orthogonal biophysical approaches. First, XL-MS revealed contacts spanning the local repeats, and long-range contacts within tau RD. Heat denaturation experiments indicated a marked drop in long-range contacts, consistent with the unfolding of secondary and tertiary structures. Concurrently, intramolecular FRET studies of tau RD experimentally followed unfolding of the ^{306}VQIVYK^{311} repeat 2/3 interface in real time by measuring distances between fluorescent probes that flank ^{306}VQIVYK^{311}. Under native conditions the local structure was collapsed, but expanded to random coil with heat denaturation. Conversely, tau RD with a P301L substitution was less

thermostable and rapidly lost FRET signal. This suggests that P301L is more susceptible to

conformational changes that expose the ^{306}VQIVYK^{311} amyloid motif. No structural differences were

seen between wildtype and tau RD-P301L at non-denaturing temperatures. While these differences are

subtle, we observe that P301L-mediated structural rearrangement only manifest under moderate stress

conditions (i.e. heat). This may explain the elusiveness of a biophysical basis of this cluster of

pathogenic mutations.

We further examined whether these structural perturbations influenced aggregation propensity.

WT tau fragments containing ^{306}VQIVYK^{311} did not aggregate spontaneously. Single point substitutions

of 6 disease-associated mutations immediately N-terminal to ^{306}VQIVYK^{311} consistently induced

spontaneous aggregation propensity *in vitro*. The fragments which aggregated *in vitro* formed classical

amyloid-like fibrils by EM and seeded soluble tau RD aggregation in tau biosensor cells. Alternative

splicing around the ^{306}VQIVYK^{311} sequence modulated spontaneous aggregation, and suggested why

3R tau aggregates less readily than 4R. To further probe the effect of local structure on aggregation

propensity, we stabilized a β-hairpin structure using a tryptophan zipper or isoelectric forces. After

deconvolution of the CD signal by BeStSel, we identified that essentially all the β-sheet signal from the

peptides tested was anti-parallel, consistent with β-hairpin structure. The magnitude of these changes

are modest as detected by CD; however, the functional consequences are significant.

Disease-associated mutations found near tau's amyloid motif, such as P301L or P301S have no

definitive biophysical mechanism but are nevertheless widely used in cell and animal models[16,17].

Mutations are generally categorized into splice or missense mutants. Two of the pro-aggregation

missense mutations studied (N296Δ, S305N) are reported in the literature as splice mutants with no

notable effects on protein aggregation or microtubule binding[40,41]. Using our peptide model system, we

observed that these missense mutations spontaneously aggregate *in vitro* and seed tau RD in cell

models. We therefore provide a rationale for the toxic-gain-of-function for several mutations and expand

on the previously reported splice effects of N296Δ and S305N. This peptide model system compliments

in vivo data by disentangling the toxic effects of mutations exerted through either mRNA or protein-

based mechanisms.

Studying the missense mutations in tau has generated valuable disease models[17,29]; however,

the majority of human tauopathies have no observed genetic mutation in *tau*[30]. Critical proline residues

N-terminal to the amyloid motif can isomerize into *cis* or *trans* rotamers spontaneously or through

unidentified cellular mechanisms. We observe that P301 *cis* and *trans* rotamers have distinct

aggregation propensities *in vitro*. In fact, the aggregation kinetics for a *trans* rotamer of P301 are on par

with some disease mutants (N296Δ, V300I). The concept of proline isomerization triggering

aggregation into amyloid is not entirely novel, as this is an accepted mechanism of β2-microglobulin

aggregation in kidney dialysis amyloidosis[42]. Other proline residues outside of tau repeat domain are

proposed to undergo proline isomerization[38]. We postulate a new mechanism whereby WT tau

aggregation could be controlled *in vivo*: specific prolyl isomerization events – possibly triggered by

cellular proline isomerases – could trigger spontaneously aggregation by modulating inter-repeat

structural elements.

Previous reports have been focused on studying intra-repeat interactions, with the assumption

that each repeat functions independently within tau RD. Peptide models have shown a relationship

between the length of a peptide fragment and the seeding capacity of tau[22]. Whereas previous work

defined the minimal sequence necessary to act as a fully functional seed, our model defines a minimal

sequence necessary for inhibiting aggregation. This work suggests that inter-repeat contacts play a

crucial role as structural elements that modulate aggregation propensity. The composition of these

inter-repeat sequences, governed by alternative splicing or missense mutations, directly impacts

stability of local structures and aggregation propensity. It is tempting to speculate that local structure

surrounding each of the four inter-repeat regions play independent roles in the exposure of amyloid

sequences. This modular nature of the tau-RD region may explain how these independent regions can

lead to different tau assemblies. A more comprehensive structure-function analysis of other sites may

help explain how each repeat contributes to the formation of different tau structures.

The expression levels of the two major isoform types of tau in the CNS – 3R and 4R – are

similar in the adult brain[18]. However, the 3R:4R ratio of aggregate deposits is disproportionally shifted

towards 4R in most tauopathies[19]. Mutations in tau that affect alternative splicing and generate excess

4R isoforms correlate with some genetic tauopathies[20,21]. The N-terminal flanking sequences leading

into ^{306}VQIVYK^{311} differ by four amino acids between the two isoforms. We find that these two isoforms

have drastically different aggregation propensities in the presence of disease-associated mutations ($t_{1/2}$

= 7 hours vs $t_{1/2}$ > 200 hours, respectively). Chimeras of R1R3 / R2R3 transition from aggregation-

resistant to aggregation-prone as they lose R1 N-terminal flanking character. The ability of an R1

leading strand to mitigate ^{306}VQIVYK^{311} aggregation may explain why 4-repeat tau correlates more

closely with pathology. Thus, inter-repeat contacts may explain aggregation propensies of tau isoforms

in disease. Encouraging data for a tau vaccine targeting a ^{300}HXPGGG^{304} sequence suggests it's

possible to utilize inter-repeat regions to select between pathogenic and non-pathogenic conformations

of tau[43].

Taking these observations together, we propose that sequences N-terminal to tau's amyloid

motif forms local contacts consistent with a β-hairpin-like compact structure, which shields the amyloid

motif and mitigates aggregation (**Figure 8**). This represents a simple model of tau aggregation that

unifies key observations throughout tau literature.

Algorithms that identify potential amyloid nucleating regions, such as TANGO, have indicated

that nearly 75% of aggregation nucleating regions in the human proteome use two or more

"gatekeeper" residues, with proline being the most common single gatekeeping residue[44]. These

gatekeeping residues are more likely than average to be the site of disease-associated missense

mutations. These observations tie in closely with our findings of gatekeeping residues near tau's

amyloid motif, and suggest that local flanking sequences and their structural contacts may play an

important role in mitigating aggregation propensity.

Finally, the identification and characterization of metastable compact structures near

^{306}VQIVYK^{311} may itself prove to be a valuable therapeutic target. One might be able to shift the

structural rearrangement of tau amyloid motif from exposed (aggregation prone) to buried (inert) using

small molecules, antibodies, or cellular co-factors. Our results indicate that subtle changes in local

structure have immense functional ramifications; therefore, small molecules that shift this structural

equilibrium modestly may have significant benefits.

Peptide Synthesis. All peptides were synthesized as ordered by Genscript with N-terminal acetylation and C-terminal amidation modifications. Peptides were purified to >95% purity by FPLC *via* an Agilent ZORBAX StableBond 250 mm C8 column.

ThT Fluorescence Assays. Peptides were disaggregated as previously described[45], lyophilized, and resuspended in 2x PBS (273 mM NaCl, 5.4 mM KCl, 20 mM Na_2HPO_4, 3.6 mM KH_2PO_4, pH 7.4). 25 mM ThT was added to 200uL of 200 μM peptide in a 96-well clear bottom plate. ThT kinetic scans were run every 5 minutes on a Tecan M1000 plate reader at 446 nm Ex (5 nm bandwidth), 482 nm Em (5 nm bandwidth). Blank wells containing buffer and ThT were subtracted from experimental values.

Tau intramolecular FRET. Tau RD was expressed and purified as previously described[46]. 10-fold excess TCEP was added to 100 μM Tau RD, the atmosphere was deoxygenated with N_2 gas, and incubated at RT for 30 minutes. To label, C_5 maleimide Alexa-488 and C_2 maleimide Alexa-647 were dissolved in DMSO and added in 5-fold excess to Tau RD, and incubated at 4^o C overnight. Alexa-labelled Tau RD was purified using an 8 kDa cut-off mini dialysis exchanged into 1x PBS overnight. FRET efficiencies were calculated as a function of acceptor emission, given $E = (I_{AD}\varepsilon_{AA}-I_{AA}\varepsilon_{AD})/I_{AA}\varepsilon_{DD}$ where I_{AA} is Acceptor intensity following Acceptor excitation, I_{AD} is Acceptor intensity following Donor excitation, and ε is the extinction coefficient of the fluorophores at given excitations and emissions. Distances in angstroms were derived from FRET energies as a function of $Å = (R_{<0>})[(1-E)/E]^{1/6}$ $R_{<0>}$ is a known value of 56 Å for the FRET pair Alexa-488 and Alexa-647, and E is the calculated energy above. Alexa 488 Ex/Em was measured at 490 nm/520 nm, Alexa 647 Ex/Em 647 nm/670 nm with a 5 nm bandwidth for both.

Model generation of tau RD using ROSETTA. The backbone NH, N, CA, CB and C=O chemical shift assignments for the tau fragment from 211-324 (generously provided by Juan Lopez and Guy Lippens) were used in CS-Rosetta to generate fragment libraries for subsequent model refinement. First, chemical shift parameters were used to predict backbone torsional angles using TALOS to generate a CS-guided fragment library representing the conformations of the protein[47]. For the *ab initio* ROSETTA calculations, the tau RD sequence was used to generate 3-mer and 9-mer fragments derived from the

protein data bank. The Rosetta energy function was used to assemble and iteratively refine 5000

structural models using each set of fragments[48,49]. Radius of gyration for each model was computed by

a python script. Ensemble wide calculation of $c\alpha$-$c\alpha$ end to end distances between residues 264-280,

295-311, 327-343 and 359-375 were carried out using a python script. All simulations were done on

UTSW's biohpc computing cluster. All plots were generated with gnuplot. Images were created using

Pymol.

Circular Dichroism. Far-UV CD measurements were performed on a JASCO J-815 spectropolarimeter

using a 1-mm path length cuvette. CD samples were prepared in parallel to ThT assays and diluted to

0.1 mg/ml in 20 µM NaF. CD spectra were analyzed using BeStSel and CONTIN.

Tau Biosensor Cells. Biosensor cells were plated into 96-well plates at 20,000 cells per well. 10 µg of

aggregated peptide material was added to 0.5 uL lipofectamine and Opti-MEM to a total volume of 10

µL, incubated at room temperature for 30 minutes, and added directly to cell media. After 3 days, cells

were harvested with 0.05% trypsin, then resuspended in Flow buffer (1x HBSS, 1% FBS, 1 mM EDTA,

1x DPBS) and analyzed by flow cytometry.

Flow Cytometry. A BD LSRFortessa was used to perform FRET flow cytometry. To measure CFP and

FRET, cells were excited with the 405 nm laser, and fluorescence was captured with a 405/50 nm and

525/50 nm filter, respectively. To measure YFP, cells were excited with a 488 laser and fluorescence

was captured with a 525/50 nm filter. To quantify FRET, we used a gating strategy where CFP bleed-

through into the YFP and FRET channels was compensated using FlowJo analysis software. We then

created a plot of FRET vs. CFP and introduced a triangular gate to assess the number of FRET-positive

cells, as previously described[16]. For each experiment, 20,000 cells per replicate were analyzed. Data

analysis was performed using FlowJo v10 software (Treestar).

Transmission electron microscopy (TEM). An aliquot of 5 µl of sample was placed onto a glow-

discharged Formvar-coated 400-mesh copper grids for 30 seconds, washed with distilled water, and

then negatively stained with 2% uranyl acetate for 1 min. Images were acquired on a Tecnai G^2 spirit

transmission electron microscope (FEI, Hillsboro, OR), serial number: D1067, equipped with a LaB_6

source at 120kV using a Gatan ultrascan CCD camera.

Crosslinking, sample processing and LC-MS/MS analysis. Preparation of tau RD was crosslinked

at a total protein concentration of 1.0 mg/mL using 100 µg of starting material. The crosslinking buffer

was 1X PBS and 1mM DTT. For each temperature (37^{o}C, 75^{o}C, 85^{o}C and 95^{o}C) five replicates

samples were prepared and equilibrated at the appropriate temperature for 30 minutes. The

crosslinking reaction was initiated by adding disuccinimidyl suberate (DSS) stock solution (25 mM DSS-

d_0 and $–d_{12}$, Creative Molecules) in DMF to a final concentration of 1 mM. Samples were further

incubated at 37^{o}C, 75^{o}C, 85^{o}C or 95^{o}C for 1 min with 350 RPM shaking. Excess reagent was quenched

by addition of ammonium hydrogen carbonate to 50 mM and incubation at 37°C for 30 min, and

subsequently flash frozen at -80°C and evaporated to dryness by lyophilization. Proteins were

resuspended in 8M urea, reduced with 2.5mM TCEP (37°C, 30 min) and alkylated with 5mM

iodoacetamide (30min, room temperature, protected from light). The sample solutions were diluted to 1

M urea with 50 mM ammonium hydrogen carbonate and trypsin (Promega) was added at an enzyme-

to-substrate ratio of 1:50. Proteolysis was carried out at 37°C overnight followed by acidification with

formic acid to 2% (v/v). Samples were then purified by solid-phase extraction using Sep-Pak tC18

cartridges (Waters) according to standard protocols. Samples were evaporated to dryness and

reconstituted in water/acetonitrile/formic acid (95:5:0.1, v/v/v) to a final concentration of approximately

0.5 µg/µl. 2µL each were injected for duplicate LC-MS/MS analyses on an Eksigent 1D-NanoLC-Ultra

HPLC system coupled to a Thermo Orbitrap Fusion Tribrid system. Peptides were separated on self-

packed New Objective PicoFrit columns (11cm x 0.075mm I.D.) containing Magic C_{18} material

(Michrom, 3µm particle size, 200Å pore size) at a flow rate of 300nL/min using the following gradient. 0-

5min = 5 %B, 5-95min = 5-35 %B, 95-97min = 35-95 %B and 97-107min = 95 %B, where A =

(water/acetonitrile/formic acid, 97:3:0.1) and B = (acetonitrile/water/formic acid, 97:3:0.1). The mass

spectrometer was operated in data-dependent mode by selecting the five most abundant precursor ions

(m/z 350-1600, charge state 3+ and above) from a preview scan and subjecting them to collision-

induced dissociation (normalized collision energy = 35%, 30ms activation). Fragment ions were

detected at low resolution in the linear ion trap. Dynamic exclusion was enabled (repeat count 1,

exclusion duration 30sec).

Analysis of mass spectrometry data. Thermo .raw files were converted to the open .mzXML format

using msconvert (proteowizard.sourceforge.net) and analyzed using an in-house version of xQuest[50].

Spectral pairs with a precursor mass difference of 12.075321 Da were extracted and searched against

the respective FASTA databases containing Tau (TAU_HUMAN P10636-8) or with a P301L

substitution. xQuest settings were as follows: Maximum number of missed cleavages (excluding the

crosslinking site) = 2, peptide length = 5-50 aa, fixed modifications = carbamidomethyl-Cys (mass shift

= 57.021460 Da), mass shift of the light crosslinker = 138.068080 Da, mass shift of mono-links =

156.078644 and 155.096428 Da, MS^1 tolerance = 10 ppm, MS^2 tolerance = 0.2 Da for common ions

and 0.3 Da for crosslink ions, search in ion-tag mode. Post-search manual validation and filtering was

performed using the following criteria: xQuest score > 18, mass error between -2.2 and +3.8ppm, %TIC

> 10, and a minimum peptide length of six aa. In addition, at least four assigned fragment ions (or at

least three contiguous fragments) were required on each of the two peptides in a crosslink. False

discovery rates (FDR's) for the identified crosslinks were estimated using xprophet[51] and estimated to

be 1.4-3.3% (**Supplemental Figure 9, Supplemental Table 1**). At each temperature, the 5 replicate

datasets were compared and only crosslinks present in 5 of the 5 datasets were used to generate a

consensus dataset (**Supplemental Table 2**). Crosslink data was visualized using Xvis[52]. Histograms

illustrating the contact order for the consensus crosslink sets were generated using gnuplot.

MD Simulations. Well-Tempered Metadynamics[53] was employed to enable accelerated conformational

sampling and to construct the associated free energy surface. Metadynamics was performed on a two-

dimensional space of parallel-beta sheet content and anti-parallel sheet content. To increase search

efficiency in oligomeric space, we have incorporated conformational symmetry constraints, which have

been shown to enable sampling of multi-polymer landscapes[25]. The initial dodecahedron simulation box

was constructed from a trimer of a randomly unfolded structure of 295-311 by adding 7587 SPCE

explicit waters and 3 neutralizing Cl ions (one for each monomer). The AMBER99sb-ildn force-field[54]

was used for all simulations. After an initial 1009 steepest descent steps of converged energy

minimization, 10 ns of NVT and 20 ns of NPT (first 10 with Berendsen[55] and the last 10 with Parrinello-

Rahman[56] barostats) equilibrations were performed. The subsequent production level trajectories are

based on 5 fs time steps using hydrogen-only virtual sites[57]. Production level trajectories were obtained

for an NPT ensemble with Parrinello-Rahman barostat, and periodic boundary conditions with Particle

Mesh Ewald (PME)[58] summation for long-range electrostatics. The tuned well-tempered metadynamics

parameters are 10, 1.4 kJ/mole, and 0.3 for bias factor, Gaussian height, collective variable space

Gaussian widths, respectively. The Gaussian perturbations were included into MD every 2.5 ps using

the PLUMED package[59] as an external patch to Gromacs-5.0.4[60]. A total of 18 µs trajectories were

generated, 9 µs for wild type and 9 µs for the P301L mutant, over a total of 6 independent runs. All

simulations were done on UTSW's biohpc computing cluster.

Statistics: All statistics were calculated using GraphPad Prism 7.0. A minimum of 3 independent ThT

experiments were run for each condition. Plots were fitted to a non-linear sigmoidal curve, from which

$t_{1/2}$ values were derived. $t_{1/2}$ error represents a 95% CI. Flow cytometry cell aggregation was conducted

a minimum of 3 independent experiments, whose values are plotted. Error bars represent a 95% CI.

Acknowledgements

This work was supported by grants from the Tau Consortium and NIH grants awarded to

1R01NS071835 (M.I.D.), the Effie Marie Cain Endowed Scholarship (L.A.J.) and the Heising-Simons

Foundation (M.M.L.). We appreciate the help of the Structural Biology Laboratory and Proteomics Core

Facility at the University of Texas Southwestern Medical Center.

Author Contributions

K.D. M.I.D., and L.A.J. conceived and designed the overall study. K.D. performed *in vitro* aggregation

assays, peptide studies, flow cytometry, and cell models. D.C. performed crosslink mass spectrometry.

D.W. performed electron microscopy. L.S. and M.L. performed molecular dynamics simulations. K.D.

and L.A.J. wrote the manuscript, and all authors contributed to its improvement.

Table 1: List of Peptide and Protein Sequences

Peptide Name	Amino Acid Sequence
VQIINK	-----------------VQIINK
VQIVYK	-----------------VQIVYK
R1R2 (264-280)	------ENLKHQPGGGKVQIINK
R2R3 (295-311)	------DNIKHVPGGGSVQIVYK
R2R3-N296Δ	------D-IKHVPGSSSVQIVYK
R2R3-V300I	------DNIKH**I**PGGGSVQIVYK
R2R3-P301L	------DNIKHV**L**GGGSVQIVYK
R2R3-P301S	------DNIKHV**S**GGGSVQIVYK
R2R3-G303V	------DNIKHVPG**V**GSVQIVYK
R2R3-S305N	------DNIKHVPGGG**N**VQIVYK
R1R3	------ENLKHQPGGGSVQIVYK
R1R3-P270L	------ENLKHQ**L**GGGSVQIVYK
R1R3-P270S	------ENLKHQ**S**GGGSVQIVYK
R1R3-G272V	------ENLKHQPG**V**GSVQIVYK
R1R3-G273R	------ENLKHQPGG**R**SVQIVYK
Trp-R2R3-P301L-Trp	**WTGK**SKDNIKHV**L**GGGSVQIVYK**EGGW**
Trp-R2R3-P301L	**WTGK**SKDNIKHV**L**GGGSVQIVYKPVDL
R2R3-P301L-Trp	SKCGSKDNIKHV**L**GGGSVQIVYK**EGGW**
R2R3-IEZip	----DDDNIKHVPGGGSVQIVYKKK
R2R3-IEZip-P301L	----DDDNIKHV**L**GGGSVQIVYKKK
R2R3-Cis[1]	------DNIKHV**P**GGGSVQIVYK
R2R3-Trans[1]	------DNIKHV**P**GGGSVQIVYK
R2R3-Ea[1]	------DNIKHV**P**GGGSVQIVYK
Tau RD[2]	R1: [244] QTAPVPMPDLKN-VKSKIGSTENLKHQPGGGK [274] R2: [275] VQIINKKLDLSN-VQSK**C**GSKDNIKHVPGGGS [305] R3: [306] VQIVYKPVDLSK-VTSK**C**GSLGNIHHKPGGGQ [336] R4: [337] VEVKSEKLDFKDRVQSKIGSLDNITHVPGGGN KKIETH [374]

526 [1]Fluorinated proline analogs were used to preferentially populate *cis, trans,* or lowered E_a barrier
527 conformers at the positions indicated as **P**.
528 [2]Cysteines labeled for Tau RD internal FRET are market as **C**.
529

530 **Table 2: List of AlzForum Disease-Associated Mutations**

Peptide Name	Amino Acid Sequence
Tau RD AlzForum Mutations	R1: 244 QTAPVPMPDLKN-V**K**SK**I**GSTEN**L**KHQP**GG**GK 274 R2: 275 VQII**N**KKLD**LS**N-VQSKCGSKD**N**IKH**VP**G**G**G**S** 305 R3: 306 VQIVYKPVD**LSK**-**V**TSKCGSLGN̄IHHK**P**GG**GQ** 336 R4: 337 **V**EVKSE**K**LDFKDRVQ**S**KIG**S**LDN**I**TH**VP**GG**G**N 368

531
532

533 **Table 3: Summary of Circular Dichroism Results**

Peptide Name	% α-helix	% anti-parallel	% parallel	% turn	% random
R1R2	0	39.5	0.9	14.5	45.1
R2R3	0	36.7	2.8	14.6	45.9
R2R3-P301L	0.3	34.0	2.6	15.4	47.7
Trp-R2R3-Trp	0	42.0	0	13.9	44.1
Trp-R2R3-P301L-Trp	0	42.5	0	13.6	43.9

534
535

	% α-helix	% anti-parallel	% parallel	% turn	% random

Figure Legends

Figure 1. **Tauopathy mutations cluster to inter repeat regions**. A. Disease-associated mutation frequency found in human tauopathies. Most mutations are found within the repeat domain (tau-RD) (repeat 1 = red; repeat 2 = green; repeat 3 = blue; repeat 4 = purple). Amyloidogenic sequences ^{275}VQIINK^{280} and ^{306}VQIVYK^{311} are shown in the inset cartoon. B. Detailed mutation frequencies found near the ^{306}VQIVYK^{311} amyloid motif. C, D. Secondary structure models of the repeat2-repeat3 interface (^{295}DNIKHVPGGGSVQIVYK^{311}) predicted from CS-ROSETTA (C) or *ab initio* (D) shown in cartoon and colored by repeat domain. The amyloid motif ^{306}VQIVYK^{311} is colored blue, the leading repeat-2 sequence is green. E. Positions of disease-associated mutations (red sphe res) mapped onto the ^{306}VQIVYK^{311}-containing *ab initio* β-hairpin structure.

Figure 2. Wildtype and mutant peptides differentially populate hairpin and extended conformations. A. Conformations obtained for a peptide fragment wild type trimer of the sequence ^{295}DNIKHVPGGGSVQIVYK^{311}. Two-dimensional root-mean-squared-differences (RMSD's) are calculated between all pairs of conformations visited during MD simulations. Snapshots of trimeric structures are depicted for select metastable basins, with each peptide monomer represented by a different color. B. The same analysis as above, but for the P301L substituted trimer. C. The free energy surface as a function of deviation from a canonical hairpin structure. Two distinct basins, corresponding to hairpin and fibril/extended sub-ensembles, are found.

Figure 3. Tau RD encodes global and local structure. A-C. Cartoon schematic of tau RD used for FRET and XL-MS studies colored according to repeat domain (repeat 1 = red; repeat 2 = green; repeat 3 = blue; repeat 4 = purple). For FRET studies, endogenous cysteines are labelled with alexa-488 and alexa-647 (red and blue balls), which flank the ^{306}VQIVYK^{311} amyloid motif. D. Monomeric tau-RD with labeled FRET pairs was denatured at varying temperatures. E-F. Tau RD samples were incubated at 37 °C or 75 °C for one hour. After cross-linking, trypsin fragmentation, and LC-MS/MS analysis, consensus cross-

link patterns (circles) are shown as contact maps. Short range crosslinks within the N-term (blue), C-term

(red) and long range contacts across N- to C-term (purple) are shown as sectors. Crosslinks are colored

according to mean frequency across replicates. G-I. Same as D-F above, except with tau RD that contains

a P301L disease-associated substitution.

Figure 4. Tauopathy mutations drive aggregation propensity. A. Schematic of tau-RD and the derived

peptides representing minimal structural elements around ^{306}VQIVYK^{311}. B. Primary amino acid sequence

mapped onto a cartoon of the predicted minimal structural element proximal to ^{306}VQIVYK^{311}. Mutations

are shown as spheres. C. Wild type and mutant peptides were disaggregated, resuspended at 200 μM, and

allowed to aggregate in the presence of ThT at room temperature. ThT signals are an average of at least 3

independent experiments and are colored according to mutation.

Figure 5. Peptides form amyloid structures and seed *in vivo*. A. After 96 hours of *in vitro* incubation,

peptides from previous ThT experiments (**Figure 4 C**) were transduced into tau biosensor cells *via*

lipofectamine (**Methods**). FRET signal from each condition (tau-RD-CFP/tau-RD-YFP) was measured

by flow cytometry used 3 biological triplicates of at least 10,000 cells per condition. Error bars represent

a 95% CI of each condition. Solid and dashed horizontal lines represent the mean and 95% error from

untreated biosensor cells, respectively, for ease of statistical comparison. B-H. Electron microscopy

images of each peptide from previous ThT experiments (**Figure 4 C**). The black bar represents 200 nm

distance in each image I-P. Qualitative fluorescence microscopy images of tau biosensor cells immediately

prior to Flow Cytometry experiments.

Figure 6. Alternative splicing modulates aggregation propensity. A. Cartoon schematic for tau 4R and

3R splice isoforms illustrate the difference in primary amino acid sequence leading into the amyloidogenic

^{306}VQIVYK^{311} motif. B. A full combinatorial panel of R2R3-P301L and R1R3-P301L chimeras were

aggregated *in vitro*. ^{306}VQIVYK^{311} is shown in blue, amino acids common between the splice isoforms

are shown in black, amino acids unique to an R3 isoform are colored red, amino acids unique to an R4

isoform are colored green. The kinetic times to aggregate, represented as $t_{1/2}$ in hours with 95% CI, are

listed in the right-side column alongside its respective peptide.

Figure 7. Enhancing β-hairpin structure rescues spontaneous aggregation phenotypes. Cartoon

schematic representation of the tryptophan zipper motif (green bars) and controls used to stabilize a β-

hairpin structure in a R2R3-P301L peptide (**Table 1**). B. Aggregation reactions of the tryptophan zipper

peptide and controls measured by ThT fluorescence (200 μM, 25 °C). ThT signals are an average of at

least 3 independent experiments. C. Schematic of proline and fluorinated proline analogs used to generate

cis and *trans* proline conformers at the position corresponding to P301 (red ball) in peptide models. D.

ThT aggregation reactions of the *cis*, *trans*, and neutral proline analogs substituted into the R2R3 peptide

(200 μM). ThT signals are an average of at least 6 independent experiments.

Figure 8. Molecular model of tau amyloid domain structural rearrangement and subsequent

aggregation. Naïve tau monomer (left) exists with a propensity to form a relatively collapsed structure,

which buries the amyloid domain [306]VQIVYK[311]. In the presence of disease-associated mutations, proline

isomerization events, or certain splice isoforms, the equilibrium is shifted to disfavor local compact

structure. This exposes the aggregation prone [306]VQIVYK[311] amyloid motif and enhances aggregation

propensity, leading to subsequent tau pathology.

References

1.   Cleveland, D. W., Hwo, S. Y. & Kirschner, M. W. Physical and chemical properties of purified tau
611   factor and the role of tau in microtubule assembly. *J. Mol. Biol.* **116,** 227–247 (1977).

612   2.   Eliezer, D. *et al.* Residual structure in the repeat domain of tau: Echoes of microtubule binding
613   and paired helical filament formation. *Biochemistry* **44,** 1026–1036 (2005).

614   3.   Fitzpatrick, A. W. P. *et al.* Cryo-EM structures of tau filaments from Alzheimer's disease. *Nature*
615   **547,** 185–190 (2017).

616   4.   Sanders, D. W. *et al.* Distinct tau prion strains propagate in cells and mice and define different
617   tauopathies. *Neuron* **82,** 1271–1288 (2014).

618   5.   Sawaya, M. R. *et al.* Atomic structures of amyloid cross-β spines reveal varied steric zippers.
619   *Nature* **447,** 453–457 (2007).

620   6.   Von Bergen, M. *et al.* Mutations of Tau Protein in Frontotemporal Dementia Promote
621   Aggregation of Paired Helical Filaments by Enhancing Local Beta-Structure. *J. Biol. Chem.* **276,**
622   48165–48174 (2001).

623   7.   Mirbaha, H. *et al.* Inert and seed-competent tau monomers elucidate the structural origins of
624   aggregation. *bioRxiv* (2017).

625   8.   Zhao, J. *et al.* Glycan Determinants of Heparin-Tau Interaction. *Biophys. J.* **112,** 921–932 (2017).

626   9.   Kellogg, E. H. *et al.* Near-atomic model of microtubule-tau interactions. **1780,** 1–9 (2018).

627   10.   Zhang, X. *et al.* RNA Stores Tau Reversibly in Complex Coacervates. *bioRxiv* 111245 (2017).
628   doi:10.1101/111245

629   11.   Mok, S.-A. *et al.* Mapping interactions with the chaperone network reveals factors that protect
630   against tau aggregation. *Nat. Struct. Mol. Biol.* **25,** 384–393 (2018).

631   12.   Baughman, H. E. R., Clouser, A. F., Klevit, R. E. & Nath, A. HspB1 and Hsc70 chaperones engage
632   distinct tau species and have different inhibitory effects on amyloid formation. *J. Biol. Chem.*
633   **293,** 2687–2700 (2018).

634   13.   Walker, S., Ullman, O. & Stultz, C. M. Using intramolecular disulfide bonds in tau protein to
635   deduce structural features of aggregation-resistant conformations. *J. Biol. Chem.* **287,** 9591–
636   9600 (2012).

637   14.   Kadavath, H. *et al.* Folding of the Tau Protein on Microtubules. *Angew. Chemie - Int. Ed.* **54,**
638   10347–10351 (2015).

639   15.   Mutations. (2017). Available at: http://www.alzforum.org/mutations/. (Accessed: 27th
640   September 2017)

641   16.   Holmes, B. B. *et al.* Proteopathic tau seeding predicts tauopathy in vivo.
642   doi:10.1073/pnas.1411649111

643   17.   Yoshiyama, Y. *et al.* Synapse Loss and Microglial Activation Precede Tangles in a P301S
644   Tauopathy Mouse Model. *Neuron* **53,** 337–351 (2007).

645   18.   Goedert, M., Wischik, C. M., Crowther, R. A., Walker, J. E. & Klug, A. Cloning and sequencing of
646   the cDNA encoding a core protein of the paired helical filament of Alzheimer disease:
647   identification as the microtubule-associated protein tau. *Proc. Natl. Acad. Sci.* **85,** 4051–4055
648   (1988).

649   19.   Williams, D. R. Tauopathies: Classification and clinical update on neurodegenerative diseases
650   associated with microtubule-associated protein tau. *Internal Medicine Journal* **36,** 652–660
651   (2006).

652   20.   Schoch, K. M. M. *et al.* Increased 4R-Tau Induces Pathological Changes in a Human-Tau Mouse
653   Model. *Neuron* **90,** 941–947 (2016).

654   21.   Hutton, M. *et al.* Association of missense and 5'-splice-site mutations in tau with the inherited
655   dementia FTDP-17. *Nature* **393,** 702–704 (1998).

656   22.   Stöhr, J. *et al.* A 31-residue peptide induces aggregation of tau's microtubule-binding region in

657  cells. *Nat. Chem.* **9,** 874–881 (2017).

658 23. Falcon, B., Zhang, W., Murzin, A. G., Murshudov, G. & Holly, J. Structures of filaments from Pick's
659  s disease reveal a novel tau protein fold. (2018).

660 24. Mylonas, E. *et al.* Domain conformation of tau protein studied by solution small-angle X-ray
661  scattering. *Biochemistry* **47,** 10345–10353 (2008).

662 25. Lin, M. M. Leveraging symmetry to predict self-assembly of multiple polymers. *Chem. Phys. Lett.*
663  **683,** 347–351 (2017).

664 26. Joachimiak, L. A., Walzthoeni, T., Liu, C. W., Aebersold, R. & Frydman, J. The structural basis of
665  substrate recognition by the eukaryotic chaperonin TRiC/CCT. *Cell* **159,** 1042–1055 (2014).

666 27. Tam, S. *et al.* The chaperonin TRiC blocks a huntingtin sequence element that promotes the
667  conformational switch to aggregation. *Nat. Struct. Mol. Biol.* **16,** 1279–1285 (2009).

668 28. Flory, P. J. & Volkenstein, M. Statistical mechanics of chain molecules. *Biopolymers* **8,** 699–700
669  (1969).

670 29. Mocanu, M.-M. *et al.* The potential for beta-structure in the repeat domain of tau protein
671  determines aggregation, synaptic decay, neuronal loss, and coassembly with endogenous Tau in
672  inducible mouse models of tauopathy. *J. Neurosci.* **28,** 737–748 (2008).

673 30. Rizzu, P. *et al.* High Prevalence of Mutations in the Microtubule-Associated Protein Tau in a
674  Population Study of Frontotemporal Dementia in the Netherlands. *Am. J. Hum. Genet.* **64,** 414–
675  421 (1999).

676 31. Micsonai, A. *et al.* Accurate secondary structure prediction and fold recognition for circular
677  dichroism spectroscopy. *PNAS* **112,** E3095–E3103 (2015).

678 32. Furman, J. L., Holmes, B. B. & Diamond, M. I. Sensitive Detection of Proteopathic Seeding
679  Activity with FRET Flow Cytometry. *J. Vis. Exp.* (2015). doi:10.3791/53205

680 33. Kaufman, S. K. *et al.* Tau Prion Strains Dictate Patterns of Cell Pathology, Progression Rate, and
681  Regional Vulnerability In Vivo. *Neuron* 1–17 (2016). doi:10.1016/j.neuron.2016.09.055

682 34. Kier, B. L., Shu, I., Eidenschink, L. a & Andersen, N. H. Stabilizing capping motif for beta-hairpins
683  and sheets. *Proc. Natl. Acad. Sci. U. S. A.* **107,** 10466–10471 (2010).

684 35. Vogel, M., Bukau, B. & Mayer, M. P. Allosteric regulation of Hsp70 chaperones by a proline
685  switch. *Mol. Cell* **21,** 359–367 (2006).

686 36. Gustafson, C. L. *et al.* A Slow Conformational Switch in the BMAL1 Transactivation Domain
687  Modulates Circadian Rhythms. *Mol. Cell* **66,** 447–457.e7 (2017).

688 37. Torbeev, V. Y. & Hilvert, D. Both the cis-trans equilibrium and isomerization dynamics of a single
689  proline amide modulate 2-microglobulin amyloid assembly. *Proc. Natl. Acad. Sci.* **110,** 20051–
690  20056 (2013).

691 38. Pastorino, L. *et al.* The prolyl isomerase Pin1 regulates amyloid precursor protein processing and
692  amyloid-beta production. *Nature* **440,** 528–34 (2006).

693 39. Lim, J. *et al.* Pin1 has opposite effects on wild-type and P301L tau stability and tauopathy. *J. Clin.*
694  *Invest.* **118,** 1877–1889 (2008).

695 40. Hasegawa, M., Smith, M. J., Iijima, M., Tabira, T. & Goedert, M. FTDP-17 mutations N279K and
696  S305N in tau produce increased splicing of exon 10. *FEBS Lett.* **443,** 93–96 (1999).

697 41. Pastor, P. *et al.* Familial atypical progressive supranuclear palsy associated with homozygosity for
698  the delN296 mutation in the tau gene. *Ann. Neurol.* **49,** 263–267 (2001).

699 42. Eichner, T. & Radford, S. E. A Generic Mechanism of B2-Microglobulin Amyloid Assembly at
700  Neutral pH Involving a Specific Proline Switch. *J. Mol. Biol.* **386,** 1312–1326 (2009).

701 43. Kontsekova, E., Zilka, N., Kovacech, B., Skrabana, R. & Novak, M. Identification of structural
702  determinants on tau protein essential for its pathological function: novel therapeutic target for
703  tau immunotherapy in Alzheimer's disease. *Alzheimers. Res. Ther.* **6,** 45 (2014).

704 44. Reumers, J., Maurer-Stroh, S., Schymkowitz, J. & Rousseau, F. Protein sequences encode

705  safeguards against aggregation. *Hum. Mutat.* **30,** 431–437 (2009).

706 45. O'Nuallain, B. *et al.* Kinetics and Thermodynamics of Amyloid Assembly Using a High-
707   Performance Liquid Chromatography-Based Sedimentation Assay. *Methods Enzymol.* **413,** 34–74
708   (2006).

709 46. Mirbaha, H., Holmes, B. B., Sanders, D. W., Bieschke, J. & Diamond, M. I. Tau trimers are the
710   minimal propagation unit spontaneously internalized to seed intracellular aggregation. *J. Biol.*
711   *Chem.* (2015). doi:10.1074/jbc.M115.652693

712 47. Shen, Y., Delaglio, F., Cornilescu, G. & Bax, A. TALOS+: A hybrid method for predicting protein
713   backbone torsion angles from NMR chemical shifts. *J. Biomol. NMR* **44,** 213–223 (2009).

714 48. Lange, O. F. *et al.* Determination of solution structures of proteins up to 40 kDa using CS-Rosetta
715   with sparse NMR data from deuterated samples. *Proc. Natl. Acad. Sci.* **109,** 10873–10878 (2012).

716 49. Raman, S. *et al.* NMR structure determination for larger proteins using backbone-only data.
717   *Science (80-.).* **327,** 1014–1018 (2010).

718 50. Rinner, O. *et al.* Identification of cross-linked peptides from large sequence databases. *Nat.*
719   *Methods* **5,** 315–318 (2008).

720 51. Walzthoeni, T. *et al.* False discovery rate estimation for cross-linked peptides identified by mass
721   spectrometry. *Nat. Methods* **9,** 901–903 (2012).

722 52. Grimm, M., Zimniak, T., Kahraman, A. & Herzog, F. XVis: A web server for the schematic
723   visualization and interpretation of crosslink-derived spatial restraints. *Nucleic Acids Res.* **43,**
724   W362–W369 (2015).

725 53. Barducci, A., Bussi, G. & Parrinello, M. Well-tempered metadynamics: A smoothly converging
726   and tunable free-energy method. *Phys. Rev. Lett.* **100,** (2008).

727 54. Lindorff-Larsen, K. *et al.* Improved side-chain torsion potentials for the Amber ff99SB protein
728   force field. *Proteins Struct. Funct. Bioinforma.* **78,** 1950–1958 (2010).

729 55. Berendsen, H. J. C., Postma, J. P. M., Van Gunsteren, W. F., Dinola, A. & Haak, J. R. Molecular
730   dynamics with coupling to an external bath. *J. Chem. Phys.* **81,** 3684–3690 (1984).

731 56. Parrinello, M. & Rahman, A. Polymorphic transitions in single crystals: A new molecular
732   dynamics method. *J. Appl. Phys.* **52,** 7182–7190 (1981).

733 57. Feenstra, K. A., Hess, B. & Berendsen, H. J. C. Improving efficiency of large time-scale molecular
734   dynamics simulations of hydrogen-rich systems. *J. Comput. Chem.* **20,** 786–798 (1999).

735 58. Darden, T., York, D. & Pedersen, L. Particle mesh Ewald: An N·log(N) method for Ewald sums in
736   large systems. *J. Chem. Phys.* **98,** 10089–10092 (1993).

737 59. Tribello, G. A., Bonomi, M., Branduardi, D., Camilloni, C. & Bussi, G. PLUMED 2: New feathers for
738   an old bird. *Comput. Phys. Commun.* **185,** 604–613 (2014).

739 60. Abraham, M. J. *et al.* Gromacs: High performance molecular simulations through multi-level
740   parallelism from laptops to supercomputers. *SoftwareX* **1–2,** 19–25 (2015).

Figure 1

a Tau Missense Mutations

N-Term — Repeat Domain — C-Term

R1 R2 R3 R4

Tau-RD   R1 R2 R3 R4

^{306}VQIVYK^{311}

b ^{295}DNIKHVPGGGSVQIVYK^{311}

c CS-ROSETTA

d *ab initio* ROSETTA

e

Figure 2

a RMSD Matrix for Wild Type

Time (µs)

RMSD (nm) 0.7 — 3.8

b RMSD Matrix for P301L Mutant

Time (µs)

RMSD (nm) 0.7 — 3.5

c

Hairpin   Fibril

Free Energy (kJ/mole)

Wildtype
P301L

RMSD from hairpin (nm)

Figure 3

Figure 4

a

b

c

Figure 5

Figure 6

Figure 7

a

R2R3-P301L-Trp

Trp-R2R3-P301L

Trp-R2R3-P301L-Trp

R2R3-P301L

b

ThT Fluorescence (A.U.)

1000

0   24   48   72

Hours

— R2R3-P301L
— Trp-R2R3-P301L-Trp (no signal)
— R2R3-P301L-Trp
— Trp-R2R3-P301L

c

Pro   4,4-Pro

4R-Pro
trans

4S-Pro
cis

d

ThT Fluorescence (A.U.)

0   50   100   150

Hours

— R2R3-Trans
— R2R3-Cis (no ThT)
— R2R3-Ea (no ThT)

Figure 8

Buried Amyloid Motif   **Exposure of Amyloid Motif**   **Amyloid Assembly Pathology**