1
2 **Research Advance to Putzbach et al., eLife. 2017; 6.**
3
4
5
6 **CD95L mRNA is toxic to cells**
7
8

9 William Putzbach[1,4], Ashley Haluck-Kangas[1,4], Quan Q. Gao[1], Aishe A. Sarshad[3],
10 Elizabeth T. Bartom[2], Austin Stults[1], Abdul S. Qadir[1],
11 Markus Hafner[3] and Marcus E. Peter[1,2,]*

12
13 [1] Department of Medicine/Division Hematology/Oncology, Feinberg School of Medicine, and [2]
14 Department of Biochemistry and Molecular Genetics, Northwestern University, Chicago, IL
15 60611, USA; [3] Laboratory of Muscle Stem Cells and Gene Regulation, NIAMS, NIH, Bethesda,
16 MD 20892, USA.
17
18
19 *Corresponding author: Marcus Peter, E-mail: m-peter@northwestern.edu, phone: 312-503-
20 1291; FAX: 312-503-0189.
21
22 [4] Shared first authorship

24 Keywords:
25 RNAi, Fas, cancer, CRISPR, cell death, DISE

26
27
28

Abstract

CD95/Fas ligand binds to the death receptor CD95/Fas to induce apoptosis in sensitive cells. We

previously reported the CD95L mRNA is enriched in sequences that, when converted to

si/shRNAs, are toxic to cells (Putzbach et al., 2017). These si/shRNAs kill all cancer cells

through a RNAi off-target effect by targeting critical survival genes. We now report expression

of full-length CD95L mRNA, itself, is highly toxic to cells and induces a similar form of cell

death. We demonstrate that small RNAs derived from CD95L are loaded into the RNA induced

silencing complex (RISC) and that the RISC is required for the toxicity. Drosha and Dicer

knock-out cells are highly sensitive to this toxicity, suggesting that processing of CD95L mRNA

into small toxic RNAs is independent of both Dicer and Drosha. The data provide evidence that a

higher vertebrate transgene can be processed to RNAi-active small RNAs that elicit cellular

responses.

Introduction

Activation of CD95/Fas through interaction with its cognate ligand CD95L or receptor-activating antibodies induces apoptosis in sensitive cells (Suda, Takahashi, Golstein, & Nagata, 1993). Virtually all research on CD95 and CD95L has focused on the physical interaction between the two proteins and the subsequent protein-based signaling cascades (Algeciras-Schimnich et al., 2002; Fu et al., 2016; Nisihara et al., 2001; Schneider et al., 1997). However, we have recently shown that the mRNA of CD95 and CD95L harbor sequences that when converted into small interfering (si) or short hairpin (sh)RNAs, cause massive and robust toxicity in all tested cancer cells. These CD95/CD95L-derived si/shRNAs target a network of survival genes, resulting in the simultaneous activation of multiple cell death pathways through RNA interference (RNAi) in a process we called DISE (Death Induced by Survival gene Elimination) (Putzbach et al., 2017). We determined that for an si/shRNA to elicit this form of toxicity, only positions 2-7 of the guide strand, the 6mer seed sequence, are required (Putzbach et al., 2017). More recently, a screen of all 4096 6mer seeds revealed that optimal 6mer seed toxicity requires G-rich seeds targeting C-rich regions in the 3'UTRs of survival genes (Gao et al., 2018).

In this report, we show that expression of the CD95L mRNA, itself, is toxic to cells even without prior conversion to siRNAs. This toxicity is independent of the full-length CD95L protein or expression of the CD95 receptor and resembles DISE. The toxicity involves RNAi, and multiple small RNAs generated within cells from the mRNA of CD95L are loaded into the RNA-induced Silencing Complex (RISC), the key mediator of RNAi (Liu et al., 2004).

Results

CD95L mRNA is toxic to cells

By testing every possible shRNA derived from the CD95L open reading frame (ORF) or its 3'UTR, we recently found a high enrichment of toxic si/shRNAs derived from the CD95L ORF (Putzbach et al., 2017). Most recently we determined that the 6mer seed toxicity observed in many si/shRNAs is due to their nucleotide composition, with G-rich seeds being the most toxic (Gao et al., 2018). When reanalyzing the CD95L ORF-derived shRNAs, we found a significant correlation between the toxicity of the most toxic CD95L-derived shRNAs (Putzbach et al., 2017) and the seed toxicity of the same 6mer seed we recently determined in a screen of all 4096 6mer seeds (Gao et al., 2018). This suggests that CD95L-derived shRNAs kill cancer cells mainly through 6mer seed toxicity.

We therefore wondered whether expression of the CD95L ORF mRNA—without pre-
processing into artificial siRNAs—would be toxic to cells. Expression of CD95L protein in most
cells kills through induction of apoptosis. Consequently, expressing CD95L in HeyA8 cells,
which are highly sensitive to CD95 mediated apoptosis, killed cells within a few hours after
infection with a lentivirus encoding CD95L (*Figure 1A*, left panel). Interestingly, severe growth
reduction was seen without any signs of apoptosis (not shown) when a CD95L mutant, unable to
bind CD95, was expressed (CD95L^{MUT} in *Figure 1A*, left panel). This mutant carries a Y218R
point mutation, which prevents the CD95L protein from binding to CD95 (Schneider et al.,
1997), and is expressed at a similar level to wild type (wt) CD95L (*Figure 1B*). To prevent the
CD95L mRNA from producing full-length CD95L protein, we also introduced a premature stop
codon right after the start codon in the CD95L^{MUT} vector (CD95L^{MUT}NP). This construct
(containing 4 point mutations and confirmed to produce mRNA with no detectable full-length
CD95L protein, *Figure 1B*) was equally active in reducing the growth of HeyA8 cells when
compared to the CD95L^{MUT} vector (*Figure 1A*, left panel). This result suggested that the CD95L
mRNA could be toxic to HeyA8 cells without the CD95L protein inducing apoptosis. This was
confirmed by expressing the three CD95L constructs in the presence of the oligo-caspase
inhibitor zVAD-fmk (*Figure 1A*, center panel). With suppressed apoptosis, all three constructs
were now equally toxic to HeyA8 cells. Finally, we tested a HeyA8 CD95 k.o. clone confirmed
to express no CD95 protein (Putzbach et al., 2017). In these cells, without the addition of zVAD-
fmk, wt CD95L and CD95L^{MUT}NP were again equally active in severely reducing the growth of
the cells (*Figure 1A*, right panel). Together, these data suggested it is the CD95L mRNA that
killed the cells. Cell death was confirmed by quantifying nuclear fragmentation (*Figure 1C*). We
also detected a significant increase of ROS in cells expressing CD95L^{MUT}NP (*Figure 1D*), which
is a characteristic feature of DISE (Hadji et al., 2014; Patel & Peter, 2017). To exclude the
possibility that truncated CD95 protein or any part of the CD95 mRNA would play a role in this
toxicity, we deleted the CD95 gene in MCF-7 cells (*Figure 1 - figure supplement 1A-E*).
Overexpression of wild-type CD95L killed clone FA4 cells, which harbor a complete
homozygous deletion of the entire CD95 gene, as well as CD95 protein k.o. clone #21 cells that
retain some truncated mRNA expression (*Figure 1E*). To further distinguish the activities of the
CD95L mRNA from CD95L protein, we generated a CD95L expression construct in which we
introduced 308 (out of 846 nucleotides) silent mutations (CD95L SIL) (*Figure 1 - figure*
supplement 2A). The activity of this mutant construct to negatively affect cell growth of CD95

k.o. HeyA8 cells was compared to two independently cloned wt CD95L constructs (WT1 and
WT2 in ***Figure 1 - figure supplement 2B***). The mutant SIL construct was equally effective in
suppressing cell growth and produced about the same amount of CD95L mRNA (***Figure 1 -***
figure supplement 2C). However, the SIL construct only produced about 12 % of WT CD95L
protein (***Figure 1 - figure supplement 2C***), again supporting the observation that it is the CD95L
RNA and not the protein that elicits toxicity.

CD95L mRNA kills cells through DISE.
After infection with CD95L, CD95 k.o. HeyA8 cells exhibited morphological changes strikingly
similar to the changes seen in wt HeyA8 cells after introduction of a CD95L-derived shRNA
(shL3) (***Figure 2A, Video 1-4***) suggesting the cells died through a similar mechanism. To
determine the cause of cell death induced by CD95L mRNA in HeyA8 CD95 k.o. cells
molecularly, we performed an RNA-Seq analysis. We found that expression of CD95L caused
preferential downregulation of critical survival genes and not of nonsurvival genes in a control
set (***Figure 2B***). In addition, cell death induced by CD95L mRNA resulted in a substantial loss
of 11 of the 12 histones detected to be downregulated in cells treated with CD95 and CD95L-
derived sh/siRNAs (***Figure 2C***). Loss of histons is an early event during DISE (Putzbach et al.,
2017). A Metascape analysis demonstrated that nucleosome assembly, regulation of mitosis, and
genes consistent with the involvement of histones were among the most significantly
downregulated RNAs across all cells in which DISE was induced by any of the four sh/siRNAs
or by the expression of CD95L mRNA (***Figure 2D***). This suggests that CD95L mRNA kills cells
in the same way as CD95/L-derived si/shRNAs.

CD95L mRNA kills cells through RNAi
Given our previous work on CD95L-derived si/shRNA toxicity, we hypothesized that CD95L
mRNA kills cells through an RNAi-based mechanism—perhaps by being processed into small
RNAs that incorporate into the RISC. Drosha k.o. cells lacking the majority of endogenous
miRNAs, but retaining expression of Ago proteins, were shown to be hypersensitive to DISE
induced by si- and shRNAs (Putzbach et al., 2017). We interpreted this effect as being caused by
an increased pool of unoccupied RNAi machinery caused by the absence of most miRNAs.
Drosha k.o. cells were also hypersensitive to the expression of CD95L^{MUT}NP (***Figure 3A,***
p=0.014, according to a polynomial fitting model); Virtually all cells died (insert in ***Figure 3A***).

To directly determine whether the RISC is involved in the toxicity, we introduced CD95L into

CD95 k.o. HeyA8 cells after knocking down AGO2 (*Figure 3B*). Knock down of AGO2 was

efficient (insert in *Figure 3B*). Toxicity elicited by CD95L was severely blunted following

AGO2 knockdown, suggesting that AGO2 was required for CD95L mRNA to be toxic.

To test the hypothesis that Drosha k.o. cells were more sensitive because their RISC was not

occupied by large amounts of nontoxic miRNAs and to determine whether CD95L mRNA could

give rise to small RNAs that incorporate into the RISC, we pulled down AGO1-4-associated

RNAs and analyzed their composition in wt and Drosha k.o. cells after expressing the

CD95L^{MUT}NP mRNA. For the pull-down, we used a peptide—derived from the AGO-binding

partner GW182—recently described to bind to all four Ago proteins (Hauptmann et al., 2015).

As expected in wt HCT116 cells, large amounts of small RNAs (19-23nt in length) were detected

bound to the Ago proteins (*Figure 3C*). Both AGO1 and AGO2 were efficiently pulled down. In

contrast, in the Drosha k.o. cells, which cannot generate canonical miRNAs, only a low amount

of small RNAs was detected, confirming the absence of miRNAs in the RISC. Surprisingly, the

amount of pulled down Ago proteins was severely reduced despite the fact these Drosha k.o.

cells express comparable levels of AGO2 (Putzbach et al., 2017). This suggests the peptide did

not have access to the Ago proteins in Drosha k.o. cells, presumably because it only binds to Ago

proteins complexed with RNA as recently shown (Elkayam et al., 2017).

The analysis of all Ago-bound RNAs showed that in the wt cells, >98.4% of bound RNAs

were miRNAs. In contrast, only 34% of bound RNAs were miRNAs in Drosha k.o. cells (*Figure*

3D and data not shown). These include miRNAs that are processed independently of Drosha

such as miR-320a (Kim, Kim, & Kim, 2016). Consistently, this miRNA became a major RNA

species bound to Ago proteins in Drosha k.o. cells (*Figure 3D*). In both wt and Drosha k.o. cells,

a significant increase in CD95L-derived small RNAs bound to the Ago proteins was detected

compared to cells infected with pLenti empty vector. They corresponded to 0.0006% and 0.043%

of all the Ago-bound RNAs in the wt cells and Drosha k.o. cells, respectively. Toxicity of

CD95L mRNA was, therefore, not due to overloading the RISC. In the absence of most

miRNAs, the total amount of RNAs bound to Ago proteins in the Drosha k.o. cells was roughly

10% of the amount bound to Ago in wt cells (*Figure 3D*). The reduction of Ago-bound miRNAs

in Drosha k.o. cells (*Figure 3E*, top row) was paralleled by a substantial increase in binding of

other small RNAs to the Ago proteins (*Figure 3E*, bottom row). Interestingly, the amount of

Ago-bound CD95L-derived small RNAs was >100 times higher in the Drosha k.o. cells

compared to the wt cells (red columns in **_Figure 3E_**). These data support our hypothesis that

Drosha k.o. cells are more sensitive to CD95L mRNA-mediated toxicity due to their ability to

take up more toxic small CD95L-derived RNAs into the RISC in the absence of most miRNAs.

CD95L ORF is degraded into small RNA fragments that are then loaded into the RISC.

Interestingly, not only did Ago proteins in Drosha k.o. cells bind much more CD95L-derived

small RNAs than in the wt cells, but also the peak length of the most abundant Ago-bound RNA

species increased from 20 to 23 nt (**_Figure 4A_**, top panel). To determine the sites within the

CD95L mRNA that gave rise to small Ago-bound RNAs, we aligned all small Ago-bound RNAs

detected in all conditions to the CD95L ORF sequence (**_Figure 4B and C_**). We identified 22

regions in the CD95L ORF that gave rise to small RNAs that could be bound by Ago proteins

(**_Figure 4B_**). To determine whether these small RNAs were formed in the cytosol and then

loaded into the RISC, we also aligned all small RNAs in the total RNA fraction isolated from

CD95L^{MUT}NP expressing HCT116 Drosha k.o. cells with CD95L (**_Figure 4C_**). Interestingly,

very similar regions of small RNAs were found. Moreover, the mean as well as the peak of the

distribution of the read lengths of small RNAs bound to Ago proteins was smaller than in the

total small RNAs fraction (**_Figure 4A_**, center panel), suggesting these fragments were trimmed

to the appropriate length either right before they are loaded into the RISC or by the RISC itself.

This was most obvious for the small RNAs in cluster 3 (**_Figure 4B and C_**). We also noticed that

certain small RNAs were more abundant in the Ago-bound fraction when compared to total RNA

relative to all other RNAs. To determine whether this type of processing was specific for

HCT116 Drosha k.o. cells, we analyzed the Ago-bound small CD95L-derived RNAs in HeyA8

CD95 k.o. cells after expression of wt CD95L (**_Figure 4D_**) and compared them with the total

RNA fraction (**_Figure 4E_**). While we found fewer CD95L-derived reads in these cells, the

general location of some of the read clusters overlapped with the ones found in the Drosha k.o.

cells and again both the mean and peak of the distribution of RNA lengths was smaller in the

Ago-bound fraction versus the total RNA fraction (**_Figure 4A_**, bottom panel). Together, these

data suggest that CD95L mRNA can be processed into smaller RNA fragments, which are then

trimmed to a length appropriate for incorporation into the RISC.

Our data suggest that the CD95L mRNA, when overexpressed, is toxic to cells due to the

formation of Ago-bound small RNAs that are incorporated into the RISC and kill cells through

RNAi. This process is independent of Drosha. To determine whether Dicer is required for either

processing of CD95L mRNA or loading the small RNAs into the RISC, we expressed
CD95L^{MUT}NP in wt and Dicer k.o. HCT116 cells (***Figure 4F***). Dicer k.o. cells were still
sensitive to toxicity induced by CD95L mRNA expression, suggesting the toxicity of the CD95L
mRNA does not require the processing by either Drosha or Dicer. Using custom real-time qPCR
primers designed to specifically detect the small RNAs from clusters 8 and 21, we detected, in
both wt and Dicer k.o. cells over-expressing CD95L^{MUT}NP, fragments from these clusters
(***Figure 4G***), demonstrating that Dicer is not involved in processing CD95L mRNA.

All the reported small RNAs derived from CD95L corresponded to the sense strand of the
expressed mRNA, raising the question of how they could be processed into double-stranded
siRNAs in the absence of an antisense strand. To get a preliminary answer to this question, we
subjected the CD95L ORF mRNA sequence to a secondary structure prediction (***Figure 4 -***
figure supplement 1A). According to this analysis, the CD95L ORF mRNA forms a tightly
folded structure with many of the small RNAs of the 22 clusters juxtaposing each other in stem-
like structures creating regions of significant complementarity. These may provide the duplexes
needed to be processed and loaded into the RISC. Interestingly, many of the juxtaposing reads
were found in duplex structures with 3' overhangs. Three of these oligonucleotides (derived from
clusters 7, 15 and 22) when expressed as siRNAs were toxic to HeyA8, H460, M565 and 3LL
cells (***Figure 4 - figure supplement 1B***).

In summary, our data suggest that si- and/or shRNAs with certain seed sequences as they are
present in CD95 and CD95L and the entire CD95L ORF are toxic to cancer cells. The CD95L
mRNA is broken down into small RNA-active fragments that are loaded into the RISC and then
target critical survival genes. This results in cell death through 6mer seed toxicity. The process is
independent of both Drosha and Dicer. Finally, the data suggest that a high miRNA content, by
"filling up" the RISC, might render cells less sensitive to this form of cell death.

Discussion

We recently reported a novel form of cell death that was observed after expression of si/shRNAs
designed from the sequences of CD95/CD95L mRNA (Putzbach et al., 2017). More recently we
described that cells die from a loss of multiple survival genes through a mechanism we call 6mer
seed toxicity (Gao et al., 2018). The most toxic si/shRNAs derived from CD95 or CD95L were
found in the ORF of CD95L (Putzbach et al., 2017). This pointed toward the CD95L mRNA,
itself being toxic.

We now show that expression of full-length CD95L mRNA triggers toxicity that is

independent of the protein product and canonical apoptosis. This is intriguing considering

previous studies showed transgenic expression of CD95L using viruses killed multiple cancer

cells that were completely resistant to CD95 mediated apoptosis after addition of agonist anti-

CD95 antibodies (ElOjeimy et al., 2006; Hyer, Voelkel-Johnson, Rubinchik, Dong, & Norris,

2000; Sudarshan et al., 2005; Sun et al., 2012). These results were interpreted as intracellular

CD95L triggering apoptosis. However, we now provide an alternate explanation—namely, both

the CD95L protein *and* mRNA are toxic to cells through distinct mechanisms. The protein

induces apoptosis, and the mRNA induces toxicity through an RNAi-based mechanism.

We demonstrate that Dicer and Drosha are not involved in generating the Ago-bound

CD95L-derived fragments but there are several candidate RNases that are capable of processing

mRNAs. Given the differences in length distribution between the cytosolic versus Ago-bound

RNA fragments, it is likely the released CD95L-derived fragment intermediates are incorporated

into the RISC and then trimmed to the appropriate length by Ago. Indeed, a similar mechanism is

known to occur during the maturation of the erythropoietic miR-451, where the pre-miRNA is

first cleaved by AGO2 and then trimmed at the 3' end to the final mature form by the

exoribonuclease PARN (Yoda et al., 2013). Furthermore, a similar process occurs with the

recently identified class of Ago-bound RNAs called agotrons (Hansen et al., 2016), which

consist of an excised intron loaded into the RISC in a manner independent of Drosha or Dicer

pre-processing. After trimmed to the appropriate size, the guide RNAs in complex with the RISC

can regulate gene expression through RNAi.

Our data provide the first evidence of an overexpressed cDNA to be toxic via an RNAi-

dependent mechanism. It was first shown in plants that overexpressed transgenes can be

converted into RNAi active short RNA sequences (Hamilton & Baulcombe, 1999). Our data on

the effects of overexpressed CD95L RNA, while mechanistically distinct from what was

reported in plants, maybe the first example of transgene determining cell fate through the RNAi

mechanism in mammalian cells.

A major question that arises from our data is whether CD95L mRNA is toxic *in vivo*. We and

others have noticed upregulation of CD95L in multiple stress-related conditions such as after

treatment with chemotherapy ((Friesen, Fulda, & Debatin, 1999) and data not shown). While the

amount of CD95L mRNA and the level of upregulation alone may not be enough to be toxic, it

could be the combination of multiple RNA fragments, derived from multiple different mRNAs

that are generated to kill cells (Putzbach et al., 2018). We view CD95L as just one of many

RNAs that have this activity. Indeed, it is unlikely CD95L is the only gene whose mRNA is toxic

to cells, as this mRNA-based level of toxicity would be redundant with the potent killing

capacity of the CD95L protein. Also, upregulating an mRNA that, by itself, could decimate the

cells that would otherwise need to upregulate that mRNA to carry out their biological function in

the first place, such as activated T cells upregulating CD95L to mount an immune response,

would be self-defeating. Therefore, nature likely distributed this mRNA-based toxicity-inducing

capacity over many genes in the genome to prevent activating it when any one of those genes is

upregulated during specific cellular processes. It is more likely there exists an entire network of

these genes that can release toxic small RNAs when the appropriate stimulus is encountered.

Consistent with this hypothesis we recently identified other genes that contain sequences that

when converted to shRNAs kill cancer cells through 6mer seed toxicity (Patel & Peter, 2017).

Future work will be aimed at identifying additional genes and the mechanism through which they

are processed and under what conditions to kill cells.

Materials and methods

Key Resources Table

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
Gene (Homo sapiens)	CD95L	NA	NM_000639	
Gene (H. sapiens)	CD95	NA	NM_000043	
Cell line (H. sapiens)	MCF-7	ATCC	ATCC: HTB-22	Human adenocarcinoma of the mammary gland, breast; derived from metastatic site: pleural effusion
Cell line (H. sapiens)	MCF-7 CD95 ΔshR6 clone #21	this paper	NA	MCF-7 CD95 ΔshR6 clone #21 with homozygous 227 nucleotide deletion of the shR6 target site in CD95 (chr10:89,008,920-89,009,146; Human Dec. 2013 GRCh38/hg38 assembly) produced using CRISPR/Cas9 technology; verified homozygous CD95 protein knockout
Cell line (H. sapiens)	MCF-7 CD95 deletion clone FA4	this paper	NA	MCF-7 CD95 deletion clone FA4 with a homozygous deletion of the entire CD95 gene (chr10:88,990,657 - 89,015,785; Human Dec. 2013 GRCh38/hg38 assembly) produced using CRISPR/Cas9 technology; verified homozygous CD95 protein knockout
Cell line (H. sapiens)	HeyA8	PMID: 4016745	RRID: CVCL_8878	Human high grade ovarian serous adenocarcinoma; derived from parent Hey cells (RRID: CVCL_0297)

Cell line (H. sapiens)	HeyA8 shR6 k.o. clone #11, HeyA8 CD95 k.o.	PMID: 29063830	NA	HeyA8 CD95 k.o. clone with a homozygous 227 nucleotide deletion of the shR6 target site in CD95 (chr10:89,008,920-89,009,146; Human Dec. 2013 GRCh38/hg38 assembly) produced using CRISPR/Cas9 technology; verified homozygous CD95 protein knockout
Cell line (H. sapiens)	HCT116	Korean Collection for Type Cultures (KCTC)	KCTC: cat#HC19023; ATCC: CCL_247	Human colorectal carcinoma
Cell line (H. sapiens)	Drosha$^{-/-}$; Drosha$^{-/-}$ clone #40	Korean Collection for Type Cultures (KCTC); PMID: 26976605	KCTC: cat#HC19020	HCT116 clone #40 with homozygous protein knockout of Drosha; knockout achieved using CRISPR/Cas9 which resulted in a single nucleotide insertion in one allele and a 26 nucleotide deletion in the other
Cell line (H. sapiens)	Dicer$^{-/-}$; Dicer$^{-/-}$ clone #43	Korean Collection for Type Cultures (KCTC); PMID: 26976606	KCTC: cat#HC19023	HCT116 clone #43 with homozygous protein knockout of Dicer; knockout achieved using CRISPR/Cas9 which resulted in a three nucleotide insertion and 14 nucleotide deletion in one allele and a 35 nucleotide deletion in the other
Cell line (H. sapiens)	Dicer$^{-/-}$; Dicer$^{-/-}$ clone #45	Korean Collection for Type Cultures (KCTC); PMID: 26976607	KCTC: cat#HC19024	HCT116 clone #45 with homozygous protein knockout of Dicer; knockout achieved using CRISPR/Cas9 which resulted in a 53 nucleotide deletion in one allele and a 28 nucleotide deletion in the other
Cell line (H. sapiens)	293T	ATCC	ATCC: CRL-3216	Derived from HEK293 cells (ATCC: CRL-1573); express large T antigen; used for packaging viruses
Cell line (H. sapiens)	H460	ATCC	ATCC: #HTB-177	Human lung pleural effusion carcinoma
Cell line (Mus musculus)	3LL	ATCC	ATCC #CRL-1642	Mouse Lewis lung carcinoma

Cell line (Mus musculus)	M565	PMID: 25366259	NA	Mouse hepatocellular carcinoma isolated from naturally occurring tumor in a floxed CD95 background
Antibody	anti-human AGO1 (rabbit monoclonal)	Cell Signaling	Cell Signaling #5053	1:2000; for western blot; primary Ab
Antibody	anti-human AGO1 (rabbit polyclonal)	Abcam	Abcam #98056	1:2000; for western blot; primary Ab
Antibody	anti-human AGO2 (rabbit polyclonal)	Abcam	Abcam #32381	1:2000; for western blot; primary Ab
Antibody	Goat anti-rabbit, IgG-HRP	Southern Biotech	Southern Biotech: cat#SB-4030-05	1:5000; for western blot; secondary Ab
Antibody	Goat anti-rabbit, IgG-HRP	Cell Signaling	Cell Signaling: cat#7074	1:2000; for western blot; secondary Ab
Antibody	Goat anti-mouse; IgG1-HRP	Southern Biotech	Southern BioTech: cat#1070-05	1:5000; for western blot; secondary Ab
Recombinant protein reagent	LzCD95L	PMID: 14504390	NA	Leucine zipper tagged CD95L; recombinant protein
Chemical compound	CellTiter-Glo	Promega	Promega #G7570	Detects ATP release as a surrogate for cell death; read-out is fluorescence
Chemical compound	propidium iodide	Sigma-Aldrich	Sigma-Aldrich: cat#P4864	Used for subG1 flow cytometry analysis
Chemical compound	puromycin	Sigma-Aldrich	Sigma-Aldrich: cat#P9620	Used for selection of cells expressing puromycin resistance cassettes
Chemical compound	G418	Affymetrix	Affymetrix: cat#11379	Used for selection of cells expressing G418 resistance cassette
Chemical compound	2',7'-dichlorodihydrofluorescein diacetate	Thermofisher Scientific	Thermofisher Scientific #D399	Dye used for detecting ROS production

Chemical compound	zVAD-fmk	Sigma-Aldrich	Sigma-Aldrich: cat#V116	Used at 20uM; pan caspase inhibitor
Recombinant DNA reagent	pLenti-GIII-CMV-RFP-2A-Puro vector; pLenti	ABM Inc	NA	pLenti control empty lentiviral vector; carries an RFP-2a-puromycin resistance cassette
Recombinant DNA reagent	pLenti-CD95L	this paper	NA	pLenti-GIII-CMV-RFP-2A-Puro vector that expresses human wild type CD95L cDNA (NM_000639.2); used to express wt human CD95L upon infection with lentiviral particles
Recombinant DNA reagent	pLenti-CD95L^{MUT}	this paper	NA	pLenti-GIII-CMV-RFP-2A-Puro vector that expresses human CD95L cDNA (NM_000639.2) with 2 nucleotide substitutions in codon 218 (TAT -> CGT) resulting in replacement of tyrosine for arginine (Y218R mutation); unable to bind CD95
Recombinant DNA reagent	pLenti-CD95L^{MUT}NP	this paper	NA	pLenti-GIII-CMV-RFP-2A-Puro vector that expresses human CD95L cDNA (NM_000639.2) with both the Y218R mutation and a single nucleotide substitution at the second codon (CAG -> TAG), resulting in a premature stop codon right after the start codon
Recombinant DNA reagent	pLenti-CD95L^{SIL}	this paper	NA	pLenti-GIII-CMV-RFP-2A-Puro vector that expresses human CD95L cDNA (NM_000639.2) with all codons containing synonymous mutations except for select codons in the proline-rich domain to meet IDT synthesis criteria

Transfected construct	gRNA scaffold	PMID: 23287722	IDT: synthesized as gene block	455 nucleotide CRISPR/Cas9 gRNA scaffold synthesized as a gene block; contains promoter, gRNA scaffold, target sequence, and termination sequence; scaffold transcribes gRNAs that target Cas9 endonuclease to cut at target sites; target sequences consist of 19 nucleotides that are complementary to the target site of choice; co-transfected with Cas9 to catalyze cleavage.
Transfected construct	pMJ920 Cas9 plasmid	Addgene; PMID: 23386978	Addgene: cat#42234	Plasmid that expresses a human codon-optimized Cas9 tagged with GFP and HA; used to express Cas9 for CRISPR-mediated deletions.
Chemical compound	Lipofectamine 2000	ThermoFisher Scientific	ThermoFisher Scientific: cat#11668019	Transfection reagent
Chemical compound	Lipofectamine RNAiMAX	ThermoFisher Scientific	ThermoFisher Scientific: cat#13778150	Transfection reagent; used for transfection of small RNAs such as siRNAs
Commercial assay or kit	StrataClone Blunt PCR Cloning Kit	Agilent Technologies	Agilent Technologies: cat#240207	Used to blunt-end clone the gRNA scaffolds into the pSC-B plasmid
Genetic reagent	Taqman Gene expression master mix	ThermoFisher Scientific	#4369016	
Sequence-based reagent	shR6 flanking Fr primer	IDT	IDT: custom DNA oligo	Fr primer that flanks shR6 site; used to detect 227 nt shR6 deletion; 5'-GGTGTCATGCTGTGACTGTTG-3'
Sequence-based reagent	shR6 flanking Rev primer	IDT	IDT: custom DNA oligo	Rev primer that flanks shR6 site; used to detect 227 nt shR6 deletion; 5'-TTTAGCTTAAGTGGCCAGCAA-3'

Sequence-based reagent	shR6 internal Rev primer	IDT	IDT: custom DNA oligo	Rev primer that overlaps with the shR6 site; used to detect 227 nt shR6 deletion; 5'-AAGTTGGTTTACATCTGCAC-3'
Sequence-based reagent	CD95 flanking Fr primer	IDT	IDT: custom DNA oligo	Fr primer that flanks the CD95 gene; used to detect CD95 gene deletion; 5'-TGTTTAATATAGCTGGGGCTATGC-3'
Sequence-based reagent	CD95 flanking Rev primer	IDT	IDT: custom DNA oligo	Rev primer that flanks the CD95 gene; used to detect CD95 gene deletion; 5'-TGGGACTCATGGGTTAAATAGAAT-3'
Sequence-based reagent	CD95 internal Rev primer	IDT	IDT: custom DNA oligo	Rev internal primer that targets within the CD95 gene; used to detect CD95 gene deletion; 5'-GACCAGTCTTCTCATTTCAGAGGT-3'
Sequence-based reagent	siScr	IDT; Dharmacon	Dharmacon #D-001810-02-05	control non-targeting siRNA; sense: UGGUUUACAUGUUGUGUGA
Sequence-based reagent	c7/1	IDT	custom siRNA; antisense strand corresponds to cluster 7 CD95L sequence	antisense: 5'-AUUGGGCCUGGGGAUGUUU-3'; antisense strand designed with 3' deoxy AA; complementary sense strand has 3' deoxy TT and 2'-O-methylation at the first two positions
Sequence-based reagent	c7/2	IDT	custom siRNA; antisense strand corresponds to cluster 7 CD95L sequence	antisense: 5'-CCUGGGGAUGUUUCAGCUC-3'; antisense strand designed with 3' deoxy AA; complementary sense strand has 3' deoxy TT and 2'-O-methylation at the first two positions

Sequence-based reagent	c11	IDT	custom siRNA; antisense strand corresponds to cluster 11 CD95L sequence	antisense: 5'-CCAACUCAAGGUCCAUGCC-3'; antisense strand designed with 3' deoxy AA; complementary sense strand has 3' deoxy TT and 2'-O-methylation at the first two positions
Sequence-based reagent	c15/1	IDT	custom siRNA; antisense strand corresponds to cluster 15 CD95L sequence	antisense: 5'-AAACUGGGCUGUACUUUGU-3'; antisense strand designed with 3' deoxy AA; complementary sense strand has 3' deoxy TT and 2'-O-methylation at the first two positions
Sequence-based reagent	c15/2	IDT	custom siRNA; antisense strand corresponds to cluster 15 CD95L sequence	antisense: 5'-AACUGGGCUGUACUUUGUA-3'; antisense strand designed with 3' deoxy AA; complementary sense strand has 3' deoxy TT and 2'-O-methylation at the first two positions
Sequence-based reagent	c16/1	IDT	custom siRNA; antisense strand corresponds to cluster 16 CD95L sequence	antisense: 5'-CAACAACCUGCCCCUGAGC-3'; antisense strand designed with 3' deoxy AA; complementary sense strand has 3' deoxy TT and 2'-O-methylation at the first two positions
Sequence-based reagent	c16/2	IDT	custom siRNA; antisense strand corresponds to cluster 16 CD95L sequence	antisense: 5'-AACUCUAAGCGUCCCCAGG-3'; antisense strand designed with 3' deoxy AA; complementary sense strand has 3' deoxy TT and 2'-O-methylation at the first two positions

Sequence-based reagent	c21	IDT	custom siRNA; antisense strand corresponds to cluster 21 CD95L sequence	antisense: 5'-UCAACGUAUCUGAGCUCUC-3'; antisense strand designed with 3' deoxy AA; complementary sense strand has 3' deoxy TT and 2'-O-methylation at the first two positions
Sequence-based reagent	c22	IDT	custom siRNA; antisense strand corresponds to cluster 22 CD95L sequence	antisense: 5'-AAUCUCAGACGUUUUUCGG-3'; antisense strand designed with 3' deoxy AA; complementary sense strand has 3' deoxy TT and 2'-O-methylation at the first two positions
Sequence-based reagent	siScr pool	Dharmacon	D-001810-10	control non-targeting siRNA pool
Sequence-based reagent	SMARTpool siRNA targeting AGO2	Dharmacon	L-004639-00-0005	siRNA pool designed to target AGO2
Sequence based reagent (human)	GAPDH primer	Thermofisher Scientific	Hs00266705_g1	RT-qPCR; control probe
Sequence based reagent (human)	CD95L primers	Thermofisher Scientific	Hs00181226_g1; Hs00181225_m1	RT-qPCR
Sequence based reagent (human)	CD95 primers	Thermofisher Scientific	Hs00531110_m1; Hs00236330_m1	RT-qPCR
Sequence based reagent (human)	CD95L^{SIL} primer	Thermofisher Scientific	assay ID: APNKTUD	Custom RT-qPCR primer designed using the Thermofisher Scientific design tool to detect CD95L^{SIL} mRNA

Sequence based reagent (human)	Cluster 8 CD95L small RNA primer	Thermofisher Scientific	custom probe	Custom RT-qPCR primer designed using the Thermofisher Scientific design tool at https://www.thermofisher.com/order/custom- genomic-products/tools/small-rna to specifically detect small RNAs from cluster 8 of CD95L (5'-AAGGAGCTGGCAGAACTCCGAGA-3')
Sequence based reagent (human)	Cluster 21 CD95L small RNA primer	Thermofisher Scientific	custom probe	Custom RT-qPCR primer designed using the Thermofisher Scientific design tool at https://www.thermofisher.com/order/custom- genomic-products/tools/small-rna to specifically detect small RNAs from cluster 21 of CD95L (5'-TCAACGTATCTGAGCTCTCTC-3')
Sequence based reagent (human)	z30 primer	Thermofisher Scientific	ThermoFisher Scientific #4427975	RT-qPCR for small RNA; control probe
Peptide, recombinant protein	Flag-GST-T6B peptide	PMID: 26351695	NA	Peptide derived from GW182 used to pull down AGO1 to 4
Commercial assay or kit	anti-Flag M2 magnetic beads	Sigma-Aldrich	Sigma-Aldrich #M8823	

Reagents and antibodies

All reagents and antibodies were described previously (Putzbach et al., 2017) except those referenced in the following paragraphs.

Cell lines

HeyA8 (RRID:CVCL_8878) and HeyA8 CD95 knock-out cells, HCT116 (ATCC #CCL-247; RRID:CVCL_0291) and HCT116 Drosha knock-out and Dicer knock-out cells, MCF-7 cells (ATCC #HTB-22; RRID:CVCL_0031), and 293T (ATCC #CRL-3216; RRID:CVCL_0063) cells were cultured as described previously (Putzbach et al., 2017). The MCF-7 CD95 knock-out

and deletion cells were cultured in RPMI 1640 medium (Cellgro #10-040-CM), 10% heat-

inactivated FBS (Sigma-Aldrich), 1% L-glutamine (Mediatech Inc), and 1%

penicillin/streptomycin (Mediatech Inc). H460 (ATCC #HTB-177; RRID:CVCL_0459) cells

were cultured in RPMI1640 medium (Cellgro Cat#10-040) supplemented with 10% FBS (Sigma

Cat#14009C) and 1% L-Glutamine (Corning Cat#25-005). 3LL cells (ATCC #CRL-1642;

RRID:CVCL_4358) were cultured in DMEM medium (Gibco Cat#12430054) supplemented

with 10% FBS and 1% L-Glutamine. Mouse hepatocellular carcinoma cells M565 cells were

described previously (Ceppi et al., 2014) and cultured in DMEM/F12 (Gibco Cat#11330)

supplemented with 10% FBS, 1% L-Glutamine and ITS (Corning #25-800-CR). All cell lines

were authenticated using STR profiling and tested monthly for mycoplasm using PlasmoTest

(Invitrogen).

Plasmids and constructs

The pLenti-CD95L was synthesized by sub-cloning an insert containing the CD95L ORF

(NM_000639.2; synthesized by IDT as minigene with flanking 5' NheI RE site and 3' XhoI RE

sites in pIDTblue vector) into the pLenti-GIII-CMV-RFP-2A-Puro vector (ABM Inc). The insert

and the backbone were digested with NheI (NEB #R0131) and XhoI (NEB #R0146) restriction

enzymes. Subsequent ligation with T4 DNA ligase created the pLenti-CD95L vector. The

pLenti-CD95L^{MUT} vector was created by sub-cloning a CD95L cDNA insert with 2 nucleotide

substitutions in codon 218 (*TAT -> CGT)* resulting in replacement of tyrosine for arginine, which

has been described to inhibit binding to CD95 (Schneider et al., 1997) into the pLenti-GIII-

CMV-RFP-2A-Puro vector. The pLenti-CD95L^{MUT}NP vector was created by inserting a CD95L

ORF cDNA sequence containing both the Y218R mutation and a single nucleotide substitution at

the second codon (*CAG -> TAG*), resulting in a premature stop codon right after the start codon,

into the pLenti-GIII-CMV-RFP-2A-Puro vector. The pLenti-CD95L^{SIL} was created by sub-

cloning a mutant CD95L ORF cDNA sequence with codons synonymously mutated (***Figure 1 -***

figure supplement 2A) to the next most highly utilized codon in human cells (exceptions were

made within the proline rich domain to meet gene synthesis design criteria.) into the pLenti-GIII-

CMV-RFP-2A-Puro vector.

Overexpression of CD95L cDNAs

All lentiviral constructs were generated in 293T cells as described previously (Putzbach et al.,

2017). HeyA8 and MCF-7 (and all derivative cell lines) cells overexpressing wild type CD95L

and mutant CD95L cDNAs were generated by seeding cells at 100,000 cells per well in a 6-well

plate and infecting cells with lentivirus generated in 293T cells (500 µl viral supernatant per

well) with 8 µg/ml Polybrene. Media was changed next day. Selection was started either during

the evening of the same day or on following day with 3 µg/ml puromycin. HCT116, HCT116

Drosha knockout, and HCT116 Dicer knockout cells (Kim et al., 2016) overexpressing CD95L

cDNAs were generated by seeding cells at 100,000 cells per well in a 24-well plate or 500,000

cells per well in a 6-well plate and infecting cells with lentivirus generated in 293T cells (100 µl

virus per 24-well or 500 µl per 6-well) in the presence of 8 µg/ml Polybrene. Media was changed

the next day, and cells were selected with 3 µg/ml puromycin the following day. Infection with

empty pLenti was always included as a control.

To assess toxicity of overexpressing CD95L cDNAs, cells infected with these constructs

were plated in on a 96-well plate 1 day after selection in the presence of puromycin (uninfected

cells were all dead after 1 day in presence of puromycin); Cell confluency was assessed over

time using the IncuCyte as described previously (Putzbach et al., 2017).

To assess overexpression of CD95L cDNAs in apoptosis-sensitive HeyA8 cells in *Figure*

1A, infection with CD95L lentiviruses were done in 96-well plate using 50 µl of virus in the

presence of 20 µM zVAD-fmk (Sigma-Aldrich #V116) and 8 µg/ml Polybrene; media was

changed next day in the presence of 20 µM zVAD-fmk; 3 µg/ml puromycin was added the

following day. Infection with the CD95L constructs for the RT-qPCR and Western blot in

Figure 1B were done in a 6-well plate in the presence of 20 µM zVAD-fmk.

For the experiment in *Figure 3B*, HeyA8 CD95 knock-out cells were reverse transfected

in a 6-well plate; 100,000 cells were plated in wells with either the On-TargetPlus non-targeting

siRNA (Dharmacon #D-001810-10) or siAGO2 pool (Dharmacon ##L-004639-00-0005) at 25

nM complexed with 1 µl RNAiMax. After ~24 hrs, the cells were infected with either pLenti or

pLenti-CD95L (500 µl of viral supernatant [25% of total volume]). Next day, media was

replaced, and cells were expanded to 10 cm plates. The following day, 3 µg/ml puromycin

was added. When puromycin selection was complete (one day later), the 750-1,500 cells were

plated per well in a 96-well plate and put in the IncuCyte machine to assess cell confluency over

time.

CRISPR deletions

We co-transfected a Cas9-expressing plasmid (Jinek et al., 2013) and two gRNAs that target

upstream and downstream to delete an entire section of DNA as described previously (Putzbach

et al., 2017). The gRNA scaffold was used as described (Mali et al., 2013). The gRNAs were

designed using the algorithm found at http://crispr.mit.edu; only gRNAs with a score above 50

were considered.

A deletion of 227 nucleotides in exon 4 of CD95 in MCF-7 cells (ΔshR6, clone #21) was

generated using gRNAs described previously (Putzbach et al., 2017). Deletion of this site results

in a frame-shift mutation that causes a protein-level knock-out (Putzbach et al., 2017). PCR with

flanking external primers (Fr: 5'-*GGTGTCATGCTGTGACTGTTG*-3' and Rev: 5'-

TTTAGCTTAAGTGGCCAGCAA-3') and internal primers (Fr primer and the internal Rev primer

5'-*AAGTTGGTTTACATCTGCAC*-3') was used to screen for single cell clones that harbor a

homozygous deletion.

The two sequences targeted by the flanking gRNAs for the deletion of the entire CD95 gene

were 5'-GTCAGGGTTCGTTGCACAAA-3' and 5'-TGCTTCTTGGATCCCTTAGA-3'. For

detection of the CD95 gene deletion, the flanking external primers were 5'-

TGTTTAATATAGCTGGGGCTATGC-3' (Fr primer) and 5'-

TGGGACTCATGGGTTAAATAGAAT-3' (Rev primer), and the internal reverse primer was 5'-

GACCAGTCTTCTCATTTCAGAGGT-3'. After screening the clones, Sanger sequencing was

performed to confirm the proper deletion had occurred.

Real-Time quantitative PCR

The relative expression of specific mRNAs was quantified as described previously (Gao et al.,

2018). The primer/probes purchased from ThermoFisher Scientific were GAPDH

(Hs00266705_g1), human CD95L (Hs00181226_g1 and Hs00181225_m1), human CD95

(Hs00531110_m1 and Hs00236330_m1), and a custom primer/probe to detect the CD95L^{SIL}

mRNA (designed using the Thermofisher Scientific custom design tool; assay ID: APNKTUD).

Custom RT-qPCR probes designed to specifically detect small RNA species were used to

detect CD95L fragments in ***Figure 4G***. These probes were designed using ThermoFisher's

Custom TaqMan Small RNA Assay Design Tool (https://www.thermofisher.com/order/custom-

genomic-products/tools/small-rna/) to target the cluster 8 sequence (5'-

AAGGAGCTGGCAGAACTCCGAGA-3') and the cluster 21 sequence (5'-

TCAACGTATCTGAGCTCTCTC-3'). Detection of these fragments involves a two-step

amplification protocol used to detect microRNAs. In the first step, the High-Capacity cDNA

reverse transcription kit is used to selectively reverse transcribe the two clusters to be quantified

using specific primers and 20 nM RNA input following the manufacturer's protocol. The cDNA

is diluted 1:5. The qPCR reaction mixture is composed of the diluted cDNA, the custom probes,

and the Taqman Universal PCR Master Mix (Applied Biosystems #43240018). Reactions were

performed in triplicate. Ct values were determined using the Applied Biosystems 7500 Real

Time PCR system with a thermocycle profile of 50°C for two min (step one), 95°C for 10 min

(step two), and then 40 cycles of 95°C for 15 s (step three) and 60°C for 1 min (step four). The

ΔΔCt values between the small RNA of interest and the control were calculated to determine

relative abundance of the small RNA. Samples were normalized to Z30 (ThermoFisher Scientific

#4427975).

Western blot analysis

Detection of human CD95, CD95L and Ago proteins was done via Western blot as described

previously (Putzbach et al., 2017).

CD95 surface staining

Flow cytometry was used to quantify the level of membrane-localized CD95 as described

previously (Putzbach et al., 2017).

Cell death quantification (DNA fragmentation) and ROS production

The percent of subG1 nuclei (fragmented DNA) was determined by PI staining/flow cytometry

as described previously (Putzbach et al., 2017). ROS production was quantified using the cell-

permeable indicator 2',7'-dichlorodihydrofluorescein diacetate (ThermoFisher Scientific #D399)

as previously described (Hadji et al., 2014).

Assessing cell growth and fluorescence over time

After treatment/infection, cells were seeded in a 96-well plate at least in triplicate. Images were

captured at indicated time points using an IncuCyte ZOOM live cell imaging system (Essen

BioScience) with a 10x objective lens. Percent confluence and total fluorescent integrated

intensity was calculated using the IncuCyte ZOOM software (version 2015A).

Infection of cells for Ago-pull down and small RNA-Seq analysis

HeyA8 ΔshR6 clone #11 cells were seeded at 75,000 cells per well on 6-well plates, and the

HCT116 and HCT116 Drosha knock-out cells were both seeded at 500,000 per well on 6-well

plates. The HeyA8 ΔshR6 clone #11 cells were infected with 0.5 mL of empty pLenti or pLenti-

CD95L-WT viral supernatant per well. The HCT116 and HCT116 Drosha knockout cells were

infected with 0.5 mL empty pLenti or pLenti-CD95L^{MUT}NP viral supernatant per well. Media

was changed the next day and the cells were pooled and expanded to multiple 15 cm dishes.

Selection with 3 μg/mL puromycin began the following day. The next day, the HeyA8 ΔshR6

clone #11 infected cells were seeded at 600,000 cells per dish in multiple 15 cm dishes; the

HCT116 and HCT116 Drosha knock-out cells were seeded at 5 million cells per dish in multiple

15 cm dishes. Two days later, each of the samples was pelleted and split in two: one pellet was

lysed and processed for small RNA sequencing, and the other pellet was flash frozen in liquid

nitrogen. The pellets were stored at -80°C until they could be used for the Ago pull-down

experiment. The purpose of splitting the sample was so that we could compare the total cellular

pool of small RNAs to the fraction that was bound to the RISC. This way, the processing

CD95L-derived fragments from the full-length mRNA in the cytosol to the final mature RISC-

bound form could be mapped. This was all done in duplicate.

RNA-Seq analysis

Total RNA was isolated using the miRNeasy Mini Kit (Qiagen, #74004) following the

manufacturer's instructions. An on-column digestion step using the RNase-free DNase Set

(Qiagen #79254) was included. Both small and large mRNA libraries were generated and

sequenced as described previously (Putzbach et al., 2017). Reads were trimmed with TrimGalore

and then aligned to the hg38 assembly of the human genome with Tophat. Raw read counts were

assigned to genes using HTSeq and differential gene expression was analyzed with the R

Bioconductor EdgeR package (Robinson, McCarthy, & Smyth, 2010).

Ago pull down and RNA-Seq analysis of bound small RNAs

Cell pellets were harvest at 50 hours after plating (122 hours after infection) and were flash

frozen in liquid nitrogen. The pellets were stored at -80°C until ready for further processing.

Between 10 and 25 x 10^6 cells were lysed in NP40 lysis buffer (20 mM Tris, pH 7.5, 150 mM

NaCl, 2 mM EDTA, 1% (v/v) NP40, supplemented with phosphatase inhibitors) on ice for 15

minutes. The lysate was sonicated 3 times for 30 s at 60% amplitude (Sonics, VCX130) and

cleared by centrifugation at 12,000g for 20 minutes. AGO1-4 were pulled down by using 500 µg

of Flag-GST-T6B peptide (Hauptmann et al., 2015) and with 60 µl anti-Flag M2 magnetic beads

(Sigma-Aldrich) for 2 hrs at 4°C. The pull-down was washed 3 times in NP40 lysis buffer.

During the last wash, 10% of beads were removed and incubated at 95°C for 5 minutes in 2x

SDS-PAGE sample buffer. Samples were run on a 4-12% SDS-PAGE and transferred to

nitrocellulose membrane. The pull-down efficiency was determined by immunoblotting against

AGO1 (Cell Signaling #5053; RRID:AB_10695871 and Abcam #98056; RRID:AB_10680548)

and AGO2 (Abcam #32381; RRID:AB_867543). To the remaining beads 500 µl TRIzol reagent

were added and the RNA extracted according to the manufacturer's instructions. The RNA pellet

was diluted in 20 µl of water. The sample was split, and half of the sample was dephosphorylated

with 0.5 U/µl of CIP alkaline phosphatase at 37°C for 15 min and subsequently radiolabeled with

0.5 µCi γ-^{32}P-ATP and 1 U/µl of T4 PNK kinase for 20 min at 37°C. The AGO1-4 interacting

RNAs were visualized on a 15% urea-PAGE. The remaining RNA was taken through a small

RNA library preparation as previously described (Hafner et al., 2012). Briefly, RNA was ligated

with 3' adenylated adapters and separated on a 15% denaturing urea-PAGE. The RNA

corresponding to insert size of 19-35 nt was eluted from the gel, ethanol precipitated followed by

5' adapter ligation. The samples were separated on a 12% Urea-PAGE and extracted from the

gel. Reverse transcription was performed using Superscript III reverse transcriptase and the

cDNA amplified by PCR. The cDNA was sequenced on Illumina HiSeq 3000. Adapter

sequences: Adapter 1 – NNTGACTGTGGAATTCTCGGGTGCCAAGG; Adapter 2 –

NNACACTCTGGAATTCTCGGGTGCCAAGG, Adapter 3 –

NNACAGAGTGGAATTCTCGGGTGCCAAGG, Adapter 4 –

NNGCGATATGGAATTCTCGGGTGCCAAGG, Adapter 47 –

NNTCTGTGTGGAATTCTCGGGTGCCAAGG, Adapter 48 –

NNCAGCATTGGAATTCTCGGGTGCCAAGG, Adapter 49 –

NNATAGTATGGAATTCTCGGGTGCCAAGG, Adapter 50 –

NNTCATAGTGGAATTCTCGGGTGCCAAGG. RT primer sequence:

GCCTTGGCACCCGAGAATTCCA; PCR primer sequences:

CAAGCAGAAGACGGCATACGAGATCGTGATGTGACTGGAGTTCCTTGGCACCCGAG

AATTCCA. To identify CD95L-derived small RNAs among the sequenced reads, a BLAST

database was generated from each set of reads, and blastn was used to query the CD95L ORF

(derived from NM_000639.2) against reads from cells infected with pLenti-CD95L and to query

the CD95L^{MUT}NP ORF sequence against reads from cells infected with CD95L^{MUT}NP. The only

reads considered further were those matching a CD95L sequence with an e-value of less than

0.05 and 100% identity across the entire length of the read. This resulted in the loss of a few

reads less than 19/20 nt in length. The filtered BLAST hits were converted to a bed formatted

file, describing the locations of reads relative to the relevant CD95L sequence, and the R

package Sushi was used to plot the bed files and generate Figures 4B-E.

Assessing toxicity of CD95L-derived small RNAs

To determine whether guide RNAs derived from the over-expressed CD95L mRNA could evoke

toxicity, the small CD95L-derived RNA reads (corresponding to different clusters shown in

Figure 4C) bound to AGO from the HCT116 Drosha knock-out cells were converted to siRNAs.

First, all reads less than 18 nucleotides were filtered out, as these do not efficiently incorporate

into the RISC. siRNAs were designed with antisense strands identical to these CD95L-derived

sequences that mapped to areas of the CD95L mRNA secondary structure (*Figure 4 - figure*

supplement 1A) that are predicted to form duplexes. These sequences were designed as 19

nucleotide oligos with a 3' deoxy AA. The complementary sense strand was designed with a 3'

deoxy TT and 2'-O-methylation at the first two positions to prevent its incorporation into the

RISC. These oligos were ordered from IDT and annealed to form the final siRNAs. The

sequences of the antisense strands (corresponding to the CD95L mRNA-derived cluster

fragments) were as follows: 5'-AUUGGGCCUGGGGGAUGUUU-3' (c7/1), 5'-

CCUGGGGAUGUUUCAGCUC-3' (c7/2), 5'-CCAACUCAAGGUCCAUGCC-3' (c11), 5'-

AAACUGGGCUGUACUUUGU-3' (c15/1), 5'- AACUGGGCUGUACUUUGUA-3' (c15/2),

5'- CAACAACCUGCCCCUGAGC-3' (c16/1), 5'- AACUCUAAGCGUCCCCAGG-3' (c16/2),

5'- UCAACGUAUCUGAGCUCUC-3' (c21), and 5'- AAUCUCAGACGUUUUUCGG-3'

(c22).

These eight siRNAs were reverse transfected into HeyA8, H460, M565, and 3LL cells using

RNAiMAX transfection reagent (ThermoFisher Scientific) at 10 nM in triplicate as previously

described (Murmann et al., 2018). The non-targeting (NT) and siL3 siRNAs, as described

previously (Putzbach et al., 2017), were used as a negative and positive control, respectively.

Cell death was quantified via ATP release 96 hours after transfection using CellTiter-Glo

(Promega). The % viability was calculated in relation to the RNAiMAX-only treatment

structure (*Figure 4 - figure supplement 1B*).

Statistical analyses

Continuous data were summarized as means and standard deviations (except for all IncuCyte

experiments where standard errors are shown) and dichotomous data as proportions. Continuous

data were compared using t-tests for two independent groups and one-way ANOVA for 3 or

more groups. For evaluation of continuous outcomes over time, two-way ANOVA was used with

one factor for the treatment conditions of primary interest and a second factor for time treated as

a categorical variable to allow for non-linearity.

The effects of treatment on wild-type versus Drosha knock-out cells were statistically

assessed by fitting regression models that included linear and quadratic terms for value over

time, main effects for treatment and cell type, and two- and three-way interactions for treatment,

cell-type and time. The three-way interaction on the polynomial terms with treatment and cell

type was evaluated for statistical significance since this represents the difference in treatment

effects over the course of the experiment for the varying cell types.

GSEA used in *Figure 2B* was performed using the GSEA v2.2.4 software from the Broad

Institute (www.http://software.broadinstitute.org/gsea); 1000 permutations were used. The

Sabatini gene lists were set as custom gene sets to determine enrichment of survival genes versus

the nonsurvival control genes in downregulated genes from the RNA-Seq data as done

previously (Putzbach et al., 2017); p-values below 0.05 were considered significantly enriched.

Genes with an average normalized read expression (across both pair of duplicates) below 3 were

excluded so as to only include genes that are truly expressed. The GO enrichment analysis shown

in *Figure 2D* was performed with all genes that after alignment and normalization were found to

be at least 1.5 fold downregulated with an adjusted p-value of <0.05 using the software available

on www.Metascape.org and default running parameters. The other data sets used in this analysis

(HeyA8 cells transfected with a toxic siRNA targeting CD95L siL3 and 293T infected with toxic

shRNAs targeting CD95L shL1 and shL3 and HeyA8 cells infected with a toxic shRNA

targeting CD95 shR6) were previously described (Putzbach et al., 2017).

All statistical analyses were conducted in Stata 14 or R 3.3.1.

Data availability

RNA sequencing data generated for this study is available in the GEO repository: GSE103631

(https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE103631, reviewer access token:

etgbqyaenvirjqn) and GSE114425

(https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE114425, reviewer access token:

edgdoaocjberbyr).

Acknowledgements

We are grateful to Siquan Chen for testing small CD95L-derived siRNAs. M.H. and A.A.S were supported by the Intramural Research Program of NIAMS. A.A.S. acknowledges support by the Swedish Research Council postdoctoral fellowship. This work was funded by training grant T32CA009560 (to W.P. and A.H.K.), R50CA221848 (to E.T.B.), and R35CA197450 (to M.E.P.).

Author contributions

W.P. planned the study and performed experiments. A.H.K., Q.Q.G., A.S.Q., and A.S. performed experiments, D.M.S. provided biostatistics support, E.B. provided biocomputational support, A.A.S. performed the Ago pull down experiments, M.H. provided assistance and discussions on the mechanism of RNAi and the RISC, and M.E.P. directed the study and M.E.P. and W.P. wrote the manuscript.

Competing financial interests

The authors declare no competing financial interests.

Figure legends

Figure 1.

The CD95L RNA is toxic to cells.

(**A**) *Left*: Schematic of the different CD95L mutants used. *Right:* Percent cell confluence over

time of HeyA8 parental cells in the absence (*left panel*) or in the presence of 20 μM zVAD-fmk

(*center panel*), or CD95 k.o. cells (*right panel*) after expression of CD95L constructs. Data are

representative of one to three independent experiments. Values were calculated from samples

done in triplicate or quadruplicate shown as mean ± SE. (**B**) *Left*: Western blot analysis of

HeyA8 cells overexpressing different CD95L mutant RNAs. Cells expressing CD95L^{MUT} or

CD95L were pretreated with 20 μM zVAD-fmk. Note the small amount of truncated CD95L in

cells infected with CD95L MUT-NP does not have CD95 binding activity. Very similar data

were obtained when the constructs were expressed in either CD95 k.o. HeyA8 cells (clone #11)

or NB 7 cells, both without treatment with zVAD (data not shown). *Right*: RT-qPCR analysis for

CD95L of the same samples. Data are representative of two independent experiments. Each bar

represents mean ± S.D. of three replicates. (**C**, **D**) Quantification of cell death (**C**) and ROS

production (**D**) in cells expressing either pLenti (v) or pLenti-CD95L (L) at different time points

(days after infection). Data are representative of two independent experiments. Each bar

represents mean ± SE of three replicates. * p<0.05, ** p<0.001, *** p<0.0001, unpaired t-test.

(**E**) Confluency over time of the MCF-7 complete CD95 k.o. FA4 clone (right) or a MCF-7 clone

#21 in which we deleted the shR6 site resulting in an out-of-frame shift after infection with either

vector control or wt CD95L. Data are representative of two independent experiments. Each data

point represents mean ± SE of three replicates.

Figure 1 - figure supplement 1.

Generation of complete CD95 k.o. MCF-7 cells.

(**A**) Schematic of the genomic locations and sequences of the gRNAs used to excise the entire

CD95 gene in MCF-7 cells. PAM site is underlined. (**B**) PCR with flanking (*top panels*) and

internal (*bottom panels*) primers used to confirm the absence of the CD95 gene in MCF-7 clones.

Parental (Par.) cells and three clones infected with Cas9 only (Cas9) and two complete k.o.

clones (F2 and FA4) are shown. (**C**) RT-qPCR analysis of the indicated clones using primers

spanning either exon 1/2 or exon 2/3 of the CD95 gene. (**D**) Surface staining for CD95 of one wt

and one k.o. clone. (**E**) Western blot analysis of all clones.

Figure 1 - figure supplement 2.

Toxicity of CD95L mRNA is independent of CD95L protein expression.

(**A**) Schematic showing the positions of the silent mutation of the CD95L^{SIL} compared to wild

type CD95L. (**B**) Percent cell confluence over time of HeyA8 CD95 k.o. cells over-expressing

empty pLenti, wild-type CD95L (from two separately cloned viruses), or the CD95L^{SIL}. (**C**) RT-

qPCR analysis and Western blot (*inset*) of wild type CD95L and CD95L^{SIL} mutant mRNAs in the

over-expressing cells shown in **B**.

Figure 2.

Toxicity induced by CD95L overexpression is reminiscent of DISE.

(**A**) Phase-contrast images of HeyA8 and HeyA8 CD95 k.o. cells after infection with pLKO-

shScr/shL3 or pLenti/pLenti-CD95L, respectively, at the indicated time point. (**B**) Gene set

enrichment analysis for the 1846 survival genes (*top panel*) and the 416 nonsurvival genes

(*bottom panel*) identified in the Sabatini study (Putzbach et al., 2017; Wang et al., 2015) of

mRNAs downregulated in CD95L expressing HeyA8 CD95 k.o. cells compared to HeyA8
CD95 k.o. cells infected with pLenti virus. p-values indicate the significance of enrichment. (**C**)
The genes downregulated in all cells including the ones downregulated following introduction of
one of four si/shRNAs (Putzbach et al., 2017) derived from either CD95 or CD95L (see ***Figure***
2D) and the ones overexpressing CD95L ORF as described in **B**. (**D**) Metascape analysis of 5
RNA Seq data sets analyzed. The boxed GO term clusters were highly enriched in all 5 data sets.
624
Figure 3.
Small RNAs generated in cells expressing CD95L mRNA are loaded into the RISC.
(**A**) Percent cell confluence over time of HCT116 parental (*left*) or Drosha k.o. (*right*) cells after
infection with CD95^{MUT}NP. Data are representative of three independent experiments. Each data
point represents the mean ± SE of three replicates. *Inset:* Phase contrast images of Drosha k.o.
cells 9 days after infection with either empty vector or CD95L^{MUT}NP. (**B**) Percent cell
confluence of HeyA8 CD95 k.o. cells transfected with either non-targeting siRNA (siCtr) or a
pool of 4 siRNAs targeting AGO2 following subsequent infection with either empty pLenti (EV)
or pLenti CD95L. *Inset:* Western blot showing successful knock-down of human AGO2. (**C**)
Top: autoradiograph on RNAs pulled down with the Ago binding peptide. *Bottom*: Western blot
analysis of pulled down Ago proteins. v, pLenti; L, pLenti-CD95L expressing cells. (**D**) Pie
charts showing the relative ratio of small RNAs pulled down with the Ago proteins in wt and
Drosha k.o. cells. Depicted are all the amounts of all small RNAs that contributed at least 0.01%
to the total RNA content. Only in the Drosha k.o. cells was a significant amount of CD95L
derived Ago bound reads found. They represented the 75th most abundant small RNA species
(arrow). The average number of total sequenced reads (of two duplicates) are shown for each
condition. (**E**) *Top*: Number of reads (normalized per million) of the top six most abundant small
RNAs in the RISC of either HCT116 wt-pLenti or -pLenti-CD95L cells. *Bottom*: Number of
reads (per million) of the top five genes with small RNAs most abundant in the RISC of either
HCT116 Drosha k.o. pLenti, or -pLenti-CD95L cells and of CD95L derived. Note: miR-21 is not
included as it is already shown in the top row. Bottom right panel: Abundance of Ago bound
CD95L derived small RNAs. Shown in all panels is the abundance of RNAs in the four samples.
Rep 1 and rep 2, replicate 1 and 2.
648
Figure 4
The entire CD95L mRNA gives rise to small RNAs that bind to the RISC.
(**A**) Length distribution of CD95L derived reads in various analyses. (**B, C**) Read alignment with
CD95L^{MUT}NP ORF of analyses of small RNAs pulled down with Ago proteins from HCT116 wt
(**B**, top) and Drosha k.o. (**B**, bottom) cells and of total small RNAs from HCT116 Drosha k.o.
cells (**C**) after infection with CD95L^{MUT}NP. (**D, E**) Read alignment with wt CD95L ORF of
analyses of small RNAs pulled down with Ago proteins (**D**) or total small RNAs (**E**) from
HeyA8 CD95 k.o. cells after infection with wt CD95L. (**F**) Percent cell confluence over time of
HCT116 parental (*top*) or Dicer k.o. (clone #43) (*bottom*) cells after infection with CD95^{MUT}NP.
(Dicer k.o. clone #45, gave a similar result, data not shown). Data are representative of two
independent experiments. Each data point represents the mean ± SE of three replicates. (**G**) RT-
qPCR analysis of clusters 8 and 21 in HCT116 parental, Dicer k.o. (clone #43), and Drosha k.o.
cells after infection with CD95^{MUT}NP. Each bar represents mean ± S.D. of three replicates. v,
vector, L, CD95L expressing cells.
663
Figure 4 - figure supplement 1

Predicted secondary structure of CD95L ORF and toxicity of CD95L-derived small
RNAs after conversion to siRNAs.
(**A**) The CD95L^{MUT}NP RNA was subjected to a RNA secondary structure analysis
(http://rna.tbi.univie.ac.at) using default settings. The locations of 22 reads representative of the
22 read clusters are shown. Regions with potential duplex formation are boxed. The
oligonucleotides that were found to be toxic when expressed as siRNAs are circled. (**B**) Toxicity
of the eight siRNAs designed using the CD95L-derived small RNA fragments bound to Ago as
the antisense strand sequences 96 hours post-transfection in the indicated cell lines. Each data
point represents the mean ± SE of three replicates.
674
675
676
677 **Supplementary Videos:**
678
679 **Video 1: CD95 k.o. HeyA8 cells (clone 11) infected with pLenti control virus.**
680
681 **Video 2: CD95 k.o. HeyA8 cells (clone 11) infected with pLenti-CD95Lvirus.**
682
683 **Video 3: HeyA8 cells infected with pLKO-shScr.**
684
685 **Video 4: HeyA8 cells infected with pLKO-shL3.**
686

687 **References**
688
Algeciras-Schimnich, A., Shen, L., Barnhart, B. C., Murmann, A. E., Burkhardt, J. K., & Peter,
M. E. (2002). Molecular ordering of the initial signaling events of CD95. *Mol Cell Biol,*
22(1), 207-220.
Ceppi, P., Hadji, A., Kohlhapp, F., Pattanayak, A., Hau, A., Xia, L., . . . Peter, M. E. (2014).
CD95 and CD95L promote and protect cancer stem cells. *Nature Commun, 5*, 5238.
Elkayam, E., Faehnle, C. R., Morales, M., Sun, J., Li, H., & Joshua-Tor, L. (2017). Multivalent
Recruitment of Human Argonaute by GW182. *Mol Cell, 67*(4), 646-658 e643.
doi:10.1016/j.molcel.2017.07.007
ElOjeimy, S., McKillop, J. C., El-Zawahry, A. M., Holman, D. H., Liu, X., Schwartz, D. A., . . .
Norris, J. S. (2006). FasL gene therapy: a new therapeutic modality for head and neck
cancer. *Cancer gene therapy, 13*(8), 739-745. doi:10.1038/sj.cgt.7700951
Friesen, C., Fulda, S., & Debatin, K. M. (1999). Cytotoxic drugs and the CD95 pathway.
Leukemia, 13(11), 1854-1858.
Fu, Q., Fu, T. M., Cruz, A. C., Sengupta, P., Thomas, S. K., Wang, S., . . . Chou, J. J. (2016).
Structural Basis and Functional Role of Intramembrane Trimerization of the Fas/CD95
Death Receptor. *Mol Cell, 61*(4), 602-613. doi:10.1016/j.molcel.2016.01.009
Gao, Q. Q., Putzbach, W., Murmann, A. E., Chen, S., Ambrosini, G., Peter, J. M., . . . Peter, M.
E. (2018). 6mer Seed Toxicity Determines Strand Selection in miRNAs. *BioRxive*.
doi:https://doi.org/10.1101/284406
Hadji, A., Ceppi, P., Murmann, A. E., Brockway, S., Pattanayak, A., Bhinder, B., . . . Peter, M.
E. (2014). Death induced by CD95 or CD95 ligand elimination. *Cell Reports, 10*, 208-
222.
Hafner, M., Renwick, N., Farazi, T. A., Mihailovic, A., Pena, J. T., & Tuschl, T. (2012).
Barcoded cDNA library preparation for small RNA profiling by next-generation
sequencing. *Methods, 58*(2), 164-170. doi:10.1016/j.ymeth.2012.07.030
Hamilton, A. J., & Baulcombe, D. C. (1999). A species of small antisense RNA in
posttranscriptional gene silencing in plants. *Science, 286*(5441), 950-952.
Hansen, T. B., Veno, M. T., Jensen, T. I., Schaefer, A., Damgaard, C. K., & Kjems, J. (2016).
Argonaute-associated short introns are a novel class of gene regulators. *Nature*
communications, 7, 11538. doi:10.1038/ncomms11538
Hauptmann, J., Schraivogel, D., Bruckmann, A., Manickavel, S., Jakob, L., Eichner, N., . . .
Meister, G. (2015). Biochemical isolation of Argonaute protein complexes by Ago-APP.
Proc Natl Acad Sci U S A, 112(38), 11841-11845. doi:10.1073/pnas.1506116112
Hyer, M. L., Voelkel-Johnson, C., Rubinchik, S., Dong, J., & Norris, J. S. (2000). Intracellular
Fas ligand expression causes Fas-mediated apoptosis in human prostate cancer cells
resistant to monoclonal antibody-induced apoptosis. *Mol Ther, 2*(4), 348-358.
Kim, Y. K., Kim, B., & Kim, V. N. (2016). Re-evaluation of the roles of DROSHA, Export in 5,
and DICER in microRNA biogenesis. *Proc Natl Acad Sci U S A, 113*(13), E1881-1889.
doi:10.1073/pnas.1602532113
Liu, J., Carmell, M. A., Rivas, F. V., Marsden, C. G., Thomson, J. M., Song, J. J., . . . Hannon,
G. J. (2004). Argonaute2 is the catalytic engine of mammalian RNAi. *Science,*
305(5689), 1437-1441.
Mali, P., Yang, L., Esvelt, K. M., Aach, J., Guell, M., DiCarlo, J. E., . . . Church, G. M. (2013).
RNA-guided human genome engineering via Cas9. *Science, 339*(6121), 823-826.
doi:10.1126/science.1232033

Murmann, A. E., Gao, Q. Q., Putzbach, W. E., Patel, M., Bartom, E. T., Law, C. Y., . . . Peter,
M. E. (2018). Small interfering RNAs based on huntingtin trinucleotide repeats are
highly toxic to cancer cells. *EMBO Rep, 19*(3). doi:10.15252/embr.201745336

Nisihara, T., Ushio, Y., Higuchi, H., Kayagaki, N., Yamaguchi, N., Soejima, K., . . . Yagita, H.
(2001). Humanization and epitope mapping of neutralizing anti-human Fas ligand
monoclonal antibodies: structural insights into Fas/Fas ligand interaction. *J Immunol,*
167(6), 3266-3275.

Patel, M., & Peter, M. E. (2017). Identification of DISE-inducing shRNAs by monitoring cellular
responses. *Cell Cycle*, 0. doi:10.1080/15384101.2017.1383576

Putzbach, W., Gao, Q. Q., Patel, M., Haluck-Kangas, A., Murmann, A. E., & Peter, M. E.
(2018). DISE - A Seed Dependent RNAi Off-Target Effect that Kills Cancer Cells.
Trends in Cancer, 4, 10-19.

Putzbach, W., Gao, Q. Q., Patel, M., van Dongen, S., Haluck-Kangas, A., Sarshad, A. A., . . .
Peter, M. E. (2017). Many si/shRNAs can kill cancer cells by targeting multiple survival
genes through an off-target mechanism. *Elife, 6*, e29702.

Robinson, M. D., McCarthy, D. J., & Smyth, G. K. (2010). edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. *Bioinformatics, 26*(1),
139-140. doi:10.1093/bioinformatics/btp616

Schneider, P., Bodmer, J. L., Holler, N., Mattmann, C., Scuderi, P., Terskikh, A., . . . Tschopp, J.
(1997). Characterization of Fas (Apo-1, CD95)-Fas ligand interaction. *J Biol Chem,*
272(30), 18827-18833.

Suda, T., Takahashi, T., Golstein, P., & Nagata, S. (1993). Molecular cloning and expression of
the Fas ligand, a novel member of the tumor necrosis factor family. *Cell, 75*(6), 1169-
1178.

Sudarshan, S., Holman, D. H., Hyer, M. L., Voelkel-Johnson, C., Dong, J. Y., & Norris, J. S.
(2005). In vitro efficacy of Fas ligand gene therapy for the treatment of bladder cancer.
Cancer gene therapy, 12(1), 12-18. doi:10.1038/sj.cgt.7700746

Sun, H., Liu, Y., Bu, D., Liu, X., Norris, J. S., & Xiao, S. (2012). Efficient growth suppression
and apoptosis in human laryngeal carcinoma cell line HEP-2 induced by an adeno-
associated virus expressing human FAS ligand. *Head Neck, 34*(11), 1628-1633.
doi:10.1002/hed.21985

Wang, T., Birsoy, K., Hughes, N. W., Krupczak, K. M., Post, Y., Wei, J. J., . . . Sabatini, D. M.
(2015). Identification and characterization of essential genes in the human genome.
Science, 350(6264), 1096-1101. doi:10.1126/science.aac7041

Yoda, M., Cifuentes, D., Izumi, N., Sakaguchi, Y., Suzuki, T., Giraldez, A. J., & Tomari, Y.
(2013). Poly(A)-specific ribonuclease mediates 3'-end trimming of Argonaute2-cleaved
precursor microRNAs. *Cell Rep, 5*(3), 715-726. doi:10.1016/j.celrep.2013.09.029

A

CD95

chr10:88,990,657   -   89,015,785 (25,129 base pairs)

GTCAGGGGTTCGTTGCACAAATGG

gRNA#1

Human Dec. 2013 (GRCh38/hg38) Assembly

TGCTTCTTGGATCCCTTAGAAGG

gRNA#2

B

| | Par. | Cas9 | | | CD95 k.o. | |
| | | #1 | #2 | #3 | F2 | FA4 |

764bp — Ext. pr.

481bp — Int. pr.

C

Rel fold change

Exons 1-2   Exons 2-3

Par.   #1   #2   #3   F2   FA4

Cas9   CD95 k.o.

D

MCF-7 Cas9 #1   MCF-7 CD95 k.o. FA4

% of Max

IgG
anti-CD95

18.1   0.66

CD95

E

| | Par. | Cas9 | | | CD95 k.o. | |
| kDa | | #1 | #2 | #3 | F2 | FA4 |

51 —
48 —   ⇐ CD95

42 —   Actin

A

B

C

A

HeyA8 - shScr   HeyA8 - shL3   HeyA8 Cl11 - pLenti   HeyA8 Cl11 - pLenti-CD95L

135 hrs   135 hrs   164 hrs   164 hrs

B

CD95L overexpression

C

HIST2H3D HIST1H2AB HIST1H1D HIST1H1C
HIST3H2BB HIST1H4A HIST1H3J HIST1H2BN
HIST1H2BF HIST1H4D HIST1H3A

D

Figure 4 - figure suppl. 1