A new *in vitro* assay measuring direct interaction of nonsense suppressors with the

eukaryotic protein synthesis machinery

Martin Y. Ng[1], Haibo Zhang[1¥], Amy Weil[1†], Vijay Singh[2], Ryan Jamiolkowski[2], Alireza

Baradaran-Heravi[3], Michel Roberge[3], Allan Jacobson[4], Westley Friesen[5], Ellen Welch[5], Yale E.

Goldman[2], and Barry S. Cooperman[1*]

[1]Department of Chemistry, University of Pennsylvania, Philadelphia, PA 19104

[2]Department of Physiology, Perelman School of Medicine, University of Pennsylvania,

Philadelphia, PA 19104

[3]Department of Biochemistry and Molecular Biology, University of British Columbia, Vancouver,

BC, Canada V6T 1Z3

[4]Department of Microbiology and Physiological Systems, University of Massachusetts Medical

School, Worcester, MA 01655

[5]PTC Therapeutics, 100 Corporate Court, South Plainfield, NJ 07080

* To whom to address inquiries

¥ Present address: Spark Therapeutics, 3737 Market Street, Philadelphia, PA, 19104

† Present address: AdMed, Inc., 122 Union Square Drive, New Hope, PA, 18938

Abstract

Nonsense suppressors (NonSups) treat premature termination codon (PTC) disorders by inducing

the selection of near cognate tRNAs at the PTC position, allowing readthrough of the PTC and

production of full-length protein. Studies of NonSup-induced readthrough of eukaryotic PTCs

have been carried out using animals, cells or crude cell extracts. In these studies, NonSups can

promote readthrough directly, by binding to components of the protein synthesis machinery, or

indirectly, by inhibiting nonsense-mediated mRNA decay or by other mechanisms. Here we utilize

a highly-purified *in vitro* system (Zhang et al., 2016. *eLife* **5**: e13429) to measure exclusively direct

NonSup-induced readthrough. Of 17 previously identified NonSups, 13 display direct effects,

apparently via at least two different mechanisms. We can monitor such direct effects by single

molecule FRET (smFRET). Future smFRET experiments will permit elucidation of the

mechanisms by which NonSups stimulate direct readthrough, aiding ongoing efforts to improve

the clinical usefulness of NonSups.

Introduction

Premature termination codons (PTCs) arise as a consequence of nonsense mutations and lead to the replacement of an amino acid codon in mRNA by one of three stop codons, UAA, UGA or UAG (Brenner, *et al.*, 1965; Shalev and Baasov, 2014; Keeling, *et al.*, 2014), resulting in inactive truncated protein products. Nonsense mutations constitute ~20% of transmitted or *de novo* germline mutations (Salvatori, *et al.*, 2009; Goldmann, *et al.*, 2012; Stenson, *et al.*, 2017). Globally, there are ~7000 genetically transmitted disorders in humans and ~11% of all human disease mutations are nonsense mutations (Loudon, 2013). Clearly, millions of people worldwide would benefit from effective therapies directed toward PTC suppression. Clinical trials have begun to evaluate the treatment of PTC disorders with therapeutic agents called nonsense suppressors (NonSups) (Peltz, *et al.*, 2013; McDonald, *et al.*, 2017; Zainal Abidin, *et al.*, 2017). NonSups induce the selection of near cognate tRNAs at the PTC position, and insertion of the corresponding amino acid into the nascent polypeptide, a process referred to as "readthrough", which restores the production of full length functional proteins, albeit at levels considerably reduced from wild-type. Even low rates of readthrough can improve clinical outcomes when essential proteins are completely absent. Examples of such essential proteins include Cystic Fibrosis Transmembrane Regulator (CFTR) (Brodlie, *et al.*, 2015), dystrophin, and the cancer tumor suppressors adenomatous polyposis coli (APS) (Floquet, *et al.*, 2011; Zilberberg, *et al.*, 2010) protein and p53 (Miyaki, *et al.*, 2002; Floquet, *et al.*, 2011; Roy, *et al.*, 2016; Baradaran-Heravi, *et al.*, 2016).

In vitro, ex vivo, and *in vivo* experiments and clinical trials have identified a diverse structural set of NonSups as candidates for PTC suppression therapy (Figure 1), including aminoglycosides (Shalev and Baasov, 2014; Bidou, *et al.*, 2017; Oishi, *et al.*, 2015; Duscha, *et al.*, 2014; Floquet, *et al.*, 2012; Sangkuhl, *et al.*, 2004; Fuchshuber-Moraes, *et al.*, 2011; Cogan, *et al.*,

2014; Baradaran-Heravi*, et al.*, 2017), ataluren (Peltz*, et al.*, 2013; Roy*, et al.*, 2016; Welch*, et al.*,

2007) and ataluren-like molecules (Du*, et al.*, 2009; Du*, et al.*, 2013; Gómez-Grau*, et al.*, 2015)

and others (Zilberberg*, et al.*, 2010; Arakawa*, et al.*, 2003; Hamada*, et al.*, 2015; Caspi*, et al.*,

2016; Mutyam*, et al.*, 2016). To date, only one NonSup, ataluren (known commercially as

Translarna), has been approved in the EU for clinical use, but this approval is limited to treatment

of patients with nonsense-mediated Duchenne muscular dystrophy. The clinical utility of other

NonSups, such as aminoglycosides, is restricted, in part, by their toxic side effects. A critical

barrier to development of NonSups that are more clinically useful is the paucity of information

regarding the precise mechanisms by which these molecules stimulate readthrough. All prior

results measuring nonsense suppressor-induced readthrough (NSIRT) of eukaryotic PTCs have

been carried out using animals, intact cells or crude cell extracts. In such systems, NonSups can

promote readthrough directly, by binding to one or more of the components of the protein synthesis

machinery, or indirectly, either by inhibiting nonsense-mediated mRNA decay (NMD) (He and

Jacobson, 2015), or by modulating processes altering the cellular activity levels of protein

synthesis machinery components (Feng*, et al.*, 2014; Keeling, 2016). These assays thus measure a

quantity we define as TOTAL-NSIRT. This multiplicity of possible mechanisms of nonsense

suppression within TOTAL-NSIRT has complicated attempts to determine the precise

mechanisms of action of specific NonSups and limited the use of rational design in identifying

new, more clinically useful NonSups.

Recently, we developed a highly purified, eukaryotic cell-free protein synthesis system

(Zhang*, et al.*, 2016) that we apply here to examine the direct effects of the NonSups on the protein

synthesis machinery, which we define as DIRECT-NSIRT. Our results allow us to distinguish

NonSups acting directly on the protein synthesis machinery from those that act indirectly and

suggest that NonSups having DIRECT-NSIRT effects can be divided into at least two distinctive

structural groups that induce nonsense suppression by different mechanisms. We also demonstrate

the potential of using single molecule fluorescence resonance energy transfer (smFRET)

experiments to elucidate the details of such mechanisms.

Materials and Methods

<u>Nonsense suppressors (Figure 1)</u>. The following NonSups were obtained from commercial sources:

gentamicin mixture and G418 (Sigma), nourseothricin sulfate, a mixture of streptothricins D and

F (Gold Biotechnology), doxorubicin (Fisher Scientific), escin and tylosin (Alfa Aesar),

azithromycin (APExBIO). Gentamicins B and B1 were prepared as described (Baradaran-Heravi,

et al., 2017). PTC Therapeutics supplied the following NonSups: ataluren sodium salt, RTC13,

GJ071, GJ072, and gentamicin X2. Negamycin was a gift from Alexander Mankin, University of

Illinois at Chicago. NB84 and NB124 (Bidou, *et al.*, 2017), currently available as ELX-02 and

ELX-03, respectively, from Eloxx Pharmaceuticals (Waltham, MA), were gifts from Timor

Baasov (Technion, Haifa).

<u>Ribosomes and factors</u>. Shrimp (*A. salina*) ribosome subunits were prepared from dried frozen

commercial cysts as described (Zhang, *et al.*, 2016; Iwasaki and Kaziro, 1979) with slight

modifications. Shrimp cysts (Pentair Aquatic Ecosystems) (425 g) were ground open using a

blender in the presence of buffer M (30 mM HEPES-KOH, pH 7.5, 50 mM KCl, 10 mM $MgCl_2$,

8.5% mannitol, 0.5 mM EDTA, 2 mM DTT, 1 mM PMSF, 1:3000 RNasin (New England Biolabs)

(500 mL), and two Protease Inhibitor Complete minitablets (Roche). Cyst debris was removed by

two centrifugations at 30,000 x g for 15 min at 4°C. 80S ribosomes in the supernatant were

precipitated by adding 175 mL of 4.5% PEG 20k (Ben-Shem, *et al.*, 2011) and resuspended in 60

mL of dissociation buffer 1 (20 mM HEPES-KOH, pH 7.5, 500 mM KCl, 2 mM $MgCl_2$, 6.8%

sucrose 2 mM DTT, 1:1000 RNasin, 2 protease minitablets). Puromycin was added to a final

concentration of 2 mM, and the resulting solution was incubated on ice for 30 min, then at 37 °C

for 15 min. 40S and 60S subunits (approximately 6,000 A_{260} units) were then resolved by a 10-

30% hyperbolic sucrose gradient centrifugation for 16 h in a Beckman Ti15 zonal rotor at 376,000

x g in dissociation buffer 2 (20 mM HEPES, pH 7.5, 0.5 M KCl, 5 mM $MgCl_2$, 3 mM EDTA, 2

mM DTT) at 4 °C. Carrier 70S ribosomes were isolated from S30 of *E. coli* cells by three

consecutive ultracentrifugations through a 1.1 M sucrose cushion in a buffer of 20 mM Tris, pH

7.5, 500 mM NH_4Cl, 10 mM Mg Acetate, 0.5 mM EDTA, 3 mM 2-mercaptoethanol. Elongation

factors eEF2 (Jørgensen, *et al.*, 2002) and eEF1A (Thiele, *et al.*, 1985) were isolated from Baker's

yeast as described. Yeast 6xHis-tagged release factors (full-length eRF1 and amino acids 166-685

of eRF3) were expressed in *E. coli* and purified using a TALON cobalt resin. Both release factors

were a generous gift from Alper Celik (University of Massachusetts Medical School).

<u>tRNA and mRNA</u>. $tRNA^{Lys}$, $tRNA^{Val}$, $tRNA^{Gln}$, and $tRNA^{Met}$ were isolated from *E. coli* bulk tRNA

(Roche). $tRNA^{Arg}$, $tRNA^{Trp}$ and $tRNA^{Leu}$ were isolated from Baker's yeast bulk tRNA (Roche),

using hybridization with immobilized complementary oligoDNA as described previously

(Barhoom, *et al.*, 2013; Liu, *et al.*, 2014). Yeast $tRNA^{Phe}$ (Sigma) and all isoacceptor tRNAs

mentioned above where charged with their cognate amino acids as described (Pan, *et al.*, 2007;

Pan, *et al.*, 2009). CrPV-IRES (Zhang, *et al.*, 2016) was modified by Genscript, Inc to encode the

initial mRNA sequence UUCAAAGUGAGA<u>UGG</u>CUAAUG (denoted Trp-IRES). A point

mutation was introduced into Trp-IRES to convert the UGG codon for Trp into a UGA stop codon

(UUCAAAGUGAGAUGACUAAUG, denoted Stop-IRES). These two sequences were inserted

into pUC57-Kan plasmid and amplified in TOP 10 competent cells. Plasmids were extracted using

QIAGEN Plasmid Kits and linearized. Trp-IRES and STOP-IRES were produced by in-vitro

transcription.

<u>POST4 and POST5 Complex Preparation</u>. 80S-IRES complex was first formed by incubating 0.8

µM 40S, 1.1 µM 60S and 0.8 µM IRES in Buffer 4 (40 mM Tris-HCl pH 7.5, 80 mM NH_4Cl, 5

mM $Mg(OAc)_2$, 100 mM KOAc, 3 mM 2-mercaptoethanol) at 37 °C for 2 min. Post-translocation

complex with FKVR-tRNA^{Arg} in the ribosomal P-site tRNA (POST4) was formed by incubating

0.4 µM 80S-IRES with 0.4 µM each of the first four aminoacylated tRNAs, 0.4 µM eEF1A, 1.0

µM eEF2, 1 mM GTP at 37 °C for 25 min in Buffer 4. POST4 was then purified by

ultracentrifugation in 1.1 M sucrose with Buffer 4 at 540,000 x g for 90 min at 4°C. POST4 pellet

was resuspended in Buffer 4. Post-translocation complex with FKVRQ-tRNA^{Gln} in the ribosomal

P-site tRNA (POST5) was prepared in identical fashion, except that the first five aminoacylated

tRNAs were added prior to the 25 min incubation at 37 °C. POST-4 amd POST-5 complexes could

be prepared and stored in small aliquots at -80 ^{0}C for at least three months with no discernible loss

of activity.

In vitro tRNA-Quant and PEP-Quant Readthrough Assays. POST5 complex (0.02 µM) was mixed

with Trp-tRNA^{Trp}, Leu-tRNA^{Leu}, and [^{35}S]-Met-tRNA^{Met} (0.08 µM each), elongation factors

eEF1A (0.08 µM), eEF-2 (1.0 µM) and release factors eRF1 (0.010 µM) and eRF3 (0.020 µM)

and incubated at 37 °C in Buffer 4 for 20 min, in the absence or presence of NonSups.

For the tRNA-Quant Assay, reaction mixture aliquots (40 µL) were quenched with 150 µL

of 0.5 M MES buffer (pH 6.0). Following addition of carrier 70S *E. coli* ribosomes (100 pmol, 3

µL of 33 µM 70S), all ribosomes were pelleted by ultracentrifugation through a 1.1 M sucrose

solution in Buffer 4 (350 µL) at 540,000 x g for 70 min at 4°C. The ribosome pellet was

resuspended in Buffer 4, and co-sedimenting FKVRQWL[^{35}S]M-tRNA^{Met} was determined.

For the Pep-Quant Assay, reaction mixture aliquots (80 µL) were quenched with 0.8 M

KOH (9 µL of 8M KOH) and the base-quenched samples were incubated at 37 °C for 1 h to

completely release octapeptide FKVRQWL[^{35}S]M from tRNA^{Met}. Acetic acid (9 µL) was then

added to lower the pH to 2.8. Samples were next lyophilized, suspended in water, and centrifuged

to remove particulates. The particulates contained no ^{35}S. The supernatant was analyzed by thin

layer electrophoresis (TLE) as previously described (Youngman*, et al.*, 2004), using the same

running buffer. The identity of FKVRQWLM was confirmed by the co-migration of the ^{35}S

radioactivity with authentic samples obtained from GenScript (Piscataway, NJ). The ^{35}S

radioactivity in the octapeptide band was used to determine the amount of octapeptide produced.

In both assays, the assay background was determined as ^{35}S either co-sedimenting (tRNA-

Quant) or comigrating (PEP-Quant) in the absence of added Trp-tRNA^{Trp}. These levels were 0.09

± 0.01 (sd, n = 80) octapeptide/POST5 for the tRNA-Quant assay and 0.04 ± 0.01 (sd, n = 8)

octapeptide/POST5 for the PEP-Quant assay. Some NonSups are poorly soluble in water and were

added to reaction mixtures from concentrated solutions made up in either DMSO (RTC13, GJ071,

GJ072, azithromycin) or methanol (escin). The level of organic solvent in the assay medium was

≤ 0.5%. For these NonSups the small amount of readthrough induced in the presence of added

Trp-tRNA^{Trp} by added organic solvent (Table S1) was additionally subtracted as background.

Readthrough levels presented in Figures 3 and S3 are all background subtracted. Although PEP-

Quant has a lower procedural background, it is time consuming to perform, and less precise than

tRNA-Quant. Both assays show a basal level of readthrough, in the absence of added NonSup, of

0.08 ± 0.02 (sd, n=40) octapeptide/POST5 above background, with some day-to-day variation.

<u>smFRET experiments</u>. Fluorescent ternary complexes (TCs) were prepared by incubating 1 μM

eEF-1A, 3 μM GTP, and 1 μM charged tRNAs labeled with either Cy3 or Cy5 (Chen*, et al.*, 2011)

at 37 °C for 15 min in Buffer 4. For experiments measuring only PRE6 complex formation, POST4

complex, containing FVKR-tRNA^{Arg} in the P-site and formed from ribosomes programmed with

either Trp-IRES or Stop-IRES biotinylated at the 5' end (Chen*, et al.*, 2011), was incubated with

15 nM Gln-TC(Cy5), 1 μM eEF-2 and 2 mM GTP in buffer 4 for 5 min at room temperature. The

resulting POST5 complex was immobilized on a streptavidin/biotin-PEG coated glass surface

(Chen*, et al.*, 2011). After two minutes of incubation, unbound reaction components were washed

out of the channel and 15 nM Trp-TC(Cy3) was added, with or without a NonSup, into the channel

to make a PRE6 complex. Unbound Trp-TC(Cy3) was washed out of the channel with Buffer 4

containing a deoxygenation enzyme system of 100 μg/mL glucose oxidase, 3 mg/mL glucose, and

48 μg/mL catalase to minimize photobleaching. Cy3 and Cy5 fluorescence intensities were

collected with 100 ms time resolution using alternating laser excitation (ALEX) between 532 nm

and 640 nm lasers on an objective-type total internal reflection fluorescence microscope described

previously (Chen*, et al.*, 2011). For experiments measuring both PRE6 complex and POST6

complex formation, Trp-IRES-PRE6 complex was formed as described above and 1 μM eEF-2

was injected while recording the FRET between FVKRQW-tRNA^{Trp}(Cy3) and tRNA^{Gln}(Cy5).

<u>Ataluren ^{19}F NMR Spectroscopy</u>. Various concentrations of ataluren solutions (0.03, 0.1 and 2.0

mM) were prepared in buffer 4 with 10% D_2O. The ^{19}F NMR spectrum of each solution was

recorded on a Bruker DMX 360 MHz NMR spectrometer with a 5 mm Quattro Nucleus Probe.

Data were analyzed with mNova software.

Results

The *in vitro* ribosomal readthrough assay. Structural studies (Fernández*, et al.*, 2014; Koh*, et al.*,

2014; Muhs*, et al.*, 2015; Murray*, et al.*, 2016; Abeyrathne*, et al.*, 2016) have shown that, prior to

polypeptide chain elongation, the cricket paralysis virus (CrPV) IRES structure occupies all three

tRNA binding sites (E, P, and A) on the 80S ribosome. We have recently demonstrated that the

first two cycles of peptide elongation proceed very slowly due to very low rates of pseudo-

translocation and translocation, but that, following translocation of tripeptidyl-tRNA, subsequent

elongation cycles proceed more rapidly (Zhang*, et al.*, 2016). Based on these results we

constructed an assay to directly monitor readthrough of the sixth codon, when the faster elongation

rate is well established. For this purpose, we prepared the two CrPv IRES coding sequences,

STOP-IRES and Trp-IRES (Figure 2). STOP-IRES contains the stop codon UGA at position 6 and

has a peptide coding sequence designed to give a high amount of readthrough, based on previous

studies showing that readthrough at the UGA stop codon proceeds in higher yields than at either

the UAA and UAG stop codons (Dabrowski*, et al.*, 2015) and that such readthrough is further

increased by both a downstream CUA codon (encoding Leu) at positions +4 - +6 (Stiebler*, et al.*,

2014; Loughran*, et al.*, 2014) and an upstream AA sequence at positions -1 and -2 (Dabrowski*, et

al.*, 2015). In TRP-IRES UGA is replaced by UGG which is cognate to tRNA^{Trp}, the most efficient

natural tRNA suppressor of the UGA codon (Blanchet*, et al.*, 2014; Roy*, et al.*, 2015). Trp-IRES

encodes the octapeptide FKVRQWLM, which permits facile quantification of octapeptide

synthesis by ^{35}S-Met incorporation.

For the results reported below, we first prepared two POST5 translocation complexes, each

containing FKVRQ-tRNA^{Gln} in the P-site, using ribosomes programmed with either STOP-IRES

or TRP-IRES. We then used the tRNA-Quant assay, which is rapid and precise, to determine the

amount of FKVRQWLM-tRNA^{Met} formed on incubating each POST5 complex with a mixture of

Trp-tRNA^{Trp}, Leu-tRNA^{Leu}, [^{35}S]-Met-tRNA^{Met}, elongation factors eEF1A and eEF2 and release

factors eRF1 and eRF3. We verified the validity of the tRNA-Quant assay by demonstrating that

it gives results that are very similar to those obtained with the PEP-Quant assay (Figure S1), in

which, following base treatment, the amount of FKVRQWLM octapeptide is determined following

a TLE separation procedure (Zhang, *et al.*, 2016; Youngman, *et al.*, 2004).

<u>Induction of readthrough by aminoglycosides (AGs)</u>. Results with the eight AGs examined are all

consistent with a single tight site of AG binding to the ribosome (Garreau de Loubresse, *et al.*,

2014) resulting in increased readthrough, with EC$_{50}$s falling in the range of $0.14 - 4$ µM and

fractional readthrough efficiencies of Stop-IRES varying from $0.1 - 0.3$ (Table 1), as compared

with an efficiency of 1.00 ± 0.02 (n = 24) for conversion of POST5 to POST8 complex with Trp-

IRES (Figure 3A, Table 1). These results are consistent with results on readthrough obtained in

intact cells showing a) G418, gentamicin B1 (Baradaran-Heravi, *et al.*, 2017), NB84, NB124

(Bidou, *et al.*, 2017) and gentamicin X2 (Friesen, *et al.*, 2018) to be much more effective than the

gentamicin mixture currently used as an approved antibiotic; b) gentamicin B1 to be much more

effective than gentamicin B, despite their differing by only a single methyl group (Figure 1)

(Baradaran-Heravi, *et al.*, 2017); c) gentamicin B1 to be more effective than streptothricin (Figure

S2C); d) NB84, NB124 (Bidou, *et al.*, 2017), and gentamicin X2 (Friesen, *et al.*, 2018), to have

similar potencies, measured by either EC$_{50}$ or readthrough efficiency.

Induction of readthrough by ataluren-like compounds. The NonSups ataluren, GJ072, and RTC 13

share similar structures, containing a central aromatic heterocycle having two or three substituents,

at least one of which is aromatic (Figure 1). They also show similar S-shaped readthrough activity

saturation curves (Figure 3B), with EC_{50} values between 0.17 – 0.35 mM and plateau readthrough

efficiencies ranging from 0.10 – 0.16 (Table 1). These S-shaped curves yield Hill n values of ~ 4,

which suggest multi-site binding of ataluren-like NonSups to the protein synthesis machinery.

Formation of NonSup aggregates in solution that induce readthrough could also give rise to S-

shaped curves, but we consider this to be unlikely based on the constancy of the chemical shift and

line shape of ataluren's ^{19}F NMR peak over a concentration range of 0.03 – 2.0 mM (see

Supplementary Information).

Induction of readthrough by other NonSups. Two other reported NonSups, negamycin (Taguchi*,

et al.*, 2017) and doxorubicin (Mutyam*, et al.*, 2016) also display readthrough activity in the tRNA-

Quant assay (Figure 3B). The results with each fit a simple saturation curve. Both NonSups have

similar readthrough efficiencies (0.10 – 0.13) but a 50-fold difference in EC_{50} values, with

doxorubicin having the much lower value (Table 1). Several other compounds that have

readthrough activity in cellular assays, tylosin (Zilberberg*, et al.*, 2010), azithromycin (Caspi*, et

al.*, 2016), GJ071 (Du*, et al.*, 2013) and escin (Mutyam*, et al.*, 2016) showed little or no activity

in the tRNA-Quant assay in the concentration range 30 – 600 μM (Figure S3). In addition, escin

at high concentration inhibits both basal readthrough elongation and normal elongation, the latter

measured with Trp-IRES programmed ribosomes, with the effect on basal readthrough being much

more severe (Figure S4).

<u>Single molecule assay of readthrough activity</u>. Two fluorescent labeled tRNAs, when bound

simultaneously to a ribosome, at either the A- and P-sites in a pretranslocation complex, or the P-

and E-sites in a postranslocation complex, are close enough to generate a FRET signal (Chen, *et*

al., 2011; Blanchard, *et al.*, 2004). We observed tRNA-tRNA FRET in the pretranslocation

complex (Trp-IRES-PRE6), formed by incubating the Cy5-labeled Trp-IRES-POST5 with Cy3-

labeled eEF1A.GTP.Trp-tRNA^{Trp}, and having tRNA^{Gln}(Cy5) in the P-site and FKVRQW-

tRNA^{Trp}(Cy3) in the A-site (Figure 4). Addition of eEF2.GTP converted Trp-IRES-PRE6 to a Trp-

IRES-POST6 complex, containing tRNA^{Gln}(Cy5) in the E-site and FKVRQW-tRNA^{Trp}(Cy3) in

the P-site, which is accompanied by an increase in Cy3:Cy5 FRET efficiency (Figure 4).

Repetition of this experiment with Stop-IRES-POST5 in the absence of eEF2 decreased the

number of pretranslocation complexes (STOP-IRES-PRE6) formed to 24% of that seen with Trp-

IRES (Figure 4). This value was increased in a dose-dependent manner by addition of either G418

or gentamicin B1 (Figure 4B), with relative potencies similar to those displayed in Table 1. G418

and gentamicin did not significantly affect formation of Trp-IRES-PRE6 from Trp-IRES-POST5

on addition of Cy3-labeled eEF1A.GTP.Trp-tRNA^{Trp}. The agreement between the ensemble and

single molecule assays demonstrates our ability to monitor NonSup-induced readthrough by

smFRET, which, in subsequent studies, will allow determination of the effects of NonSups on the

dynamics of the nascent peptide elongation cycle that commences with suppressor tRNA

recognition of a premature stop codon.

Discussion

Here we utilize a straightforward *in vitro* assay, tRNA-Quant, to measure direct nonsense

suppressor-induced readthrough (DIRECT-NSIRT) of a termination codon. In the tRNA-Quant

assay, the arrival of the UGA termination codon into the 40S subunit portion of the tRNA A-site

has two possible outcomes: termination of peptide synthesis via eRF1/eRF3-catayzed hydrolysis

of the P-site-bound FKVRQ-tRNA^{Gln} or readthrough via productive A-site binding of near-cognate

Trp-tRNA^{Trp} followed by productive binding of the cognate Leu-tRNA^{Leu} and Met- tRNA^{Met}

leading to FKVRQWLM-tRNA^{Met} formation. NonSups increase the readthrough percentage by

binding to one or more of the specific components of the protein synthesis apparatus present in the

assay. Our working hypothesis is that Direct-NSIRT is an important, perhaps dominant, part of

Total-NSIRT for NonSups showing parallel effects in tRNA-Quant and cellular assays, such as

those included in Table 1. In contrast, biological activities of NonSups showing strong

readthrough activity in cellular assays but little readthrough activity in tRNA-Quant (Figure S3),

are likely be dominated by indirect effects.

Our results suggest that aminoglycosides and ataluren-like compounds stimulate

readthrough by different mechanisms, AGs via binding to a single tight site on the ribosome and

ataluren-like compounds via weaker, multi-site binding. (Figures 3, S2; Table 1). The EC_{50} values

found in intact cells differ considerably from those measured by tRNA-Quant, being much higher

for AGs (Bidou*, et al.*, 2017; Baradaran-Heravi*, et al.*, 2017), and much lower for ataluren (Peltz*,*

et al., 2013; Roy*, et al.*, 2016), RTC13 (Du*, et al.*, 2013) and GJ072 (Du*, et al.*, 2013). We attribute

these differences to the fact that positively charged aminoglycosides are taken up poorly into cells,

while uptake is favored for the hydrophobic ataluren-like molecules. Thus, vis-à-vis the culture

medium, intracellular concentration would be expected to be lower for AGs and higher for

ataluren, RTC13 and GJ072. Although the NonSups doxorubicin and negamycin have only modest

activities (Figure 3, Table 1) each of these compounds have potential interest for future

development. Doxorubicin has a relatively low EC_{50}, is clinically approved for use in cancer

chemotherapy, and is the subject of ongoing efforts to identify doxorubicin congeners having

lower toxicity than doxorubicin itself (Kizek*, et al.*, 2012; Edwardson*, et al.*, 2015). Negamycin

exhibits low acute toxicity and there are ongoing efforts to increase its readthrough activity via

structure – function studies (Taguchi*, et al.*, 2017).

A critical barrier to further development of NonSups that are clinically useful is the paucity

of information regarding the mechanisms by which they stimulate readthrough and misreading.

Aminoglycosides have well-characterized binding sites in both prokaryotic (Lin*, et al.*, 2018) and

eukaryotic ribosomes (Garreau de Loubresse*, et al.*, 2014), proximal to the small subunit decoding

center, that have been linked to their promotion of misreading. Similarly, the functionally

important prokaryotic ribosome binding site of negamycin has also been identified within a

conserved small subunit rRNA region that is proximal to the decoding center (Lin*, et al.*, 2018;

Spahn*, et al.*, 2001), and it is not unlikely that this site is also present in eukaryotic ribosomes.

However, nothing is known about the readthrough-inducing sites of action within the protein

synthesis apparatus of the ataluren-like NonSups (Figure 3B) or of doxorubicin. Indeed, it has even

been suggested that ataluren may not target the ribosome (Pibiri*, et al.*, 2015). Although

aminoglycosides have been the subject of detailed mechanism studies of their effects on

prokaryotic misreading (Liu*, et al.*, 2014; Pape*, et al.*, 2000; Gromadski and Rodnina, 2004;

Cochella*, et al.*, 2006; Tsai*, et al.*, 2013; Zhang*, et al.*, 2018), questions remain over their precise

modes of action, and detailed mechanistic studies on aminoglycoside stimulation of eukaryotic

readthrough and misreading are completely lacking. Virtually nothing is known about the

mechanisms of action of negamycin, doxorubicin, and the ataluren-like NonSups in stimulating

eukaryotic readthrough. Single molecule FRET is a method of choice for obtaining detailed

information about processive biochemical reaction mechanisms (Roy*, et al.*, 2008), particularly in

the study of protein synthesis (Perez and Gonzalez, 2011; Aitken*, et al.*, 2010; Wang*, et al.*, 2011;

Chen*, et al.*, 2013), because it permits determination of distributions, variations and fluctuations

among different ribosome conformational states and of complex multistep reaction trajectories.

Here we demonstrate the feasibility of using smFRET observations for detailed examination of

NonSup-stimulation of readthrough (Figure 4) and misreading, which, combined with other

mechanistic studies, should aid in achieving the understanding needed to improve the clinical

usefulness of NonSups.

Acknowledgements

We thank Drs. Alexander Mankin (University of Illinois at Chicago), Timor Baasov (Technion,

Haifa) and Alper Celik (University of Massachusetts Medical School) for reagents used in these

studies and Carla Zimmerman for technical assistance. This work was supported by research grants

to BSC from PTC Therapeutics and the Orphan Disease Center, University of Pennsylvania

(MDBR-17-108-CF1282X), to YEG (NIH GM118139), to MR (Michael Smith Foundation for

Health Research) and AJ (NIH GM27757 and GM122468).

References

1   Brenner, S., Stretton, A. O. W. & Kaplan, S. 1965. Genetic Code: The 'Nonsense'
Triplets for Chain Termination and their Suppression. *Nature* **206**: 994,
doi:10.1038/206994a0.

2   Shalev, M. & Baasov, T. 2014. When Proteins Start to Make Sense: Fine-tuning
Aminoglycosides for PTC Suppression Therapy. *MedChemComm* **5**: 1092-1105,
doi:10.1039/C4MD00081A.

3   Keeling, K. M., Xue, X., Gunn, G. & Bedwell, D. M. 2014. Therapeutics Based on Stop
Codon Readthrough. *Annu. Rev. Genom. Hum. Genet.* **15**: 371-394, doi:10.1146/annurev-
genom-091212-153527.

4   Salvatori, F. *et al.* 2009. Production of β-globin and adult hemoglobin following G418
treatment of erythroid precursor cells from homozygous β(0)39 thalassemia patients. *Am.*
J. Hematol. **84**: 720-728, doi:10.1002/ajh.21539.

5   Goldmann, T., Overlack, N., Möller, F., Belakhov, V., van Wyk, M., Baasov, T.,
Wolfrum, U. & Nagel-Wolfrum, K. 2012. A comparative evaluation of NB30, NB54 and
PTC124 in translational read-through efficacy for treatment of an USH1C nonsense
mutation. *EMBO Molecular Medicine* **4**: 1186-1199, doi:10.1002/emmm.201201438.

6   Stenson, P. D., Mort, M., Ball, E. V., Evans, K., Hayden, M., Heywood, S., Hussain, M.,
Phillips, A. D. & Cooper, D. N. 2017. The Human Gene Mutation Database: towards a
comprehensive repository of inherited mutation data for medical research, genetic
diagnosis and next-generation sequencing studies. *Hum. Genet.* **136**: 665-677,
doi:10.1007/s00439-017-1779-6.

7   Loudon, J. A. 2013. Repurposing amlexanox as a 'run the red light cure- all' with read-
through – a 'no-nonsense' approach to eersonalised medicine. *J. Bioanal. Biomed.* **5**:
079-096, doi:doi: 10.4172/1948-593X.1000086.

8   Peltz, S. W., Morsy, M., Welch, E. M. & Jacobson, A. 2013. Ataluren as an Agent for
Therapeutic Nonsense Suppression. *Annu. Rev. Med.* **64**: 407-425, doi:10.1146/annurev-
med-120611-144851.

9   McDonald, C. M. *et al.* 2017. Ataluren in patients with nonsense mutation Duchenne
muscular dystrophy (ACT DMD): a multicentre, randomised, double-blind, placebo-
controlled, phase 3 trial. *Lancet* **390**: 1489-1498, doi:10.1016/S0140-6736(17)31611-2.

10   Zainal Abidin, N., Haq, I. J., Gardner, A. I. & Brodlie, M. 2017. Ataluren in cystic
fibrosis: development, clinical studies and where are we now? *Expert Opin.*
Pharmacother. **18**: 1363-1371, doi:10.1080/14656566.2017.1359255.

11   Brodlie, M., Haq, I. J., Roberts, K. & Elborn, J. S. 2015. Targeted therapies to improve
CFTR function in cystic fibrosis. *Genome Med.* **7**: 101, doi:10.1186/s13073-015-0223-6.

12 Floquet, C., Rousset, J.-P. & Bidou, L. 2011. Readthrough of Premature Termination
Codons in the Adenomatous Polyposis Coli Gene Restores Its Biological Activity in
Human Cancer Cells. *PLoS One* **6**: e24125, doi:10.1371/journal.pone.0024125.

13 Zilberberg, A., Lahav, L. & Rosin-Arbesfeld, R. 2010. Restoration of APC gene function
in colorectal cancer cells by aminoglycoside- and macrolide-induced read-through of
premature termination codons. *Gut* **59**: 496-507, doi:10.1136/gut.2008.169805.

14 Miyaki, M., Iijima, T., Yasuno, M., Kita, Y., Hishima, T., Kuroki, T. & Mori, T. 2002.
High incidence of protein-truncating mutations of the p53 gene in liver metastases of
colorectal carcinomas. *Oncogene* **21**: 6689, doi:10.1038/sj.onc.1205887.

15 Floquet, C., Deforges, J., Rousset, J.-P. & Bidou, L. 2011. Rescue of non-sense mutated
p53 tumor suppressor gene by aminoglycosides. *Nucleic Acids Res.* **39**: 3350-3362,
doi:10.1093/nar/gkq1277.

16 Roy, B. *et al.* 2016. Ataluren stimulates ribosomal selection of near-cognate tRNAs to
promote nonsense suppression. *Proc. Natl. Acad. Sci. USA* **113**: 12508-12513,
doi:10.1073/pnas.1605336113.

17 Baradaran-Heravi, A. *et al.* 2016. Novel small molecules potentiate premature
termination codon readthrough by aminoglycosides. *Nucleic Acids Res.* **44**: 6583-6598,
doi:10.1093/nar/gkw638.

18 Bidou, L., Bugaud, O., Belakhov, V., Baasov, T. & Namy, O. 2017. Characterization of
new-generation aminoglycoside promoting premature termination codon readthrough in
cancer cells. *RNA Biol.* **14**: 378-388, doi:10.1080/15476286.2017.1285480.

19 Oishi, N. *et al.* 2015. XBP1 mitigates aminoglycoside-induced endoplasmic reticulum
stress and neuronal cell death. *Cell Death Dis.* **6**: e1763, doi:10.1038/cddis.2015.108.

20 Duscha, S. *et al.* 2014. Identification and Evaluation of Improved 4′-O-(Alkyl) 4,5-
Disubstituted 2-Deoxystreptamines as Next-Generation Aminoglycoside Antibiotics.
mBio **5**, doi:10.1128/mBio.01827-14.

21 Floquet, C., Hatin, I., Rousset, J.-P. & Bidou, L. 2012. Statistical Analysis of
Readthrough Levels for Nonsense Mutations in Mammalian Cells Reveals a Major
Determinant of Response to Gentamicin. *PLoS Genet.* **8**: e1002608,
doi:10.1371/journal.pgen.1002608.

22 Sangkuhl, K., Schulz, A., Römpler, H., Yun, J., Wess, J. & Schöneberg, T. 2004.
Aminoglycoside-mediated rescue of a disease-causing nonsense mutation in the V2
vasopressin receptor gene in vitro and in vivo. *Hum. Mol. Genet.* **13**: 893-903,
doi:10.1093/hmg/ddh105.

23 Fuchshuber-Moraes, M., Sampaio Carvalho, R., Rimmbach, C., Rosskopf, D., Alex
Carvalho, M. & Suarez-Kurtz, G. 2011. Aminoglycoside-induced suppression of

CYP2C19*3 premature stop codon. *Pharmacogenet. Genomics* **21**: 694-700,
doi:10.1097/FPC.0b013e328349daba.

24   Cogan, J., Weinstein, J., Wang, X., Hou, Y., Martin, S., South, A. P., Woodley, D. T. &
Chen, M. 2014. Aminoglycosides Restore Full-length Type VII Collagen by Overcoming
Premature Termination Codons: Therapeutic Implications for Dystrophic Epidermolysis
Bullosa. *Mol. Ther.* **22**: 1741-1752, doi:doi.org/10.1038/mt.2014.140.

25   Baradaran-Heravi, A. *et al.* 2017. Gentamicin B1 is a minor gentamicin component with
major nonsense mutation suppression activity. *Proc. Natl. Acad. Sci. USA* **114**: 3479-
3484, doi:10.1073/pnas.1620982114.

26   Welch, E. M. *et al.* 2007. PTC124 targets genetic disorders caused by nonsense
mutations. *Nature* **447**: 87, doi:10.1038/nature05756.

27   Du, L. *et al.* 2009. Nonaminoglycoside compounds induce readthrough of nonsense
mutations. *J. Exp. Med.* **206**: 2285-2297, doi:10.1084/jem.20081940.

28   Du, L. *et al.* 2013. A New Series of Small Molecular Weight Compounds Induce Read
Through of All Three Types of Nonsense Mutations in the ATM Gene. *Mol. Ther.* **21**:
1653-1660, doi:10.1038/mt.2013.150.

29   Gómez-Grau, M. *et al.* 2015. Evaluation of Aminoglycoside and Non-Aminoglycoside
Compounds for Stop-Codon Readthrough Therapy in Four Lysosomal Storage Diseases.
PLoS One **10**: e0135873, doi:10.1371/journal.pone.0135873.

30   Arakawa, M. *et al.* 2003. Negamycin Restores Dystrophin Expression in Skeletal and
Cardiac Muscles of *mdx* Mice. *J. Biochem.* **134**: 751-758, doi:10.1093/jb/mvg203.

31   Hamada, K., Taguchi, A., Kotake, M., Aita, S., Murakami, S., Takayama, K., Yakushiji,
F. & Hayashi, Y. 2015. Structure–Activity Relationship Studies of 3-epi-
Deoxynegamycin Derivatives as Potent Readthrough Drug Candidates. *ACS Med. Chem.*
Lett. **6**: 689-694, doi:10.1021/acsmedchemlett.5b00121.

32   Caspi, M. *et al.* 2016. A flow cytometry-based reporter assay identifies macrolide
antibiotics as nonsense mutation read-through agents. *J. Mol. Med.* **94**: 469-482,
doi:10.1007/s00109-015-1364-1.

33   Mutyam, V. *et al.* 2016. Discovery of Clinically Approved Agents That Promote
Suppression of Cystic Fibrosis Transmembrane Conductance Regulator Nonsense
Mutations. *Am. J. Respir. Crit. Care Med.* **194**: 1092-1103, doi:10.1164/rccm.201601-
0154OC.

34   He, F. & Jacobson, A. 2015. Nonsense-Mediated mRNA Decay: Degradation of
Defective Transcripts Is Only Part of the Story. *Annu. Rev. Genet.* **49**: 339-366,
doi:10.1146/annurev-genet-112414-054639.

35 Feng, T. *et al.* 2014. Optimal Translational Termination Requires C4 Lysyl
456   Hydroxylation of eRF1. *Mol. Cell* **53**: 645-654, doi:10.1016/j.molcel.2013.12.028.

457 36 Keeling, K. M. 2016. Nonsense Suppression as an Approach to Treat Lysosomal Storage
458   Diseases. *Diseases* **4**: 32, doi:10.3390/diseases4040032.

459 37 Zhang, H., Ng, M. Y., Chen, Y. & Cooperman, B. S. 2016. Kinetics of initiating
460   polypeptide elongation in an IRES-dependent system. *eLife* **5**: e13429,
461   doi:10.7554/eLife.13429.

462 38 Iwasaki, K. & Kaziro, Y. 1979. [60] Polypeptide chain elongation factors from pig liver.
463   Moldave, K. & L. Grossman, editors. *Methods Enzymol.* New York: Academic Press, **60**:
464   657-676.

465 39 Ben-Shem, A., Garreau de Loubresse, N., Melnikov, S., Jenner, L., Yusupova, G. &
466   Yusupov, M. 2011. The Structure of the Eukaryotic Ribosome at 3.0 Å Resolution.
467   *Science* **334**: 1524-1529, doi:10.1126/science.1212642.

468 40 Jørgensen, R., Carr-Schmid, A., Ortiz, P. A., Kinzy, T. G. & Andersen, G. R. 2002.
469   Purification and crystallization of the yeast elongation factor eEF2. *Acta Crystallogr.*
470   *Sect. D. Biol. Crystallogr.* **58**: 712-715, doi:doi:10.1107/S0907444902003001.

471 41 Thiele, D., Cottrelle, P., Iborra, F., Buhler, J. M., Sentenac, A. & Fromageot, P. 1985.
472   Elongation factor 1 alpha from Saccharomyces cerevisiae. Rapid large-scale purification
473   and molecular characterization. *J. Biol. Chem.* **260**: 3084-3089.

474 42 Barhoom, S., Farrell, I., Shai, B., Dahary, D., Cooperman, B. S., Smilansky, Z., Elroy-
475   Stein, O. & Ehrlich, M. 2013. Dicodon monitoring of protein synthesis (DiCoMPS)
476   reveals levels of synthesis of a viral protein in single cells. *Nucleic Acids Res.* **41**: e177-
477   e177, doi:10.1093/nar/gkt686.

478 43 Liu, J. *et al.* 2014. Monitoring Collagen Synthesis in Fibroblasts Using Fluorescently
479   Labeled tRNA Pairs. *J. Cell. Physiol.* **229**: 1121-1129, doi:doi:10.1002/jcp.24630.

480 44 Pan, D., Kirillov, S. V. & Cooperman, B. S. 2007. Kinetically Competent Intermediates
481   in the Translocation Step of Protein Synthesis. *Mol. Cell* **25**: 519-529,
482   doi:10.1016/j.molcel.2007.01.014.

483 45 Pan, D., Qin, H. & Cooperman, B. S. 2009. Synthesis and functional activity of tRNAs
484   labeled with fluorescent hydrazides in the D-loop. *RNA* **15**: 346-354,
485   doi:10.1261/rna.1257509.

486 46 Youngman, E. M., Brunelle, J. L., Kochaniak, A. B. & Green, R. 2004. The Active Site
487   of the Ribosome Is Composed of Two Layers of Conserved Nucleotides with Distinct
488   Roles in Peptide Bond Formation and Peptide Release. *Cell* **117**: 589-599,
489   doi:10.1016/S0092-8674(04)00411-8.

47   Chen, C. *et al.* 2011. Single-Molecule Fluorescence Measurements of Ribosomal
Translocation Dynamics. *Mol. Cell* **42**: 367-377, doi:10.1016/j.molcel.2011.03.024.

48   Fernández, Israel S., Bai, X.-C., Murshudov, G., Scheres, Sjors H. W. & Ramakrishnan,
V. 2014. Initiation of Translation by Cricket Paralysis Virus IRES Requires Its
Translocation in the Ribosome. *Cell* **157**: 823-831, doi:10.1016/j.cell.2014.04.015.

49   Koh, C. S., Brilot, A. F., Grigorieff, N. & Korostelev, A. A. 2014. Taura syndrome virus
IRES initiates translation by binding its tRNA-mRNA–like structural element in the
ribosomal decoding center. *Proc. Natl. Acad. Sci. USA* **111**: 9139-9144,
doi:10.1073/pnas.1406335111.

50   Muhs, M., Hilal, T., Mielke, T., Skabkin, Maxim A., Sanbonmatsu, Karissa Y., Pestova,
Tatyana V. & Spahn, Christian M. T. 2015. Cryo-EM of Ribosomal 80S Complexes with
Termination Factors Reveals the Translocated Cricket Paralysis Virus IRES. *Mol. Cell*
57: 422-432, doi:10.1016/j.molcel.2014.12.016.

51   Murray, J., Savva, C. G., Shin, B.-S., Dever, T. E., Ramakrishnan, V. & Fernández, I. S.
2016. Structural characterization of ribosome recruitment and translocation by type IV
IRES. *eLife* **5**: e13567, doi:10.7554/eLife.13567.

52   Abeyrathne, P. D., Koh, C. S., Grant, T., Grigorieff, N. & Korostelev, A. A. 2016.
Ensemble cryo-EM uncovers inchworm-like translocation of a viral IRES through the
ribosome. *eLife* **5**: e14874, doi:10.7554/eLife.14874.

53   Dabrowski, M., Bukowy-Bieryllo, Z. & Zietkiewicz, E. 2015. Translational readthrough
potential of natural termination codons in eucaryotes – The impact of RNA sequence.
RNA Biol. **12**: 950-958, doi:10.1080/15476286.2015.1068497.

54   Stiebler, A. C., Freitag, J., Schink, K. O., Stehlik, T., Tillmann, B. A. M., Ast, J. &
Bölker, M. 2014. Ribosomal Readthrough at a Short UGA Stop Codon Context Triggers
Dual Localization of Metabolic Enzymes in Fungi and Animals. *PLoS Genet.* **10**:
e1004685, doi:10.1371/journal.pgen.1004685.

55   Loughran, G., Chou, M.-Y., Ivanov, I. P., Jungreis, I., Kellis, M., Kiran, A. M., Baranov,
P. V. & Atkins, J. F. 2014. Evidence of efficient stop codon readthrough in four
mammalian genes. *Nucleic Acids Res.* **42**: 8928-8938, doi:10.1093/nar/gku608.

56   Blanchet, S., Cornu, D., Argentini, M. & Namy, O. 2014. New insights into the
incorporation of natural suppressor tRNAs at stop codons in Saccharomyces cerevisiae.
Nucleic Acids Res. **42**: 10061-10072, doi:10.1093/nar/gku663.

57   Roy, B., Leszyk, J. D., Mangus, D. A. & Jacobson, A. 2015. Nonsense suppression by
near-cognate tRNAs employs alternative base pairing at codon positions 1 and 3. *Proc.*
Natl. Acad. Sci. USA **112**: 3038-3043, doi:10.1073/pnas.1424127112.

58   Garreau de Loubresse, N., Prokhorova, I., Holtkamp, W., Rodnina, M. V., Yusupova, G.
& Yusupov, M. 2014. Structural basis for the inhibition of the eukaryotic ribosome.
Nature **513**: 517, doi:10.1038/nature13737.

59   Friesen, W. *et al.* 2018. The minor gentamicin complex component, X2, is a potent
premature stop codon readthrough molecule with therapeutic potential. *Submitted for*
publication.

60   Taguchi, A., Hamada, K. & Hayashi, Y. 2017. Chemotherapeutics overcoming nonsense
mutation-associated genetic diseases: medicinal chemistry of negamycin. *J. Antibiot.* **71**:
205, doi:10.1038/ja.2017.112.

61   Blanchard, S. C., Kim, H. D., Gonzalez, R. L., Puglisi, J. D. & Chu, S. 2004. tRNA
dynamics on the ribosome during translation. *Proc. Natl. Acad. Sci. U. S. A.* **101**: 12893-
12898, doi:10.1073/pnas.0403884101.

62   Kizek, R., Adam, V., Hrabeta, J., Eckschlager, T., Smutny, S., Burda, J. V., Frei, E. &
Stiborova, M. 2012. Anthracyclines and ellipticines as DNA-damaging anticancer drugs:
Recent advances. *Pharmacol. Ther.* **133**: 26-39, doi:10.1016/j.pharmthera.2011.07.006.

63   Edwardson, D. W., Narendrula, R., Chewchuk, S., Mispel-Beyer, K., Mapletoft, J. P. J. &
Parissenti, A. M. 2015. Role of Drug Metabolism in the Cytotoxicity and Clinical
Efficacy of Anthracyclines. *Curr. Drug Metab.* **16**: 412-426,
doi:10.2174/1389200216888150915112039.

64   Lin, J., Zhou, D., Steitz, T. A., Polikanov, Y. S. & Gagnon, M. G. 2018. Ribosome-
Targeting Antibiotics: Modes of Action, Mechanisms of Resistance, and Implications for
Drug Design. *Annu. Rev. Biochem* **87**: null, doi:10.1146/annurev-biochem-062917-
011942.

65   Spahn, C. M. T., Beckmann, R., Eswar, N., Penczek, P. A., Sali, A., Blobel, G. & Frank,
J. 2001. Structure of the 80S Ribosome from Saccharomyces cerevisiae—tRNA-
Ribosome and Subunit-Subunit Interactions. *Cell* **107**: 373-386, doi:10.1016/S0092-
8674(01)00539-6.

66   Pibiri, I., Lentini, L., Melfi, R., Gallucci, G., Pace, A., Spinello, A., Barone, G. & Di
Leonardo, A. 2015. Enhancement of premature stop codon readthrough in the CFTR gene
by Ataluren (PTC124) derivatives. *Eur. J. Med. Chem.* **101**: 236-244,
doi:10.1016/j.ejmech.2015.06.038.

67   Pape, T., Wintermeyer, W. & Rodnina, M. V. 2000. Conformational switch in the
decoding region of 16S rRNA during aminoacyl-tRNA selection on the ribosome. *Nat.*
Struct. Biol. **7**: 104, doi:10.1038/72364.

68   Gromadski, K. B. & Rodnina, M. V. 2004. Streptomycin interferes with conformational
coupling between codon recognition and GTPase activation on the ribosome. *Nat. Struct.*
Mol. Biol. **11**: 316, doi:10.1038/nsmb742.

69   Cochella, L., Brunelle, J. L. & Green, R. 2006. Mutational analysis reveals two
independent molecular requirements during transfer RNA selection on the ribosome. *Nat.*
Struct. Mol. Biol. **14**: 30, doi:10.1038/nsmb1183.

70   Tsai, A., Uemura, S., Johansson, M., Puglisi, Elisabetta V., Marshall, R. A., Aitken,
Colin E., Korlach, J., Ehrenberg, M. & Puglisi, Joseph D. 2013. The Impact of
Aminoglycosides on the Dynamics of Translation Elongation. *Cell Reports* **3**: 497-508,
doi:10.1016/j.celrep.2013.01.027.

71   Zhang, J., Pavlov, M. Y. & Ehrenberg, M. 2018. Accuracy of genetic code translation
and its orthogonal corruption by aminoglycosides and Mg^{2+} ions. *Nucleic Acids Res.* **46**:
1362-1374, doi:10.1093/nar/gkx1256.

72   Roy, R., Hohng, S. & Ha, T. 2008. A practical guide to single-molecule FRET. *Nat.*
Methods **5**: 507, doi:10.1038/nmeth.1208.

73   Perez, C. E. & Gonzalez, R. L. 2011. In vitro and in vivo single-molecule fluorescence
imaging of ribosome-catalyzed protein synthesis. *Curr. Opin. Chem. Biol.* **15**: 853-863,
doi:10.1016/j.cbpa.2011.11.002.

74   Aitken, C. E., Petrov, A. & Puglisi, J. D. 2010. Single Ribosome Dynamics and the
Mechanism of Translation. *Ann. Rev. Biophys.* **39**: 491-513,
doi:10.1146/annurev.biophys.093008.131427.

75   Wang, L., Wasserman, M. R., Feldman, M. B., Altman, R. B. & Blanchard, S. C. 2011.
Mechanistic insights into antibiotic action on the ribosome through single-molecule
fluorescence imaging. *Ann. N.Y. Acad. Sci.* **1241**: E1-E16, doi:doi:10.1111/j.1749-
6632.2012.06839.x.

76   Chen, C., Zhang, H., Broitman, S. L., Reiche, M., Farrell, I., Cooperman, B. S. &
Goldman, Y. E. 2013. Dynamics of translation by single ribosomes through mRNA
secondary structures. *Nat. Struct. Mol. Biol.* **20**: 582, doi:10.1038/nsmb.2544.
587

593 **TABLES**

Table 1. Non-Sup Induced Readthrough			
NonSup	EC_{50} (µM)	Readthrough fraction[a]	Hill n
Aminoglycosides			
Gentamicin B1	0.14 ± 0.02	0.27 ± 0.01	-
Gentamicin X2	0.42 ± 0.08	0.31 ± 0.02	-
NB124	0.52 ± 0.05	0.21 ± 0.01	-
Gentamicin B	0.54 ± 0.15	0.081 ± 0.005	-
NB84	0.68 ± 0.06	0.19 ± 0.01	-
G418	0.99 ± 0.09	0.32 ± 0.01	-
Streptothricin	1.5 ± 0.3	0.26 ± 0.01	-
Commercial Gentamicin (mixture)	4.2 ± 0.6	0.29 ± 0.02	-
Ataluren-Like			
GJ072	98 ± 4	0.16 ± 0.01	4.2 ± 0.7
RTC13	270 ± 15	0.10 ± 0.01	3.1 ± 0.5
Ataluren	350 ± 20	0.15 ± 0.01	4.3 ± 0.8
Other			
Doxorubicin	9.8 ± 1.8	0.11 ± 0.01	-
Negamycin	490 ± 70	0.13 ± 0.01	-

[a]Plateau octapeptide formed/POST5

FIGURES

Figure 1. The structures of the nonsense suppressors (NonSups) studied in this work.

621 **Figure 2**. Coding sequences of Trp-IRES and Stop-IRES

Figure 3. Readthrough as a function of nonsense suppressor concentration. A. Aminoglycosides.

B. Ataluren-like NonSups and Others. The highest doxorubicin employed was 30 μM because

higher concentrations led to significant ribosome and Met-tRNA^{Met} particle formation (Figure S5).

None of the NonSups in Figure 3 showed appreciable inhibition of octapeptide formation from

pentapeptide by ribosomes programmed with Trp-IRES at concentrations equal to twice their EC_{50}

values.

Figure 4. smFRET Experiments. A. eEF2-induced translocation of the 80S-Trp-IRES-PRE6

complex to form 80S-Trp-IRES-POST6 complex followed by release of tRNA^{Gln}. The Trp-IRES-

PRE6 complex contains tRNA^{Gln}(Cy5) in the P-site and FKVRQW-tRNA^{Trp}(Cy3) in the A-site.

The cartoon at top shows the state progression during translocation. i. Single molecule traces.

Green and red traces show tRNA^{Trp}(Cy3) emission and tRNA^{Gln}(Cy5) sensitized emission,

respectively, following eEF2 injection, excited at 532 nm. ii. ALEX intensity signal from direct

excitation of tRNA^{Gln}(Cy5) at 640 nm. iii. FRET efficiency between tRNA^{Trp}(Cy3) and

tRNA^{Gln}(Cy5) showing a transient increase following eEF2 on conversion of PRE6 complex to

POST6 complex. B. Dose-dependent effect of G418 and GmB1 (gentamicin B1) on PRE6

complex formation from 80S-Stop-IRES-POST5 complex as compared with PRE6 complex

formation from 80S-Trp-IRES-POST5 complex in the absence of either G418 or GmB1.

SUPPLEMENTARY INFORMATION

The tRNA-Quant and PEP-Quant assays give similar results (Figure S1).

<u>Identification of streptothricin as a nonsense suppressor in yeast and human cells.</u> An in-house

collection of 664 antimicrobial compounds was screened for suppression of two nonsense alleles

in *Saccharomyces cerevisiae* using a modification of a published procedure (Baradaran-Heravi*, et*

al., 2016). Exponentially growing B0133-3B yeast cells harboring *met8-1* (TAG) and *trp5-48*

(TAA) nonsense alleles were seeded in 96-well plates at $A_{600} = 0.01$ in Synthetic Complete medium

containing 5 μM Met and 5 μM Trp. This strain is unable to grow in the presence of these low

concentrations of Met and Trp, unlike prototrophic strains. Antimicrobial compounds were added

individually to the wells using a Biorobotics TAS1 robot equipped with a 0.7 mm diameter 96-pin

tool, at a final concentration of ~15 μM. The plates were incubated at 30°C for 42 h and yeast

growth was determined by measuring A_{600}. Paromomycin was added at 10 μM to four wells as a

positive control. In this assay, yeast growth requires efficient suppression of both *met8-1* and *trp5-*

48. A single compound, streptothricin, enabled robust yeast growth (Figure S2A), with an EC_{50} of

5 μM (Figure S2B). Streptothricin was also assayed for nonsense suppression in human cells using

a previously described assay (Baradaran-Heravi *et al.*, 2016). Briefly, human breast carcinoma

HDQ-P1 cells homozygous for the *TP53* R213X (TGA) nonsense mutation were exposed to

different concentrations of streptothricin for 72 h and the production of full-length p53, the

readthrough product, and truncated p53 was determined by automated capillary electrophoresis

western analysis. Streptothricin showed weak readthrough activity, detectable at concentrations of

200 μM and above. By contrast, gentamicin B1 showed much higher levels of readthrough at lower

concentrations (Figure S2C).

<u>Assay heterogeneity</u>. The reaction mixtures used for the tRNA-Quant and Pep-Quant assays is

heterogeneous, with ribosomes derived from shrimp cysts or Hela cells, yeast elongation factors,

and yeast and *E. coli* charged tRNAs. However, such heterogeneity is not problematic. IRESs can

initiate translation on ribosomes from many eukaryotic organisms (Koh*, et al.*, 2014), including

shrimp (Cevallos and Sarnow, 2005), indicating that the molecular mechanism is not species-

specific. CrPV IRES can initiate translation on ribosomes from yeast (Thompson*, et al.*, 2001) to

human (Spahn*, et al.*, 2004). Furthermore, the structures of eukaryotic elongation factors are very

strongly conserved (Soares*, et al.*, 2009; Jørgensen*, et al.*, 2002), and charged tRNAs from one

species form fully functional complexes with both eEF1A and ribosomes from different ones

(Jackson*, et al.*, 2001; Ferguson*, et al.*, 2015).

<u>Concentration dependence of Ataluren ^{19}F NMR chemical shift</u>. A decrease in chemical shift of

an ^{19}F NMR peak provides an indication of molecular aggregation in solution (Iijima*, et al.*, 1999;

Ohta*, et al.*, 2003; Suzuki*, et al.*, 2013). We sought to determine whether aggregation was

responsible for the sigmoidal readthrough saturation curve for ataluren (Figure 3B) by examining

the chemical shift of its ^{19}F NMR peak at three concentrations, 0.03, 0.1 and 2.0 mM, that bracket

the range employed in the readthrough assay. We found that both the chemical shift (1.0 ppm

downfield from a KF standard) and the line shape of the ^{19}F NMR peak were identical at all three

concentrations, evidence that aggregation is unlikely to be the cause of the readthrough saturation

curve induced by ataluren.

<u>NonSups having low activity in the tRNA-Quant assay (Figure S3)</u>

<u>Escin inhibition of octapeptide synthesis from POST-5 complexes.</u> Escin concentrations ≥ 300 μM

inhibit octapeptide synthesis from POST5 complexes for ribosomes programmed with either Stop-

IRES (basal readthrough) or ribosomes programmed with Trp-IRES (normal octapeptide

synthesis), with inhibition being much more pronounced on basal readthrough (Figure S4). This

difference is not currently understood. One possibility under consideration is that inhibition arises

from a destabilization of peptidyl-tRNA binding to the A-site of a PRE6 complex, and that

such binding is weaker for ribosomes programmed with Stop-IRES vs. Trp-IRES.

<u>Doxorubicin-induced particle formation by both tRNA and ribosomes.</u> Doxorubicin

concentrations above 100 μM induced Met-tRNA^{Met} particle formation in accord with prior results

(Agudelo*, et al.*, 2016). High doxorubicin also induced particle formation by ribosome-IRES

complex (Figure S5). To determine the extent of particle formation, various concentrations of

doxorubicin were added to aliquots (250 μL) containing 0.1 μM 80S:IRES complex, 0.1 μM

[^{35}S]Met-tRNA^{Met}, 0.1 μM eEF1A and 1 mM GTP. The mixture was incubated at 37 °C for 20

min. Particles were removed by centrifugation at 17,000 X g for 25 min at 4 °C. The supernatant

was then layered on top of 350 μL Buffer 4 with 1.1 M sucrose and was ultracentrifuged at 540,000

x g for 70 min at 4°C to separate 80S-IRES from [^{35}S]Met-tRNA^{Met}. Virtually all of the A$_{260}$ units

and [^{35}S] radioactivity (~98% in each case) of the low speed supernatant were found in the high

speed pellet and supernatant, respectively. Accordingly, measurements of A$_{260}$ units and [^{35}S]

radioactivity in the low speed supernatant were used to determine the amounts of 80S-IRES and

[^{35}S]Met-tRNA^{Met} remaining in solution after doxorubicin-induced particle formation.

713
714   SUPPLEMENTARY REFERENCES
715

1   Baradaran-Heravi, A. *et al.* 2016. Novel small molecules potentiate premature
termination codon readthrough by aminoglycosides. *Nucleic Acids Res.* **44**: 6583-6598,
doi:10.1093/nar/gkw638.

2   Koh, C. S., Brilot, A. F., Grigorieff, N. & Korostelev, A. A. 2014. Taura syndrome virus
IRES initiates translation by binding its tRNA-mRNA–like structural element in the
ribosomal decoding center. *Proc. Natl. Acad. Sci. USA* **111**: 9139-9144,
doi:10.1073/pnas.1406335111.

3   Cevallos, R. C. & Sarnow, P. 2005. Factor-Independent Assembly of Elongation-
Competent Ribosomes by an Internal Ribosome Entry Site Located in an RNA Virus
That Infects Penaeid Shrimp. *J. Virol.* **79**: 677-683, doi:10.1128/jvi.79.2.677-683.2005.

4   Thompson, S. R., Gulyas, K. D. & Sarnow, P. 2001. Internal initiation in *Saccharomyces*
cerevisiae mediated by an initiator tRNA/eIF2-independent internal ribosome entry site
element. *Proc. Natl. Acad. Sci. USA* **98**: 12972-12977, doi:10.1073/pnas.241286698.

5   Spahn, C. M. T., Jan, E., Mulder, A., Grassucci, R. A., Sarnow, P. & Frank, J. 2004.
Cryo-EM Visualization of a Viral Internal Ribosome Entry Site Bound to Human
Ribosomes: The IRES Functions as an RNA-Based Translation Factor. *Cell* **118**: 465-
475, doi:10.1016/j.cell.2004.08.001.

6   Soares, D. C., Barlow, P. N., Newbery, H. J., Porteous, D. J. & Abbott, C. M. 2009.
Structural Models of Human eEF1A1 and eEF1A2 Reveal Two Distinct Surface Clusters
of Sequence Variation and Potential Differences in Phosphorylation. *PLoS One* **4**: e6315,
doi:10.1371/journal.pone.0006315.

7   Jørgensen, R., Carr-Schmid, A., Ortiz, P. A., Kinzy, T. G. & Andersen, G. R. 2002.
Purification and crystallization of the yeast elongation factor eEF2. *Acta Crystallogr.*
Sect. D. Biol. Crystallogr. **58**: 712-715, doi:doi:10.1107/S0907444902003001.

8   Jackson, R. J., Napthine, S. & Brierley, I. 2001. Development of a tRNA-dependent in
vitro translation system. *RNA* **7**: 765-773.

9   Ferguson, A. *et al.* 2015. Functional Dynamics within the Human Ribosome Regulate the
Rate of Active Protein Synthesis. *Mol. Cell* **60**: 475-486,
doi:10.1016/j.molcel.2015.09.013.

10   Iijima, H., Koyama, S., Fujio, K. & Uzu, Y. 1999. NMR Study of the Transformation of
Perfluorinated Surfactant Solutions. *Bull. Chem. Soc. Jpn.* **72**: 171-177,
doi:10.1246/bcsj.72.171.

11   Ohta, A., Murakami, R., Urata, A., Asakawa, T., Miyagishi, S. & Aratono, M. 2003.
Aggregation Behavior of Fluorooctanols in Hydrocarbon Solvents. *J. Phys. Chem. B* **107**:
11502-11509, doi:10.1021/jp035134o.

12   Suzuki, Y., Brender, J. R., Soper, M. T., Krishnamoorthy, J., Zhou, Y., Ruotolo, B. T.,
Kotov, N. A., Ramamoorthy, A. & Marsh, E. N. G. 2013. Resolution of Oligomeric
Species during the Aggregation of Aβ(1-40) Using (19)F NMR. *Biochemistry* **52**: 1903-
1912, doi:10.1021/bi400027y.

13   Agudelo, D., Bourassa, P., Bérubé, G. & Tajmir-Riahi, H. A. 2016. Review on the
binding of anticancer drug doxorubicin with DNA and tRNA: Structural models and
antitumor activity. *J. Photochem. Photobiol. B: Biol.* **158**: 274-279,
doi:10.1016/j.jphotobiol.2016.02.032.

794 **SUPPLEMENTARY TABLES**

Table S1. DMSO and MeOH each induce added basal readthrough	
Volume % DMSO	Basal readthrough fraction[a] above $-$Trp-RNA^{Trp} background
0	0.078 ± 0.013
0.1	0.099 ± 0.004
0.33	0.12 ± 0.01
0.67	0.14 ± 0.01
1.0	0.15 ± 0.01
Volume % MeOH	
0	0.099 ± 0.012
0.17	0.10 ± 0.01
0.5	0.12 ± 0.01
1.0	0.14 ± 0.01
[a]normalized to octapeptide synthesis by Trp-IRES in the absence of either DMSO or MeOH	

SUPPLEMENTARY FIGURES

Supplementary Figure 1. tRNA-Quant vs. Pep-Quant

Supplemental Figure 2. Nonsense suppression by streptothricin in yeast and human cells. A.

Scatter plot of the nonsense suppression activity of 664 antibiotics in yeast. B. Concentration

dependence of nonsense suppression by streptothricin in yeast. C. Concentration dependence of

p53 PTC readthrough in HDQ-P1 cells. Formation of full-length p53 (FL-p53) and truncated p53

(TR-p53) was determined by automated capillary electrophoresis western analysis and the results

displayed as pseudoblots. FL-p53 and TR-p53 chemiluminescence signal was normalized to that

of the protein loading control vinculin and expressed relative to the amount of TR-p53 detected

in untreated cells.

Supplementary Figure 3. Low activity NonSups. With the exception of escin (See Supplementary

Figure 4), none of the NonSups in this Figure at 600 µM showed appreciable inhibition of

octapeptide formation from pentapeptide by ribosomes programmed with Trp-IRES

846 **Supplementary Figure 4**. High escin concentrations inhibit both basal readthrough (ribosomes

847 programmed with Stop-IRES) and normal octapeptide synthesis (ribosomes programmed with

848 Trp-IRES).

Supplementary Figure 5. High doxorubicin concentrations induce particle formation by both

Met-tRNA^{Met} and 80S-IRES complexes.