General Functional Connectivity: shared features of resting-state and task fMRI drive reliable and heritable individual differences in functional brain networks

Maxwell L. Elliott[1*]
Annchen R. Knodt[1]
Megan Cooke[1]
M. Justin Kim[2]
Tracy R. Melzer[3,4]
Ross Keenan[3,5]
David Ireland[6]
Sandhya Ramrakha[6]
Richie Poulton[6]
Avshalom Caspi[1,7-9]
Terrie E. Moffitt[1,7-9]
Ahmad R. Hariri[1]

1. Department of Psychology & Neuroscience, Duke University Box 104410, Durham, NC 27708, USA
2. Department of Psychology, University of Hawaii at Manoa, Honolulu, HI 96822
3. New Zealand Brain Research Institute, Christchurch, New Zealand
4. Department of Medicine, University of Otago, Christchurch, New Zealand
5. Christchurch Radiology Group, Christchurch, New Zealand
6. Dunedin Multidisciplinary Health and Development Research Unit, Department of Psychology, University of Otago, 163 Union St E, Dunedin, 9016, NZ
7. Social, Genetic, & Developmental Psychiatry Research Centre, Institute of Psychiatry, Psychology, & Neuroscience, King's College London, De Crespigny Park, Denmark Hill, London SE5 8AF, UK
8. Department of Psychiatry & Behavioral Sciences, Duke University School of Medicine, Durham, NC 27708, USA
9. Center for Genomic and Computational Biology, Duke University Box 90338, Durham, NC 27708, USA

*Correspondence: maxwell.elliott@duke.edu

Abstract: Intrinsic connectivity, measured using resting-state fMRI, has emerged as a fundamental tool in the study of the human brain. However, due to practical limitations, many studies do not collect enough resting-state data to generate reliable measures of intrinsic connectivity necessary for studying individual differences. Here we present general functional connectivity (GFC) as a method for leveraging shared features across resting-state and task fMRI and demonstrate in the Human Connectome Project and the Dunedin Study that GFC offers better test-retest reliability than intrinsic connectivity estimated from the same amount of resting-state data alone. Furthermore, at equivalent scan lengths, GFC displays higher heritability on average than resting-state functional connectivity. We also show that predictions of cognitive ability from GFC generalize across datasets, performing as well or better than resting-state or task data alone. Collectively, our work suggests that GFC can improve the reliability of intrinsic connectivity estimates in existing datasets and, subsequently, the opportunity to identify meaningful correlates of individual differences in behavior. Given that task and resting-state data are often collected together, many researchers can immediately derive more reliable measures of intrinsic connectivity through the adoption of GFC rather than solely using resting-state data. Moreover, by better capturing heritable variation in intrinsic connectivity, GFC represents a novel endophenotype with broad applications in clinical neuroscience and biomarker discovery.

Introduction

Functional magnetic resonance imaging (fMRI) has proven invaluable for identifying the neural architecture of human behavior and cognition (Betti et al., 2013; Fox et al., 2007; Huth et al., 2016). Accordingly, fMRI has been widely adopted in myriad studies seeking to deepen our understanding of the human brain in both health and disease (Cole et al., 2011; Power et al., 2013; Satterthwaite et al., 2016). Recently, fMRI studies have expanded in both scale and scope, often collecting data in thousands of individuals in an effort to adequately power the search for neural correlates of complex human traits and predictive biomarkers for mental illness (Casey et al., 2018; Elliott et al., 2018; Miller et al., 2016; Swartz et al., 2015; Thompson et al., 2014). In this context, most studies have focused on the acquisition of resting-state functional MRI to map the intrinsic connectivity of neural networks. This choice is often prompted by three considerations. First, intrinsic connectivity networks appear to be more heritable than task-elicited activity (Elliott et al., 2017; Ge et al., 2017; Winkler et al., 2010). Second, resting-state data are relatively easy to collect from informative populations of children, the elderly, and mentally ill patients (Fox, 2010; Greicius, 2008; Shehzad et al., 2009). Third, intrinsic connectivity plays a primary role in shaping task-based brain activity and associated behaviors (Cole et al., 2016, 2014; Krienen et al., 2014; Tavor et al., 2016).

While analyses of resting-state data have revealed insights about average, group-level features of the human brain (Buckner et al., 2008; Power et al., 2011; Yeo et al., 2011), progress has lagged in identifying individualized signatures, which are critical if ongoing large-scale studies are to be successful in the search for risk and disease biomarkers. This lack of progress partially reflects the generally poor test-retest reliability of intrinsic connectivity estimates in many resting-state datasets (Birn et al., 2013; Guo et al., 2012). The reliability of intrinsic connectivity measures must

be improved, as unreliability limits the ability to predict individual differences that could become

clinical biomarkers (Nunnally Jr., 1970; Vul et al., 2009). In this paper we propose general

functional connectivity (GFC), a simple method for combining resting-state and task fMRI data to

improve the reliability of intrinsic connectivity.

Historically researchers have collected 5-10 minutes of resting-state data in response to both

published recommendations (Shehzad et al., 2009; Van Dijk et al., 2010) and practical limitations

(e.g., scan time, cost and subject stamina). While such data may be adequate for detecting patterns

of intrinsic connectivity common across individuals, more than 25 minutes of data are needed to

reliably detect individual differences in these same patterns (Anderson et al., 2011; Hacker et al.,

2013; Laumann et al., 2015). This suggests that intrinsic connectivity may simply be a noisy

measure of stable individual differences and that lengthier measurement is needed to identify the

"true" signal (Braga and Buckner, 2017; Gordon et al., 2017; Gratton et al., 2018; Mueller et al.,

2013). However, due to the high cost of scan time and the limited ability of many individuals

including children, the elderly, and the mentally ill to lie still without falling asleep for tens of

minutes of scanning, this level of protracted measurement is simply not feasible for a majority of

studies (Power et al., 2012; Satterthwaite et al., 2013; Tagliazucchi and Laufs, 2014). To illustrate

the point, two recent meta-analyses of resting-state intrinsic connectivity in depression (Kaiser et

al., 2015) and schizophrenia (Dong et al., 2018) revealed average resting-state scan times of 6.53

minutes (k = 23 studies) and 6.24 minutes (k = 36 studies), respectively. It is likely that intrinsic

connectivity based on these brief scans has low reliability, reducing the ability to discover

individual differences and related biomarkers. If measures of intrinsic connectivity are to realize

their potential as a clinical tool (Fox, 2010; Matthews et al., 2006) and contribute to personalized

medicine (Collins and Varmus, 2015; Hamburg and Collins, 2010), there must first be substantial

improvement in their reliability, and this should be ideally achieved within the practical limitations

of study design and participant burden.

Although few studies collect enough resting-state data to achieve high levels of reliability, many

collect multiple task scans in addition to a resting-state scan. While it has been implicitly assumed

that task and resting-state scans are two separate measures of brain function, to be analyzed and

studied independently, a growing body of evidence suggests that the intrinsic connectivity

measured by each may have substantial overlap, sharing over 80% of the same variance (Cole et

al., 2014; Geerligs et al., 2015; Tavor et al., 2016). In fact, it has long been known that intrinsic

networks extracted from task scans are quite similar to those extracted from resting-state scans,

and task scans have been used for intrinsic connectivity analyses when resting-state data are absent

(Arfanakis et al., 2000; Fair et al., 2007; Fox et al., 2006; Smith et al., 2009). Further, mental

operations that occur during task states have been linked to resting-state functional networks

(Bzdok et al., 2016; Leech et al., 2012), and task scans have been shown to accentuate individual

differences in network dynamics (Finn et al., 2017; Greene et al., 2018; Satterthwaite et al., 2018).

Despite this evidence for a complementary relationship between resting-state and task fMRI, they

are typically investigated separately. Even when researchers have collected both task and resting-

state fMRI in the same subjects, analyses of intrinsic connectivity are almost always performed on

a single short resting-state scan despite the low test-retest reliability of short scan data. If it could

be demonstrated that the reliability of intrinsic connectivity can be improved by adding task to

resting-state fMRI, many researchers could immediately benefit by adopting this approach in data

they have already collected, and also in designing future studies. The field of individual-

differences neuroscience would also benefit from the resulting increase in reliability, replicability,

and power that would follow the widespread adoption of combining task and resting-state fMRI.

Here we hypothesized that measures of intrinsic connectivity will be more reliable when task and

resting-state scans are combined into a longer dataset (i.e., GFC) compared to short resting-state

scans alone. Additionally, we hypothesized that this boost in reliability with GFC will increase the

amount of heritable variation detectable in intrinsic connectivity, because the detection of

heritability requires reliable measurement. Finally, we hypothesized that GFC would measure a

common backbone of intrinsic connectivity common across task and resting-state data allowing

for better out-of-sample prediction of individual differences. We examined two existing datasets

in 3 different ways to test these hypotheses. First, we investigated the test-retest reliability of

different combinations of task and resting-state data in both the Human Connectome Project (HCP

(Van Essen et al., 2013)), which uses state-of-the-art fMRI hardware and acquisition parameters

in a highly educated, healthy sample, and in the Dunedin Study (Poulton et al., 2015), which uses

more common fMRI hardware and acquisition parameters in a population-representative birth

cohort. Second, we estimated the heritability of GFC and resting-state functional connectivity

(RSFC) by leveraging the twin sampling design of the HCP. Third, we investigated the ability of

GFC to measure generalizable individual differences in cognitive ability across samples and across

task states by testing out-of-sample prediction of cognitive ability in the HCP and Dunedin Study

samples.

Materials and Methods

Datasets

Human Connectome Project. This is a publicly available data set that includes 1206 participants

with extensive MRI and behavioral measurement (Van Essen et al., 2013). In addition, 45 of these

subjects completed the entire scan protocol a second time (referred to hereafter as "test-retest

sample"). All participants were free of current psychiatric or neurologic illness and were 25-35

years of age. In all analyses, subjects were excluded if they had truncated scans, less than 40

minutes of combined task data, or less than 40 minutes of resting-state data after censoring. Much

of the HCP dataset consists of twins and family members. We exploited this feature of the design

for conducting heritability analyses. For these analyses, 943 participants from 420 families met

inclusion criteria (144 monozygotic families, 85 dizygotic families, and 191 full sibling families).

To avoid bias due to family confounding in the prediction analyses, only one individual per family

was retained (the family member with the least motion during scanning). This resulted in 298

subjects left to be used in our heritability analyses. Eight subjects were removed from our test-

retest analyses because they had at least one fMRI scan with truncated or missing data. Four

subjects were removed because they did not have sufficient data (i.e., > 40 minutes of rest and >

40 minutes of task) after censoring. Therefore, the test-retest reliability analyses included data

from 33 participants.

The acquisition parameters and minimal preprocessing of these data have been described

extensively elsewhere (Glasser et al., 2013). In our analyses, we used the minimally preprocessed

data in volumetric Montreal Neurological Institute (MNI) space ("fMRIVolume" pipeline).

Briefly, participants underwent extensive MRI measurement that included T1 and T2 weighted

structural imaging, diffusion tensor imaging, and nearly two hours of resting-state and task fMRI.

One hour of resting-state fMRI was collected on each participant in four 15-minute scans (1200

time-points each) split-up into two scanning sessions over two days. In each scan session the two

resting-state scans were followed by task fMRI (Smith et al., 2013). Across the two sessions, each

participant completed seven fMRI tasks described extensively elsewhere (Barch et al., 2013).

Briefly, tasks were designed to identify functionally relevant "nodes" in the brain and included

working memory (810 timepoints, 10:02 minutes), gambling (506 timepoints, 6:24 minutes), motor function (568 timepoints, 7:06 minutes), language (632 timepoints, 7:54 minutes), social cognition (548 timepoints, 6:54 minutes), relational processing (464 timepoints, 6:52 minutes) and emotional processing (352 timepoints, 4:32 minutes). Altogether, 4800 timepoints totaling 60 minutes of resting-state fMRI and 3880 timepoints totaling 48:30 minutes of task fMRI were collected on each participant.

Dunedin Study. This is a longitudinal investigation of health and behavior in a complete birth cohort of 1,037 individuals (91% of eligible births; 52% male) born between April 1972 and March 1973 in Dunedin, New Zealand (NZ), and eligible based on residence in the province and participation in the first assessment at age three. The cohort represents the full range of socioeconomic status on NZ's South Island and matches the NZ National Health and Nutrition Survey on key health indicators (e.g., BMI, smoking, GP visits) (Poulton et al., 2015). The cohort is primarily white; fewer than 7% self-identify as having non-Caucasian ancestry, matching the South Island (Poulton et al., 2015). Assessments were carried out at birth and ages 3, 5, 7, 9, 11, 13, 15, 18, 21, 26, 32, and 38 years, when 95% of the 1,007 study members still alive took part. Neuroimaging data collection is ongoing in participants who are now aged 45 years. We currently have collected completed neuroimaging data for 756 study members. Data were excluded if the study member did not complete the rest scan and all four task scans or if they did not have sufficient data after censoring and bandpass filtering was applied to the resting-state and each task scan separately (see below for censoring details). Data for 591 subjects survived these fMRI exclusion criteria and were included in analyses. Additionally, 20 study members completed the entire scan protocol a second time (mean days between scans = 79). Of these one failed to have sufficient

degrees of freedom after censoring and bandpass filtering. Therefore, the test-retest reliability

analyses included data from 19 participants.

Each participant was scanned using a Siemens Skyra 3T scanner equipped with a 64-channel

head/neck coil at the Pacific Radiology imaging center in Dunedin, New Zealand. High resolution

structural images were obtained using a T1-weighted MP-RAGE sequence with the following

parameters: TR = 2400 ms; TE = 1.98 ms; 208 sagittal slices; flip angle, 9°; FOV, 224 mm; matrix

=256×256; slice thickness = 0.9 mm with no gap (voxel size 0.9×0.875×0.875 mm); and total scan

time = 6:52 minutes. Functional MRI was collected during resting-state and four tasks with a series

of 72 interleaved axial T2-weighted functional slices acquired using a 3-fold multi-band

accelerated echo planar imaging sequence with the following parameters: TR = 2000 ms, TE = 27

ms, flip angle = 90°, field-of-view = 200 mm, voxel size = 2mm isotropic, slice thickness = 2 mm

without gap.

8:16 minutes (248 TRs) of resting-state fMRI was collected immediately before the four task fMRI

scans. During the resting-state scan participants were instructed to stay awake with their eyes open

while looking at a grey screen. Participants completed an emotion processing task (6:40 minutes,

200 TRs), a color Stroop task (6:58 minutes, 209 TRs), a monetary incentive delay (MID) task

(7:44 minutes, 232 TRs) and an episodic memory task (5:44 minutes, 172 TRs). All four tasks are

described in detail in the supplement.

fMRI pre-processing

Minimal preprocessing was first applied to all data. For the HCP dataset this was done with the

HCP minimal preprocessing pipeline (Glasser et al., 2013) and included correction for B0

distortion, realignment to correct for motion, registration to the participant's structural scan,

normalization to the 4D mean, brain masking, and non-linear warping to MNI space.

Minimal preprocessing steps were applied to the Dunedin Study dataset using custom processing

scripts. Anatomical images were skull-stripped, intensity-normalized, and nonlinearly warped to a

study-specific average template in the standard MNI stereotactic space using the ANTs SyN

registration algorithm (Avants et al., 2008; Klein et al., 2009). Time-series images were despiked,

slice-time-corrected, realigned to the first volume in the time-series to correct for head motion

using AFNI tools (Cox, 1996), coregistered to the anatomical image using FSL's Boundary Based

Registration (Greve and Fischl, 2009), spatially normalized into MNI space using the non-linear

ANTs SyN warp from the anatomical image, and resampled to 2mm isotropic voxels. Dunedin

Study images were additionally corrected for B0 distortions.

Time-series images for each dataset were further processed to limit the influence of motion and

other artifacts. Voxel-wise signal intensities were scaled to yield a time-series mean of 100 for

each voxel. Motion regressors were created using each participant's 6 motion correction

parameters (3 rotation and 3 translation) and their first derivatives (Jo et al., 2013; Satterthwaite et

al., 2013) yielding 12 motion regressors. Five components from white matter and cerebrospinal

fluid were extracted using CompCorr (Behzadi et al., 2007) and used as nuisance regressors, along

with the mean global signal. In the Dunedin Study dataset, images were bandpass filtered to retain

frequencies between .008 and .1 Hz. In the HCP dataset, images only underwent highpass filtering

with a cutoff of .008 Hz. High frequency signals were retained because removing high frequency

signals would have resulted in excessive loss of degrees of freedom due to the very low TR (.75

seconds) (Bright et al., 2017; Caballero-Gaudes and Reynolds, 2017). In the HCP dataset, we

followed the empirically derived thresholds of .39 mm frame-wise displacement or 4.9 units above

the median DVARS as recommended (Burgess et al., 2016). In the Dunedin Study dataset, we

investigated a range of framewise-displacement cutoffs using QC-RSFC plots in order to derive

the optimal threshold for removing motion artifact as recommended (Power et al., 2014). This

investigation led to thresholds of 0.35 mm framewise-displacement and 1.55 standardized

DVARS. In both datasets, nuisance regression, bandpass filtering, censoring, global-signal

regression, and smoothing of 6mm FWHM for each time-series were performed in a single

processing step using AFNI's 3dTproject. Identical time-series processing was applied to resting-

state and task time-series data with one exception.

To remove functional connectivity predominantly driven by task-evoked coactivation, signal due

to task structure was added as an additional nuisance covariate to all task scans and removed from

their time-series (Fair et al., 2007) (see supplemental for details on task modeling and regression).

However, we examined the impact of task structure on our estimates of intrinsic connectivity by

conducting parallel analyses without the regression of task structure. While task regression had a

small but consistent effect of lowering ICCs for intrinsic connectivity estimates (see supplemental

Table S1), the ability to predict cognitive ability was slightly higher when task regression was

applied (see supplemental Table S2). Thus, we focus on analyses with task regression in our

primary results.

Effects of Global Signal Regression

Global signal regression (GSR) was performed to adopt a conservative approach to motion artifact

reduction and used in our main analyses (Power et al., 2018). However, because use of GSR is still

debated (Murphy and Fox, 2017), we also preprocessed our data with parallel preprocessing
methods without GSR (see supplemental figure S1).

Functional Connectivity processing

We investigated whole-brain intrinsic connectivity using two parcellation schemes. In the
reliability and prediction analyses (described below), a 264-region parcellation was utilized. This
parcellation was derived in a large independent dataset (Power et al., 2011). BOLD data were
averaged within 5mm spheres surrounding each of the 264 coordinates in the parcellation. In the
heritability analyses we used a smaller 44-brain-region parcellation scheme due to computational
constraints imposed by the time needed to fit heritability models at each edge independently
(described below). This parcellation was derived from the 7-network parcellation scheme
described in (Yeo et al., 2011). These 7 networks were broken up into 44 spatially contiguous
regions across the two hemispheres. Average time series were extracted from all 44 regions. For
both parcellations the average time-series was extracted independently from each scan session in
every dataset. This allowed the time-series to be flexibly concatenated and recombined.
Correlation matrices were derived from these time-series using Pearson correlation, resulting in
34,716 edges in the Power et al. parcellation and 946 edges in the Yeo et al. parcellation.

Test-retest reliability

Intraclass correlation (ICC) was used to quantify the test-retest reliability of intrinsic connectivity
measures in the HCP and Dunedin Study test-retest datasets. ICC (3,1) was used in all analyses
(Chen et al., 2018). In the HCP dataset, the influence of the amount of data, as well as type of data
(resting-state or task) on ICC was investigated. In the resting-state analyses, intrinsic connectivity
matrices were calculated across a range of scan lengths (5, 10, 20, 30, and 40 minutes of post-

censoring data) mirrored in the test and retest samples. ICCs were then calculated for each edge of

this matrix, yielding 34716 (264*263/2) unique ICC values for each scan length. A more general

definition of intrinsic connectivity was derived by including task data in the intrinsic connectivity

matrices. GFC was formally investigated (Figures 1, 2 and 3) by constructing datasets from a

combination of task and resting-state data over a variety of scan lengths (5, 10, 20, 30 and 40

minutes of post-censoring data). In scan lengths up to 25 minutes, an exactly equal amount of data

from all scan types (1/8 of total scan length from each task and resting-state) was combined. After

this point, equal amounts of each task could no longer be added together because the shortest task

scan no longer had sufficient timepoints remaining in all participants. Above 25 minutes,

timepoints were selected at random from the pool of remaining timepoints. In the Dunedin Study,

two ICC matrices were constructed. The first was built from each participant's single resting-state

scan and the second from all task and resting-state scans to create the GFC matrix. Paired-sample

t-tests were used to compare mean differences in reliability between ICC matrices constructed

from traditional resting-state functional connectivity and GFC.

HCP fMRI data were collected in two phase-encoding directions (LR and RL). To minimize bias

due to phase-encoding direction on our analyses, in each participant, for each task, timepoints were

first selected from LR encoded scans until LR timepoints were exhausted. Subsequently,

timepoints were selected from remaining RL-encoded scans. This ensured that all subjects had a

roughly equivalent proportion of LR and RL timepoints within a given analysis. This also ensured

roughly equivalent proportions of LR and RL phase encoding across test and retest correlation

matrices when calculating ICCs. Selecting from LR first before RL had the additional benefit of

mimicking shorter data-acquisition scans because timepoints were utilized in the order they were

collected.

Heritability

For each edge of the connectivity matrix, we ran an extended twin model (Posthuma and Boomsma, 2000), which is an extension of the classic twin design in that it allows for the modeling of additional family members of twins. In the HCP study we had data from the families of monozygotic and dizygotic twins as well as siblings in families without a twin pair. In this model the variance of the phenotype (in this case correlations between brain regions) is partitioned into four sources: additive genetic factors (A), shared environmental factors (C), twin-specific factors (T), and unique environmental factors and error (E) based on the correlations between monozygotic twins, dizygotic twins, and their siblings. Due to their small numbers, we eliminated half-siblings from the model (N = 27) and capped family size at four individuals (7 families affected, N = 13). This left us with 1,078 individuals from 443 families (148 monozygotic families, 92 dizygotic Z families, 203 full-sibling families). Of these, 943 participants from 420 families (144 monozygotic families, 85 dizygotic families and 191 full-sibling families) had full scans from all tasks and rest and had at least 40 minutes of task data and 40 minutes of rest data after censoring. Heritability analyses were performed using OpenMx (Boker et al., 2011; Neale et al., 2016) in the R statistical computing environment. Each model also contained parameters to adjust for the effects of sex and age on the mean. Confidence intervals were obtained on the A, C, T, and E estimates using the mxCI command in OpenMx which obtains profile likelihood confidence intervals (Pek and Wu, 2015). For each edge, an estimate was determined to be significant if the lower bound of the confidence interval was greater than zero.

Connectome-based predictive modeling

The predictive utility of increasing reliability through our GFC measure was evaluated by testing

the ability of the different intrinsic connectivity matrices to predict cognitive ability in the HCP

and Dunedin Study datasets. Cognitive ability was measured in the HCP using the Raven's

Progressive Matrices (PMAT24_A_CR) (Raven, 1941). This measure was adopted because it is a

proxy for cognitive ability that has been shown to be predicted by intrinsic connectivity in the HCP

(Dubois et al., 2018; Finn et al., 2015; Noble et al., 2017). The WAIS-IV, a well-established and

validated measure of individual differences in cognitive ability (Weschler, 2008), was used in the

Dunedin Study.

Cognitive ability was predicted from intrinsic connectivity data using connectome-based

predictive modeling (CPM; (Shen et al., 2017)). This framework provides a general method to

predict any measure from intrinsic connectivity matrices. In this approach, predictors are filtered

by selecting edges that have a $p < .01$ correlation with the measure of interest. Three linear

regression predictive models are then built, one from the positive features (edges positively

correlated with the phenotype of interest), one from the negative features (edges negatively

correlated with the phenotype of interest), and one from the combination of positive and negative

features (Shen et al., 2017). Here we report results from the combined model (see supplemental

tables S3, S4 and S5 for results from all models). In our CPM analyses, we used both within-

sample and out-of-sample prediction. Within-sample models were used to directly compare RSFC

and GFC predictions of cognitive ability using a leave-one-out cross-validation scheme. Models

were trained with all participants except one and used to predict the measure in the left-out

participant. This was repeated until all participants had been left out.

Out-of-sample prediction is the gold standard to test the unbiased predictive utility of models

(Whelan and Garavan, 2014; Yarkoni and Westfall, 2017) and was therefore used to test the

generalizability of predictive models built with RSFC, GFC, and task data. To test the influence

of task state, data aggregation and scan length on predictive utility, models were trained with

RSFC, GFC, and individual tasks in the Dunedin Study dataset and then used to predict cognitive

ability from RSFC, GFC, and individual tasks in the HCP dataset. The Dunedin Study and HCP

have three parallel tasks that tap into similar behavioral circuits. The gambling task in the HCP

and the MID task in Dunedin Study both involve reward processing. The emotional processing

task in the HCP and the face matching task in the Dunedin Study both involve passive processing

of emotional facial expressions. Lastly, the working memory task in the HCP and the Stroop task

in the Dunedin Study both tap into executive control. We restricted our analyses to comparisons

of RSFC, GFC, and these 3 parallel tasks. The abilities of RSFC and GFC was contrasted with that

of these parallel tasks to predict cognitive ability within tasks. In both within-sample and out-of-

sample tests the Spearman correlation between predicted and true scores was adopted as an

unbiased effect size measure of predictive utility. Model predictions of cognitive ability were

assessed for significance using a parametric test for significance of correlations. All p-values from

correlations with cognitive ability were corrected for multiple comparisons using the false

discovery rate (Benjamini and Hochberg, 1995). Differences in predictive utility between models

were compared using Steiger's z (Steiger, 1980). All confidence intervals for CPM prediction

estimates were generated with bootstrap resampling, using AFNI's 1dCorrelate tool.

Results

What is the test-retest reliability of GFC?

We used data from the HCP to evaluate the test-retest reliability of both intrinsic connectivity

derived from resting-state scans alone and of GFC derived from combinations of task and resting-

state scans. Test-retest reliability, as measured by ICC, ranges from 0-1, and is commonly

classified according to the following cutoffs: 0 - 0.4 = poor, 0.4 - 0.6 = moderate, 0.6 - 0.75 = good

and 0.75 - 1 = excellent (Cicchetti, 1994). In the HCP, when 5 minutes (post-censoring) of resting-

state data were used to measure intrinsic connectivity defined within a common atlas of 264

regions (Power et al., 2011), the reliability was generally poor (mean ICC = .28, 95% CI [.28, .28];

71% of edges were poor , 23% moderate, 6% good, and less than 1% excellent). As more resting-

state data were added, the test-retest reliability continued to increase up to the limits of the dataset

at 40 minutes (mean ICC = .54, 95% CI [.54, .54]; 26% of edges poor, 32% moderate, 25% good,

and 18% excellent) (Figure 1A and left panel of Figure 2).

394

395
396 **Figure 1.** Test-retest reliability of individual differences in global intrinsic network connectivity increases as the
397 amount of data used to estimate either RSFC (A) or GFC (B) increases. Stacked bar plots from the HCP dataset
398 displaying the proportion of functional connections (i.e., edges) across neural networks as defined by (Power et al.,
399 2011) that meet criteria for excellent, good, moderate, and poor reliability as indexed by ICCs.
400
401
GFC, defined as exactly equal parts of task and resting-state data in the HCP, followed a similar

pattern of increasing reliability with increasing data: 5 minutes of data exhibited poor reliability

(mean ICC = .28, 95% CI [.28, .28]; 70% of edges were poor, 22% moderate, 7% good, and less

than 1% excellent) but 40 minutes of data showed good reliability (mean ICC = .56, 95% CI [.55,

.56]; 25% of edges were poor, 27% moderate, 24% good, and 23% excellent) (see Figures 1B,

right panel of Figure 2). In addition, resting-state scans were not required to generate a reliable

measure of GFC, as this pattern of increasing reliability held when task data alone were used to

measure intrinsic connectivity (see supplemental Table S1). Furthermore, data from individual

task scans alone generally resulted in poor reliability (see supplemental figures S2 and S3). A

paired sample t-test comparing ICCs across all edges revealed that GFC is significantly more

reliable than resting-state functional connectivity when both are estimated with 40 minutes of data

$(t(34715) = 140.01, p < .001)$. While significant, this difference is not likely meaningful as it

represents only a 0.02 larger mean ICC in GFC. The low p-value is driven by the large number of

degrees of freedom in the t-test comparing a mean ICC estimate from nearly 35,000 edges. Thus,

we see the mean ICC of RSFC and GFC as statistically separable but practically equivalent.

This general improvement in reliability with increasing data was not unique to the HCP but was

also observed in the population-representative Dunedin Study. The Dunedin Study resting-state

scan of 8:16 minutes showed generally low reliability (mean ICC = .30, 95% CI [.30, .30]; 65%

of edges were poor, 23% moderate, 10% good, and 2% excellent). The average reliability of the

intrinsic connectivity estimates, however, improved by 49% when using GFC created in the

Dunedin Study by combining the 8:16 minutes of resting-state data with 27 minutes (before

censoring) of task data (mean ICC = .45, 95% CI [.45, .45]; 41% of edges were poor, 26%

moderate, 19% good and 14% excellent) (Supplemental Figure S3).

Figure 2. Test-retest reliability of the intrinsic connectivity of canonical neural networks derived from either RSFC
or GFC from the HCP scales with the amount of data available for analysis. Intra-Class Correlation (ICC) matrices
for RSFC (left panel) and GFC (right panel) demonstrate comparable gains in reliability with increasing amounts of
data across common intrinsic networks (Power et al., 2011). 5 and 10 minutes are written in red because these are
common scan lengths for resting-state scans. 30 and 40 minutes are written in blue because many researchers have
collected this amount of fMRI data when resting-state and task scans are combined. To the bottom left of ICC matrices
is the color key for the ICCs, with a histogram indicating the density of ICCs for the corresponding graph.
434

Influence of the global signal on reliability

GSR is still a controversial preprocessing step. Therefore, the main reliability analyses in the HCP

were rerun without GSR to test the influence of GSR on ICCs. Mean ICCs for 5 minutes of data

without GSR were poor for both RSFC (mean ICC = .31, 95% CI [.31, .31]; 68% of edges were

poor, 27% moderate, 4% good, and < 1% excellent) and GFC (mean ICC = .35, 95% CI [.35, .35];

60% of edges were poor, 31% moderate, 8% good, and 1% excellent) (see supplemental figure

S1). In parallel fashion to the data processed with GSR, ICCs increased as scan length increased

up to 40 minutes for both RSFC (mean ICC = .61, 95% CI [.61, .61]; 8% of edges were poor, 34%

moderate, 41% good, and 16% excellent) and GFC (mean ICC = .63, 95% CI [.63, .63]; 8% of

edges were poor, 30% moderate, 40% good, and 22% excellent). Despite the improvement in mean

ICC with additional data with and without GSR, ICCs were significantly higher when GSR was

not applied: 40 minutes of GFC had an average ICC that was 0.07 higher without GSR than when

GSR was applied. The same pattern was present in RSFC: 40 minutes of resting-state without GSR

was more reliable on average than with GSR (rest with GSR = .61, rest without GSR = .54). While

higher reliabilities without GSR could be a sign of improved detection of true individual

differences in intrinsic connectivity, they may also be a sign of a stable propensity to move within

the scanner, creating correlated motion artifact in test and retest data (Power et al., 2016; van Dijk

et al., 2012). Because the global signal is highly susceptible to motion artifact and there is no

widely accepted method to isolate and remove just the artifactual component of the global signal

while retaining the neural component, we adopted a conservative approach by applying GSR signal

in all further analyses to ensure removal of motion related artifact present in the global signal

(Power et al., 2018).

457
458   *Is there network specificity to improvements in reliability?*
459
We next investigated network specificity in reliability by looking specifically at the reliability of

edges within each functional network rather than the reliability of the entire correlation matrix in

aggregate. As shown in Figure 3, there was clear network heterogeneity in reliability. While all

networks improved in reliability from 10 minutes of data to 40 minutes, some networks improved

more. With both RSFC and GFC, the limbic network had the lowest reliability at all scan lengths,

improving from 10 minutes of data (RSFC: mean ICC = .22, 95% CI [.19, .25]; GFC: mean ICC

= .30, 95% CI [.25, .34]) to 40 minutes of data (RSFC: mean ICC = .37, 95% CI [.33, .41]; GFC:

mean ICC = .39, 95% CI [.35, .44]), but failing to escape a poor mean ICC. In contrast, networks

that over-represent heteromodal association cortex (Margulies et al., 2016; Mesulam, 1998),

namely the default mode network and fronto-parietal network, on average achieved moderate

reliability with just 10 minutes of data for both RSFC (mean fronto-parietal network ICC = .53,

95% CI [.52, .53]; mean default mode network ICC = .57, 95% CI [.57, .58]) and GFC (mean

fronto-parietal network ICC = .55, 95% CI [.54, .56]; mean default mode network ICC = .57, 95%

CI [.57, .58]). Good reliability was achieved for these networks with 20 minutes of data for both

RSFC (mean fronto-parietal network ICC = .62, 95% CI [.61, .62]; mean default mode network

ICC = .66, 95% CI [.66, .67]) and GFC (mean fronto-parietal network ICC = .64, 95% CI [.63,

.64]; mean default mode network ICC = .65, 95% CI [.65, .66]). With 40 minutes of data, these

networks had mean reliabilities near the excellent range for both RSFC (mean fronto-parietal

network ICC = .73, 95% CI [.72, .73]; mean default mode network ICC = .75, 95% CI [.75, .75])

and GFC (mean fronto-parietal network ICC = .74, 95% CI [.74, .75]; mean default mode network

ICC = .77, 95% CI [.77, .77]). While there were some small differences in the network-specific

reliabilities between RSFC and GFC (e.g., see dorsal attention network in Figure 3), network

specific improvements with increasing scan length were similar for RSFC and GFC.

484
485 **Figure 3.** The improvement in test-retest reliability of intrinsic connectivity as a function of adding data varies across
486 canonical functional networks. In the left panel mean ICC for RSFC are displayed for within-network connections
487 across 7 previously defined networks (Yeo et al., 2011) for a variety of scan lengths. In the right panel the same data
488 are displayed for GFC. There are clear and consistent differences between networks. The limbic network consistently
489 has the lowest mean ICC, while the default mode network consistently has the highest mean ICC.

490
491
What is the heritability of RSFC and GFC at different scan lengths?

As shown in Figure 4A, the heritability of intrinsic connectivity depended on scan length. The

amount of variance for individual differences in RSFC attributable to the additive genetics (A)

component of the ACE model increased consistently as the amount of data increased from 5

minutes (mean A component = .09, 95% CI [.09, .1]) to 40 minutes (mean A component = .22,

95% CI [.21, .23]). This represents an increase in the amount of variance in RSFC attributable to

additive genetics of 138% as a function of increased scan length. This increase in variance

explained by additive genetics was also present with GFC from 5 minutes (mean A component =

.14, 95% CI [.13, .14]) to 40 minutes (mean A component = .28, 95% CI [.27, .29]). This

represented an increase in the amount of variance in GFC attributable to additive genetics of 107%

as a function of increased scan length. In addition, at equivalent scan lengths, GFC had

significantly higher mean heritability across edges than RSFC (e.g., 5 minutes: $t(945) = 10.11$, p

$< .001$) and 40 minutes: ($t(945) = 14.75$, $p < .001$)). Correspondingly, higher mean heritability in

GFC than RSFC led to a greater percentage of heritable edges in GFC (see Figure 4B). In the ACE

modeling, this pattern of increased heritability with increased scan time was associated with

simultaneously decreased amount of variance explained by the E component, which comprises

both non-shared environment and measurement error (Figure 4C). The E component decreased for

both RSFC and GFC as data length increased from 5 minutes (RSFC mean E component = .81,

95% CI [.81, .82]; GFC mean E component = .77, 95% CI [.76, .77]) to 40 minutes (RSFC mean

E component = .63, 95% CI [.63, .64]; GFC mean E component = .61, 95% CI [.60, .61]).

While the A and E components of the ACE modeling changed substantially with scan length, the

shared-environment (C) (Figure 4D) and twin-specific (T) (supplemental Figure S4) components

were relatively stable with scan length. The variance attributable to the C and T components was

similar for 5 minutes of RSFC (mean C component = .04, 95% CI [.03, .04]; mean T component

= .06, 95% CI [.05, .06]) and 5 minutes of GFC (mean C component = .04, 95% CI [.03, .04];

mean T component = .06, 95% CI [.06, .07]) as well as with 40 minutes of RSFC (mean C

component = .06, 95% CI [.05, .06]; mean T component = .09, 95% CI [.08, .09]) and 40 minutes

of GFC (mean C component = .05, 95% CI [.05, .06]; mean T component = .06, 95% CI [.06,

.07]).

Exploring the connection between intrinsic connectivity and cognitive ability

CPM was used to predict cognitive ability from both RSFC and GFC in both the HCP and the

Dunedin Study. In the first set of analyses, within-sample predictions were made using leave-one-

out cross validation in the HCP and Dunedin Study datasets separately (Figure 5, left and center

panels). In the HCP dataset, we found that individual differences in cognitive ability could be

predicted from 40 minutes of RSFC (r^2 = 4.2%, 95% CI [1.0%, 9.7%]) and from 40 minutes of

GFC (r^2 = 7.6%, 905% CI [2.7%, 14.2%]) (see Figure 5 and supplemental Table S3). These results

were replicated and extended in the Dunedin Study using scan lengths that are more representative

of existing datasets than the HCP. Cognitive ability could be predicted from the 8-minute RSFC

data (r^2 = 6.8%, 95% CI [3.5%, 11.3%]) as well as from 33 minutes of GFC (r^2 = 10.2%, 95% CI

[5.9%, 15.0%]; all available data; supplemental Table S4). Although in comparison with RSFC,

GFC had 81% greater predictive utility in the HCP dataset and 50% greater predictive utility in

the Dunedin Study dataset, the statistical comparison of the correlations was not statistically

significant in either the HCP (GFC r^2 = 7.6%, RSFC r^2 = 4.2%, Steiger's z = 1.317, p = .188) or

the Dunedin Study (GFC r^2 = 10.2%, RSFC r^2 = 6.8%, Steiger's z = 1.72, p = 0.085). In addition,

scan length had little effect on within-sample predictive utility (see supplemental figure S5).

While leave-one-out cross validation has been widely used with CPM (Finn et al., 2015; Noble et

al., 2017), the gold standard test of a model's generalizable predictive utility is its ability to predict

a phenotype in an independent sample (Bzdok and Yeo, 2017; Whelan and Garavan, 2014;

Yarkoni and Westfall, 2017). This is particularly important for GFC, as it purports to measure a

generalizable feature of intrinsic connectivity and thus correlations between individual differences

in GFC and phenotypes of interest should be shared across independent samples with different

batteries of fMRI tasks. Therefore, we used CPM to train two models (one with rest and one with

GFC) to predict cognitive ability in the larger Dunedin Study dataset and tested generalizability

by measuring the ability of these models to predict cognitive ability in the HCP dataset. The models

built with RSFC (r^2 = 5.6%, 95% CI [1.8%, 11.9%], p,q < .001) and GFC (r^2 = 9.5%, 95% CI

[4.0%, 16.3%], p,q < .001) from the Dunedin Study dataset both successfully predicted cognitive

ability in the HCP dataset (Figure 5 right panel and Figure 6 left panel). However, GFC had greater

out-of-sample predictive utility than RSFC (GFC r^2 = 9.5%, RSFC r^2 = 5.6%, Steiger's z = 2.57,

p = 0.010). This provides evidence that GFC is a more generalizable measure across independent

samples than RSFC.

566
567 **Figure 5.** GFC is better than RSFC at predicting cognitive ability both within and between samples. Results from
568 CPM models predicting cognitive ability from RSFC and GFC. The x-axis displays predictions from leave-one-out
569 cross validation within sample and out-of-sample models trained using the Dunedin Study dataset and tested using the
570 HCP dataset. Predictive utility is displayed as % variance explained (r^2).
571

Does GFC predict cognitive ability better than intrinsic connectivity measured from specific tasks?

Recent research has suggested that intrinsic connectivity estimates derived from task data are better

at predicting cognitive ability than estimates derived from resting-state data (Greene et al., 2018).

Our above finding that GFC outperforms RSFC at predicting cognitive ability is in line with this

observation. However, given that many researchers collect different tasks and resting-state scans

with different "task" demands (e.g., eyes-closed, fixation, etc.), we next investigated if GFC could

perform as well or better than specific tasks at out-of-sample prediction of cognitive ability. We

constructed five prediction models that were trained to predict cognitive ability using the Dunedin

Study dataset (the population-representative sample with the gold-standard WAIS IQ test) and

tested for generalizability using the HCP dataset (see supplemental figure S6 for the converse,

models trained in HCP and tested in the Dunedin Study). Two models were carried over from the

previous analyses (RSFC and GFC) and three additional task-specific models were added. These

consisted of models trained and tested on three parallel tasks available in both the Dunedin Study

and HCP (see methods for details). All models trained using the Dunedin Study dataset were able

to significantly predict cognitive ability in the HCP dataset (all FDR q < .05; Figure 6 and

supplemental Table S5). Prediction of cognitive ability from RSFC data (r^2 = 5.6%, 95% CI [1.8%,

11.9%], p,q < .001), GFC data (r^2 = 9.5%, 95% CI [4.0%, 16.3%], p,q < .001), emotional

processing data (r^2 = 9.8%, 95% CI [4.3%, 17.1%], p < .001, q = .001), reward processing data (r^2

= 9.1%, 95% CI [3.6%, 16.5%], p < .001, q = .001) and executive function data (r^2 = 4.6%, 95%

CI [1.0%, 10.8%], p < .001, q = .001) all successfully generalized from the Dunedin Study dataset

to the HCP dataset.

Comparing the relative performance of models trained with each type of data revealed that GFC

was generalizable and performed as well or better at out-of-sample prediction of cognitive ability

compared to all tasks (Figure 6). Predictions of cognitive ability from GFC performed better than

RSFC (GFC r^2 = 9.5%, RSFC r^2 = 5.6%, Steiger's z = 2.57, p = 0.01) and executive function data

(GFC r^2 = 9.5%, executive function r^2 = 4.6%, Steiger's z = 2.19, p = .03), and performed as well

as emotional processing data (GFC r^2 = 9.5%, emotion processing r^2 = 9.8%, Steiger's z = -.10, p

= .92) and reward data (GFC r^2 = 9.5%, reward r^2 = 10.2%, Steiger's z = -.31, p = .76). Out-of-

sample prediction was largely unaffected by the amount of data used in constructing the intrinsic

connectivity matrices but was adversely affected when task regression was removed (see

supplemental Table S2).

Figure 6. Out-of-sample prediction of cognitive ability for GFC is better than RSFC and as good as task-derived intrinsic connectivity. All models were trained on intrinsic connectivity data from the Dunedin Study and tested using data from the HCP. Models were trained and tested with the same type of data. With task data this meant that models were trained and tested with tasks that have a comparable (parallel) task in both the HCP and Dunedin study. Predictive utility is displayed as % variance explained (r^2).

Discussion

Here we present GFC as a method for readily improving the reliability of estimating individual differences in intrinsic connectivity by combining resting-state and task data. Across two independent datasets we found that scan length has a significant impact on the reliability of

intrinsic connectivity estimates regardless of whether the data came solely from resting-state scans

(i.e., RSFC) or from a combination of resting-state and task scans (i.e., GFC). GFC, the addition

of task scans to short resting-state scans, substantially increased the reliability of intrinsic

connectivity over and above short resting-state scans alone. Moreover, we found that the improved

reliability from aggregating data resulted in higher estimates of the heritable variation in intrinsic

connectivity, and that this gain was larger for GFC than RSFC. In addition, GFC consistently

performed better than RSFC and performed as well or better than intrinsic connectivity from

individual task scans at predicting individual differences in a complex human trait, namely

cognitive ability.

These findings have several implications for current and future neuroimaging research because

they demonstrate that 1) reliable and heritable measurement of individual differences in the

functional architecture of the brain is not only achievable in niche, specialty datasets with hours of

resting-state scans, but also in many existing datasets if GFC is adopted; 2) mapping individual

differences in behavior and cognition may be improved by adopting GFC, as demonstrated by

generalizable predictive utility when studying cognitive ability; and 3) to the extent that

commensurate gains in reliability and heritability may be achieved in other datasets, our results

provide a reference that researchers can use to roughly estimate the gain in reliability they may

achieve by adopting GFC (Figures 1, 2, S2 and S3). We next elaborate on each of these points.

Data aggregation across task and resting-state scans boosts reliability

We found that GFC, derived from the combination of resting-state and task scans, can be a reliable

measure of individual differences in intrinsic connectivity. Consistent with previous studies (Birn

et al., 2013; Finn et al., 2015; Laumann et al., 2015), we showed that the test-retest reliability of

intrinsic connectivity is highly dependent on the amount of data collected. This is true of RSFC as

well as GFC. With only 5-10 minutes of data, both RSFC and GFC show poor reliability in the

HCP and Dunedin Study datasets. Only with 30-40 minutes of data do RSFC and GFC begin to

broadly display good reliability. These findings follow directly from past studies that have shown

reliability depends on scan length (Anderson et al., 2011; Birn et al., 2013; Hacker et al., 2013),

that task and resting-state data share a large proportion of variance (Cole et al., 2014; Geerligs et

al., 2015), and that both task and resting-state data measure common individual differences in

intrinsic connectivity (Gratton et al., 2018). Some functional networks, however, can achieve good

reliability with relatively little data. For example, the default mode and fronto-parietal networks

achieved good reliability for within-network connections with just 20 minutes of RSFC or GFC

(see figure 3). In contrast, the limbic network failed to achieve even fair levels of reliability with

up to 40 minutes of data. These findings suggest that the amount of data needed to derive reliable

estimates of intrinsic connectivity for individual differences research will vary depending on the

network of interest.

Importantly, the current reality is that most neuroimaging studies have only 5-10 minutes of

resting-state data, which are further limited by sampling variability, motion-artifacts, and other

sources of noise (Gordon et al., 2017; Gratton et al., 2018; Power et al., 2014). Given the reliability

estimates reported here, it is likely that in many studies, individual differences in intrinsic

connectivity will be unmeasurable with resting-state data alone (Anderson et al., 2011; Hacker et

al., 2013). Fortunately, it is common practice for researchers to collect 10-40 minutes of task fMRI

in addition to resting-state. Therefore, if researchers adopt a broader definition of intrinsic

connectivity, such as GFC, they can combine task and resting-state data to achieve a reliable

measure of individual differences in intrinsic connectivity.

Data aggregation across task and resting-state scans boosts heritability

We found that individual differences in intrinsic connectivity in both RSFC and GFC were significantly attributable to additive genetic influences. With 40 minutes of data, 22% of the variance in RSFC and 28% of the variance in GFC was attributable to additive genetic effects. With equivalent amounts of data, GFC produced higher heritability estimates than RSFC, and the influence of additive genetic effects was detectable in a larger proportion of functional connections (Figure 4). In addition, we found that scan length had an impact on the heritability estimates of intrinsic connectivity in both RSFC and GFC. The average amount of variance in intrinsic connectivity that was attributable to additive genetic effects more than doubled in both RSFC (increase of 138%) and GFC (increase of 107%) as scan length increased from 5 to 40 minutes. While the heritability of intrinsic connectivity has been investigated before (Adhikari et al., 2018; Ge et al., 2017), the link between scan length and heritability has not been described. Although a reliable measure does not have to be heritable, a measure cannot have high heritability without high reliability (Ge et al., 2017). That is, measurement reliability places a ceiling on the heritability estimate of a phenotype. Therefore, our finding that the heritability of intrinsic connectivity increases with increasing reliability suggests that the low reliability of intrinsic connectivity in short fMRI scans limits estimations of the true underlying heritability of intrinsic connectivity in many datasets. More specifically, our finding suggests that researchers should consider the amount of fMRI data available for estimation of intrinsic connectivity when planning and evaluating imaging genetics research.

Our results have additional implications for genetically-informed fMRI research. Neuroimaging measures have been considered promising intermediate phenotypes that would bring researchers

closer to the mechanisms through which genetics lead to heritable psychological traits (Braskie et

al., 2011; Hariri and Weinberger, 2003; Hasler and Northoff, 2011; Meyer-Lindenberg and

Weinberger, 2006). To uncover these links, large datasets like ENIGMA and the UK Biobank

(Sudlow et al., 2015; Thompson et al., 2014), have been used to conduct genome wide association

studies (GWAS) of MRI-based phenotypes. While these investigations have had success in finding

genetic variants linked to structural MRI measures (Adams et al., 2016; Hibar et al., 2017, 2015;

Stein et al., 2012), they have largely failed to find significant genetic correlates of functional

intrinsic connectivity measures (Elliott et al., 2017). One reason for this failure may be the low

reliability and heritability of the short resting-state scans typically used to derive individual

differences measures of RSFC in these studies. Our results suggest that future GWAS may be

better powered to find genetic variants linked to the functional architecture of the brain by adopting

GFC rather than RSFC from short resting-state scans.

How general is general functional connectivity?

Previous research has shown that the predictive utility of intrinsic connectivity can be higher when

derived from task data than when derived from resting-state data (Greene et al., 2018). Some task

states in particular may reveal individual differences in intrinsic connectivity that aid in predictive

utility (Finn et al., 2017). While these findings provide insight into individual differences in

intrinsic connectivity and a new use for task data, they pose a problem for replication and

cumulative aggregation of findings across datasets. While most fMRI studies collect task data,

very few collect the same tasks. Our findings here suggest that GFC may provide a solution to this

problem. First, we found that like task-derived intrinsic connectivity, GFC can outperform RSFC

when predicting within-sample and out-of-sample variability in cognitive ability (Figures 5 and

6). Second, we found that GFC performed as well or better than specific tasks alone at predicting

cognitive ability out-of-sample. Critically, in these comparisons GFC was constructed with a

combination of different tasks in the training and testing datasets suggesting that GFC may

measure a common, generalizable backbone of intrinsic connectivity that can be applied and

aggregated across independent datasets with different task batteries (supplemental Tables S6 and

S7). While further research is needed to replicate these findings, they nevertheless suggest that

GFC may provide a generalizable, practical framework that researchers with resting-state and task

data can use to drive a cumulative, predictive neuroscience of individual differences (Button et al.,

2013; Szucs and Ioannidis, 2017).

Is general functional connectivity too heterogeneous?

An understandable reluctance in implementing GFC could be that it would introduce heterogeneity

in the estimation of intrinsic connectivity. Whereas RSFC is thought to be a common framework

with generalizability across datasets, the combination of different sets of tasks in different datasets

may introduce additional variability. Below, we provide three reasons why we think this is less

problematic than it first appears. We further provide specific recommendations and guidelines for

adopting and estimating GFC and point to features of our analyses that support the validity of GFC

as a robust and reliable measure of individual differences in intrinsic connectivity.

First, resting-state data are by nature heterogenous, and the resting-state is its own type of task

(Buckner et al., 2013). Researchers collect resting-state data under different conditions in which

participants close their eyes, have their eyes open, or fixate. These differences in resting-state

conditions influence intrinsic connectivity (Patriat et al., 2013). Despite these differences, all

approaches are collectively referred to as resting-state scans. In addition, factors like thought

content (Christoff et al., 2009; Hurlburt et al., 2015), caffeine intake (Wong et al., 2012), recent

cognitive tasks (Waites et al., 2005), and falling asleep (Deco et al., 2014) can introduce further

heterogeneity that can bias intrinsic connectivity estimates. It has even been estimated that

approximately 30% of participants fall asleep within the first three minutes of a resting-state scan

(Tagliazucchi and Laufs, 2014). Nevertheless, resting-state scans measure a common set of

functional networks (Yeo et al., 2011), and display trait-like features driven by stable factors such

as genetics (Glahn et al., 2010) and structural connectivity (Honey et al., 2009). For these reasons

resting-state data represent substantial promise as an individual-differences measure if enough data

are collected to average out sampling variability (Gratton et al., 2018). But, as already stated, few

researchers collect enough data to fulfill this promise. Connectivity estimates from task data are

also shaped by many of these stable factors (Cole et al., 2014; Krienen et al., 2014), and data from

different tasks predominantly measure overlapping individual differences that are only weakly

influenced by task demands (Gratton et al., 2018). Admittedly, GFC will vary, to some extent,

across samples because each study collects a unique combination of resting-state and task data.

However, RSFC will vary too. Moreover, we found that disparate task and resting-state scans

collectively measure reliable, heritable individual differences with generalizable out-of-sample

predictive utility in the form of GFC.

Second, the benefits of GFC converged across two different samples, with different scanners,

scanning parameters, and demographics. The HCP represents a healthy sample of highly-educated

individuals (Van Essen et al., 2013), while the Dunedin Study represents a population-

representative birth cohort with a broad range of mental and physical health conditions,

socioeconomic status, and full representation of variability in many complex traits (Poulton et al.,

2015). Across samples, the data came from 11 different tasks (HCP = 7, Dunedin Study = 4), many

with unrelated cognitive demands, different scan lengths, stimuli, behavioral requirements, and

instructions. The data were processed with different preprocessing schemes and software

("minimally preprocessed" in the HCP and custom scripts in the Dunedin Study). While the

amount of scan time was tightly controlled in the HCP analyses (i.e., equal time after censoring),

in the Dunedin Study motion censoring led to unequal scan lengths across participants as is the

case in many analyses of intrinsic connectivity. Despite these many differences between datasets,

the results demonstrate that GFC can achieve good test-retest reliability, inter-rater reliability

(different preprocessing schemes), out-of-sample reliability (convergence across datasets and out-

of-sample prediction), and parallel forms reliability (different tasks in each sample) (Dubois and

Adolphs, 2016). While we present evidence in favor of the generalizability of GFC, future research

should further investigate the heterogeneity in GFC to find combinations of tasks that most

efficiently estimate individual differences in intrinsic connectivity (Finn et al., 2017; Shah et al.,

2016).

Third, reliability fundamentally limits the ability to detect associations between any two measures

(Nunnally Jr., 1970; Vul et al., 2009). Therefore, any investigation mapping intrinsic connectivity

to individual differences in behavior, cognition, or disease is fundamentally limited by the

reliability of the intrinsic connectivity measures. Relatedly, statistical power to detect true effects

depends on the anticipated effect size, which is in turn dependent on reliability of each

measurement (Kanyongo et al., 2007; Williams and Zimmerman, 1989). Given that many studies

only have 5-10 minutes of resting-state data and consequently poor reliability of intrinsic

connectivity measures, they have drastically reduced statistical power (Button et al., 2013;

Ioannidis, 2008, 2005), and likely tenuous correlates of individual differences unless they adopt a

framework like GFC. Previous research has found that collectively a large number of unreliable

edges can achieve multivariate reliability that results in predictive utility (Noble et al., 2017).

While multivariate predictive utility may be achievable in studies with poor univariate edge-wise

reliability (Figure S5), the interpretability and clinical utility of these studies will be diminished

by low univariate reliability. This is because interpretability and clinical utility depend on isolating

specific brain areas or functional connections that can be said to be "most" important to the

phenotype of interest. This cannot be done without decent univariate reliability because accurate

estimation of true parameters in a prediction model depends on reliability (Cremers et al., 2017;

Vul et al., 2009; Yarkoni and Westfall, 2017). For example, if researchers want to characterize

biomarkers of mental illness that could be targeted with interventions like transcranial magnetic

stimulation (Fox et al., 2014) or real-time neurofeedback (Caria et al., 2012), they will need to

isolate specific circuits that with confidence can be said to be "altered" and relevant to the disease

process in mental illness. Actionable identification of these biomarkers will only be possible with

adequate univariate reliabilities. GFC provides a practical solution to this problem, while also

addressing power and reliability issues that have been identified more generally in neuroscience

(Button et al., 2013; Szucs and Ioannidis, 2017) and beyond (Aarts et al., 2015; Errington et al.,

2014; Ioannidis, 2005; Ioannidis et al., 2001; Munafò et al., 2017). GFC provides a tool for

researchers to repurpose existing and future datasets to better contribute to a cumulative science

of individual differences (Dubois and Adolphs, 2016) with clinical relevance (Fox, 2010). For

these reasons, we think that in many studies the gains in reliability offered by GFC will outweigh

potential task-induced "heterogeneity."

Recommendations

Based on the above summary, we generally recommend that researchers seeking to maximize the

reliability and generalizable predictive utility of GFC should aggregate as much data as possible,

across all rest and task data, that they have at their disposal. Our results further suggest that with a

minimum of 20 minutes of GFC data, researchers can achieve good reliability in some networks

of broad interest (e.g., default, frontoparietal and dorsal attention networks). Furthermore, a strict

balance of the amount of data from each task and resting-state scan is not necessary to achieve the

benefits of GFC. This is demonstrated by the increased reliability of GFC despite unequal task

lengths in the Dunedin Study dataset and in the HCP dataset by continued improvement in

reliability despite timepoints being selected at random from the remaining data after reaching a

minimum of 25 minutes of data. However, we urge caution in applying GFC to datasets that are

heavily skewed (> 50%) towards resting-state or one specific task. Although we expect that

deriving GFC from this type of dataset will boost reliability, the generalizability of these findings

to GFC derived from other datasets may be limited because of the overrepresentation of a specific

type of data. Lastly, our results suggest that researchers developing new protocols may benefit

from collecting multiple short task scans rather than a single longer task scan (supplemental tables

S1 and S5).

Conclusion

Here we propose GFC as a method for improving the reliability of estimating individual

differences in the intrinsic architecture of functional networks. We demonstrate that, when the

amount of data available for analysis is held constant, the combination of resting-state and task

data is at least as reliable as resting-state alone. Additionally, when the same amount of data is

available, GFC produces higher heritability estimates than RSFC. Many researchers who have less

than 25 minutes of resting-state data but have additionally collected task data on the same

participants can immediately boost reliability, heritability, and power by adopting GFC as a

measure of intrinsic connectivity when studying individual differences. Our findings also suggest

that future data collection should consider naturalistic fMRI (Hasson et al., 2010, 2004; Huth et

al., 2016; Lahnakoski et al., 2012; Vanderwal et al., 2017) and engaging tasks in addition to

resting-state scans when generating data for estimating intrinsic connectivity. This may be

especially true when studying children, elderly, or mentally ill participants, as these groups often

cannot lie still for 25 minutes or more of resting-state scanning (Power et al., 2012; Satterthwaite

et al., 2013; Yuan et al., 2009). Our current findings also suggest that researchers need not think

of collecting resting-state and task data as a zero-sum competition for scan time when designing

studies. Instead, our findings demonstrate that task and resting-state data provide complementary

measures that together can lead to more reliable, heritable measurement of individual differences

in intrinsic connectivity. We think that collecting adequate amounts (i.e., > 25 minutes) of high-

quality fMRI data should be emphasized over the traditional task-rest dichotomy, when individual

differences are of interest. If fMRI is going to be a part of a cumulative neuroscience of individual

differences with clinical relevance, psychometric measurement quality, especially reliability, must

be at the forefront of study design.

Acknowledgement. The Dunedin Multidisciplinary Health and Development Study is supported

by the NZ HRC and NZ MBIE. This research was supported by National Institute on Aging

grant R01AG032282, R01AG049789, and UK Medical Research Council grant MR/P005918/1.

Additional support was provided by the Jacobs Foundation. MLE is supported by the National

Science Foundation Graduate Research Fellowship under Grant No. NSF DGE-1644868. Thank

you to members of the Advisory Board for the Dunedin Neuroimaging Study. We thank the

Dunedin Study members, unit research staff, and Study founder Phil Silva. The study protocol

was approved by the institutional ethical review boards of the participating universities. Study

members gave informed consent before participating. The authors declare no competing financial

interests.

Author Contributions. Conceptualization, M.L.E.; Methodology, Software and Formal

Analysis, M.L.E., A.R.K and M.C.; Data Curation, M.L.E., A.R.K., A.C., and T.E.M.;

Resources, T.R.M., R.K., S.R., R.P., A.C., T.E.M. and A.R.H., Writing, M.L.E., A.R.K., M.J.K.,

M.C., T.R.M., R.K., D.I., S.R., R.P., A.C., T.E.M. and A.R.H.; Supervision, A.C., T.E.M. and

A.R.H.; Funding Acquisition, R.P., A.C., T.E.M. and A.R.H.

References

Aarts, A.A., Anderson, J.E., Anderson, C.J., Attridge, P.R., Attwood, A., Axt, J., Babel, M., Bahník, Š., Baranski, E., Barnett-Cowan, M., Bartmess, E., Beer, J., Bell, R., Bentley, H., Beyan, L., Binion, G., Borsboom, D., Bosch, A., Bosco, F.A., Bowman, S.D., Brandt, M.J., Braswell, E., Brohmer, H., Brown, B.T., Brown, K., Brüning, J., Calhoun-Sauls, A., Callahan, S.P., Chagnon, E., Chandler, J., Chartier, C.R., Cheung, F., Christopherson, C.D., Cillessen, L., Clay, R., Cleary, H., Cloud, M.D., Conn, M., Cohoon, J., Columbus, S., Cordes, A., Costantini, G., Alvarez, L.D.C., Cremata, E., Crusius, J., DeCoster, J., DeGaetano, M.A., Penna, N.D., Den Bezemer, B., Deserno, M.K., Devitt, O., Dewitte, L., Dobolyi, D.G., Dodson, G.T., Donnellan, M.B., Donohue, R., Dore, R.A., Dorrough, A., Dreber, A., Dugas, M., Dunn, E.W., Easey, K., Eboigbe, S., Eggleston, C., Embley, J., Epskamp, S., Errington, T.M., Estel, V., Farach, F.J., Feather, J., Fedor, A., Fernández-Castilla, B., Fiedler, S., Field, J.G., Fitneva, S.A., Flagan, T., Forest, A.L., Forsell, E., Foster, J.D., Frank, M.C., Frazier, R.S., Fuchs, H., Gable, P., Galak, J., Galliani, E.M., Gampa, A., Garcia, S., Gazarian, D., Gilbert, E., Giner-Sorolla, R., Glöckner, A., Goellner, L., Goh, J.X., Goldberg, R., Goodbourn, P.T., Gordon-McKeon, S., Gorges, B., Gorges, J., Goss, J., Graham, J., Grange, J.A., Gray, J., Hartgerink, C., Hartshorne, J., Hasselman, F., Hayes, T., Heikensten, E., Henninger, F., Hodsoll, J., Holubar, T., Hoogendoorn, G., Humphries, D.J., Hung, C.O.Y., Immelman, N., Irsik, V.C., Jahn, G., Jäkel, F., Jekel, M., Johannesson, M., Johnson, L.G., Johnson, D.J., Johnson, K.M., Johnston, W.J., Jonas, K., Joy-Gaba, J.A., Kappes, H.B., Kelso, K., Kidwell, M.C., Kim, S.K., Kirkhart, M., Kleinberg, B., Knežević, G., Kolorz, F.M., Kossakowski, J.J., Krause, R.W., Krijnen, J., Kuhlmann, T., Kunkels, Y.K., Kyc, M.M., Lai, C.K., Laique, A., Lakens, D., Lane, K.A., Lassetter, B., Lazarević, L.B., Le Bel, E.P., Lee, K.J., Lee, M., Lemm, K., Levitan, C.A., Lewis, M., Lin, L., Lin, S., Lippold, M., Loureiro, D., Luteijn, I., MacKinnon, S., Mainard, H.N., Marigold, D.C., Martin, D.P., Martinez, T., Masicampo, E.J., Matacotta, J., Mathur, M., May, M., Mechin, N., Mehta, P., Meixner, J., Melinger, A., Miller, J.K., Miller, M., Moore, K., Möschl, M., Motyl, M., Müller, S.M., Munafo, M., Neijenhuijs, K.I., Nervi, T., Nicolas, G., Nilsonne, G., Nosek, B.A., Nuijten, M.B., Olsson, C., Osborne, C., Ostkamp, L., Pavel, M., Penton-Voak, I.S., Perna, O., Pernet, C., Perugini, M., Pipitone, R.N., Pitts, M., Plessow, F., Prenoveau, J.M., Rahal, R.M., Ratliff, K.A., Reinhard, D., Renkewitz, F., Ricker, A.A., Rigney, A., Rivers, A.M., Roebke, M., Rutchick, A.M., Ryan, R.S., Sahin, O., Saide, A., Sandstrom, G.M., Santos, D., Saxe, R., Schlegelmilch, R., Schmidt, K., Scholz, S., Seibel, L., Selterman, D.F., Shaki, S., Simpson, W.B., Sinclair, H.C., Skorinko, J.L.M., Slowik, A., Snyder, J.S., Soderberg, C., Sonnleitner, C., Spencer, N., Spies, J.R., Steegen, S., Stieger, S., Strohminger, N., Sullivan, G.B., Talhelm, T., Tapia, M., Te Dorsthorst, A., Thomae, M., Thomas, S.L., Tio, P., Traets, F., Tsang, S., Tuerlinckx, F., Turchan, P., Valášek, M., Van't Veer, A.E., Van Aert, R., Van Assen, M., Van Bork, R., Van De Ven, M., Van Den Bergh, D., Van Der Hulst, M., Van Dooren, R., Van Doorn, J., Van Renswoude, D.R., Van Rijn, H., Vanpaemel, W., Echeverría, A.V., Vazquez, M., Velez, N., Vermue, M., Verschoor, M., Vianello, M., Voracek, M., Vuu, G., Wagenmakers, E.J., Weerdmeester, J., Welsh, A., Westgate, E.C., Wissink, J., Wood, M., Woods, A., Wright, E., Wu, S., Zeelenberg, M., Zuni, K., 2015. Estimating the reproducibility of psychological science. Science (80-.). 349. https://doi.org/10.1126/science.aac4716

Adams, H.H.H., Hibar, D.P., Chouraki, V., Stein, J.L., Nyquist, P.A., Rentería, M.E., Trompet, S., Arias-Vasquez, A., Seshadri, S., Desrivières, S., Beecham, A.H., Jahanshad, N., Wittfeld, K., Van Der Lee, S.J., Abramovic, L., Alhusaini, S., Amin, N., Andersson, M.,

Arfanakis, K., Aribisala, B.S., Armstrong, N.J., Athanasiu, L., Axelsson, T., Beiser, A.,
Bernard, M., Bis, J.C., Blanken, L.M.E., Blanton, S.H., Bohlken, M.M., Boks, M.P.,
Bralten, J., Brickman, A.M., Carmichael, O., Chakravarty, M.M., Chauhan, G., Chen, Q.,
Ching, C.R.K., Cuellar-Partida, G., Braber, A. Den, Doan, N.T., Ehrlich, S., Filippi, I., Ge,
T., Giddaluru, S., Goldman, A.L., Gottesman, R.F., Greven, C.U., Grimm, O., Griswold,
M.E., Guadalupe, T., Hass, J., Haukvik, U.K., Hilal, S., Hofer, E., Hoehn, D., Holmes, A.J.,
Hoogman, M., Janowitz, D., Jia, T., Kasperaviciute, D., Kim, S., Klein, M., Kraemer, B.,
Lee, P.H., Liao, J., Liewald, D.C.M., Lopez, L.M., Luciano, M., Macare, C., Marquand, A.,
Matarin, M., Mather, K.A., Mattheisen, M., Mazoyer, B., McKay, D.R., McWhirter, R.,
Milaneschi, Y., Mirza-Schreiber, N., Muetzel, R.L., Maniega, S.M., Nho, K., Nugent, A.C.,
Loohuis, L.M.O., Oosterlaan, J., Papmeyer, M., Pappa, I., Pirpamer, L., Pudas, S., Pütz, B.,
Rajan, K.B., Ramasamy, A., Richards, J.S., Risacher, S.L., Roiz-Santiañez, R., Rommelse,
N., Rose, E.J., Royle, N.A., Rundek, T., Sämann, P.G., Satizabal, C.L., Schmaal, L.,
Schork, A.J., Shen, L., Shin, J., Shumskaya, E., Smith, A. V., Sprooten, E., Strike, L.T.,
Teumer, A., Thomson, R., Tordesillas-Gutierrez, D., Toro, R., Trabzuni, D., Vaidya, D.,
Van Der Grond, J., Van Der Meer, D., Van Donkelaar, M.M.J., Van Eijk, K.R., Van Erp,
T.G.M., Van Rooij, D., Walton, E., Westlye, L.T., Whelan, C.D., Windham, B.G., Winkler,
A.M., Woldehawariat, G., Wolf, C., Wolfers, T., Xu, B., Yanek, L.R., Yang, J., Zijdenbos,
A., Zwiers, M.P., Agartz, I., Aggarwal, N.T., Almasy, L., Ames, D., Amouyel, P.,
Andreassen, O.A., Arepalli, S., Assareh, A.A., Barral, S., Bastin, M.E., Becker, D.M.,
Becker, J.T., Bennett, D.A., Blangero, J., Van Bokhoven, H., Boomsma, D.I., Brodaty, H.,
Brouwer, R.M., Brunner, H.G., Buckner, R.L., Buitelaar, J.K., Bulayeva, K.B., Cahn, W.,
Calhoun, V.D., Cannon, D.M., Cavalleri, G.L., Chen, C., Cheng, C.Y., Cichon, S.,
Cookson, M.R., Corvin, A., Crespo-Facorro, B., Curran, J.E., Czisch, M., Dale, A.M.,
Davies, G.E., De Geus, E.J.C., De Jager, P.L., De Zubicaray, G.I., Delanty, N., Depondt, C.,
Destefano, A.L., Dillman, A., Djurovic, S., Donohoe, G., Drevets, W.C., Duggirala, R.,
Dyer, T.D., Erk, S., Espeseth, T., Evans, D.A., Fedko, I.O., Fernández, G., Ferrucci, L.,
Fisher, S.E., Fleischman, D.A., Ford, I., Foroud, T.M., Fox, P.T., Francks, C., Fukunaga,
M., Gibbs, J.R., Glahn, D.C., Gollub, R.L., Göring, H.H.H., Grabe, H.J., Green, R.C.,
Gruber, O., Gudnason, V., Guelfi, S., Hansell, N.K., Hardy, J., Hartman, C.A., Hashimoto,
R., Hegenscheid, K., Heinz, A., Le Hellard, S., Hernandez, D.G., Heslenfeld, D.J., Ho,
B.C., Hoekstra, P.J., Hoffmann, W., Hofman, A., Holsboer, F., Homuth, G., Hosten, N.,
Hottenga, J.J., Pol, H.E.H., Ikeda, M., Ikram, M.K., Jack, C.R., Jenkinson, M., Johnson, R.,
Jönsson, E.G., Jukema, J.W., Kahn, R.S., Kanai, R., Kloszewska, I., Knopman, D.S.,
Kochunov, P., Kwok, J.B., Lawrie, S.M., Lemaître, H., Liu, X., Longo, D.L., Longstreth,
W.T., Lopez, O.L., Lovestone, S., Martinez, O., Martinot, J.L., Mattay, V.S., McDonald, C.,
McIntosh, A.M., McMahon, K.L., McMahon, F.J., Mecocci, P., Melle, I., Meyer-
Lindenberg, A., Mohnke, S., Montgomery, G.W., Morris, D.W., Mosley, T.H., Mühleisen,
T.W., Müller-Myhsok, B., Nalls, M.A., Nauck, M., Nichols, T.E., Niessen, W.J., Nöthen,
M.M., Nyberg, L., Ohi, K., Olvera, R.L., Ophoff, R.A., Pandolfo, M., Paus, T., Pausova, Z.,
Penninx, B.W.J.H., Pike, G.B., Potkin, S.G., Psaty, B.M., Reppermund, S., Rietschel, M.,
Roffman, J.L., Romanczuk-Seiferth, N., Rotter, J.I., Ryten, M., Sacco, R.L., Sachdev, P.S.,
Saykin, A.J., Schmidt, R., Schofield, P.R., Sigurdsson, S., Simmons, A., Singleton, A.,
Sisodiya, S.M., Smith, C., Smoller, J.W., Soininen, H., Srikanth, V., Steen, V.M., Stott,
D.J., Sussmann, J.E., Thalamuthu, A., Tiemeier, H., Toga, A.W., Traynor, B.J., Troncoso,
J., Turner, J.A., Tzourio, C., Uitterlinden, A.G., Hernández, M.C.V., Van Der Brug, M.,
Van Der Lugt, A., Van Der Wee, N.J.A., Van Duijn, C.M., Van Haren, N.E.M., Van't Ent,
D., Van Tol, M.J., Vardarajan, B.N., Veltman, D.J., Vernooij, M.W., Völzke, H., Walter,

H., Wardlaw, J.M., Wassink, T.H., Weale, M.E., Weinberger, D.R., Weiner, M.W., Wen, W., Westman, E., White, T., Wong, T.Y., Wright, C.B., Zielke, H.R., Zonderman, A.B., Deary, I.J., Decarli, C., Schmidt, H., Martin, N.G., De Craen, A.J.M., Wright, M.J., Launer, L.J., Schumann, G., Fornage, M., Franke, B., Debette, S., Medland, S.E., Ikram, M.A., Thompson, P.M., 2016. Novel genetic loci underlying human intracranial volume identified through genome-wide association. Nat. Neurosci. 19, 1569–1582. https://doi.org/10.1038/nn.4398

Adhikari, B.M., Jahanshad, N., Shukla, D., Glahn, D.C., Blangero, J., Fox, P.T., Reynolds, R.C., Cox, R.W., Fieremans, E., Veraart, J., Novikov, D.S., Nichols, T.E., Hong, L.E., Thompson, P.M., Kochunov, P., 2018. Comparison of heritability estimates on resting state fMRI connectivity phenotypes using the ENIGMA analysis pipeline. Hum. Brain Mapp. https://doi.org/10.1002/hbm.24331

Anderson, J.S., Ferguson, M.A., Lopez-Larson, M., Yurgelun-Todd, D., 2011. Reproducibility of single-subject functional connectivity measurements. Am. J. Neuroradiol. 32, 548–555. https://doi.org/10.3174/ajnr.A2330

Arfanakis, K., Cordes, D., Haughton, V.M., Moritz, C.H., Quigley, M.A., Meyerand, M.E., 2000. Combining independent component analysis and correlation analysis to probe interregional connectivity in fMRI task activation datasets. Magn. Reson. Imaging 18, 921–930. https://doi.org/10.1016/S0730-725X(00)00190-9

Avants, B.B., Epstein, C.L., Grossman, M., Gee, J.C., 2008. Symmetric diffeomorphic image registration with cross-correlation: Evaluating automated labeling of elderly and neurodegenerative brain. Med. Image Anal. 12, 26–41. https://doi.org/10.1016/j.media.2007.06.004

Barch, D.M., Burgess, G.C., Harms, M.P., Petersen, S.E., Schlaggar, B.L., Corbetta, M., Glasser, M.F., Curtiss, S., Dixit, S., Feldt, C., Nolan, D., Bryant, E., Hartley, T., Footer, O., Bjork, J.M., Poldrack, R., Smith, S., Johansen-Berg, H., Snyder, A.Z., Van Essen, D.C., 2013. Function in the human connectome: Task-fMRI and individual differences in behavior. Neuroimage 80, 169–189. https://doi.org/10.1016/j.neuroimage.2013.05.033

Behzadi, Y., Restom, K., Liau, J., Liu, T.T., 2007. A component based noise correction method (CompCor) for BOLD and perfusion based fMRI. Neuroimage 37, 90–101. https://doi.org/10.1016/j.neuroimage.2007.04.042

Benjamini, Y., Hochberg, Y., 1995. Controlling the false discovery rate: a practical and powerful approach to multiple testing. J. R. Stat. Soc. https://doi.org/10.2307/2346101

Betti, V., DellaPenna, S., de Pasquale, F., Mantini, D., Marzetti, L., Romani, G.L., Corbetta, M., 2013. Natural scenes viewing alters the dynamics of functional connectivity in the human brain. Neuron 79, 782–797. https://doi.org/10.1016/j.neuron.2013.06.022

Birn, R.M., Molloy, E.K., Patriat, R., Parker, T., Meier, T.B., Kirk, G.R., Nair, V.A., Meyerand, M.E., Prabhakaran, V., 2013. The effect of scan length on the reliability of resting-state fMRI connectivity estimates. Neuroimage 83, 550–558. https://doi.org/10.1016/j.neuroimage.2013.05.099

Boker, S., Neale, M., Maes, H., Wilde, M., Spiegel, M., Brick, T., Spies, J., Estabrook, R., Kenny, S., Bates, T., Mehta, P., Fox, J., 2011. OpenMx: An Open Source Extended Structural Equation Modeling Framework. Psychometrika 76, 306–317. https://doi.org/10.1007/s11336-010-9200-6

Braga, R.M., Buckner, R.L., 2017. Parallel Interdigitated Distributed Networks within the Individual Estimated by Intrinsic Functional Connectivity. Neuron 95, 457–471.e5. https://doi.org/10.1016/j.neuron.2017.06.038

Braskie, M.N., Ringman, J.M., Thompson, P.M., 2011. Neuroimaging measures as

1006   endophenotypes in Alzheimer's disease. Int. J. Alzheimers. Dis.
1007   https://doi.org/10.4061/2011/490140
1008 Bright, M.G., Tench, C.R., Murphy, K., 2017. Potential pitfalls when denoising resting state
1009   fMRI data using nuisance regression. Neuroimage 154, 159–168.
1010   https://doi.org/10.1016/j.neuroimage.2016.12.027
1011 Buckner, R.L., Andrews-Hanna, J.R., Schacter, D.L., 2008. The Brain's Default Network. Ann.
1012   N. Y. Acad. Sci. 1124, 1–38. https://doi.org/10.1196/annals.1440.011
1013 Buckner, R.L., Krienen, F.M., Yeo, B.T.T., 2013. Opportunities and limitations of intrinsic
1014   functional connectivity MRI. Nat. Neurosci. https://doi.org/10.1038/nn.3423
1015 Burgess, G.C., Kandala, S., Nolan, D., Laumann, T.O., Power, J.D., Adeyemo, B., Harms, M.P.,
1016   Petersen, S.E., Barch, D.M., 2016. Evaluation of Denoising Strategies to Address Motion-
1017   Correlated Artifacts in Resting-State Functional Magnetic Resonance Imaging Data from
1018   the Human Connectome Project. Brain Connect. 6, 669–680.
1019   https://doi.org/10.1089/brain.2016.0435
1020 Button, K.S., Ioannidis, J.P.A., Mokrysz, C., Nosek, B.A., Flint, J., Robinson, E.S.J., Munafò,
1021   M.R., 2013. Power failure: Why small sample size undermines the reliability of
1022   neuroscience. Nat. Rev. Neurosci. 14, 365–376. https://doi.org/10.1038/nrn3475
1023 Bzdok, D., Varoquaux, G., Grisel, O., Eickenberg, M., Poupon, C., Thirion, B., 2016. Formal
1024   Models of the Network Co-occurrence Underlying Mental Operations. PLoS Comput. Biol.
1025   12. https://doi.org/10.1371/journal.pcbi.1004994
1026 Bzdok, D., Yeo, B.T.T., 2017. Inference in the age of big data: Future perspectives on
1027   neuroscience. Neuroimage 155, 549–564. https://doi.org/10.1016/j.neuroimage.2017.04.061
1028 Caballero-Gaudes, C., Reynolds, R.C., 2017. Methods for cleaning the BOLD fMRI signal.
1029   Neuroimage 154, 128–149. https://doi.org/10.1016/j.neuroimage.2016.12.018
1030 Caria, A., Sitaram, R., Birbaumer, N., 2012. Real-time fMRI: A tool for local brain regulation.
1031   Neuroscientist. https://doi.org/10.1177/1073858411407205
1032 Casey, B.J., Cannonier, T., Conley, M.I., Cohen, A.O., Barch, D.M., Heitzeg, M.M., Soules,
1033   M.E., Teslovich, T., Dellarco, D. V., Garavan, H., Orr, C.A., Wager, T.D., Banich, M.T.,
1034   Speer, N.K., Sutherland, M.T., Riedel, M.C., Dick, A.S., Bjork, J.M., Thomas, K.M.,
1035   Chaarani, B., Mejia, M.H., Hagler, D.J., Daniela Cornejo, M., Sicat, C.S., Harms, M.P.,
1036   Dosenbach, N.U.F., Rosenberg, M., Earl, E., Bartsch, H., Watts, R., Polimeni, J.R.,
1037   Kuperman, J.M., Fair, D.A., Dale, A.M., 2018. The Adolescent Brain Cognitive
1038   Development (ABCD) study: Imaging acquisition across 21 sites. Dev. Cogn. Neurosci.
1039   https://doi.org/10.1016/j.dcn.2018.03.001
1040 Chen, G., Taylor, P.A., Haller, S.P., Kircanski, K., Stoddard, J., Pine, D.S., Leibenluft, E.,
1041   Brotman, M.A., Cox, R.W., 2018. Intraclass correlation: Improved modeling approaches
1042   and applications for neuroimaging. Hum. Brain Mapp. 39, 1187–1206.
1043   https://doi.org/10.1002/hbm.23909
1044 Christoff, K., Gordon, A.M., Smallwood, J., Smith, R., Schooler, J.W., 2009. Experience
1045   sampling during fMRI reveals default network and executive system contributions to mind
1046   wandering. Proc. Natl. Acad. Sci. 106, 8719–8724.
1047   https://doi.org/10.1073/pnas.0900234106
1048 Cicchetti, D. V., 1994. Guidlines, Criteria, and Rules of Thumb for Evalauting Normed and
1049   Standardized Assessment Instruments in Psychology. Psychol. Assess. 6, 284–290.
1050   https://doi.org/10.1037/1040-3590.6.4.284
1051 Cole, M.W., Anticevic, A., Repovs, G., Barch, D., 2011. Variable global dysconnectivity and
1052   individual differences in schizophrenia. Biol. Psychiatry 70, 43–50.
1053   https://doi.org/10.1016/j.biopsych.2011.02.010

Cole, M.W., Bassett, D.S., Power, J.D., Braver, T.S., Petersen, S.E., 2014. Intrinsic and task-evoked network architectures of the human brain. Neuron 83, 238–251. https://doi.org/10.1016/j.neuron.2014.05.014

Cole, M.W., Ito, T., Bassett, D.S., Schultz, D.H., 2016. Activity flow over resting-state networks shapes cognitive task activations. Nat. Neurosci. 19, 1718–1726. https://doi.org/10.1038/nn.4406

Collins, F.S., Varmus, H., 2015. A New Initiative on Precision Medicine. N. Engl. J. Med. 372, 793–795. https://doi.org/10.1056/NEJMp1500523

Cox, R.W., 1996. AFNI: software for analysis and visualization of functional magnetic resonance neuroimages. Comput. Biomed. Res. 29, 162–73. https://doi.org/10.1006/cbmr.1996.0014

Cremers, H.R., Wager, T.D., Yarkoni, T., 2017. The relation between statistical power and inference in fMRI 1–20. https://doi.org/10.1371/journal.pone.0184923

Deco, G., Hagmann, P., Hudetz, A.G., Tononi, G., 2014. Modeling resting-state functional networks when the cortex falls asleep: Local and global changes. Cereb. Cortex 24, 3180–3194. https://doi.org/10.1093/cercor/bht176

Dong, D., Wang, Y., Chang, X., Luo, C., Yao, D., 2018. Dysfunction of Large-Scale Brain Networks in Schizophrenia: A Meta-analysis of Resting-State Functional Connectivity. Schizophr. Bull. 44, 168–181. https://doi.org/10.1093/schbul/sbx034

Dubois, J., Adolphs, R., 2016. Building a Science of Individual Differences from fMRI. Trends Cogn. Sci. 20, 425–443. https://doi.org/10.1016/j.tics.2016.03.014

Dubois, J., Galdi, P., Han, Y., Paul, L.K., Adolphs, R., Angeles, L., Systems, I., Dubois, J., 2018. Resting-state functional brain connectivity best predicts the personality dimension of openness to experience. bioRxiv 215129. https://doi.org/10.1101/215129

Elliott, L., Sharp, K., Alfaro-Almagro, F., Douaud, G., Miller, K., Marchini, J., Smith, S., 2017. The genetic basis of human brain structure and function: 1,262 genome-wide associations found from 3,144 GWAS of multimodal brain imaging phenotypes from 9,707 UK Biobank participants. bioRxiv.

Elliott, M.L., Romer, A., Knodt, A.R., Hariri, A.R., 2018. A Connectome-wide Functional Signature of Transdiagnostic Risk for Mental Illness. Biol. Psychiatry 84, 452–459. https://doi.org/10.1016/j.biopsych.2018.03.012

Errington, T.M., Iorns, E., Gunn, W., Tan, F.E. lisabet., Lomax, J., Nosek, B.A., 2014. An open investigation of the reproducibility of cancer biology research. Elife 3. https://doi.org/10.7554/eLife.04333

Fair, D.A., Schlaggar, B.L., Cohen, A.L., Miezin, F.M., Dosenbach, N.U.F., Wenger, K.K., Fox, M.D., Snyder, A.Z., Raichle, M.E., Petersen, S.E., 2007. A method for using blocked and event-related fMRI data to study "resting state" functional connectivity. Neuroimage 35, 396–405. https://doi.org/10.1016/j.neuroimage.2006.11.051

Finn, E.S., Scheinost, D., Finn, D.M., Shen, X., Papademetris, X., Constable, R.T., 2017. Can brain state be manipulated to emphasize individual differences in functional connectivity? Neuroimage 160, 140–151. https://doi.org/10.1016/j.neuroimage.2017.03.064

Finn, E.S., Shen, X., Scheinost, D., Rosenberg, M.D., Huang, J., Chun, M.M., Papademetris, X., Constable, R.T., 2015. Functional connectome fingerprinting: Identifying individuals using patterns of brain connectivity. Nat. Neurosci. 18, 1664–1671. https://doi.org/10.1038/nn.4135

Fox, M.D., 2010. Clinical applications of resting state functional connectivity. Front. Syst. Neurosci. 4. https://doi.org/10.3389/fnsys.2010.00019

Fox, M.D., Buckner, R.L., Liu, H., Chakravarty, M.M., Lozano, A.M., Pascual-Leone, A., 2014.

Resting-state networks link invasive and noninvasive brain stimulation across diverse
psychiatric and neurological diseases. Proc. Natl. Acad. Sci.
https://doi.org/10.1073/pnas.1405003111
Fox, M.D., Snyder, A.Z., Vincent, J.L., Raichle, M.E., 2007. Intrinsic Fluctuations within
Cortical Systems Account for Intertrial Variability in Human Behavior. Neuron 56, 171–
184. https://doi.org/10.1016/j.neuron.2007.08.023
Fox, M.D., Snyder, A.Z., Zacks, J.M., Raichle, M.E., 2006. Coherent spontaneous activity
accounts for trial-to-trial variability in human evoked brain responses. Nat. Neurosci. 9, 23–
25. https://doi.org/10.1038/nn1616
Ge, T., Holmes, A.J., Buckner, R.L., Smoller, J.W., Sabuncu, M.R., 2017. Heritability analysis
with repeat measurements and its application to resting-state functional connectivity. Proc.
Natl. Acad. Sci. 114, 5521–5526. https://doi.org/10.1073/pnas.1700765114
Geerligs, L., Rubinov, M., Cam-CAN, Henson, R.N., 2015. State and Trait Components of
Functional Connectivity: Individual Differences Vary with Mental State. J. Neurosci. 35,
13949–13961. https://doi.org/10.1523/JNEUROSCI.1324-15.2015
Glahn, D.C., Winkler, A.M., Kochunov, P., Almasy, L., Duggirala, R., Carless, M.A., Curran,
J.C., Olvera, R.L., Laird, A.R., Smith, S.M., Beckmann, C.F., Fox, P.T., Blangero, J., 2010.
Genetic control over the resting brain. Proc Natl Acad Sci U S A 107, 1223–1228.
https://doi.org/10.1073/pnas.0909969107
Glasser, M.F., Sotiropoulos, S.N., Wilson, J.A., Coalson, T.S., Fischl, B., Andersson, J.L., Xu,
J., Jbabdi, S., Webster, M., Polimeni, J.R., Van Essen, D.C., Jenkinson, M., 2013. The
minimal preprocessing pipelines for the Human Connectome Project. Neuroimage 80, 105–
124. https://doi.org/10.1016/j.neuroimage.2013.04.127
Gordon, E.M., Laumann, T.O., Gilmore, A.W., Newbold, D.J., Greene, D.J., Berg, J.J., Ortega,
M., Hoyt-Drazen, C., Gratton, C., Sun, H., Hampton, J.M., Coalson, R.S., Nguyen, A.L.,
McDermott, K.B., Shimony, J.S., Snyder, A.Z., Schlaggar, B.L., Petersen, S.E., Nelson,
S.M., Dosenbach, N.U.F., 2017. Precision Functional Mapping of Individual Human Brains.
Neuron 95, 791–807.e7. https://doi.org/10.1016/j.neuron.2017.07.011
Gratton, C., Laumann, T.O., Nielsen, A.N., Greene, D.J., Gordon, E.M., Gilmore, A.W., Nelson,
S.M., Coalson, R.S., Snyder, A.Z., Schlaggar, B.L., Dosenbach, N.U.F., Petersen, S.E.,
2018. Functional Brain Networks Are Dominated by Stable Group and Individual Factors,
Not Cognitive or Daily Variation. Neuron 439–452.
https://doi.org/10.1016/j.neuron.2018.03.035
Greene, A.S., Gao, S., Scheinost, D., Constable, R.T., 2018. Task-induced brain state
manipulation improves prediction of individual traits. Nat. Commun. 9.
https://doi.org/10.1038/s41467-018-04920-3
Greicius, M., 2008. Resting-state functional connectivity in neuropsychiatric disorders. Curr.
Opin. Neurol. 21, 424–30. https://doi.org/10.1097/WCO.0b013e328306f2c5
Greve, D.N., Fischl, B., 2009. Accurate and robust brain image alignment using boundary-based
registration. Neuroimage 48, 63–72. https://doi.org/10.1016/j.neuroimage.2009.06.060
Guo, C.C., Kurth, F., Zhou, J., Mayer, E.A., Eickhoff, S.B., Kramer, J.H., Seeley, W.W., 2012.
One-year test-retest reliability of intrinsic connectivity network fMRI in older adults.
Neuroimage 61, 1471–1483. https://doi.org/10.1016/j.neuroimage.2012.03.027
Hacker, C.D., Laumann, T.O., Szrama, N.P., Baldassarre, A., Snyder, A.Z., Leuthardt, E.C.,
Corbetta, M., 2013. Resting state network estimation in individual subjects. Neuroimage 82,
616–633. https://doi.org/10.1016/j.neuroimage.2013.05.108
Hamburg, M.A., Collins, F.S., 2010. The Path to Personalized Medicine - Perspective. N. Engl.
J. Med. 363, 301–304. https://doi.org/10.1056/NEJMp1006304

Hariri, A.R., Weinberger, D.R., 2003. Imaging genomics. Br. Med. Bull. https://doi.org/10.1093/bmb/65.1.259

Hasler, G., Northoff, G., 2011. Discovering imaging endophenotypes for major depression. Mol. Psychiatry. https://doi.org/10.1038/mp.2011.23

Hasson, U., Malach, R., Heeger, D.J., 2010. Reliability of cortical activity during natural stimulation. Trends Cogn. Sci. https://doi.org/10.1016/j.tics.2009.10.011

Hasson, U., Nir, Y., Levy, I., Fuhrmann, G., Malach, R., 2004. Intersubject Synchronization of Cortical Activity during Natural Vision. Science (80-.). 303, 1634–1640. https://doi.org/10.1126/science.1089506

Hibar, D.P., Adams, H.H.H., Jahanshad, N., Chauhan, G., Stein, J.L., Hofer, E., Renteria, M.E., Bis, J.C., Arias-Vasquez, A., Ikram, M.K., Desrivières, S., Vernooij, M.W., Abramovic, L., Alhusaini, S., Amin, N., Andersson, M., Arfanakis, K., Aribisala, B.S., Armstrong, N.J., Athanasiu, L., Axelsson, T., Beecham, A.H., Beiser, A., Bernard, M., Blanton, S.H., Bohlken, M.M., Boks, M.P., Bralten, J., Brickman, A.M., Carmichael, O., Chakravarty, M.M., Chen, Q., Ching, C.R.K., Chouraki, V., Cuellar-Partida, G., Crivello, F., Den Braber, A., Doan, N.T., Ehrlich, S., Giddaluru, S., Goldman, A.L., Gottesman, R.F., Grimm, O., Griswold, M.E., Guadalupe, T., Gutman, B.A., Hass, J., Haukvik, U.K., Hoehn, D., Holmes, A.J., Hoogman, M., Janowitz, D., Jia, T., Jørgensen, K.N., Karbalai, N., Kasperaviciute, D., Kim, S., Klein, M., Kraemer, B., Lee, P.H., Liewald, D.C.M., Lopez, L.M., Luciano, M., MacAre, C., Marquand, A.F., Matarin, M., Mather, K.A., Mattheisen, M., McKay, D.R., Milaneschi, Y., Muñoz Maniega, S., Nho, K., Nugent, A.C., Nyquist, P., Loohuis, L.M.O., Oosterlaan, J., Papmeyer, M., Pirpamer, L., Pütz, B., Ramasamy, A., Richards, J.S., Risacher, S.L., Roiz-Santiañez, R., Rommelse, N., Ropele, S., Rose, E.J., Royle, N.A., Rundek, T., Sämann, P.G., Saremi, A., Satizabal, C.L., Schmaal, L., Schork, A.J., Shen, L., Shin, J., Shumskaya, E., Smith, A. V., Sprooten, E., Strike, L.T., Teumer, A., Tordesillas-Gutierrez, D., Toro, R., Trabzuni, D., Trompet, S., Vaidya, D., Van Der Grond, J., Van Der Lee, S.J., Van Der Meer, D., Van Donkelaar, M.M.J., Van Eijk, K.R., Van Erp, T.G.M., Van Rooij, D., Walton, E., Westlye, L.T., Whelan, C.D., Windham, B.G., Winkler, A.M., Wittfeld, K., Woldehawariat, G., Wolf, C., Wolfers, T., Yanek, L.R., Yang, J., Zijdenbos, A., Zwiers, M.P., Agartz, I., Almasy, L., Ames, D., Amouyel, P., Andreassen, O.A., Arepalli, S., Assareh, A.A., Barral, S., Bastin, M.E., Becker, D.M., Becker, J.T., Bennett, D.A., Blangero, J., Van Bokhoven, H., Boomsma, D.I., Brodaty, H., Brouwer, R.M., Brunner, H.G., Buckner, R.L., Buitelaar, J.K., Bulayeva, K.B., Cahn, W., Calhoun, V.D., Cannon, D.M., Cavalleri, G.L., Cheng, C.Y., Cichon, S., Cookson, M.R., Corvin, A., Crespo-Facorro, B., Curran, J.E., Czisch, M., Dale, A.M., Davies, G.E., De Craen, A.J.M., De Geus, E.J.C., De Jager, P.L., De Zubicaray, G.I., Deary, I.J., Debette, S., Decarli, C., Delanty, N., Depondt, C., Destefano, A., Dillman, A., Djurovic, S., Donohoe, G., Drevets, W.C., Duggirala, R., Dyer, T.D., Enzinger, C., Erk, S., Espeseth, T., Fedko, I.O., Fernández, G., Ferrucci, L., Fisher, S.E., Fleischman, D.A., Ford, I., Fornage, M., Foroud, T.M., Fox, P.T., Francks, C., Fukunaga, M., Gibbs, J.R., Glahn, D.C., Gollub, R.L., Göring, H.H.H., Green, R.C., Gruber, O., Gudnason, V., Guelfi, S., Håberg, A.K., Hansell, N.K., Hardy, J., Hartman, C.A., Hashimoto, R., Hegenscheid, K., Heinz, A., Le Hellard, S., Hernandez, D.G., Heslenfeld, D.J., Ho, B.C., Hoekstra, P.J., Hoffmann, W., Hofman, A., Holsboer, F., Homuth, G., Hosten, N., Hottenga, J.J., Huentelman, M., Pol, H.E.H., Ikeda, M., Jack, C.R., Jenkinson, M., Johnson, R., Jönsson, E.G., Jukema, J.W., Kahn, R.S., Kanai, R., Kloszewska, I., Knopman, D.S., Kochunov, P., Kwok, J.B., Lawrie, S.M., Lemaître, H., Liu, X., Longo, D.L., Lopez, O.L., Lovestone, S., Martinez, O., Martinot, J.L., Mattay, V.S., McDonald, C., McIntosh, A.M., McMahon, F.J., McMahon, K.L., Mecocci, P., Melle,

I., Meyer-Lindenberg, A., Mohnke, S., Montgomery, G.W., Morris, D.W., Mosley, T.H., Mühleisen, T.W., Müller-Myhsok, B., Nalls, M.A., Nauck, M., Nichols, T.E., Niessen, W.J., Nöthen, M.M., Nyberg, L., Ohi, K., Olvera, R.L., Ophoff, R.A., Pandolfo, M., Paus, T., Pausova, Z., Penninx, B.W.J.H., Pike, G.B., Potkin, S.G., Psaty, B.M., Reppermund, S., Rietschel, M., Roffman, J.L., Romanczuk-Seiferth, N., Rotter, J.I., Ryten, M., Sacco, R.L., Sachdev, P.S., Saykin, A.J., Schmidt, R., Schmidt, H., Schofield, P.R., Sigursson, S., Simmons, A., Singleton, A., Sisodiya, S.M., Smith, C., Smoller, J.W., Soininen, H., Steen, V.M., Stott, D.J., Sussmann, J.E., Thalamuthu, A., Toga, A.W., Traynor, B.J., Troncoso, J., Tsolaki, M., Tzourio, C., Uitterlinden, A.G., Hernández, M.C.V., Van Der Brug, M., Van Der Lugt, A., Van Der Wee, N.J.A., Van Haren, N.E.M., Van 'T Ent, D., Van Tol, M.J., Vardarajan, B.N., Vellas, B., Veltman, D.J., Völzke, H., Walter, H., Wardlaw, J.M., Wassink, T.H., Weale, M.E., Weinberger, D.R., Weiner, M.W., Wen, W., Westman, E., White, T., Wong, T.Y., Wright, C.B., Zielke, R.H., Zonderman, A.B., Martin, N.G., Van Duijn, C.M., Wright, M.J., Longstreth, W.T., Schumann, G., Grabe, H.J., Franke, B., Launer, L.J., Medland, S.E., Seshadri, S., Thompson, P.M., Ikram, M.A., 2017. Novel genetic loci associated with hippocampal volume. Nat. Commun. 8. https://doi.org/10.1038/ncomms13624

Hibar, D.P., Stein, J.L., Renteria, M.E., Arias-Vasquez, A., Desrivières, S., Jahanshad, N., Toro, R., Wittfeld, K., Abramovic, L., Andersson, M., Aribisala, B.S., Armstrong, N.J., Bernard, M., Bohlken, M.M., Boks, M.P., Bralten, J., Brown, A.A., Mallar Chakravarty, M., Chen, Q., Ching, C.R.K., Cuellar-Partida, G., Den Braber, A., Giddaluru, S., Goldman, A.L., Grimm, O., Guadalupe, T., Hass, J., Woldehawariat, G., Holmes, A.J., Hoogman, M., Janowitz, D., Jia, T., Kim, S., Klein, M., Kraemer, B., Lee, P.H., Olde Loohuis, L.M., Luciano, M., MacAre, C., Mather, K.A., Mattheisen, M., Milaneschi, Y., Nho, K., Papmeyer, M., Ramasamy, A., Risacher, S.L., Roiz-Santiañez, R., Rose, E.J., Salami, A., Sämann, P.G., Schmaal, L., Schork, A.J., Shin, J., Strike, L.T., Teumer, A., Van Donkelaar, M.M.J., Van Eijk, K.R., Walters, R.K., Westlye, L.T., Whelan, C.D., Winkler, A.M., Zwiers, M.P., Alhusaini, S., Athanasiu, L., Ehrlich, S., Hakobjan, M.M.H., Hartberg, C.B., Haukvik, U.K., Heister, A.J.G.A.M., Hoehn, D., Kasperaviciute, D., Liewald, D.C.M., Lopez, L.M., Makkinje, R.R.R., Matarin, M., Naber, M.A.M., Reese McKay, D., Needham, M., Nugent, A.C., Pütz, B., Royle, N.A., Shen, L., Sprooten, E., Trabzuni, D., Van Der Marel, S.S.L., Van Hulzen, K.J.E., Walton, E., Wolf, C., Almasy, L., Ames, D., Arepalli, S., Assareh, A.A., Bastin, M.E., Brodaty, H., Bulayeva, K.B., Carless, M.A., Cichon, S., Corvin, A., Curran, J.E., Czisch, M., De Zubicaray, G.I., Dillman, A., Duggirala, R., Dyer, T.D., Erk, S., Fedko, I.O., Ferrucci, L., Foroud, T.M., Fox, P.T., Fukunaga, M., Raphael Gibbs, J., Göring, H.H.H., Green, R.C., Guelfi, S., Hansell, N.K., Hartman, C.A., Hegenscheid, K., Heinz, A., Hernandez, D.G., Heslenfeld, D.J., Hoekstra, P.J., Holsboer, F., Homuth, G., Hottenga, J.J., Ikeda, M., Jack, C.R., Jenkinson, M., Johnson, R., Kanai, R., Keil, M., Kent, J.W., Kochunov, P., Kwok, J.B., Lawrie, S.M., Liu, X., Longo, D.L., McMahon, K.L., Meisenzahl, E., Melle, I., Mohnke, S., Montgomery, G.W., Mostert, J.C., Mühleisen, T.W., Nalls, M.A., Nichols, T.E., Nilsson, L.G., Nöthen, M.M., Ohi, K., Olvera, R.L., Perez-Iglesias, R., Bruce Pike, G., Potkin, S.G., Reinvang, I., Reppermund, S., Rietschel, M., Romanczuk-Seiferth, N., Rosen, G.D., Rujescu, D., Schnell, K., Schofield, P.R., Smith, C., Steen, V.M., Sussmann, J.E., Thalamuthu, A., Toga, A.W., Traynor, B.J., Troncoso, J., Turner, J.A., Valdés Hernández, M.C., Van 'T Ent, D., Van Der Brug, M., Van Der Wee, N.J.A., Van Tol, M.J., Veltman, D.J., Wassink, T.H., Westman, E., Zielke, R.H., Zonderman, A.B., Ashbrook, D.G., Hager, R., Lu, L., McMahon, F.J., Morris, D.W., Williams, R.W., Brunner, H.G., Buckner, R.L., Buitelaar, J.K., Cahn, W., Calhoun, V.D.,

Cavalleri, G.L., Crespo-Facorro, B., Dale, A.M., Davies, G.E., Delanty, N., Depondt, C.,
Djurovic, S., Drevets, W.C., Espeseth, T., Gollub, R.L., Ho, B.C., Hoffmann, W., Hosten,
N., Kahn, R.S., Le Hellard, S., Meyer-Lindenberg, A., Müller-Myhsok, B., Nauck, M.,
Nyberg, L., Pandolfo, M., Penninx, B.W.J.H., Roffman, J.L., Sisodiya, S.M., Smoller, J.W.,
Van Bokhoven, H., Van Haren, N.E.M., Völzke, H., Walter, H., Weiner, M.W., Wen, W.,
White, T., Agartz, I., Andreassen, O.A., Blangero, J., Boomsma, D.I., Brouwer, R.M.,
Cannon, D.M., Cookson, M.R., De Geus, E.J.C., Deary, I.J., Donohoe, G., Fernández, G.,
Fisher, S.E., Francks, C., Glahn, D.C., Grabe, H.J., Gruber, O., Hardy, J., Hashimoto, R.,
Hulshoff Pol, H.E., Jönsson, E.G., Kloszewska, I., Lovestone, S., Mattay, V.S., Mecocci, P.,
McDonald, C., McIntosh, A.M., Ophoff, R.A., Paus, T., Pausova, Z., Ryten, M., Sachdev,
P.S., Saykin, A.J., Simmons, A., Singleton, A., Soininen, H., Wardlaw, J.M., Weale, M.E.,
Weinberger, D.R., Adams, H.H.H., Launer, L.J., Seiler, S., Schmidt, R., Chauhan, G.,
Satizabal, C.L., Becker, J.T., Yanek, L., Van Der Lee, S.J., Ebling, M., Fischl, B.,
Longstreth, W.T., Greve, D., Schmidt, H., Nyquist, P., Vinke, L.N., Van Duijn, C.M., Xue,
L., Mazoyer, B., Bis, J.C., Gudnason, V., Seshadri, S., Ikram, M.A., Martin, N.G., Wright,
M.J., Schumann, G., Franke, B., Thompson, P.M., Medland, S.E., 2015. Common genetic
variants influence human subcortical brain structures. Nature 520, 224–229.
https://doi.org/10.1038/nature14101
Honey, C.J., Honey, C.J., Sporns, O., Sporns, O., Cammoun, L., Cammoun, L., Gigandet, X.,
Gigandet, X., Thiran, J.P., Thiran, J.P., Meuli, R., Meuli, R., Hagmann, P., Hagmann, P.,
2009. Predicting human resting-state functional connectivity from structural connectivity.
Proc. Natl. Acad. Sci. U. S. A. 106, 2035–40. https://doi.org/10.1073/pnas.0811168106
Hurlburt, R.T., Alderson-Day, B., Fernyhough, C., Kï¿½hn, S., 2015. What goes on in the
resting-state? A qualitative glimpse into resting-state experience in the scanner. Front.
Psychol. 6. https://doi.org/10.3389/fpsyg.2015.01535
Huth, A.G., De Heer, W.A., Griffiths, T.L., Theunissen, F.E., Gallant, J.L., 2016. Natural speech
reveals the semantic maps that tile human cerebral cortex. Nature 532, 453–458.
https://doi.org/10.1038/nature17637
Ioannidis, J.P.A., 2008. Why Most Discovered True Associations Are Inflated. Epidemiology 19,
640–648. https://doi.org/10.1097/EDE.0b013e31818131e7
Ioannidis, J.P.A., 2005. Why most published research findings are false. PLoS Med.
https://doi.org/10.1371/journal.pmed.0020124
Ioannidis, J.P.A., Ntzani, E.E., Trikalinos, T.A., Contopoulos-Ioannidis, D.G., 2001. Replication
validity of genetic association studies. Nat. Genet. 29, 306–309.
https://doi.org/10.1038/ng749
Jo, H.J., Gotts, S.J., Reynolds, R.C., Bandettini, P.A., Martin, A., Cox, R.W., Saad, Z.S., 2013.
Effective preprocessing procedures virtually eliminate distance-dependent motion artifacts
in resting state FMRI. J. Appl. Math. 2013. https://doi.org/10.1155/2013/935154
Kaiser, R.H., Andrews-Hanna, J.R., Wager, T.D., Pizzagalli, D.A., 2015. Large-Scale Network
Dysfunction in Major Depressive Disorder A Meta-analysis of Resting-State Functional
Connectivity. JAMA psychiatry 72, 603–611.
https://doi.org/10.1001/jamapsychiatry.2015.0071
Kanyongo, G.Y., Brook, G.P., Kyei-Blankson, L., Gocmen, G., 2007. Reliability and Statistical
Power: How Measurement Fallibility Affects Power and Required Sample Sizes for Several
Parametric and Nonparametric Statistics. J. Mod. Appl. Stat. Methods 6, 81–90.
https://doi.org/10.22237/jmasm/1177992480
Klein, A., Andersson, J., Ardekani, B.A., Ashburner, J., Avants, B., Chiang, M.C., Christensen,
G.E., Collins, D.L., Gee, J., Hellier, P., Song, J.H., Jenkinson, M., Lepage, C., Rueckert, D.,

Thompson, P., Vercauteren, T., Woods, R.P., Mann, J.J., Parsey, R. V., 2009. Evaluation of 14 nonlinear deformation algorithms applied to human brain MRI registration. Neuroimage 46, 786–802. https://doi.org/10.1016/j.neuroimage.2008.12.037

Krienen, F.M., Yeo, B.T.T., Buckner, R.L., 2014. Reconfigurable task-dependent functional coupling modes cluster around a core functional architecture. Philos. Trans. R. Soc. Lond. B. Biol. Sci. 369, 20130526-. https://doi.org/10.1098/rstb.2013.0526

Lahnakoski, J.M., Glerean, E., Salmi, J., Jääskeläinen, I.P., Sams, M., Hari, R., Nummenmaa, L., 2012. Naturalistic fMRI Mapping Reveals Superior Temporal Sulcus as the Hub for the Distributed Brain Network for Social Perception. Front. Hum. Neurosci. 6. https://doi.org/10.3389/fnhum.2012.00233

Laumann, T.O., Gordon, E.M., Adeyemo, B., Snyder, A.Z., Joo, S.J., Chen, M.Y., Gilmore, A.W., McDermott, K.B., Nelson, S.M., Dosenbach, N.U.F., Schlaggar, B.L., Mumford, J.A., Poldrack, R.A., Petersen, S.E., 2015. Functional System and Areal Organization of a Highly Sampled Individual Human Brain. Neuron 87, 658–671. https://doi.org/10.1016/j.neuron.2015.06.037

Leech, R., Braga, R., Sharp, D.J., 2012. Echoes of the Brain within the Posterior Cingulate Cortex. J. Neurosci. 32, 215–222. https://doi.org/10.1523/JNEUROSCI.3689-11.2012

Margulies, D.S., Ghosh, S.S., Goulas, A., Falkiewicz, M., Huntenburg, J.M., Langs, G., Bezgin, G., Eickhoff, S.B., Castellanos, F.X., Petrides, M., Jefferies, E., Smallwood, J., 2016. Situating the default-mode network along a principal gradient of macroscale cortical organization. Proc. Natl. Acad. Sci. U. S. A. 113, 12574–12579. https://doi.org/10.1073/pnas.1608282113

Matthews, P.M., Honey, G.D., Bullmore, E.T., 2006. Applications of fMRI in translational medicine and clinical practice. Nat. Rev. Neurosci. https://doi.org/10.1038/nrn1929

Mesulam, M.M., 1998. From sensation to cognition. Brain. https://doi.org/10.1093/brain/121.6.1013

Meyer-Lindenberg, A., Weinberger, D.R., 2006. Intermediate phenotypes and genetic mechanisms of psychiatric disorders. Nat. Rev. Neurosci. https://doi.org/10.1038/nrn1993

Miller, K.L., Alfaro-Almagro, F., Bangerter, N.K., Thomas, D.L., Yacoub, E., Xu, J., Bartsch, A.J., Jbabdi, S., Sotiropoulos, S.N., Andersson, J.L.R., Griffanti, L., Douaud, G., Okell, T.W., Weale, P., Dragonu, I., Garratt, S., Hudson, S., Collins, R., Jenkinson, M., Matthews, P.M., Smith, S.M., 2016. Multimodal population brain imaging in the UK Biobank prospective epidemiological study. Nat. Neurosci. 19, 1523. https://doi.org/10.1038/nn.4393

Mueller, S., Wang, D., Fox, M.D., Yeo, B.T.T., Sepulcre, J., Sabuncu, M.R., Shafee, R., Lu, J., Liu, H., 2013. Individual Variability in Functional Connectivity Architecture of the Human Brain. Neuron 77, 586–595. https://doi.org/10.1016/j.neuron.2012.12.028

Munafò, M.R., Nosek, B.A., Bishop, D.V.M., Button, K.S., Chambers, C.D., Percie Du Sert, N., Simonsohn, U., Wagenmakers, E.J., Ware, J.J., Ioannidis, J.P.A., 2017. A manifesto for reproducible science. Nat. Hum. Behav. 1, 1–9. https://doi.org/10.1038/s41562-016-0021

Murphy, K., Fox, M.D., 2017. Towards a consensus regarding global signal regression for resting state functional connectivity MRI. Neuroimage 154, 169–173. https://doi.org/10.1016/j.neuroimage.2016.11.052

Neale, M.C., Hunter, M.D., Pritkin, J., Zahery, M., Brick, T.R., Kirkpatrick, R.M., Estabrook, R., Bates, T.C., Maes, H.H., Boker, S.M., 2016. OpenMx 2.0: Extended Structural Equation and Statistical Modeling. Psychometrika 81, 535–549. https://doi.org/10.1007/s11336-014-9435-8

Noble, S., Spann, M.N., Tokoglu, F., Shen, X., Constable, R.T., Scheinost, D., 2017. Influences on the Test–Retest Reliability of Functional Connectivity MRI and its Relationship with

Behavioral Utility. Cereb. Cortex 1–15. https://doi.org/10.1093/cercor/bhx230

Nunnally Jr., J.C., 1970. Introduction to psychological measurement., Introduction to
psychological measurement. McGraw-Hill, New York, NY, US.

Patriat, R., Molloy, E.K., Meier, T.B., Kirk, G.R., Nair, V.A., Meyerand, M.E., Prabhakaran, V.,
Birn, R.M., 2013. The effect of resting condition on resting-state fMRI reliability and
consistency: A comparison between resting with eyes open, closed, and fixated.
Neuroimage 78, 463–473. https://doi.org/10.1016/j.neuroimage.2013.04.013

Pek, J., Wu, H., 2015. Profile Likelihood-Based Confidence Intervals and Regions for Structural
Equation Models. Psychometrika 80, 1123–1145. https://doi.org/10.1007/s11336-015-9461-
1

Poldrack, R.A., Baker, C.I., Durnez, J., Gorgolewski, K.J., Matthews, P.M., Munafò, M.R.,
Nichols, T.E., Poline, J.B., Vul, E., Yarkoni, T., 2017. Scanning the horizon: Towards
transparent and reproducible neuroimaging research. Nat. Rev. Neurosci. 18, 115–126.
https://doi.org/10.1038/nrn.2016.167

Posthuma, D., Boomsma, D.I., 2000. A note on the statistical power in extended twin designs.
Behav. Genet. 30, 147–158.

Poulton, R., Moffitt, T.E., Silva, P.A., 2015. The Dunedin Multidisciplinary Health and
Development Study: overview of the first 40 years, with an eye to the future. Soc.
Psychiatry Psychiatr. Epidemiol. https://doi.org/10.1007/s00127-015-1048-8

Power, J.D., Barnes, K.A., Snyder, A.Z., Schlaggar, B.L., Petersen, S.E., 2012. Spurious but
systematic correlations in functional connectivity MRI networks arise from subject motion.
Neuroimage 59, 2142–2154. https://doi.org/10.1016/j.neuroimage.2011.10.018

Power, J.D., Cohen, A.L., Nelson, S.M., Wig, G.S., Barnes, K.A., Church, J.A., Vogel, A.C.,
Laumann, T.O., Miezin, F.M., Schlaggar, B.L., Petersen, S.E., 2011. Functional Network
Organization of the Human Brain. Neuron 72, 665–678.
https://doi.org/10.1016/j.neuron.2011.09.006

Power, J.D., Mitra, A., Laumann, T.O., Snyder, A.Z., Schlaggar, B.L., Petersen, S.E., 2014.
Methods to detect, characterize, and remove motion artifact in resting state fMRI.
Neuroimage 84, 320–341. https://doi.org/10.1016/j.neuroimage.2013.08.048

Power, J.D., Plitt, M., Gotts, S.J., Kundu, P., Voon, V., Bandettini, P.A., Martin, A., 2018.
Ridding fMRI data of motion-related influences: Removal of signals with distinct spatial
and physical bases in multiecho data. Proc. Natl. Acad. Sci. 115, 201720985.
https://doi.org/10.1073/pnas.1720985115

Power, J.D., Plitt, M., Laumann, T.O., Martin, A., 2016. Sources and implications of whole-brain
fMRI signals in humans. Neuroimage 146, 0–1.
https://doi.org/10.1016/j.neuroimage.2016.09.038

Power, J.D., Schlaggar, B.L., Lessov-Schlaggar, C.N., Petersen, S.E., 2013. Evidence for hubs in
human functional brain networks. Neuron 79, 798–813.
https://doi.org/10.1016/j.neuron.2013.07.035

Raven, J.C., 1941. Standardization of progressive matrices. Br. J. Med. Psychol. 19, 137–150.
https://doi.org/10.1111/j.2044-8341.1941.tb00316.x

Satterthwaite, T.D., Cook, P.A., Bruce, S.E., Conway, C., Mikkelsen, E., Satchell, E., Vandekar,
S.N., Durbin, T., Shinohara, R.T., Sheline, Y.I., 2016. Dimensional depression severity in
women with major depression and post-traumatic stress disorder correlates with fronto-
amygdalar hypoconnectivty. Mol. Psychiatry 21, 894–902.
https://doi.org/10.1038/mp.2015.149

Satterthwaite, T.D., Elliott, M.A., Gerraty, R.T., Ruparel, K., Loughead, J., Calkins, M.E.,
Eickhoff, S.B., Hakonarson, H., Gur, R.C., Gur, R.E., Wolf, D.H., 2013. An improved

framework for confound regression and filtering for control of motion artifact in the preprocessing of resting-state functional connectivity data. Neuroimage 64, 240–256. https://doi.org/10.1016/j.neuroimage.2012.08.052

Satterthwaite, T.D., Xia, C.H., Bassett, D.S., 2018. Personalized Neuroscience: Common and Individual-Specific Features in Functional Brain Networks. Neuron 98, 243–245. https://doi.org/10.1016/j.neuron.2018.04.007

Shah, L.M., Cramer, J.A., Ferguson, M.A., Birn, R.M., Anderson, J.S., 2016. Reliability and reproducibility of individual differences in functional connectivity acquired during task and resting state. Brain Behav. 6, 1–15. https://doi.org/10.1002/brb3.456

Shehzad, Z., Kelly, A.M.C., Reiss, P.T., Gee, D.G., Gotimer, K., Uddin, L.Q., Lee, S.H., Margulies, D.S., Roy, A.K., Biswal, B.B., Petkova, E., Castellanos, F.X., Milham, M.P., 2009. The resting brain: Unconstrained yet reliable. Cereb. Cortex 19, 2209–2229. https://doi.org/10.1093/cercor/bhn256

Shen, X., Finn, E.S., Scheinost, D., Rosenberg, M.D., Chun, M.M., Papademetris, X., Constable, R.T., 2017. Using connectome-based predictive modeling to predict individual behavior from brain connectivity. Nat. Protoc. 12, 506–518. https://doi.org/10.1038/nprot.2016.178

Smith, S.M., Beckmann, C.F., Andersson, J., Auerbach, E.J., Bijsterbosch, J., Douaud, G., Duff, E., Feinberg, D.A., Griffanti, L., Harms, M.P., Kelly, M., Laumann, T., Miller, K.L., Moeller, S., Petersen, S., Power, J., Salimi-Khorshidi, G., Snyder, A.Z., Vu, A.T., Woolrich, M.W., Xu, J., Yacoub, E., Uğurbil, K., Van Essen, D.C., Glasser, M.F., 2013. Resting-state fMRI in the Human Connectome Project. Neuroimage 80, 144–168. https://doi.org/10.1016/j.neuroimage.2013.05.039

Smith, S.M., Fox, P.T., Miller, K.L., Glahn, D.C., Fox, P.M., Mackay, C.E., Filippini, N., Watkins, K.E., Toro, R., Laird, A.R., Beckmann, C.F., 2009. Correspondence of the brain's functional architecture during activation and rest. Proc. Natl. Acad. Sci. https://doi.org/10.1073/pnas.0905267106

Steiger, J.H., 1980. Tests for comparing elements of a correlation matrix. Psychol. Bull. 87, 245–251. https://doi.org/10.1037/0033-2909.87.2.245

Stein, J.L., Medland, S.E., Vasquez, A.A., Hibar, D.P., Senstad, R.E., Winkler, A.M., Toro, R., Appel, K., Bartecek, R., Bergmann, Ø., Bernard, M., Brown, A.A., Cannon, D.M., Chakravarty, M.M., Christoforou, A., Domin, M., Grimm, O., Hollinshead, M., Holmes, A.J., Homuth, G., Hottenga, J.J., Langan, C., Lopez, L.M., Hansell, N.K., Hwang, K.S., Kim, S., Laje, G., Lee, P.H., Liu, X., Loth, E., Lourdusamy, A., Mattingsdal, M., Mohnke, S., Maniega, S.M., Nho, K., Nugent, A.C., O'brien, C., Papmeyer, M., Pütz, B., Ramasamy, A., Rasmussen, J., Rijpkema, M., Risacher, S.L., Roddey, J.C., Rose, E.J., Ryten, M., Shen, L., Sprooten, E., Strengman, E., Teumer, A., Trabzuni, D., Turner, J., Van Eijk, K., Van Erp, T.G.M., Van Tol, M.J., Wittfeld, K., Wolf, C., Woudstra, S., Aleman, A., Alhusaini, S., Almasy, L., Binder, E.B., Brohawn, D.G., Cantor, R.M., Carless, M.A., Corvin, A., Czisch, M., Curran, J.E., Davies, G., De Almeida, M.A.A., Delanty, N., Depondt, C., Duggirala, R., Dyer, T.D., Erk, S., Fagerness, J., Fox, P.T., Freimer, N.B., Gill, M., Göring, H.H.H., Hagler, D.J., Hoehn, D., Holsboer, F., Hoogman, M., Hosten, N., Jahanshad, N., Johnson, M.P., Kasperaviciute, D., Kent, J.W., Kochunov, P., Lancaster, J.L., Lawrie, S.M., Liewald, D.C., Mandl, R., Matarin, M., Mattheisen, M., Meisenzahl, E., Melle, I., Moses, E.K., Mühleisen, T.W., Nauck, M., Nöthen, M.M., Olvera, R.L., Pandolfo, M., Pike, G.B., Puls, R., Reinvang, I., Rentería, M.E., Rietschel, M., Roffman, J.L., Royle, N.A., Rujescu, D., Savitz, J., Schnack, H.G., Schnell, K., Seiferth, N., Smith, C., Steen, V.M., Hernández, M.C.V., Van Den Heuvel, M., Van Der Wee, N.J., Van Haren, N.E.M., Veltman, J.A., Völzke, H., Walker, R., Westlye, L.T., Whelan, C.D., Agartz, I., Boomsma, D.I., Cavalleri,

G.L., Dale, A.M., Djurovic, S., Drevets, W.C., Hagoort, P., Hall, J., Heinz, A., Jack, C.R.,
Foroud, T.M., Le Hellard, S., Macciardi, F., Montgomery, G.W., Poline, J.B., Porteous,
D.J., Sisodiya, S.M., Starr, J.M., Sussmann, J., Toga, A.W., Veltman, D.J., Walter, H.,
Weiner, M.W., Bis, J.C., Ikram, M.A., Smith, A. V., Gudnason, V., Tzourio, C., Vernooij,
M.W., Launer, L.J., Decarli, C., Seshadri, S., Andreassen, O.A., Apostolova, L.G., Bastin,
M.E., Blangero, J., Brunner, H.G., Buckner, R.L., Cichon, S., Coppola, G., De Zubicaray,
G.I., Deary, I.J., Donohoe, G., De Geus, E.J.C., Espeseth, T., Fernéndez, G., Glahn, D.C.,
Grabe, H.J., Hardy, J., Hulshoff Pol, H.E., Jenkinson, M., Kahn, R.S., Mcdonald, C.,
Mcintosh, A.M., Mcmahon, F.J., Mcmahon, K.L., Meyer-Lindenberg, A., Morris, D.W.,
Müller-Myhsok, B., Nichols, T.E., Ophoff, R.A., Paus, T., Pausova, Z., Penninx, B.W.,
Potkin, S.G., Sämann, P.G., Saykin, A.J., Schumann, G., Smoller, J.W., Wardlaw, J.M.,
Weale, M.E., Martin, N.G., Franke, B., Wright, M.J., Thompson, P.M., 2012. Identification
of common variants associated with human hippocampal and intracranial volumes. Nat.
Genet. 44, 552–561. https://doi.org/10.1038/ng.2250

Sudlow, C., Gallacher, J., Allen, N., Beral, V., Burton, P., Danesh, J., Downey, P., Elliott, P.,
Green, J., Landray, M., Liu, B., Matthews, P., Ong, G., Pell, J., Silman, A., Young, A.,
Sprosen, T., Peakman, T., Collins, R., 2015. UK Biobank: An Open Access Resource for
Identifying the Causes of a Wide Range of Complex Diseases of Middle and Old Age.
PLoS Med. 12. https://doi.org/10.1371/journal.pmed.1001779

Swartz, J.R., Knodt, A.R., Radtke, S.R., Hariri, A.R., 2015. A neural biomarker of psychological
vulnerability to future life stress. Neuron 85, 505–511.
https://doi.org/10.1016/j.neuron.2014.12.055

Szucs, D., Ioannidis, J.P.A., 2017. Empirical assessment of published effect sizes and power in
the recent cognitive neuroscience and psychology literature. PLoS Biol. 15.
https://doi.org/10.1371/journal.pbio.2000797

Tagliazucchi, E., Laufs, H., 2014. Decoding Wakefulness Levels from Typical fMRI Resting-
State Data Reveals Reliable Drifts between Wakefulness and Sleep. Neuron 82, 695–708.
https://doi.org/10.1016/j.neuron.2014.03.020

Tavor, I., Parker Jones, O., Mars, R.B., Smith, S.M., Behrens, T.E., Jbabdi, S., 2016. Task-free
MRI predicts individual differences in brain activity during task performance. Science 352,
216–20. https://doi.org/10.1126/science.aad8127

Thompson, P.M., Stein, J.L., Medland, S.E., Hibar, D.P., Vasquez, A.A., Renteria, M.E., Toro,
R., Jahanshad, N., Schumann, G., Franke, B., Wright, M.J., Martin, N.G., Agartz, I., Alda,
M., Alhusaini, S., Almasy, L., Almeida, J., Alpert, K., Andreasen, N.C., Andreassen, O.A.,
Apostolova, L.G., Appel, K., Armstrong, N.J., Aribisala, B., Bastin, M.E., Bauer, M.,
Bearden, C.E., Bergmann, Ø., Binder, E.B., Blangero, J., Bockholt, H.J., Bøen, E., Bois, C.,
Boomsma, D.I., Booth, T., Bowman, I.J., Bralten, J., Brouwer, R.M., Brunner, H.G.,
Brohawn, D.G., Buckner, R.L., Buitelaar, J., Bulayeva, K., Bustillo, J.R., Calhoun, V.D.,
Cannon, D.M., Cantor, R.M., Carless, M.A., Caseras, X., Cavalleri, G.L., Chakravarty,
M.M., Chang, K.D., Ching, C.R.K., Christoforou, A., Cichon, S., Clark, V.P., Conrod, P.,
Coppola, G., Crespo-Facorro, B., Curran, J.E., Czisch, M., Deary, I.J., de Geus, E.J.C., den
Braber, A., Delvecchio, G., Depondt, C., de Haan, L., de Zubicaray, G.I., Dima, D.,
Dimitrova, R., Djurovic, S., Dong, H., Donohoe, G., Duggirala, R., Dyer, T.D., Ehrlich, S.,
Ekman, C.J., Elvsåshagen, T., Emsell, L., Erk, S., Espeseth, T., Fagerness, J., Fears, S.,
Fedko, I., Fernández, G., Fisher, S.E., Foroud, T., Fox, P.T., Francks, C., Frangou, S., Frey,
E.M., Frodl, T., Frouin, V., Garavan, H., Giddaluru, S., Glahn, D.C., Godlewska, B.,
Goldstein, R.Z., Gollub, R.L., Grabe, H.J., Grimm, O., Gruber, O., Guadalupe, T., Gur,
R.E., Gur, R.C., Göring, H.H.H., Hagenaars, S., Hajek, T., Hall, G.B., Hall, J., Hardy, J.,

Hartman, C.A., Hass, J., Hatton, S.N., Haukvik, U.K., Hegenscheid, K., Heinz, A., Hickie, I.B., Ho, B.C., Hoehn, D., Hoekstra, P.J., Hollinshead, M., Holmes, A.J., Homuth, G., Hoogman, M., Hong, L.E., Hosten, N., Hottenga, J.J., Hulshoff Pol, H.E., Hwang, K.S., Jack, C.R., Jenkinson, M., Johnston, C., Jönsson, E.G., Kahn, R.S., Kasperaviciute, D., Kelly, S., Kim, S., Kochunov, P., Koenders, L., Krämer, B., Kwok, J.B.J., Lagopoulos, J., Laje, G., Landen, M., Landman, B.A., Lauriello, J., Lawrie, S.M., Lee, P.H., Le Hellard, S., Lemaître, H., Leonardo, C.D., Li, C. shan, Liberg, B., Liewald, D.C., Liu, X., Lopez, L.M., Loth, E., Lourdusamy, A., Luciano, M., Macciardi, F., Machielsen, M.W.J., MacQueen, G.M., Malt, U.F., Mandl, R., Manoach, D.S., Martinot, J.L., Matarin, M., Mather, K.A., Mattheisen, M., Mattingsdal, M., Meyer-Lindenberg, A., McDonald, C., McIntosh, A.M., McMahon, F.J., McMahon, K.L., Meisenzahl, E., Melle, I., Milaneschi, Y., Mohnke, S., Montgomery, G.W., Morris, D.W., Moses, E.K., Mueller, B.A., Muñoz Maniega, S., Mühleisen, T.W., Müller-Myhsok, B., Mwangi, B., Nauck, M., Nho, K., Nichols, T.E., Nilsson, L.G., Nugent, A.C., Nyberg, L., Olvera, R.L., Oosterlaan, J., Ophoff, R.A., Pandolfo, M., Papalampropoulou-Tsiridou, M., Papmeyer, M., Paus, T., Pausova, Z., Pearlson, G.D., Penninx, B.W., Peterson, C.P., Pfennig, A., Phillips, M., Pike, G.B., Poline, J.B., Potkin, S.G., Pütz, B., Ramasamy, A., Rasmussen, J., Rietschel, M., Rijpkema, M., Risacher, S.L., Roffman, J.L., Roiz-Santiañez, R., Romanczuk-Seiferth, N., Rose, E.J., Royle, N.A., Rujescu, D., Ryten, M., Sachdev, P.S., Salami, A., Satterthwaite, T.D., Savitz, J., Saykin, A.J., Scanlon, C., Schmaal, L., Schnack, H.G., Schork, A.J., Schulz, S.C., Schür, R., Seidman, L., Shen, L., Shoemaker, J.M., Simmons, A., Sisodiya, S.M., Smith, C., Smoller, J.W., Soares, J.C., Sponheim, S.R., Sprooten, E., Starr, J.M., Steen, V.M., Strakowski, S., Strike, L., Sussmann, J., Sämann, P.G., Teumer, A., Toga, A.W., Tordesillas-Gutierrez, D., Trabzuni, D., Trost, S., Turner, J., Van den Heuvel, M., van der Wee, N.J., van Eijk, K., van Erp, T.G.M., van Haren, N.E.M., van 't Ent, D., van Tol, M.J., Valdés Hernández, M.C., Veltman, D.J., Versace, A., Völzke, H., Walker, R., Walter, H., Wang, L., Wardlaw, J.M., Weale, M.E., Weiner, M.W., Wen, W., Westlye, L.T., Whalley, H.C., Whelan, C.D., White, T., Winkler, A.M., Wittfeld, K., Woldehawariat, G., Wolf, C., Zilles, D., Zwiers, M.P., Thalamuthu, A., Schofield, P.R., Freimer, N.B., Lawrence, N.S., Drevets, W., 2014. The ENIGMA Consortium: Large-scale collaborative analyses of neuroimaging and genetic data. Brain Imaging Behav. 8, 153–182. https://doi.org/10.1007/s11682-013-9269-5

Van Dijk, K.R.A., Hedden, T., Venkataraman, A., Evans, K.C., Lazar, S.W., Buckner, R.L., 2010. Intrinsic Functional Connectivity As a Tool For Human Connectomics: Theory, Properties, and Optimization. J. Neurophysiol. 103, 297–321. https://doi.org/10.1152/jn.00783.2009

van Dijk, K.R.A., Sabuncu, M.R., Buckner, R.L., 2012. The influence of head motion on intrinsic functional connectivity MRI. Neuroimage 59, 431–438. https://doi.org/10.1016/j.neuroimage.2011.07.044

Van Essen, D.C., Smith, S.M., Barch, D.M., Behrens, T.E.J., Yacoub, E., Ugurbil, K., 2013. The WU-Minn Human Connectome Project: An overview. Neuroimage 80, 62–79. https://doi.org/10.1016/j.neuroimage.2013.05.041

Vanderwal, T., Eilbott, J., Finn, E.S., Craddock, R.C., Turnbull, A., Castellanos, F.X., 2017. Individual differences in functional connectivity during naturalistic viewing conditions. Neuroimage 157, 521–530. https://doi.org/10.1016/j.neuroimage.2017.06.027

Vul, E., Harris, C., Winkielman, P., Pashler, H., 2009. Puzzlingly High Correlations in fMRI Studies of Emotion, Personality, and Social Cognition1. Perspect. Psychol. Sci. 4, 274–290. https://doi.org/10.1111/j.1745-6924.2009.01125.x

Waites, A.B., Stanislavsky, A., Abbott, D.F., Jackson, G.D., 2005. Effect of prior cognitive state
on resting state networks measured with functional connectivity. Hum. Brain Mapp. 24, 59–
68. https://doi.org/10.1002/hbm.20069
Weschler, D., 2008. Wechsler Adult Intelligence Scale - Fourth Edition. Stat. Solut. 1–3.
Whelan, R., Garavan, H., 2014. When optimism hurts: Inflated predictions in psychiatric
neuroimaging. Biol. Psychiatry 75, 746–748.
https://doi.org/10.1016/j.biopsych.2013.05.014
Williams, R.H., Zimmerman, D.W., 1989. Statistical power analysis and reliability of
measurement. J. Gen. Psychol. 116, 359–369.
https://doi.org/10.1080/00221309.1989.9921123
Winkler, A.M., Kochunov, P., Blangero, J., Almasy, L., Zilles, K., Fox, P.T., Duggirala, R.,
Glahn, D.C., 2010. Cortical thickness or grey matter volume? The importance of selecting
the phenotype for imaging genetics studies. Neuroimage 53, 1135–1146.
https://doi.org/10.1016/j.neuroimage.2009.12.028
Wong, C.W., Olafsson, V., Tal, O., Liu, T.T., 2012. Anti-correlated networks, global signal
regression, and the effects of caffeine in resting-state functional MRI. Neuroimage 63, 356–
364. https://doi.org/10.1016/j.neuroimage.2012.06.035
Yarkoni, T., Westfall, J., 2017. Choosing Prediction Over Explanation in Psychology: Lessons
From Machine Learning. Perspect. Psychol. Sci. 12, 1100–1122.
https://doi.org/10.1177/1745691617693393
Yeo, B.T.T., Krienen, F.M., Sepulcre, J., Sabuncu, M.R., Lashkari, D., Hollinshead, M.,
Roffman, J.L., Smoller, J.W., Zollei, L., Polimeni, J.R., Fischl, B., Liu, H., Buckner, R.L.,
2011. The organization of the human cerebral cortex estimated by intrinsic functional
connectivity. J. Neurophysiol. 106, 1125–1165. https://doi.org/10.1152/jn.00338.2011.
Yuan, W., Altaye, M., Ret, J., Schmithorst, V., Byars, A.W., Plante, E., Holland, S.K., 2009.
Quantification of head motion in children during various fMRI language tasks. Hum. Brain
Mapp. 30, 1481–1489. https://doi.org/10.1002/hbm.20616
1561