Targeting light-gated chloride channels to neuronal somatodendritic domain reduces their

excitatory effect in the axon

Jessica E. Messier[1,3], Hongmei Chen[1,3], Zhao-Lin Cai[1,3], Mingshan Xue[1,2,3,*]

[1]Department of Neuroscience, [2]Department of Molecular and Human Genetics, Baylor College

of Medicine, Houston Texas, 77030, USA

[3]The Cain Foundation Laboratories, Jan and Dan Duncan Neurological Research Institute at

Texas Children's Hospital, Houston Texas, 77030, USA

*Correspondence: Mingshan Xue (mxue@bcm.edu)

Abstract

Light-gated chloride channels are emerging as promising optogenetic tools for inhibition of

neural activity. However, their effects depend on the transmembrane chloride electrochemical

gradient and may be complex due to the heterogeneity of this gradient in different developmental

stages, neuronal types, and subcellular compartments. Here we characterized a light-gated

chloride channel, GtACR2, in mouse cortical neurons. We found that GtACR2 activation

inhibited the soma, but unexpectedly depolarized the presynaptic terminals resulting in

neurotransmitter release. Other light-gated chloride channels had similar effects. Reducing the

chloride concentrations in the axon and presynaptic terminals diminished the GtACR2-induced

neurotransmitter release, indicating an excitatory effect of chloride channels in these

compartments. A novel hybrid somatodendritic targeting motif reduced the GtACR2-induced

neurotransmitter release while enhancing the somatic photocurrents. Our results highlight the

necessity of precisely determining the effects of light-gated chloride channels under specific

experimental conditions and provide a much-improved light-gated chloride channel for

optogenetic inhibition.

Introduction

Targeted manipulation of neural activity is a powerful approach in neuroscience that has

provided fundamental insights into the roles of specific neurons in nervous system functions.

Genetically encoded actuators such as light-gated ion channels or pumps enable control of neural

activity with unprecedented spatiotemporal specificity and are transforming neuroscience

research (Boyden et al., 2005; Han and Boyden, 2007; Li et al., 2005; Nagel et al., 2003; Zhang

et al., 2007). Actuators that enable neuronal activation are frequently used, but inhibitory

optogenetic tools are increasingly crucial because reversible and temporally precise suppression

of neuronal activity is key to revealing the causal roles of specific neurons in network dynamics

and behavior. The widely used light-driven inward chloride pumps and outward proton pumps,

such as *Natronomonas pharaonis* halorhodopsin (NpHR) and *Halorubrum sodomense*

archaerhodopsin (Arch), can hyperpolarize membrane potentials, independent of the

electrochemical gradients, to inhibit action potentials with millisecond precision (Chow et al.,

2010; Chuong et al., 2014; Han and Boyden, 2007; Han et al., 2011; Zhang et al., 2007).

However, their efficacies are limited because only one ion is transported per absorbed photon,

and their activation does not decrease membrane resistance. Light-gated chloride channels, such

as *Guillardia theta* anion channelrhodopsin 1 and 2 (GtACR1 and GtACR2), iC++, and iChloC,

overcome these limitations. They are highly sensitive to light, allow multiple ions to cross the

membrane per photocycle, and reduce membrane resistance, thereby potently inhibiting action

potentials (Berndt et al., 2014; Govorunova et al., 2015; 2017b; Wietek et al., 2014). Thus, light-

gated chloride channels are emerging as promising optogenetic tools for suppressing neuronal

activity (Govorunova et al., 2017a).

Determining the precise effects of light-gated ion channels or pumps under defined conditions is

a prerequisite to use them for interrogating the functions of specific neurons and circuits. This is

due to the possibility that these channels or pumps not only modulate membrane potentials but

also may affect other processes such as ion homeostasis or neurotransmitter release. For

example, activation of a light-driven inward chloride pump, eNpHR3.0, can transiently change

the reversal potential of $GABA_A$ receptors and alter the inhibitory synaptic inputs (Raimondo et

al., 2012). Prolonged activation of a light-driven outward proton pump, eArch3.0, can increase

presynaptic calcium concentrations and spontaneous neurotransmitter release (Mahn et al.,

2016). Despite these potential confounds, these inhibitory optogenetic molecules are increasingly

essential tools for the targeted silencing of neuronal populations, as long as these confounds are

understood and controlled (Allen et al., 2015). Therefore, it is also crucial to thoroughly

characterize light-gated chloride channels, as their effect depends on the difference between the

membrane potential and the reversal potential for chloride, both of which can vary in different

neuronal types and subcellular compartments (Marty and Llano, 2005; Trigo et al., 2008).

To this end, we investigated the effects of activating GtACR2 in mouse cortical excitatory and

inhibitory neurons. Much to our surprise, wide-field light activation of GtACR2 not only

inhibited the soma, but also caused neurotransmitter release onto neighboring neurons. A similar

phenomenon was observed with GtACR1, iC++, and iChloC. We further showed that GtACR2

activation in the axon and presynaptic terminals directly depolarized the membrane to induce

neurotransmitter release due to high chloride concentrations in these compartments. These data

explain the recent observations that photostimulation of neurons expressing GtACR1 or GtACR2

can paradoxically release neurotransmitters or generate action potentials (Mahn et al., 2016;

Malyshev et al., 2017). To reduce the excitatory effect of GtACR2, we screened a panel of

somatodendritic targeting motifs to reduce the trafficking of GtACR2 to the axon and

presynaptic terminals. We created a hybrid motif (Kv2.1C-linker-TlcnC) that is most effective in

concentrating GtACR2 in the somatodendritic domain. Activation of somatodendritically

targeted GtACR2 resulted in larger photocurrents at the soma and less neurotransmitter release

than wild type GtACR2. These results demonstrate that restricting localization of light-gated

chloride channels to the somatodendritic domain improves the inhibitory efficacy of these

optogenetic tools.

Results

Light activation of GtACR2 in mouse cortical neurons causes neurotransmitter release

To examine the efficacy of GtACR2 in mouse cortical excitatory neurons, we expressed a

GtACR2-EYFP fusion protein (referred to as GtACR2 below) together with a red fluorescent

protein, tdTomato, in layer 2/3 pyramidal neurons of the mouse visual or somatosensory cortex

by *in utero* electroporation of plasmids at embryonic day 14.5–15.5. We obtained acute coronal

brain slices from 3–8-week-old mice and observed that GtACR2 was present in the soma,

dendrites, and axon (*Figure 1A*). We performed whole-cell patch clamp recordings at the soma

of neurons expressing GtACR2 (GtACR2^{+} neurons) with a K^{+}-based pipette solution (*Figure

1B*). As previously reported, activation of GtACR2 by wide-field blue light (455 nm) potently

inhibited current-induced spiking in these neurons (***Figure 1C***). However, when we voltage

clamped the neurons to record GtACR2-mediated photocurrents, we unexpectedly found an

inward current that was superimposed on the photocurrent. This inward current resembled an

excitatory postsynaptic current (EPSC; ***Figure 1D***). To further investigate this phenomenon, we

recorded layer 2/3 pyramidal neurons that did not express GtACR2 (GtACR2⁻ neurons) with a

Cs⁺-based pipette solution (***Figure 1E***). GtACR2 activation generated inward currents in every

recorded GtACR2⁻ neuron that was voltage clamped at the reversal potential for GABAergic

inhibition (-60 mV). The onsets of these inward currents followed the onset of the blue light by

3.19 ± 0.26 ms (mean $\pm$ s.e.m., $n = 25$). These currents were abolished by the glutamatergic

receptor antagonists, NBQX and CPP (***Figure 1F***), or the voltage-gated sodium channel blocker,

tetrodotoxin (TTX; ***Figure 1G***), indicating that they were indeed monosynaptic EPSCs caused

by the glutamate transmitter released from GtACR2⁺ neurons. Activation of GtACR2 also

produced inhibitory postsynaptic currents (IPSCs) in GtACR2⁻ neurons that were voltage

clamped at the reversal potential for glutamatergic excitation (+10 mV). These IPSCs were

disynaptic because they were abolished by NBQX and CPP (***Figure 1-supplement 1***), indicating

that activating GtACR2 in pyramidal neurons can release sufficient glutamate to recruit

inhibitory interneurons.

To determine if the phenomenon of GtACR2-induced neurotransmitter release is also present in

inhibitory neurons, we expressed GtACR2 in parvalbumin-expressing (Pv) neurons by injecting

a Flpo recombinase-dependent adeno-associated virus (AAV) into the visual cortex of *Pvalb-2A-*

Flpo mice ($Pvalb^{Flpo/+}$) (Madisen et al., 2015) at postnatal day 1 (***Figure 1H***). Using acute brain

slices from 3–6-week-old mice, we found that activation of GtACR2 in Pv neurons generated

IPSCs in every recorded GtACR2⁻ layer 2/3 pyramidal neuron, and the IPSCs were abolished by

Gabazine, a GABA$_A$ receptor antagonist, or TTX (***Figure 1I–K***). The onsets of the IPSCs

followed the onset of the blue light by 2.47 ± 0.15 ms (mean $\pm$ s.e.m., $n = 17$), indicating that

they were monosynaptic IPSCs caused by the GABA transmitter released from GtACR2$^+$ Pv

neurons. Furthermore, repetitive activation of GtACR2 in layer 2/3 pyramidal neurons or Pv

neurons with a high-frequency train of light pulses produced reliable EPSCs or IPSCs,

respectively (***Figure 1-supplement 2A,B***).

These findings were unexpected, because the Nernst equilibrium potential of chloride becomes

lower than the action potential threshold in rodent cortical neurons after the second postnatal

week, and activation of chloride channels should not promote neurotransmitter release onto

neighboring cells (Ben-Ari, 2002; Owens et al., 1996). We thus sought to identify the cause of

this paradoxical neurotransmitter release and considered three possibilities. First, GtACR2 may

conduct cations to depolarize neurons. Second, an increase in the intracellular chloride or strong

hyperpolarization induced by GtACR2 activation may lead to rebound spikes. Third, GtACR2-

mediated chloride currents may be excitatory.

A GtACR2-mediated excitatory chloride conductance causes neurotransmitter release

First, it was reported that GtACR2 did not conduct physiological cations (Govorunova et al.,

2015), but some other light-gated chloride channels retained certain cation conductance (Berndt

et al., 2014; Wietek et al., 2014). Thus, we sought to verify that in cortical neurons, GtACR2 has

a similar reversal potential as a known chloride channel. To accomplish this, we determined the

reversal potential of GtACR2-mediated photocurrents in comparison with that of IPSCs

mediated by the endogenous GABA$_A$ receptors. We used *in utero* electroporation to express

GtACR2 in layer 2/3 pyramidal neurons and a Cre recombinase-dependent AAV to express a red

light-gated cation channelrhodopsin, ReaChR, in Pv neurons of *Pvalb-2A-Cre* mice (*Pvalb$^{Cre/+}$*)

(Madisen et al., 2010). We performed whole-cell voltage clamp recordings at the soma of a

GtACR2$^+$ layer 2/3 pyramidal neuron and a nearby GtACR2$^-$ pyramidal neuron simultaneously

(***Figure 2-supplement 1A***). In the GtACR2$^+$ neuron, we sequentially recorded the IPSCs induced

by activating ReaChR in Pv neurons via 617-nm light and the GtACR2-mediated photocurrents

activated by 455-nm light. Both IPSCs and photocurrents were recorded at different membrane

potentials to determine their reversal potentials in the same neuron (***Figure 2-supplement 1B,C***).

617-nm light does not activate GtACR2 (Govorunova et al., 2015), whereas 455-nm light

partially activates ReaChR (Lin et al., 2013). Thus, to avoid Pv neuron-mediated IPSCs

contaminating GtACR2-mediated photocurrents, we monitored the IPSCs in the GtACR2$^-$

neuron at the membrane potential of +10 mV to ensure that the intensity of the 455-nm light was

not sufficient to activate Pv neurons and generate IPSCs (***Figure 2-supplement 1B***). We found

that the reversal potentials of GtACR2-mediated photocurrents and GABAergic IPSCs were

similar for each neuron and well below the action potential threshold (***Figure 2-supplement 1D***),

indicating that GtACR2 does not conduct cations to cause neurotransmitter release.

Second, we tested if GtACR2-induced neurotransmitter release could be due to rebound

depolarization. An increase in the intracellular chloride caused by GtACR2 activation may

trigger rapid efflux of chloride after the blue light illumination terminates. However, this

possibility is unlikely because when GtACR2 was activated by a long pulse of blue light (e.g., 5

ms), neurotransmitter release occurred before the light illumination ended (see examples in

162 *Figure 1F,G*). Another possibility is that the strong hyperpolarization induced by GtACR2

163 activates hyperpolarization-activated I_h currents, which may depolarize the membrane potential

164 above the action potential threshold. However, pharmacological inhibition of I_h currents slightly

165 increased the amplitudes of GtACR2-induced EPSCs (*Figure 2-supplement 2A,B*), most likely

166 because inhibiting I_h currents increases neuronal membrane resistances (Robinson and

167 Siegelbaum, 2003). Thus, GtACR2-induced neurotransmitter release is not caused by rebound

168 depolarization.

170 Third, although GtACR2-mediated photocurrents are inhibitory at the soma, it is possible that the

171 chloride concentrations are higher in some other cellular compartments, such that the

172 electrochemical gradient causes chloride to exit the cell upon GtACR2 channel opening,

173 resulting in depolarization of the membrane potential. To test this hypothesis, we

174 pharmacologically inhibited the activity of Na^+-K^+-$2Cl^-$ cotransporter 1 (NKCC1) with

175 bumetanide (50 or 100 μM) to decease the intracellular chloride concentrations, as NKCC1 is

176 responsible for transporting chloride into neurons (Ben-Ari, 2017). When we activated GtACR2

177 in layer 2/3 pyramidal neurons, the resulting EPSCs in GtACR2$^-$ pyramidal neurons were

178 diminished by bath application of bumetanide (*Figure 2A,B*), indicating that GtACR2-induced

179 neurotransmitter release requires high concentrations of intracellular chloride. An alternative

180 interpretation of this result would be that bumetanide blocks GtACR2 itself. To test this

181 possibility, we simultaneously recorded the photocurrents and EPSCs in GtACR2$^+$ and GtACR2$^-$

182 pyramidal neurons, respectively. While bumetanide diminished the EPSCs in GtACR2$^-$ neurons,

183 it had no effect on the photocurrents in GtACR2$^+$ neurons (*Figure 2-supplement 3A,B*), thereby

184 ruling out the possibility that bumetanide affects GtACR2 itself. Furthermore, when a cation

185 channel, channelrhodopsin-2 (ChR2) (Boyden et al., 2005; Li et al., 2005; Nagel et al., 2003),

186 was expressed and activated in layer 2/3 pyramidal neurons, the resulting EPSCs in ChR2⁻

187 neurons were not affected by bumetanide (***Figure 2C,D***), indicating that reducing the

188 intracellular chloride concentration only affects chloride-mediated, and not cation-mediated,

189 EPSCs. Finally, bumetanide also diminished the IPSCs resulting from GtACR2 activation in Pv

190 neurons (***Figure 2E,F***). Together, these results demonstrate that light activation of GtACR2

191 generates an excitatory chloride conductance in certain neuronal compartments to trigger

192 neurotransmitter release.

194 **Expression and activation of GtACR2 in adult neurons causes neurotransmitter release**

195 Since we expressed GtACR2 by *in utero* electroporation or neonatal AAV injection, we sought

196 to determine if the excitatory effect of GtACR2 was caused by the long-term expression of

197 GtACR2 throughout development that somehow altered the chloride homeostasis. To selectively

198 express GtACR2 in adult neurons, we *in utero* electroporated a Flpo-dependent plasmid into

199 layer 2/3 pyramidal neurons that will express GtACR2 only if Flpo is present (***Figure 2-***

200 ***supplement 2C,D***). A Flpo-expressing AAV was then injected into the electroporated mice at

201 postnatal week 4 or 9 to turn on the GtACR2 expression. We obtained acute coronal brain slices

202 1–3 weeks after injecting Flpo-expressing AAV and found that light activation of GtACR2,

203 again, produced bumetanide-sensitive EPSCs in GtACR2⁻ neurons (***Figure 2-supplement 2E***).

204 Thus, it is unlikely that GtACR2 expression during neuronal development alters the chloride

205 homeostasis to render GtACR2 excitatory, as acute expression of GtACR2 in mature neurons

206 had the same effect.

Activation of other light-gated chloride channels causes neurotransmitter release

To determine if activation of other light-gated chloride channels can trigger neurotransmitter release, we examined iC++ and iChloC, two engineered blue light-gated chloride channels that were converted from cation channelrhodopsins (Berndt et al., 2015; Wietek et al., 2015), and GtACR1, another natural anion channelrhodopsin from *Guillardia theta* (Govorunova et al., 2015). We *in utero* electroporated plasmids to express iC++ in layer 2/3 pyramidal neurons of the visual cortex and obtained acute coronal brain slices from 3–9-week-old mice. Similar to GtACR2, light activation of iC++ generated EPSCs in iC++$^-$ layer 2/3 pyramidal neurons that were abolished by NBQX and CPP (amplitude reduced by $97.3 \pm 0.9\%$, mean $\pm$ s.e.m., $n = 4$). Bumetanide diminished the iC++-induced EPSCs (***Figure 2G,H***) without affecting the iC++-mediated photocurrents (***Figure 2-supplement 3C,D***). We also expressed iC++ in Pv neurons by injecting a Cre-dependent AAV into *Pvalb*$^{Cre/+}$ mice and found that light activation of iC++ caused bumetanide-sensitive IPSCs in iC++$^-$ layer 2/3 pyramidal neurons (***Figure 2I,J***). Similarly, when we expressed iChloC in Pv neurons, light activation of iChloC resulted in IPSCs in 10 out of 17 recorded iChloC$^-$ layer 2/3 pyramidal neurons (183 ± 42 pA, mean $\pm$ s.e.m., $n = 10$), presumably because iChloC generated smaller photocurrents than iC++ (iChloC, 292 ± 62 pA, $n = 7$; iC++, 2182 ± 291 pA, $n = 15$; recorded at the membrane potential of $+10$ mV; mean $\pm$ s.e.m., $P < 0.0001$, t-test with Welch's correction). Finally, activation of GtACR1 in layer 2/3 pyramidal neurons produced EPSCs onto every recorded GtACR1$^-$ pyramidal neuron (149 ± 47 pA, mean $\pm$ s.e.m., $n = 11$). Altogether, these results demonstrate that activation of different light-gated chloride channels in neurons can trigger neurotransmitter release.

GtACR2 activation directly depolarizes the presynaptic terminals

We hypothesized that the most likely neuronal compartments rendering GtACR2 excitatory were

the distal axon and presynaptic terminals because of the following previous findings. First,

activation of presynaptic $GABA_A$ or glycine receptors enhanced neurotransmitter release at

several synapses of the hippocampus, cerebellum, and brainstem (Jang et al., 2006; Pugh and

Jahr, 2011; Stell et al., 2007; Turecek and Trussell, 2001). Second, the chloride concentrations

were 4–5 times higher in the presynaptic terminals of the Calyx of Held than the parent soma

(Price and Trussell, 2006). Third, there appeared to be an axo-somato-dendritic gradient in which

the reversal potentials of GABA from the axon to the soma and dendrites of cortical neurons

become progressively more negative (Khirug et al., 2008). To test our hypothesis, we expressed

GtACR2 in layer 2/3 pyramidal neurons of the visual cortex in one hemisphere as described

above and obtained acute coronal slices from the contralateral hemisphere (*Figure 3A*). GtACR2

was present in the long-range callosal projections in the contralateral hemisphere (*Figure 3-*

supplement 1A), which enabled us to activate GtACR2 in the axon and presynaptic terminals

that were severed from their parent somas. Light activation of GtACR2 in the callosal

projections generated EPSCs in layer 2/3 pyramidal neurons of the contralateral cortex, which

were diminished by TTX (*Figure 3-supplement 1B,C*) or bumetanide (*Figure 3B*). These results

demonstrate that activation of GtACR2 in the axon and presynaptic terminals is sufficient to

trigger neurotransmitter release.

If GtACR2-mediated chloride currents are excitatory in the presynaptic terminals, then GtACR2

should be similar to ChR2, whose activation can directly depolarize the presynaptic membrane in

the absence of action potentials to trigger neurotransmitter release (Petreanu et al., 2009). To test

this prediction, we recorded EPSCs or IPSCs in GtACR2$^-$ neurons while activating GtACR2 in

layer 2/3 pyramidal neurons or Pv neurons, respectively (*Figure 3C,E*). As described above,

bath application of TTX abolished the EPSCs and IPSCs. However, when we further blocked

voltage-gated potassium channels by 4-aminopyridine (4-AP) and tetraethylammonium (TEA) to

prolong membrane depolarization (Petreanu et al., 2009), the EPSCs and IPSCs were partially

recovered (*Figure 3D,F*). These results indicate that in the absence of action potentials, light

activation of GtACR2 is sufficient to depolarize the presynaptic membrane to open voltage-gated

calcium channels and trigger neurotransmitter release.

We further tested if GtACR2-induced axonal depolarization could evoke antidromic action

potentials by performing extracellular loose-patch or whole-cell current clamp recordings at the

somas of GtACR2$^+$ pyramidal neurons. We observed antidromic spikes in 9 out of 88 neurons

recorded in loose-patch configuration and 21 out of 56 neurons recorded in whole-cell

configuration in response to blue light stimulation (*Figure 3G,H*). In the whole-cell current

clamp recordings, although the chloride concentration in the patch pipette solution sets the

Nernst equilibrium potential of chloride around -85 mV (see Methods), blue light induced a

depolarization following the initial hyperpolarization. This observation is consistent with the

notion that the depolarization antidromically propagated from the distal axon to the soma

(*Figure 3H*). The antidromic spikes were not affected by NBQX and CPP, but were abolished by

TTX (*Figure 3H*), indicating that the spikes were generated within the GtACR2$^+$ neurons, rather

than by excitatory inputs from other neurons. Antidromic spikes were only observed in a subset

of neurons, likely because the GtACR2 expression levels are heterogeneous in different neurons

and the hyperpolarization initiated at the soma can orthodromically propagate to counteract the

antidromic spikes. Similarly, TTX-sensitive antidromic spikes were observed in a subset of

GtACR2$^+$ Pv neurons (3 out of 11 neurons in loose-patch configuration and 7 out of 17 neurons

in whole-cell configuration, *Figure 3I,J*). These results show that GtACR2-induced axonal

depolarization can be sufficient to elicit antidromic action potentials.

Targeting GtACR2 to the somatodendritic domain reduces light-induced neurotransmitter

release

Activating GtACR2 and other light-gated chloride channels inhibits the soma but depolarizes the

presynaptic terminals to release neurotransmitters. This dichotomic effect can confound the

utilization of these channels as inhibitory optogenetic tools for suppressing neuronal activity. We

reasoned that reducing the trafficking of light-gated chloride channels into the axon and

presynaptic terminals should reduce or eliminate their depolarizing action. Thus, we sought to

restrict GtACR2 within the somatodendritic domain of neurons by fusing GtACR2 with a

number of reported somatodendritic targeting motifs including a 26-amino acid Myosin Va-

binding domain of Melanophilin (MBD) (Lewis et al., 2009), a 32-amino acid cytoplasmic C-

terminal motif of Neuroligin 1 (Nlgn1C) (Rosales et al., 2005), a 16-amino acid dileucine-

containing motif of potassium channel Kv4.2 (Kv4.2LL) (Rivera et al., 2003), the N-terminal

150 residues of kainate receptor subunit 2 (KA2N) (Shemesh et al., 2017), the C-terminal 17

residues of Telencephalin (TlcnC) (Mitsui et al., 2005), and a 65-amino acid cytoplasmic C-

terminal motif of potassium channel Kv2.1 (Kv2.1C) (Lim et al., 2000). Each of these GtACR2

variants (*Figure 4-supplement 1*), along with tdTomato, were expressed in layer 2/3 pyramidal

neurons of the visual cortex by *in utero* electroporation (*Figure 4A*). Since GtACR2 was also

tagged with EYFP or EGFP, we compared the EYFP or EGFP fluorescence in layer 5, which

only contains the axons of layer 2/3 pyramidal neurons, with the EYFP or EGFP fluorescence in

layer 2/3 to estimate the distribution of GtACR2 between the axon and somatodendritic domain.

We normalized the EYFP or EGFP fluorescence ratio between layer 5 and layer 2/3 by the

tdTomato fluorescence ratio between layer 5 and layer 2/3 to control for variations in the

collateral axons. Among tested motifs, TlcnC and Kv2.1C were most effective in targeting

GtACR2 to the soma and dendrites (*Figure 4B*). As these two motifs may engage different

trafficking mechanisms, we combined them to create two hybrid motifs, Kv2.1C-TlcnC and

Kv2.1C-linker-TlcnC. Kv2.1C-linker-TlcnC turned out to be the best in restricting GtACR2

within the somatodendritic domain (*Figure 4A,B,C*). Interestingly, GtACR2-EYFP-Kv2.1C and

GtACR2-EYFP-Kv2.1C-linker-TlcnC showed less intracellular aggregation than wild type

GtACR2 (*Figure 4C*), suggesting that the somatodendritic targeting motifs also enhance the

surface expression of GtACR2. Finally, the EYFP fluorescence in the callosal projections in the

contralateral hemisphere was also reduced for GtACR2-EYFP-Kv2.1C-linker-TlcnC as

compared to GtACR2-EYFP (*Figure 4D,E*), demonstrating that Kv2.1C-linker-TlcnC decreased

the trafficking of GtACR2 into the distal axon.

To determine how targeting GtACR2 to the somatodendritic domain affects its photocurrent and

ability to trigger neurotransmitter release, we compared the somatodendritically targeted

GtACR2 variants, GtACR2-EYFP-Kv2.1C or GtACR2-EYFP-Kv2.1C-linker-TlcnC, with wild

type GtACR2 in the same litters of mice. We first recorded GtACR2$^+$ layer 2/3 pyramidal

neurons and found that the blue light-activated photocurrents of GtACR2-EYFP-Kv2.1C or

GtACR2-EYFP-Kv2.1C-linker-TlcnC were 2.1- or 2.7-fold of GtACR2-EYFP photocurrents,

respectively (*Figure 5A–C*). We then recorded the EPSCs in GtACR2$^-$ layer 2/3 pyramidal

neurons in response to different strengths of blue light stimulation. The EPSCs evoked by

activating GtACR2-EYFP-Kv2.1C or GtACR2-EYFP-Kv2.1C-linker-TlcnC were reduced by

52–60% or 65–77%, respectively, as compared to those evoked by activating GtACR2-EYFP

(*Figure 5D–H*). Furthermore, the EPSCs in layer 2/3 pyramidal neurons of the contralateral

cortex were reduced by 73% for GtACR2-EYFP-Kv2.1C-linker-TlcnC as compared to GtACR2-

EYFP when photostimulating the callosal projections (*Figure 5I,J*). These results show that the

somatodendritic targeting motifs, especially Kv2.1C-linker-TlcnC, shift GtACR2 from the axon

towards the soma and dendrites, thereby reducing the excitatory action in the axon and

presynaptic terminals while enhancing the inhibitory currents at the soma and dendrites.

Discussion

Optogenetic suppression of neuronal activity and synaptic outputs is an essential approach for

dissecting the roles of specific neurons in brain functions. Light-gated chloride channels,

particularly GtACR1 and GtACR2, are increasingly used due to their large photocurrents and

high sensitivity to light (Forli et al., 2018; Mardinly et al., 2018; Mauss et al., 2017; Mohamed et

al., 2017; Mohammad et al., 2017). To use these tools to their full potentials, it is necessary that

we understand their function and importantly, their limitations. The experiments here reveal that

wide-field activation of these channels in visual cortical neurons suppresses action potentials at

the soma but also triggers neurotransmitter release at the presynaptic terminals, therefore voiding

inhibition of neuronal activity. As demonstrated, the excitatory action of chloride channels is due

to the higher concentrations of chloride in the axon and presynaptic terminals relative to the

somatodendritic domain. However, it is worth noting that the heterogeneity of chloride gradients

across subcellular compartments may be cell-type dependent. For instance, gramicidin perforated

patch recordings from the axonal blebs of layer 5 pyramidal neurons in the prefrontal cortex of

rats revealed that the reversal potential of $GABA_A$ receptors was more negative than the resting

membrane potential of the axon, which would indicate a hyperpolarizing effect of chloride

channels in this axon (Xia et al., 2014). Thus, it is critical to precisely determine the effect of

light-gated chloride channels, specific to each experimental design, when using them to

manipulate neuronal activity.

To create a better inhibitory optogenetic tool, we tested a number of somatodendritic targeting

motifs to confine GtACR2 in the soma and dendrites. We generated a hybrid motif, Kv2.1C-

linker-TlcnC, that was more effective than the widely used Kv2.1C (Baker et al., 2016; Mardinly

et al., 2018; Wu et al., 2013). The Kv2.1C motif was recently used in a bioRxiv preprint (Mahn

et al., 2017) to target GtACR2 to the somatodendritic domain. This GtACR2 variant achieved a

greater reduction of neurotransmitter release from the contralateral callosal projections of medial

prefrontal cortical neurons than what we observed with our GtACR2-EYFP-Kv2.1C and

GtACR2-EYFP-Kv2.1C-linker-TlcnC in visual cortical neurons. This quantitative difference is

likely due to different expression levels, light stimulation strengths, and neuronal types, which

reiterates the importance of characterizing the specific effect of light-gated chloride channels for

a given experiment.

Although the excitatory effect of light-gated chloride channels is undesired for neuronal

silencing, a potential utilization of their dual actions at the presynaptic terminals and soma is to

activate specific projections of neurons while minimizing the effect of antidromic spikes. For

example, long-range projection neurons often target multiple brain areas, and sometimes it is

desired to selectively excite the axonal terminals projecting to one particular area. If ChR2 is

used, local activation of ChR2-expressing axonal terminals may generate antidromic spikes,

which will affect other projections (Kim et al., 2017). However, if GtACR2 is used, one can

simultaneously activate GtACR2 in the soma and axonal terminals. Axonal depolarization will

result in neurotransmitter release, but the antidromic spikes will be reduced or suppressed by the

hyperpolarization originating from the somatodendritic domain, which reduces the likelihood of

activating other projections.

While GtACR2-EYFP-Kv2.1C-linker-TlcnC can still traffic to the axon to cause

neurotransmitter release, it is thus far the most improved light-gated chloride channel for

optogenetic inhibition. Since it can generate much larger photocurrents in the somatodendritic

domain than what is necessary to suppress action potentials, one approach to use this tool is to

reduce the overall GtACR2 expression level to further decrease its presence in the axon while

generating sufficient inhibitory photocurrents at the soma. Another approach is to selectively

photostimulate GtACR2 at the soma by two-photon excitation (Forli et al., 2018; Mardinly et al.,

2018). However, it is difficult to apply this photoactivation approach to freely moving animals or

deep brain areas. Therefore, it is imperative that we further engineer the channels to eliminate

their excitatory action in the axonal terminals. Future strategies include generating more

effective somatodendritic targeting motifs, creating outwardly rectifying channels, and the

combination of both strategies. The GtACR2-EYFP-Kv2.1C-linker-TlcnC reported here will

serve as the foundation for future improvement, which will enhance the available toolkit for

optogenetic inhibition.

Materials and methods

Mice

All procedures to maintain and use mice were approved by the Institutional Animal Care and

Use Committee at Baylor College of Medicine. Mice were maintained on 14-h:10-h light:dark

cycle with regular mouse chow and water ad libitum. Experiments were performed during the

light cycle. ICR female mice were purchased from Baylor College of Medicine Center for

Comparative Medicine or Charles River Laboratories. C57BL6/J, *Pvalb-2A-Cre*, and *Pvalb-2A-*

Flpo mice were obtained from Jackson Laboratory (stock numbers 000664, 012358, and 022730,

respectively). Both male and female mice were used in the experiments. The mice were used at

the age of 3–9 weeks for experiments, except for the conditional expression of GtACR2 in

adults, where mice were used at the age of 10–12 weeks.

DNA constructs

Plasmids pLenti-UbiC-GtACR2-EYFP (Addgene #67877) and pLenti-UbiC-GtACR1-EYFP

(Addgene #67795) were obtained from Dr. John Spudich, pAAV-CaMKIIα-iC++-TS-EYFP and

pAAV-EF1α-DIO-iC++-TS-EYFP from Dr. Karl Deisseroth, pAAV-EF1α-DIO-iChloC-T2A-

mCherry from Drs. Matthew Caudill and Massimo Scanziani, pCAG-tdTomato from Anirvan

Ghosh, and pCAG-Cre from Addgene (#13775). Plasmid pCAG-Flpo (Addgene #60662) was

previously described (Xue et al., 2014). pCAG-hChR2(H134R)-EYFP was created by replacing

the EGFP in pCAG-EGFP (Addgene #11150) with the hChR2(H134R)-EYFP from pAAV-

EF1α-DIO-hChR2(H134R)-EYFP (Addgene #20298). pAAV-EF1α-DIO-ReaChR-P2A-

dTomato was created by replacing the oChIEF(E163A T199C) in pAAV-EF1α-DIO-

oChIEF(E163A T199C)-P2A-dTomato (Addgene #51094) with the ReaChR from pAAV-hSyn-

FLEX-ReaChR-Citrine (Addgene #50955). pAAV-EF1α-FRT-FLEX-GtACR2-EYFP and

pAAV-EF1α-FRT-FLEX-GtACR1-EYFP were created by replacing the mNaChBac-T2A-

tdTomato in pAAV-EF1α-FRT-FLEX-mNaChBac-T2A-tdTomato (Addgene #60658) with the

GtACR2-EYFP and GtACR1-EYFP from pLenti-UbiC-GtACR2-EYFP and pLenti-UbiC-

GtACR1-EYFP, respectively. Motifs MBD, Nlgn1C, Kv4.2LL, TlcnC, Kv2.1C, Kv2.1C-TlcnC,

and Kv2.1C-linker-TlcnC (see *Figure 4-supplement 1*) were generated by PCR and added to the

C-terminus of the GtACR2-EYFP to create pAAV-EF1α-FRT-FLEX-GtACR2-EYFP-MBD,

pAAV-EF1α-FRT-FLEX-GtACR2-EYFP-Nlgn1C, pAAV-EF1α-FRT-FLEX-GtACR2-EYFP-

Kv4.2LL, pAAV-EF1α-FRT-FLEX-GtACR2-EYFP-TlcnC, pAAV-EF1α-FRT-FLEX-

GtACR2-EYFP-Kv2.1C, pAAV-EF1α-FRT-FLEX-GtACR2-EYFP-Kv2.1C-TlcnC, and pAAV-

EF1α-FRT-FLEX-GtACR2-EYFP-Kv2.1C-linker-TlcnC, respectively. pAAV-EF1α-FRT-

FLEX-GtACR2-KA2N-EGFP was created by replacing the EYFP in pAAV-EF1α-FRT-FLEX-

GtACR2-EYFP with the KA2N-EGFP from pAAV-hSyn-soCoChR-EGFP (Addgene #107708,

obtained from Dr. Edward Boyden).

***In utero* electroporation**

Female ICR mice were crossed with male C57BL6/J, *Pvalb-2A-Cre*, or *Pvalb-2A-Flpo* mice to

obtain timed pregnancies. *In utero* electroporation was performed as previously described (Xue

et al., 2014) with a square-wave pulse generator (Gemini X2, BTX Harvard Bioscience). To

express GtACR2, GtACR1, iC++, or ChR2 in layer 2/3 pyramidal neurons, pLenti-UbiC-

GtACR2-EYFP, pLenti-UbiC-GtACR1-EYFP, pAAV-CaMKIIα-iC++-TS-EYFP, or pCAG-

hChR2(H134R)-EYFP (all 2μg/μl) was used, respectively. In a few experiments, pAAV-EF1α-

FRT-FLEX-GtACR2-EYFP (2 μg/μl) with pCAG-Flpo (0.2 μg/μl), pAAV-EF1α-FRT-FLEX-

GtACR1-EYFP (2 μg/μl) with pCAG-Flpo (0.2 μg/μl), or pAAV-EF1α-DIO-iC++-TS-EYFP (2

μg/μl) with pCAG-Cre (0.2 μg/μl) was used to express GtACR2, GtACR1, or iC++,

respectively. To express somatodendritically targeted GtACR2 variants in layer 2/3 pyramidal

neurons and compare them with wild type GtACR2, the pAAV-EF1α-FRT-FLEX constructs

described above were used (all 2 μg/μl) with pCAG-Flpo (0.2 μg/μl). pCAG-tdTomato

(0.1μg/μl) was included in all experiments. The plasmid concentrations stated above were final

concentrations in the plasmid mix. Transfected pups were identified by the transcranial

fluorescence of tdTomato with a MZ10F stereomicroscope (Leica) 1–2 days after birth.

AAV production and injection

All recombinant AAV serotype 9 vectors were produced by the Baylor College of Medicine

Gene Vector Core except AAV9-hSyn-Flpo (Addgene #60663), which was produced by the

Penn Vector Core and was previously described (Xue et al., 2014). To express GtACR2,

ReaChR, iC++, or iChloC in Pv neurons, 200–250 nl of the following recombinant AAV

serotype 9 vectors at their respective titer were injected into the visual cortex of $Pvalb^{Flpo/+}$ (for

GtACR2) or $Pvalb^{Cre/+}$ (for ReaChR, iC++, or iChloC) mice at postnatal day 1 as previously

described (Xue et al., 2014): AAV9-EF1α-FRT-FLEX-GtACR2-EYFP (3.8×10^{13} genome

copies/ml), AAV9-EF1α-DIO-ReaChR-P2A-dTomato (7.0×10^{13} genome copies/ml), AAV9-

EF1α-DIO-iC++-TS-EYFP (3.71×10^{13} genome copies/ml), and AAV9-EF1α-DIO-iChloC-2A-

mCherry (3.7×10^{14} genome copies/ml). To conditionally express GtACR2 in juvenile and adult

neurons, mice previously electroporated with plasmid pAAV-EF1α-FRT-FLEX-GtACR2-EYFP

into layer 2/3 pyramidal neurons were injected with 200 nl of AAV9-hSyn-Flpo (1.2×10^{12}

genome copies/ml) at postnatal day 23, 60, or 64. Injection was performed as previously

described (Xue et al., 2014) with an UltraMicroPump III and a Micro4 controller (World

Precision Instruments).

Brain slice electrophysiology

Mice were anesthetized by an intraperitoneal injection of a ketamine and xylazine mix (80 mg/kg

and 16 mg/kg, respectively) and transcardially perfused with cold (0–4°C) slice cutting solution

containing 80 mM NaCl, 2.5 mM KCl, 1.3 mM NaH_2PO_4, 26 mM $NaHCO_3$, 4 mM $MgCl_2$, 0.5

mM $CaCl_2$, 20 mM D-glucose, 75 mM sucrose and 0.5 mM sodium ascorbate (315 mosmol, pH

7.4, saturated with 95% O_2/5% CO_2). Brains were removed and sectioned in the cutting solution

with a VT1200S vibratome (Leica) to obtain 300-μm coronal slices. Slices were incubated in a

custom-made interface holding chamber saturated with 95% O_2/5% CO_2 at 34 °C for 30 min and

then at room temperature for 20 min to 10 h until they were transferred to the recording chamber.

Recordings were performed on submerged slices in artificial cerebrospinal fluid (ACSF)

containing 119 mM NaCl, 2.5 mM KCl, 1.3 mM NaH_2PO_4, 26 mM $NaHCO_3$, 1.3 mM $MgCl_2$,

2.5 mM $CaCl_2$, 20 mM D-glucose and 0.5 mM sodium ascorbate (305 mosmol, pH 7.4, saturated

with 95% O_2/5% CO_2, perfused at 3 ml/min) at 30–32°C. For whole-cell recordings, we used a

K^+-based pipette solution containing 142 mM K^+-gluconate, 10 mM HEPES, 1 mM EGTA, 2.5

mM $MgCl_2$, 4 mM ATP-Mg, 0.3 mM GTP-Na, 10 mM Na_2-phosphocreatine (295 mosmol, pH

7.35) or a Cs^+-based pipette solution containing 121 mM Cs^+-methanesulfonate, 1.5 mM $MgCl_2$,

10 mM HEPES, 10 mM EGTA, 4 mM Mg-ATP, 0.3 mM Na-GTP, 10 mM Na_2-Phosphocreatine,

and 2 mM QX314-Cl (295 mosmol, pH 7.35). Membrane potentials were not corrected for liquid

junction potential (experimentally measured as 12.5 mV for the K^+-based pipette solution and

9.5 mV for the Cs^+-based pipette solution).

Neurons were visualized with video-assisted infrared differential interference contrast imaging

and fluorescent neurons were identified by epifluorescence imaging under a water immersion

objective (40×, 0.8 numerical aperture) on an upright SliceScope Pro 1000 microscope

(Scientifica) with an infrared IR-1000 CCD camera (DAGE-MTI). Data were low-pass filtered at

4 kHz and acquired at 10 kHz with an Axon Multiclamp 700B amplifier and an Axon Digidata

1550 Data Acquisition System under the control of Clampex 10.7 (Molecular Devices). Data

were analyzed offline using AxoGraph X (AxoGraph Scientific). For the photostimulation of

GtACR2-, iC++-, iChloC-, or ChR2-expressing neurons, blue light was emitted from a

collimated light-emitting diode (LED) of 455 nm, whereas for the photostimulation of GtACR1-

or ReaChR-expressing neurons, red light was emitted from a LED of 617 nm. The LEDs were

driven by a LED driver (Mightex) under the control of an Axon Digidata 1550 Data Acquisition

System and Clampex 10.7. Light was delivered through the reflected light fluorescence

illuminator port and the 40× objective.

Synaptic currents and photocurrents were recorded in the whole-cell voltage clamp mode with

the Cs^+-based patch pipette solution. Only recordings with series resistance below 20 MΩ were

included. EPSCs and IPSCs were recorded at the reversal potential for IPSCs (-60 mV) and

EPSCs (+10 mV), respectively, unless stated otherwise. Photocurrents were recorded at +10 mV

unless stated otherwise. For light pulse stimulation, pulse duration (0.5–10 ms) and intensity

(2.5–23.6 mW/mm^2) were adjusted for each recording to evoke small (to minimize voltage-

clamp errors) but reliable monosynaptic EPSCs or IPSCs. Disynaptic IPSCs were evoked using

the same light pulses that were used for evoking the corresponding monosynaptic EPSCs. Light

pulses were delivered at 30-s interstimulus intervals. Antidromic spikes in GtACR2$^+$ neurons

were recorded with the K+-based patch pipette solution in whole-cell current clamp mode or

with ACSF as the patch pipette solution in the loose-patch current clamp mode.

For pharmacology experiments, the baseline synaptic currents were recorded for at least 3

minutes in the absence of any drug. The drugs were then added to the ACSF at the following

concentrations: TTX (1 µM), NBQX (10 µM), (*RS*)-CPP (10 µM), SR95531 (Gabazine, 10 µM),

ZD7288 (20 µM), bumetanide (50 or 100 µM), TEA (1.5 mM), and 4-AP (1.5 mM). The

synaptic currents were recorded for at least 3 minutes in the presence of drugs. For ZD7288,

which did not inhibit GtACR2-induced neurotransmitter release, the efficacy of the drug was

monitored by examining the I_h current of cortical layer 5 pyramidal neurons.

Fluorescent microscopy

Fluorescent images were taken from live brain slices, except for the conditional expression of

GtACR2 in adults, where images were taken from fixed brain slices. Live brain slices were

prepared as described for slice electrophysiology. For the fixed brain slices, mice were

anesthetized by an intraperitoneal injection of a ketamine and xylazine mix (80 mg/kg and 16

mg/kg, respectively) and transcardially perfused with phosphate buffered saline (PBS, pH 7.4)

followed by 4% paraformaldehyde in PBS (pH 7.4). Brains were further fixed overnight in 4%

paraformaldehyde, cryoprotected with 30% sucrose in PBS, and frozen in optimum cutting-

temperature medium until sectioning. A HM 450 Sliding Microtome (Thermo Scientific) was

used to section the brains to obtain 30–50-mm coronal slices. Images were acquired on an Axio

Zoom.V16 Fluorescence Stereo Zoom Microscope (Zeiss) and processed using National

Institutes of Health ImageJ.

To determine the EYFP (or EGFP) fluorescence ratio between layer 5 and layer 2/3, one or two

350 μm-wide rectangular regions that were perpendicular to the pia and spanned all 6 cortical

layers were selected in the most transfected regions of each slice. The mean EYFP (or EGFP)

fluorescence was measured for layer 5 and layer 2/3 within the selected area. The mean

tdTomato fluorescence was measured similarly for layer 5 and layer 2/3. The mean background

fluorescence was measured from a nearby rectangular region (140.5 μm by 90.8 μm) where no

cellular EYFP (or EGFP) and tdTomato fluorescence was present. The normalized EYFP (or

EGFP) fluorescence ratio between layer 5 and layer 2/3 was calculated by

$\dfrac{Layer\ 5_{EYFP} - Background_{EYFP}}{Layer\ 2/3_{EYFP} - Background_{EYFP}} \Big/ \dfrac{Layer\ 5_{tdTomato} - Background_{tdTomato}}{Layer\ 2/3_{tdTomato} - Background_{tdTomato}}.$

To determine the ratio of EYFP fluorescence to tdTomato fluorescence in the callosal

projections, one or two rectangular regions that contained the tdTomato-labeled axons were

selected in each slice to measure the mean EYFP and tdTomato fluorescence. The mean

background fluorescence was measured in a nearby cortical area spanning the same layers. The

ratio of EYFP fluorescence to tdTomato fluorescence was calculated by

$\dfrac{Fluorescence_{EYFP} - Background_{EYFP}}{Fluorescence_{tdTomato} - Background_{tdTomato}}.$

Statistics

All reported sample numbers (*n*) represent biological replicates that are the numbers of recorded

neurons for electrophysiology or the numbers of analyzed regions of interest (ROI) for

fluorescent images. Statistical analyses were performed with Prism 7 (GraphPad Software). We

first determined whether the data were normally distributed by performing the D'Agostino &

Pearson test, Shapiro-Wilk test, and KS test. If all data within one experiment passed all three

normality tests, we then performed the statistical test that assumes a Gaussian distribution.

Otherwise, we performed the statistical test that assumed a non-Gaussian distribution. All

statistical tests were two-tailed with an alpha of 0.05.

Wilcoxan matched-pairs signed rank test was used for Figure 1F,K; Figure 2B,D,F,J; Figure 2-

supplement 2B,E; Figure 2-supplement 3B (EPSCs); Figure 3B; and Figure 3-supplement 1C.

Paired *t* test was used for Figure 1G,J; Figure 2H; Figure 2-supplement 1D; and Figure 2-

supplement 3B (photocurrents),D. *t* test with Welch's correction was used for Figure 4E and

Figure 5C (WT vs. Kv2.1C). Mann-Whitney test was used for Figure 5C (WT vs. Kv2.1C-

linker-TlcnC),F,G,H,J. One-way ANOVA with Greenhouse-Geisser correction and Tukey

multiple comparisons was used for Figure 3D,F. Kruskal-Wallis test with Dunn's multiple

comparisons was used for Figure 4B.

Acknowledgments

We thank Elena Govorunova and John Spudich for discussions and the pLenti-UbiC-GtACR2-

EYFP and pLenti-UbiC-GtACR1-EYFP plasmids, Anirvan Ghosh for the pCAG-tdTomato

plasmid, Karl Deisseroth for the pAAV-CaMKIIα-iC++-TS-EYFP and pAAV-EF1α-DIO-iC++-

TS-EYFP plasmids, Matthew Caudill and Massimo Scanziani for the pAAV-EF1α-DIO-iChloC-

T2A-mCherry plasmid, Zhuo-Hua Pan for the pAAV-EF1α-DIO-hChR2(H134R)-EYFP-

Kv2.1C plasmid, Edward Boyden for the pAAV-hSyn-soCoChR-EGFP plasmid, and Hsiao-

Tuan Chao and Matthew Caudill for comments on an earlier version of the manuscript. This

work was supported in part by a Whitehall Foundation Research Grant (#2015-05-54, to M.X.).

J.E.M is part of the Baylor College of Medicine Medical Scientist Training Program and a

McNair M.D./Ph.D. Student Scholar supported by the McNair Medical Institute at the Robert

and Janice McNair Foundation. M.X. is a Caroline DeLuca Scholar.

Competing interests

The authors declare no competing financial interests.

Figures and legends

Figure 1

Figure 1. Light activation of GtACR2 causes excitatory and inhibitory neurons to release

neurotransmitter.

(**A**) A representative fluorescent image of the visual cortex showing GtACR2-EYFP and

tdTomato expression in a subset of layer 2/3 pyramidal neurons. Note the strong EYFP

fluorescence in layer 5 that contains the axons of layer 2/3 pyramidal neurons. L, layer; WM,

white matter.

(**B**) Schematic of slice experiments in (C,D). GtACR2 in a subset of layer 2/3 pyramidal neurons.

(**C**) Membrane potentials (upper panel) in response to somatic current injections (lower panel)

from a GtACR2$^+$ neuron. Blue light activation of GtACR2 suppressed the action potentials

evoked by current injections ($n = 5$).

(**D**) Blue light-induced membrane currents recorded at membrane potentials of -75, -77, -78, -79,

and -80 mV from the same GtACR2$^+$ neuron in (C). Note the EPSC-like inward currents

(arrows) superimposed on the GtACR2-mediated photocurrents ($n = 2$).

(**E**) Schematic of slice experiments in (F,G). GtACR2 in a subset of layer 2/3 pyramidal neurons.

(**F**) Left, photoactivation of GtACR2 generated EPSCs in a GtACR2$^-$ neuron, which were

abolished by the glutamate receptor antagonists, NBQX and CPP. Right, summary graph of

similar experiments ($n = 17$, $P < 0.0001$).

(**G**) Left, photoactivation of GtACR2 generated EPSCs in a GtACR2$^-$ neuron, which were

abolished by the voltage-gated sodium channel blocker, TTX. Right, summary graph of similar

experiments ($n = 8$, $P = 0.02$).

(**H**) A representative fluorescent image of the visual cortex showing GtACR2-EYFP expression

in Pv neurons.

(**I,J,K**) As in (E,F,G), but for GtACR2 in Pv neurons. GtACR2 activation-induced IPSCs were

abolished by the GABA$_A$ receptor antagonist, Gabazine (J, $n = 7$, $P = 0.02$) or TTX (K, $n = 10$, P

$= 0.002$).

Figure 1-supplement 1

Figure 1-supplement 1. Recruitment of inhibitory neurons by activating GtACR2 in excitatory neurons.

Left, schematic of slice experiments. GtACR2 in a subset of layer 2/3 pyramidal neurons. Right, photoactivation of GtACR2 generated an EPSC (inward current) and IPSC (outward current) in a GtACR2$^-$ neuron. The IPSC was abolished by the glutamatergic receptor antagonists, NBQX and CPP (red trace), indicating its disynaptic nature ($n = 3$).

Figure 1-supplement 2

Figure 1-supplement 2. Repetitive activation of GtACR2 reliably causes neurotransmitter

release.

(**A**) Left, schematic of slice experiments. GtACR2 in a subset of layer 2/3 pyramidal neurons.

Right, a 20-Hz train of blue-light pulses activated GtACR2 to generate EPSCs in a GtACR2⁻

neuron (*n* = 11).

(**B**) As in (A), but for GtACR2 in Pv neurons. A 20-Hz train of blue-light pulses activated

GtACR2 to generate IPSCs in a GtACR2⁻ neuron (*n* = 14).

Figure 2

Figure 2. Reducing intracellular chloride concentrations diminishes the neurotransmitter

release induced by activation GtACR2 or iC++.

(**A**) Schematic of slice experiments in (B). GtACR2 in a subset of layer 2/3 pyramidal neurons.

(**B**) Left, photoactivation of GtACR2 generated EPSCs in a GtACR2⁻ neuron, which were

diminished by decreasing intracellular chloride concentrations with an NKCC1 blocker,

bumetanide. Right, summary graph of similar experiments ($n = 10$, $P = 0.002$).

(C,D) As in (A,B), but for ChR2. ChR2 activation-induced EPSCs were not affected by

bumetanide ($n = 8$, $P = 0.9$).

(E,F) As in (A,B), but for GtACR2 in Pv neurons. GtACR2 activation-induced IPSCs were

diminished by bumetanide ($n = 9$, $P = 0.004$).

(G,H) As in (A,B), but for iC++. iC++-induced EPSCs were diminished by bumetanide ($n = 6$, P

$= 0.009$).

(I,J) As in (E,F), but for iC++. iC++-induced IPSCs were diminished by bumetanide ($n = 8$, $P =$

0.008).

Figure 2-supplement 1

Figure 2-supplement 1. The reversal potential of GtACR2 at the soma is similar to that of

GABA$_A$ receptor.

(**A**) Schematic of slice experiments in (B). GtACR2 in a subset of layer 2/3 pyramidal neurons

and ReaChR in Pv neurons.

(**B**) In a GtACR2$^+$ neuron, GABAergic IPSCs induced by activating ReaChR in Pv neurons (left

panel) and GtACR2-mediated photocurrents (right panel) were sequentially recorded at different

membrane potentials. The example traces show the currents at -50, -60, -65, -66, -68, -70, and -

75 mV. A GtACR2$^-$ neuron was simultaneously recorded at the membrane potential of +10 mV

to ensure that the blue light did not generate IPSCs (right panel).

(**C**) The amplitudes of IPSCs and GtACR2 photocurrents were plotted as a function of the

membrane potentials for the GtACR2$^+$ neuron in (B). Each set of data were fit with linear

regression and the reversal potentials were determined by the V_m-axis intercepts.

(**D**) Summary graph of similar experiments in (B,C) where the reversal potential of IPSCs was

plotted against that of GtACR2 photocurrents for each GtACR2$^+$ neuron. All data points fall

close to the dotted unity line and there is no significant difference between the reversal potentials

of IPSCs and photocurrents ($n = 8$, $P = 0.4$).

Figure 2-supplement 2

Figure 2-supplement 2. GtACR2-induced neurotransmitter release is not due to rebound

depolarization or its long-term expression throughout development.

(**A**) Schematic of slice experiments in (B). GtACR2 in a subset of layer 2/3 pyramidal neurons.

(**B**) Left, photoactivation of GtACR2 generated EPSCs in a GtACR2⁻ neuron, which were not

decreased by blocking the I_h current with ZD7288 (20 μM). Right, summary graph of similar

experiments showing that the EPSC amplitudes were slightly increased by ZD7288 ($n = 12$, $P =$

0.03).

(**C**) Schematics of conditional expression of GtACR2 in a subset of layer 2/3 pyramidal neurons.

A Flpo-dependent plasmid (FRT-FLEX-GtACR2) was electroporated at embryonic day 15 (E15)

and AAV9-hSyn-Flpo was injected into the electroporated mice at different postnatal days (P23,

P60, or P64). Green bars indicate the approximate GtACR2 expression time windows.

(**D**) Left, representative fluorescent images of the visual cortex obtained at postnatal day 73

showing that without injection of AAV9-hSyn-Flpo, transfected neurons (labeled by tdTomato)

did not express GtACR2-EYFP (left panel, $n = 5$ mice). Note that the image from the EYFP

channel was overexposed to show no EYFP$^+$ neurons. Right, representative fluorescent images

of the visual cortex obtained at postnatal day 40 showing that the expression of GtACR2-EYFP

in transfected neurons was turned on by the injection of AAV9-hSyn-Flpo at postnatal day 23

(right panel, $n = 7$ mice).

(**E**) Photoactivation of GtACR2 generated EPSCs in GtACR2$^-$ neurons when GtACR2 was

expressed during 3 different time windows as indicated in (C). Bumetanide was applied in a

subset of experiments and the EPSCs were diminished ($n = 12$, $P = 0.0005$).

Figure 2-supplement 3

Figure 2-supplement 3. Bumetanide does not affect GtACR2 and iC++-mediated

photocurrents.

(**A**) Schematic of slice experiments in (B). GtACR2 in a subset of layer 2/3 pyramidal neurons.

(**B**) Left, photoactivation of GtACR2 generated photocurrents in a GtACR2$^+$ neuron (top panel,

recorded at the membrane potential of +10 mV) and EPSCs in a simultaneously recorded

GtACR2$^-$ neuron (bottom panel, recorded at the membrane potential of -60 mV). Bumetanide

diminished the EPSCs (circle symbols) without affecting the photocurrents (square symbols).

Right, summary graph of similar experiments ($n = 6$; EPSCs, $P = 0.03$; photocurrents, $P = 0.3$).

(**C,D**) As in (A,B), but for iC++. Bumetanide diminished the EPSCs without affecting the

photocurrents ($n = 3$; EPSCs, $P = 0.03$; photocurrents, $P = 0.7$).

Figure 3

Figure 3. GtACR2 activation directly depolarizes the distal axon and presynaptic terminals

and can result in antidromic spikes.

(**A**) Schematic of slice experiments in (B). GtACR2 in a subset of layer 2/3 pyramidal neurons in

the contralateral hemisphere.

(**B**) Left, photoactivation of GtACR2 in the callosal axons, severed from their somas, generated

EPSCs in a GtACR2⁻ neuron, which were diminished by bumetanide. Right, summary graph of

similar experiments ($n = 15$, $P < 0.0001$).

(**C**) Schematic of slice experiments in (D). GtACR2 in a subset of layer 2/3 pyramidal neurons.

(**D**) Left, photoactivation of GtACR2 generated EPSCs in a GtACR2$^-$ neuron, which were

abolished by TTX, but partially recovered by the addition of voltage-gated potassium channel

blockers, 4-AP (1.5 mM) and TEA (1.5 mM). Right, summary graph of similar experiments ($n =$

10; TTX vs. control, $P = 0.02$, average EPSC amplitude in TTX was 2% of control; TTX + 4-AP

+ TEA vs. TTX, $P = 0.0004$, average EPSC amplitude in TTX + 4-AP + TEA was 35% of

control).

(**E,F**) As in (C,D), but for Pv neurons. GtACR2 activation-induced IPSCs were abolished by

TTX, but partially recovered by 4-AP and TEA ($n = 9$; TTX vs. control, $P = 0.006$, average IPSC

amplitude in TTX was 0.9% of control; TTX + 4-AP + TEA vs. TTX, $P = 0.008$, average IPSC

amplitude in TTX + 4-AP + TEA was 23% of control).

(**G**) Schematic of slice experiments in (H). GtACR2 in a subset of layer 2/3 pyramidal neurons.

(**H**) Photoactivation of GtACR2 generated antidromic spikes in GtACR2$^+$ pyramidal neurons,

which were not affected by NBQX and CPP, but were abolished by TTX in both loose-patch

(left panel) and whole-cell (right panel) recordings. In the whole-cell recordings, the resting

membrane potentials of those neurons that generated antidromic spikes are -68.1 ± 1.7 mV

(mean ± s.e.m., $n = 21$).

(**I,J**) As in (G,H), but for Pv neurons. The antidromic spikes in GtACR2$^+$ Pv neurons were

abolished by TTX. In the whole-cell recordings, the resting membrane potentials of those

neurons that generated antidromic spikes are -60.0 ± 2.4 mV (mean ± s.e.m., $n = 7$).

Figure 3-supplement 1

Figure 3-supplement 1. Activation of GtACR2 in the callosal axons causes

neurotransmitter release.

(**A**) GtACR2-EYFP and tdTomato were expressed in a subset of layer 2/3 pyramidal neurons in

the visual cortex of one hemisphere. Representative fluorescent images of the contralateral

hemisphere showing GtACR2-EYFP (left panel) in the callosal projections labeled by tdTomato

(right panel). L, layer; WM, white matter.

(**B**) Schematic of slice experiments in (C). GtACR2-EYFP in a subset of layer 2/3 pyramidal

neurons in the contralateral hemisphere.

(**C**) Left, photoactivation of GtACR2 in the callosal axons generated EPSCs in a GtACR2⁻

neuron, which were abolished by TTX. Right, summary graph of similar experiments ($n = 9$, $P =$

0.004).

Figure 4

Figure 4. Targeting GtACR2 to neuronal somatodendritic domain.

(**A**) Wild type (WT) GtACR2 and its variants tagged with EYFP or EGFP were expressed along

with tdTomato in a subset of layer 2/3 pyramidal neurons. Representative fluorescent images of

the cortices showing the distribution of GtACR2. Note the strong EYFP fluorescence in layer 5

for WT GtACR2 and weak EYFP fluorescence for some somatodendritically targeted GtACR2

variants (e.g., Kv2.1C-linker-TlcnC).

(**B**) The EYFP or EGFP fluorescence ratio between layer 5 and layer 2/3 was normalized by the

tdTomato fluorescence ratio between layer 5 and layer 2/3. A reduction in the normalized EYFP

or EGFP fluorescence ratio indicates a shift of the EYFP or EGFP distribution from the axon to

somatodendritic domain.

(**C**) Representative fluorescent images of electroporated neurons showing that GtACR2-EYFP-

Kv2.1C and GtACR2-EYFP-Kv2.1C-linker-TlcnC were more concentrated on the membranes at

the soma and proximal dendrites (arrows) than GtACR2-EYFP. Note the intracellular

aggregation of GtACR2-EYFP (arrow head).

(**D**) Representative fluorescent images of the contralateral hemisphere showing less GtACR2-

EYFP-Kv2.1C-linker-TlcnC in the callosal projections than GtACR2-EYFP. The callosal

projections were labeled by tdTomato.

(**E**) The ratio of EYFP fluorescence to tdTomato fluorescence in the callosal projections in the

contralateral hemisphere.

The numbers of analyzed slices were indicated in the panel. The columns and error bars are

mean $\pm$ s.e.m. ** $P < 0.01$, **** $P < 0.0001$.

Figure 4-supplement 1. Sequences of somatodendritic targeting motifs.

Black, amino acid sequences of somatodendritic targeting motifs; blue, GtACR2; green, EYFP

or EGFP; magenta, linker.

Motif	Sequence
MBD	*GtACR2*-*EYFP*-GSGSGTRGSGS- RDQPLNSKKKKRLLSFRDVDFEEDSD
Nlgn1C	*GtACR2*-*EYFP*-VVLRTACPPDYTLAMRRSPDDIPLMTPNTITM
Kv4.2LL	*GtACR2*-*EYFP*-FETQHHHLLHCLEKTT
KA2N	*GtACR2*-GGSGGTGGSGGT- MPAELLLLLIVAFANPSCQVLSSLRMAAILDDQTVCGRGERLA LALAREQINGIIEVPAKARVEVDIFELQRDSQYETTDTMCQILP KGVVSVLGPSSSPASASTVSHICGEKEIPHIKVGPEETPRLQYLR FASVSLYPSNEDVSLAVS-GASGGT-*EGFP*
TlcnC	*GtACR2*-*EYFP*-AESPADGEVFAIQLTSS
Kv2.1C	*GtACR2*-*EYFP*- QSQPILNTKEMAPQSKPPEELEMSSMPSPVAPLPARTEGVIDMR SMSSIDSFISCATDFPEATRF
Kv2.1C-TlcnC	*GtACR2*-*EYFP*- QSQPILNTKEMAPQSKPPEELEMSSMPSPVAPLPARTEGVIDMR SMSSIDSFISCATDFPEATRF-AESPADGEVFAIQLTSS
Kv2.1C-linker-TlcnC	*GtACR2*-*EYFP*- QSQPILNTKEMAPQSKPPEELEMSSMPSPVAPLPARTEGVIDMR SMSSIDSFISCATDFPEATRF-GSGSGSGSGS- AESPADGEVFAIQLTSS

Figure 5

Figure 5. Activation of somatodendritically targeted GtACR2 variants generate larger photocurrents but cause less neurotransmitter release than wild type GtACR2.

(**A**) Schematic of slice experiments in (B,C). GtACR2-EYFP, GtACR2-EYFP-Kv2.1C, or GtACR2-EYFP-Kv2.1C-linker-TlcnC in a subset of layer 2/3 pyramidal neurons.

(**B**) Somatic photocurrents recorded at the membrane potential of +10 mV from GtACR2-EYFP-, GtACR2-EYFP-Kv2.1C-, or GtACR2-EYFP-Kv2.1C-linker-TlcnC-expressing neurons in response to a blue light pulse of 1-ms and 2.5-mW/mm^2.

(**C**) Summary graph showing the photocurrent peak amplitudes of GtACR2-EYFP-Kv2.1C and GtACR2-EYFP-Kv2.1C-linker-TlcnC were larger than those of GtACR2-EYFP.

(**D,E**) As in (A,B), but for recording EPSCs in GtACR2$^-$ layer 2/3 pyramidal neurons in response to a blue light pulse of 5-ms and 2.9-mW/mm^2.

(**F,G,H**) Summary graphs showing the EPSC amplitudes in response to a blue light pulse of 1-

ms and 2.5 mW/mm^2 (F), 5-ms and 2.9 mW/mm^2 (G), or 5-ms and 16.1 mW/mm^2 (H).

(**I**) Schematic of slice experiments in (J). GtACR2-EYFP or GtACR2-EYFP-Kv2.1C-linker-

TlcnC in a subset of layer 2/3 pyramidal neurons in the contralateral hemisphere.

(**J**) Summary graphs showing the EPSC amplitudes in response to a blue light pulse of 5-ms and

23.6 mW/mm^2.

The numbers of recorded neurons were indicated in the panel. The columns and error bars are

mean ± s.e.m. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$, **** $P < 0.0001$.

References

Allen, B.D., Singer, A.C., Boyden, E.S., 2015. Principles of designing interpretable optogenetic
behavior experiments. Learn. Mem. 22, 232–238. doi:10.1101/lm.038026.114
Baker, C.A., Elyada, Y.M., Parra, A., Bolton, M.M., 2016. Cellular resolution circuit mapping
with temporal-focused excitation of soma-targeted channelrhodopsin. Elife 5, 11981.
doi:10.7554/eLife.14193
Ben-Ari, Y., 2017. NKCC1 Chloride Importer Antagonists Attenuate Many Neurological and
Psychiatric Disorders. Trends Neurosci 40, 536–554. doi:10.1016/j.tins.2017.07.001
Ben-Ari, Y., 2002. Excitatory actions of gaba during development: the nature of the nurture. Nat
Rev Neurosci 3, 728–739. doi:10.1038/nrn920
Berndt, A., Lee, S.Y., Ramakrishnan, C., Deisseroth, K., 2014. Structure-guided transformation
of channelrhodopsin into a light-activated chloride channel. Science 344, 420–424.
doi:10.1126/science.1252367
Berndt, A., Lee, S.Y., Wietek, J., Ramakrishnan, C., Steinberg, E.E., Rashid, A.J., Kim, H., Park,
S., Santoro, A., Frankland, P.W., Iyer, S.M., Pak, S., Ährlund-Richter, S., Delp, S.L.,
Malenka, R.C., Josselyn, S.A., Carlén, M., Hegemann, P., Deisseroth, K., 2015. Structural
foundations of optogenetics: Determinants of channelrhodopsin ion selectivity. Proc Natl
Acad Sci USA. doi:10.1073/pnas.1523341113
Boyden, E.S., Zhang, F., Bamberg, E., Nagel, G., Deisseroth, K., 2005. Millisecond-timescale,
genetically targeted optical control of neural activity. Nat Neurosci 8, 1263–1268.
doi:10.1038/nn1525
Chow, B.Y., Han, X., Dobry, A.S., Qian, X., Chuong, A.S., Li, M., Henninger, M.A., Belfort,
G.M., Lin, Y., Monahan, P.E., Boyden, E.S., 2010. High-performance genetically targetable
optical neural silencing by light-driven proton pumps. Nature 463, 98–102.
doi:10.1038/nature08652
Chuong, A.S., Miri, M.L., Busskamp, V., Matthews, G.A.C., Acker, L.C., Sørensen, A.T.,
Young, A., Klapoetke, N.C., Henninger, M.A., Kodandaramaiah, S.B., Ogawa, M.,

Ramanlal, S.B., Bandler, R.C., Allen, B.D., Forest, C.R., Chow, B.Y., Han, X., Lin, Y., Tye,
K.M., Roska, B., Cardin, J.A., Boyden, E.S., 2014. Noninvasive optical inhibition with a
red-shifted microbial rhodopsin., Nature neuroscience. doi:10.1038/nn.3752
Forli, A., Vecchia, D., Binini, N., Succol, F., Bovetti, S., Moretti, C., Nespoli, F., Mahn, M.,
Baker, C.A., Bolton, M.M., Yizhar, O., Fellin, T., 2018. Two-Photon Bidirectional Control
and Imaging of Neuronal Excitability with High Spatial Resolution In Vivo. Cell Rep 22,
3087–3098. doi:10.1016/j.celrep.2018.02.063
Govorunova, E.G., Sineshchekov, O.A., Janz, R., Liu, X., Spudich, J.L., 2015. Natural light-
gated anion channels: A family of microbial rhodopsins for advanced optogenetics. Science
349, 647–650. doi:10.1126/science.aaa7484
Govorunova, E.G., Sineshchekov, O.A., Li, H., Spudich, J.L., 2017a. Microbial Rhodopsins:
Diversity, Mechanisms, and Optogenetic Applications. Annu. Rev. Biochem. 86, 845–872.
doi:10.1146/annurev-biochem-101910-144233
Govorunova, E.G., Sineshchekov, O.A., Rodarte, E.M., Janz, R., Morelle, O., Melkonian, M.,
Wong, G.K.S., Spudich, J.L., 2017b. The Expanding Family of Natural Anion
Channelrhodopsins Reveals Large Variations in Kinetics, Conductance, and Spectral
Sensitivity. Sci Rep 7, 43358. doi:10.1038/srep43358
Han, X., Boyden, E.S., 2007. Multiple-color optical activation, silencing, and desynchronization
of neural activity, with single-spike temporal resolution. PLoS ONE 2, e299.
doi:10.1371/journal.pone.0000299
Han, X., Chow, B.Y., Zhou, H., Klapoetke, N.C., Chuong, A., Rajimehr, R., Yang, A., Baratta,
M.V., Winkle, J., Desimone, R., Boyden, E.S., 2011. A high-light sensitivity optical neural
silencer: development and application to optogenetic control of non-human primate cortex.
Front. Syst. Neurosci. 5, 18. doi:10.3389/fnsys.2011.00018
Jang, I.S., Nakamura, M., Ito, Y., Akaike, N., 2006. Presynaptic GABAA receptors facilitate
spontaneous glutamate release from presynaptic terminals on mechanically dissociated rat
CA3 pyramidal neurons. Neuroscience 138, 25–35. doi:10.1016/j.neuroscience.2005.11.001
Khirug, S., Yamada, J., Afzalov, R., Voipio, J., Khiroug, L., Kaila, K., 2008. GABAergic
Depolarization of the Axon Initial Segment in Cortical Principal Neurons Is Caused by the
Na-K-2Cl Cotransporter NKCC1. J Neurosci 28, 4635–4639.
doi:10.1523/JNEUROSCI.0908-08.2008
Kim, C.K., Adhikari, A., Deisseroth, K., 2017. Integration of optogenetics with complementary
methodologies in systems neuroscience. Nat Rev Neurosci 18, 222–235.
doi:10.1038/nrn.2017.15
Lewis, T.L., Mao, T., Svoboda, K., Arnold, D.B., 2009. Myosin-dependent targeting of
transmembrane proteins to neuronal dendrites. Nat Neurosci 12, 568–576.
doi:10.1038/nn.2318
Li, X., Gutierrez, D.V., Hanson, M.G., Han, J., Mark, M.D., Chiel, H., Hegemann, P.,
Landmesser, L.T., Herlitze, S., 2005. Fast noninvasive activation and inhibition of neural
and network activity by vertebrate rhodopsin and green algae channelrhodopsin. Proc Natl
Acad Sci USA 102, 17816–17821. doi:10.1073/pnas.0509030102
Lim, S.T., Antonucci, D.E., Scannevin, R.H., Trimmer, J.S., 2000. A novel targeting signal for
proximal clustering of the Kv2.1 K+ channel in hippocampal neurons. Neuron 25, 385–397.
Lin, J.Y., Knutsen, P.M., Muller, A., Kleinfeld, D., Tsien, R.Y., 2013. ReaChR: a red-shifted
variant of channelrhodopsin enables deep transcranial optogenetic excitation. Nat Neurosci
16, 1499–1508. doi:10.1038/nn.3502

Madisen, L., Garner, A.R., Shimaoka, D., Chuong, A.S., Klapoetke, N.C., Li, L., van der Bourg,
A., Niino, Y., Egolf, L., Monetti, C., Gu, H., Mills, M., Cheng, A., Tasic, B., Nguyen, T.N.,
Sunkin, S.M., Benucci, A., Nagy, A., Miyawaki, A., Helmchen, F., Empson, R.M., Knöpfel,
T., Boyden, E.S., Reid, R.C., Carandini, M., Zeng, H., 2015. Transgenic mice for
intersectional targeting of neural sensors and effectors with high specificity and
performance. Neuron 85, 942–958. doi:10.1016/j.neuron.2015.02.022
Madisen, L., Zwingman, T.A., Sunkin, S.M., Oh, S.W., Zariwala, H.A., Gu, H., Ng, L.L.,
Palmiter, R.D., Hawrylycz, M.J., Jones, A.R., Lein, E.S., Zeng, H., 2010. A robust and high-
throughput Cre reporting and characterization system for the whole mouse brain. Nat
Neurosci 13, 133–140. doi:10.1038/nn.2467
Mahn, M., Prigge, M., Ron, S., Levy, R., Yizhar, O., 2016. Biophysical constraints of
optogenetic inhibition at presynaptic terminals. Nat Neurosci 19, 554–556.
doi:10.1038/nn.4266
Mahn, M., Gibor, L., Malina, K.C-K., Patil, P., Printz, Y., Oring, S., Levy, R., Lampl, I., Yizhar,
O., 2017. High-efficiency optogenetic silencing with soma-targeted anion-conducting
channelrhodopsins. bioRxiv. doi: https://doi.org/10.1101/225847
Malyshev, A.Y., Roshchin, M.V., Smirnova, G.R., Dolgikh, D.A., Balaban, P.M., Ostrovsky,
M.A., 2017. Chloride conducting light activated channel GtACR2 can produce both
cessation of firing and generation of action potentials in cortical neurons in response to light.
Neurosci. Lett. 640, 76–80. doi:10.1016/j.neulet.2017.01.026
Mardinly, A.R., Oldenburg, I.A., Pégard, N.C., Sridharan, S., Lyall, E.H., Chesnov, K.,
Brohawn, S.G., Waller, L., Adesnik, H., 2018. Precise multimodal optical control of neural
ensemble activity. Nat Neurosci 466, 123. doi:10.1038/s41593-018-0139-8
Marty, A., Llano, I., 2005. Excitatory effects of GABA in established brain networks. Trends
Neurosci 28, 284–289. doi:10.1016/j.tins.2005.04.003
Mauss, A.S., Busch, C., Borst, A., 2017. Optogenetic Neuronal Silencing in Drosophila during
Visual Processing. Sci Rep 7, 13823. doi:10.1038/s41598-017-14076-7
Mitsui, S., Saito, M., Hayashi, K., Mori, K., Yoshihara, Y., 2005. A novel phenylalanine-based
targeting signal directs telencephalin to neuronal dendrites. Journal of Neuroscience 25,
1122–1131. doi:10.1523/JNEUROSCI.3853-04.2005
Mohamed, G.A., Cheng, R.-K., Ho, J., Krishnan, S., Mohammad, F., Claridge-Chang, A.,
Jesuthasan, S., 2017. Optical inhibition of larval zebrafish behaviour with anion
channelrhodopsins. BMC Biol 15, 103. doi:10.1186/s12915-017-0430-2
Mohammad, F., Stewart, J.C., Ott, S., Chlebikova, K., Chua, J.Y., Koh, T.-W., Ho, J., Claridge-
Chang, A., 2017. Optogenetic inhibition of behavior with anion channelrhodopsins. Nat
Methods 14, 271–274. doi:10.1038/nmeth.4148
Nagel, G., Szellas, T., Huhn, W., Kateriya, S., Adeishvili, N., Berthold, P., Ollig, D., Hegemann,
P., Bamberg, E., 2003. Channelrhodopsin-2, a directly light-gated cation-selective membrane
channel. Proc Natl Acad Sci USA 100, 13940–13945. doi:10.1073/pnas.1936192100
Owens, D.F., Boyce, L.H., Davis, M.B., Kriegstein, A.R., 1996. Excitatory GABA responses in
embryonic and neonatal cortical slices demonstrated by gramicidin perforated-patch
recordings and calcium imaging. J Neurosci 16, 6414–6423.
Petreanu, L., Mao, T., Sternson, S.M., Svoboda, K., 2009. The subcellular organization of
neocortical excitatory connections. Nature 457, 1142–1145. doi:10.1038/nature07709
Price, G.D., Trussell, L.O., 2006. Estimate of the chloride concentration in a central
glutamatergic terminal: a gramicidin perforated-patch study on the calyx of Held. Journal of

Neuroscience 26, 11432–11436. doi:10.1523/JNEUROSCI.1660-06.2006

Pugh, J.R., Jahr, C.E., 2011. Axonal GABAA receptors increase cerebellar granule cell
excitability and synaptic activity. Journal of Neuroscience 31, 565–574.
doi:10.1523/JNEUROSCI.4506-10.2011

Raimondo, J.V., Kay, L., Ellender, T.J., Akerman, C.J., 2012. Optogenetic silencing strategies
differ in their effects on inhibitory synaptic transmission. Nat Neurosci 15, 1102–1104.
doi:10.1038/nn.3143

Rivera, J.F., Ahmad, S., Quick, M.W., Liman, E.R., Arnold, D.B., 2003. An evolutionarily
conserved dileucine motif in Shal K+ channels mediates dendritic targeting. Nat Neurosci 6,
243–250. doi:10.1038/nn1020

Robinson, R.B., Siegelbaum, S.A., 2003. Hyperpolarization-activated cation currents: from
molecules to physiological function. Annu. Rev. Physiol. 65, 453–480.
doi:10.1146/annurev.physiol.65.092101.142734

Rosales, C.R., Osborne, K.D., Zuccarino, G.V., Scheiffele, P., Silverman, M.A., 2005. A
cytoplasmic motif targets neuroligin-1 exclusively to dendrites of cultured hippocampal
neurons. The European journal of neuroscience 22, 2381–2386. doi:10.1111/j.1460-
9568.2005.04400.x

Shemesh, O.A., Tanese, D., Zampini, V., Linghu, C., Piatkevich, K., Ronzitti, E.,
Papagiakoumou, E., Boyden, E.S., Emiliani, V., 2017. Temporally precise single-cell-
resolution optogenetics. Nat Neurosci 20, 1796–1806. doi:10.1038/s41593-017-0018-8

Stell, B.M., Rostaing, P., Triller, A., Marty, A., 2007. Activation of presynaptic GABA(A)
receptors induces glutamate release from parallel fiber synapses. Journal of Neuroscience 27,
9022–9031. doi:10.1523/JNEUROSCI.1954-07.2007

Trigo, F.F., Marty, A., Stell, B.M., 2008. Axonal GABAA receptors. The European journal of
neuroscience 28, 841–848. doi:10.1111/j.1460-9568.2008.06404.x

Turecek, R., Trussell, L.O., 2001. Presynaptic glycine receptors enhance transmitter release at a
mammalian central synapse. Nature 411, 587–590. doi:10.1038/35079084

Wietek, J., Beltramo, R., Scanziani, M., Hegemann, P., Oertner, T.G., Simon Wiegert, J., 2015.
An improved chloride-conducting channelrhodopsin for light-induced inhibition of neuronal
activity in vivo. Sci Rep 5, 14807. doi:10.1038/srep14807

Wietek, J., Wiegert, J.S., Adeishvili, N., Schneider, F., Watanabe, H., Tsunoda, S.P., Vogt, A.,
Elstner, M., Oertner, T.G., Hegemann, P., 2014. Conversion of channelrhodopsin into a
light-gated chloride channel. Science 344, 409–412. doi:10.1126/science.1249375

Wu, C., Ivanova, E., Zhang, Y., Pan, Z.-H., 2013. rAAV-mediated subcellular targeting of
optogenetic tools in retinal ganglion cells in vivo. PLoS ONE 8, e66332.
doi:10.1371/journal.pone.0066332

Xia, Y., Zhao, Y., Yang, M., Zeng, S., Shu, Y., 2014. Regulation of action potential waveforms
by axonal GABAA receptors in cortical pyramidal neurons. PLoS ONE 9, e100968.
doi:10.1371/journal.pone.0100968

Xue, M., Atallah, B.V., Scanziani, M., 2014. Equalizing excitation-inhibition ratios across visual
cortical neurons. Nature 511, 596–600. doi:10.1038/nature13321

Zhang, F., Wang, L.-P., Brauner, M., Liewald, J.F., Kay, K., Watzke, N., Wood, P.G., Bamberg,
E., Nagel, G., Gottschalk, A., Deisseroth, K., 2007. Multimodal fast optical interrogation of
neural circuitry. Nature 446, 633–639. doi:10.1038/nature05744