Multiplexed Primer Extension Sequencing Enables High Precision Detection of Rare Splice

Isoforms

Hansen Xu[+], Benjamin J. Fair[+], Zach Dwyer, Michael Gildea, and Jeffrey A. Pleiss*

Department of Molecular Biology and Genetics, Cornell University, Ithaca, NY 14853

[+] - equal contribution

* - correspondence: jpleiss@cornell.edu

ABSTRACT

Targeted RNA-sequencing aims to focus coverage on areas of interest that are inadequately

sampled in standard RNA-sequencing experiments. Here we present a novel approach for

targeted RNA-sequencing that uses complex pools of reverse transcription primers to enable

sequencing enrichment at user-selected locations across the genome. We demonstrate this

approach by targeting pre-mRNA splice junctions in *S. cerevisiae*, revealing high-precision

detection of splice isoforms, including rare pre-mRNA splicing intermediates.

Main text

RNA sequencing (RNA-Seq) has greatly expanded our understanding of the variety of splice isoforms that can be generated within a cell. Identification of the small subset of reads that span exon-exon junctions within transcripts has enabled the unambiguous detection of vast numbers of novel splice isoforms in scores of organisms[1,2]. Yet in spite of the power presented by this approach, the sequencing depth necessary to quantitatively detect many splicing events is significantly higher than most experiments generate. While this limitation of whole-transcriptome profiling has been addressed in part by methods that utilize antisense probes[3,4] or PCR enrichment[5] to target sequencing coverage to genomic regions of interest, a deeper understanding of the basic mechanisms by which splicing is regulated, and the pathological consequences of its mis-regulation, will be facilitated by methods that enable higher resolution and precise detection of splicing states within cells. Towards this end, we have designed and implemented a novel targeted sequencing method that enhances splice junction detection and allows for genome-wide resolution of splicing intermediates. Building upon the historically validated use of primer extension as a tool for assessing splicing status, we demonstrate the ability to multiplex primer extension assays and evaluate the products by deep sequencing, an approach we hereafter refer to as Multiplexed Primer Extension sequencing, or MPE-seq (Fig1A).

The method we developed harbors two straightforward yet key features. First, user-selected primers are used to generate complementary DNA (cDNA) during a reverse transcription reaction, enabling targeting of RNA regions of interest. Each primer is appended with a next-generation sequencing adapter, as well as a unique molecular identifier (UMI) to alleviate artifacts associated with PCR amplification during library preparation[6]. The use of

elevated temperatures during the reverse transcription reaction minimizes non-specific primer

annealing (FigS1), while the inclusion of derivatized nucleotides allows for the efficient

purification of extended products and removal of excess primers. Secondly, a strand-extension

step similar to template-switching[7] is used to append the second sequencing adapter onto the 3'

terminus of the cDNA molecules. Coupling this approach with paired-end sequencing allows for

the simultaneous querying of the 5' and 3' ends of the cDNAs from targeted regions (see

Methods for full details).

As an initial demonstration of MPE-seq, we examined pre-mRNA splicing in the budding

yeast *Saccharomyces cerevisiae*. For each of the 309 annotated introns in the yeast genome,

primers were systematically designed within a 50nt window immediately downstream of the 3'

splice site, ensuring that short extensions would be sufficient to cross the splice junctions.

Primers were pooled at equimolar concentration and MPE-seq libraries were generated using

total cellular RNA from wildtype yeast and sequenced to a depth of only ~5 million reads. As a

comparative reference, standard RNA-seq libraries were generated using poly-A selected RNA

and sequenced to ~40 million reads. Whereas the standard RNA-seq libraries yielded read

coverage that comprised full gene bodies across the transcriptome, MPE-seq coverage was

focused on the selected genes, precisely targeted to the regions upstream of the designed primers

(Fig1B). Just over 75% of sequenced fragments from MPE-seq mapped to targeted regions

(Fig2A, SupplementalInformation_Table1), resulting on average in a greater than 100-fold

enrichment in sequencing depth at these regions when compared with RNA-seq (Fig2B, FigS2).

The fold-enrichment was similar across transcripts with a wide range of expression levels

(FigS2), and from these data we extrapolate that a standard RNA-seq experiment would require

~500 million sequencing reads to achieve a similar level of coverage over the targeted regions as

these 5 million MPE-seq reads provided.

Given the increased read depth achieved over targeted regions using MPE-seq, we asked

how well rare splice events were sampled. Because each primer extension event corresponded to

a single RNA molecule, determining relative isoform expression was simplified because read

counts did not need to be adjusted for mapping space as in standard analyses of RNA-seq data

(see Methods). Importantly, the levels of intron retention determined from replicate libraries

using MPE-seq showed superior internal reproducibility compared with the larger, replicate

RNA-seq libraries (Fig2C), likely reflecting the sampling noise associated with RNA-seq data

with reduced sequencing depth over the targeted regions. Moreover, while MPE-seq is not

amenable to *de novo* discovery of novel splicing events across the entire genome, it did allow for

the identification of scores of rare, previously unannotated splicing events at the targeted regions

(SupplementalInformation_Table3), consistent with the significantly increased sensitivity of this

approach. Nevertheless, while MPE-seq provided increased sensitivity and reproducibility of

splicing measurements, the intron retention levels determined from MPE-seq in a wildtype strain

only modestly correlated with those determined by RNA-seq (FigS3A, FigS3B). Notably, this

correlation improved when comparing how these techniques measured *changes* in splicing

between samples assayed by the same methodology (FigS3C), presumably reflecting inherent

biases[8] (in fragmentation, ligation, PCR amplification, library size selection, etc.) present in one

or both approaches that are internally well controlled.

With the increased resolution provided by this approach, we sought to determine whether

we could detect splicing intermediates using MPE-seq. By identifying the locations of reverse

transcription stops, primer extension reactions have historically been used to map a variety of

biological features, including among others transcription start sites (TSSs)[9], and the locations of

branch sites within the lariat intermediate species of the pre-mRNA splicing reaction[10,11]

(Fig3A). Because the approach we developed anticipated the possibility of mapping the 3' ends

of the cDNA molecules, we examined the locations of those generated by MPE-seq. As

expected, the 3' ends of many cDNAs accumulated at the TSSs as determined by an orthologous

method[12] (FigS4), indicating that reverse transcription generally proceeded to the 5' terminus of

the RNA. Importantly, we also observed many cDNAs which terminated at or near the annotated

branchpoint motifs within introns, with decreased read coverage upstream of the motifs,

consistent with the inability of reverse transcriptase to read past the branched adenosine in the

lariat intermediate species (Fig3B, Fig3C). This drop in read coverage was not apparent in MPE-

seq libraries generated from a strain harboring a conditional mutation in Prp2, an RNA helicase

required for catalyzing the 1^{st} step of splicing[13], corroborating that many of these cDNAs

originate from lariat intermediates. We noted that these lariat-intermediate derived cDNAs

contain a unique signature of mismatches incorporated by reverse transcriptase at the branched

adenosine (FigS5), which may serve as a unique tag for *de novo* identification of branch sites in

organisms with less well annotated branch sites.

The ability of MPE-seq to differentiate between unspliced isoforms in unfractionated

cellular RNA provides a unique opportunity to investigate the relative efficiencies with which

transcripts undergo the 1^{st} and 2^{nd} chemical steps in the splicing pathway. Our data revealed that

~10% of unspliced pre-mRNAs at steady state conditions were present in the lariat intermediate

form genome-wide (see FigS6, methods), albeit with significant variation between individual

pre-mRNAs (Fig3D). Remarkably, a strong correlation was observed between the relative levels

of pre-mRNA and lariat intermediate species for a given transcript (FigS7A). Correlations were

also observed between splice intermediate levels and transcript expression level, branch motif

strength, and host gene function (FigS7B, FigS7C); however no correlation was seen when

considering the ratio of pre-1^{st} step intermediate to lariat intermediate, a metric which we expect

to reflect the relative catalysis rates of the 1^{st} and 2^{nd} step of splicing. A complete understanding

of the determinants of *in vivo* splicing efficiency will require measurements of the kinetics of

these individual steps rather than their steady state levels. The ability of MPE-seq to robustly

differentiate pre-mRNA isoforms provides a powerful new opportunity to do just this.

Whereas our initial experiments were performed using individually synthesized

oligonucleotides as primers, we sought to increase the utility of this approach by examining

methods that would facilitate an increase in the number of targeted regions. Many commercial

sources exist that allow for the cost-effective, array-based synthesis of pools of thousands of

individual oligonucleotide sequences, so we developed an approach for the pooled synthesis of

an equivalent set of the previously described primer sequences in order to test the effectiveness

with which they could be used in MPE-seq. By appending a common sequence onto the 3' end of

the desired primers and then using a protocol that included PCR amplification, restriction

digestion, and targeted strand degradation (FigS8A, FigS8B), we readily prepared a sufficient

quantity of single-stranded oligonucleotides with which to generate MPE-seq libraries.

Importantly, MPE-seq libraries generated using primers synthesized by this approach also

showed strong enrichment for the targeted regions, with levels on par with what we observed

using individually synthesized oligonucleotide primers (FigS8C). Moreover, genome-wide

splicing efficiencies determined from MPE-seq libraries generated using primers from this

pooled synthesis were highly correlated with splice efficiencies derived from individually

synthesized oligos (FigS8D), confirming the capacity of this approach to generate large numbers

of unique primers for use in MPE-seq experiments.

Our work here demonstrates the capacity of MPE-seq to facilitate examinations of pre-

mRNA splicing status in a targeted, cost-effective way that improves the precision and

sensitivity of splice isoform detection. The improved sensitivity of this approach is perhaps best

exemplified by our ability to detect the lariat intermediate products of the pre-mRNA splicing

pathway. Though other methods have reported large-scale detection of upstream exon splicing

intermediates[14–16], MPE-seq uniquely detects the lariat intermediate from unfractionated cellular

RNA, and does not require laborious purification of complexes from cellular extracts. Our

demonstration that oligonucleotides derived from pooled commercial syntheses can be used in

this approach expands the types of applications to which MPE-seq could be applied in a cost-

effective manner. While we see no *de facto* limitation to the species or number of unique primer

sequences that could be used for MPE-seq, with increasing numbers of primers comes an

increasing potential for their cross-reactivity with undesirable RNA targets, highlighting the

importance of specificity in primer design. Similarly, the level of enrichment provided by this

approach would vary as a function not only of the number of regions being targeted, but also of

the distribution of the expression levels of those targets. We expect that the improved sensitivity,

precision and flexibility of this approach will enable a higher-resolution understanding of the

pre-mRNA splicing pathway. Likewise, primer extension assays have also been used to assay

RNA secondary structure after *in vitro*[17] or *in vivo*[18] chemical probing, and we expect that MPE-

seq could be readily adapted to RNA structure interrogation and other approaches where primer-

extension assays or targeted RNA sequencing is applicable.

Online Methods

Strain Maintenance and Growth Conditions. Unless otherwise indicated, all experiments used

the wild type (WT) *S. cerevisiae* strain BY4741 (*MATa, his2Δ1, leu2Δ0, met15Δ0, ura3Δ0*).

Single colonies were inoculated into liquid YPD media and grown overnight at 30°C. Overnight

cultures were then inoculated into fresh liquid YPD, seeding cultures at OD_{600} ~0.05. Cells were

collected by vacuum filtration once cultures reached OD_{600} ~0.7 immediately followed by flash

freezing in liquid nitrogen. Cell pellets were then stored at -80°C. For the temperature sensitive

mutant *prp2-1*[19] we grew cultures as described above except at 25°C. Once cultures reached

OD_{600} ~0.7, an equal volume of fresh 50°C YPD media was added to shift cells to the non-

permissive temperature of 37°C. The cultures were then maintained at 37°C for 15 minutes

before cell collection as described above.

MPE-seq Primer and Oligo Design.

Gene specific reverse transcription primer design. For each of the 309 annotated spliceosomal

introns within the budding yeast genome (annotations obtained from UCSC SacCer3), a reverse

transcription primer was designed within the first 50 nucleotides downstream of the intron.

Targeting to this region ensured that short-read sequencing of the products generated from

reverse transcription with these primers would cross the upstream exon-exon or exon-intron

boundaries, enabling determination of the splicing status. Primers were designed using

OligoWiz, a program initially developed for microarray probe design, but which enables the

selection of primer sequences optimized for target specificity relative to a designated genomic

background[20]. We used the standalone version of OligoWiz with default parameters for short

(24-26bp) oligo design to obtain optimal sequences within each 50bp window. To the 5' end of

each of these sequences was appended two additional sequence elements: a random 7-nucleotide

unique molecular index (UMI) which allows for the detection and removal of amplification

artifacts arising from library preparation[6]; and the P5XX region of the Illumina sequencing

primer to enable the sequencing of the reverse transcription products. Each of these primers was

individually synthesized by Integrated DNA Technologies (IDT), the full sequences of which are

provided in (Supplemental Information Table 5).

Complex oligo mix amplification method: The above described oligo primers were batch

synthesized as a pool on a OligoMix microarray by LC Sciences. These oligos are synthesized at

vastly lower quantities than is required for cDNA synthesis in MPE-seq. To generate a quantity

of primer pool that is sufficient, PCR amplification, along with several processing steps were

used (FigS8A). This was enabled by addition of two key sequence elements appended onto the

3' end of the individually synthesized oligo primers detailed above. From the 5' to 3' direction:

1.) a SapI restriction site, and 2.) a PCR amplification sequence (Supplemental information Table

5). First, the oligos were amplified in a standard PCR using Phusion polymerase with 14

amplification cycles. This 400 µL PCR reaction contained: 1% of the pooled oligonucleotides

from LCSciences as a template, a forward primer (oHX093) containing a C3 spacer at its 5' end,

and a reverse amplification primer (oHX094) containing a biotin-label at its 5' end (see

Supplemental Information Table 6). Cycling conditions were as follows: denaturation at 95°C for

10 sec; annealing at 60°C for 20 sec; and extension at 72°C for 30 sec. Upon completion of this

initial reaction, the entire reaction was used as a template to seed a larger (40 mL) PCR reaction.

For efficient amplification, this large reaction was performed in four 96-well plates with 100µL

in each well. Reaction conditions were identical to those described for the first reaction, and a

total of 15 cycles were performed for this second amplification. Reactions were purified and

concentrated by isopropanol precipitation. To generate single stranded primers for use in MPE-

seq, the double-stranded amplicons were first digested using SapI (NEB R0569) in a 150 µL

reaction containing 30 µL of enzyme. The reaction was incubated at 37°C overnight, after which

the reaction products were concentrated by ethanol precipitation. Next, the 5' to 3' lambda

exonuclease (NEB M0262) was used to preferentially degrade the two strands containing

unmodified 5' ends. This reaction was performed at 37°C for 2 hours according to the

manufacturer's protocol. The products of this reaction were then purified using Zymo columns

using 7X volume binding buffer (2 M guanidinium-HCl, 75% isopropanol). After this step, the

remaining DNA consisted of the desired single stranded RT primer, and an undesired single

stranded section containing the SapI site plus the amplification primer. Making use of the 5'

biotin tag on the amplification primer, these undesired oligos were removed by affinity capture

with streptavidin beads. Specifically, this was accomplished by using 50 µL of Dynabeads

MyOne Streptavidin C1 according to the manufacturer's protocol. The unbound supernatant

fraction was retained as it contains the desired products. The recovered material was precipitated

and verified using 6% native PAGE stained with SyBr Gold.

1st Strand extension template oligo design. The oligos were designed with three key features

from the 5' to 3' end of the oligo: 1.) A portion of the Nextera P7XX sequencing adapter. Of the

entirety of the P7XX adapter, the region 3' of the I7 barcode was used. This allowed for

barcoding and amplification of the sequencing libraries. 2.) a dN9 anchor on the 3' end to

randomly anneal to cDNA products. 3.) A 3' carbon block modification. This was done so that

the oligo may only be used as a template to append the Nextera sequencing adapter to the end of

cDNAs, rather than as a primer. The full sequence of this primer can be found in (Supplemental

Information Table 6).

MPE-Seq Library Prep

cDNA synthesis. RNA was isolated following a hot acid phenol extraction protocol[21]. A total of

10ug of total RNA was used to generate each library. cDNA was synthesized by mixing 1 µg of

the gene specific primer pool described above with each RNA sample in 50 mM Tris-HCl (pH

8.5), 75 mM KCl in a 20 µL volume. The primers were then annealed in a thermocycler with the

following cycle; 70°C for 1 minute, 65°C for 5 minutes, hold at 47°C. An equivalent volume of

MMLV reverse transcriptase enzyme mix containing 1 mM dATP, 1 mM dGTP, 1 mM dCTP,

0.4 mM aminoallyl-dUTP, 0.6 mM dTTP, 50 mM Tris-HCl (pH 8.5), 150 mM KCl, 6 mM

$MgCl_2$, 10 mM DTT was pre-heated to 47°C and added to the primer-annealed RNA mix

resulting in a total reaction volume of 40uL. Maintaining the samples at 47°C was essential for

reducing off-target cDNA synthesis. Reactions were incubated at 47°C for 3 hours, followed by

heat inactivation at 85°C for 5 minutes. Remaining RNA was hydrolyzed by addition of ½

volume of 0.3 M NaOH, 0.03 M EDTA and incubated at 65°C for 15 minutes. After

neutralization with ½ (original) volume of 0.3 M HCl, the cDNA was purified with a Zymo-5

column using 7X volume of binding buffer (2 M guanidinium HCl, 75% isopropanol). Purified

cDNA samples were dried in a SpeedVac until all liquid had evaporated.

NHS ester biotin coupling. Dried cDNA samples were resuspended in 18 µL of fresh 0.1 M

Sodium Bicarbonate (pH 9.0), to which 2 µL of 0.1 mg/µL NHS-biotin (ThermoFisher 20217)

was added. Reactions were incubated at 65°C for 1 hour followed by purification of biotin

coupled cDNA from unreacted NHS-biotin by using Zymo-5 columns using 7X volume of

binding buffer (2 M guanidinium HCl, 75% isopropanol).

Streptavidin-biotin purification. 20 µL of Dynabeads MyOne Streptavidin C1 (ThermoFisher

65602) per sample were pre-washed twice in 500 µL of 1X bind and wash buffer (5 mM Tris-HCl

(pH 7.5), 0.5 mM EDTA and 1 M NaCl) as per manufacturer's protocol. Washed beads were

resuspended in 50 µL of 2X bind and wash buffer per sample and 50 µL was combined with each

50 µL purified cDNA sample. Biotin-streptavidin binding was allowed to proceed for 30 minutes

at room temperature with rotation. Bound material was washed twice with 500 µL of 1X bind and

wash buffer, followed by an additional wash with 100 µL of 1X SSC. To ensure purification of

only single-stranded cDNAs, beads were then incubated with 0.1 M NaOH for two consecutive

room temperature washes for 10 minutes and 1 minute, respectively. Finally, the bound material

was washed 3 times with 100 µL 1X TE. cDNA was eluted from the beads by heating samples to

90°C for 2 minutes in the presence of 100 µL of 95% formamide, 10 mM EDTA. The eluate was

then purified using Zymo-5 columns as described above. cDNA was eluted from columns in 40

µL H_2O.

First strand extension. Primers were annealed to purified cDNA by combining: 1uL 1st strand

extension oligo (100 µM), 5 µL 10x NEB buffer 2, 40 µL purified cDNA sample, and 1 µL of 10

mM (each) dNTP mix. Samples were then incubated at 65°C for 5 minutes, followed by cooling

to room temperature on the bench top. To each sample was added 3 µL of Klenow exo- fragment

(NEB M0212) and reactions were incubated for 5 minutes at room temperature, after which they

were moved to 37°C for 30 minutes. Samples were subsequently purified via streptavidin beads

following the protocol described above. Samples were then purified and concentrated via Zymo-5

columns. Samples were then eluted in 33 µL H_2O.

PCR amplification. Amplification of the reaction products was accomplished by using 10 µL of

the purified material generated in the 1st strand extension reaction as a template in a PCR

reaction. Illumina Nextera (i5) and (i7) indexing primers were used in a standard 50 µL PCR

reaction with Phusion polymerase (ThermoFisher F530S). Cycling conditions were as follows:

denaturation at 95°C for 10 sec; annealing at 62°C for 20 sec; and extension at 72°C for 30 sec.

Libraries typically required between 14 and 20 cycles of amplification, depending upon the

amount of starting material. Libraries were then purified via PAGE. Each 50 μL PCR reaction

was run on a 6% native poly-acrylamide gel and DNA was resolved by staining with SyBr gold.

Libraries were size selected from 200bp to 800bp and DNA was extracted from gel fragments

via passive diffusion overnight in 0.3 M sodium acetate (pH 5.3). Libraries were then ethanol

precipitated and quantified.

cDNA synthesis temperature experiment. Due to the target specific nature of MPE-seq cDNA

synthesis, any reverse transcription (RT) events at non-target sites will reduce the fraction of on-

target reads. Indeed, these off-target events contribute significantly to the class of nonspecific

reads in a typical MPE-seq experiment (Fig2A). One way to reduce off-target RT events is

through increasing the specificity of the RT primers. We assessed this by testing the effect of

increased temperature during the RT reaction on off-target sequencing reads. MPE-seq libraries

were generated using the above described protocol with one primary difference: Increased

reaction temperatures required the use of a thermostable enzyme. For this reason, Superscript III

(ThermoFisher) was used along with the manufacturer supplied buffer (reaction concentration:

50 mM Tris-HCl (pH 8.3), 75 mM KCl, 3 mM MgCl2). Primer annealing and reactions were

carried out at 47°C, 51°C, and 55°C in replicate experiments.

MPE-Seq Data Anlysis

Sequencing and alignment. MPE-seq libraries were sequenced on the NextSeq platform by the

BRC Genomics Facility at Cornell university using 60bp (P5) +15bp (P7) paired-end chemistry.

PCR duplicates were removed from the dataset by filtering out non-unique reads with respect to

all base calls in both reads, including the 7bp UMI. In other words, for each set of identical

paired-end reads, a single read-pair was retained for analysis. MPE-seq Reads were aligned to

the yeast genome (Reference genome assembly R64-1-1[22]) using the STAR aligner[23] with the

following alignment parameters: {--alignEndsType EndToEnd --alignIntronMin 20 --

alignIntronMax 1000 --alignMatesGapMax 400 --alignSplicedMateMapLmin 16 --

alignSJDBoverhangMin 1 --outSAMmultNmax 1 --outFilterMismatchNmax 3 --

clip3pAdapterSeq CTGTCTCTTATACACATCTCCGAGCCCACGAGAC --clip5pNbases 7 0}.

Alignment files were filtered to exclude read mappings deriving from inserts less than 30 bases

(including the primer). We believe these short fragments represent unextended reverse

transcription primers that were retained in the sequencing libraries. These small fragments can

sometimes erroneously map to splice junctions or target introns, even though we believe they are

not derived from cellular RNA.

RNA-seq libraries were sequenced on an Illumina HiSeq2500 by the BRC Genomics Facility at

Cornell University using 100bp single end reads. Reads were aligned using the STAR aligner

with the following alignment parameters: {--alignEndsType EndToEnd --alignIntronMin 20 --

alignIntronMax 1000 --alignSJDBoverhangMin 1 --outSAMmultNmax 1 --

outFilterMismatchNmax 3 --clip3pAdapterSeq

CTGTCTCTTATACACATCTCCGAGCCCACGAGAC}.

When applicable, replicate libraries were combined prior to alignment. However, to assess

technical reproducibility of MPE-seq, replicate libraries were subsampled to varying read depths,

aligned separately, and compared to RNA-seq libraries also subsampled to varying read depths.

Estimating Splice isoform abundances from MPE-Seq data.

For each intron, the relative abundance of unspliced and spliced isoforms was determined by

counting spliced and unspliced reads. Spliced reads (S) were counted using the SJ.out.tab file

created by the aligner. Unspliced reads were counted using bedtools[24] to count the number of

reads that cover any part of the intron, considering only the first read of the paired-end reads.

Unspliced read counts were further categorized as deriving from a lariat intermediate (L) or pre-

1st step RNA (P) by considering the mapping location of the second read of the paired end reads,

which we observed to often terminate near the TSS, or in the case of a lariat-intermediate-

derived cDNA, near the branchpoint-A of the intron. Based on paired end mapping locations,

each fragment was categorized into one of six categories (See FigS6) and the counts within those

six categories were used to calculate S, P, and L as follows:

322
$$S = C_1 + C_2 \quad P = C_3\left(1 + \frac{C_5 + C_6}{C_3 + C_4}\right) \quad L = C_4\left(1 + \frac{C_5 + C_6}{C_3 + C_4}\right)$$

Locations of branchpoints (Supplementary Information Table 7) were determined by

consolidating the most used branchpoint from lariat sequencing data[25] and previously described

branch locations based on sequence motif searches[26].

Heatmaps and meta-gene plots: To generate metagene plots which illustrate read coverage

around features of interest, we used the deepTools ComputeMatrix command[27] in conjunction

with a BigWig coverage file of the 3' terminating bases and a bedfile containing TSS-positions as

determined by PRO-cap[12] or a bedfile containing the annotated branchpoint regions detailed

above. Importantly, this bedfile was filtered to only include branchpoint regions that would

produce a lariat intermediate that is within the size range captured by library size-selection of

MPE-seq libraries (see column "AttemptedLariatQuantification?" in Supplemental Information

Table 2).

RNA-seq Experiments

Library prep: For each RNA-seq library, 1 µg of total RNA was input into the "NEBNext Ultra

Directional RNA Library Prep Kit for Illumina". Libraries were prepared following the

manufacturer's protocol.

Estimating Splice isoform abundances from RNA-seq data: Similar to MPE-seq data, spliced

reads from target introns were counted using the SJ.out.tab file created by the aligner. Unspliced

reads were counted using the bedtools software package[24] to count the number of reads which

overlapped an intron. Spliced and unspliced read counts for each intron were then length

normalized for the feature's potential mapping space. The potential mapping space for a spliced

read is equal to 2 x read length minus the minimum splice junction overhang length. The

potential mapping space for an unspliced read is equal to the 2 x read length minus minimum

splice junction overhang length plus length of the intron. Reads counts assigned to each feature

were then divided by the length. Fraction unspliced was calculated for each intron as the quotient

of length normalized unspliced reads and spliced reads. Relative transcript expression was

calculated via transcripts per million (TPM) normalization[28], only considering exonic reads and

exonic gene-lengths.

Acknowledgements

We thank members of the Pleiss and A. Grimson laboratories for critical feedback on this work. This work was funded by a Research Scholars Grant from the American Cancer Society and NIGMS grant (GM098634) to J.A.P.

<u>Author Contributions</u>

H.X., B.F., Z.D., M.G. and J.A.P. designed research. H.X., Z.D., and M.G. performed research. H.X., B.F., and Z.D. analyzed data. All authors wrote the paper.

<u>Competing interests statement</u>

The authors have no competing interests to declare.

1.   Merkin, J., Russell, C., Chen, P. & Burge, C. B. Evolutionary dynamics of gene and

isoform regulation in Mammalian tissues. *Science* **338,** 1593–9 (2012).

2.   Barbosa-Morais, N. L. *et al.* The evolutionary landscape of alternative splicing in

vertebrate species. *Science (80-.).* **338,** 1587–1593 (2012).

3.   Mercer, T. R. *et al.* Targeted RNA sequencing reveals the deep complexity of the human

transcriptome. *Nat. Biotechnol.* **30,** 99–104 (2011).

4.   Mercer, T. R. *et al.* Targeted sequencing for gene discovery and quantification using RNA

CaptureSeq. *Nat. Protoc.* **9,** 989–1009 (2014).

5.   Blomquist, T. M. *et al.* Targeted RNA-Sequencing with Competitive Multiplex-PCR

Amplicon Libraries. *PLoS One* **8,** e79120 (2013).

6.   Kivioja, T. *et al.* Counting absolute numbers of molecules using unique molecular

identifiers. *Nat. Methods* **9,** 72–74 (2012).

7.   Zhu, Y. Y., Machleder, E. M., Chenchik, A., Li, R. & Siebert, P. D. Reverse transcriptase

template switching: A SMART^{TM} approach for full-length cDNA library construction.

BioTechniques **30,** 892–897 (2001).

8.   Zheng, W., Chung, L. M. & Zhao, H. Bias detection and correction in RNA-Sequencing

data. *BMC Bioinformatics* **12,** (2011).

9.   Carey, M. F., Peterson, C. L. & Smale, S. T. The primer extension assay. *Cold Spring*

Harb. Protoc. **8,** 164–173 (2013).

10.   Coombes, C. E. & Boeke, J. D. An evaluation of detection methods for large lariat RNAs.

RNA **11,** 323–31 (2005).

11.   Padgett, R. A. *et al.* Nonconsensus branch-site sequences in the in vitro splicing of

transcripts of mutant rabbit beta-globin genes. *Proc. Natl. Acad. Sci. U. S. A.* **82,** 8349–

8353 (1985).

12. Booth, G. T., Wang, I. X., Cheung, V. G. & Lis, J. T. Divergence of a conserved

elongation factor and transcription regulation in budding and fission yeast. *Genome Res.*

26, 799–811 (2016).

13. Kim, S. H. & Lin, R. J. Spliceosome activation by PRP2 ATPase prior to the first

transesterification reaction of pre-mRNA splicing. *Mol. Cell. Biol.* **16,** 6810–6819 (1996).

14. Chen, W. *et al.* Transcriptome-wide interrogation of the functional intronome by

spliceosome profiling. *Cell* **173,** 1031–1044 (2018).

15. Burke, J. *et al.* Spliceosome profiling visualizes the operations of a dynamic RNP in vivo

at nucleotide resolution. *Cell* **173,** 1014–1030 (2018).

16. Nojima, T. *et al.* Mammalian NET-Seq Reveals Genome-wide Nascent Transcription

Coupled to RNA Processing. *Cell* **161,** 526–540 (2015).

17. Lucks, J. B. *et al.* Multiplexed RNA structure characterization with selective 2'-hydroxyl

acylation analyzed by primer extension sequencing (SHAPE-Seq). *Proc. Natl. Acad. Sci.*

108, 11063–11068 (2011).

18. Rouskin, S., Zubradt, M., Washietl, S., Kellis, M. & Weissman, J. S. Genome-wide

probing of RNA structure reveals active unfolding of mRNA structures in vivo. *Nature*

505, 701–5 (2014).

19. Hartwell, L. H., McLaughlin, C. S. & Warner, J. R. Identification of ten genes that control

ribosome formation in yeast. *MGG Mol. Gen. Genet.* **109,** 42–56 (1970).

20. Wernersson, R. & Nielsen, H. B. OligoWiz 2.0 - Integrating sequence feature annotation

into the design of microarray probes. *Nucleic Acids Res.* **33,** (2005).

21. Collart, M. A. & Oliviero, S. in *Current Protocols in Molecular Biology* (2001).

doi:10.1002/0471142727.mb1312s23

22.   Engel, S. R. *et al.* The reference genome sequence of Saccharomyces cerevisiae: then and

now. *G3 (Bethesda).* **4,** 389–98 (2014).

23.   Dobin, A. *et al.* STAR: Ultrafast universal RNA-seq aligner. *Bioinformatics* **29,** 15–21

(2013).

24.   Quinlan, A. R. & Hall, I. M. BEDTools: A flexible suite of utilities for comparing

genomic features. *Bioinformatics* **26,** 841–842 (2010).

25.   Mayerle, M. *et al.* Structural toggle in the RNaseH domain of Prp8 helps balance splicing

fidelity and catalytic efficiency. *Proc. Natl. Acad. Sci.* **114,** 4739–4744 (2017).

26.   Grate, L. & Ares, M. Searching yeast intron data at Ares lab web site. *Methods in*

Enzymology **350,** 380–392 (2002).

27.   Ramírez, F. *et al.* deepTools2: a next generation web server for deep-sequencing data

analysis. *Nucleic Acids Res.* **44,** W160–W165 (2016).

28.   Conesa, A. *et al.* A survey of best practices for RNA-seq data analysis. *Genome Biology*

17, (2016).

Figure legends:

Figure1: MPE-seq uses complex pools of reverse transcription primers to target sequencing

to regions of interest

(A) MPE-seq comprises the following steps: (1) Primers containing sequencing adapter

overhangs and a target-specific region are designed for genomic regions of interest. (2)

Primers are pooled and used to reverse transcribe total cellular RNA in the presence of

amino-allyl dUTP (aa-dUTP), which allows (3) biotin coupling and purification of

cDNAs from other reaction components. (4) A second sequencing adapter is appended at

the 3' terminus of the cDNA through first strand extension using Klenow. Libraries are

further purified (5), PCR amplified (6) and sequenced (7).

(B) Genome browser screenshot of a targeted region in MPE-seq (pink) and conventional

RNA-seq (purple)

Figure2: MPE-seq enrichment enables high-precision measurements of splicing

(A) The percentage of reads mapped to target and off-target regions is depicted for MPE-seq

and conventional RNA-seq. In MPE-seq a small fraction of reads were categorized as

"Unextended primer" which corresponds to short primer extension products (0-5 bases

extended past the primer) and thus they were not categorized as cDNAs derived from

RNA targets.

(B) Each point represents the fold enrichment of a target region in MPE-seq over

conventional RNA-seq.

(C) Scatter plots depict intron-retention measurements in replicate libraries in MPE-seq and

conventional RNA-seq at matched or greater read depth.

Figure3: MPE-seq enables genome-wide profiling of lariat intermediates

(A) Schematic depicting cDNA products derived from pre 1st step (P), lariat intermediate (L)

and spliced mRNA (S) isoforms.

(B) Meta-intron coverage plot surrounding predicted branchpoints in a wild-type (Prp2) and

step1 splicing mutant strain (*prp2-1*). The region between the +10 position downstream

of the annotated branchpoint and the 3' splice site (3'ss) was re-scaled for each intron.

(C) Heatmap plots showing the relative coverage at each intron for which reads were

detected.

(D) Estimates of the relative abundance of each isoform for each targeted intron for which

reads were detected.

FigureS1: Elevated temperatures in reverse transcription reactions increase specificity

The fraction of on-target and off-target reads from replicate MPE-seq libraries generated

from reverse transcription reactions performed at various temperatures. A small fraction of reads

were categorized as "Unextended primer" which corresponds to short primer extension products

(0-5 bases extended past the primer) and thus they were neither categorized as cDNAs derived

from RNA targets or unamappable.

FigS2: Expression measurements as determined by MPE-seq and RNA-seq

(A) A scatter plot depicts gene expression measurements (RNA-seq) in replicate datasets.

Genes containing splice-events that were among those chosen for targeted sequencing are

depicted in red. These targeted genes range in expression levels by orders of magnitude.

(B) A scatter plot depicts gene expression measurements in replicate MPE-seq datasets (red).

Similar to conventional RNA-seq, expression measurements in MPE-seq are highly

reproducible between replicates, even for the small proportion of mis-priming events that

map to off-target locations (grey).

(C) A scatter plot depicts gene expression measurements in RNA-seq and MPE-seq. The

right shift of targeted genes reflects successful enrichment of targets by orders of

magnitude. The observation that even highly expressed genes as measured by RNA-seq

are proportionally highly expressed in MPE-seq suggests that primers are not limiting

during reverse transcription.

FigS3: Splicing measurements as determined by MPE-seq and RNA-seq

(A) A scatter plot depicts intron-retention measurements in MPE-seq and conventional RNA-

seq using wildtype (Prp2) RNA.

(B) A scatter plot depicts intron-retention measurements in MPE-seq and conventional RNA-

seq using RNA from a splicing mutant strain (*prp2-1*).

(C) A scatter plot depicts the fold-change (*prp2-1/*Prp2) in intron-retention as measured by

MPE-seq and conventional RNA-seq

FigS4: Transcription start site profiling by MPE-seq

Metagene profile of 3' ends mapped by MPE-seq, centered around transcription start sites

(TSS) as determined by PRO-cap, an orthologous method for mapping transcription start sites.

The high abudance of read ends that pile up at TSSs indicates that MPE-seq can be used to

profile cDNA termini.

FigS5: Lariat-intermediate derived cDNAs contain a unique signature of mismatches

incorporated by reverse transcriptase at the branched adenosine

(A) Genome browser screenshot of 3' reads from paired-end sequenced fragments

497   illustrates the unique signature of non-templated base incorporation by reverse

498   transcriptase at a branched-adenosine vs the 5' RNA terminus

499   (B) Genome-wide quantification of the mismatch frequencies at 3' cDNA termini near

500   the TSS (left) versus at the annotated branchpoint (right).

FigS6: Schematic for assigning reads to splice intermediate isoforms

To quantify the abundance of pre 1^{st} step, lariat intermediate, and spliced isoforms for

each targeted splice event, we categorized fragments into six classes based on paired-end read

alignments. Fragments containing a splice junction (C_1 and C_2) are indicative of a spliced RNA

(S). Fragments that are unspliced and traverse the branchpoint region (C_3) are classified as pre-1^{st}

step RNA (P). Fragments that are unspliced but terminate within a -3 to +5bp window from the

previously determined branchpoint (C_4) are classified as lariat intermediate (L). Fragments that

are unspliced and either terminate downstream of the branchpoint (C_5) or the terminus could not

be mapped (C_6) are ambiguous between P and L. Therefore, for accounting purposes, the counts

for these fragments were coerced into P and L classifications based on the ratio of P and L

determined by unambiguous mappings (C_3 and C_4). See methods for more details.

FigS7: Transcript features that correlate with the abundance of lariat intermediate

(A) Scatter plot depicting the correlation between the abundance of pre-1^{st} step RNA and the

515   abundance of lariat intermediate

(B) The abundance of pre-1st step RNA and lariat intermediate RNA is significantly

correlated with classification of introns into those that are in ribosomal protein genes

(RPG) and non-RPG. However, the abundance of lariat intermediate relative to pre-1st

step RNA, a metric of the efficiency of the second step of splicing, does not correlate.

(C) Spearman correlations of various features to the abundance of pre-1st step RNA, lariat

intermediate, the abundance of lariat intermediate relative to pre-1st step RNA. Errorbars

indicate 95% confidence intervals as estimated by Fisher transformation of Spearman's

correlation coefficient.

FigS8: Array-based oligonucleotide synthesis can be used to generate primer pools for use

in MPE-seq

(A) Obtaining adequate amounts of primer pools for MPE-seq from cost-effective array-

based oligonucleotide synthesis can be achieved in four steps: (1) PCR amplification of

the oligonucleotide synthesis pool using a 5' blocked sense primer and a biotinylated

antisense primer. (2) Restriction digestion to cleave off the PCR primer handle. (3)

Lambda exonuclease digestion of free 5' ends. (4) Streptavidin purification of

biotinylated PCR handle. The unbound fraction is the desired primer pool product.

(B) Steps during the amplification and purification of array-synthesized primer pools are

monitored via native gel electrophoresis. The control lane represents a pool of

individually synthesized MPE-seq primers which did not require amplification and

purification.

(C) The percentage of reads mapped to target and off-target regions is depicted for MPE-seq

using array-synthesized primers. In MPE-seq a small fraction of reads were categorized

as "Unextended primer" which corresponds to short primer extension products (0-5 bases

extended past the primer) and thus they were not categorized as cDNAs derived from

RNA targets.

(D) A scatter plot depicts intron-retention measurements in MPE-seq libraries which used

individually synthesized primer pools and array-based synthesis of primer pools

A

1 Primer Pool Design
Target specific sequence · UMI · First sequencing adapter · 5′ · Hundreds of targets

2 Reverse Transcription
aa-dUTP · RNA · 5′ · aa-dUTP incorporated · 5′

3 cDNA Purification
Biotin coupling · Streptavidin bead · Unextended primers removed · 5′

4 First Strand Extension
5′ · 3′ blocking group · Random 9mer · Second sequencing adapter · Second sequencing adapter appended · 5′

5 cDNA Purification
Streptavidin bead · Strand extension oligonucleotide removed · 5′

6 PCR Amplification
5′ · Sample barcode · 5′ · 5′

7 Sequencing
ATGGGGATTAAGCTAATGCCCATTAAGCTAATGCCCATTAAGCTAC
CAAGCCGTTGCCAATGCAATGCCTCAATGCCCATTAAGCT
CATTAAGCTAATGCCCATTAAGCTACGATGTACAAGCCGT

©2017 *LiYau*

B

YHL007C YHL006C YHL003C YHL002C-A YHL001W YHR001W YHR002W YHR004C YHR005C tT(AGU)H YHR006W

[0 - 5428]

[0 - 236]

YHL001W

MPE-seq [0 - 5428]
RNA-seq [0 - 128]

A

cDNA termini
at TSS

cDNA termini
at branchA

B

TSS

branchA

Reference
Genome Match

Mismatch

A

C

G

T

C_1
C_2
C_3
C_4
C_5
C_6

5′   A   3′

R2   R1
Fragment mapped
as paired-end alignments

Fragment mapped
as single-end alignment

Spliced segment
in alignment

A Array synthesis of oligos

First sequencing adapter | UMI | Gene specific variable region | SapI digest site | Reverse amplification primer region

1 PCR Amplification
Blocked 5′ end
Forward amplification primer
biotin
Reverse amplification primer

2 SapI Digestion
SapI digestion

3 Lambda Exonuclease Digestion
Lambda exonuclease

4 Streptavidin Purification
Streptavidin bead
Removed

Complex Primer Pool
~300 different gene specific variable regions

©2017 *Li Yao*

B
100 bp ladder | ① | ① | ② | ③ | ④ | Contol
Column cleaned up
300
200
100

C
31
45
2
23
Off-target
On-target
PCR-duplicate
Unmapped
Unextended primer
UMI

D
Array ($^{Unspliced}/_{Spliced}$)
Individual ($^{Unspliced}/_{Spliced}$)
$R^2 = 0.82$