Transcriptome analysis of early stage of neurogenesis reveals temporal-specific gene network for preplate neuron differentiation and Cajal-Retzius cell specification

Jia Li[1,2,3†], Lei Sun[4†], Xue-Liang Peng[1], Xiao-Ming Yu[2], Shao-Jun Qi[1,2], Zhi John Lu[5], Jing-Dong J. Han[6], Qin Shen[1*].

†Equal contributors

[1] Current address: Brain and Spinal Cord Innovative Research Center of Tongji Hospital, School of Life Sciences and Technology, Tongji University, Shanghai 200065, People's Republic of China

[2] School of Medicine, Tsinghua University, Beijing 100084, People's Republic of China

[3] PTN graduate program, School of Life Sciences, Peking University, Beijing 100871, People's Republic of China

[4] PTN graduate program, School of Life Sciences, Tsinghua University, Beijing 100084, People's Republic of China

[5] MOE Key Laboratory of Bioinformatics, Center for Synthetic and Systems Biology, School of Life Sciences, Tsinghua University, Beijing 100084, People's Republic of China

[6] Key Laboratory of Computational Biology, CAS Center for Excellence in Molecular Cell Science, Collaborative Innovation Center for Genetics and Developmental Biology, Chinese Academy of Sciences-Max Planck Partner Institute for Computational Biology, Shanghai Institutes for Biological Sciences, Chinese Academy of Sciences, Shanghai 200031, People's Republic of China

*Correspondence: shenqin@tongji.edu.cn

Abstract

Background: Neurogenesis in the developing cerebral cortex begins with the generation of the

preplate, which consists of early born neurons including Cajal-Retzius (CR) cells and subplate

neurons. CR cells later segregate to layer 1, the most superficial layer, and play an important role

in organizing cortical lamination. However, the molecular mechanisms underlying the generation

of preplate and CR cell differentiation remain poorly understood.

Results: Using the Ebf2-EGFP transgenic mouse line in which EGFP labels the preplate first then

persists in CR cells, we compared differentially expressed gene profile in Ebf2-EGFP-positive and

Ebf2-EGFP-negative cell population by high-throughput RNA-seq at different stages during early

cortical development. Bioinformatic analysis revealed the temporal gene expression dynamics of

Ebf2-EGFP+ cells and identified a number of coding genes and lncRNAs that were involved in

early neuronal differentiation. Furthermore, single cell RNA-seq analysis unraveled eight

subpopulations of Ebf2-EGFP+ cells at E15.5, which were classified into three developing states

using pseudotiming analysis. We also found that the upstream regulatory regions of Ebf2+

cell-enriched genes including lncRNAs were associated with active histone modifications,

indicating epigenetic involvements in early differentiating neuron gene expression. Furthermore,

we characterized the functional role of Ebf2+ cell-enriched lncRNAs in early neuron

differentiation using assays including primary cell culture, wholemount culture and *In Utero*

Electroporation of layer 1.

Conclusions: We have provided a genetic and epigenetic atlas of early neurogenesis and identified

genes associated with CR neuron identity. Our work will help understand the molecular

mechanisms governing the first differentiation step of cortical development.

Keywords: neurogenesis, preplate, Cajal-Retzius neuron, cerebral cortex, neural development,

RNA-seq, single-cell RNA-seq, histone modification, ChIP-seq, lncRNAs

Background

The mammalian cortical neurogenesis occurs on a precise time schedule during development.

Neural stem cells (NSCs), the neuroepithelial cells and radial glial cells in the ventricular zone

(VZ) first undergo symmetric cell divisions to expand, then begin to produce neurons [1]. The

first-born neurons are generated as early as E10 in mice and form the preplate or primordial

plexiform layer, which situates above the ventricular zone and beneath the pial surface [1-3]. The

preplate is then split into the marginal zone (MZ, layer 1) and the subplate by immigrating cortical

plate neurons around E13.5 in mouse and the seventh to eighth week of gestation in human [4-6],

setting up of the upper and lower borders of the cortical plate. The formation and splitting of

preplate are the first steps in the establishment of the highly organized cortical layers.

The preplate and its derivatives play a critical role in regulating subsequent inside-out

neuronal migration, cortical lamination and the establishment of topographic connections [7].

Subplate neurons contribute to the guidance of corticofugal and thalamocortical axons during

early development and are important for the functional maturation and plasticity of the cortical

circuitry [8, 9]. Cajal-Retzius neurons are the major cell type in layer 1, the most superficial

cortical layer. Discovered more than a century ago by Retzius and Ramon y Cajal, CR neurons

have been studied extensively [3, 10]. Best known for their expression of Reln, CR neurons

regulate the inside-out migration of cortical neurons, correct layer formation, maintenance of

radial glia and cortical patterning [11]. Loss of Reln or mislocalization of CR cells contributes to

defective migration and disordered cortical layer formation that are associated with human brain

malformation such as lissencephaly [12] and dystroglycanopathy [13]. Although functional

importance of these early differentiating neurons is well recognized, characterization of

cell-intrinsic, dynamic gene expression patterns in these cells remains incomplete. Recent studies

have identified the genes enriched in subplate neurons by microarray and RNA-seq analysis

[14-18], providing a comprehensive gene expression profile of subplate at different developmental

stages, which offers valuable insight on molecular mechanisms underlying neurodevelopmental

disorders [17]. However, how gene expression is dynamically regulated during the transition from

the preplate to CR neurons remains unclear.

We previously have shown that in the Ebf2-EGFP transgenic mice (GENSAT project) [19],

the GFP signal is expressed specifically in early differentiating preplate neurons and persistent in

CR cells [20]. After preplate splitting, virtually all of the EGFP-positive cells exhibit

morphological and molecular features of CR neurons in the embryonic and postnatal brain [20],

consistent with the pattern of the Ebf2[GFPiCre] [21], making this line suitable for tracing the

development and profiling genetic and epigenetic pattern of early born neurons including CR

neurons. In addition, single-cell gene expression profiling allows further dissection of the

heterogeneity and developmental trajectory within a cell population [22, 23]. While multiple types

of neurons in hippocampus and adult cortex have been identified by single-cell RNA-sequencing

[23, 24], a full characterization of early differentiating neurons at single-cell level has not been

done before. In this study, we used FACS-purified Ebf2-EGFP-expressing cells to probe the

dynamic molecular features of preplate and CR neurons during the early stages of forebrain

development. Whole genome transcriptome analysis identified the initial difference between the

preplate and progenitor cells at E11.5, reflecting the molecular choice during the first cell fate

decision in neural differentiation. We also recognized clusters of genes that displayed temporal

changes corresponding to different developmental trajectories. Furthermore, single-cell

sequencing of E15.5 CR neurons identified novel subpopulations. Genome-wide overview of

active (H3K27ac) and repressive (H3K27me3) histone modifications in preplate neurons, CR

neurons and NPCs revealed that their occupancies around TSS and distal region were associated

with expression of early differentiating neurons-enriched genes and lncRNAs. By bioinformatic

functional enrichment analysis, we also identified and tested the function of genes and lncRNAs

that were enriched in CR neurons in neuronal differentiation. Hence, our study provides unique

molecular markers to extend the repertoire of preplate and CR neuron labels and reveals

remarkable dynamics in gene expression during early cortical development.

Results

Characterization of Ebf2-EGFP-expressing cells in the early cerebral cortex

We purified Ebf2-EGFP expressing cells by FACS from the embryonic mouse forebrain at E11.5,

E13.5 and E15.5, to compare the developmental gene expression dynamics during preplate

formation and differentiation of early neurons. At E11.5, Ebf2-EGFP+ cells are mostly preplate

neurons including both subplate neurons and CR neurons, and at E13.5, Ebf2-EGFP expression is

diminishing in subplate neurons but maintained in CR neurons. By E15.5 Ebf2-EGFP is

specifically expressed in CR neurons [20]. At E11.5, Ebf2-expressing cells consisted of 17.1% ±

1.07% of the total cells. At E13.5 and E15.5, the percentage of Ebf2-expressing cells gradually

decreased to 8.37 ± 2.29% and 3.9 ± 1.59%, respectively (Figure 1A). 100% of FACS-sorted

Ebf2-EGFP+ cells were β-III tubulin positive at all stages, indicating these cells have committed

neuronal fate decision, consistent with our previous in vivo analysis [20]. Accordingly, after

culturing sorted and unsorted cells for 4 days, almost all of the Ebf2-EGFP-positive cells remained

single and extended processes, while the Ebf2-EGFP-negative cells proliferated and generated

large clones similar to the unsorted cell group, indicating a good separation of differentiating

neurons from progenitor cells (Figure S1A).

Identification of Ebf2-EGFP+ cell-enriched genes in the developing mouse forebrain

Next, we collected FACS-selected Ebf2-EGFP positive cells and Ebf2-EGFP negative cells from

the three early neurogenic stages and isolated total RNA for whole-genome transcriptome

sequencing. Two biological replicates for each cell type (Ebf2-EGFP+ and Ebf2-EGFP-) were

sequenced and expressed transcripts were identified as described in method. To verify the

reproducibility of our samples, we analyzed the Pearson's correlation coefficient (Pearson's r) for

the replicates and there was a strong association between different preparations of FACS-sorted

cells from the same stage (Figure S1B).

To validate the RNA-seq results, we selected several protein-coding genes, including CR

neurons markers (Ebf2 and Calb2) and genes associated with CR neurons fate regulation (Foxg1,

Tbr1, Ebf1, and Ebf3) [10, 20, 25-28], and verified by qPCR that the temporal dynamic expression

levels of these genes were in accordance with that detected by RNA-seq (Figure S1C).

Furthermore, the temporal expression trends were consistent with the mRNA expression pattern

revealed by in situ hybridization (ISH) from the data available in Allen Brain Atlas data (Figure

S1C).

We used DE-Seq to analyze the dynamic expression profiles during the early cortical

differentiation period. 4859 genes exhibited significant difference in expression level between the

two cell populations during the time course (FDR<0.05, p-values<0.05 by Fisher's exact test) and

were selected as differential expressed genes (DEGs) for further analysis (Figure 1B; Additional

file 1).

We analyzed DEGs expressed in any of the developmental stages sequenced (E11.5, E13.5

and E15.5). Genes with expression level (FPKM) more than 1, and 2-fold higher in Ebf2-EGFP

positive population compared to that in Ebf2-EGFP negative population were defined as

Ebf2-EGFP positive cells enriched genes, Ebf2+ genes in short. We found that there were 1044

genes at E11.5, 643 genes at E13.5, and 541 at E15.5 enriched in Ebf2-EGFP positive cells,

respectively. Considerable overlap was found between the different stages (Figure 1C). In total,

1513 genes displayed enriched expression in Ebf2-EGFP positive population at least at one stage,

and 1894 such genes in Ebf2-EGFP negative population. Of these, 166 genes were consistently

enriched in Ebf2-EGFP positive population throughout all three embryonic stages, and 420 genes

were differentially expressed in Ebf2-EGFP negative population (Figure 1C). Of the 166 DEGs

included widely-used CR neuron markers such as Reln and Trp73, and genes known to be

expressed in the CR neurons such as Calb2 (Calretinin), Tbr1, Ebf3, Cdkn1a [10, 25, 28, 29]. We

further defined the DEGs with FPKM ≥ 5 in Ebf2-EGFP+ cells, and fold change ≥ 2 at all three

stages as CR-specific genes. After excluding 10 subplate genes reported in previous studies [14,

16] and 14 CR-specific genes reported in previous studies [25], 77 are novel genes with no

reported association with CR neurons. Among the 77 CR-specific genes, 56 genes have valid in

situ data from public resources. We also noticed that there were genes showing high expression

levels in both Ebf2-EGFP+ cells and Ebf2-EGFP- cells although their expression levels meet the

fold change criteria. To obtain the genes specifically expressed in Ebf2-EGFP+ cells but not in

Ebf2-EGFP- cells, we added a cutoff of FPKM ≤ 15 as CR-specific genes should be expressed at

low level in Ebf2-EGFP- cells at these stages. We found 33 genes and confirmed that 8 with

specific expression in layer 1 cells (Figure 1D; Figure S1D-S1E; Additional file 1). Hence, these

genes could serve as CR-specific markers for future study.

Remarkably, the expressions of these 8 genes were regulated by overexpression or

knockdown of Ebf2, and Foxg1 knockdown in E11.5 cortical cell culture (Figure 1E; Figure

S1F-S1G). These genes included transcription factors, synapse molecules, and axon guidance

molecules, such as Nhlh2, Sncb, Slit1. For example, Nhlh2 was upregulated or inhibited by

overexpression and knockdown of Ebf2, respectively. While Sncb and Ppp2r2c responded to Ebf2

overexpression but not knockdown, the expression of Slit1 and Brinp1 were decreased by Ebf2

knockdown, suggesting they might also be regulated by molecules involved in developmental

pathways other than Ebf2 (Figure 1E; Figure S1F-S1G).

Cortical CR neurons are considered to have multiple extra-cortex origins [30]. We and others

have shown previously that Foxg1 suppressed CR neuron fate [27], therefore we hypothesized that

CR-specific genes identified by our RNA-seq analysis could be regulated by Foxg1 expression.

Indeed, silencing of Foxg1 using shRNA up-regulated Ebf2 and CR-specific genes (Nhlh2,

Ppp2r2c) in the mRNA level (Figure 1G).

Biological processes related to Ebf2-EGFP+ neuron-enriched (FC>2, FPKM>1) genes

We performed Gene Ontology analysis using the DAVID software to predict the potential function

of the DEGs that were enriched in the Ebf2-EGFP+ or Ebf2-EGFP- cell population, respectively

(Figure 2A-2D; Additional file 2). The GO terms associated with neuronal differentiation and

functions were highly enriched in DEGs expressed in the Ebf2-EGFP+ cell population, including

"synapse", "neuron projection", "neuron differentiation", "cell morphogenesis involved in

neuron", "axonogenesis" and "neuron migration" (Figure 2A). In contrast, the GO terms enriched

in the Ebf2-EGFP- cell population were mostly correlated with progenitor cell properties, such as

"forebrain development", "regulation of cell proliferation", "growth factor binding", "regulation

of neurogenesis" and "dorsal/ventral pattern formation" (Figure 2B).

In addition, we utilized the KEGG pathway repository to help determine the relative

representation of specific pathways in our global RNA-seq expression data. Significantly enriched

KEGG pathways of the global DEGs pointed out many signaling pathways implicated in normal

neural development (Figure 2E; Additional file 2). For example, genes falling under the MAPK

Signaling Pathway, Wnt Signaling Pathway and DNA replication were enriched. The Focal

Adhesion and Axon Guidance pathway were also enriched, indicating regulation of cell

morphology and neuronal differentiation were involved. Moreover, we performed the KEGG

pathway analysis focused on DEGs enriched in Ebf2-EGFP+ cell population (Figure 2F;

Additional file 2). As expected, the major groups of KEGG categories enriched in this data set

were Calcium Signaling Pathway, Neuroactive Ligand-receptor interaction, Gap Junction, GnRH

Signaling Pathway, signifying differentiation of neuronal properties. Interestingly, as Ebf2 is an

important factor in GnRH neuron differentiation [31], the enrichment of GnRH Signaling pathway

suggests that it would regulate similar pathway genes in cortical neurons.

Temporal transcriptome analysis of Ebf2-EGFP+ cells revealed early neuronal specification

and CR differentiation dynamics

The high enrichment of these GO analysis terms and KEGG pathways imply that the

transcriptional difference between the two cell populations is consistent with their different cell

type identities. The components of Ebf2-EGFP+ and Ebf2-EGFP- cell population change as

development proceeds. At E11.5, Ebf2-EGFP+ cells are mostly preplate neurons including both

subplate neurons and CR neurons, while by E15.5 Ebf2-EGFP is specifically expressed in CR

neurons [20]. In contrast, at E11.5 most if not all of the Ebf2-EGFP- cells are progenitor cells

while at E13.5 and E15.5 the Ebf2-EGFP- population also includes cortical plate neurons.

Therefore, the genes that are highly enriched in Ebf2-EGFP+ cells at all three stages, particularly

those with increased enrichment, could be specifically involved in CR neuron differentiation,

while the genes that are highly enriched in Ebf2-EGFP+ cells at E11.5 but not at E13.5 or E15.5

could be involved in regulating general neuronal differentiation processes. Hence, we further

compared temporal changes in global DEGs at the three different stages. Genes with expression

level (FPKM) in Ebf2-EGFP positive population 2 folds of that in Ebf2-EGFP negative population

were defined as "up-regulated". If the ratio of FPKM (Ebf2-EGFP+/ Ebf2-EGFP-) was between 2

and 1, these genes were defined as "unchanged". If the ratio was between 0 and 1, these genes

were defined as "down-regulated". Theoretically, we expected 27 possible combination of

expression change patterns, while our analysis showed 26 classes without the pattern of "E11up -

E13down - E15up". These 26 classes could be classified into 6 groups representing various

aspects of early neurogenesis, and 6 typical classes were shown in Figure 2G and Figure S2A

(Additional file 3). We hypothesized that CR-specific genes are consistently higher at the three

stages, while the difference in genes involved in global neuronal differentiation would be down

after E13.5 as the Ebf2-EGFP negative population also includes newly generated cortical plate

neurons. Indeed, as we expect, the genes in the group "E11up - E13up - E15up" include the

well-known CR marker genes, such as Reln, Lhx5, Trp73 and Ebf2, CR-specific genes, and

mature neuron genes related to axon regeneration like Cacna2d2 [32]. In contrast, genes in the

"E11down - E13down - E15down" keep higher expression in Ebf2-EGFP- cells, showing that they

may function on regulating NSC proliferation, such as Sox2, Pax6 and Hes5. Genes that are first

expressed in progenitor cells then in cortical neurons other than preplate and CR neurons may also

be included in this group, such as Tle1 [33]. The "E11up - E13unchange - E15down" group

includes subplate genes identified previously, such as Syt4, Igfbpl1, Myt1 [14, 16], reflecting that

subplate genes are highly expressed in the preplate on E11.5, and decreased in layer1 when the

preplate splits later on E13.5 and E15.5. Interestingly, we found that most genes in the "E11down

- E13unchange - E15up" group initially have higher expression in the cortical hem or MGE at

E11.5, then appear in layer 1 at E13.5 and E15.5, implying that these genes may be involved in

migration of CR neurons from various CR origins. For example, Zic1 and Zic2 are expressed in

the progenitor cells in the septum and cortical hem, and Bmp7 is expressed in the cortical hem,

meninges, and choroid plexus, typical CR neuron origins and location [34, 35]. The genes in the

"E11down - E13unchange - E15down" group include Foxg1, Bcl6 and Miat, which are expressed

along the lineage of neural progenitor cells [36, 37]. The "E11unchange - E13unchange -

E15unchange" group is enriched with genes associated with basic cellular organelles or activities

that required for various eukaryotic cells. For example, Atp6v0c and Atp6v0e2 in this group are

vacuolar ATPase and play essential roles in maintaining PH in the cellular compartments, protein

degradation and endocytic pathway [38, 39], while Rab11fip5 is reported to participate in protein

trafficking between different organelle membranes [40]. Taken together, our analysis revealed that

temporal gene expression dynamics reflects distinct developmental processes, allowing us to

identify novel factors that play specific roles in early cortical neurogenesis and CR development.

Differentially expressed lncRNAs in the Ebf2-EGFP+ cells and Ebf2-EGFP- cells

From our RNA-seq analysis, we identified 2417 long non-coding RNA transcripts in cortical cells

using the Cufflinks method. 200 lncRNAs were differentially expressed between the Ebf2-EGFP+

and the Ebf2-EGFP- population (FDR<0.05) and termed DElncRNAs (Figure 3A). 58 lncRNAs

were highly expressed in Ebf2-EGFP+ cells compared to Ebf2-EGFP- cells, while 64 of lncRNAs

showed the opposite pattern. We further filtered DElncRNAs with these criterions: 1) specifically

expressed in Ebf2-EGFP+ cells with FPKM at least 2-fold of that in Ebf2-EGFP negative cells

(Figure 3B); these lncRNAs also showed temporal specific expression patterns. 2) lncRNAs from

1) with FPKM ≤ 15 in Ebf2-EGFP- cells during all three stages, and FPKM ≥ 5 in Ebf2-EGFP+

cells at E15.5. We identified 9 lncRNAs that meet the criterions and these had never been studied

before in the context of neural development (Additional file 4). We called these lncRNAs as

ln-CR# for simplicity. The qPCR results to validate the mRNA expression level were generally in

agreement with the RNA-seq data (Figure 3C; Figure S3A). Notably, we detected strong

expression of ln-CR2 in E11.5 Ebf2-EGFP+ cells, and comparably lower or even no signals in

Ebf2-EGFP- cells, indicating Ebf2-EGFP+ cell-specific expression. In situ hybridization on E11.5,

E13.5 and E15.5 mouse embryonic brain sections revealed that ln-CR2 was selectively expressed

in the cortical layer next to the pial surface, as well as the TE (thalamus eminence), one of the

extra-cortical origins of CR neurons, indicating cell-type specificity (Figure 3D). Furthermore, we

performed GO analysis (see method) of DElncRNAs to predict their potential functions (Figure

3E). We found significant enrichments of neuronal development process-related GO terms, such

as "nervous system development", "neuron migration", "negative regulation of neuron apoptotic

process" and "neuron differentiation", neuron identities-related GO terms such as "neuron

projection", "axon" and "dendrite", also transcription factors regulation-related GO terms such as

"transcription factor complex", "transcription factor binding". The functional annotation of

DElncRNAs identified from our sequencing data suggests that these lncRNAs are potential

players in the processes of neurogenic lineage progression, neuronal maturation and transcription

factor regulation. Overexpression of Ebf2 altered the expression levels of some CR-specific

DElncRNAs but not others (Figure 3F), suggesting other regulatory factors might control their

expression. We also performed knockdown Foxg1, which only significantly up-regulated ln-CR5.

Some lncRNAs like ln-CR1 also had the trends of up regulation, while others had not, suggesting

that CR-specific lncRNAs might be involved in different aspects of early neurogenesis (Figure

S3B).

Overview of subtypes and differentiation states of CR neurons at the single-cell level

As early differentiating neurons located in the layer 1 of the embryonic cerebral cortex consist of

multiple cell types including CR neurons from different origins, we performed single-cell

RNA-seq (10X genomic Chromium) of 3000 E15.5 Ebf2-EGFP+ cells purified by FACS to

characterize individual gene expression profile and developmental states of early differentiating

neurons. After filtering out potential noises as multiplet cells and mitochondria genes, then

normalization and PCA analysis, the acquired gene expression data sets were selected for the

following analysis (Figure S4A-S4C). By K-Means clustering, Ebf2-EGFP positive cells at this

stage were molecularly clustered into eight major subpopulations, each showing distinct gene

expression pattern and with highly differentially expressed markers (Figure 4A). These eight

clusters can be visualized using the t-SNE plot (Figure 4B and 4C). We noticed that while Cluster

0, 3 and 4 had highly overlapped gene expression, Cluster 2 was the most segregated group from

the others. Interestingly, we found that 46% of E13.5 CR genes and 50.6% of P2 CR genes

identified from a previous study [25] were included in Cluster 2 (29 genes, and 44 genes,

respectively), indicating high enrichment of CR genes in this cluster.

To characterize the developmental state of the Ebf2+ neurons, we performed pseudo-time

reconstruction using Monocle analysis [41], which is an algorithm that cells with similar genes

expression pattern can be put together to identify sequential transcriptional changes in response to

various cell states along pseudo-time points. We chose genes that were expressed across more than

10 cells and their average expression levels were more than 1 for this analysis (Figure 4D; Figure

S4D). We found that the eight major cell subpopulations clustered from t-SNE plot corresponded

to the temporal cortical neuron development progress from early to late stages (Figure 4E). Genes

in Cluster 5 could be the first activated population during the pseudotemporal ordering,

sequentially followed by Cluster 3 and 7 genes, which displayed high expression levels at the

beginning of time period, later dropped to low levels. For example, Cluster 5 gene Neurod1 is

known to be expressed in committed neuronal progenitor cells at the upper SVZ and in newborn

neurons at the lower IZ, which plays a prominent role in cell fate specification [42, 43]. Cluster 3

gene Rorb is a marker of layer 4 neurons and also labels a subset of layer 5 neurons [44] (Figure

4E-4F). Thus, Cluster 5, 3 and 7 genes might be involved in initial neuronal differentiation. In

contrast, genes in Cluster 2 were expressed at low expression levels at the start side, but high

expression levels at the end of pseudotime. The top differential expressed genes in this cluster

such as Reln, Nhlh2, Lhx5 may represent a more committed CR neuron state. Genes in Cluster 7,

0, 6, 1, 4 displayed higher expression levels during the middle period of pseudotemporal ordering

than the start or the terminal. Genes in these clusters were correlated with regulation or negative

regulation of neuron differentiation (Tle4, Rnd3, Pou3f3) [45, 46] (Figure 4F; Figure S4E),

implying a function in progressive specification of neuronal cell fate. Taken together, our

scRNA-seq results unravel a previously unreported molecular trajectory of E15.5 CR neurons,

developing from general neuronal fate to committed CR neurons through suppressing other

cortical layer-associated genes.

Epigenetic regulation of Ebf2-EGFP+ cell-enriched genes and lncRNAs

Histone modification is a critical determinant of transcriptionally active or silent chromatin states

[47]. To gain a comprehensive view of histone modification in the early differentiating neurons, we performed H3K27ac and H3K27me3 chromatin immunoprecipitation followed by high-throughput sequencing (ChIP-seq) in Ebf2-EGFP+ cells during the early embryonic development (E11.5, E13.5, E15.5) and Ebf2-EGFP- cells at E11.5 (mostly neural stem cells and progenitor cells at this stage). We analyzed the occupancy of these histone modifications combined with RNA-seq data on these cell types in the followed analysis. First, our data show strong concordance in H3K27ac and H3K27me3 bindings across two biological replicates (Figures S5A-S5B). According to a general strategy used to classify REs (regulatory elements) [48-50], we classified H3K27ac and H3K27me3 bound TSS (± 5kb) by the pattern of histone modification as TSS activation (H3K27ac), repression (H3K27me3) and bivalent (H3K27ac, H3K27me3). Interestingly, we observed bivalent H3K27ac and H3K27me3 bindings at 58.3% of TSS sites of all detected DEGs in Ebf2-EGFP- cells (0.9% had only H3K27ac bindings, 4.5% had only H3K27me3 bindings). While the majority of genes (72.4% had only H3K27ac bindings, 0.5% had only H3K27me3 bindings, 16.4% had both bindings) in Ebf2-EGFP+ cells carried TSS activating modifications with strong H3K27ac enrichments and weak H3K27me3 enrichments (Figure 5A). Such observations were consistent with previous studies that bivalent histone modification patterns form a poised state to be ready for initiation of further developmental events in the ES cells or progenitor cells, while monovalent states occurred in the differentiated cells mean more organized states in the established specific cell identity [50, 51]. As development proceeded from E11.5 to E15.5, H3K27ac enrichments around TSS in Ebf2-EGFP+ cells became stronger. In contrast, H3K27ac enrichments around TSS in E11.5 neural progenitor cells were lowest compared to others. However, H3K27me3 bindings did not show temporal changes in Ebf2-EGFP+ cells (Figure 5A; Figure S5C-S5D). These patterns suggested that the epigenetic modifications were different between early differentiating neurons and neural progenitor cells and dynamically changed as Ebf2-EGFP+ cells differentiate.

We found that in Ebf2-EGFP+ cells, H3K27ac bindings were located in intergenic (61.3%), gene regions (62.2%, including 3'UTRs, 5'UTRs, Exons, Introns) and promoter regions (11.8%), while H3K27me3 bindings were located in intergenic (65.6%), gene regions (43.3%, including 3'UTRs, 5'UTRs, Exons, Introns) and promoter regions (3.9%) (Figure 5B; Figure S5E-S5F). Intergenic regions contained the highest percentage of both H3K27ac and H3K27me3 peaks,

likely implying potential distal regulatory elements. In addition, more peaks of H3K27ac were

located in promoter (11.8%) and 5'UTRs (11.0%) than H3K27me3 (promoter 3.9% and 5'UTRs

2.8%) in Ebf2-EGFP+ cells (Figure S5E-S5F), consistent with H3K27ac's role in transcription

regulation. In E11.5, genomic distribution patterns of H3K27ac and H3K27me3 peaks in

Ebf2-EGFP- cells were similar to that in Ebf2-EGFP+ cells (Figure S5G-S5J).

In the 1513 Ebf2+ genes enriched in Ebf2-EGFP+ cells, 97.2% had H3K27ac bindings and

63.8% had H3K27me3 bindings. 63.7% (964 genes) of them had both histone marks, 33.4% had

only H3K27ac bindings while 0.1% had only H3K27me3 bindings (Figure 5C). We further

investigated the distribution of these histone marks around the TSS and distal regions of the Ebf2+

genes. We found that for the H3K27ac bindings, 88% (1334) of Ebf2+ genes in Ebf2-EGFP+ cells

had peaks around TSS, 42% (639) had peaks around distal region (within 100kb). Interestingly,

the majority of Ebf2+ genes either had H3K27ac peaks only on TSS (55%, 831 genes) or on both

TSS and distal region (33%, 503 genes), while 9% (136) had enrichments only on distal region

(Figure 5D). Most CR marker genes and CR-specific expressed genes were occupied by H3K27ac

modification on TSS regions alone or together with distal sites. Conversely, more than half (56.3%,

851) of Ebf2+ genes had H3K27me3 peaks on distal regions, 18.1% (274) on TSS, 10.5% (159)

on both TSS and distal regions. We found that most CR marker genes and CR-specific expressed

genes such as Calb2, Tbr1, Ebf1, Lhx1, Lhx5, Zic5, Klf7 and Sncb, fall in the 45.8% (692) of

Ebf2+ genes that have H3K27me3 bindings only on distal regions. 7.6% (115) of Ebf2+ genes had

bindings only on TSS (Figure 5E), showing that more H3K27me3 enrichments on distal region of

Ebf2+ genes than on TSS. Notably, H3K27ac and H3K27me3 bindings on Ebf2+ genes

distributed on TSS and distal regions in E11.5 Ebf2-EGFP- cells were quite different from that in

Ebf2-EGFP+ cells. In E11.5 Ebf2-EGFP- cells, only 6% (90) and 6.4% (96) Ebf2+ genes had

H3K27ac peaks around TSS and distal region, respectively. However, there were 65.5% (989) and

25.8% (390) of Ebf2+ genes with H3K27me3 peaks around TSS and distal region, respectively

(Figure 5D-5E). Low H3K27ac and high H3K27me3 bindings around Ebf2+ genes in NPCs

implied that these genes were repressed, while they were activated in the Ebf2-EGFP+ cells.

Therefore, H3K27ac enrichments on TSS and distal sites, along with H3K27me3 enrichments on

distal region in Ebf2-EGFP+ cells might contribute to maintain or promote Ebf2+ genes to be

highly expressed in CR neurons.

The epigenomic data demonstrate that chromatin state round TSS is a strong indicator of RE

function [50]. Our results revealed that chromatin state at H3K27ac or H3K27me3-bound REs

were associated with differential expression of the nearest CR-specific genes. The dynamic

epigenomic signatures mentioned above in the regions around the genes and lncRNAs also

correlate with the changes in their expression levels in Ebf2-EGFP+ cells or NPC cells. For

example, B3gat1 and ln-CR2 expression increased in Ebf2-EGFP+ cells from E11.5 to E15.5,

accompanied by increasingly stronger H3K27ac enrichments at TSS in Ebf2-EGFP+ cells during

these stages. Meanwhile, these two genes are expressed at low levels in Ebf2-EGFP- cells on

E11.5, corresponding to H3K27me3 enrichments at TSS of these genes in Ebf2-EGFP- cells

(Figure 5F). Moreover, ln-CR2, which is increased in Ebf2-EGFP+ cells from E13.5 to E15.5,

exhibits a change in histone modification with no H3K27ac peaks on the TSS region on E13.5 but

high H3K27ac peaks on E15.5 (Figure 5F).

Functional analysis of CR-specific lncRNAs from *In Vitro* and *In Vivo* aspects

To further investigate what roles these lncRNA candidates may play during early neuronal

differentiation, we performed in vitro and in vivo functional assays using lentivirus-mediated

shRNA silencing and overexpression. We cultured single E11.5 cortical progenitor cells in

adherent conditions for 5 days (DIV5), then fixed the culture and stained for neuronal markers

β-III tubulin (immature neurons) and Reln (CR neurons) (Figure 6A). In the control culture there

were 29.89% Tuj1+ neurons, of which 3.16% were Reln+ (Figure 6B). Overexpression of ln-CR1,

ln-CR2 and ln-CR3 increased β-III tubulin+ cells and reduced total cell number (Figure 6C-6D).

Notably, Reln+ neurons were increased to 11.56% by ln-CR1 overexpression, 6.09% and 6.83%

by ln-CR2 and ln-CR3 overexpression, respectively (Figure 6B). Interestingly, the number of

small clones with less than 4 cells and all Tuj1+ increased in CAG-ln-CR1, CAG-ln-CR2 and

CAG-ln-CR3 compared to CAG-control, while the number of clones with more than 8 cells was

much less than that in control (Figure 6C and 6E). These results suggest that overexpression of

CR-specific lncRNAs promotes neuronal differentiation. Moreover, we also observed that

overexpression of ln-CR1 altered the morphology of Reln+ neurons, increased whole processes

length, branch depth and branch levels, while no significant differences were detected in neuron

soma size (Figure 6F).

During the postnatal stage, CR neurons develop more complex morphology. We further

analyzed the effect of ln-CR1 on cell morphology in an ex-vivo culture system using the

wholemount cortical hemispheres isolated from E15.5 Ebf2-EGFP+ embryos and cultured to P5.5

(DIV10). Compared to control, we found that overexpression of ln-CR1 increased the

morphological complexity of CR neurons, leading to more filopodia, spine-like structures (arrows)

and bouton-like structures (arrowheads) (Figure 6H; Figure S6A-S6B). We observed similar

results in P3.5 embryos that were ln-CR1 overexpressed by in utero electroporation into the layer

1 of the E15.5 cerebral cortex.

To uncover how overexpression of these CR-specific lncRNAs molecularly affected CR

neuron development, we assessed the mRNA level of several known CR neuron-related genes via

qPCR (Figure 6G). We found that CR marker genes including Reln, Ebf2, Calb2 and Trp73, and

genes that are expressed in CR neurons such as Ebf1, Ebf3, Tbr1, Nhlh2 and Lhx1, were

selectively altered when overexpressing CR-specific lncRNAs candidates ln-CR1, ln-CR2 and

ln-CR3 (Figure 6G), implying that these lncRNAs might be involved in the molecular pathways

for CR neuronal differentiation.

Discussion

In this study, we characterized the dynamic transcriptomic profile of early neurogenesis and

identified gene expression signatures associated with preplate neuronal differentiation and

subsequent establishment of CR cell properties. We also demonstrate that E15.5 CR cells could be

classified into eight subpopulations with three distinct maturation stages by single cell RNA-seq

analysis, which has not been reported before. Furthermore, we provide the first neuron subtype-

specific profile of histone modifications, revealing active, repressive or bivalent histone

modification states in the early differentiating neurons and neural progenitor cells that are

correlated with gene expression level. We have identified novel lncRNAs expressed in CR cells

and some of them are functionally important for CR cell differentiation. Our findings provide a

comprehensive molecular depiction of the early neuronal differentiation events, shedding new

light on the mechanism underlying preplate formation and differentiation of CR cells, the first step

of cortical neurogenesis.

Molecular transition from preplate to CR neurons

As cortical neurogenesis proceeds, the preplate is physically separated into the subplate and the layer 1 by migrating cortical plate neurons. We first identified over 1044 coding genes and 40 lncRNAs with higher expression in the newly differentiated preplate neurons (Ebf2-EGFP+) at E11.5 (Figure 1C; Figure 3B; Additional file 1 and Additional file 4). At E13.5, 370 genes and continue to show enrichment in preplate neurons, while additional 20 genes also have higher expression in subplate neurons. A subset of these genes continue to be differentially expressed in the Ebf2-EGFP+ cells at E15.5, while the total DEGs at E15.5 are much less than earlier stages, suggesting further specification of the CR cell fate. Indeed, we found fewer overlaps between the confirmed subplate neuron-specific genes at this age [17] (28 genes overlap, 2.3% of total DEGs at E15.5), confirming the molecular segregation of preplate neurons into the subplate and layer 1 by E15.5. The fact that the number of DEGs is the highest at E11.5 reflects the degree of the difference between the two populations: At E11.5, all Ebf2-EGFP+ cells are early differentiating neurons, while the Ebf2-EGFP- cells are mostly cortical progenitor cells. As the Ebf2-negative population includes both cortical plate neurons and neural progenitor cells at E13.5 and E15.5, genes common to neurons might not have significant changes.

This dynamic transcriptional profile may underlie the progression of preplate differentiation to the acquisition of unique molecular identity of CR neurons and subplate. For example, 166 genes are enriched in E11.5 and highly expressed in preplate, then later become restricted in CR neurons, while 851 genes which are also enriched in E11.5 but no longer specifically expressed in Layer 1 at E15.5.

Our transcriptome analysis also revealed a surprising enrichment of genes categorized in the GO terms such as "neuron projection", "neuron differentiation", "axonogenesis", even "synapse" and "synaptic transmission" as early as E11.5. Terms such as "gate channel activity" and "ion channel activity" also appear before E15.5. Interestingly, whole-cell patch-clamp recording showed an inward sodium current in putative CR cells as early as E12 [52]. In addition, dynamic expression of K+ channels and Ca2+ channels were also described in pre- and postnatal CR cells [11, 52-54]. We found that Ebf2-EGFP-positive cells expressed neurotransmitter receptors including ionotropic and metabotropic glutamate receptors, AMPA glutamate receptors, at E11.5. Interestingly, it has been reported that at very early stages of neural development, subplate neurons

form transient synapses with axons coming from the subcortical regions to guide later formation

of cortico-subcortical connectivity [55, 56]. Our results suggest that populations of early

differentiating neurons, including Cajal-Retzius cells, are equipped with machineries suited for

electric or chemical synaptic connections shortly after they are born. It's possible that functional

connections are established earlier than currently recognized. Alternatively, expression of these

synapse-related proteins may not necessarily confer electrophysiological properties, but rather

prime early differentiating neurons to be ready for synaptic connection once they encounter with

other cortical plate neurons.

Disease-related genes are highly expressed during early differentiation period [17, 57].

Abnormality in the process of preplate formation and splitting are associated with neurological

disorders including mental retardation, epilepsy, and autism [7]. Interestingly, schizophrenia has a

highly susceptible stage between 6-8 weeks gestation, the time when preplate differentiation

occurs [58]. These findings suggest the importance of proper regulation during the early

differentiation process.

Single-cell RNA-seq analysis reveals subpopulations of CR neurons

CR neurons were reported to be generated from many areas of developing pallium and were

redistributed on the cortical surface via tangential migration. However, it is not known whether

CR neurons from different origins maintain their original identity or develop to the same type of

cells with identical gene expression profile. In the first case, we expect to identify subpopulations

with distinct expression level of regional genes. For example, Trp73, a gene highly expressed in

cortical hem-derived and TE-derived CR cells [59], would select the cells originated from the

cortical hem or TE. Interestingly, we found that Trp73 is only expressed in one cell subpopulation,

with pattern similar to other CR genes such as Ebf2, CXCR4 (Additional file 6). We also found

that genes that are expressed in various CR cell origin territories were enriched in the same cluster.

For example, Trp73 (hem), Lhx1 (Septum), Emx2 (VZ) were all included Cluster 2 (Additional

file 6) [59-61]. Therefore, our results revealed loss of the original regional identity of CR neurons

once they migrate into the cerebral cortex, suggesting that molecular and functional maturation of

CR neurons may follow a general path regardless of their origins.

Pseudo-time analysis of single cell RNA-seq data has offered a way to further dissect the

transitional difference in gene expression within a heterogeneous population. Here, we reported

that by pseudo-time analysis, CR neurons could be classified into 3 subgroups. The first group

consisted of activated cluster 5, 3 and 7 genes, which were highly expressed at the beginning of

pseudotemporal period. We found that genes in this group are associated with characteristics of

cells located in the upper VZ and lower cortical plate such as NeuroD1, representing transitioning

from neural progenitor cells to newborn neurons [42-44]. The second group is enriched with

cluster 7, 6, 0, 1, 4 genes that were activated during the middle of pseudotime trajectory. These

genes were categorized under "neural differentiation" "neurite outgrowth" and "migration" terms

using DAVID analysis (data not shown). The third group along the pseudo-time line consisted of

mainly cluster 2 genes, which includes many CR marker genes, synapse and axonogenesis-related

genes. Interestingly, we found that genes first expressed in neural progenitor cells then in cortical

plate neurons, such as Tle4 and Pou3f3, were expressed higher first, then down-regulated in the

third group, suggesting when cells undergo fate decisions to become CR neurons, genes related to

other neuronal identity would need to be suppressed at the later differentiation stage. We found

that Zic2, a CR-enriched gene, has binding site in the promoter of Tle4 and neuroD1, raising the

possibility that genes in the third group might regulate non-CR genes to allow CR-specific traits to

be established. Further studies will help resolve the biological relevance of these analysis-acquired

developmental trajectory of gene expression. Overall, our study provides a comprehensive profile

of gene expression for the early born neurons in the cerebral cortex at population and single cell

level, unravelling the molecular dynamics during early neurogenesis.

Methods

Mouse

All animal protocols used in this study were approved by the IACUC (Institutional Animal Care

and Use Committee) of Tsinghua University and performed in accordance with guidelines of the

IACUC. The laboratory animal facility has been accredited by AAALAC (Association for

Assessment and Accreditation of Laboratory Animal Care International). For timed mice mating,

the day of the vaginal plug was considered E0.5.

Mapping and transcripts assembling

Sequencing reads were mapped to mouse genome using bowtie with parameter "-v 2 -X 350 -fr -m

10 -S -k 1". After mapping sequencing reads to the mouse genome, we assembled the transcripts

using Cufflinks [62]. We assembled transcripts into known gene models or novel gene models

guided by known gene annotations. Known gene annotations were downloaded from Ensembl

(Mus_musculus.GRCm38.74). The assembled transcripts were then merged by the Cuffmerge

utility provided by the Cufflinks package. Then the merged transcripts were concurrently

annotated by Cuffcompare program in Cufflinks suite, of which known transcripts were used as

reference to screen for novel lncRNAs. Only transcripts with tag "intergenic" and "intronic" were

retained. Transcripts shorter than 200nt or with FPKM lower than 0.3 were filtered out from the

transcripts set.

Clustering analysis and functional enrichment analysis

K-means clustering algorithm based on Euclidean distance was used to distinguish different gene

expression patterns, which was visualized by Java TreeView software

(http://bioinformatics.oxfordjournals.org/content/20/17/3246.abstract?keytype=ref&ijkey=31ed76

d5EFKWk). Then, we used findGO.pl program in Homer to perform functional enrichment (GO

annotation, KEGG and Wiki Pathway) analysis for each gene set with different expression patterns

[63]. GO analysis was also performed by DAVID website (https://david.ncifcrf.gov/tools.jsp;

version 6.8).

LncRNA differential expression

We used DE-seq to identify differential expressed lncRNAs. As input, the DESeq package

expected count data as obtained from RNA-Seq in the form of a rectangular table of integer values.

The table cell in the i-throw and the j-th column of the table told how many reads had been

mapped to gene i in sample j. The count values must be raw counts of sequencing reads. As a first

processing step, we needed to estimate the effective library size. This step was sometimes also

called normalization. The effective library size information was called the size factors vector,

since the package only needed to know the relative library sizes. If the counts of non-differentially

expressed genes in one sample were, on average, twice as high as in another (because the library

was sequenced twice as deeply), the size factor for the first sample should be twice that of the

other sample. The function estimateSizeFactors estimated the size factors from the count data. The

inference in DESeq relied on an estimation of the typical relationship between the data's variance

and their mean, or, equivalently, between the data's dispersion and their mean. The dispersion

could be understood as the square of the coeffcient of biological variation. The variance seen

between counts was the sum of two components: the sample-to-sample variation just mentioned,

and the uncertainty in measuring a concentration by counting reads. The latter, known as shot

noise or Poisson noise, was the dominating noise source for lowly expressed genes. The former

dominated for highly expressed genes. The sum of both, shot noise and dispersion, was considered

in the differential expression inference. Having estimated the dispersion for each gene, it was

straight-forward to look for differentially expressed genes. To contrast two conditions, e.g., to see

whether there was differential expression between conditions "positive" and "negative", we

simply called the function nbinom Test. It returned a data frame with the p-values and other useful

information. Genes and lncRNAs with q-value lower than 0.05 were regarded as differential

expressed.

Functional analysis of lncRNAs

To identify the function of lncRNAs, we calculated the expression correlation of lncRNAs

with coding genes. We assumed that co-expressed genes were more likely to be correlated in

function. Based on expression profiles in lncRNAs and protein coding genes, we calculated the

correlation between them. FPKM of genes and lncRNAs from RNA-seq data sets were used to

calculate the correlation matrix. Before calculating the correlation matrix, we filtered out some

low-expressed genes and lncRNAs for the accuracy of further characterization. For protein coding

genes with FPKM ranked in top 75% in at least one data set were retained. For lncRNAs, the

filtering processes were as follows: (1) expressed (FPKM>0) in at least 1 condition, (2) FPKM

ranked in top 75% in at least one datasets, (3) co-efficient of variation (cv) of FPKM ranked in top

75% in all ncRNA candidates. Cv represented for coefficient of variance, the standard deviation

divided by mean. Cv showed the extent of variability in relation to mean of the population.

Correlations between gene-gene and gene-lncRNA were calculated. We chose to calculate

Pearson correlation in the following analysis. The Pearson correlation coefficient was a measure

of the linear correlation between two variables X and Y, giving a value between +1 and −1

inclusive, where 1 was total positive correlation, 0 was no correlation, and −1 was negative

correlation. It was widely used in the sciences as a measure of the degree of linear dependence

between two variables. Pearson's correlation coefficient between two variables was defined as the

covariance of the two variables divided by the product of their standard deviations.

$$r = \frac{\sum (X - \overline{X})(Y - \overline{Y})}{\sqrt{\sum (X - \overline{X})^2 (Y - \overline{Y})^2}}$$

The calculation of correlation was done by SciPy, a fundamental python library for scientific

computing. The two-sided p-value for a hypothetical test whose null hypothesis was that two sets

of data were uncorrelated. The p-value roughly indicated the probability of an uncorrelated system

producing datasets that had a Pearson correlation at least as extreme as the one computed from

these datasets.

Based on the assumption that functionally correlated genes were likely to co-express in

various conditions, the functional prediction or annotation of lncRNAs could be performed. One

of the most direct way was to look at the functional annotations of genes in its immediate

neighborhood. For each lncRNAs, we chose top 1000 correlated genes to perform Gene Ontology

analysis. We used a R package GOstats to perform GO analysis. The enriched GO term could be

used to illustrate the possible function of lncRNAs.

Quantitative RT-PCR

Total RNA was isolated from purified Ebf2-EGFP positive or negative cells, extracted using Trizol

reagent (Cat#15596018, Life Technologies), and reverse transcribed. qPCR was performed using

PrimeScript RT regent Kit (Perfect Real Time, Cat#DRR047A, Takara). Gene expression levels

were detected using a BioRad CFX384 Touch™ real-time PCR detection system (Bio-Rad

Laboratories) and determined with ΔΔCt method, normalized to an internal control GAPDH

mRNA for quantification. All reactions were replicated 3 times. qRT-PCR primers for mouse gene

sequence were included in the Additional file 7.

Tissue collection and RNA In Situ hybridization

Mouse brains at various developmental stages were collected and fixed in 4% paraformaldehyde

(PFA) at 4°C for 4h. After briefly rinsing in PBS, brains were dehydrated in 30% sucrose in PBS

at 4°C until the time they sank, and embedded in OCT (Sakura Finetek, Torrance, CA). Frozen

brains were cryosectioned to 16 μm and mounted to coverslip. In situ hybridization was performed

according to instructions as previously described (Wigard et al., 2006; Gregor et al., 2007; Robert

et al., 2007). Digoxigenin-labeled RNA probes were designed based on lncRNA sequences

determined by Cufflinks and Ensembl. Images were taken on a Zeiss Slide Scanner (Axio Scan.

Z1) fluorescent microscope.

In utero electroporation of cortical layer 1

The E15.5 embryonic cortical layer 1 were electroporated as described previously [64]. Briefly,

embryonic day 15.5 (E15.5) mice were anesthetized and placed on a heating pad. Then the uterus

were carefully pulled out and injected with proper volume of plasmid mix (0.5-1μl/embryo, 2-3

μg/μl) into the gap under the skull and above the pial surface with a glass capillary. The

Electroporator (CUY21VIVO-SQ; BEX) was set to five 50ms, 36V pulses with an interval of

999ms, and embryos' heads were electroporated with a 5-mm electrode (LF650P5; BEX). After

that, the embryos were placed back into their original location of abdominal cavity, and the

abdomen wall and skin were closed. Then waited for a while until the mice recovered from

surgery. For phenotype observation, the mice progenies were sacrificed on postnatal day 3.5 or 5.5

(P3.5 or P5.5), and both Ebf2-EGFP and RFP signals positive pups were selected for the following

PFA fixation dealing and immunohistochemistry. Confocal images were taken using LSM780

(Zeiss) Laser Scanning System on a Zeiss microscope. At least three independent experiments,

three animals each experiment were analyzed for each condition.

Wholemount culture neuronal processes analysis

For neuronal processes and spine analysis, Ebf2-EGFP+ wholemounts of E15.5 cerebral cortex

were infected by lenti-virus CAG-RFP, H1-RFP, or CAG-ln-CR1, CAG-ln-CR2, CAG-ln-CR3.

Next, wholemounts were cultured for 12 days in a 35°C bioincubator. The culture medium was

replaced every 3 days. After culture, wholemounts were fixed and stained with chicken anti-GFP

(GFP-1010, Aves lab lnc.) and rat anti-chicken-Alexa488 (Jackson Immuno Research lab).

Confocal microscopy was performed with a LSM780 (Zeiss) Laser Scanning System on a Zeiss

microscope, and wholemount tissues were analyzed by taking 1μm serial optical sections with

ZEN_2011_Lite_x64 software (Zeiss). Images were acquired using a 1024x1024 scan format with

a 63x objective.

Immunocytochemistry

Cell culture media were removed and cells were fixed in 4% (w/v) PFA for 30min at RT, rinsed by

1×PBS three times. Fixed cells were blocked in 5% (v/v) BSA with 0.3% (v/v) Triton X-100 in

1×PBS for 1h and rinsed by 1×PBS three times. Immunostainings were performed for 4h at RT or

overnight at 4°C with rotation by incubating with primary antibodies, then washed by 1×PBS

three times. Cells were incubated with secondary antibodies and DAPI diluted in 1×PBS for 1h in

the dark, followed by washing with 1×PBS three times. Images were taken on Zeiss fluorescent

microscopy (AxioVision). Primary antibodies used: GFP (GFP-1010, Aves lab lnc.), Reln

(MAB5364, Millipore), Tuj1 (T8660, Sigma).

Chromatin-immunoprecipitation and sequencing (ChIP-Seq)

ChIP assays were performed using mice primary Ebf2-EGFP positive cells at E11.5, E13.5, E15.5

and Ebf2-EGFP negative cells at E11.5, with antibody against H3K27ac (Ab4729, Abcam) and

H3K27me3 (Ab6002, Abcam), experimental procedures were as previously described [65]. Briefly,

Ebf2-EGFP positive and negative cells (~1-5 million cells) were sorted and collected from FACS

(BD Influx^{TM}), and fixed by 0.4% (v/v) formaldehyde at room temperature (RT) for 10 min. The

reaction was stopped by adding 0.125 M glycine for 5 min. After rinsing cells three times with

cold PBS, the nuclei were resuspended in Nuclei Lysis Buffer (50 mM Tri-Cl, pH8.1, 10 mM

EDTA, 1% SDS, 1 mM DTT, 1 mM PMSF and Sigma protease inhibitor cocktail, RT) and

incubated on ice for 10 min. Nuclear suspension were sonicated 6" on, 15" off, 20 cycles (power

set at 25%) on ice. The fragmented chromatin was diluted for 5 folds in ChIP Dilution Buffer

(16.7 mM Tri-Cl, pH8.1, 167 mM NaCl, 1.2 mM EDTA, 1.1% Triton X-100, 1 mM PMSF, and

Sigma protease inhibitor cocktail). Then 5 ug IP antibody was added and incubated overnight at

4 °C. Chromatin-protein complex was pulled down with Dynabeads protein G (Cat#10004D,

Invitrogen) for 2h at 4°C with rotation and followed by 5 sequential washes as the following: 1.

low salt wash buffer (20 mM Tri-Cl, pH8.1, 150 mM NaCl, 1 mM EDTA, 1% Triton X-100, 0.1%

SDS, 0.1% Na-deoxycholate); 2. high salt wash buffer (20 mM Tri-Cl, pH8.1, 500 mM NaCl, 1

mM EDTA, 1% Triton X-100, 0.1% SDS, 0.1% Na-deoxycholate); 3. LiCl wash buffer (10 mM

Tri-Cl, pH8.1, 250 mM LiCl, 1 mM EDTA, 0.5% NP-40, 0.5% Deoxycholic acid (Na salt)); and

the last two washes with TE buffer (10 mM Tri-Cl, pH8.1, 1 mM EDTA). Immunoprecipitated

complex was eluted from the antibody in elution buffer (1% SDS, 50 mM TrisCl (pH 8.1), 1 mM

EDTA) by votexing once at 65℃ for 20min and once at RT. Eluted chromatins were incubated at

65℃ overnight for reverse crosslinks and chromatin fragmented purification with PCR column

purification. ChIP-Seq libraries were prepared using NEB Next Ultra™ II ChIP-Seq Library Prep

Kit (Cat#E7645S & Cat#7335S, NEB) according to the manufacturer's instructions. ChIP-seq data

were aligned to mouse genome mm10, performing Bowtie v. 1.1.2 on the Galaxy platform with

the following options: -v 3 –M 1 -5 15. Resulting output SAM files were converted to BAM

format, and finally converted to BED files. Peaks were called using MACS (Model-Based

Analysis of ChIP-Seq), with a p-value cutting off of 1e-5.

Acknowledgments

We thank Dr. Zhengang Yang and Dr. Chunyang Wang at Fudan University Collaborative

Innovation Center for Brain Science for their assistance of In Situ hybridization experiments in

this study. We thank Yuchuan Wang, Chao Di and Xinqiang Ding for their help of bioinformatic

analysis. We thank for all members of Shen lab for their helpful discussion during the preparation

of manuscript. We are also grateful to Center of Biomedical Analysis at Tsinghua University.

Funding

This work was supported by grants from the National Natural Science Foundation of China

(31170247 to Q.S.).

Availability of data and materials

The sequencing data generated in this study will be deposited in GEO repository before

manuscript accepted.

Author's contribution

JL and QS designed the experiments. JL, LS, X-L P and S-J Q performed the experiments, JL and X-M Y analyzed sequencing data. JL and QS wrote the manuscript. QS supervised the project.

Ethics approval and consent to participate

The study was approved by the IACUC (Institutional Animal Care and Use Committee) of Tsinghua University. All animal experiments were performed in accordance with their guidelines.

Consent for publication

The authors declare that they have no competing interests.

Competing interests

Not applicable.

Figure Legends

Fig 1. Transcriptome sequencing of Ebf2-EGFP+ and Ebf2-EGFP- cells in the early embryonic stage

(A) FACS purification of Ebf2-EGFP+ and Ebf2-EGFP- cells from early embryonic stages, E11.5, E13.5 and E15.5 respectively. The percentage of Ebf2-EGFP+ cells at each stage was calculated and shown in the graph.

(B) Expression levels of all differential expressed genes (FDR<0.05) in Ebf2-EGFP+ & Ebf2-EGFP- samples at the three early embryonic developmental stages are shown.

(C) Venn diagrams displaying the extent of overlapping between DEGs at the three early embryonic stages.

(D) Expression of selected CR-specific coding genes are found to be specifically expressed in cortical layer 1 of E14.5 mouse embryos (In situ hybridization images from Eurexpress Anatomy Atlas and GenePaint.org).

(E) qRT-PCR analysis comparing the expression levels of CR-specific genes in NSC culture assay

after virus infection with control (H1) or Ebf2-overexpression. Data represent mean ± SEM (n=3

independent experiments, *P<0.05, T test).

Fig 2. Functional prediction of DEGs in the Ebf2-EGFP+ cells and Ebf2-EGFP- cells

(A) Enriched functional assessments (including biological processes, cellular components and

molecular function) of highly expressed DEGs in Ebf2-EGFP+ or Ebf2-EGFP- cell population

(fold change $\geqslant$ 2).

(B-D) Enriched functional assessments of highly expressed DEGs in E11.5, E13.5 or E15.5

Ebf2-EGFP+ cell population, respectively.

(E-F) Significantly (p<0.001, FDR<0.001) enriched KEGG pathways within DEGs from all

RNA-seq samples and DEGs enriched in Ebf2-EGFP+ samples.

(G) Temporal dynamic expression pattern of DEGs at the three different stages (y axis = Fold

Change between Ebf2-EGFP+ cells and Ebf2-EGFP- cells). Noted the dynamic pattern is related

to functions of associated genes.

Fig 3. Functional prediction of differentially expressed lncRNAs in the Ebf2-EGFP+ and

Ebf2-EGFP- cells

(A) Heatmap representing all differential expressed lncRNAs (FDR<0.05, p-value<0.05) in

Ebf2-EGFP+ & Ebf2-EGFP- cell samples at the three embryonic stages.

(B) Venn diagram displaying the extent of overlapping between DElncRNAs at the three

embryonic stages.

(C) The expression pattern of DElncRNAs between Ebf2-EGFP+ or Ebf2-EGFP- cells is

consistent revealed by RNA-seq and qRT-PCR.

(D) In situ hybridization showing the endogenous expression level of CR-specific lncRNA ln-CR2

at E11.5, E13.5 and E15.5. Regions with specific expression are labeled and higher magnification

images of boxed areas are shown below. Abbreviations: DP, dorsal pallium; VP, ventral pallium;

MP, medial pallium; Th, thalamus; Str, striatum; TE, thalamus eminence.

(E) Enriched functional assessments (including biological processes, cellular components and

molecular function) of highly expressed DElncRNAs in Ebf2-EGFP+ or Ebf2-EGFP- cell

population (fold change ≥ 2).

(F) qRT-PCR analysis comparing the expression levels of lncRNAs in NSC culture assay after

virus infection with control (H1) or Ebf2-overexpression. Data represent mean ± SEM (n=4

independent experiments, *P<0.05, T test).

Fig 4. Single cell overview of Cajal-Retzius neuron subtypes

(A) Heatmap of differentially expressed genes in E15.5 Ebf2-EGFP+ cells sub-clusters.

(B) The K-means algorithm-based plot showing eight major populations.

(C) Visualization of the eight major sub-classes of cells using t-SNE clustering. Each dot

represents a single cell. Dot color represents expression level (from grey to blue, expression levels

from low to high).

(D) Pseudotime trajectories of Ebf2-EGFP+ cells reconstructed by Monocle. Cell states are

indicated by colors and numbers in the black circle.

(E) Heatmap shows gene dynamics along pseudotemporal cortical neuron differentiation. X-axis

shows pseudotime ordering from early to late, Y-axis shows a single gene expression level

dynamic per row. Gene expression level from high to low is indicated by colors from red to blue.

(F) Expression level dynamics across pseudotime ordering of Lhx5 (in Cluster 2) and Neurod1

(in Cluster 5) are shown.

Fig 5. Genomic characterization of histone modification in the early differentiating neurons

and NPCs

(A) Heatmap representation of ChIP-seq signal density for H3K27ac and H3K27me3 for 10 kb

centered on predicted Gene TSSs in Ebf2-EGFP- (E11.5) cells and Ebf2-EGFP+ (E11.5, E13.5

and E15.5).

(B) Genomic features of H3K27ac and H3K27me3-bound REs in Ebf2-EGFP+ cells.

(C) Comparison of genes number with H3K27ac and H3K27me3 bindings in Ebf2-EGFP+ cells.

(D) Proportion of Ebf2-EGFP+ cell-enriched genes with TSS and distal H3K27ac bindings in

Ebf2-EGFP+ cells (upper) and NPCs (lower).

(E) Proportion of Ebf2+ genes with TSS and distal H3K27me3 bindings in Ebf2-EGFP+ cells

(upper) and NPCs (lower).

(F) The TSS regions of CR-specific genes B3gat1, and CR-specific lncRNAs ln-CR2 demonstrate

a similar pattern of histone modification during early embryonic development (bivalent in NPCs

and become more activated in Ebf2-EGFP+ cells from E11.5 to E15.5).

Fig 6. Functional validation of CR-specific lncRNAs ln-CR1, ln-CR2 and ln-CR3 *In Vitro*

and *In Vivo*

(A) Immunostaining of NSCs treated with control (CAG-control) or overexpression lentiviruses

for CR-specific lncRNAs ln-CR1 (CAG-ln-CR1), ln-CR2 (CAG-ln-CR2) and ln-CR3

(CAG-ln-CR3) after in vitro culture for 5 days. Reln (red), Tuj1 (cyan), or DAPI (blue) for CR

neurons, neurons, and nucleus, respectively. Scale bar, left 3 rows 150 μm, right 50 μm.

(B-E) Overexpression of CR-specific lncRNAs significantly increased CR neuron number,

promoted NSC differentiation and generated more neuron-like small cell clones. All the

experiments were performed 4 times and 8 randomly selected microscopy fields were counted

each time. Data represented mean ± SEM. p values indicated were calculated by Student's t test

(unpaired). **p<0.01, *p<0.05.

(F) Quantification of process branch depth, branch levels and total dendrites length and soma size

of CR neurons after overexpression of CR-specific lncRNAs compared to control. All the

experiments were performed 3 times and 8 randomly selected microscopy fields were counted

each time. Data represented mean ± SEM. p values indicated were calculated by Student's t test

(unpaired). **p<0.01, *p<0.05.

(G) Quantification of CR neuron molecular markers as Reln, Ebf2, and Calb2 at 72h after

lentiviral transduction in the NSC culture assay comparing the effect of overexpression of

CR-specific lncRNAs to control. Data represented mean ± SEM (**p<0.01, *p<0.05, n>3,

Student's T test).

(H) Comparison of dendritic spine, filopodia and processes in cortical wholemounts

overexpressing ln-CR1 (CAG-ln-CR1) to CAG-control. Scale bar, 20 μm. ln-CR1 overexpression

resulted in increased in spine and filopodia (arrows) as compared to CAG-control transfected CR

neurons. Cortical wholemounts were collected from P3 pups after *in utero* electroporation to

cortex layer 1 on E15.5.

Additional files

Fig S1. RNA-seq quality control, related to Fig 1

(A) Phase and fluorescent images of Ebf2-EGFP+ cells and Ebf2-EGFP- cells after FACS at Day

0 and Day 4 in vitro culture showing the purity and growth of sorted and unsorted cells (scale bar,

25 μm).

(B) Sample to sample Pearson's correlation (top-right quadrants) and pairwise comparison

(bottom-left quadrants) of normalized gene expression between the two biological replicates and

cell types.

(C) qRT-PCR and endogenous expression patterns (In Situ Hybridization data from Allen Brain

Atlas) to validate the RNA-seq results at different embryonic stages. CR neuron molecular

markers as Calb2, Ebf2, Tbr1, and CR neuron related genes as Ebf1, Ebf3 and Foxg1 were

selected for validation.

(D) qRT-PCR and endogenous expression patterns (In Situ Hybridization data from Allen Brain

Atlas) to validate the RNA-seq results of Nhlh2 at three embryonic stages.

(E) Expression of the CR-specific coding gene Dnm3 is found to be specifically expressed in layer

1 of E14.5 mouse brain (In Situ Hybridization data from Eurexpress Anatomy Atlas).

(F-G) qRT-PCR analysis of CR-specific genes in NSCs culture assay after lentiviral transduction

of control (H1) or Ebf2-knockdown, or Foxg1-knockdown. Data represent mean ± SEM (n=3

independent experiments, *P<0.05, T test).

Fig S2. Annotation of temporal dynamic genes, related to Fig 2

(A) Gene Ontology analysis of selected genes with distinct temporal dynamic expression pattern.

Fig S3. Functional prediction of CR-specific DElncRNAs, related to Figure 3

(A) qRT-PCR analysis to validate the RNA-seq results at different embryonic stages. CR-specific

lncRNAs ln-CR6 and Meg3 were selected for test.

(B) qRT-PCR analysis of CR-specific lncRNAs in NSCs culture assay after lentiviral transduction

of control (H1) or Foxg1-knockdown. Data represent mean ± SEM (n=4 independent experiments,

*P<0.05, T test).

Fig S4. Quality control and pseudotime reconstruction analysis of single cell RNA-seq,

related to Fig 4

(A) Violin plots show genes, UMIs and mitochondrion ratio in scRNA-seq. Left, violin plot

shows detected gene number in all sample cells. X-axis is sample name and Y-axis is gene

numbers detected in a single cell; Middle, X-axis is the same with left and Y-axis is UMIs detected

in a single cell; Right, Y-axis is mitochondrion numbers detected in a single cell.

(B) Scatter diagram of mitochondrion ratio, gene numbers compared to UMI numbers,

respectively. X-axis is UMI numbers, Y-axis is mitochondrion ratio detected in a single cell in the

left, Y-axis is gene numbers detected in a single cell in the right.

(C) Scatter diagram of all detected genes dispersion. X-axis is average expression of a gene in all

cells and Y-axis is dispersion value of that gene.

(D) Pseudotime trajectories of E15.5 Ebf2-EGFP+ cells reconstructed by Monocle. Pseudotime

points are indicated by colors and cell clusters are indicated by numbers in the black circle.

(E) Expression level dynamics of Reln (in Cluster 2) and Tle4 (in Cluster 4) across pseudotime

ordering.

Fig S5. Chromatin signatures in Ebf2-EGFP+ cells and NPCs, related to Fig 5

(A-B) Average profile of H3K27ac (A) or H3K27me3 (B) binding at called ChIP-seq peaks

comparing the two biological replicates.

(C-D) A profile plot showing normalized H3K27ac (C) or H3K27me3 (D) enrichment around TSS

of all genes at the three embryonic stages.

(E-J) Quantitative data showing genomic features of H3K27ac and H3K27me3-bound REs in

Ebf2-EGFP+ cells (E, F), E11.5 NPCs (G, H) and E11.5 Ebf2-EGFP+ cells (I, J).

Fig S6. LncRNAs function in CR neurons in ex vivo wholemount culture assay, related to Fig

6

(A) Comparison of dendritic spine, filopodia and processes of cultured cortical wholemounts

overexpressing ln-CR1 (CAG-ln-CR1), ln-CR2 (CAG-ln-CR2), ln-CR3 (CAG-ln-CR3) and

knockdown ln-CR2 (shRNA-ln-CR2) to wholemounts expressing CAG-control. Scale bar, 15 μm.

(B) Quantification of CR neuron processes branch depth, branch levels and total dendrites length

after overexpression of CR-specific lncRNAs. All the experiments were performed 3 times and 4

randomly selected microscopy fields were counted each time. Data represent mean ± SEM. p

values indicated were calculated by Student's t test (unpaired). **$p<0.01$, *$p<0.05$.

References

1.   Caviness VS. Neocortical Histogenesis in Normal and Reeler Mice: A Developmental Study
Based Upon [3H]Thymidine Autoradiography. Developmental Brain Research. 1982;4:293-302.
2.   Marin-Padilla M. Ontogenesis of the Pyramidal Cell of the Mammalian Neocortex and
Developmental Cytoarchitectonics: A Unifying Theory. J Comp Neurol. 1992;321:321-3.
3.   Hevner RF, Neogi T, Englund C, Daza RM, Fink A. Cajal–Retzius cells in the mouse:
transcription factors, neurotransmitters, and birthdays suggest a pallial origin. Developmental Brain
Research. 2003;141:39-53.
4.   Bayer SA, Altman J. Development of the Endopiriform Nucleus and the Claustrum in the Rat
Brain. Neuroscience. 1991;45:391-412.
5.   Takahashi T, Goto T, Miyama S, Nowakowski RS, Caviness VS Jr. Sequence of Neuron Origin
and Neocortical Laminar Fate: Relation to Cell Cycle of Origin in the Developing Murine Cerebral
Wall. J Neurosci. 1999;19:10357-71.
6.   Bystron I, Blakemore C, Rakic P. Development of the human cerebral cortex: Boulder Committee
revisited. Nat Rev Neurosci. 2008;9:110-22.
7.   Olson EC. Analysis of preplate splitting and early cortical development illuminates the biology of
neurological disease. Front Pediatr. 2014;2:1-9.
8.   Ghosh A, Antonini A, McConnell SK, Shatz CJ. Requirement of Subplate Neurons in the
Formation of Thalamocortical Connections. Nature. 1990;347:179-181.
9.   Molnar Z, Adams R, Blakemore C. Mechanisms Underlying the Early Establishment of
Thalamocortical Connections in the Rat. J Neurosci. 1998;18:5723-45.
10.   Del Rio JA, Martinez A, Fonseca M, Auladell C, Soriano E. Glutamate-like Immunoreactivity and
Fate of Cajal-Retzius Cells in the Murine Cortex as Identified with Calretinin Antibody. Cereb Cortex.
1995;5:13-21.
11.   Kirischuk S, Luhmann HJ, Kilb W. Cajal-Retzius cells: update on structural and functional

properties of these mystic neurons that bridged the 20th century. Neuroscience. 2014;275:33-46.

12. Hong SE, Shugart YY, Huang DT, Shahwan SA, Grant PE, Hourihane JOB, Martin NDT, Walsh CA. Autosomal recessive lissencephaly with cerebellar hypoplasia is associated with human RELN mutations. Nature Genetics. 2000;26:93-6.

13. Nakagawa N, Yagi H, Kato K, Takematsu H, Oka S. Ectopic clustering of Cajal-Retzius and subplate cells is an initial pathological feature in Pomgnt2-knockout mice, a model of dystroglycanopathy. Sci Rep. 2015;5:11163.

14. Oeschger FM, Wang WZ, Lee S, Garcia-Moreno F, Goffinet AM, Arbones ML, Rakic S, Molnar Z. Gene expression analysis of the embryonic subplate. Cereb Cortex. 2012;22:1343-59.

15. Osheroff H, Hatten ME. Gene expression profiling of preplate neurons destined for the subplate: genes involved in transcription, axon extension, neurotransmitter regulation, steroid hormone signaling, and neuronal survival. Cereb Cortex. 2009;19 Suppl 1:i126-34.

16. Hoerder-Suabedissen A, Wang WZ, Lee S, Davies KE, Goffinet AM, Rakic S, Parnavelas J, Reim K, Nicolic M, Paulsen O, Molnar Z. Novel markers reveal subpopulations of subplate neurons in the murine cerebral cortex. Cereb Cortex. 2009;19:1738-50.

17. Hoerder-Suabedissen A, Oeschger FM, Krishnan ML, Belgard TG, Wang WZ, Lee S, Webber C, Petretto E, Edwards AD, Molnar Z. Expression profiling of mouse subplate reveals a dynamic gene network and disease association with autism and schizophrenia. Proc Natl Acad Sci U S A. 2013;110:3555-60.

18. Belgard TG, Marques AC, Oliver PL, Abaan HO, Sirey TM, Hoerder-Suabedissen A, Garcia-Moreno F, Molnar Z, Margulies EH, Ponting CP. A transcriptomic atlas of mouse neocortical layers. Neuron. 2011;71:605-16.

19. Gong S, Zheng C, Doughty ML, Losos K, Didkovsky N, Schambra UB, Nowak NJ, Joyner A, Leblanc G, Hatten ME, Heintz N. A gene expression atlas of the central nervous system based on bacterial artificial chromosomes. Nature. 2003;425:917-925.

20. Chuang SM, Wang Y, Wang Q, Liu KM, Shen Q. Ebf2 marks early cortical neurogenesis and regulates the generation of cajal-retzius neurons in the developing cerebral cortex. Dev Neurosci. 2011;33:479-93.

21. Chiara F, Badaloni A, Croci L, Yeh ML, Cariboni A, Hoerder-Suabedissen A, Consalez GG, Eickholt B, Shimogori T, Parnavelas JG, Rakic S. Early B-cell factors 2 and 3 (EBF2/3) regulate early migration of Cajal-Retzius cells from the cortical hem. Dev Biol. 2012;365:277-89.

22. Macosko EZ, Basu A, Satija R, Nemesh J, Shekhar K, Goldman M, Tirosh I, Bialas AR, Kamitaki N, Martersteck EM, et al. Highly Parallel Genome-wide Expression Profiling of Individual Cells Using Nanoliter Droplets. Cell. 2015;161:1202-1214.

23. Zeisel A, Muñoz-Manchado AB, Codeluppi S, Lönnerberg P, La Manno G, Juréus A, Marques S, Munguba H, He LQ, Hjerling-Leffler J, Linnarsson S. Cell types in the mouse cortex and hippocampus revealed by single-cell RNA-seq. Science. 2015;347:1138-42.

24. Shin J, Berg DA, Zhu Y, Shin JY, Song J, Bonaguidi MA, Enikolopov G, Nauen DW, Christian KM, Ming GL, Song H. Single-Cell RNA-Seq with Waterfall Reveals Molecular Cascades underlying Adult Neurogenesis. Cell Stem Cell. 2015;17:360-72.

25. Yamazaki H, Sekiguchi M, Takamatsu M, Tanabe Y, Nakanishi S. Distinct ontogenic and regional expressions of newly identified Cajal–Retzius cell-specific genes during neocorticogenesis. PNAS. 2004;101:14509-14.

26. Hanashima C, Li SC, Shen L, Lai E, Fishell G. Foxg1 Suppresses Early Cortical Cell Fate.

Science. 2004;303:56-59.

27. Shen Q, Wang Y, Dimos JT, Fasano CA, Phoenix TN, Lemischka IR, Ivanova NB, Stifani S,
Morrisey EE, Temple S. The timing of cortical neurogenesis is encoded within lineages of individual
progenitor cells. Nat Neurosci. 2006;9:743-51.

28. Hevner RF, Shi L, Justice N, Hsueh YP, Sheng M, Smiga S, Bulfone A, Goffinet AM,
Campagnoni AT, Rubenstein JL. Tbr1 Regulates Differentiation of the Preplate and Layer 6. Neuron.
2001;29:353-366.

29. Siegenthaler JA, Miller MW. Generation of Cajal-Retzius neurons in mouse forebrain is regulated
by transforming growth factor beta-Fox signaling pathways. Dev Biol. 2008;313:35-46.

30. Meyer G. Building a human cortex: the evolutionary differentiation of Cajal-Retzius cells and the
cortical hem. J Anat. 2010;217:334-43.

31. Corradi A, Croci L, Broccoli V, Zecchini S, Previtali S, Wurst W, Amadio S, Maggi R, Quattrini A,
Consalez GG. Hypogonadotropic hypogonadism and peripheral neuropathy in Ebf2-null mice.
Development. 2003;130:401-410.

32. Tedeschi A, Dupraz S, Laskowski CJ, Xue J, Ulas T, Beyer M, Schultze JL, Bradke F. The
Calcium Channel Subunit Alpha2delta2 Suppresses Axon Regeneration in the Adult CNS. Neuron.
2016;92:419-434.

33. Buscarlet M, Perin A, Laing A, Brickman JM, Stifani S. Inhibition of Cortical Neuron
Differentiation by Groucho/TLE1 Requires Interaction with WRPW, but Not Eh1, Repressor Peptides.
Journal of Biological Chemistry. 2008;283:24881-8.

34. Inoue T, Ogawa M, Mikoshiba K, Aruga J. Zic deficiency in the cortical marginal zone and
meninges results in cortical lamination defects resembling those in type II lissencephaly. J Neurosci.
2008;28:4712-25.

35. Segklia A, Seuntjens E, Elkouris M, Tsalavos S, Stappers E, Mitsiadis TA, Huylebroeck D,
Remboutsika E, Graf D. Bmp7 regulates the survival, proliferation, and neurogenic properties of neural
progenitor cells during corticogenesis in the mouse. PLoS One. 2012;7:e34088.

36. Tiberi L, van den Ameele J, Dimidschstein J, Piccirilli J, Gall D, Herpoel A, Bilheu A, Bonnefont
J, Iacovino M, Kyba M, et al. BCL6 controls neurogenesis through Sirt1-dependent epigenetic
repression of selective Notch targets. Nat Neurosci. 2012;15:1627-35.

37. Aprea J, Prenninger S, Dori M, Ghosh T, Monasor LS, Wessendorf E, Zocher S, Massalini S,
Alexopoulou D, Lesche M, et al. Transcriptome sequencing during mouse brain development identifies
long non-coding RNAs functionally involved in neurogenic commitment. EMBO J. 2013;32:3145-60.

38. Nishi T, Forgac M. The vacuolar (H+)-ATPases--nature's most versatile proton pumps. Nat Rev
Mol Cell Biol. 2002;3:94-103.

39. Ueda T, Ugawa S, Shimada S. A novel putative M9.2 isoform of V-ATPase expressed in the
nervous system. Neuroreport. 2003;14:25-30.

40. Boal F, Hodgson LR, Reed SE, Yarwood SE, Just VJ, Stephens DJ, McCaffrey MW, Tavare JM.
Insulin promotes Rip11 accumulation at the plasma membrane by inhibiting a dynamin- and
PI3-kinase-dependent, but Akt-independent, internalisation event. Cell Signal. 2016;28:74-82.

41. Trapnell C, Cacchiarelli D, Grimsby J, Pokharel P, Li S, Morse M, Lennon NJ, Livak KJ,
Mikkelsen TS, Rinn JL. The dynamics and regulators of cell fate decisions are revealed by
pseudotemporal ordering of single cells. Nat Biotechnol. 2014;32:381-386.

42. Hevner RF, Hodge RD, Daza RA, Englund C. Transcription factors in glutamatergic neurogenesis:
conserved programs in neocortex, cerebellum, and adult hippocampus. Neurosci Res. 2006;55:223-33.

43. Lee JE, Hollenberg SM, Snider L, Turner DL, Lipnick N, Weintraub H. Conversion of Xenopus
Ectoderm into Neurons by NeuroD, a Basic Helix-Loop-Helix Protein. Science. 1995;268:836-44.

44. Schaeren-Wiemers N, Andre E, Kapfhammer JP, Becker-Andre M. The Expression Pattern of the
Orphan Nuclear Receptor RORb in the Developing and Adult Rat Nervous System Suggests a Role in
the Processing of Sensory Information and in Circadian Rhythm. Eur J Neurosci. 1997;9:2687-701.

45. Pacary E, Heng J, Azzarelli R, Riou P, Castro D, Lebel-Potter M, Parras C, Bell DM, Ridley AJ,
Parsons M, Guillemot F. Proneural transcription factors regulate different steps of cortical neuron
migration through Rnd-mediated inhibition of RhoA signaling. Neuron. 2011;69:1069-84.

46. Oishi K, Aramaki M, Nakajima K. Mutually repressive interaction between Brn1/2 and Rorb
contributes to the establishment of neocortical layer 2/3 and layer 4. Proc Natl Acad Sci U S A.
2016;113:3371-6.

47. Jenuwein T, Allis CD. Translating the Histone Code. Science. 2001;293:1074-1080.

48. Attanasio C, Nord AS, Zhu Y, Blow MJ, Biddie SC, Mendenhall EM, Dixon J, Wright C, Hosseini
R, Akiyama JA, et al. Tissue-specific SMARCA4 binding at active and repressed regulatory elements
during embryogenesis. Genome Res. 2014;24:920-9.

49. Ernst J, Kheradpour P, Mikkelsen TS, Shoresh N, Ward LD, Epstein CB, Zhang X, Wang L, Issner
R, Coyne M, et al. Mapping and analysis of chromatin state dynamics in nine human cell types. Nature.
2011;473:43-9.

50. Bernstein BE, Mikkelsen TS, Xie X, Kamal M, Huebert DJ, Cuff J, Fry B, Meissner A, Wernig M,
Plath K, et al. A bivalent chromatin structure marks key developmental genes in embryonic stem cells.
Cell. 2006;125:315-26.

51. Albert M, Kalebic N, Florio M, Lakshmanaperumal N, Haffner C, Brandl H, Henry I, Huttner WB.
Epigenome profiling and editing of neocortical progenitor cells during development. EMBO J.
2017;36:2642-58.

52. Albrieux M, Platel JC, Dupuis A, Villaz M, Moody WJ. Early expression of sodium channel
transcripts and sodium current by cajal-retzius cells in the preplate of the embryonic mouse neocortex.
J Neurosci. 2004;24:1719-25.

53. Mienville JM, Barker JL. Potassium Current Expression during Prenatal Corticogenesis in the Rat.
Neuroscience. 1997;81:163-172.

54. Kirmse K, Grantyn R, Kirischuk S. Developmental downregulation of low-voltage-activated
Ca2+ channels in Cajal-Retzius cells of the mouse visual cortex. Eur J Neurosci. 2005;21:3269-76.

55. Moore AR, Filipovic R, Mo Z, Rasband MN, Zecevic N, Antic SD. Electrical Excitability of Early
Neurons in the Human Cerebral Cortex during the Second Trimester of Gestation. Cerebral Cortex.
2009;19:1795-1805.

56. Selemon LD, Zecevic N. Schizophrenia: a tale of two critical periods for prefrontal cortical
development. Transl Psychiatry. 2015;5:e623.

57. Heather LC, Wang X, West JA, Griffin JL. A practical guide to metabolomic profiling as a
discovery tool for human heart disease. J Mol Cell Cardiol. 2013;55:2-11.

58. Knuesel I. Reelin-mediated signaling in neuropsychiatric and neurodegenerative diseases.
Progress in Neurobiology. 2010;91:257-274.

59. Meyer G, Perez-Garcia CG, Abraham H, Caput D. Expression of p73 and Reelin in the
Developing Human Cortex. J Neurosci. 2002;22:4973–86.

60. Miquelajauregui A, Varela-Echavarria A, Ceci ML, Garcia-Moreno F, Ricano I, Hoang K,
Frade-Perez D, Portera-Cailliau C, Tamariz E, De Carlos JA, et al. LIM-homeobox gene Lhx5 is

required for normal development of Cajal-Retzius cells. J Neurosci. 2010;30:10551-62.

61.   Shinozaki K, Miyagi T, Yoshida M, Miyata T, Ogawa M, Aizawa S, Suda Y. Absence of
Cajal-Retzius cells and subplate neurons associated with defects of tangential cell migration from
ganglionic eminence in Emx1/2double mutant cerebral cortex. Development. 2002;129:3479-92.

62.   Trapnell C, Williams BA, Pertea G, Mortazavi A, Kwan G, van Baren MJ, Salzberg SL, Wold BJ,
Pachter L. Transcript assembly and quantification by RNA-Seq reveals unannotated transcripts and
isoform switching during cell differentiation. Nat Biotechnol. 2010;28:511-5.

63.   Heinz S, Benner C, Spann N, Bertolino E, Lin YC, Laslo P, Cheng JX, Murre C, Singh H, Glass
CK. Simple combinations of lineage-determining transcription factors prime cis-regulatory elements
required for macrophage and B cell identities. Mol Cell. 2010;38:576-89.

64.   Levy R, Molina J, Danielpour M, Breunig JJ. Neonatal pial surface electroporation. J Vis Exp.
2014:1-7.

65.   Shen X, Kim W, Fujiwara Y, Simon MD, Liu Y, Mysliwiec MR, Yuan GC, Lee Y, Orkin SH.
Jumonji modulates polycomb activity and self-renewal versus differentiation of stem cells. Cell.
2009;139:1303-14.

Figure 1

A

Flow Cytometer Sorting

B

C

Ebf2-EGFP-

Ebf2-EGFP+

D

E

Figure 2

A Enriched in Ebf2+/Ebf2- cell population DEGs GO Terms

Ebf2+ cells

- cell cycle
- extracellular matrix
- forebrain development
- regulation of cell proliferation
- negative regulation of cell differentiation
- growth factor binding
- regulation of neurogenesis
- regulation of nervous system development
- pallium development
- dersal/ventral pattern formation

Ebf2- cells

- synapse
- neuron projection
- neuron differentiation
- cell morphogenesis involved in neuron
- axonogenesis
- cell junction
- calcium ion binding
- cell-cell signaling
- neuron migration
- cytoskeleton

-LOG10(P-value)

E All DEGs KEGG analysis

- Axon guidance
- Focal adhesion
- Cell cycle
- MAPK signaling pathway
- Wnt signaling pathway
- Gap junction
- DNA replication
- ECM-receptor interaction

■ -LOG10(P-value)

Ebf2+ cells:

B E11.5
- neuron projection
- synapse
- neuron differentiation
- axonogenesis
- synaptic transmission

C E13.5
- synapse
- neuron projection
- plasma membrane
- cell junction
- gated channel activity

D E15.5
- synapse
- ion channel activity
- neuron projection
- plasma membrane
- gated channel activity

F Enriched in Ebf2+ cell population DEGs KEGG analysis

- Axon guidance
- MAPK signaling pathway
- Calcium signaling pathway
- Neuroactive ligand-receptor interaction
- Gap junction
- GnRH signaling pathway
- Chemokine signaling pathway

G

1. E11.5-up E13.5-up E15.5-up (166 genes)
   Reln, Cacna2d2, Lhx5

2. E11.5-down E13.5-down E15.5-down (1983 genes)
   Tle1, Pax6, Sox2

3. E11.5-unchange E13.5-unchange E15.5-unchange (439 genes)
   Atp6v0c, Atp6v0e2, Rab11fip5

4. E11.5-up E13.5-unchange E15.5-down (161 genes)
   Syt4, Igfbpl1, Myt1

5. E11.5-down E13.5-unchange E15.5-up (64 genes)
   Zic1, Zic2, Bmp7

6. E11.5-down E13.5-unchange E15.5-down (43 genes)
   Foxg1, Bcl6, Miat

Figure 3

Figure 4

Figure 5

Figure 6