Actomyosin dynamics, Bmp and Notch signaling pathways drive apical extrusion of proepicardial cells

Laura Andrés-Delgado[1,2], Alexander Ernst[3], María Galardi-Castilla[1], David Bazaga[1]*, Marina Peralta[1,4]*,

Juliane Münch[5,6,7], Juan Manuel González-Rosa[1,8], Federico Tessadori[9], Jeroen Bakkers[9], José Luis de la

Pompa[5,6], Julien Vermot[4], Nadia Mercader[1,3 #]

1 Development of the epicardium and its role during regeneration laboratory, Centro Nacional de

Investigaciones Cardiovasculares Carlos III, Melchor Fernández Almagro 3, 28029 Madrid, Spain.

2 Departamento de Anatomía, Histología y Neurociencia, Facultad de Medicina, Universidad Autónoma de

Madrid, Spain.

3 Institute of Anatomy, University of Bern, 3000 Bern 9, Switzerland.

4 present address: Institute of Genetics and Molecular and Cellular Biology (IGBMC), 67404 Illkirch, France.

5 Intercellular Signaling in Cardiovascular Development and Disease Laboratory, Centro Nacional de

Investigaciones Cardiovasculares Carlos III (CNIC), Melchor Fernández Almagro 3, 28029 Madrid, Spain.

6 Ciber CV, 28029, Madrid, Spain.

7 present address: Institute of Biochemistry and Biology, Potsdam University, Karl-Liebknecht-Straße 24-25,

Potsdam D-14476, Germany.

8 present address: Cardiovascular Research Center, Massachusetts General Hospital and Harvard Medical

School, Boston, MA 02114, USA.

9 Hubrecht Institute-KNAW and UMC Utrecht, Uppsalalaan 8, 3584CT Utrecht, The Netherlands.

* equal contribution

corresponding author

ABSTRACT

The epicardium, the outer mesothelial layer enclosing the myocardium, plays key roles in heart development

and regeneration. During embryogenesis it arises from the proepicardium (PE), a cell cluster that appears in

the dorsal pericardium close to the venous pole of the heart. Little is known about how the PE emerges from

the pericardial mesothelium. Using the zebrafish model and a combination of genetic tools, pharmacological

agents and quantitative *in vivo* imaging we reveal that a coordinated collective movement of the dorsal

pericardium drives PE formation. We found that PE cells are apically extruded in response to actomyosin

activity. Our results reveal that the coordinated action of Notch/Bmp pathways is critically needed for apical

extrusion of PE cells. More generally, by comparison to cell extrusion for the elimination of unfit cells from

epithelia, our results describe a unique mechanism where extruded cell viability is maintained.

The epicardium is the outer mesothelial layer of the heart. During development, the epicardium sustains the underlying myocardium through paracrine signals that promote its growth[1,2]. The epicardium is also an important cell source during embryogenesis. Epicardial-derived progenitor cells (EPDCs) differentiate into cardiac fibroblasts as well as other cell types including adipocytes[3], mesenchymal stem cells, and smooth muscle or endothelial cells of the coronary vessels, and contribute to the formation of the annulus fibrosus and the valves[4,5]. After heart injury, EPDCs are involved in several aspects of tissue repair and regeneration, contributing to cardiac fibrosis, controlling the inflammatory response, promoting neoangiogenesis and cardiomyocyte proliferation[6].

The epicardium derives from the proepicardium (PE), a cluster of cells that emerges from the pericardium close to the venous pole of the heart tube around the time of heart looping and after the onset of heart beating[7,8]. In the zebrafish, PE formation is regulated by bone morphogenetic protein (Bmp) signaling. Accordingly, mutants for the Bmp receptor acvr1l do not form a PE, whereas Bmp2b overexpression extends PE marker gene expression[9]. During the process of PE formation, cells undergo a change in polarity, suggesting that an epithelial-mesenchymal-like transition has a role in cluster generation[10-13].

After the PE is formed, the heartbeat has an essential role in allowing PE cells to be "washed away" into the pericardial cavity. Heartbeat generates a pericardial fluid flow allowing the PE cells to reach the myocardial surface, to which they ultimately adhere and begin epicardial layer formation[14,15]. Nevertheless, while it is clear that pericardial flow triggers the release of PE cells to the pericardial cavity and is needed to form the epicardial layer, less is known about the role of mechanical forces on PE formation.

During morphogenesis, cell migration and proliferation results in the continuous rearrangement of mechanical properties of tissue layers. Collective cell migration and proliferation can lead to local cell crowding and the generation of tissue tension[16]. Conversely, changes in tissue growth can further influence cell signaling[17,18]. The actomyosin cytoskeleton plays a central role in controlling cell shape and developmental events[19-22]. It is tightly associated with membrane junction complexes and can react to extracellular signals or signals from neighboring cells by altering cell properties[22-24]. From looping, trabeculation to valvulogenesis, mechanical activity of the heart has a fundamental function in controlling cardiac morphogenesis[25-28]. Here we used the zebrafish model to study the morphogenetic events leading to PE formation. We found that cells from the dorsal pericardium (DP) collectively move towards the DP midline,

where some of them round up and extrude into the pericardial cavity. This movement of DP cells to the

midline and PE cell extrusion is dependent on actomyosin dynamics. We found that Bmp2 signaling lies

upstream of the actomyosin cytoskeletal rearrangements necessary for PE formation. Furthermore, we show

that the developing heart tube influences PE formation as endocardial Notch signaling controls Bmp

expression.

RESULTS

Constriction of the dorsal pericardium and apical extrusion leads to PE delamination. To investigate the

formation of the PE, we performed a detailed analysis of pericardial mesothelial cell movement in zebrafish

embryos. PE clusters appeared at the midline of the DP, which extends from the venous to the arterial pole of

the heart tube (Fig. 1a). To image PE formation, which begins around 52 hours post fertilization (hpf), we used

the enhancer trap line *Et(-26.5Hsa.WT1-gata2:EGFP)^{cn1}* (hereafter termed epi:GFP) in which GFP expression

is controlled by the regulatory elements of *wilms tumor 1a* (*wt1a*), and recapitulates its expression pattern[14].

Thus, pericardial and PE cells are GFP$^+$ in the epi:GFP line. GFP expression is present in all DP cells and is

particularly strong around the cell nucleus[29] and thus suitable for cell tracking. We tracked individual cells of

the DP using time-lapse video imaging from 52 to 60 hpf, and found that they became displaced and

converged at the midline (Fig. 1b,c and Supplementary Movie 1). We also measured the angle of the cell

trajectories of the constricting DP tissue in relation to the midline and found that the majority were close to 90º,

indicating that DP cells move nearly perpendicular to the midline and suggesting that this movement leads to

an active directional accumulation of cells at the midline producing DP tissue constriction (Fig. 1d; n=3

embryos).

To further characterize the morphological changes to the DP during PE formation, we quantified the changes

in the relative velocity of DP cells with respect to their neighboring cells using a customized calculator of the

velocity field divergence based on DP cell tracking in epi:GFP embryos (Supplementary Fig. 1). The

divergence field represented predominantly by purple-blue colors indicates an overall constriction in the tissue,

whereas tissue expansion is reflected by red-orange pseudostaining within the divergence field. By calculating

this factor for the imaged DP at 43 different time points every 12 min, and measuring 150 DP cell tracks in

three embryos, we confirmed that there was an overall constriction of the DP from 52 to 60 hpf, with highest

local levels of constriction at the site of PE formation (Fig. 1e and Supplementary Movie 2).

We next characterized the emergence of PE cells, again by *in vivo* imaging of the epi:GFP line. During

the displacement of DP cells towards the midline, cells close to the midline began to show a stronger GFP

signal and rounded up (Fig. 1f and Supplementary Movie 3, which is representative for the event observed in a

total of 10 different movies). We further observed that the cells surrounding these emerging PE cells came

closer together. Ultimately, one or multiple cells protruded from the DP layer and remained only loosely

attached to the neighboring DP mesothelial cells. The cells that were bordering the newly formed PE cells

subsequently converged under the rounded PE cell. These cell behaviors are landmarks of apical extrusion,

allowing cells to bulge out and leave organized epithelia[30]. Therefore, the results suggest that PE formation

occurs through the local overcrowding of cells at the DP midline, inducing apical extrusion of mesothelial

cells of the DP.

Proliferation of pericardial cells contributes to constriction at the midline. To determine whether the

overall constriction of the DP was in part due to cell proliferation, we analyzed the spatial distribution of cell

division within the DP. By measuring the distance of the cell division plane to a defined midline (a surface of

about 20 μm diameter drawn from the arterial to the venous pole of the heart tube) we found that pericardial

cells divided all over the DP (Fig. 2a,b and Supplementary Movie 4). However, cell division occurred neither

directly at the midline nor preferentially close to it (inside the surface = 0 μm 0 divisions, 0–20 μm 7 divisions,

20–80 μm 11 divisions, Fig. 2c). Categorizing orientation of cell divisions showed that most divisions were

perpendicular to the midline (Fig. 2d). We next assessed cell proliferation by immunostaining for phospho-

histone 3 (pH3) on fixed epi:GFP embryos. While proliferating cells could be found in the pericardium (Fig.

2e-g and Supplementary Fig. 2a,b), only one pH3$^+$ PE cell was observed in 4 out of 12 embryos (Fig. 2g,h

Supplementary Fig. 2c,d). Inhibiting cell proliferation from 48 hpf onwards with the pharmacological agents

nocodazole (noc) or aphidilcolin/hydroxyurea (aph/hydrU) significantly reduced the number of pH3$^+$ cells in

the pericardium from 52 ± 29 in controls to 21 ± 8 in noc and 0 ± 1 in aph/hydrU-treated embryos at 60 hpf;

(Fig. 2e,f). Noc inhibits cell proliferation by halting mitosis[31] and aph/hydrU inhibits DNA synthesis[32]. Both

treatments significantly reduced the number of PE cells from 9 ± 3 to 0 ± 1 or 2 ± 1, respectively (Fig. 2i).

However, we observed that in noc-treated embryos, DP cells still converged at the midline (Supplementary

Movie 5). Thus, cell proliferation might contribute partially to the local crowding of DP cells at the midline,

which ultimately leads to PE cluster formation (Fig. 2j).

PE formation depends on actomyosin dynamics. Cytoskeleton reorganization is a fundamental process for

the apical extrusion of apoptotic epithelial cells and for cell migration/invasion of metastatic cells[33-35]. Prior to

extrusion, cells present an overall increase in F-actin levels and a change in actomyosin localization, from

basal to apico-cortical deposition[36]. Because myosin II plays a fundamental role in actin filament motion[37-39],

we analyzed its expression in PE cells. Immunofluorescence analysis revealed that myosin II-A was highly

expressed in PE cells (Fig. 3a'). In DP cells close to the midline, myosin II-A expression was strong at the cell

boundaries (Fig. 3a''). In rounded PE cells, myosin II-A was also expressed apically and at boundaries

between PE cell pairs (Fig. 3a''', 3a''''). *In vivo* imaging of the *actb2:myl12.1-mCherry*[e1954] line, expressing

myosin II fused to mCherry under the embryonic ubiquitous promoter *actin beta 2* (*actb2*), confirmed an

accumulation of myosin II in DP cells at the midline and in the PE cluster (Supplementary Movies 6,7).

We also analyzed the localization of polymerized actin during PE cell formation at 60 hpf using

whole-mount labeling of actin with a phalloidin-coupled fluorophore (Fig. 3b'). Whereas actin was located in

the basal region of the DP cells (Fig. 3b'', 3b'''), it was polarized and accumulated apico-cortically in PE

cells, consistent with the pattern observed for myosin II (Fig. 3b''',3b''''). In vivo characterization of

actomyosin dynamics was performed in the double transgenic line *βactin:LifeAct-RFP*[e2212Tg];epi:GFP. We

observed that from 52 to 60 hpf, F-actin concentrated in DP cells at the midline where the PE appears (Fig. 3c).

Indeed, a thick actin cable was visible in epi:GFP+ cells at the time of PE formation, spanning the midline of

the DP (Supplementary Movies 8–11). Actin was concentrated basally in the DP cells that surrounded the PE

cells, whereas in PE cells actin changed its polarity and and became localized cortically and concentrated in

the contact region between the rounded up PE cells (Supplementary Movie 12). In PE cells that were close to

being released into the cavity, we observed that in a final stage prior to release a thin actin rich stalk appeared

in the PE cell still attached to the DP (Supplementary Movie 13). In sum, our data suggest that PE formation

occurs concomitantly with extensive actin reorganization: before PE formation, F-actin is located basally in

DP cells, then it becomes concentrated in PE precursor cells and it finally accumulates all around the cortex of

the rounded PE cells. We previously found that PE formation is impaired in the presence of butanedione

monoxime (BDM)[14], which interferes with myosin-ADP-Pi phosphate release, and locks myosin II into a low

affinity conformation with actin impeding myosin movement on top of actin filaments[40]. Inhibition of myosin

II with blebbistatin (BLEB), through maintenance of myosin II in an actin-detached state, also impaired PE

formation and this was dose dependent (Supplementary Fig. 3a-c). Compared to non-treated embryos, BDM

treatment led to a reduction in LifeAct-RFP expression in the DP (Fig. 3d,e), showing that F-actin is unstable

in DP cells at the midline. Results from *in vivo* imaging revealed that the arrangement of epi:GFP$^+$ cells at the

midline was inhibited by BDM and the movement of the DP was impeded (Supplementary Movies 14,15).

We reasoned that if actin polymerization was required for PE formation, pharmacological

enhancement of actin stability should counteract the effect of myosin II inhibition. To test this, we

administered jasplakinolide (jasp), which promotes actin filament polymerization and stabilization[41], to

epi:GFP animals in the presence or absence of BDM. Embryos treated with jasp and BDM from 48 hpf

onwards showed stronger actin signals than embryos treated only with BDM, suggesting that jasp rescued

actin polymerization in BDM-treated animals (Fig. 3f). Moreover, animals treated with jasp or BDM + jasp

had 8 ± 4 and 9 ± 3 PE cells, respectively (n=15 or n=14 embryos each), which resembles control conditions,

whereas BDM-treated animals had 1 ± 1 PE cells (n=13) (Fig. 3g,h). Thus, enhancement of actin filament

polymerization and stabilization in the presence of BDM correlated with PE formation, suggesting that F-actin

is necessary for PE formation.

Bmp2b influences actomyosin dynamics during PE formation. During heart tube development, myocardial

cells are involved in the secretion of signaling molecules such as Bmps[42-44], and Bmp signaling is necessary

for correct PE cluster formation in zebrafish[9]. To evaluate whether ectopic Bmp could rescue the impairment

in PE formation upon inhibition of actomyosin dynamics, we next performed experiments using the transgenic

line *hsp70:bmp2b*[45] crossed into epi:GFP, which allows *bmp2b* levels to be increased at a specific

developmental stage by heat shocking the embryos at 39°C for one hour. In control (non-transgenic for

hsp70:bmp2b) animals subjected to heat shock (HS), the PE was completely formed and visible at 60 hpf and

comprised approximately 10 cells (Fig. 4a,b). Overexpression of *bmp2b* during the embryonic stages

preceding PE formation by delivering HS pulses at 26, 32 and 48 hpf, resulted in a 2-fold increase in the

number of PE cells; from 10 ± 5 cells in control embryos to 22 ± 9 in *bmp2b*-overexpressing embryos at 60

hpf ($P < 0.0001$) (Fig. 4a,b). The total number of DP cells prior to PE formation did not differ between

bmp2b-overexpressing fish and control animals heat shocked for 26 and 32 hpf (Supplementary Fig. 3d; n=6/7

animals), suggesting that the enlarged PE clusters observed upon *bmp2b* overexpression were not a

consequence of an expanded population of DP cell precursors. PE cells emerge to a small extent from the DP

close to the venous pole of the heart (vpPE) and to a larger extent from the DP close to the atrioventricular

canal (avcPE)[14]. To assess whether Bmp2b was acting on a particular subpopulation we individually

quantified the number of cells in the avcPE and vpPE clusters and found that *bmp2b* overexpression

significantly increased the number of cells in the avcPE cluster (Supplementary Fig. 3e).

In addition, the number of epicardial cells found on the ventricular myocardial surface was higher in

bmp2b-overexpressing fish at 60 hpf (Fig. 4a,c). In line with this, antagonizing Bmp signaling by

overexpressing *noggin3* in the *hsp70l:noggin3[fr1]4* line [45] through HS at 48 hpf significantly reduced the

number of GFP cells within the PE cluster (Fig. 4a,b).

We next aimed to assess whether Bmp2b activated Bmp signaling directly in PE cells. To do this, we

evaluated the expression of its downstream effector pSmad1/5. pSmad1/5 immunostaining was found both in

DP cells and in some PE cells between 55 and 60 hpf (Fig. 4d-f). Approximately 20% of DP cells were

pSmad1/5$^+$ (a mean of 4 ± 3 positive cells from 15 ± 7 total DP cells; n=21 embryos), whereas only about 2%

of PE cells were pSmad1/5$^+$ (0 ± 1 positive cells from a total of 9 ± 4 cells; n=16 embryos). The number of

pSmad1/5$^+$ DP cells was significantly higher than the number of pSmad1/5$^+$ PE cells (P < 0.001). Moreover,

the total number of pSmad1/5+ DP cells was increased in *bmp2b*-overexpressing embryos (Fig. 4f).We futher

used a Bmp reporter line expressing Kusabira Orange (KuO) under the control of a promoter harboring several

smad binding sites, named Bmp responsive elements (BRE)[46]. Results from *in vivo* imaging of *BRE*:KuO;

epi:GFP fish revealed that DP cells and some PE cells were KuO$^+$ (n=1–2 KuO$^+$ PE cells per animal, seen in 3

animals from 52 to 58 hpf), confirming that the Bmp signaling pathway was transiently active in PE cells (Fig.

4g and Supplementary Movie 16). Therefore, an increase in Bmp activity within the DP correlates with PE

formation.

We addressed whether overexpression of *bmp2b* could rescue the impaired PE cluster formation with

BDM from 48 hpf onwards. PE formation was observed in *bmp2b*-overexpressing animals treated with BDM,

and clusters were larger than those observed in controls (23 ± 3 vs 10 ± 5 cells; Fig. 4a,b $P < 0.0001$; n=29

embryos). Overexpression of *bmp2b* also rescued PE formation in BLEB-treated animals (Supplementary Fig.

3a,b). We observed that the rescue of PE formation was dependent on *bmp2b* expression levels. Accordingly,

larger PE clusters were detected after 3 HS pulses (at 26, 32 and 48 hpf) (22 ± 9 cells; n= 25 embryos)

(Supplementary Fig. 3b) than with only one HS pulse at 48 hpf (10 ± 4 cells; n=30 embryos) (Supplementary

Fig. 3d). We also assessed at which developmental stage the effect of bmp2b was more prominent on PE

formation. A unique HS at 48 hpf rescued PE formation at 60 hpf in BDM-treated hearts (8 ± 4 cells vs 3 ± 2,

n=20 embryos) (Supplementary Fig. 3f), but a single HS at 26 hpf failed to rescue PE formation at 60 hpf in

BDM-treated embryos (6 ± 3 cells vs 4 ± 2, n=12 embryos) (Supplementary Fig. 3g). *bmp2b* overexpression

after BDM treatment also failed to rescue PE formation (Supplementary Fig. 3h, n=10).

BDM treatment impairs pericardial cell proliferation[14]. As we observed that DP proliferation events

preceded PE formation, we questioned whether *bmp2b* overexpression in BDM-treated embryos might

increase DP cell proliferation. However, the number of proliferating epi:GFP[+] pericardial cells was very low

and not different between BDM-treated animals overexpressing or not *bmp2b* (Supplementary Fig. 2b,d, n=23).

We therefore explored how the actomyosin network is altered by *bmp2b* overexpression to understand the

mechanisms through which Bmp2b might neutralize the effect of BDM on the actomyosin cytoskeleton. BDM

treatment strongly reduced the amount of myosin II-A in DP cells (n=8 animals) and overexpression of *bmp2b*

in BDM-treated animals rescued the apical polarization of myosin II-A in PE cells (n=11) to levels similar to

those observed in control embryos (n=10) (Fig. 5a), suggesting a recovery of actomyosin dynamics by the

Bmp pathway. Thus, we investigated how actin polymerization is affected upon Bmp2b overexpression.

Examination of F-actin levels by immunostaining at 60 hpf revealed that *bmp2b* overexpression increased

these levels significantly. In the presence of BDM, actin levels in PE cells were lower than those in controls;

however, they were significantly rescued upon *bmp2b* overexpression (Fig. 5b,c).

Our results show that actin filaments are necessary for PE formation. In agreement with this, treatment

with the actin polymerization and elongation inhibitor, cytochalasin D (cytD) at 2 μM from 48 hpf onwards,

decreased the size of PE clusters from 9 ± 3 cells (in non-treated fish) to 4 ± 3 cells (n=10–14 embryos in each

group) at 60 hpf. Of note, *bmp2b* overexpression was unable to rescue PE formation in the presence of cytD

(Fig. 5d,e), indicating that the inhibitory effect of cytD on F-actin assembly is downstream of the effect of

Bmp2b on the actin cytoskeleton in PE cells.

We next evaluated whether the effect of Bmp inhibition on PE formation could be rescued by

increasing actin stabilization. Thus, the number of cells in PE clusters of animals treated with the Bmp

receptor-I inhibitor LDN-193189 (LDN)[47] was compared with those of animals treated with a combination of

LDN and jasp. LDN treatment reduced the number of PE cells per cluster compared with untreated controls

and reverted the increase in PE size upon *bmp2b* overexpression (Fig. 5f,g). The mean number of PE cells was

2 ± 2 cells (from a total of 24 animals) for the LDN treated group and 3 ± 2 (from a total of 9 animals) for the

LDN + jasp group (Fig. 5g). In both cases, the mean number of PE cells was lower than that usually observed

in untreated controls, which was 9 ± 2 cells. By contrast, when we stabilized actin filaments with jasp from 48

hpf onwards and 4 h prior to LDN administration, the number of PE cells was significantly increased to 6 ± 3

(from a total of 11 animals) compared with the LDN group. Taken together, these results show that the Bmp

signaling pathway influences actomyosin polymerization and the stabilization of the F-actin network partially

compensates for the negative effect of Bmp signaling inhibition on PE formation.

To gain deeper insight into the mechanisms of Bmp2b action, we analyzed how DP cell displacement

is altered in BDM-treated animals in a background of *bmp2b* overexpression. We imaged epi:GFP animals

from 52 to 60 hpf and tracked epi:GFP[+] cells in the DP. In *bmp2b*-overexpressing animals, the DP constricts to

the midline (Fig. 6a), as observed in controls (Fig. 1c). Upon BDM treatment, the typically observed crowding

of GFP[+] cells at the midline was not apparent (Fig. 6b). However, *bmp2b* overexpression rescued DP cell

displacement towards the midline upon BDM treatment (Fig. 6c). We next quantified DP cell displacement

using divergence fields. BDM treatment led to expansion, whereas control and *bmp2b* overexpression led to

DP tissue constriction (Fig. 6d,e). Accordingly, cell displacement tracking revealed a movement towards the

midline in control (n=4 embryos) and *bmp2b*-overexpressing embryos (n=4), whereas DP cells were

predominantly displaced away from the midline in BDM-treated embryos (n=5) (Fig. 6f). Again, *bmp2b*

overexpression in BDM-treated animals increased the direction of displacement and favored the movement

towards midline (n=3).   Altogether, the results suggest that bmp signaling acts through actomyosin

cytoskeleton to allow the displacement of DP cells towards the midline, which ultimately leads to PE

formation.

Endocardial/endothelial Notch signaling acts upstream of Bmp2b to control PE formation. Notch and

Bmp signaling pathways are connected during the development of several organs including the heart[48-52].

Moreover, loss of function of Notch signaling alters epicardium formation and Notch signaling regulates

smooth muscle differentiation of epicardium-derived cells[53]. To study the link between Notch and Bmp2

signaling pathways during PE formation in the zebrafish, we used the transgenic line *UAS:NICD-myc*[KCA4 54] to

overexpress the intracellular active domain of the Notch receptor (NICD) under a HS-inducible promoter

hsp70:Gal4[KCA3]. When we performed HS in this line at 48 hpf, PE formation was unaltered at 60 hpf when

compared with non-transgenic zebrafish (8 ± 4 cells *vs* 8 ± 3 cells, respectively) (Fig. 7a,b). However, the

overexpression of *NICD* led to the maintenance of a PE as late as 80 hpf, a time point at which a PE cluster is

typically no longer visible (Supplementary Fig. 4a,b) (6 ± 3 cells *vs* 0 ± 1 cells in controls; $P < 0.0001$).

To determine the cell type in which the activity of NICD is necessary, we crossed *UAS:NICD-myc*[KCA4]

transgenic fish with *wt1b:Gal4* animals to drive NICD expression in DP and PE cells, or with *fli1a:gal4*[ubs3Tg]

[55], to overexpress NICD in endothelial and endocardial cells (Fig. 7a,b). Whereas activation of the Notch

pathway in *wt1b*[+] cells did not affect PE formation, its activation in *fli1a*[+] cells resulted in the significant

increase in PE cell numbers, which was already apparent at 60 hpf, as well as a maintenance of the PE cluster

at 80 hpf (Supplementary Fig. 4a,b) (14 ± 8 cells *vs* 0 ± 1 cells in controls; $P < 0.0001$). Therefore, Notch-over

activation in endothelial/endocardial cells affects PE cluster formation. Next, we tested whether *NICD*

overexpression was also able to rescue PE formation under BDM treatment, as observed for *bmp2b*. At 60 hpf,

PE formation was not rescued by *NICD* overexpression (Fig. 7c,d) but at 80 hpf a PE cluster was observed in

BDM-treated animals (13 ± 8 cells *vs* 0 ± 1 cells in BDM only; $P < 0.0001$;Supplementary Fig. 4c,d). *NICD*

overexpression did not increase cell proliferation in the DP, as assessed by quantification of pH3[+] cells and it

also did not rescue the number of proliferating cells under BDM treatment (Fig. 7e). The effect of Notch1

inhibition for PE formation was also assessed with the Notch inhibitor, RO-4929097 (RO)[56]. RO

administration reduced PE cluster size (2 ± 2 cells *vs* 8 ± 3 cells in controls), which occurred concomitant with

the loss of actin cytoskeleton polarization in DP cells at the midline (Fig. 7 b,f). These findings suggest that

paracrine signals from the underlying endothelium or endocardium guide PE formation. Indeed, we observed

that at the site of cluster formation at the DP, the PE is situated on top of *kdrl*[+] endothelial sinus venosus horns

(Fig. 7g and Supplementary Movie 17). Moreover these endothelial cells show GFP expression in

kdrl:mCherry transgenic fish crossed to *ET33-mi60a* animals[57]. This enhancer trap line, herein called *lfng:*GFP,

expresses GFP under the control of the *lunatic fringe* (*lfng*) gene regulatory region, a modulator for Notch

signaling[58,59] (Fig. 7h). *lfng:*GFP marks endothelial cells from the sinus venosus and the endocardium (Fig.

7f,g). PE cells do not present high levels of *lnfg:*GFP expression (Fig. 7h,i). As a further proof-of-concept that

Notch activity is present in endocardial cells, we lineage traced Notch responsive cells using *Tp1:CreERT2*[60] .

In this line, *CreERT2* is driven by RBPƙ binding sites. Crossing Tg(*Tp1:CreERT2)* into Tg(*ubb:LOXP-EGFP-*

LOXP-mCherry) and treating embryos with 4-hydroxytamoxifen (4-OHT) at 48 hpf allows for the permanent

labeling of cells that receive Notch signaling at the time of 4-OHT administration by mCherry expression. At

80 hpf, GFP[+] cells were detected only in the endocardial cells at the atrioventricular canal, but not in the

epicardium (Fig. 7j). Fluorescent *in situ* mRNA hybridization to detect *notch1a* revealed that its coexpression

colocalizes with *fli1a*:GFP[61], an endocardial reporter line (Fig. 7k). Furthermore, the Notch pathway target

Hey2 is expression in the endocardium of the embryonic heart tube (Fig. 7l). Altogether our data indicate that

endocardial/ endothelial Notch pathway activation is required for PE formation. The fact that the rescue of PE

formation under BDM treatment by *NICD* overexpression occurred with a delay in comparison to that seen

with Bmp2b activity suggests that the Notch pathway might act upstream of *bmp2b* in the control of PE

formation.

To further study a possible effect of Notch on Bmp signaling during PE formation we examined the

presence of PE cells and pSmad1/5[+] DP cells upon endothelial/endocardial Notch overactivation in

fli1a:gal4;UAS:NICD animals. At 60 hpf we observed that animals with ectopically-activated Notch signaling

in *fli1a*[+], but not in *wt1b*[+] cells, revealed pSmad1/5[+]DP and PE cells, suggesting an upregulation of Bmp

signaling in pericardial cells in response to endothelial Notch activity (Fig. 8a-c). At 80 hpf, this effect is

exacerbated (Supplementary Fig. 4e-g).

We also analyzed the impact of the increase in PE cells upon *NICD* and *bmp2b* overexpression on

epicardium formation. Whereas epicardial cell numbers were unaffected by the overexpression of *NICD*,

bmp2b overexpression increased the epicardial cell number at 80 hpf as compared with untreated animals

(Supplementary Fig. 4h). Similarly, we wanted to assess whether PE formation was stable over longer time

periods: at 5 days postfertilization (dpf) the PE was not present (Supplementary Fig. 4i). To further address the

relationship between Notch and Bmp signaling pathways, we combined various Notch and Bmp gain- and

loss-of-function scenarios. First, we added LDN from 48 hpf to heat-shocked $hsp70:Gal4^{KCA3};UAS:NICD$-

myc^{KCA4} embryos (Fig. 8d). At 60 hpf, PE cell formation was not rescued by $NICD$ overexpression in presence

of LDN (2 ± 3 cells vs 9 ± 2 cells in untreated animals) (Fig. 8e). The reduction in PE formation upon RO

treatment from 48 hpf onwards was reverted by the overexpression of $bmp2b$ where a cluster similar to that

seen in a wildtype condition was present (2 ± 2 cells vs 8 ± 5 cells, $P < 0.0001$) (Fig. 8d,e).

Next, we evaluated how the different treatments affected pSmad1/5 levels in the DP (Fig. 8f). As

expected, $bmp2b$ overexpression increased the number of pSmad1/5$^+$ cells and LDN treatment reduced this.

RO administration also led to a reduction in the number of pSmad1/5$^+$ cells and this was rescued by $bmp2b$

overexpression (n=10), but HS-induced $NICD$ overexpression could not rescue the number of pSmad1/5$^+$ cells

upon LDN treatment. Given that pSmad1/5 levels increased upon Notch overactivation, we analyzed whether

bmp expression levels were altered in the heart tube. At 60 hpf, RO treatment reduced $bmp4$ expression levels

(Fig. 8g, n=22/26 for control and 12/20 for RO-treated animals). Under endothelial $NICD$ overexpression,

$bmp4$ expression is increased in the heart tube (n=9/13). At 80 hpf, we also detected increased $bmp4$ and

additionally $bmp2b$ expression under endocardial $NICD$ overexpression, and reduced levels in RO-treated

animals (Supplementary Fig. 4j, n=20 embryos for each condition). Fluorescent in $situ$ hybridization against

$bmp4$ was performed to analyze in which cell type this gene is expressed. We found colocalization of $bmp4$

expression with the myocardial marker Myosin Heavy Chain in control and endocardial $NICD$ overexpression

(Fig. 8h).

One possible explanation for the increase of PE clusters observed upon $bmp2b$ or $NICD$

overexpression is that less cells are released from the PE and hence that PE clusters change in size in the

different tested conditions. Thus, we counted the number of PE cell release events during 8 h in $vivo$ and no

differences were observed between control, $bmp2b$ or $NICD$ overexpression (Fig. 8i). Similarly, the number of

PE cells released per event did not differ between groups (Fig. 8j).

Overall, these experiments indicate that the Notch signaling pathway acts in the endothelium and

endocardium to regulate Bmp signaling in the myocardium, which is necessary for PE formation (Fig. 8k).

DISCUSSION

We propose a model in which Notch activity in endocardial cells leads to the expression of *bmp2b* and *bmp4* in the embryonic heart tube myocardium, which subsequently signals to PE precursor cells promoting PE cluster formation through its effect on the actomyosin cytoskeleton (Fig. 8k). These findings illustrate the importance of an intact actomyosin scaffold for generating the interactions and forces between DP cells necessary for the apical extrusion of PE cells, the source of the epicardial layer. Thus, PE formation is a highly dynamic morphogenetic event in which the combined action of mechanical events and signaling pathways converge.

At the midline of the DP, individual DP cells delaminate from the pericardial layer towards the cavity, and therefore PE cluster formation is based on apical extrusion. Apical extrusion is a cellular mechanism that has been reported to control epithelial layer homeostasis in the intestine while preserving its barrier function[62-64]. Live cell extrusion has also been reported to occur in epithelia during embryogenesis to control cell number[30]. Accordingly, an increase in cellular density leads to the elimination of supernumerary cells towards the apical side[30,65]. Extruded cells then undergo apoptosis upon loss of cell contact with their neighbors[66,67]. Neighboring wild-type cells also play an active role in this process by accumulating cytoskeleton as intermediate filaments at the interface with transformed cells[68]. Instances where extruded cells survive, as observed here during epicardium formation, have been documented for transformed cells: specifically in epithelial cancer models where apically-extruded transformed cells overexpress the oncogenes *Src* or *Ras*, which prevent cell death[69-72].

Here we describe a physiological mechanism of apical cell extrusion that results in cell survival and is part of a natural process occurring during heart development that is required for the generation of epicardial precursor cells. To our knowledge, this is the first description of apical extrusion occurring during embryogenesis that leads to living extruded cells, but it may represent only one example of a series of morphogenetic events in which this mechanism is involved. An additional process where apical extrusion is likely to play a role is during the emergence of hematopoietic stem cells from the dorsal aorta[73,74]. The gene regulatory mechanism involved in promoting the survival of extruded PE cells, until they attach to the

myocardium, remains to be elucidated. It is plausible, that rapid attachment of extruded PE cells to the

myocardial surface promotes their cell survival.

Our results reveal a previously unidentified role for pericardial cell movements in PE formation. We

found that these motions are dependent on the actomyosin cytoskeleton and lead to an overall constriction of

the DP tissue at the midline. Interestingly, actomyosin-dependent cell rearrangements at the DP influence

outflow track development in the mouse [75].

Cell proliferation contributed partially to cell crowding at the midline and influenced PE formation.

However, ectopic *bmp2b* overexpression could promote PE formation in the absence of cell proliferation,

which raises the question of the extent to which cell proliferation is a limiting step for DP cell movement

towards the midline. During gastrulation, cell rearrangements are not always regulated by proliferation as they

can be controlled by global adherent junction-remodeling arising from myosin-II-mediated local contractile

forces between cells[76]. This is reminiscent of the movement of DP cells, which we observed during PE cluster

generation that also is dependent on myosin II activity.

The interplay between cell signaling and mechanics during development is a major focus of research[77].

An important question to be answered is how Bmp2 signaling and actomyosin dynamics are linked to control

PE formation. Enhanced Bmp2b signaling could overcome the negative effect of BDM on PE formation by

blocking myosin II activity. Our results suggest that Bmp2b counteracts the tissue compliance caused by

inhibition of the actomyosin network. They could also indicate that Bmp2b stabilizes F-actin: since actin

intensity levels were higher upon Bmp2 overexpression, a higher dose of BDM would be required to disrupt

the actomyosin network. Another possibility is that Bmp2 rescues the F-actin tension by modulating the

expression of other non-conventional myosins that are not inhibited by BDM. Indeed, Bmp signaling has been

recently shown to regulate epithelial morphogenesis by controlling F-actin rearrangements, apical constriction,

cell elongation and epithelial bending[78]. Along this line, interkinetic nuclear migration during morphogenesis

of the retina is driven by actomyosin forces that are blocked when myosin II is inhibited, and the reduction in

the velocity of migration and cytoskeleton dynamics by BDM can be rescued by BDM-insensitive myosins[79].

Furthermore Bmp2 has been shown to control the expression of the *non-muscle myosin Va* gene, thereby

promoting cellular migration[80]. Bmp2 signaling has been recently suggested to regulate *Prrx1* expression in

the lateral plate mesoderm, which in turn regulates the expression of *palladin*, an actin bundling protein[27]. This

signaling cascade might be important for PE formation.

Our results suggest a direct effect of Bmp2b signaling on actomyosin dynamics, but it is also

conceivable that Bmp2b acts on DP cells by changing their cell-cell adhesion or cell-basal membrane

interactions, which would allow them to overcome the absence of a functional actomyosin network. E-

cadherin and tight junction-associated proteins are involved in polarization and apical extrusion of transformed

epithelial cells[81]. Integrin-paxillin signaling also plays an important role in PE formation[12], and Bmp signaling

has been previously shown to affect integrin signaling. Bmp2 can induce spreading of a myoblast cell line by

activating integrins, which increases cell adhesion dynamics and leads to reorganization of the cytoskeleton[82].

Moreover, integrins *per se* are Bmp targets[83]. Thus, a possible scenario can be envisaged in which ectopic

bmp2 promotes PE formation by promoting integrin-basement membrane interactions. Studies determining

how the Bmp pathway affects the actomyosin cytoskeleton and whether it relates to the redistribution in

junction components such as E-cadherin or integrins during PE formation will help to reveal the process of PE

cell detachment from the DP.

Bmp signaling has been described to be necessary for PE specification in the zebrafish[9]. In chicken PE

explant assays, a balance between Bmp and Fgf signaling determines the specification of precardiac mesoderm

into either PE or cardiomyocytes[84,85]. Furthermore, Bmp is one of the factors needed to drive differentiation of

human pluripotent stem cells into an epicardial lineage[86,87]. A relationship between Notch and Bmp2 signaling

during cardiovascular development has already been reported in the mouse[51,88]. *N1ICD* gain of function

experiments showed that myocardial Notch1 overexpression negatively regulates Bmp2[52]. Consistently, Notch

signaling abrogation leads to increased Bmp2 signaling in the PE[53]. Thus, whereas the Notch/Bmp2 axis

promotes PE formation in the zebrafish, it represses it in the mouse. In mammals, the PE is formed by an outer

mesothelial layer and an inner mesenchymal core[89], whereas in the zebrafish, PE cells are derived from the

pericardial mesothelium[14]. It is possible that this distinct PE architecture contributes to a different effect of

Notch and Bmp2 signaling.

In conclusion, our findings illustrate the importance of an intact actomyosin scaffold for generating the

interactions and forces between DP cells necessary for the apical extrusion of PE cells, the source of the

epicardial layer. Collectively, our results reveal an exquisite orchestration between heart tube maturation and

PE formation and may represent a paradigm for the coordinated action of signaling molecules and mechanical

forces in controlling tissue morphogenesis.

MATERIALS AND METHODS

Zebrafish strains and husbandry

All experiments were approved by the Community of Madrid "Dirección General de Medio Ambiente" in

Spain; and the "Amt für Landwirtschaft und Natur" from the Canton of Bern, Switzerland. Animals were

housed and experiments performed in accordance with Spanish and Swiss bioethical regulations for the use of

laboratory animals. Fish were maintained at a water temperature of 28 ºC. The following fish were used: wild-

type AB strain; Et(-26.5Hsa.WT1-gata2:EGFP)[cn1] (epi:GFP)[14]; *Tg(myl7:mRFP)[90]*, *Tg(hsp70l:bmp2b)*[fr13] and

Tg(*hsp70l::noggin3*)[fr14] [45], Tg(*kdrl::mCherry*)[ci5] (from Elke Ober´s laboratory), Tg(*uas::myc-Notch1a-*

intra)[kca3], Tg(*hsp70l::Gal4*)[kca4] [91]; Tg(*fli1a:gal4*)[ubs3Tg] [55], Tg(*βactin:LifeActin:RFP*)[e2212Tg] [92], Tg(*actb2:myl12.1-*

mCherry)[e1954] [93], Tg(*BRE-AAVmlp:dmKO2*)[mw40] [46], Et(*krt4:EGFP*)[sqet33mi60A] [57], Tg(*Tp1:CreERT2*)[s951] [60], Tg(-

3.5ubb:LOXP-EGFP-LOXP-mCherry) [cz1702] [94] Tg(*fli1a:eGFP*)[61] and Tg(*UAS:mRFP*)[95].

The Et(-*26.5Hsa.WT1-gata2:EGFP*)[cn1] line contains a reporter construct flanked by FRT sites, in which

cardiac actin drives the expression of RFP. The cassette was removed by injection of flipase into one-cell stage

embryos. We named this new line Et(-*26.5Hsa.WT1-gata2:EGFP*)[cn14].

For experiments to overexpress NICD in the endocardium/endothelium the following triple transgenic line was

used: Tg(*fli1a:gal4*);(*hsp70:gal4*);(*UAS:NICD*) and the control was Tg(*hsp70:gal4*);(*UAS:NICD*), without

the HS step.

Heat shock

Heat shock was performed to the embryos at 39ºC in preheated water for 1 h.

Generation of the *TgBAC(wt1b:GAL4)* transgenic line

The translational start codon of *wt1b* in the BAC clone CH73-186G17 was replaced with the *galff-polyA-Kan[R]*

cassette by Red/ET recombineering technology (GeneBridges) as described[96]. To generate the targeting PCR

product, *wt1b*-specific primers were designed to contain 50 nucleotide homology arms around the ATG with

~20 nucleotide ends to amplify the *galff-polyA-Kan[R]* cassette To facilitate transgenesis, the BAC-derived *loxP*

site was replaced with the *iTol2-Amp[R]* cassette[97] using the same technology. The final BAC was purified with

the HiPure Midiprep kit (Invitrogen) and co-injected with Tol2 mRNA into *Tg(UAS:GFP)* embryos[95]. The full

name of this transgenic line is *TgBAC(wt1b:GALFF)*.

Primers used to generate the *wt1b*-GALFF targeting PCR product were

wt1b_HA1_Gal4-For:

gacattttgaactcagatattctagtgtttttgcaacccagaaaatccgtcaccATGAAGCTACTGTCTTCTATCGAAC

and wt1b_HA2-KanR-Rev:

gcgctcaggtctctgacatccgatcccatcgggccgcacggctctgtcagTCAGAAGAACTCGTCAAGAA

(lower case indicates homology arms**).**

Immunofluorescence

Embryos were fixed overnight in 4% paraformaldehyde in PBS, washed in 0.01% PBS-Tween-20 (Sigma) and

permeabilized with 0.5% Triton-X100 (Sigma) in PBS for 20 min. Several washing steps were followed by

blocking for 2 h with 5% goat serum, 5% BSA, 20 mM MgCl2 in PBS followed by overnight incubation with

the primary antibody at 4ºC. Secondary antibodies were diluted 1:500 in PBS and incubated for 3 h. Nuclei

were counterstained with DAPI (Invitrogen) for 30 min. After several washes, embryos were mounted in

Vectashield (Vector).

The antibodies and stains for immunofluorescence detection were as follows: anti-myosin heavy chain (MF20,

DSHB) at a 1:20 dilution, anti-pH3 (Millipore) at 1:100, anti-GFP (aveslab) at 1:1000, anti-pSmad1/5 (Cell

Signaling Technology) at 1:100, Phalloidin-488 (Thermo Fisher) at 1:100, anti-myosin IIA (Sigma) at 1:100

and anti-mRFP (Abcam) at 1:250. Secondary antibodies were the following: anti-mouse-Cy3 (The Jackson

Laboratory), anti-mouse IgG2b 568 (Invitrogen), anti-chicken 488 (Life Technologies), anti-rabbit 647

(Thermo Scientific), all diluted at 1:500.

Embryos were imaged with a Zeiss 780 confocal microscope fitted with a 20× objective 1.0 NA with a

dipping lens. Z-stacks were taken every 3–5 µm. Maximal projections of images were 3D reconstructed in

whole-mount views using IMARIS software (Bitplane Scientific Software). The pericardial ventral tissue was

digitally removed to provide a clearer view of the heart. Optical sections of 1–3 z-slices were also

reconstructed.

Quantification of DP and PE cells

PE cells have been described to emerge from two main regions of the DP: the avcPE appears close to the atrio-

ventricular canal, and the vpPE around the venous pole. We counted each cell in each z plane using DAPI and

GFP expression using the line epi:GFP. We took care not to count any cell twice. Cells with a round

morphology at the vp or avc region were counted as PE cells, Cells with a flat morphology in the DP were

counted as DP cells. See Supplementary Fig. 5 and Supplementary Movie 18 for further explanation.

Actin mean intensity measurement

Images from 3 embryos of each condition at 60 hpf were acquired at the same conditions with a Zeiss 780

confocal microscope. Z-stacks were taken every 5 μm. Acquired images were in monochrome 8-bits. Z-slides

where the PE was present were opened in ImageJ (NIH) and squared areas of 3.32 x 3.32 pixels were drawn in

the channel of the PE cluster. The mean intensity was calculated for actin (phalloidin staining) in this PE

region. The intensity brightness sum was normalized to the number of pixels in the selected area to obtain the

actin mean intensity brightness in the PE. Brightness was in arbitrary units from 0–255 (where 0 is a pixel with

no staining pixel and 255 a saturated pixel). Mean intensity values were represented as spots in a histogram.

Pharmacological treatments

Embryos were manually dechorionated and incubated with compounds from 48 hpf onwards (unless otherwise

stated). The following compounds were used: aphidilcolin (150 μM), hydroxyurea (20 μM), nocodazole (0.01

mg/ml), BDM (10–20 mM), LDN-193189 (20mM), cytocalasin D (2 μM) (all from Sigma), BLEB (25–50

μM; Abcam), jasplakinolide (0.15 μM; Thermofisher), RO-4929097 (10 μM; Selleckchem), 4-OHT (5 μM;

Sigma).

In situ hybridization

ISH on whole mount embryos was performed as described[98] using riboprobes against full coding sequence of

bmp4 or *bmp2b* cDNAs as well as *notch1b*[99]. Embryos at 60 hpf or 80 hpf were fixed in 4% PFA overnight,

dehydrated in methanol series and stored at -20°C until its use. On day 1, embryos was bleached in 1,5% of

H_2O_2 in methanol, rehydrated, washed in TBS with 0.1% Tween20 (TBST), digested with proteinase K 10 µg

ml^{-1} for 17 min, rinsed in TBST, blocked the endogenous alkaline phosphatase with 0,1M triethanolamine pH

8 with 25 µl/ml of acetic anhydride for 20 min, washed in TBST, re-fixed in 4% PFA for 20 min. After

washing again in TBST they were pre-hybridized at 68°C for at least 1 h. The antisense riboprobe was added

at 0.5 µg ml^{-1}. After overnight hybridization, two washed with 50%Formamide/5xSSC plus 2% Tween20, four

washed with 2xSSC plus 0,2% Tween20, all at 68°C. Then, embryos were transferred to RT, washed in TBST

and incubated with 10% heat inactivated goat serum, 1,2% of blocking reagent (Roche, 11096176001) in

maleic acid buffer (MABT). Then, embryos were incubated overnight with 1:4000 dilution of anti

digoxigenin-AP antibody (Roche, 11093274910) in blocking solution. After overnight, embryos were washed

in MABT and developed in BM-Purple until signal was detected.

ISH on paraffin sections were done following commercial RNAscope protocol (Advanced Cell Diagnostics)

[100].

For fluorescent *in situ* hybridization combined with immunostaining in paraffin sections were deparaffinized,

post-fixed 20 minutes with PFA 4%, washed with PBS, treated with proteinase K 10 µm ml^{-1} for 10 minutes at

37 ºC, washed with PBS, post-fixed with PFA 4% for 5 minutes, washed with PBS, treated with HCl 0.07 N

for 15 minutes, washed with PBS, treated with 0.25% acetic anhydride in triethanolamine 0.1 M pH 8 for 10

minutes, washed with PBS, washed with RNase free water and then hybridized with the probe in pre-

hybridization buffer over night at 65 ºC. The following day sections where washed twice with post-

hybridization buffer 1 (50% Formamide, 5xSSC, 1% SDS) for 30 minutes at 65 ºC and twice more with post-

hybridization buffer 2 (50% Formamide, 2xSSC, 1% SDS). Then, sections were washed with MABT buffer at

room temperature, and incubated at least 2 hours in blocking solution at room temperature. They were next

incubated overnight with anti-digoxigenin-POD antibody (1:500) in blocking solution. The third day, they

were washed with MABT for several hours, after that sections were incubated with 1:200 tyramides (Perkin-

Elmer, NEL701A001KT), washed with PBST. Following with the immunofluorescence incubated the sections

with Myosin Heavy Chain antibody (DSHB hybridoma bank, MF20 1:20) overnight at 4°C. The fourth day

washed the slices and incubated with secondary antibody and counterstained with DAPI.

In vivo imaging

Embryos were transferred to fish water containing 0.2 mg/ml tricaine (Sigma) and 0.0033% 1-phenyl-2-

thiourea (Sigma), and immobilized in 0.7% agarose (NuSieve GT Agarose, Lonza) in a 35-mm Petridish with

a glass cover (MatTek Corporation). Zebrafish hearts were scanned bidirectionally at 30 frames per second

(fps) with an SP5 confocal microscope (Leica) using a 20× glycerol immersion objective with 0.7 NA. Videos

were acquired every 5 µm, with a line average of 6 and a pinhole of 1.9 AU. Around 65 z-stack videos were

acquired per heart every 10-15 min. GFP, red and brightfield channels were acquired simultaneously.

High resolution *in vivo* imaging was performed with the Zeiss LSM880 airy scan fastmode, using a

40x/1.1 water immersion objective lens. Sampling was performed with 1x Nyquist Coefficient parameters.

Airy scan processing was performed in ZEN 2.3.

Realigned 4D data sets were displayed and analyzed using Imaris (Bitplane AG) or ImageJ.

Drift correction

Small movements of the plate and the embryo in the agarose were manually suppressed using a drift

correction, selecting a particle, usually a noise voxel, following the movement to be corrected in two

consecutive frames. Moreover, a posterior drift correction was applied to suppress the intrinsic movement of

the developing embryo. Microscope and growth drift was corrected in Imaris. The venous pole was identified

as a stable reference structure and tracked in each frame. Subsequently, the "Correct drift" tool was applied to

the resulting track.

Cell tracking

Cell tracking was performed using a built-in Imaris tool, allowing for an automatic creation of 4D trajectories

of dorsal pericardial cells. A region of interest was manually selected and after applying a background

subtraction algorithm – Gaussian filter – an autoregressive algorithm is applied to perform cell tracking. Such

autoregressive algorithm works under the assumption that particles move from frame to frame in a quasi-

predictive fashion, interpolating the data from previous frames. This approach is pertinent in cells embedded in

a tissue and hence was chosen. 4D trajectories were further filtered out in terms of duration and overall

displacement to remove false-positive trajectories that would add noise to further calculations.

Divergence of velocity field calculator tool

The Algorithm to obtain the divergence of the dorsal pericardium velocity field was determined as follows:

Divergence of the velocity field associated with the movements of the dorsal pericardium was calculated as an

indicator of the expansion of the tissue. The divergence is a mathematical operator that, when applied to a

velocity field, indicates how much a set of particles expand or constrict in space. In this particular case, the set

of particles is given by the dorsal pericardial cells, hence giving an estimate of the expansion/constriction of

the tissue and characterizing it kinematically. It is mathematically described in Equation 1.

$$\nabla \cdot v = \frac{\partial v_x}{\partial x} + \frac{\partial v_y}{\partial y} + \frac{\partial v_z}{\partial z}$$

To do so, a customized Matlab code was created. Taking as an input the 4D trajectories calculated

with Imaris a 2D grid was created, which was used to interpolate the z-position of the dorsal pericardial cells,

hence resembling the geometry of the dorsal pericardium. The spacing of the grid satisfies the Nyquist criteria,

such that $\Delta x_{grid} < 2*$size of the cell nucleus. This interpolation was calculated using a Delaunay triangulation

method. In a similar procedure, each velocity component was interpolated across the mesh using a spline

method. The interpolated velocity field and z-position were used to calculate the divergence of the velocity

field.

Since the divergence calculated takes into account 3 dimensions, but the geometry of the tissue is a

laminar 2D surface embedded in a 3D space, 2D divergence were taken in the XY, YZ, XZ planes and

algebraically operated to obtain the 3D divergence as described in Equation 2. $\nabla \cdot v = \frac{1}{2}\sum_{i=1}^{3}\sum_{j=1}^{3}\nabla \cdot v_{i,j} =$

$\frac{1}{2}\sum_{i=1}^{3}\sum_{j=1}^{3}(\frac{\partial v_i}{\partial i} + \frac{\partial v_j}{\partial j})$ (Equation 2). After calculation of the divergence field each node of the z-interpolated

surface is assigned a divergence value, with positive values indicating an expansion and negative values a

constriction.

Additionally, the code provides the mean divergence of the surface for each time frame, according to Equation

3. $\langle \nabla \cdot v \rangle = \frac{1}{S}\iint \nabla \cdot v \, dxdy$ (Equation 3).

Angle calculation and directionality to midline

To calculate the angle of cell movement to the midline, epi:GFP embryos were *in vivo* imaged from 52 hpf.

The midline is a defined line that runs from the venous pole to the outflow tract of the heart tube. Dorsal

pericardium epi:GFP$^+$ cells were tracked using Imaris software during the time-lapse, and the movement

vector was calculated regarding the first and the last position of the cell. The angles that form the cell tracks

vectors with the midline were calculated in ImageJ. The directionality of the cell tracks in relation to the

midline was calculated with regards to the relative distance of the first and last cell track point to the midline

vector. Distance of first track point to midline > distance of last track point = cell movement towards midline;

distance of first track point to midline < distance of last track point = cell movement against midline.

Statistical analysis

Student's t test for comparisons between two groups or one-way ANOVA for comparisons between more than

two groups was used when normal distribution could be assumed. When the normality assumption could not

be verified with a reliable method, the Kruskal-Wallis test was used. Model assumptions of normality and

homogeneity were checked with conventional residual plots. The specific test used in each comparison is

indicated in the figure legend. Calculations were made with Microsoft Excel and GraphPad. P-values are

indicated either in the figure legends, the main text or summarized in Supplementary Table 1.

ACKNOWLEDGMENTS

We are grateful to the Animal facility and Microscopy Units from CNIC, and the Anatomy, Histology and

Neuroscience department from the UAM, MIC-Bern and J. Langa and R. Baal for fish husbandry of the

University of Bern. We thank Mathias Hammerschmidt, Salim Seyfried, Carl-Philipp Heisenberg and Nikolay

Ninov for sharing transgenic lines. N.M. was funded by the Spanish Ministry of Economy and

Competitiveness through grant BFU2014–56970–P (Plan Estatal de Investigación Científica y Técnica y de

Innovación 2013-2016. Programa Estatal de I+D+i Orientada a los Retos de la Sociedad Retos Investigación:

Proyectos I+D+i 2016, del Ministerio de Economía competitividad e Industria), and co-funding by Fondo

Europeo de Desarrollo Regional (FEDER). N.M. is also supported by the Swiss National Science Foundation

grant ANR-SNF 310030E-164245, the ERC starting grant 337703 and European Industrial Doctorate Program

EID 722427. L.A-D was funded through the postdoctoral fellowship Ayudas Postdoctorales 2013. JL de la

Pompa was supported by grants SAF2016-78370-R, CB16/11/00399 (CIBER CV) and RD16/0011/0021

(TERCEL) from the Spanish Ministry of Economy, Industry and Competitiveness (MEIC). The CNIC is

supported by the Ministry of Economy, Industry and Competitiveness (MEIC) and the Pro CNIC Foundation,

and is a Severo Ochoa Center of Excellence (MEIC award SEV-2015-0505).

BIBLIOGRAPHY

1   Olivey, H. E. & Svensson, E. C. Epicardial-myocardial signaling directing coronary
vasculogenesis. *Circ Res* **106**, 818-832, doi:10.1161/circresaha.109.209197 (2010).
2   Perez-Pomares, J. M. & de la Pompa, J. L. Signaling during epicardium and coronary vessel
development. *Circ Res* **109**, 1429-1442, doi:10.1161/circresaha.111.245589 (2011).
3   Chau, Y. Y. *et al.* Visceral and subcutaneous fat have different origins and evidence supports
a mesothelial source. *Nat Cell Biol* **16**, 367-375, doi:10.1038/ncb2922 (2014).
4   Duenas, A., Aranega, A. E. & Franco, D. More than Just a Simple Cardiac Envelope; Cellular
Contributions of the Epicardium. *Front Cell Dev Biol* **5**, 44, doi:10.3389/fcell.2017.00044
(2017).
5   Wessels, A. *et al.* Epicardially derived fibroblasts preferentially contribute to the parietal
leaflets of the atrioventricular valves in the murine heart. *Dev Biol* **366**, 111-124,
doi:10.1016/j.ydbio.2012.04.020 (2012).
6   Kennedy-Lydon, T. & Rosenthal, N. Cardiac regeneration: epicardial mediated repair. *Proc
Biol Sci* **282**, 20152147, doi:10.1098/rspb.2015.2147 (2015).
7   Maya-Ramos, L., Cleland, J., Bressan, M. & Mikawa, T. Induction of the Proepicardium. *J
Dev Biol* **1**, 82-91, doi:10.3390/jdb1020082 (2013).
8   Schulte, I., Schlueter, J., Abu-Issa, R., Brand, T. & Manner, J. Morphological and molecular
left-right asymmetries in the development of the proepicardium: a comparative analysis on
mouse and chick embryos. *Dev Dyn* **236**, 684-695, doi:10.1002/dvdy.21065 (2007).
9   Liu, J. & Stainier, D. Y. Tbx5 and Bmp signaling are essential for proepicardium
specification in zebrafish. *Circ Res* **106**, 1818-1828, doi:<u>CIRCRESAHA.110.217950 [pii]</u>
<u>10.1161/CIRCRESAHA.110.217950</u> (2010).
10   Hirose, T. *et al.* PAR3 is essential for cyst-mediated epicardial development by establishing
apical cortical domains. *Development* **133**, 1389-1398, doi:10.1242/dev.02294 (2006).
11   Serluca, F. C. Development of the proepicardial organ in the zebrafish. *Dev Biol* **315**, 18-27,
doi:S0012-1606(07)01431-5 [pii] 10.1016/j.ydbio.2007.10.007 (2008).
12   Tandon, P., Wilczewski, C. M., Williams, C. E. & Conlon, F. L. The Lhx9-integrin pathway
is essential for positioning of the proepicardial organ. *Development* **143**, 831-840,
doi:10.1242/dev.129551 (2016).
13   Wu, M. *et al.* Epicardial spindle orientation controls cell entry into the myocardium. *Dev Cell*
19, 114-125, doi:10.1016/j.devcel.2010.06.011 (2010).
14   Peralta, M. *et al.* Heartbeat-driven pericardiac fluid forces contribute to epicardium
morphogenesis. *Curr Biol* **23**, 1726-1735, doi:10.1016/j.cub.2013.07.005 (2013).
15   Plavicki, J. S. *et al.* Multiple modes of proepicardial cell migration require heartbeat. *BMC
Dev Biol* **14**, 18, doi:10.1186/1471-213x-14-18 (2014).

16 Tada, M. & Heisenberg, C. P. Convergent extension: using collective cell migration and cell
intercalation to shape embryos. *Development* **139**, 3897-3904, doi:10.1242/dev.073007
(2012).
17 Aegerter-Wilmsen, T. *et al.* Integrating force-sensing and signaling pathways in a model for
the regulation of wing imaginal disc size. *Development* **139**, 3221-3231,
doi:10.1242/dev.082800 (2012).
18 Hiscock, T. W. & Megason, S. G. Orientation of Turing-like Patterns by Morphogen
Gradients and Tissue Anisotropies. *Cell Syst* **1**, 408-416, doi:10.1016/j.cels.2015.12.001
(2015).
19 Heisenberg, C. P. & Bellaiche, Y. Forces in tissue morphogenesis and patterning. *Cell* **153**,
948-962, doi:10.1016/j.cell.2013.05.008 (2013).
20 Levayer, R. & Lecuit, T. Biomechanical regulation of contractility: spatial control and
dynamics. *Trends Cell Biol* **22**, 61-81, doi:10.1016/j.tcb.2011.10.001 (2012).
21 Martin, A. C., Kaschube, M. & Wieschaus, E. F. Pulsed contractions of an actin-myosin
network drive apical constriction. *Nature* **457**, 495-499, doi:10.1038/nature07522 (2009).
22 Munjal, A. & Lecuit, T. Actomyosin networks and tissue morphogenesis. *Development* **141**,
1789-1793, doi:10.1242/dev.091645 (2014).
23 Lecuit, T. & Yap, A. S. E-cadherin junctions as active mechanical integrators in tissue
dynamics. *Nat Cell Biol* **17**, 533-539, doi:10.1038/ncb3136 (2015).
24 Martin, A. C., Gelbart, M., Fernandez-Gonzalez, R., Kaschube, M. & Wieschaus, E. F.
Integration of contractile forces during tissue invagination. *J Cell Biol* **188**, 735-749,
doi:10.1083/jcb.200910099 (2010).
25 Andres-Delgado, L. & Mercader, N. Interplay between cardiac function and heart
development. *Biochim Biophys Acta* **1863**, 1707-1716, doi:10.1016/j.bbamcr.2016.03.004
(2016).
26 Collins, M. M. & Stainier, D. Y. Organ Function as a Modulator of Organ Formation:
Lessons from Zebrafish. *Curr Top Dev Biol* **117**, 417-433, doi:10.1016/bs.ctdb.2015.10.017
(2016).
27 Ocana, O. H. *et al.* A right-handed signalling pathway drives heart looping in vertebrates.
Nature **549**, 86-90, doi:10.1038/nature23454 (2017).
28 Steed, E., Boselli, F. & Vermot, J. Hemodynamics driven cardiac valve morphogenesis.
Biochim Biophys Acta **1863**, 1760-1766, doi:10.1016/j.bbamcr.2015.11.014 (2016).
29 Peralta, M., Gonzalez-Rosa, J. M., Marques, I. J. & Mercader, N. The Epicardium in the
Embryonic and Adult Zebrafish. *J Dev Biol* **2**, 101-116, doi:10.3390/jdb2020101 (2014).
30 Eisenhoffer, G. T. *et al.* Crowding induces live cell extrusion to maintain homeostatic cell
numbers in epithelia. *Nature* **484**, 546-549, doi:10.1038/nature10999 (2012).
31 Vasquez, R. J., Howell, B., Yvon, A. M., Wadsworth, P. & Cassimeris, L. Nanomolar
concentrations of nocodazole alter microtubule dynamic instability in vivo and in vitro. *Mol
Biol Cell* **8**, 973-985 (1997).
32 Mutomba, M. C. & Wang, C. C. Effects of aphidicolin and hydroxyurea on the cell cycle and
differentiation of Trypanosoma brucei bloodstream forms. *Mol Biochem Parasitol* **80**, 89-102
(1996).
33 Kadeer, A. *et al.* Plectin is a novel regulator for apical extrusion of RasV12-transformed cells.
Sci Rep **7**, 44328, doi:10.1038/srep44328 (2017).
34 Saitoh, S. *et al.* Rab5-regulated endocytosis plays a crucial role in apical extrusion of
transformed cells. *Proc Natl Acad Sci U S A* **114**, E2327-E2336,
doi:10.1073/pnas.1602349114 (2017).
35 Wu, S. K., Lagendijk, A. K., Hogan, B. M., Gomez, G. A. & Yap, A. S. Active contractility at
E-cadherin junctions and its implications for cell extrusion in cancer. *Cell Cycle* **14**, 315-322,
doi:10.4161/15384101.2014.989127 (2015).

36 Martin, A. C. & Goldstein, B. Apical constriction: themes and variations on a cellular
mechanism driving morphogenesis. *Development* **141**, 1987-1998, doi:10.1242/dev.102228
(2014).
37 Noguchi, T. Q., Morimatsu, M., Iwane, A. H., Yanagida, T. & Uyeda, T. Q. The role of
structural dynamics of actin in class-specific myosin motility. *PLoS One* **10**, e0126262,
doi:10.1371/journal.pone.0126262 (2015).
38 Schaub, S., Bohnet, S., Laurent, V. M., Meister, J. J. & Verkhovsky, A. B. Comparative maps
of motion and assembly of filamentous actin and myosin II in migrating cells. *Mol Biol Cell*
18, 3723-3732, doi:10.1091/mbc.E06-09-0859 (2007).
39 Wilson, C. A. *et al.* Myosin II contributes to cell-scale actin network treadmilling through
network disassembly. *Nature* **465**, 373-377, doi:10.1038/nature08994 (2010).
40 Ostap, E. M. 2,3-Butanedione monoxime (BDM) as a myosin inhibitor. *J Muscle Res Cell*
Motil **23**, 305-308 (2002).
41 Holzinger, A. Jasplakinolide: an actin-specific reagent that promotes actin polymerization.
Methods Mol Biol **586**, 71-87, doi:10.1007/978-1-60761-376-3_4 (2009).
42 de Pater, E. *et al.* Bmp signaling exerts opposite effects on cardiac differentiation. *Circ Res*
110, 578-587, doi:10.1161/circresaha.111.261172 (2012).
43 Palencia-Desai, S. *et al.* Myocardium and BMP signaling are required for endocardial
differentiation. *Development* **142**, 2304-2315, doi:10.1242/dev.118687 (2015).
44 Wu, C. C. *et al.* Spatially Resolved Genome-wide Transcriptional Profiling Identifies BMP
Signaling as Essential Regulator of Zebrafish Cardiomyocyte Regeneration. *Dev Cell* **36**, 36-
49, doi:10.1016/j.devcel.2015.12.010 (2016).
45 Chocron, S., Verhoeven, M. C., Rentzsch, F., Hammerschmidt, M. & Bakkers, J. Zebrafish
Bmp4 regulates left-right asymmetry at two distinct developmental time points. *Dev Biol* **305**,
577-588, doi:10.1016/j.ydbio.2007.03.001 (2007).
46 Collery, R. F. & Link, B. A. Dynamic smad-mediated BMP signaling revealed through
transgenic zebrafish. *Dev Dyn* **240**, 712-722, doi:10.1002/dvdy.22567 (2011).
47 Cuny, G. D. *et al.* Structure-activity relationship study of bone morphogenetic protein (BMP)
signaling inhibitors. *Bioorg Med Chem Lett* **18**, 4388-4392, doi:10.1016/j.bmcl.2008.06.052
(2008).
48 Garside, V. C., Chang, A. C., Karsan, A. & Hoodless, P. A. Co-ordinating Notch, BMP, and
TGF-beta signaling during heart valve development. *Cell Mol Life Sci* **70**, 2899-2917,
doi:10.1007/s00018-012-1197-9 (2013).
49 Jain, R., Rentschler, S. & Epstein, J. A. Notch and cardiac outflow tract development. *Ann N*
Y Acad Sci **1188**, 184-190, doi:10.1111/j.1749-6632.2009.05099.x (2010).
50 de la Pompa, J. L. & Epstein, J. A. Coordinating tissue interactions: Notch signaling in
cardiac development and disease. *Dev Cell* **22**, 244-254, doi:10.1016/j.devcel.2012.01.014
(2012).
51 Luna-Zurita, L. *et al.* Integration of a Notch-dependent mesenchymal gene program and
Bmp2-driven cell invasiveness regulates murine cardiac valve formation. *J Clin Invest* **120**,
3493-3507, doi:10.1172/jci42666 (2010).
52 Luxan, G., D'Amato, G., MacGrogan, D. & de la Pompa, J. L. Endocardial Notch Signaling in
Cardiac Development and Disease. *Circ Res* **118**, e1-e18, doi:10.1161/circresaha.115.305350
(2016).
53 del Monte, G. *et al.* Differential Notch signaling in the epicardium is required for cardiac
inflow development and coronary vessel morphogenesis. *Circ Res* **108**, 824-836,
doi:10.1161/CIRCRESAHA.110.229062 (2011).
54 Scheer, N., Groth, A., Hans, S. & Campos-Ortega, J. A. An instructive function for Notch in
promoting gliogenesis in the zebrafish retina. *Development* **128**, 1099-1107 (2001).

| 757 | 55 | Herwig, L. *et al.* Distinct cellular mechanisms of blood vessel fusion in the zebrafish embryo. *Curr Biol* **21**, 1942-1948, doi:10.1016/j.cub.2011.10.016 (2011). |

55 Herwig, L. *et al.* Distinct cellular mechanisms of blood vessel fusion in the zebrafish embryo.
Curr Biol **21**, 1942-1948, doi:10.1016/j.cub.2011.10.016 (2011).

56 Munch, J., Grivas, D., Gonzalez-Rajal, A., Torregrosa-Carrion, R. & de la Pompa, J. L. Notch
signalling restricts inflammation and serpine1 expression in the dynamic endocardium of the
regenerating zebrafish heart. *Development* **144**, 1425-1440, doi:10.1242/dev.143362 (2017).

57 Poon, K. L., Liebling, M., Kondrychyn, I., Garcia-Lecea, M. & Korzh, V. Zebrafish cardiac
enhancer trap lines: new tools for in vivo studies of cardiovascular development and disease.
Dev Dyn **239**, 914-926, doi:10.1002/dvdy.22203 (2010).

58 Bruckner, K., Perez, L., Clausen, H. & Cohen, S. Glycosyltransferase activity of Fringe
modulates Notch-Delta interactions. *Nature* **406**, 411-415, doi:10.1038/35019075 (2000).

59 Moloney, D. J. *et al.* Fringe is a glycosyltransferase that modifies Notch. *Nature* **406**, 369-375,
doi:10.1038/35019000 (2000).

60 Ninov, N. *et al.* Metabolic regulation of cellular plasticity in the pancreas. *Curr Biol* **23**,
1242-1250, doi:10.1016/j.cub.2013.05.037 (2013).

61 Lawson, N. D. & Weinstein, B. M. In vivo imaging of embryonic vascular development using
transgenic zebrafish. *Dev Biol* **248**, 307-318 (2002).

62 Knodler, L. A. *et al.* Dissemination of invasive Salmonella via bacterial-induced extrusion of
mucosal epithelia. *Proc Natl Acad Sci U S A* **107**, 17733-17738,
doi:10.1073/pnas.1006098107 (2010).

63 Ritchie, J. M. *et al.* Inflammation and disintegration of intestinal villi in an experimental
model for Vibrio parahaemolyticus-induced diarrhea. *PLoS Pathog* **8**, e1002593,
doi:10.1371/journal.ppat.1002593 (2012).

64 Simovitch, M. *et al.* EspM inhibits pedestal formation by enterohaemorrhagic Escherichia
coli and enteropathogenic E. coli and disrupts the architecture of a polarized epithelial
monolayer. *Cell Microbiol* **12**, 489-505, doi:10.1111/j.1462-5822.2009.01410.x (2010).

65 Marinari, E. *et al.* Live-cell delamination counterbalances epithelial growth to limit tissue
overcrowding. *Nature* **484**, 542-545, doi:10.1038/nature10984 (2012).

66 Gibson, M. C. & Perrimon, N. Extrusion and death of DPP/BMP-compromised epithelial
cells in the developing Drosophila wing. *Science* **307**, 1785-1789,
doi:10.1126/science.1104751 (2005).

67 Rosenblatt, J., Raff, M. C. & Cramer, L. P. An epithelial cell destined for apoptosis signals its
neighbors to extrude it by an actin- and myosin-dependent mechanism. *Curr Biol* **11**, 1847-
1857 (2001).

68 Kajita, M. *et al.* Filamin acts as a key regulator in epithelial defence against transformed cells.
Nat Commun **5**, 4428, doi:10.1038/ncomms5428 (2014).

69 Hogan, C. *et al.* Characterization of the interface between normal and transformed epithelial
cells. *Nat Cell Biol* **11**, 460-467, doi:10.1038/ncb1853 (2009).

70 Kajita, M. *et al.* Interaction with surrounding normal epithelial cells influences signalling
pathways and behaviour of Src-transformed cells. *J Cell Sci* **123**, 171-180,
doi:10.1242/jcs.057976 (2010).

71 Leung, C. T. & Brugge, J. S. Outgrowth of single oncogene-expressing cells from suppressive
epithelial environments. *Nature* **482**, 410-413, doi:10.1038/nature10826 (2012).

72 Vidal, M., Larson, D. E. & Cagan, R. L. Csk-deficient boundary cells are eliminated from
normal Drosophila epithelia by exclusion, migration, and apoptosis. *Dev Cell* **10**, 33-44,
doi:10.1016/j.devcel.2005.11.007 (2006).

73 Bertrand, J. Y. *et al.* Haematopoietic stem cells derive directly from aortic endothelium
during development. *Nature* **464**, 108-111, doi:10.1038/nature08738 (2010).

74 Zovein, A. C. *et al.* Fate tracing reveals the endothelial origin of hematopoietic stem cells.
Cell Stem Cell **3**, 625-636, doi:10.1016/j.stem.2008.09.018 (2008).

75   Francou, A., De Bono, C. & Kelly, R. G. Epithelial tension in the second heart field promotes
mouse heart tube elongation. *Nat Commun* **8**, 14770, doi:10.1038/ncomms14770 (2017).
76   Krieg, M. *et al.* Tensile forces govern germ-layer organization in zebrafish. *Nat Cell Biol* **10**,
429-436, doi:10.1038/ncb1705 (2008).
77   Miller, C. J. & Davidson, L. A. The interplay between cell signalling and mechanics in
developmental processes. *Nat Rev Genet* **14**, 733-744, doi:10.1038/nrg3513 (2013).
78   Jidigam, V. K., Srinivasan, R. C., Patthey, C. & Gunhaga, L. Apical constriction and
epithelial invagination are regulated by BMP activity. *Biol Open* **4**, 1782-1791,
doi:10.1242/bio.015263 (2015).
79   Norden, C., Young, S., Link, B. A. & Harris, W. A. Actomyosin is the main driver of
interkinetic nuclear migration in the retina. *Cell* **138**, 1195-1208,
doi:10.1016/j.cell.2009.06.032 (2009).
80   Zhang, M. *et al.* BMP-2 overexpression augments vascular smooth muscle cell motility by
upregulating myosin Va via Erk signaling. *Oxid Med Cell Longev* **2014**, 294150,
doi:10.1155/2014/294150 (2014).
81   Grieve, A. G. & Rabouille, C. Extracellular cleavage of E-cadherin promotes epithelial cell
extrusion. *J Cell Sci* **127**, 3331-3346, doi:10.1242/jcs.147926 (2014).
82   Fourel, L. *et al.* beta3 integrin-mediated spreading induced by matrix-bound BMP-2 controls
Smad signaling in a stiffness-independent manner. *J Cell Biol* **212**, 693-706,
doi:10.1083/jcb.201508018 (2016).
83   Kurosawa, T. *et al.* Nephronectin Expression Is Up-Regulated by BMP-2. *Biol Pharm Bull* **39**,
1211-1215, doi:10.1248/bpb.b15-00941 (2016).
84   Kruithof, B. P. *et al.* BMP and FGF regulate the differentiation of multipotential pericardial
mesoderm into the myocardial or epicardial lineage. *Dev Biol* **295**, 507-522,
doi:10.1016/j.ydbio.2006.03.033 (2006).
85   van Wijk, B. *et al.* Epicardium and myocardium separate from a common precursor pool by
crosstalk between bone morphogenetic protein- and fibroblast growth factor-signaling
pathways. *Circ Res* **105**, 431-441, doi:10.1161/circresaha.109.203083 (2009).
86   Iyer, D. *et al.* Robust derivation of epicardium and its differentiated smooth muscle cell
progeny from human pluripotent stem cells. *Development* **143**, 904, doi:10.1242/dev.136143
(2016).
87   Witty, A. D. *et al.* Generation of the epicardial lineage from human pluripotent stem cells.
Nat Biotechnol **32**, 1026-1035, doi:10.1038/nbt.3002 (2014).
88   Wang, Y. *et al.* Endocardial to myocardial notch-wnt-bmp axis regulates early heart valve
development. *PLoS One* **8**, e60244, doi:10.1371/journal.pone.0060244 (2013).
89   Schlueter, J. & Brand, T. Origin and fates of the proepicardium. *Aswan Heart Centre Science*
& Practice Series **2011**, 11 (2011).
90   Rohr, S., Otten, C. & Abdelilah-Seyfried, S. Asymmetric involution of the myocardial field
drives heart tube formation in zebrafish. *Circ Res* **102**, e12-19,
doi:10.1161/CIRCRESAHA.107.165241 (2008).
91   Scheer, N., Riedl, I., Warren, J. T., Kuwada, J. Y. & Campos-Ortega, J. A. A quantitative
analysis of the kinetics of Gal4 activator and effector gene expression in the zebrafish. *Mech*
Dev **112**, 9-14 (2002).
92   Maitre, J. L. *et al.* Adhesion functions in cell sorting by mechanically coupling the cortices of
adhering cells. *Science* **338**, 253-256, doi:10.1126/science.1225399 (2012).
93   Behrndt, M. *et al.* Forces driving epithelial spreading in zebrafish gastrulation. *Science* **338**,
257-260, doi:10.1126/science.1224143 (2012).
94   Mosimann, C. & Zon, L. I. Advanced zebrafish transgenesis with Tol2 and application for
Cre/lox recombination experiments. *Methods Cell Biol* **104**, 173-194, doi:10.1016/b978-0-12-
374814-0.00010-0 (2011).

95   Asakawa, K. *et al.* Genetic dissection of neural circuits by Tol2 transposon-mediated Gal4
gene and enhancer trapping in zebrafish. *Proc Natl Acad Sci U S A* **105**, 1255-1260,
doi:10.1073/pnas.0704963105 (2008).
96   Bussmann, J. & Schulte-Merker, S. Rapid BAC selection for tol2-mediated transgenesis in
zebrafish. *Development* **138**, 4327-4332, doi:10.1242/dev.068080 (2011).
97   Suster, M. L., Abe, G., Schouw, A. & Kawakami, K. Transposon-mediated BAC transgenesis
in zebrafish. *Nat Protoc* **6**, 1998-2021, doi:10.1038/nprot.2011.416 (2011).
98   Jowett, T. & Lettice, L. Whole-mount in situ hybridizations on zebrafish embryos using a
mixture of digoxigenin- and fluorescein-labelled probes. *Trends Genet* **10**, 73-74 (1994).
99   Westin, J. & Lardelli, M. Three novel Notch genes in zebrafish: implications for vertebrate
Notch gene evolution and function. *Dev Genes Evol* **207**, 51-63, doi:10.1007/s004270050091
(1997).
100   Wang, F. *et al.* RNAscope: a novel in situ RNA analysis platform for formalin-fixed,
paraffin-embedded tissues. *J Mol Diagn* **14**, 22-29, doi:10.1016/j.jmoldx.2011.08.002 (2012).
870

Figure Legends

Fig. 1 Collective cell movements and apical extrusion lead to PE delamination. (**a**) Scheme of lateral view
of a 60 hpf zebrafish embryo. Zoom on heart below overview. Lateral view is rotated by -90°. Legend for
structures are on right side of the panel. Shown are the dorsal and ventral pericardium, ventricle, atrium,
outflow tract venous pole, midline and proepicardium (PE). (**b**) 3D ventral view of the heart region in a
epi:GFP; myl7:mRFP double transgenic larvae. epi:GFP[+] cells in the dorsal pericardium (DP) and PE are
shown in green, *myl7:mRFP* in red (myocardium). Maximal projection from an *in vivo* time-lapse at different
time-points is shown. White arrows point to PE clusters (see also Movie 1). (**c**) First and last frame of an *in
vivo* time-lapse DP cells from an *epi:GFP* larvae; the midline is shown by a discontinuous white line. Blue
dots indicate the tracked cells. Full colored tracks label the first time frame in purple and the last in red.
Arrows indicate overall direction of tracked cells towards the midline. (**d**) The half-rose diagram shows the
number of DP cells from 3 animals for which the angle of the track was measured relative to the midline. (**e**)
3D reconstruction of the divergence field of the DP on top of images from **b**. Blue regions represent
constriction and red expansion of the tissue. Black arrows point to PE clusters (see also Movie 2). (**f**) epi:GFP
in vivo time-lapse. A maximum intensity projection of 22 μm is shown. Left panel shows an overview of the
vp and DP at 52 hpf. Right panels show zoomed frames of the time-lapse from 52 to 67 hpf. Colored
arrowheads indicate emerging PE cells. DP cells surrounding emerging PE cells are labeled with colored
asterisks (shown is 1 out of 10 acquisitions, see also Movie 3). ap, arterial pole; at, atrium; hpf; hours post
fertilization; v, ventricle; vp, venous pole. Scale bar: 50 μm (in **f** overview 20μm and time-lapse 10μm).

Fig. 2 Proliferation of pericardial cells is necessary for PE formation. (**a**) Six frames of an epi:GFP *in vivo*
time-lapse (see also Movie 4). The dorsal pericardium (DP) was isolated and presented from a top view.

Division planes are indicated with red ($45° \leq \alpha \leq 90°$) and green lines ($0° \leq \alpha \leq 45°$). A representation of the

midline is shown in yellow. Micrometer values show the measured distance of the division to the midline (**b**)

Scheme showing the principle of measurements for distance and angles of division planes to midline. Graphs

show number of divisions during *in vivo* time-lapse relative to the midline (3 embryos with 20 cell divisions;

P-value= 0.2, unpaired Student's T-test) (**c**) and angle of midline to division plane relative to distance to the

midline (**d**) The division plane was more often perpendicular than parallel to the midline P-value= 0.0144,

unpaired Student's T-test. (**e**) Whole mount immunofluorescence with anti-GFP (green), Myosin Heavy Chain

(red) and anti-pH3 (white). DAPI counterstains nuclei (blue). Shown are optical transversal sections of a

control heart and hearts form larvae treated with the proliferation inhibitors aphidilcolin/hydroxyurea

(aph/hydrU) or nocodazol (noc). Zoomed view of the dotted PE area is shown on the right. Arrowheads point

to the PE. (**f**) Quantification of total number of pH3$^+$ cells in the pericardium. (**g**) Percentage of pH3$^+$ cells in

the DP compared with the PE. (**h**) Zoomed view of a unique pH3$^+$ PE cell in a control embryo. (**i**) PE cell

number from conditions shown in **e**.   (**j**) Scheme of apical PE extrusion mechanism: 1) DP is a flattened

mesothelium; 2) DP cells move towards the midline; 3) PE cells rounds up at the midline; 4) proliferation of

DP cells at the border site also contributes to the constriction and PE cells finally extrude. at, atrium; DP,

dorsal pericardium; hpf, hours post fertilization; PE, proepicardium; v, ventricle. Scale bar: 50 μm (in **h**

zoomed images, 20μm). All data are means ± s.d., one-way ANOVA followed by Kruskal-Wallis significant

difference test was used in **f** and **g**; (Unpaired two-tailed Student's *t*-test was used in panel **i**). * $P < 0.05$; ** P

< 0.01, *** $P < 0.001$.

Fig. 3 PE formation depends on actomyosin dynamics. (**a**) Upper panel shows a maximum projection of a

60 hpf zebrafish heart immunostained for Myosin Heavy Chain (MHC, white), myosin II-A (red) and GFP

(green). Nuclei are counterstained with DAPI (blue). The three bottom zoomed images are different views of

the PE region. Yellow arrows point to myosin II-A concentration sites. DP indicates a flat dorsal pericardial

cell. (**b**) Immunostaining as in **a**, but with phalloidin-488 to visualize F-actin (red). White arrows point to actin

concentration sites. (**c, d**) Ventral view of the heart region in a *epi:GFP; lifeActin-RFP* larvae at different time

points of an *in vivo* time-lapse. Ventral pericardium and heart tube was post-process eliminated and the heart

tube shape was drawn with a red discontinuous line. (**c**) Untreated control; white arrows mark sites of actin

concentration in the DP. (**d**) BDM-treated larvae. (**e**) Scheme at 58 hpf of the actin concentration at DP and PE

clusters during PE formation in untreated or BDM-treated larvae. Actin is in red, DP and PE cells in green. (**f**)

Maximal projection of hearts labeled for actin in fish untreated or treated with jasp and BDM. (**g**) Top panels

show a maximum projection of a 60 hpf zebrafish heart, and bottom panels show an optical section of epi:GFP

animals immunostained for MHC (red) and nuclei counterstained with DAPI (blue). Larvae were either

untreated or treated with combinations of jasp and BDM. White arrowheads indicate the PE. (**h**)

Quantification of PE cell number in **g**. at, atrium; BDM, Butanedione Monoxime; DP, dorsal pericardium; hpf,

hours post fertilization; jasp, jasplakinolide; PE, proepicardium; v, ventricle; vp; venous pole. Scale bar: 50

μm (in **a**''-**a**'''''4 and **b**''-**b**'''''is 10μm). Data are mean ± s.d., one-way ANOVA followed by Kruskal-Wallis
significant difference test. ***$P < 0.001$.

Fig. 4 Bmp2b rescues PE formation upon Myosin II inhibition. (**a**) Top panels show a maximum projection
of a 60 hpf zebrafish heart, and bottom panels show an optical section of epi:GFP animals immunostained for
GFP in green and Myosin Heavy Chain (MHC) in red and nuclei counterstained with DAPI (blue). Untreated
fish were compared with those overexpressing *bmp2b* with and without BDM or with those overexpressing
Noggin 3. Arrowheads point the PE. Arrows mark epicardial cells. (**b**) Quantification of PE cell number in **a**.
(**c**) Quantification of epicardial cell number at 60 hpf in *bmp2b*-overexpressed versus non-overexpressed
animals. (**d**) Optical sections through epi:GFP hearts at 60 hpf showing pSmpad1/5 staining. Zoomed views
are also shown. epi:GFP animals were immunostained for GFP in green, MHC in red, pSmad1/5 in white.
Nuclei were counterstained with DAPI (blue). Arrowheads point to PE cells and asterisks mark DP cells. (**e**)
Percentage of pSmad1/5$^+$ cells in the DP compared to the PE. (**f**) Total number of pSmad1/5$^+$ cells in the
pericardium at 60 hpf. (**g**) Optical section from an *in vivo* time-lapse of an BMP reporter line *BRE*:KuO/
epi:GFP embryo. Yellow arrowheads point to the double positive cells in the PE. at, atrium; BDM,
Butanedione monoxime; DP, dorsal pericardium; epic, epicardium; hpf, hours post fertilization; PE,
proepicardium; v, ventricle. Scale bar: 50 μm (in **a** and **d** zoomed images, 20μm). Data are mean ± s.d., one-
way ANOVA followed by Kruskal-Wallis significant difference test. Unpaired two-tailed Student's *t*-test was
used in panel **c** and **f.** * $P < 0.05$; ** $P < 0.01$, *** $P < 0.001$.

Fig. 5 Bmp2b controls actin rearrangements necessary for PE formation. (**a**) Heart and pericardial cavity
from untreated zebrafish compared with Bmp2-overexpressing and BDM-treated fish at 60 hpf. Zoomed view
of the DP-PE area is shown. Pericardium and PE are in green, myosin II-A is in red and nuclei were
counterstained with DAPI (blue). Yellow arrows mark myosin II-A apicocortical accumulation in PE cells. (**b**)
Maximum projection of 60 hpf zebrafish hearts; optical sections of the hearts are shown in the middle panels,
and zoomed images (bottom panels) are views of the PE region. Untreated fish were compared with BDM-
treated and *bmp2b*-overexpressing fish. Pericardium and PE are in green and actin is in red. Nuclei are
counterstained with DAPI. Arrowheads mark the PE cluster. Arrows point to actin concentration sites in the
PE cluster. (**c**) Quantification of actin intensity (arbitrary units) in PE cells in **b**. (**d**) Top images are maximum
projections of 60 hpf hearts from epi:GFP control or *bmp2b*-overexpressing fish treated with cytochalasin D
(cytD), immunostained for GFP (green), myosin heavy chain (MHC, red) and phalloidin-Alexa488 (actin,
white). Nuclei are counterstained with DAPI (blue). (**e**) Quantification of PE cell number from embryos as
shown in **d**. (**d**) Top images are maximum projections and optical sections of animals treated with LDN-193189 (LDN) and
jasplakinolide (jasp). epi:GFP$^+$ cells are in green, the MHC$^+$ heart tube in red, nuclei are counterstained with
DAPI (blue). (**g**) Quantification of PE cell number in **f**. at, atrium; BDM, Butanedione Monoxime; hpf, hours
post fertilisation; PE, proepicardium; v, ventricle. Scale bar: 50 μm except for **a** (10 μm) and zoomed views in

d and **f** (20μm). Data are mean ± s.d., one-way ANOVA followed by Kruskal-Wallis significant difference test.

*P < 0.05, **P < 0.01, ***P < 0.001.

Fig. 6 Effect of Bmp2b and myosin II inhibition on dorsal pericardial layer constriction. (**a-c**) First and

last frame of an epi:GFP *in vivo* time-lapse, (a) *bmp2b*-overexpressing, (**b**) BDM-treated or (**c**) Bmp2b

overexpression in BDM-treated larvae; midline is shown by a discontinuous white line. Colored tracks label

first time-frame in purple and the last in red. Arrows indicate overall direction of tracked cells. (**d**) Ventral

view of *epi:GFP;myl7:mRFP* hearts. Shown are maximal projections of: untreated, *bmp2b*-overexpressing,

BDM-treated and *bmp2b*-overexpression in BDM-treated fish. Bottom images show 3D reconstruction of the

divergence field of the DP; blue regions represent constriction and red-orange regions of tissue expansion.

Arrows point to PE cells; dotted lines draw the midline. (**e**) Mean divergence of the tracked cells within the DP

from *in vivo* time-lapses. (**f**) Percentage of cells that move towards or away from the midline. Data in **e** and **f**

are mean ± s.d., one-way ANOVA followed by Kruskal-Wallis significant difference test was used and is

shown with black line and asterisk the summary, ** *P* < 0.01. Unpaired two-tailed Student's *t*-test was also

used and shown with red lines and asterisks, * *P* < 0.05; ** *P* < 0.01, *** *P* < 0.001. ap, arterial pole; at,

atrium; BDM, Butanedione Monoxime; hpf, hours post fertilization; PE, proepicardium; v, ventricle; vp,

venous pole. Scale bar: 50 μm.

Fig. 7 Effect of Notch signaling activation on PE formation. (**a**) Maximal projection and optical sections at

60 hpf of control embryos and embryos overexpressing NICD (*UAS:NICD*) upon heat shock induction in all

cells (*hsp70:Gal4*), in endocardial and endothelial (*fli1a:Gal4*) pericardial and proepicardial cells (*wt1b:Gal4*).

Immunostaining was performed for GFP (green), MHC (red) and nuclei were counterstained with DAPI (blue).

Arrowheads point to the PE. (**b**) Quantification of PE cell number in **a**. (**c**) Maximal projection and optical

sections at 60 hpf of larvae BDM treatment. (**d**) Quantification of PE cell number in c. (**e**) Quantification of

pH3$^+$ cells in the pericardium. (**f**) Maximum projection of 60 hpf zebrafish hearts; optical sections are shown

in the middle panels, zoomed images (bottom panels) are views of the PE region. Untreated fish were

compared with RO-treated animals and larvae overexpressing *NICD*. GFP labels the pericardium and PE

(green), actin labeled with phalloidin is shown in red. Nuclei are counterstained with DAPI (blue). Arrows

point to actin concentration sites in the PE cluster. (**g**) epi:GFP crossed with the *kdrl:mCherry* line. Maximal

projection and optical sections of a representative frame from an *in vivo* time-lapse. GFP$^+$ pericardium and PE

are shown in green, the mCherry$^+$ endocardium in red. (**h**) *lnfg:GFP* crossed with the *kdrl:mCherry* line.

Maximal projection and optical section of a representative frame from an *in vivo* time-lapse. (**i**) Optical

sections through the avc and vp regions of *wt1b:Gal4*; *UAS:mCherry*; *lnfg:GFP* larvae at 60 hpf. PE cells

(red), *lnfg*:GFP (green), MHC$^+$ heart tube (white), cell nuclei (DAPI, blue). Zoomed views including single

channels are shown in the bottom panels. Arrowheads point to mCherry$^+$/GFP$^-$ PE cells. (**j**)

Tp1:CreERT2;ubb:switch embryos were treated with 4-Hydroxytamoxifen to trace the fate of Notch1

responsive cells with mCherry. At 80 hpf, mCherry$^+$ cells are observed in the endocardium. (**k,l**) Fluorescent

in situ mRNA hybridization for *notch1a* (k, green, n=3/3 embryos) and RNAscope detection of *hey2* (**l**)

followed by immunofluorescence for *fli1a:*GFP (red) and MHC (gray) on 60 hpf heart sections (2/2 embryos).

Nuclei are counterstained with DAPI (blue). at, atrium; avc, atrio-ventricular canal; DP, dorsal pericardium;

hpf; hours post fertilization; MHC, Myosin Heavy Chain; PE, proepicardium; v, ventricle; vp, venous pole.

Scale bar: 50 μm except for **i** middle images, 20 μm and zoomed images in **k** (10 μm) and **l** (25 μm). Data are

mean ± s.d., one-way ANOVA followed by Kruskal-Wallis significant difference test in **b** and Student's *t*-test

in **d**. *** $P < 0.001$.

1011

Fig. 8 Endothelial Notch signaling acts upstream of Bmp to control PE formation. (**a**) Maximal

projections and optical sections of larvae at 60 hpf overexpressing NICD in DP/PE cells (*wt1b:Gal4*) or in the

endothelium (*fli1a:Gal4*). Zoomed views are shown below. epi:GFP$^+$ cells are in green, MHC in red,

pSmad1/5 in white and nuclei were counterstained with DAPI (blue). Arrowheads mark PE cluster and

asterisks pSmad1/5$^+$ PE cells. (**b**) Quantification of pSmad 1/5$^+$ DP cell numbers. (**c**) Quantification of

pSmad1/5$^+$ PE cell numbers. (**d**) Maximal projections and optical sections of LDN193189 (LDN) or

RO4929097 (RO)-treated animals at 60 hpf in non-transgenic versus *bmp2b* or *NICD*-overexpressing animals

using *hsp70:Gal4*. DP and PE are in green, heart tube in red, pSmad1/5 in white. Nuclei were counterstained

with DAPI. Arrowheads indicate the PE. Asterisks point to pSmad1/5$^+$ DP cells. (**e**) Quantification of PE cell

number as shown in **d**. (**f**) Quantification of DP pSmad1/5$^+$ cell number in **d**. (**g**) Whole mount *in situ*

hybridization for *bmp4* in control, RO-treated and endothelial *NICD* overexpressing animals. White arrows

point to the venous pole, black arrows to the atrioventricular canal (avc) of the heart tube. (**h**) *in situ*

hybridization for *bmp4* on sections from control (2/3 embryos) and endothelial *NICD* overexpressing animals

through the heart. *bmp4* expression is enhanced upon *NICD* overexpression (2/3 embryos, arrowhead). (**i**)

Number of PE cells shown during each event of cell release per larvae observed from 55-64 hpf. (**j**) Number of

events of cell release. (**k**) Model of the interaction between Notch, BMP and the actomyosin network on PE

formation. A section of a 60 hpf larvae close to the venous pole was drawn. at, atrium; DP, dorsal pericardium;

hpf; hours post fertilization; HS, heat shock; MHC, Myosin Heavy Chain; PE, proepicardium; v, ventricle.

Scale bar: 50 μm except for zoomed images in **a** and **h** (20 μm). Data are mean ± s.d., one-way ANOVA

followed by Kruskal-Wallis significant difference test. *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$.

1032

1033

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

a **Bmp2b** b **BDM** c **BDM + Bmp2b**

ap
midline
vp
52 hpf

60 hpf

time

epi:GFP

d **control** **Bmp2b** **BDM** **BDM + Bmp2b**

60 hpf
myl7:mRFP epi:GFP

v at

+ 6h BDM

+ 6h BDM

divergence

constriction < −10⁻⁶ s⁻¹   0   > 10⁻⁶ s⁻¹ expansion

e

mean divergence

0.4
0.2
0
−0.2
−0.4

expansion
constriction

control Bmp2b BDM BDM + Bmp2b

** ** ** **

f

% of DP cells

towards midline
away from midline

control Bmp2b BDM BDM + Bmp2b

** *** *** *

Figure 7

Figure 8