Intraspecific variation and warming have comparable effects on eco-

evolutionary dynamics.

A. Raffard[1,2*], J. Cucherousset[3], F. Santoul[2], L. Di Gesu[1] and S. Blanchet[1,3*]

[1] CNRS, Université Toulouse III Paul Sabatier, Station d'Écologie Théorique et

Expérimentale du CNRS à Moulis, UMR-5321, F-09200 Moulis, France.

[2] EcoLab, Université de Toulouse, CNRS, INPT, UPS, Toulouse, France.

[3] Laboratoire Évolution and Diversité Biologique (EDB UMR 5174), Université de

Toulouse, CNRS, IRD, UPS, 118 route de Narbonne, Toulouse 31062, France.

Classification: *Biological Sciences / Ecology*

Abstract

Rapid evolutionary divergences within species can affect the way organisms shape

their environment, which in turn can affect the evolutionary trajectories of species.

These eco-evolutionary feedbacks have recently been proved, but their relevance

compared to that of key environmental drivers in the dynamics of biological systems

is still unknown. Here, we filled this gap in knowledge by quantifying the effects of

intraspecific variation on ecological processes and evolutionary trajectories and by

comparing these effects to those induced by a 2°C warming. We used a common

gardening experiment with simultaneous variations in the genetic and phenotypic

characters of a widespread freshwater fish species (the European minnow, *Phoxinus*

phoxinus) and in the ecosystem temperature. We showed that intraspecific variation

led to eco-evolutionary feedbacks that were as strong as the eco-evolutionary

consequences of ecosystem warming. Specifically, we found that variations in two

heritable phenotypic traits (body size and growth rate) in adult minnows led to

ecological changes in the environment that subsequently modulated the evolutionary

trajectories of juvenile minnows. Importantly, the eco-evolutionary consequences of

intraspecific variation were different and independent from those induced by

warming. We conclude that eco-evolutionary feedbacks are not biologically

negligible and that intraspecific variation is an indisputable driver of both ecological

and evolutionary dynamics.

Significance statement

Global changes are directly modifying the functioning of ecosystems and the

evolution of species. Phenotypic and genetic variation observed within species

(intraspecific variation) has recently also been shown to alter biological dynamics and

is threatened by global changes. It is hence of utmost importance to quantify the

relative importance of intraspecific variation and global changes (here considering

warming) on biological dynamics. Here, we demonstrate that intraspecific variation

affects ecosystem functioning as intensively as a warming of 2°C. The ecosystem

changes induced by intraspecific variation were strong enough to generate feedbacks

in the evolution of organisms, with a magnitude similar to that of warming.

Intraspecific variation should be a major target of conservation plans because it

affects biological dynamics as much as contemporary warming.

Introduction

Reciprocal interactions between ecological and evolutionary dynamics occurring over contemporary time scales (*eco-evolutionary dynamics*) have been increasingly studied in the last two decades[1–4]. Theory predicts that evolutionary diversification within a species can affect ecological processes such as primary productivity, resulting in environmental changes that can act as new selective pressures modulating the evolution of organisms[5–7]. These reciprocal interactions can generate eco-evolutionary feedbacks linking species evolution and ecosystem functioning, providing an integrative and temporally dynamic framework for understanding biological systems[8]. Although the study of eco-evolutionary feedback loops has long been conceptual[3,5,8], recent experimental studies have demonstrated the existence of these process[9–12]. However, the relative importance of eco-evolutionary feedbacks in the dynamics of natural ecosystems has been questioned[2,3].

To answer this question, it is important to determine whether eco-evolutionary feedbacks are negligible compared to key environmental drivers affecting both ecological and evolutionary dynamics, such as temperature, nutrient availability, predation or parasitism[2,13]. Recent investigations have revealed that intraspecific variation (emerging from evolutionary diversification) can affect ecological processes with an intensity similar to that of key environmental drivers[14–16]. However, whether the effects of intraspecific variation on evolutionary dynamics (mediated by the effects of intraspecific variation on ecological processes) are similar to or stronger than those of indisputable environmental drivers is still unknown. Addressing this question is fundamental to determining the relative contribution of eco-evolutionary feedbacks in driving the responses of biological systems to varying environmental conditions[13].

Here, we experimentally quantified the consequences of intraspecific variation

on ecological processes and subsequent evolutionary trajectories (i.e., eco-

evolutionary feedbacks) and then compared these consequences to the ecological and

evolutionary consequences of warming. Temperature is a key abiotic factor that

strongly varies at the landscape scale, directly affects key ecological functions such

primary productivity and ecosystem respiration[17–19], and imposes a strong selective

pressure on organism traits[20–22]. We ran a two-phase "common gardening experiment"

(*sensu*[8,11], Fig. 1) and manipulated (i) intraspecific variation in a freshwater fish

(European minnow, *Phoxinus phoxinus*) by selecting individuals from six

evolutionary and functionally distinct populations (i.e., differences in genotypes and

functional traits, see Methods and Fig. S1 and S2) and (ii) water temperature by

setting mesocosms varying by 2°C throughout the experiment (Fig. S3). An increase

in temperature of 2°C represents the general warming expectations for freshwaters

over the next 40 years[23]. During the first experimental phase (*ecological effects*, 10-

weeks), we compared the strengths of the effects of intraspecific variation among

adult minnows, to the strengths of the effects of warming on prey community

structure and ecosystem functions (Fig. 1). Adults were then removed from the

mesocosms and replaced by juveniles with a common origin for the second

experimental phase (*evolutionary effects*, 11-weeks). We tested how the ecological

variations induced during the first phase (due to intraspecific variation and/or

warming) affected the evolutionary trajectories (fitness and performance) of juveniles.

Results

In the first phase, we found that the effects of intraspecific variation in adult

minnows on ecological processes (measured over all ecological parameters) were at

least as strong as those of warming on ecological processes (mean effect size (MES) ± standard error = 0.103 ± 0.018 and MES ± SE = 0.078 ± 0.036 for intraspecific variation and warming, respectively; $t = 0.624$, d.f = 18, $p = 0.540$, Fig. 2). Nonetheless, the effects were heterogeneous across ecological parameters (Fig. 2b, Fig. S4). For instance, intraspecific variation had the strongest ecological effect on the abundance of Cladocera, whereas warming had a particularly strong ecological effect on decomposition rate (Fig. 2b, Fig. S5 and S6). A single interaction term between warming and intraspecific variation was significant (i.e., for benthic primary productivity, $F = 10.831$, d.f = 5,52, $p = 0.022$), indicating that the ecological effects of intraspecific variation were not temperature-dependent for most ecological parameters. The body mass and growth rate of minnows, two functionally important traits differing among minnow populations (although the later was also affected by experimental temperature, Fig. S2), were included in a path analysis testing the direct and indirect relationships among trait variation, warming and ecological parameters. We found that intraspecific variation in these two functional traits affected ecological processes as much as warming and that body mass was the most influential functional trait (Fig. 3). We further found that the intraspecific trait variation acted both directly and indirectly on ecological parameters (Fig. 3b). For instance, adult body mass affected the abundance of Copepoda directly, subsequently leading to an indirect effect on the abundance of Cladocera (Table 1, Fig. 3a). The ecological effects of warming were mainly direct (67%), although some indirect effects were also observed (Fig. 3b). For instance, warming directly increased Bivalvia abundance, positively affecting the abundance of Copepoda and the size of Cladocera, hence representing an indirect effect of warming on the zooplankton community (Fig. 3).

In the second phase (Fig. 1), we found that the strength of the effect sizes of

intraspecific variation and warming on the fitness proxies (survival, growth rate and

body condition) were similar (MES $\pm$ SE = 0.044 $\pm$ 0.004 and MES $\pm$ SE = 0.032 $\pm$

0.016 for intraspecific variation and warming, respectively, t = 0.665, d.f = 4, p =

0.542, Fig. 2). Notably, the average effect sizes of intraspecific variation and warming

on the evolutionary parameters were half the intensity of those on ecological

parameters (Fig. 2), indicating that evolutionary dynamics were less affected than

ecological processes by the initial treatments. Nonetheless, we observed eco-

evolutionary feedbacks since the juvenile growth rate was related to the ecological

parameters (benthic primary productivity, decomposition rate and Cladocera

abundance), which were controlled by intraspecific variation (Fig. 3a). Juvenile

survival was also related to juvenile growth rate (density-dependent growth rate) and

was indirectly related to warming. We also identified a direct relationship between

adult trait variation and juvenile growth rate (Fig. 3), which was unexpected given

that the adults were removed from the tanks before the juveniles were introduced.

This result indicates interspecific variation had unmeasured indirect effects (mediated

by ecological changes) on evolutionary dynamics. Juvenile survival was positively

related to the body size of Cladocera and negatively related to temperature (i.e.,

survival increased as temperature decreased). Juvenile body condition covaried with

both juvenile survival and growth rate and was lower when the abundance of

Bivalvia, which was directly affected by temperature, was high (Fig. 3).

Discussion

We found that the ecological consequences of intraspecific variation and

warming were similar in strength but acted on different ecological processes. Adult

minnows from evolutionary and phenotypically distinct populations modulated both

the abundance and the size of their prey (zooplankton Cladocera), probably because

these populations have different prey consumption and selectivity characters. In

contrast, warming strongly accelerated leaf litter decomposition, probably because

warmer temperatures stimulate bacterial activity[18,20]. The evolutionary effects of

intraspecific variation (resulting from eco-evolutionary feedbacks) and warming were

also similar in strength, regardless of the fitness traits investigated. For instance, the

survival of juvenile minnows was higher the low-temperature than the high-

temperature treatment (which is expected given that minnows inhabit relatively cold

rivers[24]), whereas the growth rate of juveniles differed depending on the adult

minnow population introduced at the onset of the experiment. Notably, the effects of

warming on the evolutionary trajectories of juveniles might be overestimated

compared to the effects of intraspecific variation, since the former represents the

cumulative results of both second-phase direct effects and first-phase indirect effects

mediated by ecological changes. Overall, our results demonstrate that intraspecific

variation can affect the ecological and evolutionary dynamics of biological systems as

much as warming does, although in different directions. Thus, eco-evolutionary

feedbacks occur in this type of biological system and significantly modulate the

whole biological dynamics of ecosystems.

Here, the eco-evolutionary feedback comprised indirect effects of intraspecific

variation among adult minnows on the evolutionary trajectory of juveniles, which

were mediated by the direct consequences of adult minnows on the ecological theatre.

Currently, very few studies have demonstrated the existence of eco-evolutionary

feedback, and most of have focused on model organisms[9–12]. By focusing on a non

model organism, our study extends the taxonomic scope of eco-evolutionary feedback

loops and suggests that this process does not concern only species with strong eco-

evolutionary divergences[8]. We further identified two heritable traits (body mass and

growth rate[25,26]) that partially initiate these eco-evolutionary feedbacks and vary

between adult minnows originating from environmentally and evolutionary distinct

populations. For instance, the zooplankton community was strongly impacted by trait

variability, probably through diet specialization. In turn, juveniles performed better

(i.e., higher growth rate) in the mesocosms with a higher abundance of zooplankton,

confirming that eco-evolutionary feedbacks could arise from a change in prey

availability[9]. Previous studies have identified growth rate and body mass as important

traits for ecological processes[27–29], and we here provide novel insights into the indirect

evolutionary consequences of these traits.

Interestingly, intraspecific variation and warming acted additively but not

interactively on ecological and evolutionary dynamics. Indeed, we identified only one

significant interaction between intraspecific variation and warming on benthic

primary productivity, indicating that the effect of intraspecific variation on benthic

primary productivity dynamics was temperature-dependent. This finding confirms

that the ecological consequences of intraspecific variation are often independent from

the abiotic context[15], which might also be the case for eco-evolutionary feedbacks.

This independence is surprising, since local adaptation for specific fitness traits and/or

for reaction norms often leads to strong context dependency in the responses of

organisms to local abiotic conditions[30,31], and we may have observed cascading

interactive effects of intraspecific variation on ecological and evolutionary

dynamics[32,33]. This finding is important because the absence of strong interactive

effects reduces biological complexities and may therefore improve our ability to

forecast the ecological and evolutionary consequences of environmental and

biodiversity changes[34].

In conclusion, we demonstrated for the first time that the magnitude of eco-

evolutionary feedbacks was as strong as the effects of warming on ecological and

evolutionary dynamics; thus, eco-evolutionary feedbacks are not biologically

negligible. Intraspecific variation in major heritable traits such as growth rate or body

mass is commonplace in the wild and can arise via various evolutionary processes

including natural selection and genetic drift[35]. Our study proves that intraspecific

variation is an indisputable driver of biological dynamics (at both the ecological and

evolutionary scale) that should not be considered noise in ecosystems. Current

environmental changes are rapid and can directly affect ecosystem functioning[18].

These changes can also directly modulate the distribution of intraspecific variation in

landscapes and thereby indirectly effect the eco-evolutionary dynamics of biological

systems[10,11]. These results reinforce recent reports that changes in intraspecific

variations of wild populations (e.g., harvest[36] or pollution[37]) could be as harmful as

considerable environmental changes (e.g., warming) to biological dynamics and that

this facet of biodiversity should therefore be conserved adequately[38,39].

Methods

Study species

European minnow (*Phoxinus phoxinus*) was used as the model species. *P. phoxinus* is

a small-bodied (maximum length: ~80 mm, mean generation time: ~2 years) cyprinid

fish species widely distributed in Western Europe. *P. phoxinus* lives in relatively cold

waters, mainly in streams and rivers but also in mountain lakes[24,39]. It is a generalist

species that feeds on small invertebrates, algae, zooplankton and small fish larvae[40–]

[42].

In September 2016, we collected adult minnows by electrofishing in six rivers

in southwestern France (Fig. S1). We selected populations that were isolated

geographically (minimal riparian distance among sites = 64 km, mean ± SD = 343 km

± 182) and had distinct environments (Fig. S7) to favor both genetic and phenotypic

divergences among populations. Accordingly, the mean genetic divergence among

populations was F_{st} = 0.162 (measured using 17 microsatellites, min-max = 0.043-

0.313), indicating a high evolutionary distinctiveness among the populations. The

body mass (a highly heritable[25,26] and important functional trait[43,44], Fig. S2) of the

sampled populations also varied, as did two other important functional traits[20,45–47]:

the metabolic (min-max = 0.1388-0.2737 mg O^2.g^{-1}.h^{-1}, F = 14.599, d.f = 5,188, p <

0.001) and ammonium excretion (min-max = 17.02-43.48 μg NH^{4+}.g^{-1}.L^{-1}.h^{-1}, F =

4.695, d.f = 5,175, p < 0.001, unpublished data) rates. All fish collections and

husbandry for adults and juveniles were conducted in accordance with sampling

permits obtained from local authorities (25-08-2016, 24-05-2016, 09-273, SA-013-

PB-092, A09-3). Fish from different populations were reared separately for ~6

months in 1100 L outdoor tanks to minimize previous environmental effects on

phenotypes. During rearing, the fish were fed with a mixture of pelletized food and

dead chironomids until the start of the experiment.

Phase 1: effects of intraspecific variation and temperature on ecological processes

The experiment consisted of 72 replicated mesocosms placed in a greenhouse with a

12:12 h light-dark photoperiod. Mesocosms were filled with 100 L of tap water and 1

cm of gravel covering the bottom of each tank. Tanks were covered with a 1 cm

plastic mesh net to prevent fish escapes. Nutrients were added to the mesocosms

using 5 mL of solution containing nitrogen and phosphorus (ratio N: P: K = 3.3: 1.1:

5.8) on December 2^{nd} 2016. Each mesocosm was then inoculated with 200 mL of a

concentrated solution of phytoplankton from a unique lake origin (Lake Lamartine,

France 43°30'21.5"N, 1°20'32.7"E) on December 12^{th} 2016. Two months later

(February 15^{th} 2017), an additional 200 mL of concentrated solution of zooplankton

from the same lake was added to each mesocosm. Finally, we inoculated each

mesocosm with sediment and macroinvertebrates (i.e., mainly Gastropoda and

Bivalvia) from Lake Lamartine.

Each tank was assigned to one of twelve treatments according to a full-

factorial design with intraspecific variation (i.e., population origin, six levels

corresponding to each population) and temperature (two levels: low and high

temperature) as the main factors (Fig. 1). Each treatment was replicated six times.

Water temperature was controlled and adjusted using a *Blue Marine®* water chiller

and a stainless steel coil placed in each tank through which a flux of water

(independent from the water of the tanks) flowed at either 18°C or 21°C. Natural

seasonal temperature variations occurred; on average, the low and high water

temperature treatments differed by 2.08°C according to seasonal variations (Fig. S3).

In March 2017, adult fish were weighed to the nearest 0.01 g and a single fish

was introduced to each mesocosm. This individual-based approach prevented the

experimental ecosystems from collapsing due to the over-density of top consumers

and allowed the ecological effects of individual phenotypes to be measured. After 73

days (Fig. 1), each fish was removed, weighed and euthanized in a solution of

benzocaine at 25 mg.L^{-1}. The growth rate (%.day^{-1}) of the adults was calculated

as $SGR = \frac{ln(Wf) - ln(Wi)}{T} * 100$, where *Wf* and *Wi* are the final and initial body masses,

respectively, and *T* is time interval between two measurements (in days).

Concomitantly, we measured multiple community and ecosystem parameters to

evaluate differences in ecological processes among treatments.

(*i*) Pelagic algae stock was assessed as a proxy of pelagic primary

productivity. Measurements were performed using a portable spectrometer

(AlgaeTorch, bbe Moldaenke®) to assess the chlorophyll-a concentration (μg/L) in

the water column. Two measurements were taken in each mesocosm and were

averaged for the analyses.

(*ii*) Benthic algae stock was assessed as a proxy of the benthic primary

productivity using a portable spectrometer (BenthoTorch, bbe Moldaenke®). The

chlorophyll-a concentration (μg/cm^2) was measured on two tiles (20 x 20 cm) placed

in the mesocosms the day before the start of the experiment. These measurements

were averaged for analyses.

(*iii*) The abundance of filamentous algae was quantified. Filamentous algae

cover (%) was visually estimated by two operators, and values were averaged for

analyses.

(*iv*) Zooplankton community was assessed by filtering 5 L of water through a

200 μm sieve. Samples was conserved in a 70% ethanol solution and subsequently

identified to the order or family levels, including Copepoda (i.e., Cyclopoida and

Calanoida) and Cladocera (i.e., Daphniidae, Chydoridae and Bosminidae).

Zooplankton size was assessed by measuring 10 individuals of each order and family

level from each mesocosm to the nearest 0.001 mm using ImageJ®.

(*v*) Decomposition rate was measured by quantifying the mass loss of black

poplar (*Populus nigra,* a dominant riparian tree in southern France) abscised leaves[48].

One day before the start of the experiment, 4 g of air-dried leaves were put in each

mesocosm within a coarse mesh (1 x 1 cm) bag. At the end of the phase 1, the

remaining leaf material was removed from the mesocosms, rinsed with tap water,

oven dried at 60°C for three days and weighed to the nearest 0.001 g to assess the loss

of biomass. The decomposition rate was calculated as $k = -\frac{ln\,(X)}{t}$ [48], where X is the

proportion of litter remaining after phase 1 and t is the elapsed time in days.

(*vi*) Macroinvertebrates (> 1 mm, essentially molluscs) were collected from

the mesh bags used to measure decomposition rates, conserved in a 70% ethanol

solution, and identified as Bivalvia or Gastropoda.

(*vii*) Abiotic parameters of the water [pH, specific conductance (μS), oxygen

concentration (mg.L^{-1}) and turbidity (NTU)] were measured with a multiparameters

probe (YSI Pro DSS Water Quality Meter®). We summarized these parameters using

principal component axis (PCA) (package ade4 in R[49]). We selected the first axis of

the PCA as the synthetic variable. This axis explained 60% of the variance and was

correlated to the oxygen concentration (loading component: -0.95), pH (-0.93),

specific conductance (0.70) and, to a lesser extent, turbidity (0.25).

Phase 2: effect of ecological differences on juvenile evolution

After the removal of adult fish on June 13th 2017, 45 juvenile minnows were

introduced to each mesocosm. We used juveniles from a unique origin (i.e., fish farm,

Amorvif EURL) to control for potential genetic effects. Juveniles were introduced as

soon as possible after hatching to increase the possibility of differential mortality

and/or ontogenetic plasticity. Therefore, juveniles were introduced when they were

only two weeks old as stage III larvae[50] (Fig. S8). They were not manipulated (i.e.,

weighted and/or measured) before being randomly introduced in the mesocosms to

limit potential mortality. The juveniles were removed from the mesocosms 79 days

later, and we measured several proxies for their fitness. Individuals were counted to

assess survival, weighed to the nearest 0.001 g to assess growth rate (assuming all

juveniles had the same initial body mass, we used the final body mass of juveniles as

a measure of growth rate), and measured in length to the nearest 0.1 mm (using

ImageJ) to assess the body condition, which was calculated as the residuals of the

relationship between individual body mass and length.

Statistical analyses

Two adult individuals died before the end of phase 1, so we discarded these two

replicates from the analyses. Moreover, we identified six tanks in which crayfish had

been inadvertently introduced; we discarded these six replicates because crayfish are

known to have disproportionally strong impacts on ecosystems[48]. As such, the final

analyses were run on 64 replicates.

First, we compared the magnitude of the effects of intraspecific variation and

temperature on ecological (phase 1) and evolutionary (phase 2) dynamics. To do so,

we used a meta-analytic approach consisting of first running linear models linking

each ecological or evolutionary parameter (dependent variables) to the explicative

variables, i.e., intraspecific variation (categorical factor, six levels), temperature

(categorical factor, two levels) and the resulting two-term interaction. The interaction

term was removed when nonsignificant because it prevents the interpretation of

simple terms[51]. From these linear models, we calculated the standardized effect sizes

eta squared[52] (η^2) as follows: $\eta^2 = SS_x/SS_{tot}$, where SS_x is the sum of squares for the

effect of interest (intraspecific variation, temperature or the interaction term, if

significant) and SS_{tot} is the total sum of squares. Sums of squares were extracted from

type II analysis of variance when the interaction was not in the model and from type

III analysis of variance when the interaction was significant[53]. Finally, the MES

values of intraspecific variation and temperature across the ecological or evolutionary

parameters were compared using t-test.

Next, we assessed the direct and indirect links between intraspecific variation,

temperature, and the ecological and evolutionary parameters using a causal analysis.

Since we aimed at identifying the mechanisms by which the mesocosms diverged, we

included the body mass and the growth rate of adult fish from phase 1 because these

two traits are known to drive ecological processes[43]. We used path analyses[54–56] to set

a full model based on biologically rational paths and the visual inspection of the

variance-covariance matrix, and all variables were scaled to the mean to facilitate the

comparison. This full model was then simplified by removing sequentially weak

and/or nonsignificant paths until reaching a model that was correct statistically (i.e., a

model that best fit the observed covariance matrix based on the maximum likelihood

χ^2 statistic[55]), while leading to the lowest Akaike Information Criteria (AIC) value.

We finally extracted the absolute values of path coefficients from the final model to

tease apart the direct and indirect effects of body mass, growth rate and temperature

on the ecological and evolutionary parameters. Statistical analyses were performed

using R software[57], and path analyses were run using Amos[58].

References

1. Hendry, A. P. *Eco-Evolutionary Dynamics*. (2016).
2. Schoener, T. W. The Newest Synthesis: Understanding the Interplay of
Evolutionary and Ecological Dynamics. *Science* **331,** 426–429 (2011).
3. Thompson, J. N. Rapid evolution as an ecological process. *Trends Ecol. Evol.* **13,**
329–332 (1998).
4. Whitham, T. G. *et al.* Community and ecosystem genetics: a consequence of the
extended phenotype. *Ecology* **84,** 559–573 (2003).

5. Kokko, H. & López-Sepulcre, A. The ecogenetic link between demography and
evolution: can we bridge the gap between theory and data? *Ecol. Lett.* **10,**
773–782 (2007).

6. Post, D. M. & Palkovacs, E. P. Eco-evolutionary feedbacks in community and
ecosystem ecology: interactions between the ecological theatre and the
evolutionary play. *Philos. Trans. R. Soc. B-Biol. Sci.* **364,** 1629–1640 (2009).

7. Odling-Smee, J., Erwin, D. H., Palkovacs, E. P., Feldman, M. W. & Laland, K. N.
Niche Construction Theory: A Practical Guide for Ecologists. *Q. Rev. Biol.*
88, 3–28 (2013).

8. Matthews, B. *et al.* Toward an integration of evolutionary biology and ecosystem
science: Integration of evolutionary biology and ecosystem science. *Ecol. Lett.*
14, 690–701 (2011).

9. Best, R. J. *et al.* Transgenerational selection driven by divergent ecological
impacts of hybridizing lineages. *Nat. Ecol. Evol.* **1,** 1757–1765 (2017).

10. Brunner, F. S., Anaya-Rojas, J. M., Matthews, B. & Eizaguirre, C. Experimental
evidence that parasites drive eco-evolutionary feedbacks. *Proc. Natl. Acad.*
Sci. **114,** 3678–3683 (2017).

11. Matthews, B., Aebischer, T., Sullam, K. E., Lundsgaard-Hansen, B. & Seehausen,
O. Experimental Evidence of an Eco-evolutionary Feedback during Adaptive
Divergence. *Curr. Biol.* **26,** 483–489 (2016).

12. Turcotte, M. M., Reznick, D. N. & Daniel Hare, J. Experimental Test of an Eco-
Evolutionary Dynamic Feedback Loop between Evolution and Population
Density in the Green Peach Aphid. *Am. Nat.* **181,** S46–S57 (2013).

13. Rudman, S. M. *et al.* What genomic data can reveal about eco-evolutionary
dynamics. *Nat. Ecol. Evol.* **2,** 9–15 (2018).

14. Burkle, L. A., Souza, L., Genung, M. A. & Crutsinger, G. M. Plant genotype,
nutrients, and G X E interactions structure floral visitor communities.
Ecosphere **4,** UNSP 113 (2013).

15. El-Sabaawi, R. W. *et al.* Intraspecific phenotypic differences in fish affect
ecosystem processes as much as bottom-up factors. *Oikos* **124,** 1181–1191
(2015).

16. Des Roches, S. *et al.* The ecological importance of intraspecific variation. *Nat.*
Ecol. Evol. **2,** 57–64 (2018).

17. Yvon-Durocher, G., Jones, J. I., Trimmer, M., Woodward, G. & Montoya, J. M.
Warming alters the metabolic balance of ecosystems. *Philos. Trans. R. Soc. B*
Biol. Sci. **365,** 2117–2126 (2010).

18. Yvon-Durocher, G. *et al.* Five Years of Experimental Warming Increases the
Biodiversity and Productivity of Phytoplankton. *PLoS Biol* **13,** e1002324
(2015).

19. Yvon-Durocher, G. & Allen, A. P. Linking community size structure and
ecosystem functioning using metabolic theory. *Philos. Trans. R. Soc. B Biol.*
Sci. **367,** 2998–3007 (2012).

20. Brown, J. H., Gillooly, J. F., Allen, A. P., Savage, V. M. & West, G. B. Toward a
metabolic theory of ecology. *Ecology* **85,** 1771–1789 (2004).

21. Parmesan, C. Ecological and Evolutionary Responses to Recent Climate Change.
Annu. Rev. Ecol. Evol. Syst. **37,** 637–669 (2006).

22. Rey, O., Danchin, E., Mirouze, M., Loot, C. & Blanchet, S. Adaptation to Global
Change: A Transposable Element–Epigenetics Perspective. *Trends Ecol. Evol.*
31, 514–526 (2016).

23. IPCC. Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change (eds Core Writing Team, Pachauri, R.K., Meyer, L.A.). (2014).

24. Keith, P., Persat, H., Feunteun, E. & Allardi, J. *Les poissons d'eau douce de France.* (2011).

25. Heath, D. D., Fox, C. W. & Heath, J. W. Maternal effects on offspring size: variation through early development of chinook salmon. *Evolution* **53,** 1605–1611 (1999).

26. Thériault, V., Garant, D., Bernatchez, L. & Dodson, J. J. Heritability of life-history tactics and genetic correlation with body size in a natural population of brook charr (*Salvelinus fontinalis*). *J. Evol. Biol.* **20,** 2266–2277 (2007).

27. Rudolf, V. H. W. & Rasmussen, N. L. Population structure determines functional differences among species and ecosystem processes. *Nat. Commun.* **4,** (2013).

28. Rudolf, V. H. & Rasmussen, N. L. Ontogenetic functional diversity: Size structure of a keystone predator drives functioning of a complex ecosystem. *Ecology* **94,** 1046–1056 (2013).

29. Woodward, G. *et al.* Body size in ecological networks. *Trends Ecol. Evol.* **20,** 402–409 (2005).

30. Kawecki, T. J. & Ebert, D. Conceptual issues in local adaptation. *Ecol. Lett.* **7,** 1225–1241 (2004).

31. Blanquart, F., Kaltz, O., Nuismer, S. L. & Gandon, S. A practical guide to measuring local adaptation. *Ecol. Lett.* **16,** 1195–1205 (2013).

32. Gillooly, J. F. Effects of Size and Temperature on Metabolic Rate. *Science* **293,** 2248–2251 (2001).

33. Rosenblatt, A. E. & Schmitz, O. J. Climate Change, Nutrition, and Bottom-Up and Top-Down Food Web Processes. *Trends Ecol. Evol.* (2016). doi:10.1016/j.tree.2016.09.009

34. Beckage, B., Gross, L. J. & Kauffman, S. The limits to prediction in ecological systems. *Ecosphere* **2,** art125 (2011).

35. Lowe, W. H., Kovach, R. P. & Allendorf, F. W. Population Genetics and Demography Unite Ecology and Evolution. *Trends Ecol. Evol.* **32,** 141–152 (2017).

36. Palkovacs, E. P., Moritsch, M. M., Contolini, G. M. & Pelletier, F. Ecology of harvest-driven trait changes and implications for ecosystem management. *Front. Ecol. Environ.* **16,** 20–28 (2018).

37. Brodin, T., Fick, J., Jonsson, M. & Klaminder, J. Dilute Concentrations of a Psychiatric Drug Alter Behavior of Fish from Natural Populations. *Science* **339,** 814–815 (2013).

38. Paz-Vinas, I. *et al.* Systematic conservation planning for intraspecific genetic diversity. (2018). doi:10.1101/105544

39. Mimura, M. *et al.* Understanding and monitoring the consequences of human impacts on intraspecific variation. *Evol. Appl.* **10,** 121–139 (2016).

40. Collin, H. & Fumagalli, L. Evidence for morphological and adaptive genetic divergence between lake and stream habitats in European minnows (Phoxinus phoxinus, Cyprinidae). *Mol. Ecol.* **20,** 4490–4502 (2011).

41. Frost, W. E. The Natural History of the Minnow, Phoxinus phoxinus. *J. Anim. Ecol.* **12,** 139 (1943).

42. Miró, A., Sabás, I. & Ventura, M. Large negative effect of non-native trout and minnows on Pyrenean lake amphibians. *Biol. Conserv.* **218,** 144–153 (2018).

43. Rudolf, V. H. W., Rasmussen, N. L., Dibble, C. J. & Van Allen, B. G. Resolving
the roles of body size and species identity in driving functional diversity.
Proc. R. Soc. B Biol. Sci. **281,** 20133203 (2014).
44. Raffard, A. *et al.* The functional syndrome: linking individual trait variability to
ecosystem functioning. *Proc. R. Soc. B Biol. Sci.* **284,** 20171893 (2017).
45. Bassar, R. D. *et al.* Local adaptation in Trinidadian guppies alters ecosystem
processes. *Proc. Natl. Acad. Sci.* **107,** 3616–3621 (2010).
46. McIntyre, P. B. *et al.* Fish distributions and nutrient cycling in streams: can fish
create biogeochemical hotspots. *Ecology* **89,** 2335–2346 (2008).
47. Hildrew, A. G., Raffaelli, D. & Edmonds-Browns, R. *Body Size: the Structure*
and Function of Aquatic Ecosystems. (2007).
48. Alp, M., Cucherousset, J., Buoro, M. & Lecerf, A. Phenological response of a key
ecosystem function to biological invasion. *Ecol. Lett.* **19,** 519–527 (2016).
49. Chessel, D., Dufour, A. B. & Dray, S. ade4: Analysis of Ecological Data:
Exploratory and Euclidean methods in Multivariate data analysis and
graphical display. R package version. 1–4 (2007).
50. Pinder, A. *Keys to larval and juvenile stages of coarse fishes from fresh waters in*
the British isles. (2001).
51. Nakagawa, S. & Cuthill, I. C. Effect size, confidence interval and statistical
significance: a practical guide for biologists. *Biol. Rev.* **82,** 591–605 (2007).
52. Levine, T. & Hullett, C. Eta squared, partial eta squared, and misreporting of
effect size in communication research. *Hum. Commun. Res.* **28,** 612–625
(2002).
53. Langsrud, O. ANOVA for unbalanced data: Use Type II instead of Type III sums
of squares. 5
54. Duffy, J. E., Lefcheck, J. S., Stuart-Smith, R. D., Navarrete, S. A. & Edgar, G. J.
Biodiversity enhances reef fish biomass and resistance to climate change.
Proc. Natl. Acad. Sci. **113,** 6230–6235 (2016).
55. Grace, J. B. *Structural equation modeling and natural systems.* (2006).
56. Grace, J. B. *et al.* Integrative modelling reveals mechanisms linking productivity
and plant species richness. *Nature* **529,** 390–393 (2016).
57. R Core Team. R: A Language and Environment for Statistical Computing. (2013).
58. Arbuckle, J. L. Amos (Version 23.0) [Computer Program]. (2014).
506

Acknowledgements

We warmly thank Jose M. Montoya, Delphine Legrand and Jérôme G. Prunier for

their valuable comments on a previous version of the manuscript. We thank Lucas

Mignien, Kéoni Saint-Pe and Yoann Buoro for their help during the experimental

work. AR was financially supported by a doctoral scholarship from the Université

Fédérale de Toulouse. This work was undertaken at SETE and EDB, which is part of

the "Laboratoire d'Excellence" (LABEX) entitled TULIP (ANR-10-LABX-41).

519 **Table 1:** Causal pathways between variations in intraspecific traits (body mass$_{adults}$
520 and growth rate$_{adults}$), temperature and ecological and evolutionary parameters
521 obtained from path analyses.
522

Response	Effect	Path coefficient	S.E.	p-value
Growth rate$_{adults}$	Body mass$_{adults}$	-0.674	0.093	< 0.001
Cladocera abundance	Growth rate$_{adults}$	-0.341	0.111	0.002
	Copepod abundance	0.339	0.11	0.002
Size Cladocera	Bivalve abundance	0.428	0.114	< 0.001
Copepoda abundance	Growth rate$_{adults}$	0.37	0.153	0.016
	Bivalve abundance	0.311	0.113	0.006
	Body mass$_{adults}$	0.477	0.153	0.002
Bivalvia abundance	Temperature	0.441	0.113	< 0.001
Abiotic parameters	Body mass$_{adults}$	-0.289	0.124	0.019
	Filamentous algae	-0.644	0.092	< 0.001
	Growth rate$_{adults}$	-0.194	0.124	0.116
Pelagic prod.	Prod. benthic	0.244	0.12	0.043
Benthic prod.	Growth rate$_{adults}$	-0.522	0.139	< 0.001
	Bivalvia abundance	-0.228	0.118	0.054
	Body mass$_{adults}$	-0.486	0.139	< 0.001
	Temperature	0.292	0.132	< 0.001
	Filamentous algae	0.239	0.12	0.047
Decomposition rate	Body mass$_{adults}$	0.375	0.132	< 0.001
	Temperature	0.532	0.096	< 0.001
Filamentous algae	Temperature	0.554	0.118	< 0.001
	Bivalvia abundance	-0.207	0.123	0.092
	Copepoda abundance	-0.168	0.110	0.126
Survival$_{juveniles}$	Size Cladocera	-0.195	0.123	0.112
	Temperature	-0.183	0.123	0.135

Growth rate$_{juveniles}$	Body mass$_{adults}$	0.479	0.130	< 0.001
	Growth rate$_{adults}$	0.725	0.125	< 0.001
	Prod. benthic	0.332	0.092	< 0.001
	Cladocera abundance	0.250	0.089	0.005
	Survival$_{juveniles}$	-0.566	0.084	< 0.001
	Decomposition rate	-0.234	0.093	0.012
Body condition$_{juveniles}$	Bivalve abundance	-0.175	0.073	0.016
	Survival$_{juveniles}$	0.556	0.087	< 0.001
	Growth rate$_{juveniles}$	0.941	0.083	< 0.001

Fig. 1. Experimental design used to test the ecological (phase 1) and evolutionary

(phase 2) effects of intraspecific variation among adult minnows (*Phoxinus phoxinus*)

and warming.

Fig. 2. Comparison of the ecological and evolutionary effects of intraspecific

variation and warming. (a) Eta squared (η^2) effect size of the intraspecific variation

among adults and temperature on ecological (green) and evolutionary (blue)

dynamics. Error bars represent $\pm$ 1 SE. **(b)** Delta of effect sizes (η^2) of intraspecific

variation and temperature on ecological and evolutionary parameters. Positive values

indicate a higher effect of intraspecific variation, and negative values indicate a higher

effect of temperature.

Fig. 3. Effects of intraspecific trait variation and warming on ecological and

evolutionary dynamics. (a) Representation of the simplified path analysis linking

intraspecific trait variation among adult minnows, temperature, and ecological and

evolutionary dynamics. The covariance structure of this simplified path model, which

contains both ecological and evolutionary dynamics, did not differ from that of the

data (χ^2 = 65.373, d.f = 72, p = 0.696), indicating that the data were well supported by

the model. The arrow line width is proportional to the β path coefficients (brown and

pink arrows represent positive and negative values, respectively), and the dotted lines

represent indirect effects of intraspecific variation on evolutionary parameters. **(b)**

Cumulated absolute β path coefficients extracted from the simplified path analysis

depicting the direct (filled) and indirect (hatched) relationships between intraspecific

trait variation among adult minnows, temperature, ecological dynamics (green) and

evolutionary dynamics (blue). Note that the direct effects of intraspecific variation on

evolutionary dynamics are not true direct effects (because the adult fish were removed

before phase 2 of the experiment) but instead reveal unmeasured causal relationships.

 # = abundance.

Fig. 1.

Fig. 2.

Fig. 3.