**Meta-analysis of Reward Processing in Major Depressive Disorder Reveals
Distinct Abnormalities within the Reward Circuit**

Tommy H. Ng, MPhil, Lauren B. Alloy, PhD, & David V. Smith, PhD
Department of Psychology, Temple University, Philadelphia, PA, USA

Running title: Reward Processing in Depression

Corresponding Author:

David V. Smith, Ph.D.
Assistant Professor of Psychology
Temple University
Weiss Hall, Room 825
1701 North 13th Street
Philadelphia, PA 19122
Office Phone: 215-204-1552
Fax Number: 215-504-5539
Email: david.v.smith@temple.edu

Abstract

Many neuroimaging studies have investigated reward processing dysfunction in major depressive disorder. These studies have led to the common idea that major depressive disorder is associated with blunted responses within the reward circuit, particularly in the ventral striatum. Yet, the link between major depressive disorder and reward-related responses in other regions remains inconclusive, thus limiting our understanding of the pathophysiology of major depressive disorder. To address this issue, we performed a coordinate-based meta-analysis of 41 whole-brain neuroimaging studies encompassing reward-related responses from a total of 794 patients with major depressive disorder and 803 healthy controls. Our findings argue against the common idea that major depressive disorder is primarily linked to deficits within the reward system. Instead, our results demonstrate that major depressive disorder is associated with opposing abnormalities in the reward circuit: hypo-responses in the ventral striatum and hyper-responses in the orbitofrontal cortex. The current findings suggest that dysregulated corticostriatal connectivity may underlie reward-processing abnormalities in major depressive disorder, providing an empirical foundation for a more refined understanding of abnormalities in the reward circuitry in major depressive disorder.

Keywords: coordinate-based meta-analysis, reward, major depressive disorder, neuroimaging

Introduction

Depression is a prevalent mental disorder ranked as the leading non-fatal cause of disability by the World Health Organization.[1] Therefore, it is of paramount importance to understand its underlying neurobiological mechanisms. Over the past decade, theorists have proposed that anhedonia, one of the core symptoms of depression, is linked to reward processing dysfunction.[2–9] In particular, many neuroimaging studies have reported that relative to healthy controls (HC), individuals with major depressive disorder (MDD) exhibit reduced activity in the ventral striatum (VS) in response to reward.[10–14]

The striatum, which can be divided into dorsal and ventral sections, is the primary input zone for basal ganglia.[16,17] It receives afferent projections from the midbrain, amygdala, and prefrontal cortex (PFC), such as the orbitofrontal cortex (OFC), dorsolateral prefrontal cortex (dlPFC), ventromedial prefrontal cortex (vmPFC), and anterior cingulate cortex (ACC).[16,17] It also projects to such regions as the ventral pallidum, ventral tegmental area, and substantia nigra.[17] Many of the regions linked to the striatum, particularly prefrontal regions, have been associated with the computation and representation of reward value,[18–24] as well as the regulation of affect and reward-related behavior in animals and healthy individuals.[25–29] The striatum also has been proposed to play an important role in the onset and course of MDD, with longitudinal studies demonstrating that blunted activation in the VS during reward anticipation predicts the emergence of depressive symptoms and disorder[30,31] and deep-brain stimulation studies using it as a treatment target for treatment-resistant depression.[32]

Although blunted striatal response to reward in MDD is a well-established finding in the literature,[9,33–36] it is less clear how other regions, particularly the PFC, also may contribute to reward processing deficits in MDD. For instance, some studies have found that relative to HC, MDD exhibited greater activation in the OFC,[13,37] dlPFC,[12,38,39] vmPFC,[40,41]

ACC,[42,43] middle frontal gyrus,[41,43] inferior frontal gyrus,[42,44] subgenual cingulate,[40,44] and

dorsomedial prefrontal cortex[41] during the processing of rewarding stimuli. In contrast, other

studies have reported less activity in MDD in response to reward in the OFC,[37,43]

ACC,[12,13,37,44] middle frontal gyrus,[13,42,44] and frontal pole.[43] The inconsistencies may be due

to a number of factors, such as limited statistical power[45–47] and susceptibility artifacts in the

ventral portions of the PFC.[25,48,49] Therefore, the association between prefrontal regions and

MDD remains equivocal, both in terms of the *direction* (i.e., hyper- or hypo-responses) and

the *location* (e.g., OFC, dlPFC, vmPFC and/or ACC) of the effect.

Inconsistencies in the literature have prompted researchers to conduct coordinate-

based meta-analyses to identify common activation patterns implicated in MDD during

reward processing. Although prior meta-analytic efforts[34,35,50] have shown some overlapping

findings in the striatum in MDD, there is a striking degree of anatomical disagreement across

these efforts, with non-overlapping findings all throughout the brain (see Supplementary

Table 1 and Supplementary Figure 1 for a complete comparison of findings across meta-

analyses). The lack of agreement across meta-analyses can be due to methodological issues,

such as lenient thresholding, overlapping samples, software issues,[51] different inclusion

criteria, and inclusion of region-of-interest (ROI) coordinates, as detailed in a previous

review.[52] For example, two previous meta-analyses[34,35] corrected for multiple comparisons

using the false discovery rate (FDR) approach, which has been shown to be inadequate in

controlling the false positives among clusters in neuroimaging meta-analyses[53,54] and might

have contributed to the lack of agreement across studies.

To address these issues and extend extant work, we performed a coordinate-based

meta-analysis following procedures recommended by new guidelines.[52,55] The current work

differed from previous meta-analyses on reward processing in MDD in various aspects, such

as only including whole-brain studies to avoid localization bias; only including studies that

used an active control condition to isolate reward-related processes; only including

independent samples to avoid double counting the same participants; using more stringent

thresholding criteria; having the most up-to-date literature search; and only conducting a

meta-analysis when there were at least 17 eligible experiments to restrict excessive

contribution of any particular studies to cluster-level thresholding.[56]

Our primary hypothesis was that the literature would consistently show that compared

with HC, individuals with MDD would exhibit blunted activation of the striatum and

abnormal activation of the prefrontal regions (e.g., the OFC) during the processing of

rewarding stimuli. We also explored whether there were consistent neural responses to

punishing stimuli in MDD relative to HC. To examine these questions, we conducted four

separate coordinate-based meta-analyses testing spatial convergence of neuroimaging

findings for the following four contrasts: 1) positive valence (reward > punishment or neutral

stimuli; neutral stimuli > punishment) for MDD > HC; 2) negative valence (punishment >

reward or neutral stimuli; neutral stimuli > reward) for MDD > HC; 3) positive valence for

HC > MDD; and 4) negative valence for HC > MDD. Neutral stimuli were considered

rewarding relative to punishing stimuli but punishing relative to rewarding stimuli, in view of

the idea that brain activation during reward processing reflects the valence of a stimulus and

follows a linear pattern.[57,58]

The comprehensive and rigorous nature of the current meta-analysis allowed us to

investigate whether a quantitative synthesis of neuroimaging studies on reward-related

processing in MDD would unveil common activation patterns that may be difficult to discern

by individual studies due to inconsistent findings. We aimed to address two main questions.

First, which brain regions show consistent hypo-responses to reward-relevant stimuli in MDD

relative to HC? Second, which brain regions show consistent hyper-responses to reward-

relevant stimuli in MDD relative to HC?

Materials and Methods

Study Selection

The current coordinate-based meta-analysis primarily followed the guidelines for meta-analyses, whenever applicable.[55,59] We conducted a systematic literature search to identify neuroimaging studies on reward processing abnormalities in mood disorders. Potentially eligible studies published between 1/1/1997 and 8/7/2018 were identified by searching the MEDLINE, EMBASE, PsycINFO, PsycARTICLES, Scopus, and Web of Science using the grouped terms (fMRI* or PET*) AND (depress* OR bipolar* OR mania* OR manic* OR hypomania* OR hypomanic*) AND (reward* OR effort* OR decision* OR reinforce* OR habit* OR discounting* OR "prediction error" OR "delayed gratification" OR "approach motivation" OR "positive valence systems"). To enhance search sensitivity, the reference lists of the retrieved articles and review papers were further checked to identify potentially relevant articles. Although our initial goal was to investigate reward processing dysfunction in both MDD and bipolar disorder, the current meta-analysis only focused on MDD due to an inadequate number of studies on bipolar disorder (the search identified 23 studies on bipolar disorder across positive and negative valence contrasts, yielding fewer than 17 experiments for each targeted meta-analysis).

Inclusion Criteria

We included studies that (a) used a reward and/or punishment task, (b) reported comparisons between people with MDD and HC, (c) used standardized diagnostic criteria (e.g., *DSM-IV, DSM-IV-TR*, ICD-10) to determine psychiatric diagnoses, (d) used fMRI or PET in conjunction with parametric analysis or subtraction methodology contrasting an experimental condition and an active control condition (e.g., a punishment condition, a lower-intensity reward condition, or a neutral condition) to isolate reward-related processes and identify foci of task-related neural changes, (e) reported significant results of whole-brain

group analyses without small volume corrections (SVC), as non-whole-brain coordinates (e.g., ROI-based coordinates) and analyses involving SVC have been argued to bias coordinate-based meta-analyses,[55,56] (f) reported coordinates in a standard stereotactic space [Talairach or Montreal Neurological Institute (MNI) space], and (g) used independent samples. Authors were contacted if the required information were unavailable in the published reports.

The study with the largest sample size was included if there was sample overlap between studies. Reward tasks were operationalized as involving presentation of a rewarding stimulus (e.g., winning money, favorite music, positive faces), whereas punishment tasks were operationalized as involving presentation of a punishing stimulus (e.g., losing money, negative faces). The stimuli used in the included studies of the meta-analysis reflect a reward-punishment or positive-negative continuum. For example, positive faces are considered as rewards based on previous research showing that positive faces activate the reward circuitry, that they are discounted as a function of time, that they are tradable for other rewards (e.g., money), that they reinforce work, and that people are willing to work to view positive faces and exert more effort for more positive faces.[60,61]

Coordinate-Based Meta-Analysis

Given the inconsistency of findings in the literature of reward processing abnormalities in MDD, we used a coordinate-based meta-analytic approach and activation likelihood estimation[54,62] to examine whether we could identify consistent activation patterns across studies. Our main analyses focused on which brain regions show consistent hypo- or hyper-responses to reward-relevant stimuli in MDD relative to HC. Coordinate-based meta-analyses were performed using GingerALE 2.3.6 (http:// brainmap.org), which employs the activation likelihood estimation (ALE) method.[54,63] The ALE method aims to identify regions showing spatial convergence between experiments and tests against the null hypothesis that

the foci of experiments are uniformly and randomly distributed across the brain.[54] It treats

foci from individual experiments as centers for 3D Gaussian probability distributions

representing spatial uncertainty. The width of these distributions was determined based on

between-subject and between-template variability.[62] The ALE algorithm weighs the between-

subject variability by the number of participants for each study, based on the idea that

experiments of larger sample sizes are more likely to reliably report true activation effects.

Therefore, experiments with larger sample sizes are modeled by smaller Gaussian

distributions, resulting in a stronger influence on ALE scores, which indicate the probability

that at least one true peak activation lies in the voxel across the population of all possible

studies.[62] As a result, studies with larger sample sizes would be weighed more heavily

relative to studies with smaller sample sizes.

The ALE method is implemented in the following steps. First, for each included

study, a map of the activation likelihood is computed. Second, the maps are aggregated to

compute the ALE score for each voxel. Finally, a permutation test is employed to identify

voxels in which the ALE statistic is larger than expected by chance.[54,62,63] The ALE method

takes into account heterogeneity in spatial uncertainty across studies[54,62,63] and differences in

number of peak coordinates reported per cluster.[63] This approach allows random-effects

estimates of ALE, increasing generalizability of the results.[62]

Coordinate-based meta-analyses represent a departure from traditional meta-

analyses.[55] Specifically, whereas traditional meta-analyses aim to calculate pooled effect

sizes to determine the direction and magnitude of an effect based on a body of literature,

coordinate-based meta-analyses evaluate whether the location of an effect is consistent within

a body of literature. In other words, coordinate-based meta-analyses are blind to effect size

magnitude, but direction is tied to the analysis.[55] Although the coordinate-based meta-

analytic approach is limited to studies that report suprathreshold coordinates for a given

contrast, this approach will continue to be the most common method for conducting valid meta-analyses for neuroimaging studies until it becomes standard practice to release unthresholded statistical maps.

Statistical Analysis

All analyses were performed in Montreal Neurological Institute (MNI) space. Coordinates reported in Talairach space were converted to MNI using the "icbm2tal" transformation.[64] We assessed statistical significance and corrected for multiple comparisons using the permutation-based approach (N = 1000) recommended by the developers of GingerALE.[51,56] This approach utilized a cluster-forming threshold of $p < 0.001$ (uncorrected) and maintained a cluster-level family-wise error rate of 5%.[56] Additionally, we took steps to minimize within-group effects on the meta-analyses.[63] If a study reported more than one contrast (often referred to as an "experiment" in coordinate-based meta-analyses), the contrasts examining similar processes were pooled together to avoid double counting the same participants in a meta-analysis. For example, when a study reported between-group effects in response to $1.50 and $5 rewards relative to neutral or loss conditions, the coordinates derived from the two contrasts were coded as a single reward experiment.

To capture anatomical variation between individual human brains,[65] we show probabilistic anatomical labels for the locations of the maximum ALE values using the Harvard–Oxford cortical and subcortical atlases.[66] For transparency, all of the statistical maps (thresholded and unthresholded) derived from the meta-analyses are publicly available on NeuroVault at https://neurovault.org/collections/3884. Readers are free to access these maps and define brain regions using their own labels. Other study materials are available on Open Science Framework at https://osf.io/sjb4d.

Results

As shown in Figure 1, the systematic literature search identified a total of 41 neuroimaging studies that met the inclusion criteria, yielding 4 coordinate-based meta-analyses with at least 17 independent experiments. Supplementary Tables 2 and 3 show the characteristics of the included studies and their samples.

In the present meta-analytic dataset, for the MDD group, the mean number of participants was 19.9, the mean age was 36.4, the mean percentage of females was 60.9%, and the mean percentage of medication usage was 36.6%. For the HC group, the mean number of participants was 20.1, the mean age was 34.9, and the mean percentage of females was 60.3%. Types of reward or punishment used by the included studies encompass money, points, or voucher (41.5%; 17/41); faces (34.1%; 14/41); pictures (12.2%; 5/41); words, statements, captions, or paragraphs (12.2%; 5/41); and autobiographical memory (4.9%; 2/41). Both reward and punishment contrasts were reported in 41.5% (17/41) of studies; 29.3% (12/41) of studies reported punishment contrasts only; and 26.8% (11/41) of studies reported reward contrasts only.

We first synthesized results of 22 studies reporting less activity in response to reward in people with MDD than HC (i.e. HC > MDD for reward > punishment or neutral stimuli and/or neutral stimuli > punishment). As expected, results indicated that these studies reliably reported less activation in a single cluster extending bilaterally across the VS and including part of the subcallosal cortex in MDD (Table 1; Figure 2A).

In addition to examining which regions consistently showed hypo-responses to reward, we also examined which, if any, brain regions showed consistent hyper-responses to rewarding stimuli. We aggregated results of 18 studies reporting greater activity in response to reward in people with MDD than HC (i.e. MDD > HC for reward > punishment or neutral stimuli and/or neutral stimuli > punishment). Importantly, results indicated that these studies

reliably reported greater activation in the right OFC (in a region located in between the medial OFC and the lateral OFC) in MDD (Table 1; Figure 2B).

We conducted sensitivity analyses to examine whether excluding studies that used neutral stimuli > punishment as a reward contrast would affect the main results related to reward responses in MDD. After excluding the two experiments of neutral stimuli > punishment,[38,67] the results remained the same: We found significant convergence among experiments reporting blunted responses for reward in MDD relative to HC in the VS, as well as significant convergence among experiments reporting elevated responses for reward in MDD relative to HC in the OFC (Supplementary Table 4).

We also conducted exploratory analyses to examine which brain regions consistently show aberrant responses to punishment in MDD relative to HC. First, we meta-analyzed 24 studies reporting greater activity in response to punishment in people with MDD than HC (i.e. MDD > HC for punishment > reward or neutral stimuli and/or neutral stimuli > reward). The results indicated that these studies reliably reported greater activation in the left sublenticular extended amygdala in MDD (Table 2; Figure 3). Second, we synthesized 17 studies reporting less activity in response to punishment in people with MDD than HC (i.e. HC > MDD for punishment > reward or neutral stimuli and/or neutral stimuli > reward). The results indicated that these studies did not report consistent activation patterns. Together, these results suggest that relative to HC, people with MDD exhibited hyper-responses in the left sublenticular extended amygdala during processing of punishment-relevant stimuli.

Discussion

A growing number of researchers have used neuroimaging methods to enhance our understanding of the underlying pathophysiology of MDD.[68–70] Many of these studies have shown that patients with MDD exhibit blunted responses in the VS, but more disparate patterns of responses in other brain areas. Therefore, it remains unclear what brain regions,

other than the VS, are most consistently implicated in people with MDD, particularly during reward processing (see Supplementary Table 1 and Supplementary Figure 1). To address this issue, we performed a coordinate-based meta-analysis of 41 neuroimaging studies containing reward-related responses from a total of 794 patients with MDD and 803 HC. Our meta-analytic findings confirm that reward responses within the VS are consistently blunted in MDD relative to HC across studies. In contrast, we find that reward responses within the OFC are consistently elevated in MDD. Contrary to the common notion that MDD is characterized by blunted responses to reward, these findings suggest that MDD may be characterized by both hypo- and hyper-responses to reward at the neural level and highlight the need for a more fine-tuned understanding of the various components of reward processing in MDD.

Although our blunted striatal findings are in line with previous meta-analytic work documenting reward processing abnormalities in MDD,[34,35,50] we emphasize that our work differs in two key ways. First, our results implicate highly specific—yet distinct— abnormalities in the reward circuit, with hypo-responses to reward in the VS and hyper-responses to reward in the OFC. In sharp contrast, previous meta-analyses have generally reported distributed patterns of abnormalities, with little anatomical agreement across studies (see Supplementary Table 1 and Supplementary Figure 1). Second, to minimize bias, our study employed more stringent analysis methods than prior studies in this area, following recommendations by new guidelines.[52,55] For example, instead of using the FDR approach which has been shown to be inadequate in controlling the false positives among clusters in neuroimaging meta-analyses,[53,54] we corrected for multiple comparisons using the permutation-based approach. We also excluded ROI- or SVC-based studies and only included whole-brain studies that used an active control condition and independent samples. In addition, we only conducted a meta-analysis when there were at least 17 eligible experiments

to restrict excessive contribution of any particular studies to cluster-level thresholding.[56] We speculate that the enhanced rigor and methods of our study contributed to our ability to identify highly circumscribed and distinct abnormalities in the reward circuit.

A prior meta-analysis using similarly rigorous methods revealed no significant convergence of findings among neuroimaging studies comparing MDD and HC.[52] Nevertheless, we note that the previous meta-analysis differed from the current meta-analysis in at least four key ways. First, whereas the previous meta-analysis focused on emotional or cognitive processing, the current meta-analysis focused solely on reward processing. Second, the previous meta-analysis excluded participants younger than 18 years old; in contrast, the current meta-analysis included participants of all ages, boosting its power and ability to generalize its findings to MDD across ages. Finally, the previous meta-analysis excluded MDD participants in remission, whereas the current meta-analysis included them, leading to the question of whether reward processing dysfunction is not simply a state, but a trait of MDD. This issue needs to be addressed explicitly when there are enough studies available to compare current to remitted depression. Our ability to identify significant convergence highlights the significance of reward processing dysfunction in MDD and might indicate the literature on reward processing in MDD is more homogeneous than that on emotional or cognitive processing in MDD.

In our view, the most important current finding is that studies consistently report that people with MDD exhibit hyper-responses to reward in an OFC region located between the medial OFC and the lateral OFC.[71] Exposure to rewards (e.g., money and pleasant sights) evokes activity in the OFC, which has been associated with the computation of reward value[18–24] as well as the representation of current stimulus-outcome associations.[72,73] For example, whereas the lateral OFC is responsive to stimuli of both positive and negative valence, the medial OFC is particularly responsive to cues of positive valence.[74–76] Therefore,

given that MDD is traditionally linked to blunted response to reward or reduced capacity to experience pleasure,[9] the finding of hyperactivity of the OFC in response to reward in MDD may seem paradoxical. One interpretation would be that MDD is at least partly characterized by hyper-responses to reward, which fits with a set of experimental studies reporting that individuals with severe MDD found dextroamphetamine to be more rewarding than did controls.[77–79]

Alternatively, OFC hyperactivity may reflect enhanced inhibitory control over subcortical regions underlying reward-related behavior, leading to anhedonia. Optogenetic and neuroimaging studies have revealed that hyperactivity in prefrontal regions (e.g., medial PFC, vmPFC) innervated by glutamatergic neurons may causally inhibit reward-related behavior via suppressing striatal responses to dopamine neurons in the midbrain[7,26] and increasing connectivity between the medial PFC, lateral OFC, and VS.[7,26] These findings from animal studies suggest that prefrontal regions (e.g., medial PFC, vmPFC, and lateral OFC) might exert inhibitory control over subcortical regions underlying reward-relevant behavior, potentially explaining the central role of anhedonia in MDD. Nevertheless, it is important to note that given the complexity of the OFC and the inability of the meta-analysis to unravel its varied functions, these interpretations remain speculative.

The extent to which corticostriatal connectivity during reward processing is disrupted in MDD remains an open and important question.[2,80,81] Previous meta-analyses indicate that at least some people with MDD exhibit dysfunction in resting-state corticostriatal connectivity.[80,81] The current meta-analytic results will provide a springboard for future studies that seek to develop a full picture of the pathophysiology of MDD and understand the role of dysregulated corticostriatal connectivity in MDD, particularly in the context of reward processing. These endeavors may require empirical assessments of connectivity within the reward circuit using psychophysiological interaction analysis[82–84] and dynamic causal

modeling.[85] Such approaches have shown promise for revealing specific patterns of task-dependent corticostriatal interactions in samples containing healthy individuals,[86] clinical populations,[2,87] or a mix of both.[88] Nevertheless, a caveat of such approaches is that dysregulated corticostriatal connectivity may involve modulatory regions, such as the midbrain.[89]

In addition to distinct abnormalities within the reward circuit, the current study also found that MDD is associated with hyper-responses in the left sublenticular extended amygdala in response to punishment. The current finding fits with others in suggesting that amygdala hyperactivation is linked to the processing of affectively salient, especially punishing, stimuli in MDD, and may underlie negativity bias in depression.[90,91] It is also in agreement with a meta-analysis indicating increased activation in the amygdala in response to negative stimuli in MDD relative to HC[68] and a series of studies indicating that the amygdala may be a key brain region implicated in the pathophysiology of depression.[92,93] Interestingly, longitudinal studies have reported that amygdala reactivity, potentially in combination with life stress, prospectively predicts internalizing (e.g., depressive and anxiety) symptoms,[94,95] highlighting the importance of amygdala reactivity in the course of depression.

Even though the current meta-analysis reveals circumscribed patterns of abnormal responses to reward and punishment, we note that the findings should be interpreted in the context of their limitations. First, heterogeneity across studies may have added noise to the current analyses and restricted its capacity for detecting true effects. Specifically, due to the limited number of studies, the current analyses collapsed across different reward processes (e.g., anticipation and outcome), reward modalities (e.g., monetary and social), and specific contrasts that would help isolate and differentiate neural responses to salience and valence.[58] The current analyses also collapsed across different mood states, psychotropic medication usage, ages, and comorbidities. In doing so, important differences in brain activation may be

obscured and more specific questions related to brain activation—particularly questions related to neural representations of valence or salience—cannot be addressed in the current work. Future studies should examine how these factors may affect reward processing in MDD. Nevertheless, we highlight that the convergence of findings despite the heterogeneity of the included studies is striking and suggests that the current findings may reflect central abnormalities of MDD.

Second, some included studies had relatively small sample sizes and reported coordinates that were not corrected for multiple comparisons, which may lead to biased results.[45] Future work should follow the most updated guidelines for best practices in the field to avoid generating biased findings.[96] Third, most of the included studies only recruited adults with acute major depression. More studies on other ages (e.g., pre-adolescents, adolescents) and mood states (e.g., remission) are needed. Fourth, we note that the search criteria were designed to focus on studies on reward and might not have identified some studies on punishment. Therefore, the analyses and results in relation to punishment are exploratory in nature and should be interpreted with caution. Fifth, the ALE method, by nature, cannot incorporate null results.[55] As a result, the current findings could be confounded by publication bias. Sixth, coordinate-based meta-analyses inherently are blind to effect sizes. Therefore, no claims about the magnitude of the effects observed in the current study could be made. Finally, some patients in the included studies were medicated. The normalizing effects of treatment could obscure differences between MDD and HC, increasing the probability of type II errors.[97]

Notwithstanding these caveats, the current meta-analysis shows that MDD is consistently associated with opposing abnormalities in the reward circuit in response to reward: hypo-response in the VS and hyper-response in the OFC. The current meta-analytic results therefore suggest that MDD is associated with both hypo and hyper responses to

reward. MDD may stem, in part, from dysregulated connectivity between the VS and the OFC. We believe the current findings will help lay a foundation towards developing a more refined understanding and treatment of MDD and its comorbid psychiatric disorders, particularly ones that involve abnormal reward processing.[98] For example, a more refined understanding of the abnormalities in the reward circuitry in MDD may help distinguish it from other disorders exhibiting reward processing abnormalities, such as bipolar disorder, schizophrenia, and substance use disorder. Moreover, although reinforcement learning models, such as actor-critic models and prediction error models, have been utilized to understand the pathophysiology of several psychiatric disorders (e.g., schizophrenia and addiction), research on their application on MDD has been scant.[99] The current results help delineate specific abnormalities within the reward circuit and supply a foundation for refining connectivity-based and computational models of MDD. Finally, given that previous treatment targets for deep brain stimulation for treatment-resistant depression have yielded mixed results,[32] the portion of OFC implicated by the current results could be a promising treatment target.

Acknowledgements

Conflict of Interest

The authors declare no conflict of interest.

References

Friedrich MJ. Depression is the leading cause of disability around the world. *JAMA* 2017; **317**: 1517.

Admon R, Pizzagalli DA. Dysfunctional reward processing in depression. *Curr Opin Psychol* 2015; **4**: 114–118.

Alloy LB, Olino T, Freed RD, Nusslock R. Role of reward sensitivity and processing in major depressive and bipolar spectrum disorders. *Behav Ther* 2016; **47**: 600–621.

Forbes EE, Dahl RE. Altered Reward Function in Adolescent Depression: What, When, and How? *J Child Psychol Psychiatry* 2012; **53**: 3–15.

Nusslock R, Alloy LB. Reward Processing and Mood-Related Symptoms: An RDoC and Translational Neuroscience Perspective. *J Affect Disord* 2017; **216**: 3–16.

Olino TM. Future Research Directions in the Positive Valence Systems: Measurement, Development, and Implications for Youth Unipolar Depression. *J Clin Child Adolesc Psychol Off J Soc Clin Child Adolesc Psychol Am Psychol Assoc Div 53* 2016; **45**: 681–705.

Robbins TW. Illuminating anhedonia. *Science* 2016; **351**: 24–25.

Treadway MT, Zald DH. Reconsidering Anhedonia in Depression: Lessons from Translational Neuroscience. *Neurosci Biobehav Rev* 2011; **35**: 537–555.

Whitton AE, Treadway MT, Pizzagalli DA. Reward processing dysfunction in major depression, bipolar disorder and schizophrenia. *Curr Opin Psychiatry* 2015; **28**: 7–12.

Arrondo G, Segarra N, Metastasio A, Ziauddeen H, Spencer J, Reinders NR *et al.* Reduction in ventral striatal activity when anticipating a reward in depression and schizophrenia: A replicated cross-diagnostic finding. *Front Psychol* 2015; **6**. doi:10.3389/fpsyg.2015.01280.

McCabe C, Cowen PJ, Harmer CJ. Neural representation of reward in recovered depressed patients. *Psychopharmacology (Berl)* 2009; **205**: 667–677.

Pizzagalli DA, Holmes AJ, Dillon DG, Goetz EL, Birk JL, Bogdan R *et al.* Reduced caudate and nucleus accumbens response to rewards in unmedicated subjects with major depressive disorder. *Am J Psychiatry* 2009; **166**: 702–710.

Smoski MJ, Felder J, Bizzell J, Green SR, Ernst M, Lynch TR *et al.* FMRI of alterations in reward selection, anticipation, and feedback in major depressive disorder. *J Affect Disord* 2009; **118**: 69–78.

Luking KR, Pagliaccio D, Luby JL, Barch DM. Reward processing and risk for depression across development. *Trends Cogn Sci* 2016; **20**: 456–468.

Knutson B, Bhanji JP, Cooney RE, Atlas LY, Gotlib IH. Neural responses to monetary incentives in major depression. *Biol Psychiatry* 2008; **63**: 686–692.

Haber SN. Corticostriatal circuitry. *Dialogues Clin Neurosci* 2016; **18**: 7–21.

Haber SN, Knutson B. The reward circuit: Linking primate anatomy and human imaging. *Neuropsychopharmacology* 2010; **35**: 4–26.

Berridge KC, Kringelbach ML. Pleasure systems in the brain. *Neuron* 2015; **86**: 646–664.

Der-Avakian A, Markou A. The neurobiology of anhedonia and other reward-related deficits. *Trends Neurosci* 2012; **35**: 68–77.

Levy DJ, Glimcher PW. The root of all value: A neural common currency for choice. *Curr Opin Neurobiol* 2012; **22**: 1027–1038.

Padoa-Schioppa C, Conen KE. Orbitofrontal cortex: A neural circuit for economic decisions. *Neuron* 2017; **96**: 736–754.

Rangel A, Camerer C, Montague PR. A framework for studying the neurobiology of value-based decision making. *Nat Rev Neurosci* 2008; **9**: 545–556.

Saez RA, Saez A, Paton JJ, Lau B, Salzman CD. Distinct roles for the amygdala and orbitofrontal cortex in representing the relative amount of expected reward. *Neuron* 2017; **95**: 70-77.e3.

Stalnaker TA, Cooch NK, Schoenbaum G. What the orbitofrontal cortex does not do. *Nat Neurosci* 2015; **18**: 620–627.

Delgado MR, Beer JS, Fellows LK, Huettel SA, Platt ML, Quirk GJ *et al.* Viewpoints: Dialogues on the functional role of the ventromedial prefrontal cortex. *Nat Neurosci* 2016; **19**: 1545–1552.

Ferenczi EA, Zalocusky KA, Liston C, Grosenick L, Warden MR, Amatya D *et al.* Prefrontal cortical regulation of brainwide circuit dynamics and reward-related behavior. *Science* 2016; **351**: aac9698.

Peters J, Büchel C. Neural representations of subjective reward value. *Behav Brain Res* 2010; **213**: 135–141.

Phelps EA, Lempert KM, Sokol-Hessner P. Emotion and decision making: Multiple modulatory neural circuits. *Annu Rev Neurosci* 2014; **37**: 263–287.

Voorn P, Vanderschuren LJMJ, Groenewegen HJ, Robbins TW, Pennartz CMA. Putting a spin on the dorsal-ventral divide of the striatum. *Trends Neurosci* 2004; **27**: 468–474.

Stringaris A, Vidal-Ribas Belil P, Artiges E, Lemaitre H, Gollier-Briant F, Wolke S *et al.* The brain's response to reward anticipation and depression in

adolescence: Dimensionality, specificity, and longitudinal predictions in a community-based sample. *Am J Psychiatry* 2015; **172**: 1215–1223.

Morgan JK, Olino TM, McMakin DL, Ryan ND, Forbes EE. Neural response to reward as a predictor of increases in depressive symptoms in adolescence. *Neurobiol Dis* 2013; **52**: 66–74.

Naesström M, Blomstedt P, Bodlund O. A systematic review of psychiatric indications for deep brain stimulation, with focus on major depressive and obsessive-compulsive disorder. *Nord J Psychiatry* 2016; **70**: 483–491.

Heshmati M, Russo SJ. Anhedonia and the brain reward circuitry in depression. *Curr Behav Neurosci Rep* 2015; **2**: 146–153.

Zhang WN, Chang SH, Guo LY, Zhang KL, Wang J. The neural correlates of reward-related processing in major depressive disorder: A meta-analysis of functional magnetic resonance imaging studies. *J Affect Disord* 2013; **151**: 531–539.

Groenewold NA, Opmeer EM, de Jonge P, Aleman A, Costafreda SG. Emotional valence modulates brain functional abnormalities in depression: Evidence from a meta-analysis of fMRI studies. *Neurosci Biobehav Rev* 2013; **37**: 152–163.

Hanson JL, Hariri AR, Williamson DE. Blunted ventral striatum development in adolescence reflects emotional neglect and predicts depressive symptoms. *Biol Psychiatry* 2015; **78**: 598–605.

Forbes EE, May JC, Siegle GJ, Ladouceur CD, Ryan ND, Carter CS *et al.* Reward-related decision-making in pediatric major depressive disorder: An fMRI study. *J Child Psychol Psychiatry* 2006; **47**: 1031–1040.

Demenescu LR, Renken R, Kortekaas R, van Tol M-J, Marsman JBC, van Buchem MA *et al.* Neural correlates of perception of emotional facial expressions in out-patients with mild-to-moderate depression and anxiety. A multicenter fMRI study. *Psychol Med* 2011; **41**: 2253–2264.

Forbes EE, Hariri AR, Martin SL, Silk JS, Moyles DL, Fisher PM *et al.* Altered striatal activation predicting real-world positive affect in adolescent major depressive disorder. *Am J Psychiatry* 2009; **166**: 64–73.

Rizvi SJ, Salomons TV, Konarski JZ, Downar J, Giacobbe P, McIntyre RS *et al.* Neural response to emotional stimuli associated with successful antidepressant treatment and behavioral activation. *J Affect Disord* 2013; **151**: 573–581.

Keedwell PA, Andrew C, Williams SCR, Brammer MJ, Phillips ML. A double dissociation of ventromedial prefrontal cortical responses to sad and happy stimuli in depressed and healthy individuals. *Biol Psychiatry* 2005; **58**: 495–503.

Mitterschiffthaler MT, Kumari V, Malhi GS, Brown RG, Giampietro VP, Brammer MJ *et al.* Neural response to pleasant stimuli in anhedonia: An fMRI study. *Neuroreport* 2003; **14**: 177–182.

Dichter GS, Kozink RV, McClernon FJ, Smoski MJ. Remitted major depression is characterized by reward network hyperactivation during reward anticipation and hypoactivation during reward outcomes. *J Affect Disord* 2012; **136**: 1126–1134.

Kumari V, Mitterschiffthaler MT, Teasdale JD, Malhi GS, Brown RG, Giampietro V *et al.* Neural abnormalities during cognitive generation of affect in treatment-resistant depression. *Biol Psychiatry* 2003; **54**: 777–791.

Button KS, Ioannidis JPA, Mokrysz C, Nosek BA, Flint J, Robinson ESJ *et al.* Power failure: Why small sample size undermines the reliability of neuroscience. *Nat Rev Neurosci* 2013; **14**: 365–376.

Jia XZ, Zhao N, Barton B, Burciu R, Carriere N, Cerasa A *et al.* Small effect size leads to reproducibility failure in resting-state fMRI studies. *bioRxiv* 2018. doi:10.1101/285171.

Poldrack RA, Baker CI, Durnez J, Gorgolewski KJ, Matthews PM, Munafò MR *et al.* Scanning the horizon: Towards transparent and reproducible neuroimaging research. *Nat Rev Neurosci* 2017; **18**: 115–126.

Chase HW, Kumar P, Eickhoff SB, Dombrovski AY. Reinforcement learning models and their neural correlates: An activation likelihood estimation meta-analysis. *Cogn Affect Behav Neurosci* 2015; **15**: 435–459.

Ojemann JG, Akbudak E, Snyder AZ, McKinstry RC, Raichle ME, Conturo TE. Anatomic localization and quantitative analysis of gradient refocused echo-planar fMRI susceptibility artifacts. *NeuroImage* 1997; **6**: 156–167.

Keren H, O'Callaghan G, Vidal-Ribas P, Buzzell GA, Brotman MA, Leibenluft E *et al.* Reward processing in depression: A conceptual and meta-analytic review across fMRI and EEG studies. *Am J Psychiatry* 2018; **175**: 1111–1120.

Eickhoff SB, Laird AR, Fox PM, Lancaster JL, Fox PT. Implementation errors in the GingerALE software: Description and recommendations. *Hum Brain Mapp* 2017; **38**: 7–11.

Müller VI, Cieslik EC, Serbanescu I, Laird AR, Fox PT, Eickhoff SB. Altered brain activity in unipolar depression revisited: Meta-analyses of neuroimaging studies. *JAMA Psychiatry* 2017; **74**: 47–55.

Chumbley JR, Friston KJ. False discovery rate revisited: FDR and topological inference using Gaussian random fields. *NeuroImage* 2009; **44**: 62–70.

Eickhoff SB, Bzdok D, Laird AR, Kurth F, Fox PT. Activation Likelihood Estimation meta-analysis revisited. *Neuroimage* 2012; **59**: 2349–2361.

Müller VI, Cieslik EC, Laird AR, Fox PT, Radua J, Mataix-Cols D *et al.* Ten simple rules for neuroimaging meta-analysis. *Neurosci Biobehav Rev* 2017. doi:10.1016/j.neubiorev.2017.11.012.

Eickhoff SB, Nichols TE, Laird AR, Hoffstaedter F, Amunts K, Fox PT *et al.*
      Behavior, sensitivity, and power of activation likelihood estimation
      characterized by massive empirical simulation. *NeuroImage* 2016; **137**: 70–85.

57   Bartra O, McGuire JT, Kable JW. The valuation system: A coordinate-based
      meta-analysis of BOLD fMRI experiments examining neural correlates of
      subjective value. *NeuroImage* 2013; **76**: 412–427.

58   Cooper JC, Knutson B. Valence and salience contribute to nucleus accumbens
      activation. *NeuroImage* 2008; **39**: 538–547.

59   Moher D, Liberati A, Tetzlaff J, Altman DG, PRISMA Group. Preferred
      reporting items for systematic reviews and meta-analyses: The PRISMA
      statement. *Ann Intern Med* 2009; **151**: 264–269.

60   Hayden BY, Parikh PC, Deaner RO, Platt ML. Economic principles motivating
      social attention in humans. *Proc R Soc B Biol Sci* 2007; **274**: 1751–1756.

61   Tsukiura T, Cabeza R. Orbitofrontal and hippocampal contributions to memory
      for face-name associations: The rewarding power of a smile. *Neuropsychologia*
      2008; **46**: 2310–2319.

62   Eickhoff SB, Laird AR, Grefkes C, Wang LE, Zilles K, Fox PT. Coordinate-
      based activation likelihood estimation meta-analysis of neuroimaging data: A
      random-effects approach based on empirical estimates of spatial uncertainty.
      Hum Brain Mapp 2009; **30**: 2907–2926.

63   Turkeltaub PE, Eickhoff SB, Laird AR, Fox M, Wiener M, Fox P. Minimizing
      within-experiment and within-group effects in Activation Likelihood Estimation
      meta-analyses. *Hum Brain Mapp* 2012; **33**: 1–13.

64   Lancaster JL, Tordesillas-Gutiérrez D, Martinez M, Salinas F, Evans A, Zilles K
      et al. Bias between MNI and Talairach coordinates analyzed using the ICBM-
      152 brain template. *Hum Brain Mapp* 2007; **28**: 1194–1205.

65   Mazziotta JC, Toga AW, Evans A, Fox P, Lancaster J. A probabilistic atlas of
      the human brain: Theory and rationale for its development. The International
      Consortium for Brain Mapping (ICBM). *NeuroImage* 1995; **2**: 89–101.

66   Desikan RS, Ségonne F, Fischl B, Quinn BT, Dickerson BC, Blacker D *et al.* An
      automated labeling system for subdividing the human cerebral cortex on MRI
      scans into gyral based regions of interest. *NeuroImage* 2006; **31**: 968–980.

67   Wagner G, Schachtzabel C, Peikert G, Bär K-J. The neural basis of the abnormal
      self-referential processing and its impact on cognitive control in depressed
      patients. *Hum Brain Mapp* 2015; **36**: 2781–2794.

68   Hamilton JP, Etkin A, Furman DJ, Lemus MG, Johnson RF, Gotlib IH.
      Functional neuroimaging of major depressive disorder: A meta-analysis and new
      integration of base line activation and neural response data. *Am J Psychiatry*
      2012; **169**: 693–703.

Miller CH, Hamilton JP, Sacchet MD, Gotlib IH. Meta-analysis of functional neuroimaging of major depressive disorder in youth. *JAMA Psychiatry* 2015; **72**: 1045–1053.

Palmer SM, Crewther SG, Carey LM, START Project Team. A meta-analysis of changes in brain activity in clinical depression. *Front Hum Neurosci* 2014; **8**: 1045.

Hare TA, O'Doherty J, Camerer CF, Schultz W, Rangel A. Dissociating the role of the orbitofrontal cortex and the striatum in the computation of goal values and prediction errors. *J Neurosci* 2008; **28**: 5623–5630.

Pickens CL, Saddoris MP, Setlow B, Gallagher M, Holland PC, Schoenbaum G. Different roles for orbitofrontal cortex and basolateral amygdala in a reinforcer devaluation task. *J Neurosci* 2003; **23**: 11078–11084.

Sharpe MJ, Stalnaker T, Schuck NW, Killcross S, Schoenbaum G, Niv Y. An integrated model of action selection: Distinct modes of cortical control of striatal decision making. *Annu Rev Psychol* 2019; **70**: 53–76.

Lewis PA, Critchley HD, Rotshtein P, Dolan RJ. Neural correlates of processing valence and arousal in affective words. *Cereb Cortex* 2007; **17**: 742–748.

Nusslock R, Alloy LB. Reward processing and mood-related symptoms: An RDoC and translational neuroscience perspective. *J Affect Disord* 2017; **216**: 3–16.

Schmidt L, Cléry-Melin M-L, Lafargue G, Valabrègue R, Fossati P, Dubois B *et al.* Get aroused and be stronger: Emotional facilitation of physical effort in the human brain. *J Neurosci* 2009; **29**: 9450–9457.

Tremblay LK, Naranjo CA, Graham SJ, Herrmann N, Mayberg HS, Hevenor S *et al.* Functional neuroanatomical substrates of altered reward processing in major depressive disorder revealed by a dopaminergic probe. *Arch Gen Psychiatry* 2005; **62**: 1228–1236.

Tremblay LK, Naranjo CA, Cardenas L, Herrmann N, Busto UE. Probing brain reward system function in major depressive disorder: Altered response to dextroamphetamine. *Arch Gen Psychiatry* 2002; **59**: 409–416.

Naranjo CA, Tremblay LK, Busto UE. The role of the brain reward system in depression. *Prog Neuropsychopharmacol Biol Psychiatry* 2001; **25**: 781–823.

Kaiser RH, Andrews-Hanna JR, Wager TD, Pizzagalli DA. Large-scale network dysfunction in major depressive disorder: A meta-analysis of resting-state functional connectivity. *JAMA Psychiatry* 2015; **72**: 603–611.

Drysdale AT, Grosenick L, Downar J, Dunlop K, Mansouri F, Meng Y *et al.* Resting-state connectivity biomarkers define neurophysiological subtypes of depression. *Nat Med* 2017; **23**: 28–38.

Friston KJ, Buechel C, Fink GR, Morris J, Rolls E, Dolan RJ. Psychophysiological and modulatory interactions in neuroimaging. *NeuroImage* 1997; **6**: 218–229.

McLaren DG, Ries ML, Xu G, Johnson SC. A generalized form of context-dependent psychophysiological interactions (gPPI): A comparison to standard approaches. *NeuroImage* 2012; **61**: 1277–1286.

Smith DV, Gseir M, Speer ME, Delgado MR. Toward a cumulative science of functional integration: A meta-analysis of psychophysiological interactions. *Hum Brain Mapp* 2016; **37**: 2904–2917.

Friston KJ, Harrison L, Penny W. Dynamic causal modelling. *NeuroImage* 2003; **19**: 1273–1302.

Smith DV, Rigney AE, Delgado MR. Distinct reward properties are encoded via corticostriatal interactions. *Sci Rep* 2016; **6**: 20093.

Admon R, Pizzagalli DA. Corticostriatal pathways contribute to the natural time course of positive mood. *Nat Commun* 2015; **6**: 10065.

Hanson JL, Knodt AR, Brigidi BD, Hariri AR. Heightened connectivity between the ventral striatum and medial prefrontal cortex as a biomarker for stress-related psychopathology: Understanding interactive effects of early and more recent stress. *Psychol Med* 2017; : 1–9.

Murty VP, Shermohammed M, Smith DV, Carter RM, Huettel SA, Adcock RA. Resting state networks distinguish human ventral tegmental area from substantia nigra. *NeuroImage* 2014; **100**: 580–589.

Roiser JP, Elliott R, Sahakian BJ. Cognitive mechanisms of treatment in depression. *Neuropsychopharmacology* 2012; **37**: 117–136.

Eshel N, Roiser JP. Reward and punishment processing in depression. *Biol Psychiatry* 2010; **68**: 118–124.

Price JL, Drevets WC. Neurocircuitry of mood disorders. *Neuropsychopharmacology* 2010; **35**: 192–216.

Rive MM, van Rooijen G, Veltman DJ, Phillips ML, Schene AH, Ruhé HG. Neural correlates of dysfunctional emotion regulation in major depressive disorder. A systematic review of neuroimaging studies. *Neurosci Biobehav Rev* 2013; **37**: 2529–2553.

Swartz JR, Knodt AR, Radtke SR, Hariri AR. A neural biomarker of psychological vulnerability to future life stress. *Neuron* 2015; **85**: 505–511.

Mattson WI, Hyde LW, Shaw DS, Forbes EE, Monk CS. Clinical neuroprediction: Amygdala reactivity predicts depressive symptoms 2 years later. *Soc Cogn Affect Neurosci* 2016; **11**: 892–898.

Nichols TE, Das S, Eickhoff SB, Evans AC, Glatard T, Hanke M *et al.* Best practices in data analysis and sharing in neuroimaging using MRI. *Nat Neurosci* 2017; **20**: 299–303.

Dichter GS, Felder JN, Petty C, Bizzell J, Ernst M, Smoski MJ. The effects of psychotherapy on neural responses to rewards in major depression. *Biol Psychiatry* 2009; **66**: 886–897.

Diehl MM, Lempert KM, Parr AC, Ballard I, Steele VR, Smith DV. Toward an integrative perspective on the neural mechanisms underlying persistent maladaptive behaviors. *Eur J Neurosci* 2018; **48**: 1870–1883.

Maia TV, Frank MJ. From reinforcement learning models to psychiatric and neurological disorders. *Nat Neurosci* 2011; **14**: 154–162.

Figure Legends

Figure 1. Flowchart of study selection. The systematic literature search identified a total of 41 neuroimaging studies that met the inclusion criteria, yielding 4 coordinate-based meta-analyses with at least 17 independent studies; ROI, region of interest; SVC, small volume correction; MDD, major depressive disorder.

Figure 2. Opposing abnormalities in the reward circuit in response to reward in major depressive disorder. **(A)** To examine regions that consistently showed blunted response to reward, 22 studies reporting less activity in response to reward in people with major depressive disorder (MDD) than healthy controls (HC) were synthesized. Results indicated that these studies reliably report less activation in the ventral striatum (VS) in MDD. **(B)** To identify regions that consistently showed hyper-responses to reward, 18 studies reporting greater activity in response to reward in people with MDD than HC were meta-analyzed. Results indicated that these studies reliably report greater activation in the right orbitofrontal cortex (OFC) in MDD.

Figure 3. Hyper-responses to punishment in the sublenticular extended amygdala in major depressive disorder. To conduct exploratory analyses to examine which brain regions consistently show elevated response to punishment in major depressive disorder (MDD) relative to healthy controls (HC), we meta-analyzed 24 studies reporting greater activity in response to punishment in people with MDD than HC. The results indicated that these studies reliably report greater activation in the left sublenticular extended amygdala (SLEA) in MDD.

Table 1

Peak Coordinates of Group Differences in Neural Responses to Reward

Contrast	Cluster Size (mm^3)	Probabilistic Anatomical Label	x	y	z
MDD > HC	912	Frontal Orbital Cortex (26%), Frontal Pole (13%)	20	32	-12
HC > MDD	1768	Subcallosal Cortex (14%)	-2	8	-4
		Caudate (32.1%), Accumbens (11.1%)	8	6	-2

Abbreviations: MDD, major depressive disorder; HC, healthy controls. Coordinates are x,y,z values of the locations of the maximum activation likelihood estimation (ALE) values in MNI space. Probabilistic labels reflect the probability that a coordinate belongs to a given region derived from the Harvard-Oxford probabilistic atlas. For clarity, we only report labels whose likelihood exceeds 5%.

Table 2

Peak Coordinates of Group Differences in Neural Responses to Punishment

Contrast	Cluster Size (mm^3)	Probabilistic Anatomical Label	x	y	z
MDD > HC	1104	Amygdala (85.4%)	-26	-8	-14
		Amygdala (61.4%)	-16	-2	-18

Abbreviations: MDD, major depressive disorder; HC, healthy controls. Coordinates are x,y,z values of the locations of the maximum activation likelihood estimation (ALE) values in MNI space. Probabilistic labels reflect the probability that a coordinate belongs to a given region derived from the Harvard-Oxford probabilistic atlas. For clarity, we only report labels whose likelihood exceeds 5%.

A Hypo Response to Reward in the VS

L

Y = 12   Z = 1   X = -8

B Hyper Response to Reward in the OFC

Y = 32   Z = -15   X = 23

Hyper Response to Punishment in the SLEA

L

Y = -6   Z = -16   X = -26