Effect of set up protocols on the accuracy of alchemical free energy calculation over a set of ACK1 inhibitors

José M. Granadino-Roldán[a,], Antonia S. J. S. Mey[b], Juan J. Pérez González[c], Stefano Bosisio[b], Jaime Rubio-Martinez[d], Julien Michel,[b,*]*

a) *Departamento de Química Física, Facultad de Ciencias Experimentales, Universidad de Jaén, Campus "Las Lagunillas" s/n, 23071, Jaén, Spain*

b) EaStCHEM School of Chemistry, Joseph Black Building, The King's Buildings, Edinburgh, EH9 3FJ, UK

c) Department of Chemical Engineering, Universitat Politecnica de Catalunya, ETSEIB. Av. Diagonal, 647, 08028 Barcelona, Spain

d) *Departament de Química Física, Universitat de Barcelona (UB) and the Institut de Recerca en Quimica Teorica i Computacional (IQTCUB), Martí i Franqués 1, 08028 Barcelona, Spain*

* José M. Granadino-Roldán. Tel.: +34953213057. E-mail jmroldan@ujaen.es

* Julien Michel. Tel.: +441316504797. E-mail julien.michel@ed.ac.uk

ABSTRACT

Hit-to-lead virtual screening frequently relies on a cascade of computational methods that starts

with rapid calculations applied to a large number of compounds and ends with more expensive

computations restricted to a subset of compounds that passed initial filters. This work focuses

on set up protocols for alchemical free energy (AFE) scoring in the context of a Docking –

MM/PBSA – AFE cascade. A dataset of 15 congeneric inhibitors of the ACK1 protein was used

to evaluate the performance of AFE set up protocols that varied in the steps taken to prepare

input files (using previously docked and best scored poses, manual selection of poses, manual

placement of binding site water molecules). The main finding is that use of knowledge derived

from X-ray structures to model binding modes, together with the manual placement of a

bridging water molecule, improves the R^2 from 0.45 ± 0.06 to 0.76 ± 0.02 and decreases the

mean unsigned error from 2.11 ± 0.08 to 1.24 ± 0.04 kcal mol^{-1}. By contrast a brute force

automated protocol that increased the sampling time ten-fold lead to little improvements in

accuracy. Besides, it is shown that for the present dataset hysteresis can be used to flag poses

that need further attention even without prior knowledge of experimental binding affinities.

INTRODUCTION

There is continuous interest in computational methods to decrease time and costs of hit-to-lead and lead optimization efforts in preclinical drug discovery (1). A recurring topic in computational chemistry is the use of virtual *in silico* screens to find ligands for proteins (2, 3). Typically, the goal is to filter via a cascade of computational methods a large library to focus experimental efforts on a small number of molecules. Usually inexpensive methodologies are applied first to eliminate a large number of poorly suited molecules, with more expensive calculations reserved to a subset of promising ligands. This approach may be applied in the context of hit discovery where the goal is to identify a few weak binders from a library of existing molecules; or for hit-to-lead efforts where the goal is to identify analogues of a hit structure that could be prioritized for synthesis and assays. In both cases the main steps frequently involve library screening, docking, initial scoring, and re-scoring with diverse molecular simulation methods such as Molecular Mechanics Poisson Boltzmann (Generalized Born) Surface Area (MM/PBSA) (4), Linear Interaction Energy (LIE) (5) or Free energy Perturbation (FEP) (6) methods (7).

In a previous study a multistep docking and scoring protocol was benchmarked in the context of re-scoring with the MM/PB(GB)SA method (8). The set of ligands analysed in that study belonged to the same scaffold and it was assumed that the core binding mode of the conserved scaffold would not deviate from that of the experimentally X-ray resolved one. The present study investigates the suitability of alchemical free energy (AFE) methods for improving on this multistep docking and scoring protocol by means of a further re-scoring of ligands. AFE methods are increasingly used for predictions of free energies of binding in blinded competitions such as SAMPL (Statistic Assessment of Modelling of Proteins and Ligands) and D3R grand challenges (9-15). Some AFE protocols have even achieved predictions of binding energies with root mean square deviations (RMSD) under 1.5 kcal mol^{-1}, and Pearson Correlation coefficients (R) of around 0.7 or better (16-18). Nevertheless, the performance

varies significantly between different AFE protocols and targets (19-21) and it is important to

explore further the robustness of these methodologies.

Specifically, this study aimed to explore the extent to which a setup protocol motivated by

previous domain knowledge may influence the accuracy of AFE calculations, and whether

issues such as binding poses selection or binding site water placement can be overcome via an

increase of the simulation time. This was investigated using a dataset of 15 congeneric

inhibitors of the protein activated Cdc42-associated kinase (ACK1) (22), a potential cancer

target (23, 24). The compounds span a large range of activity (K_i values ranging from more than

10 μM to 0.0002 μM), as seen in Table 1, and are typical of the structural modifications

performed in hit-to-lead programs. The 15 ligands were first docked into the ACK1 ATP-

binding site, and a set of docked poses obtained for each ligand was re-scored with a 4-step

minimization protocol followed by a single-snapshot MM/PBSA re-scoring. The best scored

pose was alchemically studied and the relative binding energy was compared to the

experimental one. The alchemical calculations were also repeated with a 10-fold increase in

sampling time. The role of a possible bridging water molecule in the binding pocket was also

considered. Finally, thermodynamic cycle closures were analyzed as a way to detect incorrectly

predicted poses without knowledge of the experimental relative binding energies.

Table 1. Ligands studied in this work, along with reported K_i values(22). Compound numbering is the same as that used in reference 22.

	A	B	C	D	E	F	G	J	K_i (µM)[22]	Compound
Batch 1	O	C		N	C	C	O	H	0.01	**2**
	O	C		N	C	C	S	H	> 10	**3**
	O	C		N	C	N	N	H	7.3	**4**
	O	C		N	N	C	N	H	1.8	**5**
	O	C		C	C	C	O	H	0.07	**6**
	O	C		N	C	C	NH	H	0.006	**7**
Linker	O	C		N	C	C	O	OCH$_3$	0.005	**8**
Batch 2	O	C		N	C	C	O	O(CH$_2$)$_2$NMe$_2$	0.005	**15**
	O	C		N	C	C	NH	O(CH$_2$)$_2$NMe$_2$	0.006	**16**
	S	S		N	C	C	NH	O(CH$_2$)$_2$NMe$_2$	0.0003	**35**
	S	S		N	C	C	O	O(CH$_2$)$_2$NMe$_2$	0.0002	**36**
	S	C		N	C	C	O	O(CH$_2$)$_2$NMe$_2$	0.08	**38**
	O	O		N	C	C	O	O(CH$_2$)$_2$NMe$_2$	0.013	**39**
	C	C	OH	N	C	C	NH	O(CH$_2$)$_2$NMe$_2$	0.04	**44**
	C	C	OMe	N	C	C	NH	O(CH$_2$)$_2$NMe$_2$	0.05	**45**

MATERIALS AND METHODS

Dataset

The dataset consists of 15 ACK1 competitive inhibitors for which inhibition constants (K_i) have been reported. The structure of only one protein-ligand complex (compound **35**) was determined by X-ray crystallography(22) (Fig 1a). This dataset was further divided into two subsets: *batch 1* (6 ligands with K_i values ranging from >10 µM to 0.006 µM), and *batch 2* (9 ligands with K_i values ranging from 0.013 to 0.0002 µM).

Fig 1. (a-d) Superimposition of the X-ray diffraction derived structure of the ACK1 protein co-crystallized with ligand 35 (grey) (PDB code 4EWH). (a) with the best predicted MM/PBSA docked pose for ligand **6** (blue), (b) with ligand **7** (purple) exhibiting a different binding mode, (c) with ligand **44** from MM/PBSA prediction using the best predicted binding mode and (d) with ligand **44** using the second-best binding mode prediction. All carbon atoms of ligand **44** are colored in red. Hydrogens are omitted for clarity.

Protein and ligand setup

The ACK1 kinase domain structure was taken from the Protein Data Bank, code 4EWH (22), using chain B of the crystal structure, which was protonated with MOE v2009.1 (25). The structure has no missing residues; Tyrosine 284 was dephosphorylated with MOE following Lougheed *et al.* observation that inhibitor binding is not expected to be sensitive to the phosphorylation state of this residue (26). The protonation state of each ligand was predicted using the SDwash program in MOE v2009.1.

Docking

Docking was performed with MOE v2009.1 (25). The full docking process was done in three steps. The first one was an exhaustive conformational search of the ligands using the *Systematic* option of MOE together with the option *Enforce chair conformations* on. All other parameters were set to the standard options. A maximum of 100 conformations by compound were selected for the *Placement* step. In the second step the receptor was defined as those atoms within 9.0 Å from the ligand. The *Rotate Bonds* option was activated and the *Affinity dG function* employed together with the *Triangle Matcher* method for placement. A maximum of 30 poses for each ligand were retained. Finally, the 500 best structures were submitted to the *Refinement* step with the *Force Field* function and allowing the lateral chains of the pocket residues to move during the optimization without restriction. All other parameters were set to the standard options. The five best structures obtained for each ligand, according to their predicted binding energies, were retained for minimization and re-scoring with MM/PBSA.

MM/PBSA

A four-step minimization protocol followed by a single snapshot MM/PBSA re-scoring was performed with Amber 14 (27). Ligands were prepared with Antechamber using the GAFF force field (28) with AM1-BCC partial charges (29, 30), while the ff99SB (31) force field was used for the protein. All systems were solvated in a rectangular box of TIP3P water molecules (32). Counterions were added as necessary to neutralize the systems (33). Energy minimization

was performed under periodic boundary conditions using the particle-mesh-Ewald method for

the treatment of the long-range electrostatic interactions (34). A cut-off distance of 10 Å was

chosen to compute non-bonded interactions. The four-step minimization procedure was as

follows: 1) 5000 steepest descent (SD) steps applied to water molecule coordinates only; 2)

5000 SD steps applied also to protein atoms, with positional harmonic restraints (5 kcal mol^{-1} Å^{-}

^{2}) applied to backbone atoms only; 3) 5000 SD steps as done previously with backbone atom

restraints set to 1 kcal mol^{-1} Å^{-2} and 4) 5000 SD steps with no restraints.

For each of the energy minimized structures, a binding free energy was estimated following the

MM/PBSA method using the MM/PBSA.py program (35). No entropic contributions were

taken into account, while the variables *cavity_surften* and *cavity_offset* were assigned the values

of 0.00542 kcal mol^{-2} Å^{-2} and -1.008, respectively, using the defaults for all remaining variables.

Alchemical free energy calculations

Relative binding free energies were calculated using a single topology molecular dynamics

alchemical free energy approach (36). Alchemical free energy calculations avoid direct

computation of the free energy change associated with the reversible binding of a ligand to a

protein through an artificial morphing of a ligand *X* into another ligand *Y* by using a parameter λ

which defines the change from *X* to *Y*. Thus, the relative free energy of binding ($\Delta\Delta G_{X \to Y}$) was

given by equation 1 as:

$$\Delta\Delta G_{X \to Y} = \Delta G_{X \to Y}^{complex} - \Delta G_{X \to Y}^{free} \ (1)$$

Where $\Delta G_{X \to Y}^{free}$ is the free energy change for transforming ligand *X* into ligand *Y* in solution

whereas $\Delta G_{x \to y}^{complex}$ is the free energy change for the same transformation in the protein binding

site. A relative free energy perturbation network for both *batch 1* and *batch 2* was designed (S1

Fig and S11 Fig). The top-scored MM/PBSA pose for each ACK1 ligand was used as input for

the subsequent alchemical free energy preparation protocol using the FESetup software package

(37). The protocol used by FESetup for the automated preparation of ligands, protein and

complexes was as follows:

Ligands. Atomic charges were assigned by using the Antechamber module in AmberTools 14

(27), selecting the AM1-BCC method (29, 30), and the GAFF2 force field (28). Ligands were

solvated with TIP3P water molecules (32), with counterions added as necessary to neutralize the

system (33). Each system was energy minimized for 100 SD cycles and equilibrated at 300 K

and 1 atm pressure for 10^5 molecular dynamics (MD) steps with a 2 fs timestep using the

module Sander (27), with a positional harmonic restraint (10 kcal mol^{-1} Å^{-2}) applied to ligand

atoms. Bonds involving hydrogen atoms were constrained.

Protein. The protein was parametrized using the Amber ff14SB force field (38).

Complexes. Each ligand was combined back with the ACK1 protein model and the complex

was solvated with TIP3P water molecules (32). Counterions were also added to neutralize the

solution (33). The system was afterwards equilibrated following the procedure already

described for ligands, using now 5000 MD steps.

All alchemical free energy calculations used 11 equidistant λ windows. For each λ value MD

trajectories were computed in the NPT ensemble with a pressure of 1 atm and temperature of

300 K using the software SOMD 2016.1.0 (20, 39). SOMD has been used in several recent

studies to model the binding energetics of enzyme inhibitors (21), carbohydrate ligands (19),

and host-guest systems (13). Each λ window was sampled for 4 ns. Pressure was regulated

using a Monte Carlo barostat (40, 41) with an update frequency of 25 MD steps. Temperature

was kept constant using the Andersen thermostat (42), with a collision frequency of 10 ps^{-1}. A 2

fs time step was used with the leapfrog-Verlet integrator. All bonds involving hydrogens were

constrained to their equilibrium distances. Non-bonded interactions were evaluated setting a

cut-off distance of 12 Å. Long-range electrostatic interactions were calculated using the shifted

atom-based Barker-Watts reaction field (43), with the medium dielectric constant set to 82.0. In

order to avoid steric clashes at the beginning of each MD run due to modifications of the ligand

parameters associated with changes in λ, each structure was energy minimized for 1000 steps

prior to MD simulation.

Each simulation was repeated at least twice using different initial assignments of velocities, and

both $\Delta\Delta G_{X \to Y}$ and $\Delta\Delta G_{Y \to X}$ were calculated from independent simulations. In some cases, when

poor agreement was observed between duplicates a third run was performed.

Ligand **38** was tested as a racemic mixture for consistency with the experimental conditions.

Calculations were carried out for each enantiomer and the binding energies relative to this

ligand were given with equation 2:

$$\Delta\Delta G_{\mathbf{38} \to X} = -kT \ln\left[0.5\left(\exp\left(\frac{-\Delta\Delta G_{38R \to X}}{kT}\right) + \exp\left(\frac{-\Delta\Delta G_{38S \to X}}{kT}\right)\right)\right] (2)$$

Cycle closures were evaluated using free energy changes from both the forward (**X→Y**) and

reverse (**Y→X**) perturbations. The metrics used to evaluate the datasets were the determination

coefficient R^2, linear regression slope and the mean unsigned error (MUE). Experimental

binding affinities were calculated from the corresponding inhibition constants (22) (K_i) using

$\Delta G = RT \ln\left(\frac{K_i}{C^0}\right)$ with $C^0 = 1$ mol L^{-1}. As no uncertainties have been reported for the K_i values,

an uncertainty of 0.4 kcal mol^{-1} was assumed (44, 45).

Relative free energies were estimated using the multistate Bennett's acceptance ratio (MBAR)

method(46), as included in the software *analyse_freenrg* from the Sire software suite. Relative

free energies for complete datasets were evaluated using version 0.3.5 of the *freenrgworkflows*

python module [https://github.com/michellab/freenrgworkflows], using ligand 3 as a reference,

for which a $K_i = 10$ μM is used. For more details, see Mey *et al.* (47). All analysis scripts are

available online at https://github.com/michellab/ACK1_Data.

Alchemical free energy Protocols

Five different alchemical free energy protocols were followed. *Protocol A* uses for each ligand

the best scored pose according to MM/PBSA. This leads to a pose that differs from the X-ray

crystallographic pose of **35** for several ligands (**2**, **4**, **7**, **8**, **16**, **44** and **45**). *Protocol B* assumes

user intervention to select the pose that resembles the most the X-ray binding mode among the 5

MM/PBSA poses. *Protocols C* and *D* explore the effect of manually modelling a water

molecule inside the ACK1 ATP-binding site (see S6 Fig). This reflects user knowledge that in

other high-resolution structures of ACK1 (e.g. the 1.31 Å resolution 4HZR structure (48)) one

additional binding site water molecule between the protein and ligand is apparent. *Protocol C*

uses the same ligand poses as *Protocol A*, while *Protocol D* uses the same poses as *Protocol B*.

Finally, *Protocol E* is simply *Protocol A* with the per λ simulation time increased ten-fold. This

was done to evaluate whether the different binding mode and ATP-binding site water

rearrangements seen in *Protocols A-D* can be sampled with longer MD simulation protocols.

Protocol E is computationally expensive and was applied to *batch 1* only (ca. 10 μs of

simulation time). Figs were rendered with VMD (49), while graphs were prepared with Origin

(50) and python using plotting libraries Matplotlib version 2.0.2 (51) and Seaborn version 0.7.1

(52).

RESULTS

Batch 1

Protocol A renders (Table 2) modest results, with a R^2 of 0.36±0.07 and a strong

underestimation of relative free energies, as shown by the slope of the regression line (0.3).

Inspection of Fig 2a shows that ligands **2**, **4** and **7** are clear outliers. These ligands have a

predicted docked pose which differ more from the X-ray derived binding mode (see Figs 1b and

1c). Results for *protocol B* are shown in Table 2, and S2 and S5 Figs. This protocol gives

clearly better results, although the underestimation (slope 0.4) of relative binding free energies

remains high, and ligands **2**, **4** and **7** are still ranked poorly.

Fig 2. (a) and (b) Comparison of experimental and predicted relative free energies of

binding of *batch 1* for *protocols A* and *D*, respectively. Free energies of binding are relative to

ligand **3**. The linear regression line (dashed line) and a line with slope unity (solid line) are also

presented.

Table 2. R^2, MUE (kcal mol^{-1}) and slope metrics obtained from the comparison of experimental and predicted relative free energies of binding of *batch 1*.

Protocol	R^2	MUE	Slope
A	0.36±0.07	1.55±0.06	0.3
B	0.67±0.03	1.29±0.03	0.4
C	0.5±0.2	1.53±0.07	0.2
D	0.84±0.03	1.29±0.03	0.5
E	0.33±0.08	1.73±0.08	0.3

An analysis of the relative binding energies calculated with *protocols A* and *B* (S1 and S2 Figs), for ligands **2**, **4** and **7**, reveals that these ligands appear in the perturbations that show the highest deviations between the experimental and calculated relative binding energies. Thus, for *protocol A* deviations of more than 3.0 kcal mol^{-1} are observed for **2→3**, **7→4**, **7→6**, **7→3** and **3→7**, while for *protocol B* these deviations appear for perturbations **2→3**, **4→7**, **7→4**, **7→3** and **3→7**. An analysis of the docked structures of ligands **6** and **7** suggested that a possible explanation for the inability of the protocol to reproduce the experimental relative binding affinities is due to interactions of the extra nitrogen atom in the pyrimidine ring of ligand **7** that is missing in the pyridine ring of ligand **6** (see Fig 3). The extra N atom in the pyrimidine ring could establish a hydrogen bond with THR205 (see S6 Fig) if a bridging water was present. Indeed, several water molecules are present inside the ATP-binding pocket of 4HZR (48). That possibility was explored in *protocols C* and *D*, where a water molecule was manually placed inside the binding pocket between the nitrogen in position D of ligand **7** (see Table 1) and THR205. The final position of the water molecule is obtained after 100 steps of SD minimization fixing all other atoms. Results for *protocol C* are shown in Table 2 and S7 Fig, while those for *protocol D* appear in Table 2 and Figs 2b and 3. *Protocol D* clearly surpass all

others, with a R^2 of 0.84±0.03 and an improvement in the underestimation of relative binding

energies (slope = 0.5). A comparison of the calculated relative binding energies for ligands **3**

and **4** allows to conclude that using a different pose for ligand **4** does not seem to affect the

results (both *protocols A* and *B* for example, give an average $\Delta\Delta G_{3\rightarrow4}$ of 1.3 kcal mol^{-1}).

Inspection of the calculated trajectories show that ligand **4** rapidly converts from its initial

docked pose (*protocols A* and *C*) to one similar to that used as input for *protocols B* and *D*.

Fig 3. Calculated and experimental (in bold) relative binding affinities (in kcal mol^{-1}) for

all the perturbations run in *batch 1* with *protocol D*. The calculated values correspond to

independent repeats.

The possibility of resolving ambiguities in binding poses and binding site water content without

user intervention was next tested by increasing the simulation sampling time to 40 ns for each λ

window. The expectation was this would allow the ligand to find the correct pose and to allow

water molecules diffuse in the ATP-binding site (see S6 Fig). Results are shown in Table 2 and

S8 Fig. The increased simulation time does not translate into any improvement of the results.

The R^2, slope and MUE values are as poor or poorer as those for *protocol A*, while the outliers

remain the same. The MD trajectories show that, even with the increased simulation time,

ligand **7** is not able to change its docking pose, while ligand **4** needs under 4 ns to adopt a pose

that resembles the X-ray pose of **35**. Besides, a water molecule enters and remains in the ATP-

binding site in 7 out of 22 MD trajectories only.

Analysis of the complete dataset

The robustness of the results obtained for *batch 1* was tested by processing *batch 2* and re-

analyzing the full dataset. Ligands in *batch 2* are positively charged in the assay conditions,

whereas *batch 1* ligands are neutral. Relative free energy calculations that involve a net charge

change are still technically challenging for simulations carried out with a reaction-field cut-off.

Thus, the perturbations between ligands **8** and **15** were carried out assuming **15** is neutral.

Results for individual perturbations in *batch 2* are shown in S11 to S15 Figs.

Protocol A, as expected given the results obtained for *batch 1*, gives modest results, as can be

seen in Table 3 and Fig 4a (R^2 =0.45±0.06 and slope of 0.5). The slope has improved from 0.3

to 0.5 because the relative free energies of compounds in *batch 2* are not as under predicted as

those from *batch 1* (see S1 Table). Ligands **16**, **44** and **45** need further inspection. S11 Fig

shows that, while the experimental $\Delta\Delta G_{45\to44}$ is -0.1 kcal mol^{-1}, the calculated $\Delta\Delta G_{45\to44}$ are

1.8/1.5 (run 1/run 2) kcal mol^{-1} (the reverse perturbation was calculated as -2.0/-1.9 kcal mol^{-1}).

Similarly, while the experimental $\Delta\Delta G_{16\to45}$ is 1.2 kcal mol^{-1}, the calculated results are $\Delta\Delta G_{16\to45}$

-0.8/-1.6 (run 1/run 2) kcal mol^{-1} and $\Delta\Delta G_{45\to16}$ -2.2/-2.4 /run 1/run 2) kcal mol^{-1}.

Fig 4. (a) and (b) Comparison of experimental and predicted relative free energies of

binding of the whole set for *protocols A* and *D*, respectively. Free energies of binding are

relative to ligand **3**. The linear regression line (dashed line) and a line with slope 1 (solid line)

are also presented.

Table 3. R^2, MUE (kcal mol^{-1}) and slope metrics obtained from the comparison of

experimental and predicted relative free energies of binding of the whole set.

Protocol	R^2	MUE	Slope
A	0.45±0.06	2.11±0.08	0.5
B	0.59±0.04	2.12±0.04	0.5
C	0.55±0.05	1.82±0.06	0.5
D	0.76±0.02	1.24±0.04	0.8

Interestingly, the dihedral angle defining the relative orientation of the NH group that links the

pyrimidine and the cyclopentanol rings changes values rather quickly during the simulation. Fig

5a shows an example for the first repeat of the perturbation **44→45** at λ=0. For the simulations

involving ligand **44** an intramolecular H-bond between its aniline NH group and its cyclopentyl

hydroxyl group is established (see Fig 5c). That conformation is precisely the second-best

MM/PBSA docked one (see Fig 1c), which features that intramolecular hydrogen bond. Thus,

batch 2 protocol B includes the second-best scored MM/PBSA poses for ligands **8**, **16** and **44**.

In the case of ligands **8** and **16**, this implies using a pose that resembles the most the X-ray

binding mode, while for ligand **44** the second-best scored MM/PBSA pose differs from the best-

scored one in the aniline NH dihedral angle (see Fig 1c). The improvement, as shown in Table 3

and S9 Fig, for *protocol B* as compared with *protocol A*, is quite modest. Results are clearly

better for the **16→45** and **45→16** perturbations, with the disagreement between experimental

and calculated relative binding energy decreasing from 3.5 to 0.3 kcal mol^{-1} (compare S11 and

S12 Figs), but ligand **44** is still an outlier. Although the experimental relative binding energy for

the **45 → 44** perturbation is just -0.1 kcal mol^{-1}, ligand **45** is predicted to bind much more

strongly to ACK1 (calculated $\Delta\Delta G_{45\to 44}$ are –2.0/-1.9 and -1.2/-1.6 kcal mol^{-1} for *protocols A*

and *B*, respectively) than **44**. This suggests possible deficiencies in the force field used for **44** in

this study.

Fig 5. (a) 4 ns trajectory monitoring dihedral angle of ligand 44 (blue circles) and 45

(purple crosses) as indicated in (b) and (c) as well as probability distribution of dihedrals

over the trajectory. (b) Snapshot of the conformation of ligand 44 taken from a λ = 0

trajectory at t=0 ns indicating dihedral conformation monitored in (a) highlighted by

spheres. (c) Snapshot of the conformation of ligand 44 taken from the same trajectory

after 3 ns, showing an intramolecular hydrogen bond.

Protocols C and D, follow the same trends already explained for *batch 1*, pointing to an

improvement in the results when a water molecule is included in the ATP-binding pocket (Table

3). An encouraging R^2 of 0.76 ± 0.02 and an improvement in the underestimation of relative

binding energies (slope 0.8) is obtained, though there is still room for improvements for affinity

predictions for **44** and **16**.

Thermodynamic cycle closures analysis

Hysteresis, being defined as the difference in binding energy between the forward and reverse

perturbation (39, 53, 54), has been proposed as useful metric to identify problematic

perturbations (55, 56). Cycle closures for both *batch 1* and *batch 2* were computed to determine

whether incorrectly predicted binding poses could be detected in the absence of experimental

binding affinity data. Results are shown in Table 4.

Table 4. Calculated thermodynamic cycle closures. Cycle closures that exceed or equal a

threshold of 0.8 kcal mol^{-1} are highlighted in bold.

		Cycle closure (kcal mol^{-1})				
	Protocol	A	B	C	D	E
Cycle	3-2-5-6	0.6 ± 0.3	0.0 ± 0.3	0.6 ± 0.6	0.2 ± 0.4	**0.8 ± 0.3**
	3-2-5	0.2 ± 0.2	-0.4 ± 0.2	0.7 ± 0.5	0.0 ± 0.1	0.5 ± 0.3
	2-6-5	**-0.8 ± 0.4**	0.0 ± 0.3	0.0 ± 0.3	-0.2 ± 0.3	-0.7 ± 0.8
	3-5-6	0.4 ± 0.2	0.4 ± 0.2	-0.1 ± 0.4	0.2 ± 0.4	0.4 ± 0.1
	3-6-2	0.2 ± 0.4	0.1 ± 0.3	-0.6 ± 0.5	0.0 ± 0.2	-0.1 ± 0.9
	3-4-7-6	**1.0 ± 0.4**	0.3 ± 0.4	**1.6 ± 0.4**	0.2 ± 0.4	0.3 ± 0.7
	3-7-6	0.2 ± 0.4	0.1 ± 0.3	**0.9 ± 0.7**	0.2 ± 0.3	-0.2 ± 0.9
	4-6-7	**0.9 ± 0.6**	0.0 ± 0.3	**-1.1 ± 0.5**	0.2 ± 0.3	-0.2 ± 0.8
	3-4-7	**0.8 ± 0.4**	0.2 ± 0.4	0.7 ± 0.7	0.0 ± 0.2	0.5 ± 0.5
	3-4-6	**1.9 ± 0.4**	0.3 ± 0.2	0.5 ± 0.5	0.2 ± 0.4	0.1 ± 0.4
	45-16-44	**-2 ± 1**	**-3 ± 1**	**-2 ± 1**	**-1.8 ± 0.9**	N/A
	38-39-35-36	0.6 ± 0.5	0.6 ± 0.5	0.2 ± 0.3	0.2 ± 0.3	N/A

As could be expected, similar conclusions can be obtained when analyzing ring cycle closures

or comparing forward and reverse perturbations, although there are some cases with high

deviations between experimental and calculated relative binding energies, while exhibiting

almost null hysteresis for the forward and reverse perturbations (i.e. the perturbations between

ligands **2** and **5** in *batch 1* and those between ligands **44** and **45** in *batch 2*).

Overall it appears that a threshold of $\pm$ 0.8 kcal mol^{-1} for cycle closure errors is useful to flag

poses that need further attention even without prior knowledge of the experimental binding

affinities. Thus, for *protocol A*, **3**-**4**-**7**-**6**, **3**-**4**-**6**, **3**-**4**-**7**, **4**-**6**-**7**, **2**-**6**-**5** and **45**-**16**-**44** thermodynamic

cycle closures are indicative of problematic ligands. According to this metric, a significant

improvement when using *protocol B* (only one thermodynamic cycle closure above the

threshold) is seen, while a comparison between *protocols A* (6 cycles with hysteresis above the

threshold) and *C* (4 cycles) suggest a modest improvement. Results for *batch 2* clearly indicate

that ligands **44**, **45** and **16** (hysteresis of -2 $\pm$ 1 kcal mol^{-1} in their thermodynamic cycle for

protocol A) are much more problematic than ligands **35**, **36**, **38** and **39** (hysteresis of 0.2 $\pm$ 0.3

kcal mol^{-1} for *protocol A*). The best performing *protocol D* is unable to improve the hysteresis

for the **45**-**16**-**44** thermodynamic cycle.

DISCUSSION

This work has explored the viability of using alchemical free energy methods as a final filter in

a cascade of computational methods for hit-to-lead virtual screens in the context of a dataset of

ACK1 inhibitors. The two major limitations of AFE methods are the quality of the potential

energy function used, and the extent to which the configurational sampling performed has

captured relevant protein-ligand conformations (55, 56) In principle sufficient long simulations

will relax a protein-ligand complex to the ligand pose and protein conformation preferred by the

force field used. However, because computing time is limited in practical scenarios, AFE

simulations typically afford only a few ns per window, which can make the calculated binding

affinities sensitive to the selection of the starting conformations. This work indicates that use of

experimental data to bias the selection of poses and setup of binding site water content could

lead to significant performance improvements. While the dataset studied here is small, the

lessons from this study are likely applicable to other binding sites. Of course, as illustrated with

ligand **4**, even in cases where the MD simulations relax a previously modelled binding pose to

one that closely resembles a pose inferred from X-ray data, the free energy calculations may

still fail to reproduce the experimental binding affinities.

Careful analysis of literature structural data (48, 57, 58) was key to identify a conserved

hydration site that was not modelled in the prior docking calculations. This knowledge was

important to realize upon inspection of putative poses for some ligands in *batch 1* the feasibility

of a hydrogen bonding interaction via a bridging water molecule. Gratifyingly modelling of this

hydration site leads to significant accuracy improvements for several perturbations.

In principle, assuming an accurate potential energy function, these sampling issues could be

dealt with by simply increasing the sampling time of the MD simulations. For the present

dataset, we find that a one order of magnitude increase in sampling time was insufficient to

bring about improvements in binding poses accuracy and binding site water content. Thus, at

present it seems wise to pay attention to the starting conditions of the free energy calculations to

maximize cost effectiveness. Where experimental data is lacking, a number of molecular

modelling protocols have been proposed to determine location and energetics of important

binding site water molecules (59-68).

ACKNOWLEDGMENTS

JM acknowledges support from a University Research Fellowship from the Royal Society.

REFERENCES

1.   PhRMA. Fact Sheet "Drug Discovery and Development. Understanding theR&D
process" 2017 [Available from: http://www.phrma.org/graphic/four-facts-about-
spending-on-prescription-medicines.
2.   Whitesides GM, Krishnamurthy VM. Designing ligands to bind proteins. Q Rev
Biophys. 2006;38(4):385-95.
3.   Smith AJT, Zhang X, Leach AG, Houk KN. Beyond Picomolar Affinities:
Quantitative Aspects of Noncovalent and Covalent Binding of Drugs to Proteins. J Med
Chem. 2009;52(2):225-33.
4.   Kollman PA, Massova I, Reyes C, Kuhn B, Huo S, Chong L, et al. Calculating
Structures and Free Energies of Complex Molecules: Combining Molecular Mechanics
and Continuum Models. Acc Chem Res. 2000;33(12):889-97.
5.   Aqvist J, Marelius J. The linear interaction energy method for predicting ligand
binding free energies. Comb Chem High Throughput Screen. 2001;4(8):613-26.
6.   Zwanzig RW. High-Temperature Equation of State by a Perturbation Method. I.
Nonpolar Gases. J Chem Phys. 1954;22(8):1420-6.
7.   Michel J, Foloppe N, Essex JW. Rigorous Free Energy Calculations in Structure-
Based Drug Design. Mol Inform. 2010;29(8-9):570-8.

8.   Granadino-Roldan JM, Garzon A, Gomez-Gutierrez P, Pasamontes-Funez I,
Santos Tomas M, Rubio-Martinez J. A multistep docking and scoring protocol for
congeneric series: Implementation on kinase DFG-out type II inhibitors. Future Med
Chem. 2018;10(3):297-318.
9.   Yin J, Henriksen NM, Slochower DR, Shirts MR, Chiu MW, Mobley DL, et al.
Overview of the SAMPL5 host–guest challenge: Are we doing better? J Comput Aided
Mol Des. 2017;31(1):1-19.
10.   Muddana HS, Fenley AT, Mobley DL, Gilson MK. The SAMPL4 host-guest blind
prediction challenge: an overview. J Comput Aided Mol Des. 2014;28(4):305-17.
11.   Deng N, Flynn WF, Xia J, Vijayan RS, Zhang B, He P, et al. Large scale free energy
calculations for blind predictions of protein-ligand binding: the D3R Grand Challenge
2015. J Comput Aided Mol Des. 2016;30(9):743-51.
12.   Gaieb Z, Liu S, Gathiaka S, Chiu M, Yang H, Shao C, et al. D3R Grand Challenge 2:
blind prediction of protein-ligand poses, affinity rankings, and relative binding free
energies. J Comput Aided Mol Des. 2018;32(1):1-20.
13.   Bosisio S, Mey A, Michel J. Blinded predictions of host-guest standard free
energies of binding in the SAMPL5 challenge. J Comput Aided Mol Des. 2017;31(1):61-
70.
14.   Mey A, Juarez-Jimenez J, Hennessy A, Michel J. Blinded predictions of binding
modes and energies of HSP90-alpha ligands for the 2015 D3R grand challenge. Bioorg
Med Chem. 2016;24(20):4890-9.
15.   Athanasiou C, Vasilakaki S, Dellis D, Cournia Z. Using physics-based pose
predictions and free energy perturbation calculations to predict binding poses and
relative binding affinities for FXR ligands in the D3R Grand Challenge 2. J Comput Aided
Mol Des. 2018;32(1):21-44.
16.   Wang L, Wu Y, Deng Y, Kim B, Pierce L, Krilov G, et al. Accurate and reliable
prediction of relative ligand binding potency in prospective drug discovery by way of a
modern free-energy calculation protocol and force field. J Am Chem Soc.
2015;137(7):2695-703.
17.   Shoichet BK, Walters WP, Jiang H, Bajorath J. Advances in Computational
Medicinal Chemistry: A Reflection on the Evolution of the Field and Perspective Going
Forward. J Med Chem. 2016;59(9):4033-4.
18.   Steinbrecher TB, Dahlgren M, Cappel D, Lin T, Wang LL, Krilov G, et al. Accurate
Binding Free Energy Predictions in Fragment Optimization. J Chem Inf Mod.
2015;55(11):2411-20.
19.   Mishra SK, Calabró G, Loeffler HH, Michel J, Koča J. Evaluation of Selected
Classical Force Fields for Alchemical Binding Free Energy Calculations of Protein-
Carbohydrate Complexes. Journal of Chemical Theory and Computation.
2015;11(7):3333-45.
20.   Mey ASJS, Juárez-Jiménez J, Hennessy A, Michel J. Blinded predictions of
binding modes and energies of HSP90-α ligands for the 2015 D3R grand challenge.
Biorg Med Chem. 2016;24(20):4890-9.
21.   Mey ASJS, Jiménez JJ, Michel J. Impact of domain knowledge on blinded
predictions of binding energies by alchemical free energy calculations. J Comput Aided
Mol Des. 2018;32(1):199-210.

22. Jiao X, Kopecky DJ, Liu J, Liu J, Jaen JC, Cardozo MG, et al. Synthesis and
optimization of substituted furo[2,3-d]-pyrimidin-4-amines and 7H-pyrrolo[2,3-
d]pyrimidin-4-amines as ACK1 inhibitors. Bioorg Med Chem Lett. 2012;22(19):6212-7.
23. Mahajan K, Mahajan NP. Shepherding AKT and androgen receptor by Ack1
tyrosine kinase. J Cell Physiol. 2010;224(2):327-33.
24. Chua BT, Lim SJ, Tham SC, Poh WJ, Ullrich A. Somatic mutation in the ACK1
ubiquitin association domain enhances oncogenic signaling through EGFR regulation in
renal cancer derived cells. Mol Oncol. 2010;4(4):323-34.
25. Molecular Operating Environment (MOE) 2009.1. Chemical Computing Group
Inc., 1010 Sherboke St. West, Suite #90, Montreal, QC, Canada, H3A 2R7, 2013.
2013.08 ed.
26. Lougheed JC, Chen RH, Mak P, Stout TJ. Crystal structures of the
phosphorylated and unphosphorylated kinase domains of the Cdc42-associated
tyrosine kinase ACK1. J Biol Chem. 2004;279(42):44039-45.
27. Case DA, Babin V, Berryman JT, Betz RM, Cai Q, Cerutti DS, et al. AMBER 14.
University of California, San Francisco: University of California, San Francisco; 2014
2014.
28. Wang J, Wolf RM, Caldwell JW, Kollman PA, Case DA. Development and testing
of a general amber force field. J Comput Chem. 2004;25(9):1157-74.
29. Jakalian A, Bush BL, Jack DB, Bayly CI. Fast, efficient generation of high-quality
atomic charges. AM1-BCC model: I. Method. J Comput Chem. 2000;21(2):132-46.
30. Jakalian A, Jack DB, Bayly CI. Fast, efficient generation of high-quality atomic
charges. AM1-BCC model: II. Parameterization and validation. J Comput Chem.
2002;23(16):1623-41.
31. Hornak V, Abel R, Okur A, Strockbine B, Roitberg A, Simmerling C. Comparison
of multiple Amber force fields and development of improved protein backbone
parameters. Proteins: Struct Funct Bioinform. 2006;65(3):712-25.
32. Jorgensen WL, Chandrasekhar J, Madura JD, Impey RW, Klein ML. Comparison
of simple potential functions for simulating liquid water. J Chem Phys. 1983;79(2):926-
35.
33. Åqvist J. Ion-water interaction potentials derived from free energy perturbation
simulations. The Journal of Physical Chemistry. 1990;94(21):8021-4.
34. Darden T, York D, Pedersen L. Particle mesh Ewald: An N log(N) method for
Ewald sums in large systems. J Chem Phys. 1993;98(12):10089-92.
35. Miller BR, McGee TD, Swails JM, Homeyer N, Gohlke H, Roitberg AE.
MMPBSA.py: An Efficient Program for End-State Free Energy Calculations. Journal of
Chemical Theory and Computation. 2012;8(9):3314-21.
36. Michel J, Essex JW. Prediction of protein-ligand binding affinity by free energy
simulations: assumptions, pitfalls and expectations. J Comput Aided Mol Des.
2010;24(8):639-58.
37. Loeffler HH, Michel J, Woods C. FESetup: Automating Setup for Alchemical Free
Energy Simulations. J Chem Inf Model. 2015;55(12):2485-90.
38. Maier JA, Martinez C, Kasavajhala K, Wickstrom L, Hauser KE, Simmerling C.
ff14SB: Improving the Accuracy of Protein Side Chain and Backbone Parameters from
ff99SB. Journal of Chemical Theory and Computation. 2015;11(8):3696-713.

39.   Calabro G, Woods CJ, Powlesland F, Mey AS, Mulholland AJ, Michel J.
Elucidation of Nonadditive Effects in Protein-Ligand Binding Energies: Thrombin as a
Case Study. J Phys Chem B. 2016;120(24):5340-50.
40.   Chow K-H, Ferguson DM. Isothermal-isobaric molecular dynamics simulations
with Monte Carlo volume sampling. Comput Phys Commun. 1995;91(1):283-9.
41.   Åqvist J, Wennerström P, Nervall M, Bjelic S, Brandsdal BO. Molecular dynamics
simulations of water and biomolecules with a Monte Carlo constant pressure
algorithm. Chem Phys Lett. 2004;384(4):288-94.
42.   Andersen HC. Molecular dynamics simulations at constant pressure and/or
temperature. J Chem Phys. 1980;72(4):2384-93.
43.   Barker JA, Watts RO. Monte Carlo studies of the dielectric properties of water-
like models. Mol Phys. 1973;26(3):789-92.
44.   Mikulskis P, Genheden S, Ryde U. A Large-Scale Test of Free-Energy Simulation
Estimates of Protein–Ligand Binding Affinities. J Chem Inf Mod. 2014;54(10):2794-806.
45.   Brown SP, Muchmore SW, Hajduk PJ. Healthy skepticism: assessing realistic
model performance. Drug Discovery Today. 2009;14(7):420-7.
46.   Shirts MR, Chodera JD. Statistically optimal analysis of samples from multiple
equilibrium states. J Chem Phys. 2008;129(12):124105.
47.   Mey A, Jimenez JJ, Michel J. Impact of domain knowledge on blinded
predictions of binding energies by alchemical free energy calculations. J Comput Aided
Mol Des. 2018;32(1):199-210.
48.   Gajiwala KS, Maegley K, Ferre R, He Y-A, Yu X. Ack1: Activation and Regulation
by Allostery. PLoS One. 2013;8(1):e53994.
49.   Humphrey W, Dalke A, Schulten K. VMD: visual molecular dynamics. J Mol
Graph. 1996;14(1):33-8, 27-8.
50.   Origin (OriginLab, Northampton, MA) OriginLab, Northampton, MA.
51.   Hunter JD. Matplotlib: A 2D graphics environment. Comput Sci Eng.
2007;9(3):90-5.
52.   https://github.com/mwaskom/seaborn   [
53.   König G, Hudson PS, Boresch S, Woodcock HL. Multiscale Free Energy
Simulations: An Efficient Method for Connecting Classical MD Simulations to QM or
QM/MM Free Energies Using Non-Boltzmann Bennett Reweighting Schemes. Journal
of Chemical Theory and Computation. 2014:140303161846003.
54.   Michel J, Tirado-Rives J, Jorgensen WL. Prediction of the water content in
protein binding sites. J Phys Chem B. 2009;113(40):13337-46.
55.   Cournia Z, Allen B, Sherman W. Relative Binding Free Energy Calculations in
Drug Discovery: Recent Advances and Practical Considerations. J Chem Inf Model.
2017;57(12):2911-37.
56.   Williams-Noonan BJ, Yuriev E, Chalmers DK. Free Energy Methods in Drug
Design: Prospects of "Alchemical Perturbation" in Medicinal Chemistry. J Med Chem.
2018;61(3):638-49.
57.   Kopecky DJ, Hao X, Chen Y, Fu J, Jiao X, Jaen JC, et al. Identification and
optimization of N3,N6-diaryl-1H-pyrazolo[3,4-d]pyrimidine-3,6-diamines as a novel
class of ACK1 inhibitors. Biorg Med Chem Lett. 2008;18(24):6352-6.
58.   Lougheed JC, Chen R-H, Mak P, Stout TJ. Crystal Structures of the
Phosphorylated and Unphosphorylated Kinase Domains of the Cdc42-associated
Tyrosine Kinase ACK1. J Biol Chem. 2004;279(42):44039-45.

59.   Luccarelli J, Michel J, Tirado-Rives J, Jorgensen WL. Effects of Water Placement
on Predictions of Binding Affinities for p38alpha MAP Kinase Inhibitors. J Chem Theory
Comput. 2010;6(12):3850-6.
60.   Bodnarchuk MS, Viner R, Michel J, Essex JW. Strategies to calculate water
binding free energies in protein-ligand complexes. J Chem Inf Model. 2014;54(6):1623-
33.
61.   Michel J, Henchman RH, Gerogiokas G, Southey MWY, Mazanetz MP, Law RJ.
Evaluation of Host–Guest Binding Thermodynamics of Model Cavities with Grid Cell
Theory. Journal of Chemical Theory and Computation. 2014;10(9):4055-68.
62.   Sindhikara Daniel J, Yoshida N, Hirata F. Placevent: An algorithm for prediction
of explicit solvent atom distribution—Application to HIV-1 protease and F-ATP
synthase. J Comput Chem. 2012;33(18):1536-43.
63.   Sridhar A, Ross GA, Biggin PC. Waterdock 2.0: Water placement prediction for
Holo-structures with a pymol plugin. PLoS One. 2017;12(2):e0172743.
64.   Hu B, Lill Markus A. WATsite: Hydration site prediction program with PyMOL
interface. J Comput Chem. 2014;35(16):1255-60.
65.   Ramsey S, Nguyen C, Salomon-Ferrer R, Walker RC, Gilson MK, Kurtzman T.
Solvation thermodynamic mapping of molecular surfaces in AmberTools: GIST. J
Comput Chem. 2016;37(21):2029-37.
66.   Lazaridis T. Inhomogeneous Fluid Approach to Solvation Thermodynamics. 1.
Theory. The Journal of Physical Chemistry B. 1998;102(18):3531-41.
67.   Lazaridis T. Inhomogeneous Fluid Approach to Solvation Thermodynamics. 2.
Applications to Simple Fluids. The Journal of Physical Chemistry B. 1998;102(18):3542-
50.
68.   Bayden AS, Moustakas DT, Joseph-McCarthy D, Lamb ML. Evaluating Free
Energies of Binding and Conservation of Crystallographic Waters Using SZMAP. J Chem
Inf Mod. 2015;55(8):1552-65.

SUPPORTING INFORMATION

S1 Fig. **Calculated and experimental (in bold) relative binding affinities (in kcal mol[-1]) for**

all the perturbations essayed in *batch 1*, *protocol A*. The drawing tries to reflect the different

conformations adopted by ligands **2**, **4** and **7**. The calculated values correspond to independent

repeats.

S2 Fig. **Calculated and experimental (in bold) relative binding affinities (in kcal mol[-1]) for**

all the perturbations essayed in *batch 1*, *protocol B*. The calculated values correspond to

independent repeats.

S3 Fig. **Calculated and experimental (in bold) relative binding affinities (in kcal mol[-1]) for**

all the perturbations essayed in *batch 1*, *protocol C*. The drawing tries to reflect the different

conformations adopted by ligands **2**, **4** and **7**. The calculated values correspond to independent

repeats.

S4 Fig. **Calculated and experimental (in bold) relative binding affinities (in kcal mol^{-1}) for**

all the perturbations essayed in *batch 1*, **protocol E.** The drawing tries to reflect the different

conformations adopted by ligands **2**, **4** and **7**. The calculated values correspond to independent

repeats.

S5 Fig. **Comparison of experimental and predicted relative free energies of binding of**

batch 1 **for** *protocol B***.** Free energies of binding are relative to ligand **3**. The linear regression

line (dashed line) and a line with slope 1 (solid line) are also presented.

S6 Fig. **Snapshot taken after 2 ns of MD for the λ=0, run 1, 7→6 perturbation, showing**

the manually placed water molecule inside the ATP-binding pocket.

S7 Fig. **Comparison of experimental and predicted relative free energies of binding of**

batch 1 **for** *protocol C***.** Free energies of binding are relative to ligand **3**. The linear regression

line (dashed line) and a line with slope 1 (solid line) are also presented.

S8 Fig. **Comparison of experimental and predicted relative free energies of binding of**

batch 1 **for** *protocol E***.** Free energies of binding are relative to ligand **3**. The linear regression

line (dashed line) and a line with slope 1 (solid line) are also presented.

S9 Fig. **Comparison of experimental and predicted relative free energies of binding of the**

whole set for *protocol B***.** Free energies of binding are relative to ligand **3**. The linear regression

line (dashed line) and a line with slope 1 (solid line) are also presented.

S10 Fig. **Comparison of experimental and predicted relative free energies of binding of the**

whole set for *protocol C***.** Free energies of binding are relative to ligand **3**. The linear regression

line (dashed line) and a line with slope 1 (solid line) are also presented.

S11 Fig. **Calculated and experimental (in bold) relative binding affinities (in kcal mol^{-1}) for**

all the perturbations essayed in *batch 2***,** *protocol A***.** The drawing tries to reflect the different

conformations adopted by ligands **44** and **16**. The calculated values correspond to independent

repeats.

S12 Fig. **Calculated and experimental (in bold) relative binding affinities (in kcal mol^{-1}) for**

all the perturbations essayed in *batch 2*, *protocol B*. The calculated values correspond to

independent repeats.

S13 Fig. **Calculated and experimental (in bold) relative binding affinities (in kcal mol^{-1}) for**

all the perturbations essayed in *batch 2*, *protocol C*. The drawing tries to reflect the different

conformations adopted by ligands **44** and **16**. The calculated values correspond to independent

repeats.

S14 Fig. **Calculated and experimental (in bold) relative binding affinities (in kcal mol^{-1}) for**

all the perturbations essayed in *batch 2*, *protocol D*. The calculated values correspond to

independent repeats.

S15 Fig. **Calculated and experimental (in bold) relative binding affinities (in kcal mol^{-1}) for**

the perturbations used to link *batch 1* and *batch 2* with *protocols A*, *B*, *C* and *D*. The

calculated values correspond to independent repeats.

S1 Table. **R^2, MUE (kcal mol^{-1}) and slope obtained from the comparison of experimental**

and predicted relative free energies of binding of *batch 2*

A

B

C

2.9 Å