The maternal hormone in the male brain: sexually dimorphic distribution of prolactin signalling in

the mouse brain

Authors: Hugo Salais-López[1], Carmen Agustín-Pavón[1,2], Enrique Lanuza[2], Fernando Martínez-García[1]

1. Unitat Predepartamental de Medicina, Facultat de Ciències de la Salut, Universitat Jaume I,

Castelló de la Plana, Spain

2. Departament de Biologia Cel·lular i de Biologia Funcional, Facultat de Ciències Biològiques,

Universitat de València, València, Spain

Keywords: Neuroendocrinology; pSTAT5; Sexual dimorphism; testosterone; Sociosexual Brain

Network; Prolactin receptor

Author for correspondence/ reprints: Fernando Martínez-García, Facultat de Ciències de la Salut,

Unitat Predepartamental de Medicina, Universitat Jaume I; Av. de Vicent Sos Baynat, s/n; 12071

Castelló de la Plana, Spain. Phone: +34 964 387457. Email: femartin@uji.es

Funding

Funded by the Spanish Ministry of Economy and Competitiveness-FEDER (BFU2016-77691-C2-2-P

and C2-1-P) and the Generalitat Valenciana (PROMETEO/2016/076) and the *Universitat Jaume I de*

Castelló (UJI-B2016-45). H.S-L has been a predoctoral fellow of the FPU (*Formación de Profesorado*

Universitario) programme of the Spanish Ministry of Education, Culture and Sport.

ABSTRACT

Research of the central actions of prolactin (PRL) is virtually restricted to females, but this hormone

has also documented roles in male physiology and behaviour. Here, we provide the first description

of the pattern of PRL-derived signalling in the male mouse brain, employing the immunostaining of

phosphorylated signal transducer and activator of transcription 5 (pSTAT5) after exogenous PRL

administration. Next, we explore possible sexual dimorphism by comparing pSTAT5

immunoreactivity in PRL-supplemented males and females. We also assess the role of testosterone

in the regulation of central prolactin signalling in males by comparing intact with castrated PRL-

supplemented males.

Prolactin-supplemented males displayed a widespread pattern of pSTAT5 immunoreactivity,

restricted to brain centres showing expression of the PRL receptor. Immunoreactivity for pSTAT5

was present in several nuclei of the preoptic, anterior and tuberal hypothalamus, as well as in the

septofimbrial nucleus or posterodorsal medial amygdala of the telencephalon. Conversely, non-

supplemented control males were virtually devoid of pSTAT5-immunoreactivity, suggesting that

central PRL actions in males are limited to situations concurrent with substantial hypophyseal PRL

release (e.g. stress or mating). Furthermore, comparison of PRL-supplemented males and females

revealed a significant, female-biased sexual dimorphism, supporting the view that PRL has a

preeminent role in female physiology and behaviour. Finally, in males, castration significantly

reduced pSTAT5 immunoreactivity in some structures, including the paraventricular and

ventromedial hypothalamic nuclei and the septofimbrial region, thus indicating a region-specific

regulatory role of testosterone over central PRL signalling.

Précis

We map prolactin signalling in the male brain using pSTAT5-immunostaining after prolactin

administration. We show that prolactin signalling is both sexually dimorphic and modulated by

testosterone.

ABBREVIATIONS

3V	third ventricle
AAD	anterior amygdaloid area, dorsal part
AC/ADP	anterior commissural nucleus/anterodorsal preoptic nucleus region
ac	anterior commissure
Acb	nucleus accumbens
ACo	anterior cortical amygdaloid nucleus
AHi	amygdalohippocampal area
AHP	anterior hypothalamic area, posterior part
Aq	aqueduct (Sylvius)
Arc	arcuate hypothalamic nucleus
AVP	arginine-vasopressin
AVPe	anteroventral periventricular nucleus
Bar	Barrington´s nucleus
BLA	basolateral amygdaloid nucleus, anterior part
BLV	basolateral amygdaloid nucleus, ventral part
BMP	basomedial amygdaloid nucleus, posterior part
BSTIA	bed nucleus of the stria terminalis, intraamygdaloid division
BSTLD	bed nucleus of the stria terminalis, lateral division, dorsal part
BSTLP	bed nucleus of the stria terminalis, lateral division, posterior part
BSTLV	bed nucleus of the stria terminalis, lateral division, ventral part
BSTMA	bed nucleus of the stria terminalis, medial division, anterior part
BSTMV	bed nucleus of the stria terminalis, medial division, ventral part
BSTMPI	bed nucleus of the stria terminalis, medial division, posterointermediate part
BSTMPL	bed nucleus of the stria terminalis, medial division, posterolateral part
BSTMPM	bed nucleus of the stria terminalis, medial division, posteromedial part
cc	corpus callosum

1	Ce	central amygdala
2	CeC	central amygdaloid nucleus, capsular part
3	CeL	central amygdaloid nucleus, lateral division
4	CeM	central amygdaloid nucleus, medial division
5	CPu	caudate putamen (striatum)
6	CRF	corticotropin-releasing factor
7	CxA	cortex-amygdala transition zone
8	D3V	dorsal third ventricle
9	DEn	dorsal endopiriform nucleus
10	Dk	nucleus of Darkschewitsch
11	DLG	dorsal lateral geniculate nucleus
12	DLPAG	dorsolateral periaqueductal grey
13	DM	dorsomedial hypothalamic nucleus
14	DMPAG	dorsomedial periaqueductal grey
15	DMTg	dorsomedial tegmental area
16	DpG	deep grey layer of the superior colliculus
17	DpMe	deep mesencephalic nucleus
18	DpWh	deep white layer of the superior colliculus
19	DR	dorsal raphe nucleus
20	DTg	dorsal tegmental nucleus
21	f	fornix
22	fi	fimbria of the hippocampus
23	HDB	nucleus of the horizontal limb of the diagonal band
24	I	intercalated nuclei of the amygdala
25	ic	internal capsule
26	IF	interfascicular nucleus
27	InC	interstitial nucleus of Cajal
28	InG	intermediate grey layer of the superior colliculus
29	InWh	intermediate white layer of the superior colliculus

1	IPAC	interstitial nucleus of the posterior limb of the anterior commissure
2	La	lateral amygdaloid nucleus
3	LA	lateroanterior hypothalamic nucleus
4	LC	locus coeruleus
5	LDTg	laterodorsal tegmental nucleus
6	LGP	lateral globus pallidus
7	LH	lateral hypothalamic area
8	LHb	lateral habenular nucleus
9	lo	lateral olfactory tract
10	LOT	nucleus of the lateral olfactory tract
11	LPAG	lateral periaqueductal grey
12	LPBC	lateral parabrachial nucleus, central part
13	LPBD	lateral parabrachial nucleus, dorsal part
14	LPBE	lateral parabrachial nucleus, external part
15	LPBI	lateral parabrachial nucleus, internal part
16	LPBV	lateral parabrachial nucleus, ventral part
17	LSD	lateral septal nucleus, dorsal part
18	LSI	lateral septal nucleus, intermediate part
19	LSV	lateral septal nucleus, ventral part
20	LV	lateral ventricle
21	MCLH	magnocellular nucleus of the lateral hypothalamus
22	mcp	middle cerebellar peduncle
23	MCPO	magnocellular preoptic nucleus
24	ME	median eminence
25	MeA	medial amygdaloid nucleus, anterior part
26	MePD	medial amygdaloid nucleus, posterodorsal part
27	MePV	medial amygdaloid nucleus, posteroventral part
28	MGD	medial geniculate nucleus, dorsal part
29	MGM	medial geniculate nucleus, medial part

1	MGP	medial globus pallidus
2	MGV	medial geniculate nucleus, ventral part
3	MHb	medial habenular nucleus
4	ml	medial lemniscus
5	mlf	medial longitudinal fasciculus
6	MnPO	median preoptic nucleus
7	MnR	median raphe nucleus
8	MPA	medial preoptic area
9	MPB	medial parabrachial nucleus
10	MPBE	medial parabrachial nucleus, external part
11	MPO	medial preoptic nucleus
12	MS	medial septal nucleus
13	mt	mammilothalamic tract
14	MTu	medial tuberal nucleus
15	MZMGV	marginal zone of the medial geniculate
16	opt	optic tract
17	OT	oxytocin
18	ox	optic chiasm
19	Pa	paraventricular nucleus
20	PaLM	paraventricular nucleus, lateral magnocellular part
21	PaMM	paraventricular nucleus, medial magnocellular part
22	PaPO	paraventricular nucleus, posterior part
23	PaV	paraventricular nucleus, ventral part
24	PAG	periaqueductal greyPe periventricular hypothalamic nucleus
25	PeF	perifornical nucleus
26	PH	posterior hypothalamic area
27	PIL	posterior intralaminar thalamic nucleus
28	Pir	piriform cortex
29	PLCo	posterolateral cortical amygdaloid nucleus

1	PnC	pontine reticular nucleus, caudal part
2	PnO	pontine reticular nucleus, oral part
3	PnR	pontine raphe nucleus
4	PnV	pontine reticular nucleus, ventral part
5	PoT	posterior thalamic nuclear group, triangular part
6	PP	peripeduncular nucleus
7	PRL	prolactin
8	PSTh	parasubthalamic nucleus
9	PV	paraventricular thalamic nucleus
10	PVA	paraventricular thalamic nucleus, anterior part
11	RCh	retrochiasmatic area
12	RMg	raphe magnus nucleus
13	s5	sensory root of the trigeminal nerve
14	SCh	suprachiasmatic nucleus
15	SFi	septofimbrial nucleus
16	SFO	subfornical organ
17	SG	suprageniculate thalamic nucleus
18	SI	substantia innominata
19	SLEA	sublenticular extended amygdala
20	sm	stria medullaris of the thalamus
21	SNC	substantia nigra, pars compacta
22	SNR	substantia nigra, reticular part
23	SO	supraoptic nucleus
24	SPa	subparaventricular zone of the hypothalamus
25	st	stria terminalis
26	STh	subthalamic nucleus
27	SubCD	subcoeruleus nucleus, dorsal part
28	SubCV	subcoeruleus nucleus, ventral part
29	SuM	supramammillary nucleus

1	Te	terete hypothalamic nucleus
2	Tu	olfactory tubercle
3	VEn	ventral endopiriform nucleus
4	VLG	ventral lateral geniculate nucleus
5	VLGMC	ventral lateral geniculate nucleus, magnocellular part
6	VLGPC	ventral lateral geniculate nucleus, parvicellular part
7	VLPAG	ventrolateral periaqueductal grey
8	VLPO	ventrolateral preoptic nucleus
9	VMH	ventromedial hypothalamic nucleus
10	VMHc	ventromedial hypothalamic nucleus, central part
11	VMHdm	ventromedial hypothalamic nucleus, dorsomedial part
12	VMHvl	ventromedial hypothalamic nucleus, ventrolateral part
13	VMPO	ventromedial preoptic nucleus
14	VOLT	vascular organ of the lamina terminalis
15	VP	ventral pallidum
16	VRe	ventral reuniens thalamic nucleus
17	vsc	ventral spinocerebellar tract
18	VTA	ventral tegmental area
19	ZI	zona incerta
20		

INTRODUCTION

Prolactin (PRL) is a polypeptide hormone produced at the adenohypophysis, best-known for its role

in the development of the mammary gland and milk production [1]. To date, PRL has been attributed

more than 300 different biological actions [2] in reproduction, homeostasis, angiogenesis or immunity,

among others [3]. To fulfil this array of functions, PRL has different target tissues, including the

mammary gland, the uterus and other peripheral organs [3] and, importantly, the brain [4,5]. In the

brain, PRL is mainly involved in the regulation of a number of behaviours and physiological processes

related to female reproduction and lactation [6] . Indeed, PRL has been described as a pleiotropic

hormone responsible for adapting every important aspect of female physiology and behaviour to the

demands of motherhood [7].

The key role of prolactin in maternal physiology has traditionally biased neuroendocrinological

studies towards its functions in the brain of females, whereas the actions of this hormone in the

male brain have drawn less attention. For instance, the distribution of the PRL receptor (PRLR) has

been reported for the female [8,9] but not the male rodent brain. One of the main reasons for this bias

relates to the fact that male rodent models display low levels of circulating PRL under standard

conditions [10], reflecting a limited access and functionality of PRL in the brain. Still, several studies do

report acute rises in systemic PRL secretion in males associated to certain physiological and

pathological conditions, such as the stress response [11] or sexual behaviour [12].

Regarding the latter, PRL is known to be involved in the control of male copulatory behaviour in

rats[13], mice [14], other rodent models [15] and humans[16]. Thus, PRL is released acutely with ejaculation [12]

and it is proposed to intervene in the satiation following copulation[17] leading to the refractory

period. In fact, chronic hyperprolactinaemia suppresses copulatory behaviour in animal models[18]. In

men, hyperprolactinaemia associated to several clinical conditions [19] (including antipsychotic

treatments [20]), yields sexual dysfunction and other side-effects [21]. The inhibitory role of PRL on

sexual function is proposed to have an important central component [22,23], but the exact regions and

mechanisms through which PRL operates in the male brain are still poorly understood.

In this light, the main aim of this work is to provide a comprehensive description of the distribution

of cells sensitive to PRL in the brain of males. To do so, we have supplemented males with a

standard dose of exogenous PRL and revealed PRL-responsive cells in the brain by means of the

immunohistochemical detection of the phosphorylated form of signal transducer and activator of

transcription 5 (pSTAT5). Phosphorylation of STAT5 is a key event in the Jak/STAT signalling cascade

of the PRL receptor [3]. In contrast to other methods, e.g. in situ hybridization of PRL receptor (PRLR)

mRNA (or its immunodetection), pSTAT5 immunohistochemistry in PRL-supplemented animals

reveals those cells that are not just expressing PRL receptors or specific elements of their signalling

cascade, but those that are fully responsive to PRL.

The second goal of this work is to compare the pattern of central PRL-derived signalling between

male and female mice, in search for intersexual differences. For this purpose, we compare

qualitatively and quantitatively the patterns of our PRL-supplemented males with ovariectomised,

steroid-primed females treated with an equivalent dose of PRL. Given the key role of PRL in female

behaviour and physiology and the broad patterns of PRL-derived signalling in the female mouse

brain [4], we hypothesize that the central sensitivity to PRL will show a female-biased sexual

dimorphism. In addition, a direct comparison of the distribution of pSTAT5 in the brain of males and

females might also be useful for the functional contextualization of PRL input to the male brain.

In accordance with the involvement of PRL in reproductive physiology and behaviour, gonadal

steroids are important regulators of PRL function in the brain. In female rodents, estradiol and

progesterone have been shown to target most PRL-sensitive neurons in the brain [24] and to regulate

PRL signalling at different levels [25–27]. On the other hand, testosterone has been documented to

inhibit PRL receptor expression in the pituitary gland of male rats and pituitary PRL release in male

rats and mice [28–30]. Beyond these findings, to our knowledge there is currently no available evidence

supporting a regulatory role of testosterone in brain PRL signalling. Hence, in this work we also

explore the putative contribution of testosterone in shaping the patterns of PRL responsiveness in

the male brain. To this end, we analyse the levels of pSTAT5-ir in the main PRL-responsive regions of

the male mouse brain after permanent testosterone withdrawal through castration, as compared to

intact male mice, both supplemented with PRL.

By characterizing the patterns of PRL signalling in the male brain and clarifying their dependence on

testosterone, we intend to shed light on the roles of PRL in male behaviour and physiology, as well

as on the neural substrates for PRL-dependent, male-specific behaviours.

MATERIAL AND METHODS

Animals

For the present study, we used 15 male and 6 female mice of the CD1 strain (Charles River

Laboratories, France), aging between 8 and 24 weeks. These animals were housed in polypropylene

plastic cages under controlled temperature (24 ± 2 ºC) and lighting conditions (12h:12h; lights ON at

8 am), with *ad libitum* access to food and water. Intact males were housed individually (to avoid

inter-male aggression) whereas castrated males and females were group-housed (4 to 6 animals per

group, to avoid isolation-derived stress). Animals were treated throughout according to the

European Union Council Directive 2010/63/EU (6106/1/10 REV1) and procedures were approved by

the Committee of Ethics on Animal Experimentation of the University of Valencia, where the

experiments were performed.

Experimental Design

Male mice were randomly assigned to three experimental groups: (1) Male Control (n=3); (2)

Male+PRL (n=6) and (3) Castrated+PRL (n=6). Animals of Male Control and Male+PRL groups were

left gonadally intact, whereas males from Castrated+PRL group underwent orchidectomy. Following

the procedure by Brown and collaborators[5,31], Male+PRL and Castrated+PRL groups received an

exogenous PRL supplementation 45 minutes prior to perfusion.

In order to compare central PRL sensitivity between males and females, we also included in our

study a group of ovariectomised, steroid-primed and PRL-supplemented females, which belonged to

a sample of a previously published set of experiments [4]. The steroid-replacement schedule

conducted on these females intended to emulate the proestrus-estrus phase of the estrous cycle [32].

Given the considerable variability of pSTAT5-ir in the brain of freely-cycling females, likely due to

gonadal steroid influence on PRL signalling [4], this group provided a more stable comparison than

freely-cycling females and a more accurate comparison than estrous cycle screening.

Data from these four groups of animals (Male Control; Male+PRL; Castrated+PRL and Female+PRL)

allowed us to: a) characterize the pattern of PRL-derived signalling in the male brain by analysing the

distribution of pSTAT5-ir in group Male+PRL; b) analyse the effect of PRL supplementation on

pSTAT5-ir distribution by comparing the Male Control with Male+PRL groups; c) explore sexual

dimorphism in the central sensitivity for PRL by comparing Male+PRL with Female+PRL groups; and

finally, d) assess the modulatory role of testosterone on PRL signalling in the male brain through the

comparison of the Male+PRL and the Castrated+PRL groups.

Ovariectomy and Orchidectomy

Males of Castrated+PRL group were orchidectomised at approximately 12 weeks of age. Briefly,

animals were deeply anaesthesised with i.p. ketamine (Imalgene 500, Merial, Toulouse, France,

75mg/kg) and medetomidine (Domtor 1mg/ml, Esteve, Barcelona, Spain, 1 mg/kg) and surgery was

performed via a single midline incision on the scrotal sac. After surgery, i.p. atipamezol

hydrochloride (Antisedan, Pfizer, New York, USA, 1 mg/kg) was administered to reverse anaesthesia

and s.c. butorphanol tartrate 1% (Torbugesic, Pfizer, New York, USA, 20 μl) was regularly

administered through the postsurgical period for pain control. Castrated males had 2-3 weeks of

recovery under undisturbed housing conditions before the beginning of the experiments.

Experimental females underwent ovariectomy 9-10 weeks after birth under identical pharmacological conditions as explained for males [4]. Surgery was conducted through two incisions on both sides of the back of the animal. Ovariectomized females had at least seven days of recovery after surgery.

Hormone Treatments

Experimental females underwent a steroid replacement schedule [4] consisting of a sustained treatment with 17-β estradiol and an acute treatment with progesterone. In accordance with the experimental induction of estrus [32], estradiol was administered on a slow-release profile during 7 days, by means of the subcutaneous placement of silastic tubing implants (Dow Corning Corporation, Midland, MI, USA) filled with 20 µg/ml β-estradiol (Sigma, St Louis, MO, USA) diluted in sunflower oil. Silastic tubing had an inner diameter of 1.67 mm and an outer diameter of 2.41 mm, and implants were cut to a length of 20 mm. Implants were inserted subcutaneously on the lumbar region under isoflurane anaesthesia (Isoflo, Esteve Veterinaria, Barcelona, Spain). Animals were also administered a subcutaneous dose of 20 µl butorphanol tartrate 1% (Torbugesic, Pfizer, New York) during implant placement surgery, for pain control. By contrast, progesterone (Sigma, St Louis, MO, USA) was administered to females acutely in a 500 µg subcutaneous injection, diluted in sunflower oil, in the morning of the seventh day of estradiol treatment, specifically 3 hours and 45 minutes prior to perfusion. In order to make males and females directly comparable for the analysis of sexual dimorphism, males of the Male+PRL group received vehicle treatment parallel to that of females (implants with sunflower oil, plus s.c. oil injection 3h 45 minutes prior to perfusion).

Regarding PRL supplementation, except for the Male Control group (which received vehicle injection), all experimental groups were administered an acute 5 mg/kg i.p. dose of ovine PRL (Sigma, St Louis, MO, USA) 45 minutes before perfusion. The dose and timing of PRL administration rendered homogenous, supraphysiological circulating levels of PRL and ensured that the peak of STAT5 phosphorylation in response to a PRL challenge coincided with perfusion [5]. This is intended to

reveal the maximal potential response patterns among experimental animals and also to allow direct

comparison between experimental groups, regardless of sex (male vs female) or physiological

condition (intact vs castrated males).

Tissue Collection and Histological Processing

Animals received an overdose of sodium pentobarbital (Vetoquinol, Madrid, Spain) and were

transcardially perfused with 4% paraformaldehyde in 0.1M phosphate buffer (PB), pH 7.4. Brains

were carefully extracted and post-fixed overnight through immersion in the same fixative, then

cryoprotected by immersion in 30% sucrose in 0.01 M PB until they sank (2-3 days). Using a freezing

microtome (Microm HM-450, Walldorf, Germany) brains were sectioned in five parallel series of 40

μm thick coronal sections. Series were stored in PB-30% sucrose at -20ºC until their use.

Immunohistochemistry for pSTAT5

Immunohistochemistry was conducted in free-floating sections under light shaking at room

temperature (25ºC) unless otherwise stated. Immunohistochemistry protocol was adapted from

Brown et al.[5,31]. Tissue sections were thoroughly rinsed between stages for at least three 10-min

washes in TRIS-buffered saline, 0.05M, pH 7.6 (TBS). After thawing, sections underwent an initial

antigen retrieval step, consisting in two sequential 6-minute incubations in 0.01 M TRIS buffer (TB),

pH 10 at 85ºC, and brought quickly to room temperature in between. Tissue was then incubated in:

a) 1% hydrogen peroxide (H_2O_2) for 30 minutes, for endogenous peroxidase inhibition; b) 2% BSA,

2% goat serum and 0.3% Triton X-100 in TBS for 1h, in order to block unspecific labelling; c) rabbit

monoclonal anti-pSTAT5 primary antibody (Phospho-Stat5 Tyr694 (D47E7), Catalog number #4322,

Cell Signaling Technology, Danvers, MA) diluted 1:500 in TBS plus Triton X-100 0.1% for 72 h at 4ºC;

d) biotinylated goat anti-rabbit IgG (Vector Laboratories, Peterborough, UK; RRID AB_2313606)

1:200 in TBS for 90 minutes; and e) avidin-biotin-peroxidase complex (ABC Elite kit; Vector

Laboratories; RRID AB_2336819) in TBS for 90 minutes. Peroxidase label was developed using

0.005% 3-3'-diaminobenzidine (Sigma) and 0.01% H_2O_2 in TB pH 7.6 for about 15 minutes, obtaining

thereby a brown nuclear staining. Sections were rinsed in TB and mounted onto gelatinized slides,

dehydrated in graded ethanol, cleared with xylene and coverslipped with Entellan.

The specificity of the anti-pSTAT5 primary antibody was demonstrated by means of an antibody

competition assay (Supplementary Figure 1).

Analysis of pSTAT5 Immunoreactivity

The patterns of pSTAT5-ir distribution were mapped for the brain of intact males (Male+PRL) and

females (Female+PRL) and illustrated semiquantitatively in camera lucida drawings of selected brain

sections (Fig. 1). In addition, a quantitative assessment of the density of cells showing pSTAT5

immunoreactivity (pSTAT5-ir) in representative brain sites was performed for PRL-treated females

and males, both intact and castrated. To do so, we first selected frames of the chosen nuclei using

the stereotaxic atlas of Paxinos and Franklin (2004). Then, we obtained photomicrographs of these

frames in both hemispheres using a digital camera (Leica DFC495) attached to a microscope Leitz

DMRB (Leica AG, Germany). Thereafter, images were processed and analysed using ImageJ. Briefly,

we subtracted background light and converted the RGB colour image to greyscale by selecting the

green channel. Then, we binarised the greyscale image setting the 75% of the mode of the histogram

as a threshold, thus including every pixel below this threshold as positively labelled. Using the

ImageJ commands "fill holes", "open" (3 iterations) and "watershed", we filtered small particles

(noise or fragmented nuclei) and separated fused objects. The resulting particles were additionally

filtered by area (larger than 70 μm^2, corresponding to an approximate diameter of 9.4 μm) and

finally counted automatically. We calculated the mean (interhemispheric) density of pSTAT5-

imunoreactive cell nuclei for each specimen by dividing the counts for both hemispheres by the area

of both frames.

Statistical Analysis

We performed statistical analysis of the resulting data on the SPSS software package. After checking

for normality (Kolmogorov-Smirnov test with Lilliefors' correction) and homogeneity of variances

(Levene's test), we conducted three different statistical comparisons. First, we compared levels of

pSTAT5-ir density between intact males (Males + PRL) and ovariectomised, steroid-primed females

(Female+PRL), in search for intersexual differences. Then, we checked for the effect of orchidectomy

on male pSTAT5-ir levels by comparing intact (Male+PRL) and castrated (Castrated+PRL) males. For

both analyses, the samples fulfilling the criteria for a parametric analysis were subject to an

independent t-test, whereas the samples of the remaining nuclei were subject to a non-parametric

Mann-Whitney test. For each statistical test, we applied a significance level of 0.05.

RESULTS

The main focus of this section is to characterise the central patterns of pSTAT5-ir found in male mice

in the presence of high levels of circulating PRL (Male+PRL group), as an indicator of PRL-derived

signalling occurring in the male brain. In the same vein, we compare the distribution of pSTAT5-ir

between PRL-supplemented males (Male+PRL group) and non-supplemented males (Male Control

group). Then, we analyse intersexual differences in PRL-derived signalling in the mouse brain. For

this purpose, we compare the patterns of pSTAT5-ir in the male (Male+PRL group) and female

(Female+PRL group) brains and assess the density of pSTAT5-ir in a number of brain regions

displaying PRL-derived signalling in both males and females. Finally, we explore the effect of

testosterone withdrawal in the PRL-derived signalling of the male brain by comparing pSTAT5-ir

levels between intact and castrated male mice. For the following anatomical descriptions, we adhere

to the neuroanatomical terminology proposed by Paxinos and Franklin[33] (see also list of

abbreviations above).

A.   Patterns of pSTAT5 immunoreactivity in the male mouse brain: effect of PRL supplementation

Immunohistochemistry for pSTAT5 produced a defined staining in the examined brain tissue, which

was restricted to the cell nucleus (see Fig. 2), as shown in our previous work [4]. Labelling was

suppressed by pre-incubation of the antibody with the phosphorylated peptide used for

immunization, but not when the antibody was incubated with the equivalent, non-phosphorylated

peptide (see Suppl. Fig. 1). Therefore, we safely assume that, in our immunohistochemical

procedure, the antibody is detecting specifically pSTAT5.

Figure 1 maps, on the left side of the brain, the distribution of pSTAT5-ir observed in PRL-

supplemented males (blue dots; for the significance of red dots see below). On the right side of the

brain, black dots show the distribution of cells expressing PRL receptor (PRLR) in the male mouse

brain, as reported by the Allen Brain Institute (see Suppl. Fig. 2). Figure 2 shows photomicrographs

illustrating pSTAT5-ir in selected brain nuclei for all of our experimental groups.

Males supplemented with PRL (Male+PRL) showed moderate-to-dense pSTAT5-ir in several nuclei of

the hypothalamus and cerebral hemispheres (Fig. 2, left column), as well as in the choroid plexus

(not shown). In the telencephalon, pSTAT5-ir was mainly observed in the septum, specifically in the

area comprising the septofimbrial nucleus (SFi) and the triangular septal nucleus (TS) (Figs. 1c and

1b). In this region, the subfornical organ (SFO) also showed some immunoreactive cells. In addition,

the posterodorsal nucleus of the medial amygdala (MePD) displayed scarce pSTAT5-ir, with few

immunolabelled cells restricted to the upper corner of the nucleus, in close contact to the

intramygdaloid BST (BSTia, see Figs. 1f and 2f). Importantly, all these brain nuclei display PRLR mRNA

expression in the male mouse (Fig. 2 right side).

In the hypothalamus, pSTAT5-ir was present in the preoptic, anterior and tuberal regions. In the

preoptic hypothalamus (Figs. 1a-d and 2j), pSTAT5-ir cells were concentrated in the juxtaventricular

nuclei, namely the vascular organ of the *lamina terminalis* (VOLT) and the anteroventral

periventricular (AVPe), the ventromedial preoptic (VMPO) and the Pe nuclei. In addition, the

adjoining MPA displayed a few immunolabelled cell nuclei.

a- Br +0.50 mm

b- Br +0.20 mm

c- Br +0 mm

Figure 1. Mapping of pSTAT5 immunoreactivity and prolactin receptor expression in the brains of

male and female mice

Figure 2. Representative examples of pSTAT5 immunoreactivity in the brain of female, intact male and castrated male mice

In the anterior hypothalamus, pSTAT5-ir was once again restricted to juxtaventricular structures, the paraventricular nucleus (Pa) and the Pe (Figs. 1e and 2n), as well as the retrochiasmatic area (RCh, Fig. 1e). In the Pa, labelling was restricted to the medial aspect of the nucleus, mainly to the ventral

(PaV), medial magnocellular (PaMM) and medial parvocellular (PaMP) subdivisions, with very few

labelling observed in the lateral aspect of the nucleus (PaLM) or the subparaventricular nucleus

(SPa). In the tuberal hypothalamus (Figs. 1f and 2r), the arcuate nucleus (Arc) displayed abundant

pSTAT5-ir, with some immunolabelled cells displaced into the median eminence (ME). In some

males, the dorsomedial nucleus (DM) displayed few scattered immunolabelled cells. Finally, pSTAT5-

ir appeared in the ventromedial hypothalamic nucleus (VMH), with higher levels in the ventrolateral

subdivision (VMHvl) and sparse labelling in the rest (VMHc and VMHdm).

In contrast to the Male+PRL group, non-supplemented males (Male Control, Fig. 2, rightmost

column), were virtually devoid of pSTAT5-ir, except for some labelled cells in the ventral aspect of

the arcuate nucleus (Arc, Fig. 2t) and the anterior aspect of the periventricular nucleus (Pe, not

shown).

B. Comparative analysis of pSTAT5-ir the male and female mouse brain

In comparison to males, ovariectomised, steroid-primed females supplemented with PRL

(Female+PRL) showed a more extensive distribution of pSTAT5-ir (Fig. 1, Fig. 2 middle-right column),

which matches previous descriptions conducted on freely-cycling female mice[4]. To summarize,

labelling in our experimental females was widespread in the basal telencephalon and hypothalamus,

but was also found within thalamic, midbrain and brainstem structures. To allow a direct comparison

between sexes, Figure 2 shows the distribution of pSTAT5-ir in equivalent brain sections of male and

female mice. In addition, in the semischematic drawings of brain sections depicted in Fig. 1, pSTAT5-

ir labelling that was common to males and females is shown as blue spots, whereas labelling

exclusively present in females is illustrated as red spots. Importantly, there was not a single brain

site where labelling was exclusively found in males. In other words, after administration of

equivalent doses of exogenous PRL, males show pSTAT5-ir in a reduced number of brain centres,

whereas females show immunoreactive cells in the same nuclei, plus a set of additional brain

centres. This represents a clear case of sexual dimorphism in favour of females.

We also checked for quantitative sexual dimorphism in PRL-derived signalling, by analysing the

density of pSTAT5-ir in the main nuclei showing pSTAT5-ir in PRL-supplemented intact males and

steroid-primed ovariectomised females. We designed counting frames for the following brain nuclei:

the AVPe/VMPO region, the Pa, the Arc, the DM, the VMHvl, the area comprising the central and

dorsomedial VMH (VMHc and VMHdm, respectively), the MePD and the SFi (Fig. 1). Using ImageJ we

assessed the density of pSTAT5-ir in each nucleus (mean hemispheric cell density) and performed

separate statistical tests on each of the analysed nuclei to compare Male+PRL and Female+PRL

groups. When data fulfilled the criteria for a parametric statistical analysis (the AVPe, the Arc, the Pa

and the VMHvl), we performed t-tests for independent samples. For the remaining nuclei (the DM,

the VMHc, the MePD and the SFi), we performed a non-parametric Mann-Whitney test. Figure 3

shows the results of these analyses. Females showed a general trend towards higher levels of

pSTAT5-ir than males. This effect reached statistical significance in the Arc ($t(10)=3.043$, $p=0.012$),

the DM ($U=32.0$; $p=0.026$) and the MePD ($U=30.0$; $p=0.004$), although the AVPe/VMPO also showed

an almost significant trend towards higher pSTAT5-ir density in females ($t(9)=2.183$; $p=0.057$).

Figure 3. Quantitative analysis of pSTAT5 immunoreactivity in selected brain regions of female and

male mice

C.   Effect of testosterone withdrawal in PRL-derived signalling in the male mouse brain

Finally, we explored the putative role of testosterone in the regulation of PRL-derived signalling in

the male brain. To do so, we compared pSTAT5-ir patterns between intact, PRL-supplemented males

(Male+PRL; Fig. 2, leftmost column) and a sample of castrated, PRL-supplemented males

(Castrated+PRL; Fig. 2, rightmost column). In qualitative terms, castrated and intact male mice

displayed similar patterns of pSTAT5-ir, but castrated males showed apparently lower levels of

pSTAT5-ir. Quantitative analysis of the labelling in both groups (using the same methodology and

exploring the same subset of nuclei analysed in the previous section) allowed comparing pSTAT5-ir

between gonadally intact and castrated males by means of independent t-tests for the Arc and AVPe

and non-parametric Mann-Whitney tests for the rest of the analysed regions. The results of this

statistical comparison reveal a significant effect of testosterone withdrawal in diminishing the levels

of pSTAT5-ir in the Pa (U= 33.00; p=0.004), in both the ventrolateral and central/dorsomedial VMH

(U= 30.00, p=0.015 for both frames) and in the SFi (U= 26.00, p=0. 05). In the remaining analysed

nuclei, the density of pSTAT5-ir cells was similar in Male+PRL and Castrated+PRL animals, as no

statistical differences were found (p>0.1) (Fig. 4).

Figure 4. Effect of testosterone withdrawal in pSTAT5 immunoreactivity of the male mouse brain

DISCUSSION

The present study examined, for the first time, the responsiveness of the brain of males to PRL.

Specifically, we analysed the distribution and steroid regulation of PRL-derived signalling in the male

mouse brain, by means of the immunohistochemical detection of pSTAT5. We also compared the

pattern of pSTAT5 immunoreactivity in the brain of males and females, aiming at assessing its

possible sexual dimorphism. In this section, we will first analyse the distribution of pSTAT5 in the

brain of male mice, with or without exogenous PRL administration, and compare it to the pattern of

expression of the PRLR, as shown in in situ hybridisation material provided by the Allen Brain

Institute (see Figure 1 and supplementary Figure 2). Then, we will comment on the observed sexually

dimorphic differences in central sensitivity to PRL. Next, we will discuss the role of testosterone as a

putative regulator of central PRL signalling and on the mutual regulatory relationship of PRL and

testosterone. We will conclude by reviewing how our findings contribute to elucidate the

neuroanatomical substrate of some putative actions of PRL in the brain of males.

Prolactin signalling in the male mouse brain, PRL supplementation and PRL receptor expression

Although the actions of PRL in the female mouse brain have received considerable attention due to

the role of PRL in female reproduction and motherhood[34], this work is, to our knowledge, the first

descriptive analysis of the distribution of PRL-derived signalling in the male mouse brain. In PRL-

supplemented male mice (Male+PRL group), brain regions positively labelled for pSTAT5-ir were

located mainly in the hypothalamus (the AVPe and VMPO, MPA, Pe, Pa, Arc, DM and VMH) and, to a

lesser extent, in the telencephalon (SFi in the septum and MePD in the amygdala), with no additional

pSTAT5-ir detected in any of the remaining brain divisions (Fig. 1).

Importantly, this pattern of pSTAT5-ir was highly dependent on systemic supplementation with high

doses of exogenous PRL, whereas non-supplemented controls (Male Control group) were virtually

devoid of pSTAT5-ir, except for some faintly labelled cells in the ventral Arc (Fig. 2t) and in the

anterior periventricular hypothalamus (not shown). These results suggest, in the first place, that the

patterns of pSTAT5-ir we report in this work are specifically produced by PRL, whereas other endocrine agents signalling through the Jak/STAT pathway, such as growth hormone[35], do not contribute substantially to STAT5 phosphorylation in the studied cases. This is further supported by the fact that pSTAT5 appears only in centres that show expression of PRLR in the brain of male mice (Figure 1; Suppl. Fig. 2), whereas the distribution of growth hormone receptor does not fit the pattern of pSTAT5-ir[35,36].

In addition, the reduced pSTAT5-ir found in males not supplemented with exogenous PRL indicates that, in basal conditions, PRL signalling in the male brain is very low, what fits the circulating levels reported for the hormone for males[10,37]. In fact, the presumed location of the TIDA neurons (as revealed by tyrosine hydroxylase immunohistochemistry), the dorsal aspect of the Arc[38], displays scarce pSTAT5-ir in Male Control mice (Fig. 2t). This supports the view of very low circulating PRL levels in male mice, at least in the light period of the day [37]. In line with these observations, PRL would influence the brain of males especially under some physiological conditions in which there is substantially increased hypophyseal or extra-hypophyseal PRL release, for instance after mating[39] or as part of the stress response [40].

Sexually dimorphic PRL-derived signalling in the mouse brain

In comparison to males, female mice displayed a more extensive distribution of pSTAT5-ir, mainly in the hypothalamus and basal telencephalon, but also present in nuclei of the thalamus, midbrain and brainstem. Patterns of pSTAT5-ir in males were always a completely overlapping fraction of those of female mice. In other words, we found no single brain structure labelled in males but not in females. Furthermore, a number of nuclei with pSTAT5 expression in both sexes displayed significantly higher pSTAT5-ir density in females than in males (the AVPe/VMPO, Arc, DM and MePD), but none of the analysed nuclei showed higher levels of pSTAT5-ir in males (Fig. 3). Altogether, our study reveals a clear, female-biased sexual dimorphism in PRL-derived signalling in the mouse brain upon exogenous PRL administration. In fact, this sexually dimorphic responsiveness to PRL would be extensive to

specific functional systems of the brain, for instance to the Sociosexual Behavioural Network (SBN),

the core system for the integration of social and reproductive behaviour[45]. In contrast to females,

several nodes of the SBN (the lateral septum, BST, medial amygdala or PAG, see Figs. 1 and 2) display

no pSTAT5-ir or marginal levesl (MePD) in males, even after exogenous PRL administration, although

they do express the PRL receptor (Fig. 1 and Supplementary Fig. 2). This suggests that, in these

areas, either PRL responsiveness is regulated through different receptor activation pathways in

males and females, or PRL signalling is downregulated in males, leading to a lower implication of PRL

in the regulation of sociosexual behaviour in males. However, this does not exclude a role in some

male-specific behaviours, as we will further discuss below.

Prolactin and Testosterone: Mutual Neuroendocrine Regulation

One of the main outcomes of the present work is the identification of a putative regulatory role of

testosterone in PRL signalling in the male mouse brain. Orchidectomy and consequent testosterone

withdrawal led to a significant decrease in pSTAT5-ir density in the SFi, the Pa and both the

central/dorsomedial and lateral VMH (Fig. 4). This indicates a positive regulatory role of testosterone

on PRL signalling in those brain regions. The actual mechanism of this process is not clear. Even

though testosterone has documented inhibitory effects on hypophyseal PRL release in male rats[29]

and mice[30], the effect of castration on pSTAT5-ir cannot be attributed to differences in circulating

PRL, since intact and castrated males were supplemented with equivalent levels of exogenous PRL.

In addition, this effect is not likely explained by a facilitated access of systemic PRL to the brain, as

our results reveal that the positive effect of castration on pSTAT5-ir is not generalised but rather

specific and restricted to discrete brain sites. Therefore, our results suggest that testosterone might

upregulate PRLR expression or perhaps signalling downstream the PRLR in specific neural

populations. Further research is required to elucidate the specific action of testosterone on PRL

signalling in male mice.

An additional question of interest is the identity of the actual biological modulator of PRL signalling we report here. Testosterone can be locally metabolized into two different neuroactive steroids: it can be reduced to dihydrotestosterone[42], which binds to androgen receptors, or aromatised to estradiol[43] that then binds to oestrogen receptors. Our results make it very unlikely that this regulation occurs through the aromatisation of testosterone to estradiol. Although all the nuclei where PRL signalling is affected by castration express at least one form of oestrogen receptor[44], none of them displays aromatase activity in the male mouse brain[45]. Conversely, all of the nuclei where PRL-derived signalling decreased with castration express androgen receptors, with the Pa and VMH showing very high levels[45]. Hence, evidence suggests that the effect of testosterone on PRL signalling that we report here is likely mediated by androgen rather than oestrogen receptors.

In addition to the reported regulation of testosterone on central PRL signalling, our data, together with published evidence, suggest a reciprocal regulatory action of PRL on the hypothalamus-pituitary-gonadal (HPG) axis and ultimately on testosterone release. According to the literature, PRL appears to upregulate the HPG axis and testosterone secretion in males. For instance, Bartke and collaborators[14] reported increases in FSH and LH in hyperprolactinaemic male mice, and Chandrashekar and Bartke[46] observed that exogenous PRL infusions stimulate the increase in serum testosterone in male rats, whereas immunological disruption of endogenous prolactin eliminated this effect. This regulatory action of PRL on the HPG axis likely occurs through the kisspeptin system. Kisspeptin plays an essential role in reproduction, as it regulates gonadotropin-releasing hormone (GnRH) expressing neurons in the brain[47], and thereby activate the HPG axis. Therefore, the expression of PRL receptors by kisspeptin neurons[48] allows for a direct role of PRL on HPG axis control and testosterone secretion. In this context, our results provide support for the hypothesis that the brain kisspeptin system is subject to PRL regulation in male mice, too, since our sample of male mice displayed substantial pSTAT5-ir in the hypothalamic nuclei containing kisspeptin-positive neurons[49], the AVPe and the Arc (Figs. 2j and 2r). Nevertheless, this possibility requires experimental

proof by assessing the coexpression of kisspeptin and pSTAT5 immunoreactivity in the

aforementioned nuclei.

Altogether, our results and previously published evidence in mice and rats suggest that PRL and

testosterone have complex and reciprocal regulatory actions. On the one hand, testosterone inhibits

hypophyseal PRL release in male mice[30], but it also appears to enhance PRL signalling in some brain

nuclei (Fig. 4). On the other hand, PRL might act onto hypothalamic kisspeptin neurons to stimulate

the release of testosterone by the testes.

Prolactin signalling in the male brain: functional implications

Despite the reported female bias in the central responsiveness to PRL, our findings confirm that the

male mouse brain is indeed responsive to PRL, implying that PRL likely exerts certain actions in the

male mouse brain, too. The nature of these actions is mostly unknown, but PRL signalling is virtually

absent in the nodes of the socio-sexual brain network (SBN; with the exception of the VMH), thus

suggesting that PRL has a minor direct role in regulation of social behaviours (sexual, agonistic,

affiliative and/or parental), although a putative indirect role cannot be excluded either (for instance

through the regulation of central oxytocinergic pathways, see below). When comparing males with

females, PRL action in the SBN shows a clear female-biased dimorphism (see above) where it

certainly regulates sociosexual (including maternal) behaviours. Indeed, the differential action of

hormones in males and females is a defining feature of the SBN [41] that explains the gender-specific

endocrine regulation of most social behaviours[4,24].

In this respect, the most prominent (but poorly understood) function of PRL in males is the

regulation of sexual behaviour [3,16]. It has been shown that central action of PRL after mating

suppresses sexual behaviour as part of the satiatory mechanisms operating after ejaculation [17,22].

Similar inhibitory action of PRL onto sexual behaviour occurs under chronic hyperprolactinaemia[50–52].

Prolactin is proposed to do so by modulating the function of some of the major dopaminergic

circuits in the brain, closely involved in male sexual function[53]. This hypothesis is supported by

experimental evidence that acute elevations of circulating PRL after copulation lead to significant

decreases in dopaminergic activity in the target sites of dopaminergic pathways, including the

striatum [23,54] and the MPA region [55]. It is suggested that this decreased dopaminergic activity would

occur by PRL action onto cells giving rise to ascending dopaminergic pathways, e.g. the incerto-

hypothalamic, mesocorticolimbic and nigrostriatal projections[17]. However, according to our findings,

virtually none of the nuclei originating these dopaminergic pathways displays PRL-derived signalling

in male mice, even after injection of exogenous PRL (Fig. 2), neither PRLR expression (Fig. 1 and

supplementary Fig.2). In sum, our results suggest that the proposed action of PRL over these

dopaminergic systems should rather be indirect, through nuclei afferent to the dopaminergic cells.

In this context, the VMH is closely related to sexual behaviour, it is sexually dimorphic[56], known to

promote the expression of copulatory behaviour in females[57] and to inhibit mounting in male rats [58].

According to our results, the VMH is substantially responsive to PRL in males, showing in fact similar

levels of pSTAT5-ir as in females (Figs. 2q, 2r and 3). In addition, our data on castrated male mice

suggest that this responsiveness is also dependent on testosterone (Fig. 4). Hence, PRL could be able

to regulate male sexual behaviour also through the VMH in an androgen-dependent fashion.

Related with this, research conducted fundamentally in male rats has established that the magno-

and parvocellular oxytocin (OXT) cells in the Pa are involved both in the peripheral and central

regulation of male sexual behaviour. On the one hand, peripheral OXT released from Pa cells [59,60]

facilitates genital reflexes, erection and copulatory behaviour in response to sexual cues and sexual

interaction [61]. On the other hand, OXT also regulates sexual behaviour through its central action,

affecting key neural centres for sexual behaviour, such as the MPO [62–64] and VMH (see above). As

described before, our findings reveal substantial levels of pSTAT5-ir in the Pa of male mice,

comprising the magno- and especially the parvocellular division of the nucleus. This responsiveness

to PRL is also dependent on testosterone, since castrated males display significantly lower levels of

pSTAT5-ir in this nucleus (Fig. 3). Summarizing, PRL could regulate copulatory behaviour expression

through its testosterone-dependent action over two specific hypothalamic nuclei, the Pa (specifically

through the OXTergic populations of this nucleus) and the VMH.

The interaction between OXT and PRL at the level of the Pa can also have a role in the regulation of

parental behaviours. There is solid evidence indicating that PRL [65] and OXT are fundamental in the

control of maternal behaviour [66,67] and it is likely that they have an equivalent role in males.

Tachikawa and collaborators[68] showed that paternal behaviour can be induced in male mice (which

are usually infanticidal) by postcopulatory cohabitation with a maternal female. This behavioural

induction of paternal behaviour is dependent on OXT release in the nucleus accumbens (Acb), as it is

disrupted in knock out male mice for CD38 (a second messenger involved in the release of OXT)[69]

and rescued subsequently after the infusion of OXT in the Acb [70]. Dolen and collaborators [71]

demonstrated that the OXT innervation of the Acb arises in the Pa (see also[72]). Hence, the input to

the Acb from OXTergic Pa cells appears to be instrumental in the induction of paternal care in male

mice. Our results, in turn, suggest that OXT cells in the Pa are probably responsive to PRL in male as

well as female mice (Figs. 2m and 2n). Therefore, PRL might be involved in the induction of parental

behaviour in male mice, too. In support of this hypothesis, PRL facilitates the pup-sensitisation

process in male rats [73] in a similar fashion as in females [65] and has further roles in other experimental

models for paternal behaviour [74].

Exogenous PRL in male rats has also been recently shown to have pro-social effects, in particular in

social recognition and social learning paradigms in which OXT has also been involved [75]. The OXT

neurons in the Pa, probably sensitive to PRL according to our results, may mediate the PRL effects on

OXT-dependent social behaviours[71].

Our results indicate a modulatory role of PRL onto cells of the Arc and Pe nuclei of males, which

show abundant pSTAT5-ir (Fig. 2r-t). These nuclei contain the dopaminergic neuron populations

responsible for the inhibitory feedback control of hypophyseal PRL release[76]: the tuberoinfundibular

(TIDA) and tuberohypophyseal (THDA) neurons in the dorsomedial and rostral Arc[77], and the

periventricular hypophyseal neurons (PHDA) in the Pe[78]. In fact, it has been shown in female rats

that PRL feedback signal in these dopaminergic populations is conveyed through the Jak/STAT

pathway, specifically through STAT5b[79]. Therefore, our results in the Arc and Pe likely reflect the

(expected) occurrence of the same feedback mechanism in male mice.

Studies in lactating rodents ascribe an additional role of PRL in the regulation of energy balance and

food intake[6]. This physiological process is controlled by a complex neural network involving several

hypothalamic nuclei and different neurochemical signals[80]. According to our findings, some of the

hypothalamic centres belonging to this network display responsiveness to PRL in the male mouse

brain, namely the Arc, the VMH, the DM and the Pa. Taken together, our findings indicate that PRL

might be involved in the control of food intake not only during motherhood, but also in non-lactating

females and males, under certain physiological circumstances concurrent with high levels of

circulating PRL (e.g. acute stress[81,82]).

Prolactin also plays an important role in the modulation of anxiety and stress. Concerning anxiety,

studies in female rats indicate that PRL has anxiolytic properties when administered i.c.v. or i.v.,

whereas chronic i.c.v. antisense blocking of the PRLR expression results in increased anxiety[83]. In

contrast to females, PRL has a weak anxiolytic effect in males[83]. This intersexual difference might be

related to our findings of a sexually dimorphic responsiveness to PRL in the central structures

directly involved in the regulation of anxiety and emotionality, the Ce and the anterodorsal BST [84]. In

these nuclei, males are devoid of pSTAT5-ir (Fig. 1) but show PRLR mRNA transcript (Fig. 1 and

Supplementary Fig. 2d-e) whereas females show quite abundant immunostaining, especially during

pregnancy and lactation [4].

Concerning stress modulation, central inhibition of the expression of PRLR by i.c.v. antisense

blocking results in an enhanced activation of HPA axis[83] in male and female rats, thus demonstrating

a downregulation of the stress response by PRL. The presence of pSTAT5-ir in the Pa of males (in

equivalent levels to females, see Figs. 2m, 2n and 3) suggests that this modulatory effect of PRL on

the reactivity of the HPA axis, would probably take place through PRL action on corticotropin-releasing factor (CRF) or vasopressin (AVP) cells [85]. In fact, PRL is a well-documented regulator of the function of magno- and parvocellular vasopressinergic neurosecretory cells at the level of the hypothalamic Pa [86–88]. Although effects of PRL on these parvocellular AVP cells has been proven to be mediated by ERK/MAPK pathway [88], our data in males and females suggest that the Jak/STAT5 signalling cascade can also be involved. This would imply a non-dimorphic regulatory role of PRL on the stress response (in addition to its role on anxiety).

Conclusions

This work characterizes the distribution of PRL responsiveness in the male mouse brain and its dependence on circulating PRL levels, sexual dimorphism and gonadal steroid regulation. Male mice display specific patterns of PRL-derived signalling only in the presence of high levels of circulating PRL. These patterns comprise mainly hypothalamic nuclei and some telencephalic sites, and are also more limited in extension and density than the equivalent patterns in female mice, evidencing a clear sexual dimorphism in favour of females. Furthermore, PRL-derived signalling in the male brain is regionally dependent on testosterone input, indicating a reciprocal regulation between both hormones. This work confirms that PRL exerts central actions in males, suggesting a role of the "maternal hormone" in the regulation of male sexual behaviour, PRL release feedback control, as well as in regulation of stress response.

ACKNOWLEDGEMENTS

The authors are indebted to The Allen Brain Institute for providing open access to their invaluable material and permission to use and publish its data.

REFERENCES

1.   Riddle O, Bates R, Dykshorn S. The preparation, identification and assay of prolactin—a

hormone of anterior pituitary. *Am J Physiol* 1933;(105):191–216.

2.   Bole-feysot C, Goffin V, Edery M, Binart N, Kelly P a. Prolactin (PRL) and Its Receptor :

Actions , Signal PRL Receptor Knockout Mice. *Endocr. Rev.* 1998;19(February):225-268.

doi:10.1210/er.19.3.225.

3.   Freeman ME, Kanyicska B, Lerant A, Nagy G. Prolactin: structure, function, and regulation of

secretion. *Physiol. Rev.* 2000;80(4):1523-631.

4.   Salais-López H, Lanuza E, Agustín-Pavón C, Martínez-García F. Tuning the brain for

motherhood: prolactin-like central signalling in virgin, pregnant, and lactating female mice.

Brain Struct. Funct. 2017;222(2):895-921. doi:10.1007/s00429-016-1254-5.

5.   Brown RSE, Kokay IC, Herbison AE, Grattan DR. Distribution of prolactin-responsive neurons

in the mouse forebrain. *J. Comp. Neurol.* 2010;518:92-102. doi:10.1002/cne.22208.

6.   Grattan DR, Pi XJ, Andrews ZB, et al. Prolactin receptors in the brain during pregnancy and

lactation: implications for behavior. *Horm. Behav.* 2001;40:115-124.

doi:10.1006/hbeh.2001.1698.

7.   Grattan DR, Kokay IC. Prolactin: A pleiotropic neuroendocrine hormone. *J. Neuroendocrinol.*

2008;20:752-763. doi:10.1111/j.1365-2826.2008.01736.x.

8.   Bakowska JC, Morrell JI. Atlas of the neurons that express mRNA for the long form of the

prolactin receptor in the forebrain of the female rat. *J. Comp. Neurol.* 1997;386(April):161-

177. doi:10.1002/(SICI)1096-9861(19970922)386:2<161::AID-CNE1>3.0.CO;2-#.

9.   Bakowska JC, Morrell JI. The distribution of mRNA for the short form of the prolactin receptor

in the forebrain of the female rat. *Mol. Brain Res.* 2003;116:50-58. doi:10.1016/S0169-

328X(03)00213-4.

10. Guillou A, Romanò N, Steyn F, et al. Assessment of lactotroph axis functionality in mice:

longitudinal monitoring of PRL secretion by ultrasensitive-ELISA. *Endocrinology* 2015;(February):en.2014-1571. doi:10.1210/en.2014-1571.

11. Kirk SE, Xie TY, Steyn FJ, Grattan DR, Bunn SJ. Restraint stress increases prolactin-mediated phosphorylation of signal transducer and activator of transcription 5 in the hypothalamus and adrenal cortex in the male mouse. *J. Neuroendocrinol.* 2017;29(6). doi:10.1111/jne.12477.

12. Bronson FH, Desjardins C. Endocrine Responses to Sexual Arousal in Male Mice. *Endocrinology* 1982;111(4):1286-1291. doi:10.1210/endo-111-4-1286.

13. Drago F, Pellegrini-Quarantotti B, Scapagnini U, Gessa GL. Short-term endogenous hyperprolactinaemia and sexual behavior of male rats. *Physiol. Behav.* 1981;26(2):277-9.

14. Bartke A, Morgan WW, Clayton RN, et al. Neuroendocrine studies in hyperprolactinaemic male mice. *J. Endocrinol.* 1987;112(2):215-20.

15. Shrenker P, Bartke A. Effects of hyperprolactinaemia on male sexual behaviour in the golden hamster and mouse. *J. Endocrinol.* 1987;112(2):221-8.

16. Krüger THC, Hartmann U, Schedlowski M. Prolactinergic and dopaminergic mechanisms underlying sexual arousal and orgasm in humans. *World J. Urol.* 2005;23(2):130-138. doi:10.1007/s00345-004-0496-7.

17. Krüger THC, Haake P, Hartmann U, Schedlowski M, Exton MS. Orgasm-induced prolactin secretion: feedback control of sexual drive? *Neurosci. Biobehav. Rev.* 2002;26(1):31-44.

18. Cruz-Casallas PE, Nasello AG, Hucke EE, Felicio LF. Dual modulation of male sexual behavior in rats by central prolactin: relationship with in vivo striatal dopaminergic activity. *Psychoneuroendocrinology* 1999;24(7):681-93.

19. Rastrelli G, Corona G, Maggi M. The role of prolactin in andrology: what is new? *Rev. Endocr. Metab. Disord.* 2015;16(3):233-48. doi:10.1007/s11154-015-9322-3.

20. Riecher-Rössler A, Schmid C, Bleuer S, Birkhäuser M. Antipsychotics and hyperpolactinaemia:

pathophysiology, clinical relevance, diagnosis and therapy. *Neuropsychiatr. Klin. Diagnostik,*

Ther. und Rehabil. Organ der Gesellschaft Österreichischer Nervenärzte und Psychiater

2009;23(2):71-83.

21. Capozzi A, Scambia G, Pontecorvi A, Lello S. Hyperprolactinemia: pathophysiology and

therapeutic approach. *Gynecol. Endocrinol.* 2015;31(7):506-10.

doi:10.3109/09513590.2015.1017810.

22. Rehman J, Christ G, Alyskewycz M, Kerr E, Melman A. Experimental hyperprolactinemia in a

rat model: alteration in centrally mediated neuroerectile mechanisms. *Int. J. Impot. Res.*

2000;12(1):23-32.

23. Kalra PS, Simpkins JW, Luttge WG, Kalra SP. Effects on male sex behavior and preoptic

dopamine neurons of hyperprolactinemia induced by MtTW15 pituitary tumors.

Endocrinology 1983;113(6):2065-71. doi:10.1210/endo-113-6-2065.

24. Furigo IC, Kim KW, Nagaishi VS, et al. Prolactin-sensitive neurons express estrogen receptor-α

and depend on sex hormones for normal responsiveness to prolactin. *Brain Res.*

2014;1566:47-59. doi:10.1016/j.brainres.2014.04.018.

25. Neill JD, Freeman ME, Tillson SA. Control of the proestrus surge of prolactin and luteinizing

hormone secretion by estrogens in the rat. *Endocrinology* 1971;89(6):1448-53.

doi:10.1210/endo-89-6-1448.

26. Scully KM, Gleiberman AS, Lindzey J, Lubahn DB, Korach KS, Rosenfeld MG. Role of estrogen

receptor-alpha in the anterior pituitary gland. *Mol. Endocrinol.* 1997;11(6):674-81.

doi:10.1210/mend.11.6.0019.

27. Anderson GM, Kieser DC, Steyn FJ, Grattan DR. Hypothalamic prolactin receptor messenger

ribonucleic acid levels, prolactin signaling, and hyperprolactinemic inhibition of pulsatile

luteinizing hormone secretion are dependent on estradiol. *Endocrinology* 2008;149(4):1562-70. doi:10.1210/en.2007-0867.

28. Tong Y, Simard J, Labrie C, Zhao HF, Labrie F, Pelletier G. Inhibitory effect of androgen on estrogen-induced prolactin messenger ribonucleic acid accumulation in the male rat anterior pituitary gland. *Endocrinology* 1989;125(4):1821-8. doi:10.1210/endo-125-4-1821.

29. Gill-Sharma MK. Prolactin and male fertility: the long and short feedback regulation. *Int. J. Endocrinol.* 2009;2009:687259. doi:10.1155/2009/687259.

30. O'Hara L, Curley M, Tedim Ferreira M, Cruickshanks L, Milne L, Smith LB. Pituitary androgen receptor signalling regulates prolactin but not gonadotrophins in the male mouse. *PLoS One* 2015;10(3):e0121657. doi:10.1371/journal.pone.0121657.

31. Brown RSE, Herbison a E, Grattan DR. Differential changes in responses of hypothalamic and brainstem neuronal populations to prolactin during lactation in the mouse. *Biol. Reprod.* 2011;84(December 2010):826-836. doi:10.1095/biolreprod.110.089185.

32. Rissman EF, Early AH, Taylor JA, Korach KS, Lubahn DB. Estrogen receptors are essential for female sexual receptivity. *Endocrinology* 1997;138(1):507-10. doi:10.1210/endo.138.1.4985.

33. Franklin KBJ, Paxinos G. *The Mouse Brain in Stereotaxic Coordinates*. Compact 2. Asterdam etc.: Academic Press; 2004.

34. Grattan DR. Behavioural significance of prolactin signalling in the central nervous system during pregnancy and lactation. *Reproduction* 2002;123:497-506. doi:10.1530/reprod/123.4.497.

35. Furigo IC, Metzger M, Teixeira PDS, Soares CRJ, Donato J. Distribution of growth hormone-responsive cells in the mouse brain. *Brain Struct. Funct.* 2017. doi:10.1007/s00429-016-1221-1.

36. Burton KA, Kabigting EB, Clifton DK, Steiner RA. Growth hormone receptor messenger ribonucleic acid distribution in the adult male rat brain and its colocalization in hypothalamic somatostatin neurons. *Endocrinology* 1992;131(2):958-63. doi:10.1210/endo.131.2.1353444.

37. Sinha YN, Salocks CB, Wickes MA, Vanderlaan WP. Serum and pituitary concentrations of prolactin and growth hormone in mice during a twenty-four hour period. *Endocrinology* 1977;100(3):786-91. doi:10.1210/endo-100-3-786.

38. Li C, Chen P, Smith MS. Neuropeptide Y and tuberoinfundibular dopamine activities are altered during lactation: Role of prolactin. *Endocrinology* 1999;140(1):118-123. doi:10.1210/en.140.1.118.

39. Kamel F, Wright WW, Mock EJ, Frankel AI. The influence of mating and related stimuli on plasma levels of luteinizing hormone, follicle stimulating hormone, prolactin, and testosterone in the male rat. *Endocrinology* 1977;101(2):421-9. doi:10.1210/endo-101-2-421.

40. Kirk SE, Xie TY, Steyn FJ, Grattan DR, Bunn SJ. Restraint stress increases prolactin-mediated phosphorylation of signal transducer and activator of transcription 5 in the hypothalamus and adrenal cortex in the male mouse. *J. Neuroendocrinol.* 2017. doi:10.1111/jne.12477.

41. Newman SW. The medial extended amygdala in male reproductive behavior. A node in the mammalian social behavior network. *Ann. N. Y. Acad. Sci.* 1999;877:242-257. doi:10.1111/j.1749-6632.1999.tb09271.x.

42. Melcangi RC, Giatti S, Garcia-Segura LM. Levels and actions of neuroactive steroids in the nervous system under physiological and pathological conditions: Sex-specific features. *Neurosci. Biobehav. Rev.* 2015. doi:10.1016/j.neubiorev.2015.09.023.

43. Naftolin F, Ryan KJ, Davies IJ, et al. The formation of estrogens by central neuroendocrine tissues. *Recent Prog. Horm. Res.* 1975;31:295-319.

44. Mitra SW, Hoskin E, Yudkovitz J, et al. Immunolocalization of estrogen receptor α in the

mouse brain: Comparison with estrogen receptor β. *Endocrinology* 2003;144(5):2055-2067.
doi:10.1210/en.2002-221069.

45.   Stanić D, Dubois S, Chua HK, et al. Characterization of aromatase expression in the adult male
and female mouse brain. I. Coexistence with oestrogen receptors α and β, and androgen
receptors. *PLoS One* 2014;9(3):e90451. doi:10.1371/journal.pone.0090451.

46.   Chandrashekar V, Bartke A. Influence of endogenous prolactin on the luteinizing hormone
stimulation of testicular steroidogenesis and the role of prolactin in adult male rats. *Steroids*
1988;51(5-6):559-76.

47.   Dungan HM, Clifton DK, Steiner RA. Minireview: kisspeptin neurons as central processors in
the regulation of gonadotropin-releasing hormone secretion. *Endocrinology*
2006;147(3):1154-8. doi:10.1210/en.2005-1282.

48.   Kokay IC, Petersen SL, Grattan DR. Identification of prolactin-sensitive GABA and kisspeptin
neurons in regions of the rat hypothalamus involved in the control of fertility. *Endocrinology*
2011;152(2):526-35. doi:10.1210/en.2010-0668.

49.   Semaan SJ, Tolson KP, Kauffman AS. The development of kisspeptin circuits in the
Mammalian brain. *Adv. Exp. Med. Biol.* 2013;784:221-52. doi:10.1007/978-1-4614-6199-
9_11.

50.   Drago F, Lissandrello CO. The "low-dose" concept and the paradoxical effects of prolactin on
grooming and sexual behavior. *Eur. J. Pharmacol.* 2000;405(1-3):131-137. doi:10.1016/S0014-
2999(00)00678-6.

51.   Hernandez ME, Soto-Cid A, Rojas F, et al. Prostate response to prolactin in sexually active
male rats. *Reprod. Biol. Endocrinol.* 2006;4:28. doi:10.1186/1477-7827-4-28.

52.   Saito TR, Terada M, Oláh M, Nagy GM. The Role of Prolactin in the Regulation of Male
Copulatory Behavior. In: Nagy, György M., Toth BE, ed. *Prolactin*. InTech; 2013.

doi:10.5772/55515.

53.   Hull EM, Lorrain DS, Du J, et al. Hormone-neurotransmitter interactions in the control of

sexual behavior. *Behav. Brain Res.* 1999;105(1):105-16.

54.   Hernández ML, Fernández-Ruiz JJ, Navarro M, et al. Modifications of mesolimbic and

nigrostriatal dopaminergic activities after intracerebroventricular administration of prolactin.

J. Neural Transm. Gen. Sect. 1994;96(1):63-79.

55.   Lookingland KJ, Moore KE. Effects of estradiol and prolactin on incertohypothalamic

dopaminergic neurons in the male rat. *Brain Res.* 1984;323(1):83-91.

56.   Dugger BN, Morris JA, Jordan CL, Breedlove SM. Androgen receptors are required for full

masculinization of the ventromedial hypothalamus (VMH) in rats. *Horm. Behav.*

2007;51(2):195-201. doi:10.1016/j.yhbeh.2006.10.001.

57.   Kow LM, Pfaff DW. Mapping of neural and signal transduction pathways for lordosis in the

search for estrogen actions on the central nervous system. *Behav. Brain Res.* 1998;92(2):169-

80.

58.   Christensen LW, Nance DM, Gorski RA. Effects of hypothalamic and preoptic lesions on

reproductive behavior in male rats. *Brain Res. Bull.* 1977;2(2):137-41.

59.   Waldherr M, Neumann ID. Centrally released oxytocin mediates mating-induced anxiolysis in

male rats. *Proc. Natl. Acad. Sci. U. S. A.* 2007;104(42):16681-4.

doi:10.1073/pnas.0705860104.

60.   Nishitani S, Moriya T, Kondo Y, Sakuma Y, Shinohara K. Induction of Fos immunoreactivity in

oxytocin neurons in the paraventricular nucleus after female odor exposure in male rats:

effects of sexual experience. *Cell. Mol. Neurobiol.* 2004;24(2):283-91.

61.   Veening JG, de Jong TR, Waldinger MD, Korte SM, Olivier B. The role of oxytocin in male and

female reproductive behavior. *Eur. J. Pharmacol.* 2015;753:209-28.

doi:10.1016/j.ejphar.2014.07.045.

62. Simerly RB, Swanson LW. The organization of neural inputs to the medial preoptic nucleus of

the rat. *J. Comp. Neurol.* 1986;246(3):312-42. doi:10.1002/cne.902460304.

63. Wang J, Swann JM. Connections of the magnocellular medial preoptic nucleus (MPN mag) in

male Syrian hamsters. II. The efferents. *Neuroscience* 2014;274:102-18.

doi:10.1016/j.neuroscience.2014.05.014.

64. Gil M, Bhatt R, Picotte KB, Hull EM. Oxytocin in the medial preoptic area facilitates male

sexual behavior in the rat. *Horm. Behav.* 2011;59(4):435-43.

doi:10.1016/j.yhbeh.2010.12.012.

65. Bridges RS, Ronsheim PM. Prolactin (PRL) regulation of maternal behavior in rats:

bromocriptine treatment delays and PRL promotes the rapid onset of behavior.

Endocrinology 1990;126(2):837-848.

66. Bosch OJ. Maternal aggression in rodents: brain oxytocin and vasopressin mediate pup

defence. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* 2013;368:20130085.

doi:10.1098/rstb.2013.0085.

67. Bosch OJ, Neumann ID. Both oxytocin and vasopressin are mediators of maternal care and

aggression in rodents: From central release to sites of action. *Horm. Behav.* 2012;61(3):293-

303. doi:10.1016/j.yhbeh.2011.11.002.

68. Tachikawa KS, Yoshihara Y, Kuroda KO. Behavioral transition from attack to parenting in male

mice: a crucial role of the vomeronasal system. *J. Neurosci.* 2013;33(12):5120-6.

doi:10.1523/JNEUROSCI.2364-12.2013.

69. Jin D, Liu H-X, Hirai H, et al. CD38 is critical for social behaviour by regulating oxytocin

secretion. *Nature* 2007;446(7131):41-5. doi:10.1038/nature05526.

70.   Akther S, Korshnova N, Zhong J, et al. CD38 in the nucleus accumbens and oxytocin are

related to paternal behavior in mice. *Mol. Brain* 2013;6:41. doi:10.1186/1756-6606-6-41.

71.   Dölen G, Darvishzadeh A, Huang KW, Malenka RC. Social reward requires coordinated activity

of nucleus accumbens oxytocin and serotonin. *Nature* 2013;501(7466):179-84.

doi:10.1038/nature12518.

72.   Otero-García M, Agustín-Pavón C, Lanuza E, Martínez-García F. Distribution of oxytocin and

co-localization with arginine vasopressin in the brain of mice. *Brain Struct. Funct.*

2016;221(7):3445-73. doi:10.1007/s00429-015-1111-y.

73.   Sakaguchi K, Tanaka M, Ohkubo T, et al. Induction of brain prolactin receptor long-form

mRNA expression and maternal behavior in pup-contacted male rats: promotion by prolactin

administration and suppression by female contact. *Neuroendocrinology* 1996;63(6):559-68.

74.   Saltzman W, Ziegler TE. Functional significance of hormonal changes in mammalian fathers. *J.*

Neuroendocrinol. 2014;26(10):685-96. doi:10.1111/jne.12176.

75.   Donhoffner ME, Al Saleh S, Schink O, Wood RI. Prosocial effects of prolactin in male rats:

Social recognition, social approach and social learning. *Horm. Behav.* 2017;96:122-129.

doi:10.1016/j.yhbeh.2017.09.011.

76.   DeMaria JE, Lerant AA, Freeman ME. Prolactin activates all three populations of hypothalamic

neuroendocrine dopaminergic neurons in ovariectomized rats. *Brain Res.* 1999;837(1-2):236-

41.

77.   Björklund A, Moore RY, Nobin A, Stenevi U. The organization of tubero-hypophyseal and

reticulo-infundibular catecholamine neuron systems in the rat brain. *Brain Res.* 1973;51:171-

91.

78.   Goudreau JL, Lindley SE, Lookingland KJ, Moore KE. Evidence that hypothalamic

periventricular dopamine neurons innervate the intermediate lobe of the rat pituitary.

Neuroendocrinology 1992;56(1):100-5.

79. Ma FY, Anderson GM, Gunn TD, Goffin V, Grattan DR, Bunn SJ. Prolactin specifically activates signal transducer and activator of transcription 5b in neuroendocrine dopaminergic neurons. *Endocrinology* 2005;146(12):5112-9. doi:10.1210/en.2005-0770.

80. Elmquist JK, Elias CF, Saper CB. From lesions to leptin: hypothalamic control of food intake and body weight. *Neuron* 1999;22(2):221-32.

81. Fujikawa T, Soya H, Tamashiro KLK, et al. Prolactin prevents acute stress-induced hypocalcemia and ulcerogenesis by acting in the brain of rat. *Endocrinology* 2004;145(4):2006-13. doi:10.1210/en.2003-1446.

82. Torner L. Actions of Prolactin in the Brain: From Physiological Adaptations to Stress and Neurogenesis to Psychopathology. *Front. Endocrinol. (Lausanne).* 2016;7:25. doi:10.3389/fendo.2016.00025.

83. Torner L, Toschi N, Pohlinger A, Landgraf R, Neumann ID. Anxiolytic and anti-stress effects of brain prolactin: improved efficacy of antisense targeting of the prolactin receptor by molecular modeling. *J. Neurosci.* 2001;21(9):3207-14.

84. Davis M, Shi C. The extended amygdala: are the central nucleus of the amygdala and the bed nucleus of the stria terminalis differentially involved in fear versus anxiety? *Ann. N. Y. Acad. Sci.* 1999;877(0077-8923):281-291. Available at: pm:10415655.

85. Aguilera G, Subburaju S, Young S, Chen J. The parvocellular vasopressinergic system and responsiveness of the hypothalamic pituitary adrenal axis during chronic stress. *Prog. Brain Res.* 2008;170:29-39. doi:10.1016/S0079-6123(08)00403-2.

86. Donner N, Neumann ID. Effects of chronic intracerebral prolactin on the oxytocinergic and vasopressinergic system of virgin ovariectomized rats. *Neuroendocrinology* 2009;90(3):315-22. doi:10.1159/000225986.

87. Vega C, Moreno-Carranza B, Zamorano M, et al. Prolactin promotes oxytocin and vasopressin

release by activating neuronal nitric oxide synthase in the supraoptic and paraventricular

nuclei. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 2010;299(6):R1701-8.

doi:10.1152/ajpregu.00575.2010.

88. Blume A, Torner L, Liu Y, Subburaju S, Aguilera G, Neumann ID. Prolactin activates mitogen-

activated protein kinase signaling and corticotropin releasing hormone transcription in rat

hypothalamic neurons. *Endocrinology* 2009;150(4):1841-9. doi:10.1210/en.2008-1023.

FIGURE LEGENDS

Figure 1. Mapping of pSTAT5 immunoreactivity and prolactin receptor expression in the brains of

male and female mice

Semi-schematic camera lucida drawings of coronal sections of the mouse brain showing the

distribution patterns of pSTAT5-ir in female and male mice (left side). and the pattern of PRLR

expression in the brain of an adult male mouse specimen (right side), as determined by PRLR

expression data obtained from the Allen Brain Institute (© 2004 Allen Institute for Brain Science.

Allen Mouse Brain Atlas. Available from: mouse.brain-map.org), experiment 72340223

(http://mouse.brain-map.org/experiment/show/72340223). In the left side, pink dots represent

pSTAT5 expression exclusive of ovariectomized, steroid-primed females, whereas dark blue dots

encode overlapping expression of pSTAT5-ir in both the female and male specimens. Black dots in

the right represent PRLR expression in the male mouse brain. Each dot represents approximately 4

cell nuclei labelled for pSTAT5 or equivalent number of cells expressing the PRLR. Shaded areas

illustrate the counting frames designed for the AVPe/VMPO region (Fig. 1a), the Pa (Fig. 1e), and the

Arc, VMHvl, VMHc and VMHdm, DM and MePD (all frames in Fig. 1f) as part of the quantitative

analysis performed in this work (see below). Approximate distance to bregma is enclosed for each

section.

Figure 2. Representative examples of pSTAT5 immunoreactivity in the brain of female, intact male and castrated male mice

Photomicrographs illustrating pSTAT5 labelling in representative brain sections in an ovariectomized, steroid-primed female supplemented with PRL (I), an intact, PRL-supplemented male (II), a castrated, PRL supplemented male (III) and a intact control male lacking PRL supplementation (IV). Sections correspond to the preoptic hypothalamus (A), the paraventricular region in the anterior hypothalamus (B), the tuberal hypothalamus (C), the caudal septum (D) and the medial posterior amygdala (E). The approximate distance to bregma is indicated in each section. Scale bars correspond to 100 µm.

Figure 3. Quantitative analysis of pSTAT5 immunoreactivity in selected brain regions of female and male mice

Assessment of pSTAT5-ir density (pSTAT5-positive cell nuclei/mm^2) in the major brain regions with expression of pSTAT5-ir in both male and female mice. Bar histograms show mean interhemispheric pSTAT5-ir density ± SEM in gonadally-intact, vehicle-treated males (Male+PRL; n=6; black) and ovariectomized females treated with estradiol and progesterone (Female+PRL; n=6; grey). Counting frames for each of the analyzed nuclei are included in Fig.1. Statistical analysis was applied independently to each brain region (independent t-test for parametric data or Mann-Whitney test for non-parametric data, see Results). *P ≤ 0.05; **P ≤ 0.01; (*) P ≤ 0.06.

Figure 4. Effect of testosterone withdrawal in pSTAT5 immunoreactivity of the male mouse brain

Assessment of pSTAT5-ir density (pSTAT5-positive cell nuclei/mm^2) in gonadally-intact male mice and castrated male mice within the major brain regions showing pSTAT5 expression in the male mouse brain. Bar histograms show mean interhemispheric pSTAT5-ir density ± SEM in intact, PRL-supplemented males (Males+PRL; n=6; black) and castrated, PRL-supplemented males (Castrated+PRL; n=6; white). Counting frames for each of the analyzed nuclei are enclosed in Fig. 2.

1   Statistical analysis was applied independently to each brain region (independent t-test for

2   parametric data or Mann-Whitney test for non-parametric data). *P ≤ 0.05; **P ≤ 0.01.