Aggregation of *Vibrio cholerae* by cationic polymers enhances quorum sensing but over-

rides biofilm dissipation in response to autoinduction

Nicolas Perez-Soto[1,2], Oliver Creese[2,3], Francisco Fernandez-Trillo[2, 3,#], Anne-Marie Krachler[4,#]

[1] School of Biosciences, University of Birmingham, Edgbaston, B15 2TT Birmingham, UK

[2] Institute of Microbiology and Infection, University of Birmingham, Edgbaston B15 2TT

Birmingham, UK

[3] School of Chemistry, University of Birmingham, Edgbaston, B15 2TT Birmingham, UK

[4] University of Texas Health Science Center at Houston, McGovern Medical School, Department

of Microbiology and Molecular Genetics, 6431 Fannin St, Houston TX 77030, USA

[#] Corresponding authors: Francisco Fernandez-Trillo (f.fernandez-trillo@bham.ac.uk) and Anne-

Marie Krachler (anne.marie.krachler@uth.tmc.edu)

Keywords: *Vibrio cholerae*, cholera, quorum sensing, aggregation, biofilm, autoinducer, surface

sensing;

Abstract

Vibrio cholerae is a Gram-negative bacterium found in aquatic environments and a human pathogen of global significance. Its transition between host-associated and environmental life styles involves the tight regulation of niche-specific phenotypes such as motility, biofilm formation and virulence. *V. cholerae*'s transition from the host to environmental dispersal usually involves suppression of virulence and dispersion of biofilm communities. In contrast to this naturally occurring transition, bacterial aggregation by cationic polymers triggers a unique response, which is to suppress virulence gene expression while also triggering biofilm formation by *V. cholerae*, an artificial combination of traits that is potentially very useful to bind and neutralize the pathogen from contaminated water. Here, we set out to uncover the mechanistic basis of this polymer-triggered bacterial behavior. We found that bacteria-polymer aggregates undergo rapid autoinduction and achieve quorum sensing at bacterial densities far below those required for autoinduction in the absence of polymers. We demonstrate this induction of quorum sensing is due both to a rapid formation of autoinducer gradients and local enhancement of autoinducer concentrations within bacterial clusters, as well as the stimulation of CAI-1 and AI-2 production by aggregated bacteria. We further found that polymers cause an induction of the biofilm specific regulator VpsR and the biofilm structural protein RbmA, bypassing the usual suppression of biofilm during autoinduction. Overall, this study highlights that synthetic materials can be used to cross-wire natural bacterial responses to achieve a combination of phenotypes with potentially useful applications.

Introduction

Both natural and synthetic cationic macromolecules, such as cationic antimicrobial peptides, cationic polymers and dendrimers, have been extensively reported as antimicrobial.[1] Due to their positive charge, these polymers can efficiently bind the negatively charged envelope of Gram-negative and Gram-positive bacteria.[2] At high concentrations and charge densities, these molecules have the potential to interfere with membrane integrity and decrease bacterial viability.[1] However, antimicrobial activity is heavily dependent on the length and nature of the polymer and, more importantly, on the nature of the targeted microbe. At low concentrations, cationic polymers are still capable of causing bacterial aggregation by mediating electrostatic interactions, but do so without significantly affecting bacterial membrane integrity and growth.[3]

We and others have previously reported that bacteria clustered by sub-inhibitory concentrations of cationic polymers display interesting phenotypes resembling those of biofilm communities.[3] For instance, we have recently demonstrated that clustering of the diarrheal pathogen *Vibrio cholerae* with methacrylamides containing primary or tertiary amines leads to accumulation of biomass and extracellular DNA, and represses ToxT-regulated virulence factors including cholera toxin.[3f] However, what drives these phenotypic changes in response to polymer exposure remains unclear. Similarly, the marine bacterium *Vibrio harveyi* shows enhanced bioluminescence in response to polymer-mediated clustering.[3a, 3b, 3e] The bacterial components necessary to produce luminescence are encoded by the *luxCDABE* genes and subject to complex regulatory mechanisms. A major regulatory cascade controlling luminescence is quorum sensing. Since the regulators controlling luminescence are functionally conserved between *Vibrio* species, expression of *V. harveyi luxCDABE* genes can be used as a tool to probe quorum sensing in heterologous hosts.[4]

In *V. cholerae*, four parallel quorum sensing pathways operate,[5] each governed by a autoinducer synthase, which produces a small, freely diffusible molecule that is released in the environment and can be sensed by a corresponding sensor/kinase component that controls the activity of the LuxU/LuxO phosphorelay. Of the four pathways, the LuxS/LuxPQ system, which produces and detects the inter-species autoinducer AI-2 (S-TMHF-borate) and the CqsA/CqsS system, which produces and senses the *Vibrio* specific autoinducer CAI-1 (S-3-hydroxytridecan-4-one), are the best characterized. While CqsA and LuxS synthesize the autoinducers, the hybrid sensor/kinases CqsS and LuxQ sense and respond to their cognate autoinducers by dephosphorylating and inactivating the regulator LuxO. When extracellular autoinducer concentrations are low, LuxO is phosphorylated and activates the transcription of the small RNAs Qrr1-4, which in turn inhibit the expression of HapR, a master regulator controlling diverse cellular functions including luminescence, biofilm formation and virulence.[6] When the extracellular autoinducer concentration increases above a threshold, the sensor/kinases instead act as phosphatases, leading to dephosphorylation and inactivation of LuxO, ultimately allowing the expression of HapR. Consequently, when autoinducer concentration is high, HapR activates luminescence, but suppresses virulence and biofilm genes. This regulatory mechanism is thought to mediate the dissipation of host-associated biofilms and enable transmission of *V. cholerae* from the intestine to the environment.[7] Since in *V. harveyi*, exposure to sub-inhibitory concentrations of cationic polymers lead to enhanced luminescence, here we set out to study whether *V. cholerae* would also activate quorum sensing in response to polymer-mediated clustering, and how quorum sensing could be related to the observed phenotypic changes, including lowered virulence and enhanced biofilm formation.

Results

Polymers enhance quorum sensing in *V. cholerae*. First, we set out to test if clustering by the cationic polymers poly(*N*-[3-aminopropyl] methacrylamide), (P1, Figure 1A) and poly(*N*-[3-(dimethylamino)propyl] methacrylamide), (P2, Figure 1B) induced quorum sensing in *V. cholerae*. As previously shown and detailed here by N-SIM super-resolution microscopy, even low concentrations of polymers induced rapid clustering of bacteria without affecting bacterial viability (Figure 1C). In order to create a fast and direct read-out for quorum sensing, the cosmid pBB1, which contains the *luxCDABE* luminescence genes from *V. harveyi*,[8] was used to transform *V. cholerae* serogroup O1 biovar El Tor strain A1552 (serotype Inaba). The transformed bacteria were then grown in LB for 20 hours, washed with artificial marine water (AMW) and adjusted to an OD of 0.2 in AMW alone, or AMW containing polymers at concentrations ranging from 0.005 to 0.5 mg/ml. In the absence of polymer, luminescence as a read-out of quorum sensing first decreased due to back-dilution of the culture from a high density overnight culture to OD 0.2, and then gradually increased due to accumulation of autoinducers, peaking at 4.5 hours (Figure 1D and E, black traces). Interestingly, the quorum induction kinetics were significantly different in the presence of either P1 (Figure 1D) or P2 (Figure 1E), with initial luminescence being sustained and reaching a higher second peak at around 2.5 hours, as opposed to 4.5 hours in the absence of polymers. The magnitude of induction was higher in polymer-treated cultures (approx. 5-fold at peak quorum induction) compared to untreated cultures. Results were similar for the Ogawa serotype strain E7646, in that quorum sensing was initially sustained, and then further enhanced (approx. 4-fold at peak quorum induction) by polymer-mediated bacterial aggregation (Figure 1F, G). In both strains, the magnitude of luminescence induction was independent of the polymer concentration used, suggesting a threshold response.

To assess viable bacterial counts at the experimental end point, samples were serially

diluted in high-salt buffer to disrupt clusters as previously described,[3f] and plated. Similar numbers

of colony forming units were recovered from untreated or polymer-treated samples (Figure 1H, I),

suggesting that the presence of polymers did not affect bacterial viability or proliferation, in

agreement with our previous data.[3f] Taken together, our data demonstrate that these cationic

polymers that cluster *V. cholerae* modulate the community behavior to give an accelerated and

more robust autoinduction.

Figure 1. **Polymers enhance quorum sensing in *Vibrio cholerae*.** Chemical structures of P1 (A) and P2 (B). N-SIM

super-resolution image of LIVE/DEAD-stained *V. cholerae* A1552 clustered by P1 (C). *V. cholerae* El Tor strains

A1552 (D, E, H) or E7646 (F, G, I) containing the luminescence reporter pBB1 were adjusted to an OD_{600} of 0.2

following 16 hrs of growth and incubated in AMW alone (black) or AMW containing polymer P1 (D, F) or P2 (E, G)

at concentrations of 0.005 (green), 0.05 (blue), or 0.5 (red) mg/ml. Luminescence was recorded every 30 min for 10

hrs and plotted as means ± s.e.m from at least three biological replicates. To test the effect of polymers on bacterial

viability, samples were removed after 10 hrs, serially diluted and plated on LB (H, I).

Bacterial density shapes the kinetics of quorum induction in response to polymer.

Since quorum sensing is usually tightly linked to bacterial density, we set out to explore how initial

culture density affects quorum induction in the presence of polymers. *V. cholerae* carrying pBB1

as a quorum sensing reporter was adjusted to optical densities of 0.005, 0.05 and 0.5 in AMW

alone, or AMW containing 0.005-0.5 mg/ml P1 or P2, and luminescence was monitored (Figure

2). The onset of autoinduction was not significantly modulated by the addition of polymers P1 or

P2 to higher density cultures (OD_{600} of 0.5), but a ~ six-fold (P1) to nine-fold (P2) increase in peak

luminescence was observed (Figure 2A, D). At lower culture densities, autoinduction of the

untreated cultures was less pronounced (0.05) and eventually ceased (OD 0.005) since not enough

autoinducer accumulated to reach the threshold concentration. In the presence of polymers, the

initial quorum present in the culture was sustained, even in very dilute cultures, and the

enhancement in peak luminescence was much more pronounced compared to untreated cultures

(Figure 2B-F). Interestingly, clustering of very dilute cultures led to two peaks in luminescence,

with a gap of approximately 4 hours (Figure 2C, F). Of note, the luminescence was not a result of

bacterial growth, which was negligible under the observed conditions (AMW) and over the time

frame described, both at high and low initial densities (Figure 2G, H).

Figure 2. Bacterial density shapes the kinetics of quorum induction in response to polymer. Cultures of *V.*

cholerae A1552 containing pBB1 were grown for 16 hrs and diluted into AMW alone (black) or AMW containing

0.005 (green), 0.05 (blue) or 0.5 (red) mg/ml of P1 (A-C) or P2 (D-F). Bacterial densities were adjusted to result in

OD_{600} values of 0.5 (A, D), 0.05 (B, E), and 0.005 (C, F), respectively. Luminescence and OD_{600} were recorded every

30 min for 10 hrs and values are means ± s.e.m from at least three biological replicates. No significant growth was

detected over this time frame, either at initial densities of 0.5 (G) or 0.005 (H).

To visualize the process of bacterial clustering and luminescence induction, *V. cholerae*

A1552 containing pBB1 was incubated in the presence of P1, P2, or AMW alone in glass-bottom

plates. Bacteria were imaged every 30 minutes to simultaneously visualize clustering and

luminescence induction. With both P1 and P2 bacterial clusters were observed within minutes and

remained stable over the duration of the experiment (Figure 3). Luminescence appears to originate

from and to be restricted to bacterial clusters. Quantification of luminescence based on integrated

pixel intensities showed a polymer-mediated enhancement of autoinduction (Figure 3H), in

agreement with spectroscopic data.

Figure 3. **Polymers enhance quorum sensing in *Vibrio cholerae*.** *V. cholerae* A1552 containing pBB1 was grown for 16 hrs and then adjusted to an OD600 of 0.2 either in the presence of P1 (A-C), P2 (D-F) or AMW alone (G). Polymers were adjusted to final concentrations of 0.005 (A, D), 0.05 (B, E), or 0.5 (C, F) mg/ml in AMW. Luminescence of samples was imaged every 30 min for 15 hrs and representative images for each time point are shown (0 hrs, top left to 15 hrs (bottom right) of each panel. Luminescence intensities over time were analyzed by quantifying pixel intensities, and means ± s.e.m from at least three biological replicates are shown for P1 (left) and P2 (right panel), (H).

Polymer-mediated luminescence is not due to nutrient starvation within clusters.

While autoinduction controls HapR and luminescence via the regulator LuxO, other environmental

cues, including nutrient availability, have been reported to feed into the LuxO signaling cascade

and thus have the potential to affect luminescence. Nutrient sensing and the LuxO signaling

pathway converge at the cyclic AMP (cAMP) receptor protein, CRP.[9] During limitation of PTS

sugars such as glucose, cAMP-CRP is capable of modulating LuxO activation by affecting the

expression of autoinducer synthases.[10] Since clustering of bacteria by polymers may limit nutrient

access and induce starvation, we tested whether a CRP deletion strain would be capable of

activating luminescence in the presence of polymers. The *V. cholerae* Δ*crp* strain was transformed

with pBB1 to monitor luminescence. However, the culture produced very low levels of

luminescence, both in the presence and absence of polymers (Figure 4A, B), which is in agreement

with previous work on *V. fischeri* CRP.[11] While this suggests that cross-talk between CRP and

LuxO signaling is a dominant cue for luminescence induction in *V. cholerae* as well, this made it

unfeasible to determine whether the presence of polymers induced a state of carbon starvation,

leading to luminescence induction via CRP. Instead, we repeated the experiment using the pBB1

containing *V. cholerae* wild type strain and supplementing the AMW with additional glucose. If

clustering would limit nutrient diffusion, increasing the nutrient concentration should be able to

overcome this and revert the bacteria to a non-luminescent phenotype. However, polymers still

enhanced and sustained luminescence in the presence of 1% glucose, to a similar extent as in AMW

alone (Figure 4C), suggesting that the effect was not due to nutrient limitation in the clusters.

Figure 4. Polymer-mediated luminescence is not due to nutrient starvation within clusters. *V. cholerae* E7646

Δ*crp* containing the luminescence reporter pBB1 was grown for 16 hrs and then diluted to an OD_{600} of 0.2 in AMW

alone (black) or AMW containing P1 (A) or P2 (B) at concentrations of 0.005 (green), 0.05 (blue) or 0.5 (red) mg/ml.

Luminescence was recorded every 30 min for 10 hrs and means ± s.e.m from at least three biological replicates are

shown. (C) *V. cholerae* E7646 wild type containing pBB1 was grown for 16 hrs and adjusted to an OD_{600} of 0.2 in

AMW containing 1% glucose alone (orange), or in the presence of 0.5 mg/ml P1 (light blue) or P2 (dark blue).

Luminescence was recorded every 30 min for 10 hrs and means ± s.e.m from at least three biological replicates are

shown.

Polymer mediated enhancement of quorum sensing is dominated by CAI-1. In *V.*

cholerae, at least four parallel quorum sensing pathways converge to control the activity of the

quorum sensing regulator, LuxO and thus, quorum regulated phenotypes including biofilm

formation and virulence.[5] We set out to test whether the polymer mediated effect on quorum

sensing was specific for any one pathway. *V. cholerae* wild type strain A1552 was mixed with

equal numbers of cells of quorum sensing mutants transformed with pBB1 either deficient in the

production of AI-2 (DH231), or CAI-1 (WN1103), to give a final OD_{600} of 0.2. Additionally, the

mutants were unable to sense the presence of AI-2 or CAI-1, respectively. Cultures producing less

AI-2 produced a luminescence profile similar in shape and magnitude to quorum sensing proficient

cells in response to polymer P1 addition (Figure 5A). In contrast, cultures producing less CAI-1

still had a comparable response profile upon addition of P1, but the magnitude of the luminescence

enhancement was decreased (Figure 5B)_ compared to cultures producing less AI-2 (Figure 5A).

In co-cultures producing less of both autoinducers (containing strain BH1578), the luminescence

response to polymer was abolished (Figure 5C). Similarly, co-cultures containing a low-density

locked mutant of the down-stream quorum regulator LuxO (LuxO^{D47E}) showed no luminescence

induction upon addition of polymer P1 (Figure 5D). These data pointed at both autoinducers being

involved in the quorum sensing enhancement in response to polymer, with CAI-1 being the

dominant inducer. Additionally, the luminescence response to the polymer seems to proceed

through the canonical LuxO-dependent pathway.

Figure 5. Polymer mediated enhancement of quorum sensing is mainly driven by CAI-1. *V. cholerae* A1552 wild

type (dark) and quorum sensing mutants containing pBB1 were grown for 16 hrs, and then diluted into AMW to give

equal cell densities and a total OD_{600} of 0.2. Strains were grown together in AMW alone (black) or AMW containing

P1 at 0.005 (green), 0.05 (blue) or 0.5 (red) mg/ml. Luminescence was recorded every 30 min for 10 hrs and means ±

s.e.m from at least three biological replicates are shown. Mutants grown in co-culture with the wild type were (A)

DH231 (Δ*luxS*Δ*cqsS*) producing no AI-2, (B) WN1103 (Δ*luxQ*Δ*cqsA*) producing no CAI-1, (C) BH1578

(Δ*luxS*Δ*cqsA*) producing no AI-2 or CAI-1, and (D) BH1651 (luxO^{D47E}).

Enhanced quorum sensing is driven by enhanced production of autoinducers in

response to polymers. It has been described that at least under some conditions, quorum sensing

may be subject to positive feedback, where quorum induction leads to increased production of one

of the autoinducer synthases.[12] Therefore, the enhancement in luminescence in response to

polymers could be due to positive feedback, as a result of the polymers increasing the local

concentration of autoinducers above the threshold. Alternatively, the polymers could have a direct

effect on the production of autoinducers. We set out to test this by establishing a reporter assay

that allowed us to decouple quorum sensing from the production of autoinducers. For this assay,

we used as a luminescence reporter a *V. cholerae* strain transformed with pBB1 that could sense

both CAI-1 and AI-2, but could not produce either molecule (BH1578). This reporter strain was

exposed to supernatants from producer strains grown under different conditions, to evaluate the

effect of the polymer. Initially, we evaluated the assay by growing the reporter in the presence of

supernatants harvested from wild type *V. cholerae*, or strains incapable of producing either AI-2,

CAI-1, or both autoinducers. Supernatant harvested from the quorum proficient wild type strain

grown to high cell density triggered the highest level of luminescence in BH1578 (Figure 6A). The

luminescence triggered by the AI-2 deficient strain was slightly decreased (Figure 6B), whereas

luminescence was significantly decreased in response to supernatant from the CAI-1 deficient

strain (Figure 6C) and was abolished in response to the strain deficient in both CAI-1 and AI-2

production (Figure 6D). Hence, the assay was capable of detecting different levels of autoinducers

produced by a second strain.

We took this assay forward and harvested supernatants from wild type cells exposed to

AMW alone, or AMW containing 0.005-0.5 mg/ml polymers P1 or P2, and exposed the reporter

strain to filtered supernatants to test if the levels of autoinducers produced by the wild type strain

in the presence of polymers were different. The reporter strain was not clustered under the assay

conditions. While wild type cells exposed to AMW alone did not produce a detectable amount of

autoinducer and thus, no significant luminescence reading in the reporter strain, both polymers P1

and P2 enhanced the production of autoinducers by wild type *V. cholerae*, leading to an increase

in luminescence upon exposure of the reporter to the supernatants (Figure 6E, F). These data

demonstrate that polymer exposure leads to enhanced production of autoinducers by the bacteria.

Figure 6. Enhanced quorum sensing is driven by enhanced production of autoinducers in response to polymers.

Cultures of *V. cholerae* were adjusted to an OD of 0.2, grown for 16 hrs in LB medium, and supernatants were

harvested, filtered, and incubated with *V. cholerae* BH1578 containing pBB1. Strains used to harvest supernatants

were (A) wild type A1552 (B) DH231 ($\Delta luxS\Delta cqs$), (C) WN1103 ($\Delta luxQ\Delta cqsA$), and (D) BH1578 (($\Delta luxS\Delta cqsA$).

Luminescence was recorded every 30 min for 10 hrs and means ± s.e.m from at least three biological replicates are

shown. *V. cholerae* wild type was adjusted to an OD_{600} of 0.2 in AMW alone or AMW containing 0.005-0.5 mg/ml

P1 (E) or P2 (F) and supernatants harvested and filtered 16 hrs later. To determine their autoinducer content,

supernatants were incubated with the reporter strain *V. cholerae* BH1578 containing pBB1. Luminescence was

recorded every 30 min for 10 hrs and means ± s.e.m from at least three biological replicates are shown.

Polymer-mediated quorum induction over-rides the canonical biofilm dissipation

programme in *V. cholerae*. Contrary to many bacteria that use quorum signaling as a means to

induce biofilm formation, in *V. cholerae* autoinduction promotes repression of biofilm production

and dissemination, via the regulator HapR.[13] However, we had previously observed enhanced

accumulation of *V. cholerae* upon prolonged exposure to cationic polymers, but whether this was

accompanied by transcriptional changes at the level of biofilm production was not known. Thus,

we grew *V. cholerae* containing transcriptional fusions to promoters regulating key biofilm

components in the presence or absence of P1 and P2 (Figure 7). *V. cholerae* biofilms contain the

structural protein RbmA and require the regulator VpsR, which controls the expression of the *vps*

polysaccharide biosynthesis genes.[14] Upon exposure to either P1 or P2, *vpsR* and *rbmA* were both

significantly induced (Figure 7A, B), suggesting that upon polymer-mediated clustering, quorum

sensing does not, as usually, suppress genes involved in biofilm production, but instead their

transcription is enhanced. AphA, which is another direct target and is usually induced by VpsR, is

suppressed in the presence of polymers (Figure 7C).

Figure 7. Polymer-mediated quorum induction over-rides the canonical biofilm dissipation programme in *V.*

***cholerae*.** *V. cholerae* wild type A1552 containing pRW50T lacZ reporters for the promoters of *rbmA* (A), or *vpsR*

(B), were grown for 16 hrs and then diluted into AMW alone or containing 0.005 or 0.05 mg/ml P1 or P2 as indicated,

to give an OD_{600} of 0.2. Following 16 hrs of incubation, clustered bacteria were removed and either processed for

beta-galactosidase assays or treated with high-salt PBS to disperse the cultures for OD_{600} measurements.

Transcriptional activities were calculated and normalized to untreated cultures. Shown are means ± s.e.m and

individual measurements for three biological replicates. Statistical significance was determined by ANOVA and a

Dunnett's multiple comparison test and is depicted as (****) for p-values ≤ 0.0001, (***) p≤0.001, (**) p≤0.01 (*)

p≤0.05 and ns or not significant (p≥0.05). (C) *V. cholerae* wild type A1552 containing pRW50T lacZ reporter for the

aphA promoter was grown in AMW alone (orange) or containing P1 (light blue) at 0.05 mg/mL or P2 (dark blue) at

0.5 mg/mL for 18 hrs. Clustered bacteria were removed at indicated times and either processed for beta-galactosidase

assays or treated with high-salt PBS to disperse the cultures for OD_{600} measurements. Transcriptional activities were

calculated and normalized to the activities of untreated cultures at 2hrs. Shown are means ± s.e.m and individual

measurements for three biological replicates.

Discussion

Traditionally, work on cationic polymers has been carried out with the development of

antimicrobial materials as a main goal.[1] However, recent work by our groups and others has

demonstrated that such cationic polymers can be titrated against bacteria, to achieve a charge

balance that allows for the rapid and efficient clustering of bacteria but avoids membrane

disruption and bacterial cell death.[3]

The use of cationic polymers to induce rapid bacterial clustering in this way has proven as

an interesting path to study effects of cell aggregation and crowding on bacterial physiology. While

such behaviors are often studied in batch cultures, by incubating bacterial cultures over a prolonged

time, this means aggregation is accompanied by bacterial growth, and eventually nutrient

limitation, which makes it difficult to establish the primary cause of the observed phenotypes. In

contrast, cationic polymers induce cell aggregation rapidly, within minutes, which allows us to

study these phenomena independent of cellular proliferation and nutrient limitation.

We and others have previously observed that cationic polymers and dendrimers can, under

certain conditions, trigger bioluminescence in the marine bacterium *V. harveyi*, suggesting they

may induce or enhance quorum sensing.[3a, 3b, 3d, 3e] In a more recent study where we extended this

work to the human pathogen *V. cholerae*, we observed that polymer-mediated clustering led to

enhanced deposition of biomass and extracellular DNA, while it interfered with the induction of

virulence genes in an infection model.[3f] Since virulence and biofilm production are both regulated

by quorum sensing but are usually both regulated concurrently, the goal of this study was to test

whether cationic polymers would trigger quorum sensing in *V. cholerae* and how this would affect

down-stream transcription of biofilm genes.

We used *V. cholerae* strains heterologously expressing the *luxCDABE* luminescence genes

(on cosmid pBB1) from *V. harveyi* to be able to use luminescence as direct readout for

autoinduction. Over 16 hours, *V. cholerae* would grow to high cell densities and as a result, was

strongly luminescent. On dilution into artificial marine water, cell density and autoinducer

concentration would rapidly decrease, resulting in a decline in luminescence. After several hours,

cells would eventually accumulate sufficient autoinducer to reach the quorum threshold and induce

luminescence again. This behavior was observed in AMW alone (Figure 1) and is in agreement

with commonly observed results from such experiments.[5, 12] In contrast, when cells were diluted

into media containing polymers, they would undergo extensive clustering almost instantaneously,

and luminescence readouts never dropped, but instead, further increased immediately (Figure 1),

suggesting that clustering not only countered the dilution effect, but further increased autoinducer

concentration within the clusters. Interestingly, this behavior was observed over a broad space of

cell densities (at least two orders of magnitude), including in dilute cultures that did not by

themselves experience autoinduction (Figure 2C, F), suggesting that during clustering, polymers

create pockets containing strongly increased concentrations of autoinducers around bacterial

aggregates.

We further demonstrated that both CAI-1 and AI-2 dependent quorum sensing cascades

are activated in response to polymers and that clustering leads to an enhanced production of both

autoinducers (Figures 5 and 6). The effect of the *Vibrio*-specific autoinducer CAI-1 dominated the

clustering-driven luminescence phenotype (Figure 5), in line with previous results obtained for

batch-cultures of *V. cholerae* in rich medium.[12]

Some studies have hypothesized that luminescence could be a result of limited diffusion of

nutrients in the polymer-mediated bacterial aggregates.[3d] Catabolite repression of luminescence

has been reported for *V. fischeri*, where cAMP-CRP stimulates *luxCDABE* expression.[15] However,

this effect is alleviated by high concentrations of autoinducer.[11] In our hands, CRP was essential

for luminescence, both triggered by high cell density in the absence of polymers, in line with

previous findings for an *E. coli* Δ*crp* mutant,[11] as well as in response to polymer-induced

clustering. Additionally, supplementation of the media with excess glucose did not quench

luminescence, even in the absence of autoinduction. This suggests that nutrient limitation within

the clusters is not a major cue for luminescence induction, but further underpins that cross-talk

between nutrient sensing and quorum sensing pathways exists.

Finally, we followed up on our earlier observation that exposure to cationic polymers

causes deposition of *V. cholerae* on inorganic surfaces and release of extracellular DNA, both

hallmarks of biofilm formation.[3f] Here, we showed that this phenotype is the result of

transcriptional activation of genes involved in biofilm production in response to polymer exposure.

Biofilm induction may explain the enhanced resistance towards antimicrobials of bacteria that

have been exposed to cationic polymers, as previously described by others.[3d] The expression of

the biofilm regulator VpsR and the biofilm structural protein RbmA were both induced upon

exposure to the polymers (Figure 7). This upregulation is in contrast to the canonical biofilm

regulation where biofilm genes are repressed during autoinduction. VpsR is a master regulator of

biofilm formation and a two component system response regulator. Although no cognate histidine

kinase has been identified, VpsR is epistatic to the intracellular hybrid sensor histidine kinase

VpsS.[16] Induction of *vpsR* likely leads to the downstream induction of *rbmA* we observed here,

since *rbmA* is a direct target of VpsR regulation.[17] However, *vpsR* induction in the presence of

polymers seems to happen despite autoinduction, which should normally lead to suppression of

vpsR. What is also different from a regular biofilm response is that VpsR in the presence of

polymer, fails to upregulate one of its other direct targets, *aphA*. We showed that in contrast to this

canonical response, *aphA* is strongly suppressed by the presence of polymers (Figure 7C). When

Shikuma et al. identified VpsS as a regulator of VpsR, they established the existence of a pathway

that proceeds from VpsS through the quorum regulators LuxU and LuxO and results in the VpsR

dependent activation of biofilm production, independent of HapR.[16] It may be that in the presence

of polymers, this pathway is active and dominates the effects of the CAI-1 and AI-2 pathways on

biofilm. Unfortunately, the cognate signal activating VpsS is as yet unidentified.

Conclusions

We showed here that clustering of *V. cholerae* in response to cationic polymers leads to

autoinduction, due to a rapid increase of local autoinducer concentration in the vicinity of

aggregated bacteria. Moreover, we demonstrate that stimulation of further autoinducer synthesis

is also observed and involves at least two of the four known quorum sensing systems, CAI-1 and

AI-2. We speculate that the third quorum sensing pathway, which proceeds through the

intracellular hybrid sensor kinase VpsS [5, 16] is also activated, and leads to the production of biofilm

in response to polymer driven aggregation. Our previous work together with the data presented

here rules out membrane disruption and nutrient limitation within clusters, respectively, as cues

leading to the phenotypes observed here. Our future work will aim to further dissect the pathway(s)

triggered in response to polymer exposure, to clarify whether VpsS is indeed involved, and is

activated in response to polymers. Polymeric materials that inhibit bacterial dissemination, both

mechanically and transcriptionally, may be useful for applications to enhance waste water

treatment for decontamination of water from *V. cholerae*.

Materials and Methods

Bacterial strains and culture conditions. *V. cholerae* El Tor strains used in this study (Table 1)

were derived from A1552 [18] and E7946.[19] The *E. coli* K12 strains JCB387 [20] DH5α [21] and SM10

λpir [22] were used for general cloning and conjugation procedures. Strains were propagated at 37

ºC in lysogeny broth (LB) supplemented with 10 μg/μL tetracycline or 30 μg/μL kanamycin for

selection when required. Plasmids were introduced into *V. cholerae* strains by triparental mating

with *E. coli* DH5a carrying the desired plasmid (donor) and *E. coli* SM10 (helper strain) carrying

the conjugative machinery on pRK2013. Cultures were mixed at a volumetric ratio of 1:2:2 of

recipient:helper:donor in 250 µl and spotted onto brain-heart infusion (BHI) agar to be incubated

overnight at 37 °C. Spots of bacteria were dislodged after an overnight incubation and resuspended

in 3 mL of sterile PBS. 100 μL of serial dilutions were plated onto TCBS plates containing 10

μg/μL of tetracycline. Resulting colonies were checked by PCR in the case of pRW50T constructs,

while pBB1 transconjugants were screened for luminescence.

Beta-galactosidase assays. pRW50T derivative construction was described before.[3f] Regions

encoding *aphA, rbmA*, or *vpsR* promoters were amplified by PCR and cloned into pRW50T using

EcoRI and HindIII sites. The insertion was checked by PCR using external primers. Measurement

of β-galactosidase activity as a readout for transcriptional activity was done as previously

described [23], with some modifications to accommodate testing of aggregated bacteria. Small

cultures of reporter strains were grown in the absence or presence of polymers P1 and P2 and

incubated overnight at 37 ºC with shaking. Clustered bacteria were split in two, and either used for

transcriptional assays or washed with high salt PBS (200 mM NaCl) to disrupt aggregation and

enable OD_{600} measurements.

Luminescence assays. Luminescence assays were done using *V. cholerae* pBB1 transconjugants.

The pBB1 cosmid[24] was introduced into *V. cholerae* strains by triparental mating in the same

conditions as for pRW50T. Overnight cultures of *V. cholerae* pBB1 were adjusted to OD_{600} of 0.5,

0.1 and 0.01 in artificial marine water with 10 μg/μL of tetracycline. Polymers P1 and P2 were

added at concentrations of 0.005 mg/mL, 0.05 mg/mL and 0.5 mg/mL in DMEM or AMW, in 200

μL final volume using a dark wall clear bottom 96-well plate. Plate was incubated up to 15 hours

at 37ºC with shaking a 200 rpm, while luminescence and OD_{600} were recorded every 30 minutes

using a FLUOstar Omega plate reader. The following assays were done with bacterial cultures

with OD_{600} adjusted to 0.2. Cells were recovered after the assay and washed with high-salt PBS

containing 200 mM NaCl to disrupt charge-based aggregation, and plated onto LB with

tetracycline to determine the viability. Plates were imaged using a BioRad Gel Doc XR System

and images were processed with ImageJ.

Luminescence time-lapse imaging. Overnight culture of *V. cholerae* A1552 pBB1 was diluted to

an OD_{600} of 0.2 in artificial marine water or clear DMEM with 10 μg/μL of tetracycline, and

polymers at concentrations of 0.005 mg/mL, 0.05 mg/mL and 0.5 mg/mL. Samples were prepared

in 200 μL using a glass-bottom 96-well plate and incubated at 37 ºC with 5% CO_2 in a microscope

imaging chamber. Images were taken every 30 minutes with 10 seconds of exposure at 40X

magnification, using an Evolve 512 EMCCD camera mounted on a Nikon-Eclipse TE2000-U

microscope. Image acquisition was done using Nikon NIS-Elements software and final images

processed with ImageJ. Pixel intensity was determined from several clusters within frame using

ImageJ.

Super resolution microscopy of bacterial clusters. *V. cholerae* A1552 was incubated with 0.05

mg/ml P1 in PBS for 1 hour. To visualize membrane integrity, the sample was stained using the

LIVE/DEAD BacLight Bacterial Viability Kit (Life Technologies) for 10 minutes at room

temperature. The sample was mounted with ProLong Gold Antifade Mountant and covered with a

cover slip. Images were taken on a Nikon N-SIM super resolution microscope fitted with SR Apo

TIRF 100x lens, at 100 ms exposure. Deconvolution was carried out using the Nikon NIS elements

software.

Luminescence assays using *V. cholerae* BH1578 pBB1 as a reporter. *V. cholerae* BH1578

pBB1 was used to determine the effect of polymers on the production of autoinducers. *V. cholerae*

strains at an OD_{600} of 0.2 were clustered with polymers at concentrations of 0.005 mg/mL, 0.05

mg/mL and 0.5 mg/mL in artificial marine water. Supernatants were recovered by centrifugation,

and used to resuspend *V. cholerae* BH1578 pBB1 previously adjusted to an OD_{600} of 0.2 in 200

μL. Luminescence was recorded at 37 °C using a FLUOStar Omega plate reader. Similarly, *V.*

cholerae strains and *V. cholerae* BH1578 pBB1 were co-cultured in 200 μL final volume and

polymers added at concentrations of 0.005 mg/mL, 0.05 mg/mL and 0.5 mg/mL in artificial marine

water. Both strains were adjusted to a final OD_{600} of 0.1 each (0.2 total density). Incubation was

done at 37 °C with shaking at 200 rpm, and luminescence and OD_{600} were measured every 30

minutes using a FLUOStar Omega plate reader.

Acknowledgements

We would like to thank B. Bassler for the generous gift of quorum sensing mutants BH1578, DH23, WN1103 and the cosmid pBB1. We would like to thank H. Kaplan, the Krachler and Fernandez-Trillo labs for critical reading of the manuscript, and for suggestions on how to improve this study. This work was supported by University of Birmingham Fellowships (to A.M.K. and F.F.-T.), Wellcome Trust grant 177ISSFPP (to A.M.K. and F.F.-T), BBSRC grants BB/M021513/1 (to A.M.K.) and BB/L007916/1 (to A.M.K.), a CONICYT fellowship (to N.P.-S.), BBSRC MIBTP scholarship BB/M01116X/1 (to O.C.) and a UT Systems Science and Technology Acquisition and Retention Award (to A.M.K.).

Competing Financial Interests

The authors declare no competing financial interests.

References

1.   (a) Hartlieb, M.; Williams, E. G. L.; Kuroki, A.; Perrier, S.; Locock, K. E. S., Antimicrobial Polymers: Mimicking Amino Acid Functionali ty, Sequence Control and Three-dimensional Structure of Host-defen se Peptides. *Current Medicinal Chemistry* **2017,** *24* (19), 2115-2140; (b) Palermo, E. F.; Kuroda, K., Chemical structure of cationic groups in amphiphilic polymethacrylates modulates the antimicrobial and hemolytic activities. *Biomacromolecules* **2009,** *10* (6), 1416-28; (c) Tew, G. N.; Liu, D.; Chen, B.; Doerksen, R. J.; Kaplan, J.; Carroll, P. J.; Klein, M. L.; DeGrado, W. F., De novo design of biomimetic antimicrobial polymers. *Proceedings of the National Academy of Sciences of the United States of America* **2002,** *99* (8), 5110-4.
2.   (a) Cevc, G., Membrane electrostatics. *Biochimica et Biophysica Acta (BBA) - Reviews on Biomembranes* **1990,** *1031* (3), 311-382; (b) Sonohara, R.; Muramatsu, N.; Ohshima, H.; Kondo, T., Difference in surface properties between Escherichia coli and Staphylococcus aureus as revealed by electrophoretic mobility measurements. *Biophysical Chemistry* **1995,** *55* (3), 273-277; (c) Devi, U. V.; Puri, P.; Sharma, N. N.; Ananthasubramanian, M., Electrokinetics of Cells in Dielectrophoretic Separation: A Biological Perspective. *BioNanoScience* **2014,** *4* (3), 276-287.
3.   (a) Xue, X.; Pasparakis, G.; Halliday, N.; Winzer, K.; Howdle, S. M.; Cramphorn, C. J.; Cameron, N. R.; Gardner, P. M.; Davis, B. G.; Fernandez-Trillo, F.; Alexander, C., Synthetic polymers for simultaneous bacterial sequestration and quorum sense interference. *Angew Chem Int*

Ed Engl **2011,** *50* (42), 9852-6; (b) Lui, L. T.; Xue, X.; Sui, C.; Brown, A.; Pritchard, D. I.; Halliday,
N.; Winzer, K.; Howdle, S. M.; Fernandez-Trillo, F.; Krasnogor, N.; Alexander, C., Bacteria
clustering by polymers induces the expression of quorum-sensing-controlled phenotypes. *Nat Chem*
2013, *5* (12), 1058-65; (c) Louzao, I.; Sui, C.; Winzer, K.; Fernandez-Trillo, F.; Alexander, C.,
Cationic polymer mediated bacterial clustering: Cell-adhesive properties of homo- and copolymers.
Eur J Pharm Biopharm **2015,** *95* (Pt A), 47-62; (d) Zhang, P.; Lu, H.; Chen, H.; Zhang, J.; Liu, L.;
Lv, F.; Wang, S., Cationic Conjugated Polymers-Induced Quorum Sensing of Bacteria Cells.
Analytical chemistry **2016,** *88* (6), 2985-8; (e) Leire, E.; Amaral, S. P.; Louzao, I.; Winzer, K.;
Alexander, C.; Fernandez-Megia, E.; Fernandez-Trillo, F., Dendrimer mediated clustering of
bacteria: improved aggregation and evaluation of bacterial response and viability. *Biomater Sci* **2016,**
4 (6), 998-1006; (f) Perez-Soto, N.; Moule, L.; Crisan, D. N.; Insua, I.; Taylor-Smith, L. M.; Voelz,
K.; Fernandez-Trillo, F.; Krachler, A. M., Engineering microbial physiology with synthetic
polymers: cationic polymers induce biofilm formation in Vibrio cholerae and downregulate the
expression of virulence genes. *Chem Sci* **2017,** *8* (8), 5291-5298.
4.   Miller, M. B.; Skorupski, K.; Lenz, D. H.; Taylor, R. K.; Bassler, B. L., Parallel quorum
sensing systems converge to regulate virulence in Vibrio cholerae. *Cell* **2002,** *110* (3), 303-14.
5.   Jung, S. A.; Chapman, C. A.; Ng, W. L., Quadruple quorum-sensing inputs control Vibrio
cholerae virulence and maintain system robustness. *PLoS pathogens* **2015,** *11* (4), e1004837.
6.   Svenningsen, S. L.; Waters, C. M.; Bassler, B. L., A negative feedback loop involving small
RNAs accelerates Vibrio cholerae's transition out of quorum-sensing mode. *Genes Dev* **2008,** *22* (2),
226-38.
7.   Teschler, J. K.; Zamorano-Sanchez, D.; Utada, A. S.; Warner, C. J.; Wong, G. C.; Linington,
R. G.; Yildiz, F. H., Living in the matrix: assembly and control of Vibrio cholerae biofilms. *Nature*
reviews. Microbiology **2015,** *13* (5), 255-68.
8.   Engebrecht, J.; Silverman, M., Identification of genes and gene products necessary for
bacterial bioluminescence. *Proceedings of the National Academy of Sciences of the United States of*
America **1984,** *81* (13), 4154-8.
9.   Silva, A. J.; Benitez, J. A., Transcriptional regulation of Vibrio cholerae
hemagglutinin/protease by the cyclic AMP receptor protein and RpoS. *Journal of bacteriology* **2004,**
186 (19), 6374-82.
10.   Liang, W.; Pascual-Montano, A.; Silva, A. J.; Benitez, J. A., The cyclic AMP receptor
protein modulates quorum sensing, motility and multiple genes that affect intestinal colonization in
Vibrio cholerae. *Microbiology* **2007,** *153* (Pt 9), 2964-75.
11.   Dunlap, P. V.; Greenberg, E. P., Control of Vibrio fischeri lux gene transcription by a cyclic
AMP receptor protein-luxR protein regulatory circuit. *Journal of bacteriology* **1988,** *170* (9), 4040-6.
12.   Hurley, A.; Bassler, B. L., Asymmetric regulation of quorum-sensing receptors drives
autoinducer-specific gene expression programs in Vibrio cholerae. *PLoS Genet* **2017,** *13* (5),
e1006826.
13.   Zhu, J.; Miller, M. B.; Vance, R. E.; Dziejman, M.; Bassler, B. L.; Mekalanos, J. J., Quorum-
sensing regulators control virulence gene expression in Vibrio cholerae. *Proceedings of the National*
Academy of Sciences of the United States of America **2002,** *99* (5), 3129-34.
14.   (a) Yildiz, F. H.; Dolganov, N. A.; Schoolnik, G. K., VpsR, a Member of the Response
Regulators of the Two-Component Regulatory Systems, Is Required for Expression of vps
Biosynthesis Genes and EPS(ETr)-Associated Phenotypes in Vibrio cholerae O1 El Tor. *Journal of*
bacteriology **2001,** *183* (5), 1716-26; (b) Fong, J. C.; Karplus, K.; Schoolnik, G. K.; Yildiz, F. H.,
Identification and characterization of RbmA, a novel protein required for the development of rugose
colony morphology and biofilm structure in Vibrio cholerae. *Journal of bacteriology* **2006,** *188* (3),
1049-59; (c) Fong, J. C.; Yildiz, F. H., The rbmBCDEF gene cluster modulates development of

rugose colony morphology and biofilm formation in Vibrio cholerae. *Journal of bacteriology* **2007,**
189 (6), 2319-30.
15.   Friedrich, W. F.; Greenberg, E. P., Glucose repression of luminescence and luciferase in
Vibrio fischeri. *Arch. Microbiol.* **1983,** *134*, 87-91.
16.   Shikuma, N. J.; Fong, J. C.; Odell, L. S.; Perchuk, B. S.; Laub, M. T.; Yildiz, F. H.,
Overexpression of VpsS, a hybrid sensor kinase, enhances biofilm formation in Vibrio cholerae.
Journal of bacteriology **2009,** *191* (16), 5147-58.
17.   Zamorano-Sanchez, D.; Fong, J. C.; Kilic, S.; Erill, I.; Yildiz, F. H., Identification and
characterization of VpsR and VpsT binding sites in Vibrio cholerae. *Journal of bacteriology* **2015,**
197 (7), 1221-35.
18.   Yildiz, F. H.; Schoolnik, G. K., Role of rpoS in stress survival and virulence of Vibrio
cholerae. *Journal of bacteriology* **1998,** *180* (4), 773-84.
19.   Miller, V. L.; Dirita, V. J.; Mekalanos, J. J., Identification of Toxs, a Regulatory Gene Whose
Product Enhances Toxr-Mediated Activation of the Cholera-Toxin Promoter. *Journal of Bacteriology*
1989, *171* (3), 1288-1293.
20.   Page, L.; Griffiths, L.; Cole, J. A., Different physiological roles of two independent pathways
for nitrite reduction to ammonia by enteric bacteria. *Arch Microbiol* **1990,** *154* (4), 349-54.
21.   Taylor, R. G.; Walker, D. C.; McInnes, R. R., E. coli host strains significantly affect the
quality of small scale plasmid DNA preparations used for sequencing. *Nucleic Acids Res* **1993,** *21*
(7), 1677-8.
22.   Simon, R.; Priefer, U.; Puhler, A., A Broad Host Range Mobilization System for Invivo
Genetic-Engineering - Transposon Mutagenesis in Gram-Negative Bacteria. *Bio-Technol* **1983,** *1* (9),
784-791.
23.   Bell, A. I.; Gaston, K. L.; Cole, J. A.; Busby, S. J., Cloning of binding sequences for the
Escherichia coli transcription activators, FNR and CRP: location of bases involved in discrimination
between FNR and CRP. *Nucleic acids research* **1989,** *17* (10), 3865-74.
24.   Bassler, B. L.; Wright, M.; Showalter, R. E.; Silverman, M. R., Intercellular signalling in
Vibrio harveyi: sequence and function of genes regulating expression of luminescence. *Mol*
Microbiol **1993,** *9* (4), 773-86.
25.   Hammer, B. K.; Bassler, B. L., Regulatory small RNAs circumvent the conventional quorum
sensing pathway in pandemic Vibrio cholerae. *Proceedings of the National Academy of Sciences of*
the United States of America **2007,** *104* (27), 11145-9.
26.   Ng, W. L.; Wei, Y.; Perez, L. J.; Cong, J.; Long, T.; Koch, M.; Semmelhack, M. F.;
Wingreen, N. S.; Bassler, B. L., Probing bacterial transmembrane histidine kinase receptor-ligand
interactions with natural and synthetic molecules. *Proceedings of the National Academy of Sciences*
of the United States of America **2010,** *107* (12), 5575-80.

Tables

Table 1. Bacterial strains used in this study.

Strain	Description or genotype	Source or Reference
Vibrio cholerae		
A1552	Wild-type; O1 biovar El Tor serotype Inaba	[18]
E7956	Wild-type; O1 biovar El Tor serotype Ogawa	[19]
BH1651	*luxO*^{D47E}	[6]
BH1578	Δ*luxS*Δ*cqsA*	[25]
DH231	Δ*luxS*Δ*cqsS*	[26]
WN1103	Δ*luxQ*Δ*cqsA*	[26]
E7956 Δ*crp*	Δ*crp* Kan^R	Gift from D. Grainger
NP5005	A1552 pRW50T containing upstream region of *aphA* promoter; Tet^R	[3f]
Escherichia coli		
DH5α	Donor and maintenance of pBB1	[21]
JCB387	Donor and maintenance of pRW50T	[20]
SM10	Helper strain; λ*pir* pRK2013; Kan^R	[22]

568   **Table 2. Plasmids used in this study.**

Plasmid	Description	Source or Reference
pRW50T	pRW50 derivative with a oriT sequence from pRK2; Tet^R	Gift from D. Grainger
pRW50T-*rbmA*	pRW50T containing 273 bp of the upstream region of *rbmA*, cloned between EcoRI and HindIII restriction sites; Tet^R.	This study
pRW50T- *vpsR*	pRW50T containing 195 bp of the upstream region of *vpsR*, cloned between EcoRI and HindIII restriction sites; Tet^R	This study
pRW50T-*aphA*	pRW50T containing the upstream region of aphA, cloned between EcoRI and HindIII sites; Tet^R	[3f]
pBB1	*luxCDABE* cosmid; Tet^R	[24]

570