Running Title: PORCN-independent Wnt signaling

Research article

WNT4 and WNT3A activate cell autonomous Wnt signaling independent of PORCN or secretion

Deviyani M. Rao[1], Evelyn K. Bordeaux[1], Tomomi M. Yamamoto[2], Benjamin G. Bitler[2], Matthew J.

Sikora[1]

Affiliations

[1]Dept. of Pathology, [2]Dept. of Obstetrics and Gynecology, University of Colorado Anschutz Medical

Campus

Corresponding author

Matthew J. Sikora, PhD.; Mail Stop 8104, Research Complex 1 South, Room 5117, 12801 E. 17[th] Ave.;

Aurora, CO 80045. Tel: (303)724-4301; Fax: (303)724-3712; email: matthew.sikora@ucdenver.edu.

Twitter: @mjsikora

Authors' contributions

DMR and MJS conceived of the project and experiments. DMR and MJS designed and performed

experiments. DMR and TMY developed models for the project. DMR, EKB, BGB, and MJS contributed

to data analysis and interpretation. DMR wrote the draft manuscript; all authors read and revised the

manuscript and have read and approved of this version of the manuscript.

Key words

WNT4, WNT3A, PORCN, Wnt signaling, Wnt secretion

Summary Statement

Wnt proteins can mediate an atypical mode of cell-autonomous signaling, distinct from paracrine

signaling, that is independent of both palmitoylation by PORCN and Wnt secretion.

Abstract

The enzyme PORCN is considered essential for Wnt secretion and signaling, however, we observed

PORCN inhibition did not phenocopy the effects of WNT4 knockdown in WNT4-dependent breast

cancer cells. This suggests a unique relationship between PORCN and WNT4 signaling. To examine the

role of PORCN in WNT4 signaling, WNT4 or WNT3A were over-expressed in breast and ovarian cancer,

and fibrosarcoma cell lines. Conditioned medium from these lines, and co-culture systems, were used to

assess the dependence of Wnt secretion and activity on critical Wnt secretion proteins PORCN and WLS.

We observed that WLS was universally required for Wnt secretion and paracrine signaling. In contrast,

the dependence of WNT3A secretion and activity on PORCN varied across cell lines, and WNT4

secretion was PORCN-independent in all models. Surprisingly, WNT4 did not present paracrine activity

in any tested context. Absent the expected paracrine activity of secreted WNT4, we identified cell

autonomous Wnt signaling activation by WNT4 and WNT3A, independent of PORCN or secretion. The

PORCN-independent, cell-autonomous Wnt signaling demonstrated herein may be critical in WNT4-

driven cellular contexts, or those that are otherwise considered to have dysfunctional Wnt signaling.

Introduction

Wnt signaling is an ancestrally conserved pathway that plays fundamental roles in embryonic development and adult tissue homeostasis. Dysregulation of Wnt signaling is a causative factor for a range of human pathologies, including several forms of cancer (reviewed in (1)). As a result, inhibition of Wnt signaling has become an attractive therapeutic target in ongoing clinical trials, with some strategies targeting the upstream activation of signaling by Wnt proteins (1–3). Wnt proteins comprise a family of secreted glycoproteins that act as intercellular ligands, which stimulate a myriad of signal transduction cascades regulating cellular proliferation, stem cell renewal, cell motility, angiogenesis, and apoptosis (1, 4–6). Wnt proteins are post-translationally modified by the O-acyltransferase Porcupine (PORCN), which palmitoylates Wnt proteins at single serine residues (2, 7–9). This lipidation forms a binding motif for interaction with Wntless (WLS), which chaperones Wnt proteins to the plasma membrane for secretion (8, 10, 11). Once secreted, Wnt proteins signal in a paracrine manner, binding nearby receptor complexes. Wnts typically bind a Frizzled (FZD) receptor in conjunction with the LRP5 or LRP6 co-receptor, resulting in activation of the Disheveled second messenger proteins (DVL1/2/3 in humans) and initiation of either canonical (β-catenin-dependent) or non-canonical (β-catenin-independent) signaling (1, 4). The essential initiating step in Wnt processing is palmitoylation by PORCN, which has prompted the development of PORCN inhibitors, including IWP compounds (11), WNT974 (a.k.a. LGK974) (3), and others (2, 12). PORCN inhibitors have been shown to block Wnt secretion, inhibit downstream Wnt signaling, and suppress Wnt-driven tumor growth in animal models (3, 13, 14), with WNT974 currently in Phase I/II clinical trials for cancer treatment (NCT01351103, NCT02278133). Based on these observations, PORCN inhibitors are an attractive strategy to target Wnt-driven pathologies.

The Wnt protein WNT4 is critical in organogenesis of endocrine organs and regulation of bone mass, and underlies steroid hormone-related phenotypes in humans (15–22). WNT4 dysregulation via loss-of-function mutation results in developmental female to male sex reversal (23–26). Similarly, *WNT4* polymorphisms are associated with endocrine dysfunction, gynecological malignancies, reduced bone density with premature skeletal aging, and related phenotypes (27–33). WNT4 is also critical in mammary gland development, as *Wnt4* knockout in mouse mammary gland prevents progesterone-driven ductal elongation and branching during pregnancy (34, 35). In this context, activated progesterone receptor drives expression of *Wnt4* in mammary gland luminal cells resulting in paracrine signaling that supports maintenance of the mammary stem cell niche (6, 36–38). Despite these observed critical roles of WNT4 in both normal and malignant tissues, WNT4 signaling is crudely understood due to varied context-dependent functions. In a cell type- and tissue-specific manner, WNT4 (human or murine) has been shown to regulate either canonical or non-canonical Wnt signaling, and has been shown to either activate

or suppress signaling (described in references herein). Further, conflicting reports exist as to whether

Wnt4 can or cannot activate canonical Wnt signaling in the murine mammary gland (36, 39). As such,

WNT4 has been described as a "problem child" among Wnt proteins. It is also unclear which FZD

receptor complexes are utilized by WNT4, as WNT4 is often required for distinct, non-redundant

functions versus other Wnt proteins (reviewed in (35)). Since WNT4 has myriad downstream signaling

effects, inhibition of WNT4 upstream of Wnt effector pathways (e.g. with PORCN inhibitors) is an

attractive approach to block WNT4 signaling in a "pathway indifferent" manner to treat WNT4-related

pathologies.

We recently reported that regulation of *WNT4* expression is co-opted by the estrogen receptor in a subtype

of breast cancer, invasive lobular carcinoma (ILC) (40, 41). Estrogen-driven WNT4 is required in ILC

cells for estrogen-induced proliferation and survival, as well as anti-estrogen resistance (41). Though

WNT4-driven signaling in ILC is yet to be fully elucidated, ILC cells lack the capacity to engage

canonical Wnt signaling, as the characteristic genetic loss of E-cadherin in ILC leads to loss of β-catenin

protein (41, 42). This suggests WNT4 drives non-canonical Wnt signaling in ILC cells. Though the

specific non-canonical pathway activated by WNT4 is unknown, PORCN inhibition should be an

effective strategy to block WNT4 upstream and treat this subtype of breast cancer. However, treatment of

ILC cells with PORCN inhibitors did not suppress growth or survival. These unexpected results initiated

further studies into the mechanisms of WNT4 secretion and signaling. In this report, we show WNT4

secretion is mediated by atypical mechanisms. Our observations challenge the paradigm that PORCN-

mediated secretion is required for Wnt signaling, and suggest a novel process by which Wnt proteins,

including WNT4, can initiate non-canonical Wnt signaling.

Results

PORCN inhibition does not mimic WNT4 siRNA in lobular carcinoma cells

We hypothesized that since ILC cells are dependent on WNT4 for proliferation and survival (41),

inhibition of PORCN would phenocopy *WNT4* siRNA by blocking WNT4 secretion and downstream

signaling. Proliferation and cell death were monitored by live cell imaging of MM134 (ILC) cells either

transfected with siRNA targeting *PORCN* (siPORCN) or treated with PORCN inhibitor (PORCNi)

LGK974. Proliferation and cell death were compared to untreated cells, and cells treated with the anti-

estrogen fulvestrant (Fulv) or transfected with siRNA targeting *WNT4* (siWNT4). As we previously

reported (41), siRNA-mediated *WNT4* knockdown or Fulv halt proliferation, and WNT4 knockdown

induces cell death (**Fig. 1A**). However, neither genetic nor chemical PORCN inhibition had any effect on

cell proliferation or survival of MM134 cells (**Fig. 1A,B**). Similar results were obtained in ILC cell line

SUM44PE, as PORCN inhibitor at concentrations up to 1μM did not affect proliferation (**Supplemental**

Fig. 1). These data suggest PORCN inhibition is not sufficient to inhibit WNT4 function, and WNT4

signaling likely occurs via PORCN-independent mechanisms.

WNT4 secretion is WLS-dependent but PORCN-independent

Since targeting PORCN did not phenocopy WNT4 knockdown, we further examined the role of PORCN

in WNT4 secretion. To facilitate Wnt secretion studies we over-expressed WNT3A or WNT4 in MM134

(MM134:W3 and MM134:W4; **Fig. 2A**, **Table 1**), and measured secreted Wnt proteins in conditioned

medium. Of note, since epitope tags may alter Wnt secretion and activity (e.g. (8)), we performed all

studies with non-tagged Wnt constructs. A general workflow for experiments assessing Wnt secretion and

function, for each figure, is shown in **Supplemental Fig. 2**.

PORCN-mediated palmitoylation of Wnt proteins is commonly described as required for Wnt binding to

WLS and transport to the cell surface for secretion (see Introduction), so we examined the requisite of

PORCN (using PORCNi and siPORCN) or WLS (using siWLS) for Wnt secretion. Secreted WNT3A and

WNT4 were detected in conditioned medium from MM134:W3 and MM134:W4 respectively (**Fig. 2B**).

Consistent with the lack of effect of cell proliferation, PORCNi treatment had no effect on WNT4

secretion, and WNT3A secretion was also unaffected by PORCNi (**Fig. 2B, top**). Similarly, siPORCN

had no effect on secretion of either WNT4 or WNT3A (**Fig. 2B, bottom**; efficacy of PORCNi and

siPORCN were confirmed below). However, WLS was required for Wnt secretion, as siWLS suppressed

secretion of both WNT3A and WNT4 from MM134 (**Fig. 2B, bottom**). These data suggest that Wnt

processing and secretion may be atypical in ILC cells, but the PORCN-independent secretion of WNT4 is

a potential mechanism of PORCNi resistance (**Figure 1**).

To determine whether PORCN-independent WNT4 secretion is ILC-specific, we utilized the HT1080

fibrosarcoma cell line, a well-characterized model for Wnt secretion, signaling, and activity (8, 43).

HT1080 are derived from a bone-like tissue and thus are a relevant context for WNT4 signaling (18, 22)

(e.g. WNT4 activates DVL via non-canonical Wnt signaling in HT1080 (8)). We generated WNT3A and

WNT4 over-expressing cells from both wild-type HT1080 and PORCN-knockout HT1080 (HT1080-

PKO, clone delta-19 (43)) (**Fig. 2C, Table 1**), and assessed Wnt secretion as above. Unlike the ILC

model, WNT3A secretion from HT1080 was PORCN-dependent, as WNT3A could be detected in

conditioned medium from HT1080:W3 but not HT1080-PKO:W3 cells (**Fig. 2D**). PORCNi treatment

also blocked WNT3A secretion from HT1080:W3 (**Supplemental Fig. 3A**). In contrast, WNT4 secretion

was detected from both HT1080:W4 and HT1080-PKO:W4 (**Fig. 2D**), and PORCNi did not suppress

WNT4 secretion from HT1080:W4 (**Supplemental Fig. 3A**), supporting that WNT4 secretion is

PORCN-independent.

Endogenously expressed WNT4 was also detected in conditioned medium from HT1080 and HT1080-

PKO cell lines (**Fig. 2D**). We observed that secreted WNT4 can be resolved by electrophoresis as a

doublet. The larger species was PORCN-dependent and not detected in HT1080-PKO, suggesting that

these species represent palmitoylated versus non-palmitoylated proteins. Notably, in HT1080:W3

endogenous secreted WNT4 shifted to the larger species (**Fig. 2D**, column 2), potentially due to positive

feedback activation of PORCN activity (44). We observed loss of both species in conditioned media after

WNT4 knockdown by siRNA, confirming both secreted species as WNT4 (**Supplemental Fig. 3A**).

These data indicate that while WNT4 is modified by PORCN, PORCN is not required for WNT4

secretion. WLS knockdown suppressed secretion of both WNT3A and WNT4 from HT1080 (**Fig. 2E**),

confirming that WNT3A secretion is dependent on both WLS and PORCN , while WNT4 secretion is

WLS-dependent but PORCN-independent .

Though WNT4 secretion was PORCN-independent in both MM134 and HT1080, WNT4 appeared to be

post-translationally modified during secretion (**Fig. 2D**). Also, in both MM134 and HT1080, secreted

WNT4 migrated as a higher molecular weight species than WNT4 from cell lysate (**Supplemental Fig.

3B**). The increased molecular weight is due at least in part to glycosylation (45), as treatment with

tunicamycin (N-linked glycosylation inhibitor) decreased the apparent molecular weight of secreted

WNT4 in either cell line (**Supplemental Fig. 3B**). Notably, MM134 secreted two WNT4 species when

treated with tunicamycin, suggesting WNT4 modification can be variable and cell context-specific.

Together these data demonstrate that WNT4 secretion is PORCN-independent, and that Wnt protein

processing and signaling are more broadly atypical in ILC. PORCN-independent WNT4 secretion may

explain the disparate effects of siWNT4 versus PORCN inhibition. However, it is unclear if Wnt proteins

secreted independently of PORCN are competent to activate paracrine Wnt signaling.

WNT4 over-expression does not activate paracrine Wnt signaling

Since palmitoylation of Wnt proteins is typically necessary for activation of downstream signaling, we

assessed whether WNT4 secreted independent of PORCN could activate paracrine Wnt signaling.

Paracrine signaling was tested by treating HT1080-PKO cells ("receiver" cells, PKO reduces endogenous

paracrine Wnt signaling; see **Supplemental Fig. 2**) with conditioned medium from MM134 cells, with or

without Wnt over-expression. After 24hr treatment with conditioned medium, activation of Wnt signaling

in receiver cells was assessed via phosphorylation of DVL2, DVL3, and LRP6, and expression of β-

catenin target gene *AXIN2*. Conditioned medium from neither MM134:W4 nor MM134:W3 were able to

activate Wnt signaling in receiver cells by any measure (**Fig. 3A,B**) despite the presence of Wnt protein in

conditioned medium from these cells (**Fig. 2B**). This suggests Wnt secretion is not sufficient for paracrine

activity. We also assessed Wnt paracrine activity from wild-type vs PKO Wnt-overexpressing HT1080.

Conditioned medium from HT1080:W3 activated Wnt signaling in the receiver cells by all measures (**Fig.**

3C,D), and this was blocked by PORCN knockout (ie. from HT1080-PKO:W3), consistent with a lack of

WNT3A secretion in HT1080-PKO (**Fig. 2D**). However, conditioned medium from neither HT1080:W4

nor HT1080-PKO:W4 activated Wnt signaling in the receiver cells (**Fig. 3C,D**) despite detectable

secreted WNT4 in both contexts (**Fig. 2D**). Thus, secreted WNT4 from MM134, HT1080, or HT1080-

PKO was unable to activate paracrine Wnt signaling.

Paracrine signaling by Wnt proteins can be mediated by Wnt secretion as well as by Wnt presentation on

the cell surface and subsequent cell-cell contact (46, 47). To determine whether paracrine WNT4

signaling is initiated specifically by cell-cell contact, we assessed whether Wnt signaling could be

activated in "receiver" cells via co-culture with Wnt-overexpressing cells. HT1080-PKO cells transfected

with the TOP-FLASH reporter ("receiver" cells) were co-cultured with cells expressing WNT3A or

WNT4 (**Fig. 3E**). WNT3A-expressing cells activated TOP-FLASH in co-cultured cells in a PORCN-

dependent manner (ie. blocked by LGK974, and absent from HT1080-PKO:W3 cells). However, under

no condition was WNT4 able to activate TOP-FLASH in co-cultured cells. While WNT3A is able to

mediate paracrine Wnt signaling in both secreted and co-culture models, neither secreted nor cell surface

WNT4 mediates paracrine signaling.

Since WNT4 from neither HT1080 nor MM134 cells was able to activate paracrine Wnt signaling, we

treated HT1080-PKO cells with recombinant human Wnt proteins (rWNT3A and rWNT4; 10-500ng/mL).

rWNT3A activated Wnt signaling by all measures, while rWNT4 failed to activate Wnt signaling at any

concentration (**Fig. 3F,G**). These data suggest that HT1080 may be non-responsive to paracrine WNT4,

raising the need for orthogonal systems to validate the function of both secreted and recombinant WNT4.

To address this, we used MC3T3-E1 as an additional "receiver" cell line (**Supplemental Fig. 2**). MC3T3-

E1 cells are another bone-like model highly responsive to exogenous Wnt protein, and induce alkaline

phosphatase production upon Wnt signaling activation (measured by colorimetric assay, see Materials and

Methods (48)). Conditioned medium as above was used to treat "receiver" 3T3-E1 cells, and alkaline

phosphatase (AP) activity was used as the readout for activation of paracrine Wnt signaling. Either

rWNT3A or rWNT4 increased AP activity (**Fig. 4A**), and both rWNT3A and rWNT4 induced DVL and

LRP6 phosphorylation in 3T3-E1 cells (**Supplemental Fig. 4A**). This confirmed that 3T3-E1 respond to

paracrine WNT4, and could be used to assess the activity of secreted Wnts in conditioned medium.

Conditioned medium from HT1080:W3, but not HT1080-PKO:W3, induced AP activity in 3T3-E1 (**Fig.**

4B), consistent with a requirement of PORCN for paracrine WNT3A signaling. However, though rWNT4

induced AP activity, conditioned medium from neither HT1080:W4 nor HT1080-PKO:W4 induced AP

activity in 3T3-E1 (**Fig. 4B**). Parallel results were obtained using MM134, as conditioned medium from

MM134:W3 induced AP activity in 3T3-E1, which was blocked by PORCNi treatment, but no AP

activity was induced with MM134:W4 conditioned medium (**Fig. 4C**). We considered that the

concentration of secreted WNT4 protein may be insufficient to activate 3T3-E1, but secreted WNT3A or

WNT4 is within the range of active rWNT concentrations in this assay (**Supplemental Figure 4B-D**).

Notably, commercially available rWNT3A and rWNT4 are produced lacking the N-terminal signal

peptide, and both rWNT3A and rWNT4 migrated as a smaller peptide than the corresponding Wnt protein

secreted from either HT1080 or MM134 (**Supplemental Fig. 4B-C**). Taken together, these results

indicate secreted Wnt proteins are not equivalent to recombinant proteins, and have distinct capacities for

activating paracrine Wnt signaling. However, while secreted WNT3A activated paracrine Wnt signaling

in a context-dependent manner (based on both source and receiver cells), secreted WNT4 was unable to

act as a functional paracrine signaling ligand in any tested context.

PORCN-independent WNT4 secretion and lack of paracrine activity is observed in varied model systems

We further examined Wnt secretion and function via WNT3A or WNT4 over-expression in a second ILC

cell line (SUM44PE), an additional breast cancer cell line (HCC1428), and an ovarian cancer cell line

(PEO1) (**Table 1**). Like bone, ovarian cancer is a relevant context as WNT4 mediates Müllerian tissue

and ovary development (23, 25). Wnt expression and secretion were assessed as above, and paracrine

activity of secreted Wnt proteins was tested using the 3T3-E1 receiver model.

Secreted WNT3A and WNT4 were detected after over-expression in SUM44PE (ILC, **Fig. 5A**), and

WNT4 secretion in SUM44PE was PORCN-independent but WLS-dependent (**Fig. 5B**). Neither WNT3A

nor WNT4 induced AP activity versus parental SUM44PE cells (**Fig. 5C**). These data with SUM44PE are

consistent with our observations of PORCN-independent Wnt secretion and inactive paracrine activity of

secreted Wnt proteins in ILC. Secreted WNT4 also lacked any paracrine activity from both HCC1428

(**Fig. 5D-E**) and PEO1 (**Fig. 5F-G**). In total, we are unable to detect paracrine WNT4 activity in 5 cell

lines derived from 3 different WNT4-responsive tissues of origin (mammary gland, bone, and ovary),

further supporting that secreted WNT4 does not mediate paracrine signaling in WNT4 expressing cells.

Secreted or paracrine WNT4 are not required for ILC cell proliferation and viability

Since secreted WNT4 did not activate paracrine Wnt signaling in MM134 or SUM44PE cells, we

hypothesized that WNT4 secretion is dispensable for WNT4 function in ILC cells. To confirm this, we

examined whether blocking WNT4 secretion by WLS knockdown would phenocopy WNT4 knockdown

in MM134 cells. MM134 were transfected with siRNAs targeting WNT4, PORCN, or WLS, followed by

cell proliferation and death assays as above. Despite the loss of WNT4 secretion, siWLS had no

detrimental effect on MM134 proliferation or viability and did not phenocopy siWNT4 (**Supplemental**

Fig. 5A-B). Similarly, we attempted to rescue WNT4 knockdown with conditioned medium from

MM134 (with or without Wnt over-expression) or rWNT protein. Exogenous Wnt protein had no effect

on siWNT4-mediated growth inhibition or cell death (**Supplemental Fig. 5C-D**). These data support that

secretion may does not mediate the activation of WNT4-driven signaling.

WNT4 and WNT3A activate cell-autonomous signaling

Our observations suggest WNT4 may activate signaling not via paracrine or autocrine mechanisms, but

rather by a cell-autonomous mechanism. To examine cell-autonomous Wnt-induced signaling we

assessed DVL and LRP6 phosphorylation and *AXIN2* expression directly in HT1080 and HT1080-PKO

Wnt over-expressing cells (**Fig. S2**; **Fig. 6A,B**). Consistent with conditioned medium/receiver cell

experiments, both HT1080:W3 and HT1080:W4 had increased DVL and LRP6 phosphorylation (**Fig.**

6A), but only HT1080:W3 had increased *AXIN2* expression (**Fig. 6B**). In HT1080-PKO, over-expression

of either WNT3A or WNT4 induced cell-autonomous DVL2 and DVL3 phosphorylation (**Fig. 6A**). DVL

activation was independent of paracrine signaling, as Wnt over-expression in HT1080-PKO did not

activate LRP6 or induce *AXIN2* expression. Interestingly, this cell-autonomous DVL activation by Wnt

over-expression occurred despite the inability of HT1080 to respond to paracrine WNT4 (**Fig. 3**), and the

lack of WNT3A secretion in HT1080-PKO (**Fig. 2**). Similarly, Wnt over-expression did not activate

LRP6 in MM134 (**Supplemental Fig. 6**) further supporting that secreted WNT4 does not activate

paracrine signaling. These data support that WNT4 and WNT3A can mediate intracellular signaling

independent of secretion or activation of extracellular signaling (ie. LRP6).

As WNT4 is required for bone regeneration and cell proliferation (22), we examined if WNT4 is similarly

essential for proliferation and/or viability of HT1080 or HT1080-PKO. We hypothesized that WNT4

might be dispensable in HT1080, due to redundant Wnt signaling. However, without functional PORCN

for secretion and paracrine signaling of Wnt family members, HT1080-PKO may become reliant on cell-

autonomous PORCN-independent WNT4 signaling. WNT4 knockdown induced ~21% cell death at 48h

post-knockdown in HT1080 (**Fig. 6C**), leading to a modest suppression of proliferation (**Fig. 6D**). In

contrast, WNT4 knockdown in HT1080-PKO strongly suppressed growth, and cell death was accelerated

(~70% cell death at 48h post-knockdown). Importantly, minimal cell death was induced by WLS

knockdown in either HT1080 or HT1080-PKO, despite global suppression of Wnt secretion in the latter

with ablation of both PORCN and WLS. The sensitivity of HT1080-PKO to knockdown of WNT4, but

not WLS, supports a critical role for secretion-independent functions of WNT4, in particular in the

absence of paracrine signaling by other Wnt proteins.

To confirm that cell-autonomous PORCN-independent WNT signaling was similarly independent of

WLS, we knocked-down WLS or PORCN in HT1080-PKO Wnt over-expressing cells and assessed

DVL3 activation (**Fig. 6E**). siPORCN did not affect DVL3 activation, consistent with our above

observations. DVL3 activation in HT1080-PKO:W3 was suppressed, suggesting disparate roles of

PORCN and WLS in activating paracrine versus cell-autonomous signaling. However, siWLS did not

impact cell-autonomous DVL3 activation by WNT4, further supporting a secretion-independent cell-

autonomous signaling mechanism for WNT4.

Discussion

The Wnt modifying enzyme PORCN is commonly described as a gatekeeper for the secretion of Wnt

proteins, and thus PORCN inhibition is an approach to broadly block Wnt signaling without targeting cell

type- or tissue-specific downstream Wnt pathways. WNT4 signaling is required for survival and

proliferation of ILC cells, but we show that PORCN is dispensable, calling into question the role of

PORCN in WNT4 signaling. PORCN was not required for WNT4 secretion from a panel of cell lines, as

genetic or chemical PORCN blockade had no effect on WNT4 section. However, WNT4 was not capable

of activating paracrine Wnt signaling in any model tested, despite the ability of recombinant human

WNT4 to do so in a context-dependent manner. These data together suggest that secreted WNT4 may not

be responsible for driving signaling in WNT4-expressing cells. Instead, we determined that WNT4 and

WNT3A can activate cell-autonomous, intracellular signaling independent of secretion (**Fig. 7,**

Supplemental Fig. 7). This unique mode of Wnt signaling is likely essential for the survival and

proliferation of WNT4-dependent cells.

Our observations of PORCN-independent WNT4 signaling are supported by other studies showing a

disconnect between PORCN activity and Wnt secretion/signaling, and support that the requirement for

PORCN is context-dependent and not absolute. Nusse and colleagues reported PORCN-independent

secretion and activity of *Drosophila* WntD (49). WntD is secreted at high levels in fly tissues and cell

culture models independent of both Porcupine and Wntless, and ablation of either Porcupine or Wntless

did not affect WntD signaling. WntD utilizes the early secretory pathway protein Rab1 GTPase (RAB1A

homolog) for secretion, a distinct secretion mechanism versus Porcupine-mediated secretion of Wingless

(WNT1 homolog). WntD lacks the conserved serine residue that is palmitoylated by PORCN (49), unique

among Wnt proteins, but this supports that Wnt proteins can be secreted and signal independent of

PORCN-driven modification. PORCN-independent Wnt secretion and signaling was also observed in

human primary cells by Richards et al (50). Neither PORCNi (IWP-2) nor *PORCN* siRNA knockdown

suppressed secretion of any endogenously expressed Wnt proteins from CD8+ T-cells (Wnts 1, 3, 5B,

10B) or astrocytes (Wnts 1, 3, 6, 7A, 10A, 16). Wnt proteins secreted from PORCNi-treated cells were

functional in conditioned medium experiments, but PORCN-independent secretory mechanisms were not

characterized. Other studies have shown that PORCN may differentially regulate the activity of individual

Wnt proteins. In HEK293T cells, over-expression of PORCN and WLS together enhanced secretion of

WNT1 but suppressed WNT1-induced paracrine or autocrine activation of β-catenin (versus WLS over-

expression alone) (51). This was not observed with WNT3A, indicating PORCN specifically suppressed

paracrine activity of WNT1, despite driving WNT1 secretion. Wnt-specific Porcupine functions have also

been reported in Zebrafish (52). Knockdown of *porcn* in Zebrafish embryos suppressed secretion of

Wnt5a and resulted in defects in non-canonical (β-catenin-independent) Wnt signaling. Conversely,

canonical β-catenin-dependent Wnt signaling was not altered by *porcn* knockdown, and secretion of

Wnt3a was not impaired. Similar data regarding specifically WNT4 are limited. A clinical *WNT4*

mutation (L12P (24), discussed below) blocks WNT4 palmitoylation but not secretion, yet ultimately is

associated with WNT4 loss-of-function, consistent with a disconnect between PORCN, WNT4 secretion,

and WNT4 signaling. These studies together show Wnt proteins secreted in PORCN-independent

manners can be active in paracrine signaling models, but to our knowledge, this is the first report of

PORCN-independent Wnt activity that is also independent of secretion. Of note, Kurita et al recently

demonstrated that *Wnt4* siRNA in mouse pancreatic β-cells suppressed glucose-induced insulin secretion,

but recombinant Wnt4 had no effect on insulin secretion (53). These data parallel our findings and

support that WNT4 signals independent of PORCN, secretion, and paracrine activity.

Our data, together with the above reports on PORCN-independent Wnt signaling, highlight that the roles

of PORCN and WLS in Wnt modification, secretion, and signaling are Wnt-specific, as well as cell type-

and context-dependent. This observed context-dependence includes not only the specific "receiver" cells

in question (perhaps best understood in the context of differentially expressed FZD receptors), but also

includes the source of the Wnt protein (**Fig. 7, Supplemental Fig .7**). For example, WNT3A secreted

from HT1080 robustly activated Wnt signaling in 3T3-E1 or HT1080-PKO cells, and PORCN was

required for both secretion and paracrine activity. In contrast, WNT3A secreted from MM134 cells

activated Wnt signaling in 3T3-E1 but not HT1080-PKO cells, and PORCN was required for paracrine

activity but not secretion. Cell-type specific Wnt protein post-translational modification (e.g.

glycosylation patterns) may drive these differences (51, 54). This parallels our observation that

recombinant Wnt proteins were more promiscuously able to activate signaling than secreted Wnt proteins

(eg. rWNT4, but not secreted WNT4, activated 3T3-E1 Wnt signaling). rWnt proteins may thus represent

a specific functional context versus secreted Wnt protein.

WNT4 and WNT3A likely signal via at least three distinct mechanisms: 1) as a secreted protein with

PORCN modification; 2) as a secreted protein without PORCN modification; 3) by a cell-autonomous

mechanism independent of secretion. This may explain the myriad context-dependent signaling controlled

by WNT4, including activating canonical β-catenin activity (15, 36, 55), repressing β-catenin-driven

transcription (56, 57), or activating non-canonical Wnt signaling pathways (22, 58). The signaling

differences we observed between recombinant and secreted WNT4 indicate differential protein processing

may guide Wnt proteins to activate distinct signaling pathways. The commercial recombinant Wnt

proteins used in our study lack the N-terminal signal peptide (residues 1-22 for WNT4), however, Wnt

signal peptides may have important roles in the regulation of signaling activity (45). The WNT4 signal

peptide has uniquely high content of arginine (14%) and serine (18%) compared to other human Wnts

(average 5% and 9%, respectively), whereas WNT3A has no charged or polar residues in its signal

peptide. Mutation in the signal peptide (L12P) of WNT4 has also been linked to Mayer-Rokitansky-

Küster-Hauser syndrome (24). The L12P mutant functions as a dominant negative inhibitor and

suppresses the activity of wild-type WNT4 when co-expressed. Although the L12P mutant protein is not

palmitoylated, it is secreted and does not prevent secretion of wild-type protein (24). These observations

are consistent with our findings of PORCN-independent WNT4 secretion. The mechanism of dominant-

negative activity has not been described but suggests distinct forms of WNT4 may drive cell autonomous

Wnt signaling. Characterizing distinct WNT4 species is an important future direction.

ILC is a unique context for paracrine Wnt signaling, as we observed that both WNT3A and WNT4 were

secreted from ILC cells in a PORCN-independent manner. While WNT3A activity remained dependent

on PORCN (similar to Wnt5a in 293T cells (59)), the role of WNT4 processing in paracrine activity is

unclear. However, this atypical Wnt processing in ILC cells (also observed in relation to glycosylation)

may be related to broader Wnt signaling dysfunction. The genetic hallmark of ILC is the loss of E-

cadherin (*CDH1*) (42), which leads to dysfunction of catenin proteins, including activation of p120

catenin (60) and inactivation of β-catenin. E-cadherin loss in ILC leads to a loss of β-catenin protein in

both patient tumors and cell lines (41, 42), and as a result, a β-catenin-driven TOP-Flash reporter cannot

be activated in ILC cells (41). Catenin dysfunction was previously postulated as being linked to PORCNi

sensitivity, and ILC patients were specifically included in a trial of WNT974 (NCT01351103). This trial

opened in 2011, but by 2015 ILC patients were removed from the inclusion criteria. It is unclear whether

this is due to accrual problems or a lack of efficacy, as neither have been reported for ILC patients on this

trial, although our data suggest PORCNi are unlikely to have clinical efficacy for ILC. This highlights the

importance of defining the unique context for Wnt signaling in ILC, in particular for WNT4, based on our

prior findings (41). Our laboratory has begun to characterize WNT4-driven signaling in ILC cells, which

are likely mediated by PORCN-independent, cell-autonomous WNT4 signaling.

Our study uses diverse cell line models to investigate Wnt secretion and paracrine activity of Wnt

proteins. We used non-tagged Wnt expression constructs, eliminating previously described complications

caused by epitope tags on Wnt proteins. Wnt signaling activity was measured by complementary pathway

readouts that facilitated our study of context-dependent Wnt signaling activities. The cell line models

used for Wnt secretion were all cancer-derived, and thus further study is needed to translate our findings

to normal tissue, developmental, or *in vivo* contexts. We observed identical processing and secretion for

endogenous WNT4 as for over-expressed WNT4 (models herein express low or no endogenous WNT3A),

but over-expression was required to facilitate signaling experiments. Future studies will determine the

contribution of endogenous Wnt protein levels to activating the signaling pathways discussed herein.

The secretion and paracrine activity of Wnt proteins are heavily context-dependent, as WNT3A and

WNT4 present with differing dependence on PORCN for secretion and paracrine activity in distinct

model systems. Secretion of WNT4 is PORCN-independent, but WNT4 did not present paracrine activity

in cells that were dependent on WNT4, indicative of secretion-independent, cell-autonomous activity.

Both WNT4 and WNT3A presented cell-autonomous activity independent of secretion. Our studies

identify a PORCN-independent mode of Wnt signaling may be critical to understanding WNT4-driven

cellular contexts or those which are otherwise considered to have dysfunctional Wnt signaling.

Methods and Materials

Cell culture

HT1080 and PORCN-knockout HT1080 (HT1080-PKO; clone delta-19) were a generous gift from Dr.

David Virshup (43), and were maintained in DMEM/F12 (Corning, Corning, NY, USA, cat#10092CV) +

10% FBS (Nucleus Biologics, San Diego, CA, USA, cat#FBS1824). MDA MB 134VI (MM134; ATCC,

Manassas, VA, USA) and SUM44PE (BioIVT, Westbury, NY, USA) were maintained as described (40).

PEO1 and HCC1428 (ATCC) were maintained in DMEM/F12 + 10% FBS. MC3T3-E1 (ATCC) were

maintained in MEM Alpha without ascorbic acid (Thermo Fisher Scientific, Waltham, MA, USA,

cat#A10490-01) + 10% FBS. Wnt overexpression lines were generated by lentiviral transduction of Wnt

expression plasmids (see below) with selection of antibiotic-resistant pools, and were maintained in 2.5

μg/mL blasticidin. PEO1:W3 were established by us previously (61). All lines were incubated at 37°C in

5% CO_2. Cell lines are authenticated annually by the University of Arizona Genetics Core cell line

authentication service and confirmed to be mycoplasma negative every four months. Authenticated cells

were in continuous culture <6 months.

Reagents and plasmids

LGK974 was obtained from Cayman Chemical (Ann Arbor, MI, USA, cat#14072) and was dissolved in

DMSO. WntC59 was obtained from Tocris Biosciences (Bristol, UK, cat#5148) and was dissolved in

DMSO. Fulvestrant was obtained from Tocris Biosciences (#1047) and was dissolved in EtOH.

Tunicamycin was obtained from Cayman Chemical (# 11445) and reconstituted in DMSO. Recombinant

human WNT3A (#5036-WN-010) and WNT4 (#6076-WN-005) were obtained from R&D Systems

(Minneapolis, MN, USA) and reconstituted per the manufacturer's instructions.

Wnt plasmids used in this publication were a gift from Drs. Marian Waterman, David Virshup and Xi He

(Addgene, Watertown, MA, USA, kit #1000000022) (8). The M50 Super 8x TOPFlash plasmid was a

generous gift from Dr. Randall Moon (Addgene, plasmid #12456) ((62). Lentiviral vectors for WNT3A

and WNT4 were generated by Gateway Recombination of pENTR-STOP Wnt open reading frames to

pLX304, a kind gift from Dr. Bob Sclafani.

Transient transfection assays

HT1080 cells were transfected with Active WNT3A-V5 (G-8) and Active WNT4-V5 (G-10) from

Addgene kit #1000000022. The transfection used Lipofectamine LTX Reagent with PLUS Reagent

(Thermo Fisher Scientific, Cat# 15338100) using the manufacturer's instructions.

Protein extraction from conditioned medium

HT1080, MM134, HCC1428, or PEO1cells were plated in full medium (as above, 10% FBS). Decreased

FBS is required to reduce competition of serum proteins with secreted Wnt proteins during extraction.

24hrs post-plating, medium was changed to reduced serum medium: DMEM/F12 + 2% FBS (HT1080,

PEO1); DMEM + 2% FBS (HCC1428); DMEM/L15 + 5% FBS (MM134). Conditioned medium was

harvested 4-5 days later for HT1080 cells and 6-7 days later for MM134, HCC1428 and PEO1 cells,

typically once medium acidification was apparent by phenol red color change. SUM44PE cells are

normally cultured in low serum (DMEM/F12 + 2% CSS), so standard medium was allowed to condition

for 6-7 days before harvesting. Medium was centrifuged at 300xg for 4min to pellet any cells or debris.

The supernatant was then syringe-filtered using a 0.2μm filter. The protein concentration of the

conditioned medium was measured using a Pierce BCA Protein Assay Kit (Thermo Fisher Scientific,

cat#23225), and medium volumes, normalized to total protein, were adjusted with sterile-filtered PBS.

Strataclean resin (Agilent Technologies, Santa Clara, CA, USA, cat#400714) was added to the

conditioned media at a ratio of 10μL of resin to 100μg of medium protein, and vortexed to re-suspend the

resin. The medium+resin mixture was incubated, rotating at 4C for 30min, then centrifuged at 425xg for

1min at 4C. The supernatant was then aspirated and the resin was washed using sterile-filtered PBS. The

resin was centrifuged again at 425xg for 1min at 4C, and the supernatant was aspirated. An equal volume

of 2X Laemmli sample buffer (Bio-Rad Laboratories, Hercules, CA, USA, cat#1610737) was added to

the resin to release bound protein, and the slurry was heated at 95C for 5min. Slurry equivalent to 100ug

of conditioned medium protein was run on SDS-PAGE gels to detect secreted WNT3A or WNT4 via

immunoblotting (below).

Alkaline phosphatase assay

MC3T3-E1 cells were seeded into a 96-well plate at 10,000 cells/well. 24hrs later the cells were treated

with either conditioned media (CM) or recombinant WNTs (rWNTs). 72hrs later, alkaline phosphatase

activity was assessed, using *para*-Nitrophenyl phosphate (pNp) as a substrate and measuring absorbance

at 405nm. Assay protocol is based on Nakamura et al (63) with buffer modifications provided by the

recombinant Wnt manufacturer (R&D Systems). A more detailed protocol is available upon request.

para-Nitrophenyl Phosphate tablets were obtained from Cayman Chemical (cat# 400090).

Co-culture and Dual Luciferase Assay

HT1080-PKO cells were transfected with both a TOP-FLASH and renilla reporter plasmid. 24 hours later,

these cells were co-cultured with either HT1080 or HT1080-PKO WNT overexpressing cells, with or

without 10nM LGK974 treatment. 48 hours later, luciferase and renilla activity was assessed using

Promega's Dual Luciferase Assay (Promega, Madison, WI, USA, cat# E1910) and a BioTek Synergy 2

microplate reader (BioTek, Winooski, VT, USA) with a dual injector system according to the

manufacturer's instructions.

Proliferation and viability assays

Cells were seeded in a 96-well plate and 24hrs later cells were treated with 100nM Sytox green (Thermo

Fisher Scientific, cat# S7020). The plate was then placed into an Incucyte Zoom (Essen Bioscience, Ann

Arbor, MI, USA) for 4-5 days where they were imaged every 4hrs at 10x magnification. Cell confluence

and cell death (Sytox green-positive counts) were assessed using Incucyte S3 software (v2018A). After

time-course completion, total double-stranded DNA was measured by hypotonic lysis of cells in ultra-

pure H_2O, followed by addition of Hoechst 33258 (Thermo Fisher Scientific, #62249) at 1μg/mL in Tris-

NaCl buffer (10mM Tris, 2M NaCl; pH 7.4) at equivalent volume to lysate. Fluorescence (360nm ex /

460nm em) was measured on a Bio-Tek Synergy 2 microplate reader.

RNA interference

siRNAs were reverse transfected using RNAiMAX (ThermoFisher) according to the manufacturer's

instructions. All constructs are siGENOME SMARTpool siRNAs (GE Healthcare Dharmacon, Lafayette,

CO, USA): Non-targeting pool #2 (D-001206-14-05), Human *WNT4* (M-008659-03-0005), Human

PORCN (M-009613-00) and Human *WLS* (M-018728-01-0005). Details regarding validation of the

specific effects of the *WNT4* siRNA pool are previously described (41).

Gene expression analyses

RNA extractions were performed using the RNeasy Mini kit (Qiagen, Venlo, Netherlands); mRNA was

converted to cDNA on an Eppendorf Mastercycler Pro (Eppendorf, Hamburg, Germany) and using

Promega reagents: Oligo $(dT)_{15}$ primer (cat# C110A), Random Primers (cat# C118A), GoScript 5x

Reaction Buffer (cat# A500D), 25mM MgCl2 (cat# A351H), 10mM dNTPs (cat# U1511), RNasin Plus

RNase Inhibitor (cat# N261B) and GoScript Reverse Transcriptase (cat# A501D). qPCR reactions were

performed with PowerUp SYBR Green Master Mix (Life Technologies, cat # 100029284) on a

QuantStudio 6 Flex Real-Time PCR system. Expression data were normalized to *RPLP0*. The following

primers were used: *RPLP0*, Forward – CAGCATCTACAACCCTGAAG, Reverse –

GACAGACACTGGCAACATT; *WNT4*, Forward – GCCATTGAGGAGTGCCAGTA, Reverse –

CCACACCTGCCGAAGAGATG; *WNT3A*, Forward – ATGGTGTCTCGGGAGTT, Reverse –

TGGCACTTGCACTTGAG; *PORCN*, Forward – ACCATCCTCATCTACCTACTC, Reverse –

CCTTCATGGCCACAATCA; *AXIN2:* Forward – CTCTGGAGCTGTTTCTTACTG, Reverse –

CTCTGGAGCTGTTTCTTACTG.

Immunoblotting

Whole-cell lysates were obtained by incubating cells in lysis buffer (1% Triton X-100, 50mM HEPES pH

7.4, 140mM NaCl, 1.5mM $MgCl_2$, 1mM EGTA, 10% glycerol; supplemented with Roche

protease/phosphatase inhibitor cocktail (Sigma-Aldrich, St. Louis, MO, USA)) for 30min on ice. Cells

were centrifuged at 16000xg for 15min at 4C and the resulting supernatant was collected for analysis.

Protein concentration was measured using the Pierce BCA Protein Assay Kit (#23225). Standard methods

were used to perform SDS-PAGE. Proteins were transferred onto PVDF membranes. Antibodies were

used according to manufacturer's recommendations: WNT4 (R&D, MAB4751, clone# 55025, 1:1,000),

WNT3A (R&D, MAB13242, clone# 217804.2R, 1:1,000), DVL2 (Cell Signaling Technology, Danvers,

MA, USA, cat#3216, 1:2,000), DVL3 (Cell Signaling, 3218, 1:2,000), pLRP6 (s1490, Cell Signaling,

2568, 1:2,000), LRP6 (Cell Signaling, 2560, clone# C5C7, 1:2,000), Vinculin (Cell Signaling, cat#

13901S 1:10,000) and V5 (Novus Biologicals, Centennial, CO, USA, NB100-62264, clone #SV5-PK1,

1:2,500). Specificity of WNT4 antibody was validated using siWNT4 (Figure S3) and the recognition of

recombinant WNT4 compared to endogenous and overexpressed WNT4 (Figure 4E-F). Secondary

antibodies were used according to manufacturer's instruction and were obtained from Jackson

ImmunoResearch Laboratories (West Grove, PA, USA): Goat Anti-Mouse IgG (cat # 115-035-068), Goat

Anti-Rabbit IgG (cat# 111-035-045) and Goat Anti-Rat IgG (cat# 112-035-062). All secondary antibodies

were used at a dilution of 1:10,000. Chemiluminescence was used to detect antibodies and either film or

the LI-COR c-Digit (LI-COR Biosciences, Lincoln, NE, USA) was used to develop the immunoblots. Of

note, WNT4 MAB4751 detects a prominent non-specific band at ~50kD in immunoblots from cell

lysates. This non-specific target largely precludes detection of WNT4, but cutting immunoblot

membranes immediately below a 50kD ladder marker prevents this issue. This non-specific band was not

detected in WNT4 immunoblots from conditioned medium. Similarly, in immunoblots of conditioned

medium WNT3A MAB13242 detects a prominent non-specific band at ~60kD that precludes detection of

secreted WNT3A; cutting membranes above a 50kD ladder marker prevents this issue. This non-specific

band was not detected in WNT3A immunoblots from cell lysates.

546
547   *Statistical Considerations*

548   Prism was used for all graphical representation and statistical analyses. All western blot figures are

549   representative of at least two to three independent experiments.

550
551   *Software*

552   Prism (GraphPad Software, San Diego, CA, USA, version 7.02) and Image Studio (LI-COR Biosciences,

553   Lincoln, NE, USA, version 5.2) were used to obtain and analyze that data presented in this manuscript.

555   **Acknowledgements**

556   The authors thank Dr. Caroline Alexander (U. Wisconsin) for critical reading of the manuscript.

Competing interests

The authors have nothing to disclose.

Funding

This work was supported by R00 CA193734 (MJS) and R00 CA194318 (BGB) from the National

Institutes of Health, by a grant from the Cancer League of Colorado, Inc (MJS), and by T32 GM007635

(EKB).

Data availability

Data associated with experiments herein will be available at an Open Science Framework repository (64)

(https://doi.org/10.17605/OSF.IO/7X8NG).

569   **References**

570   1.   Nusse R, Clevers H. 2017. Wnt/β-Catenin Signaling, Disease, and Emerging Therapeutic
571   Modalities. Cell 169:985–999.

572   2.   Madan B, Ke Z, Harmston N, Ho SY, Frois AO, Alam J, Jeyaraj DA, Pendharkar V, Ghosh K,
573   Virshup IH, Manoharan V, Ong EHQ, Sangthongpitag K, Hill J, Petretto E, Keller TH, Lee MA,
574   Matter A, Virshup DM. 2016. Wnt addiction of genetically defined cancers reversed by PORCN
575   inhibition. Oncogene 35:2197–207.

576   3.   Liu J, Pan S, Hsieh MH, Ng N, Sun F, Wang T, Kasibhatla S, Schuller AG, Li AG, Cheng D, Li J,
577   Tompkins C, Pferdekamper A, Steffy A, Cheng J, Kowal C, Phung V, Guo G, Wang Y, Graham
578   MP, Flynn S, Brenner JC, Li C, Villarroel MC, Schultz PG, Wu X, McNamara P, Sellers WR,
579   Petruzzelli L, Boral AL, Seidel HM, McLaughlin ME, Che J, Carey TE, Vanasse G, Harris JL.
580   2013. Targeting Wnt-driven cancer through the inhibition of Porcupine by LGK974. Proc Natl
581   Acad Sci U S A 110:20224–9.

582   4.   van Amerongen R. 2012. Alternative Wnt pathways and receptors. Cold Spring Harb Perspect
583   Biol 4:a007914–a007914.

584   5.   Loh KM, van Amerongen R, Nusse R. 2016. Generating Cellular Diversity and Spatial Form: Wnt
585   Signaling and the Evolution of Multicellular Animals. Dev Cell 38:643–55.

586   6.   Brisken C, Hess K, Jeitziner R. 2015. Progesterone and Overlooked Endocrine Pathways in Breast
587   Cancer Pathogenesis. Endocrinology 156:3442–50.

588   7.   Miranda M, Galli LM, Enriquez M, Szabo LA, Gao X, Hannoush RN, Burrus LW. 2014.
589   Identification of the WNT1 residues required for palmitoylation by Porcupine. FEBS Lett
590   588:4815–24.

591   8.   Najdi R, Proffitt K, Sprowl S, Kaur S, Yu J, Covey TM, Virshup DM, Waterman ML. 2012. A
592   uniform human Wnt expression library reveals a shared secretory pathway and unique signaling
593   activities. Differentiation 84:203–13.

594   9.   Takada R, Satomi Y, Kurata T, Ueno N, Norioka S, Kondoh H, Takao T. 2006. Monounsaturated
595   Fatty Acid Modification of Wnt Protein : Its Role in Wnt Secretion 791–801.

596   10.   Bänziger C, Soldini D, Schütt C, Zipperlen P, Hausmann G, Basler K. 2006. Wntless, a conserved
597   membrane protein dedicated to the secretion of Wnt proteins from signaling cells. Cell 125:509–
598   22.

599   11.   Coombs GS, Yu J, Canning CA, Veltri CA, Covey TM, Cheong JK, Utomo V, Banerjee N, Zhang
600   ZH, Jadulco RC, Concepcion GP, Bugni TS, Harper MK, Mihalek I, Jones CM, Ireland CM,
601   Virshup DM. 2010. WLS-dependent secretion of WNT3A requires Ser209 acylation and vacuolar
602   acidification. J Cell Sci 123:3357–67.

603   12.   Proffitt KD, Madan B, Ke Z, Pendharkar V, Ding L, Lee MA, Hannoush RN, Virshup DM. 2013.
604   Pharmacological inhibition of the Wnt acyltransferase PORCN prevents growth of WNT-driven
605   mammary cancer. Cancer Res 73:502–7.

606   13.   Hayashi M, Baker A, Goldstein SD, Albert CM, Jackson KW, McCarty G, Kahlert UD, Loeb DM.
607   2017. Inhibition of porcupine prolongs metastasis free survival in a mouse xenograft model of
608   Ewing sarcoma. Oncotarget 8:78265–78276.

609   14.   Agarwal P, Zhang B, Ho Y, Cook A, Li L, Mikhail FM, Wang Y, McLaughlin ME, Bhatia R.
610   2017. Enhanced targeting of CML stem and progenitor cells by inhibition of porcupine
611   acyltransferase in combination with TKI. Blood 129:1008–1020.

612   15.   Li Q, Kannan A, Das A, Demayo FJ, Hornsby PJ, Young SL, Taylor RN, Bagchi MK, Bagchi IC.
613   2013. WNT4 acts downstream of BMP2 and functions via β-catenin signaling pathway to regulate
614   human endometrial stromal cell differentiation. Endocrinology 154:446–57.

615   16.   Itäranta P, Chi L, Seppänen T, Niku M, Tuukkanen J, Peltoketo H, Vainio S. 2006. Wnt-4
616   signaling is involved in the control of smooth muscle cell fate via Bmp-4 in the medullary stroma
617   of the developing kidney. Dev Biol 293:473–83.

618   17.   Tomizuka K, Horikoshi K, Kitada R, Sugawara Y, Iba Y, Kojima A, Yoshitome A, Yamawaki K,
619   Amagai M, Inoue A, Oshima T, Kakitani M. 2008. R-spondin1 plays an essential role in ovarian
620   development through positively regulating Wnt-4 signaling. Hum Mol Genet 17:1278–91.

621   18.   Chang J, Sonoyama W, Wang Z, Jin Q, Zhang C, Krebsbach PH, Giannobile W, Shi S, Wang C-
622   Y. 2007. Noncanonical Wnt-4 signaling enhances bone regeneration of mesenchymal stem cells in
623   craniofacial defects through activation of p38 MAPK. J Biol Chem 282:30938–48.

624   19.   Prunskaite-Hyyryläinen R, Skovorodkin I, Xu Q, Miinalainen I, Shan J, Vainio SJ. 2016. Wnt4
625   coordinates directional cell migration and extension of the Müllerian duct essential for ontogenesis
626   of the female reproductive tract. Hum Mol Genet 25:1059–73.

627   20.   Strochlic L, Falk J, Goillot E, Sigoillot S, Bourgeois F, Delers P, Rouvière J, Swain A, Castellani
628   V, Schaeffer L, Legay C. 2012. Wnt4 participates in the formation of vertebrate neuromuscular
629   junction. PLoS One 7:e29976.

630   21.   Caprioli A, Villasenor A, Wylie LA, Braitsch C, Marty-Santos L, Barry D, Karner CM, Fu S,
631   Meadows SM, Carroll TJ, Cleaver O. 2015. Wnt4 is essential to normal mammalian lung
632   development. Dev Biol 406:222–34.

633   22.   Yu B, Chang J, Liu Y, Li J, Kevork K, Al-Hezaimi K, Graves DT, Park N-H, Wang C-Y. 2014.
634   Wnt4 signaling prevents skeletal aging and inflammation by inhibiting nuclear factor-κB. Nat Med
635   20:1009–17.

636   23.   Bernard P, Harley VR. 2007. Wnt4 action in gonadal development and sex determination. Int J
637   Biochem Cell Biol 39:31–43.

638   24.   Philibert P, Biason-Lauber A, Rouzier R, Pienkowski C, Paris F, Konrad D, Schoenle E, Sultan C.
639   2008. Identification and functional analysis of a new WNT4 gene mutation among 28 adolescent
640   girls with primary amenorrhea and müllerian duct abnormalities: a French collaborative study. J
641   Clin Endocrinol Metab 93:895–900.

642   25.   Vainio S, Heikkilä M, Kispert A, Chin N, McMahon AP. 1999. Female development in mammals
643   is regulated by Wnt-4 signalling. Nature 397:405–9.

644   26.   Biason-Lauber A, De Filippo G, Konrad D, Scarano G, Nazzaro A, Schoenle EJ. 2007. WNT4
645   deficiency--a clinical phenotype distinct from the classic Mayer-Rokitansky-Kuster-Hauser
646   syndrome: a case report. Hum Reprod 22:224–9.

647   27.   Powell JE, Fung JN, Shakhbazov K, Sapkota Y, Cloonan N, Hemani G, Hillman KM, Kaufmann
648   S, Luong HT, Bowdler L, Painter JN, Holdsworth-Carson SJ, Visscher PM, Dinger ME, Healey
649   M, Nyholt DR, French JD, Edwards SL, Rogers PAW, Montgomery GW. 2016. Endometriosis
650   risk alleles at 1p36.12 act through inverse regulation of CDC42 and LINC00339. Hum Mol Genet
651   25:5046–5058.

652   28.   Rahmioglu N, Macgregor S, Drong AW, Hedman ÅK, Harris HR, Randall JC, Prokopenko I,
653   Nyholt DR, Morris AP, Montgomery GW, Missmer SA, Lindgren CM, Zondervan KT, Collins
654   FS, International Endogene Consortium (IEC) TGC, Nyholt DR, Morris AP, Montgomery GW,
655   Missmer SA, Lindgren CM, Zondervan KT. 2015. Genome-wide enrichment analysis between
656   endometriosis and obesity-related traits reveals novel susceptibility loci. Hum Mol Genet
657   24:1185–1199.

658   29.   Mafra F, Catto M, Bianco B, Barbosa CP, Christofolini D. 2015. Association of WNT4
659   polymorphisms with endometriosis in infertile patients. J Assist Reprod Genet 32:1359–64.

30.   Zhang G, Feenstra B, Bacelis J, Liu X, Muglia LJLMLJ, Juodakis J, Miller DE, Litterman N,

Jiang P-P, Russell L, Hinds DA, Hu Y, Weirauch MT, Chen X, Chavan AR, Wagner GP, Pavličev
M, Nnamani MC, Maziarz J, Karjalainen MK, Rämet M, Sengpiel V, Geller F, Boyd HA, Palotie
A, Momany A, Bedell B, Ryckman KK, Huusko JM, Forney CR, Kottyan LC, Hallman M,
Teramo K, Nohr EA, Davey Smith G, Melbye M, Jacobsson B, Muglia LJLMLJ. 2017. Genetic
Associations with Gestational Duration and Spontaneous Preterm Birth. N Engl J Med 377:1156–
1167.

31.   Zmuda JM, Yerges-Armstrong LM, Moffett SP, Klei L, Kammerer CM, Roeder K, Cauley JA,
Kuipers A, Ensrud KE, Nestlerode CS, Hoffman AR, Lewis CE, Lang TF, Barrett-Connor E,
Ferrell RE, Orwoll ES, Osteoporotic Fractures in Men (MrOS) Study Group. 2011. Genetic
analysis of vertebral trabecular bone density and cross-sectional area in older men. Osteoporos Int
22:1079–90.

32.   Zheng Y, Wang C, Zhang H, Shao C, Gao L-H, Li S-S, Yu W-J, He J-W, Fu W-Z, Hu Y-Q, Li M,
Liu Y-J, Zhang Z-L. 2016. Polymorphisms in Wnt signaling pathway genes are associated with
peak bone mineral density, lean mass, and fat mass in Chinese male nuclear families. Osteoporos
Int 27:1805–1815.

33.   Kuchenbaecker KB, Ramus SJ, Tyrer J, Lee A, Shen HC, Beesley J, Lawrenson K, McGuffog L,
Healey S, Lee JM, Spindler TJ, Lin YG, Pejovic T, Bean Y, Li Q, Coetzee S, Hazelett D, Miron
A, Southey M, Terry MB, Goldgar DE, Buys SS, Janavicius R, Dorfling CM, Van Rensburg EJ,
Neuhausen SL, Ding YC, Hansen TVO, Jønson L, Gerdes AM, Ejlertsen B, Barrowdale D, Dennis
J, Benitez J, Osorio A, Garcia MJ, Komenaka I, Weitzel JN, Ganschow P, Peterlongo P, Bernard
L, Viel A, Bonanni B, Peissel B, Manoukian S, Radice P, Papi L, Ottini L, Fostira F,
Konstantopoulou I, Garber J, Frost D, Perkins J, Platte R, Ellis S, Godwin AK, Schmutzler RK,
Meindl A, Engel C, Sutter C, Sinilnikova OM, Damiola F, Mazoyer S, Stoppa-Lyonnet D, Claes
K, De Leeneer K, Kirk J, Rodriguez GC, Piedmonte M, O'Malley DM, De La Hoya M, Caldes T,
Aittomäki K, Nevanlinna H, Margriet JC, Rookus MA, Oosterwijk JC, Tihomirova L, Tung N,
Hamann U, Isaccs C, Tischkowitz M, Imyanitov EN, Caligo MA, Campbell IG, Hogervorst FBL,
Olah E, Diez O, Blanco I, Brunet J, Lazaro C, Pujana MA, Jakubowska A, Gronwald J, Lubinski
J, Sukiennicki G, Barkardottir RB, Plante M, Simard J, Soucy P, Montagna M, Tognazzo S,
Teixeira MR, Pankratz VS, Wang X, Lindor N, Szabo CI, Kauff N, Vijai J, Aghajanian CA,
Pfeiler G, Berger A, Singer CF, Tea MK, Phelan CM, Greene MH, Mai PL, Rennert G, Mulligan
AM, Tchatchou S, Andrulis IL, Glendon G, Toland AE, Jensen UB, Kruse TA, Thomassen M,
Bojesen A, Zidan J, Friedman E, Laitman Y, Soller M, Liljegren A, Arver B, Einbeigi Z,
Stenmark-Askmalm M, Olopade OI, Nussbaum RL, Rebbeck TR, Nathanson KL, Domchek SM,
Lu KH, Karlan BY, Walsh C, Lester J, Hein A, Ekici AB, Beckmann MW, Fasching PA,
Lambrechts D, Van Nieuwenhuysen E, Vergote I, Lambrechts S, Dicks E, Doherty JA, Wicklund
KG, Rossing MA, Rudolph A, Chang-Claude J, Wang-Gohrke S, Eilber U, Moysich KB, Odunsi
K, Sucheston L, Lele S, Wilkens LR, Goodman MT, Thompson PJ, Shvetsov YB, Runnebaum IB,
Dürst M, Hillemanns P, Dörk T, Antonenkova N, Bogdanova N, Leminen A, Pelttari LM, Butzow
R, Modugno F, Kelley JL, Edwards RP, Ness RB, Du Bois A, Heitz F, Schwaab I, Harter P,
Matsuo K, Hosono S, Orsulic S, Jensen A, Kjaer SK, Hogdall E, Hasmad HN, Noor Azmi MA,
Teo SH, Woo YL, Fridley BL, Goode EL, Cunningham JM, Vierkant RA, Bruinsma F, Giles GG,
Liang D, Hildebrandt MAT, Wu X, Levine DA, Bisogna M, Berchuck A, Iversen ES, Schildkraut
JM, Concannon P, Weber RP, Cramer DW, Terry KL, Poole EM, Tworoger SS, Bandera E V.,
Orlow I, Olson SH, Krakstad C, Salvesen HB, Tangen IL, Bjorge L, Van Altena AM, Aben KKH,
Kiemeney LA, Massuger LFAG, Kellar M, Brooks-Wilson A, Kelemen LE, Cook LS, Le ND,
Cybulski C, Yang H, Lissowska J, Brinton LA, Wentzensen N, Hogdall C, Lundvall L,
Nedergaard L, Baker H, Song H, Eccles D, McNeish I, Paul J, Carty K, Siddiqui N, Glasspool R,
Whittemore AS, Rothstein JH, McGuire V, Sieh W, Ji BT, Zheng W, Shu XO, Gao YT, Rosen B,
Risch HA, McLaughlin JR, Narod SA, Monteiro AN, Chen A, Lin HY, Permuth-Wey J, Sellers
TA, Tsai YY, Chen Z, Ziogas A, Anton-Culver H, Gentry-Maharaj A, Menon U, Harrington P,

Lee AW, Wu AH, Pearce CL, Coetzee G, Pike MC, Dansonka-Mieszkowska A, Timorek A, Rzepecka IK, Kupryjanczyk J, Freedman M, Noushmehr H, Easton DF, Offit K, Couch FJ, Gayther S, Pharoah PP, Antoniou AC, Chenevix-Trench G. 2015. Identification of six new susceptibility loci for invasive epithelial ovarian cancer. Nat Genet 47:164–171.

34. Brisken C, Heineman A, Chavarria T, Elenbaas B, Tan J, Dey SK, McMahon JA, McMahon AP, Weinberg RA. 2000. Essential function of Wnt-4 in mammary gland development downstream of progesterone signaling. Genes Dev 14:650–4.

35. Alexander CM, Goel S, Fakhraldeen S a., Kim S. 2012. Wnt signaling in mammary glands: plastic cell fates and combinatorial signaling. Cold Spring Harb Perspect Biol 4.

36. Rajaram RD, Buric D, Caikovski M, Ayyanan A, Rougemont J, Shan J, Vainio SJ, Yalcin-Ozuysal O, Brisken C. 2015. Progesterone and Wnt4 control mammary stem cells via myoepithelial crosstalk. EMBO J 34:641–652.

37. Joshi PA, Jackson HW, Beristain AG, Di Grappa MA, Mote PA, Clarke CL, Stingl J, Waterhouse PD, Khokha R. 2010. Progesterone induces adult mammary stem cell expansion. Nature 465:803–807.

38. Meier-Abt F, Milani E, Roloff T, Brinkhaus H, Duss S, Meyer DS, Klebba I, Balwierz PJ, van Nimwegen E, Bentires-Alj M. 2013. Parity induces differentiation and reduces Wnt/Notch signaling ratio and proliferation potential of basal stem/progenitor cells isolated from mouse mammary epithelium. Breast Cancer Res 15:R36.

39. Kim YC, Clark RJ, Pelegri F, Alexander CM. 2009. Wnt4 is not sufficient to induce lobuloalveolar mammary development. BMC Dev Biol 9:55.

40. Sikora MJ, Cooper KL, Bahreini A, Luthra S, Wang G, Chandran UR, Davidson NE, Dabbs DJ, Welm AL, Oesterreich S. 2014. Invasive lobular carcinoma cell lines are characterized by unique estrogen-mediated gene expression patterns and altered tamoxifen response. Cancer Res 74:1463–74.

41. Sikora MJ, Jacobsen BM, Levine K, Chen J, Davidson NE, Lee A V, Alexander CM, Oesterreich S. 2016. WNT4 mediates estrogen receptor signaling and endocrine resistance in invasive lobular carcinoma cell lines. Breast Cancer Res 18:92.

42. Ciriello G, Gatza ML, Beck AH, Wilkerson MD, Rhie SK, Pastore A, Zhang H, McLellan M, Yau C, Kandoth C, Bowlby R, Shen H, Hayat S, Fieldhouse R, Lester SC, Tse GMK, Factor RE, Collins LC, Allison KH, Chen Y-Y, Jensen K, Johnson NB, Oesterreich S, Mills GB, Cherniack AD, Robertson G, Benz C, Sander C, Laird PW, Hoadley KA, King TA, TCGA Research Network, Perou CM. 2015. Comprehensive Molecular Portraits of Invasive Lobular Breast Cancer. Cell 163:506–519.

43. Proffitt KD, Virshup DM. 2012. Precise regulation of porcupine activity is required for physiological Wnt signaling. J Biol Chem 287:34167–78.

44. Glaeser K, Urban M, Fenech E, Voloshanenko O, Kranz D, Lari F, Christianson JC, Boutros M. 2018. ERAD-dependent control of the Wnt secretory factor Evi. EMBO J 37:e97311.

45. Willert K, Nusse R. 2012. Wnt proteins. Cold Spring Harb Perspect Biol 4:a007864.

46. Kispert A, Vainio S, McMahon AP. 1998. Wnt-4 is a mesenchymal signal for epithelial transformation of metanephric mesenchyme in the developing kidney. Development 125:4225–34.

47. Galli LM, Santana F, Apollon C, Szabo LA, Ngo K, Burrus LW. 2018. Direct visualization of the Wntless-induced redistribution of WNT1 in developing chick embryos. Dev Biol 439:53–64.

48. Rawadi G, Vayssière B, Dunn F, Baron R, Roman-Roman S. 2003. BMP-2 controls alkaline phosphatase expression and osteoblast mineralization by a Wnt autocrine loop. J Bone Miner Res 18:1842–53.

49. Ching W, Hang HC, Nusse R. 2008. Lipid-independent secretion of a Drosophila Wnt protein. J
Biol Chem 283:17092–8.

50. Richards MH, Seaton MS, Wallace J, Al-Harthi L. 2014. Porcupine is not required for the
production of the majority of Wnts from primary human astrocytes and CD8+ T cells. PLoS One
9:e92159.

51. Galli LM, Zebarjadi N, Li L, Lingappa VR, Burrus LW. 2016. Divergent effects of Porcupine and
Wntless on WNT1 trafficking, secretion, and signaling. Exp Cell Res 347:171–83.

52. Chen Q, Takada R, Takada S. 2012. Loss of Porcupine impairs convergent extension during
gastrulation in zebrafish. J Cell Sci 125:2224–34.

53. Kurita Y, Ohki T, Soejima E, Yuan X, Kakino S, Wada N, Hashinaga T, Nakayama H, Tani J,
Tajiri Y, Hiromatsu Y, Yamada K, Nomura M. 2019. A High-Fat/High-Sucrose Diet Induces
WNT4 Expression in Mouse Pancreatic β-cells. Kurume Med J 1–8.

54. Yamamoto H, Awada C, Hanaki H, Sakane H, Tsujimoto I, Takahashi Y, Takao T, Kikuchi A.
2013. The apical and basolateral secretion of Wnt11 and Wnt3a in polarized epithelial cells is
regulated by different mechanisms. J Cell Sci 126:2931–43.

55. Lyons JP, Mueller UW, Ji H, Everett C, Fang X, Hsieh J-C, Barth AM, McCrea PD. 2004. Wnt-4
activates the canonical beta-catenin-mediated Wnt pathway and binds Frizzled-6 CRD: functional
implications of Wnt/beta-catenin activity in kidney epithelial cells. Exp Cell Res 298:369–87.

56. Jordan BK, Shen JH-C, Olaso R, Ingraham HA, Vilain E. 2003. Wnt4 overexpression disrupts
normal testicular vasculature and inhibits testosterone synthesis by repressing steroidogenic factor
1/beta-catenin synergy. Proc Natl Acad Sci U S A 100:10866–71.

57. Bernard P, Fleming A, Lacombe A, Harley VR, Vilain E. 2008. Wnt4 inhibits beta-catenin/TCF
signalling by redirecting beta-catenin to the cell membrane. Biol Cell 100:167–77.

58. Louis I, Heinonen KM, Chagraoui J, Vainio S, Sauvageau G, Perreault C. 2008. The signaling
protein Wnt4 enhances thymopoiesis and expands multipotent hematopoietic progenitors through
beta-catenin-independent signaling. Immunity 29:57–67.

59. Kurayoshi M, Yamamoto H, Izumi S, Kikuchi A. 2007. Post-translational palmitoylation and
glycosylation of Wnt-5a are necessary for its signalling. Biochem J 402:515–23.

60. Schackmann RCJ, van Amersfoort M, Haarhuis JHI, Vlug EJ, Halim VA, Roodhart JML, Vermaat
JS, Voest EE, van der Groep P, van Diest PJ, Jonkers J, Derksen PWB. 2011. Cytosolic p120-
catenin regulates growth of metastatic lobular carcinoma through Rock1-mediated anoikis
resistance. J Clin Invest 121:3176–3188.

61. Yamamoto TM, McMellen A, Watson ZL, Aguilera J, Sikora MJ, Ferguson R, Nurmemmedov E,
Thakar T, George-Lucian M, Kim H, Cittelly DM, Wilson H, Behbakht K, Bitler BG. 2018.
Targeting Wnt Signaling To Overcome PARP Inhibitor Resistance. bioRxiv 378463 [Pre-print].

62. Short B. 2017. The signal hypothesis matures with age. J Cell Biol 216:1207–1207.

63. Nakamura K, Shirai T, Morishita S, Uchida S, Saeki-Miura K, Makishima F. 1999. p38 mitogen-
activated protein kinase functionally contributes to chondrogenesis induced by
growth/differentiation factor-5 in ATDC5 cells. Exp Cell Res 250:351–63.

64. Sikora MJ, Jabobsen BM, O'Connor DP, Riggins RB, Stires H, Oesterreich S. 2017. ILC-CORE
(Collaboration, Openness, REproducibility). Open Sci Framew 19 Sept.

798
799

800   **Table 1 – Model Systems**

Cell Line	Tissue Type / Tumor (WNT4-responsive tissue)	WNT Over-expression (Wnt OE)		PORCN Activity
		WNT3A	**WNT4**	
MM134	ILC/Breast Cancer (Mammary Gland)	-	-	Normal
MM134:W3		Yes	-	Normal
MM134:W4		-	Yes	Normal
HT1080	Fibrosarcoma (Bone)	-	-	Normal
HT1080:W3		Yes	-	Normal
HT1080:W4		-	Yes	Normal
HT1080-PKO	Fibrosarcoma (Bone)	-	-	Inactive
HT1080-PKO:W3		Yes	-	Inactive
HT1080-PKO:W4		-	Yes	Inactive
SUM44PE	ILC/Breast Cancer (Mammary Gland)	(as above)		Normal
HCC1428	Breast Cancer (Mammary Gland)	(as above)		Normal
PEO1	Ovarian Cancer (Ovary)	(as above)		Normal

805   **List of Abbreviations**

807   AP - Alkaline Phosphatase
808   AXIN2 - Axin related protein 2
809   CDH1 - E-cadherin
810   CM - Conditioned Media
811   DVL - Disheveled
812   Fulv - Fulvestrant
813   FZD - Frizzled
814   ILC - Invasive Lobular Carcinoma
815   LRP - Lipoprotein receptor-related protein
816   PKO - Porcupine knockout
817   PORCN - O-acyltransferase Porcupine
818   PORCNi - Porcupine inhibitor
819   rWNT - Recombinant WNT
820   sWNT - Secreted WNT
821   WLS - Wntless
822   WNT - Wingless/Integrated

Figure 1. Inhibition or knockdown of PORCN does not phenocopy knockdown of WNT4 in MM134.
(A), MM134 cells were transfected with siRNA or treated with fulvestrant (100nM), LGK974 (10nM), or 0.1% vehicle (EtOH or DMSO) at time 0, prior to live cell imaging for proliferation (phase-contrast confluence) and death (SyTOX green incorporation). Points represent mean of 6 biological replicates ±SD. (B), Total double-stranded DNA was measured from assays in (A) at timecourse completion. *, p<0.05 vs control, ANOVA with Dunnett's multiple correction. Results in A,B are representative of three independent experiments.

Figure 2. WNT4 secretion is PORCN-independent, but WLS-dependent.
(A), MM134 constitutive Wnt over-expression cell lines were created as detailed in Methods and Materials. Top, qPCR for WNT3A and WNT4. Bars represent mean of 2 technical replicates ±SD. Bottom, immunoblot for cellular expression of WNT3A and WNT4. Endogenous WNT4 could not be visualized here due to the level of over-expression (B), Top, MM134 were treated with 10nM LGK974, and medium was allowed to condition for 7 days. Bottom, MM134 were transfected with siPORCN or siWLS, and after 24hrs, medium was changed and conditioned for 7 days. Total protein was extracted from medium as above for immunoblot. (C), HT1080 constitutive Wnt over-expression cell lines were created as detailed in Methods and Materials. Left, qPCR for WNT3A and WNT4. Bars represent mean of 3 technical replicates ±SD. Right, immunoblot for cellular expression of WNT3A and WNT4. Over-expression shows endogenous WNT4 expression. (D), HT1080 medium was conditioned for 5 days as described in Materials and Methods, prior to total protein extraction for immunoblot. (E), HT1080 cells were transfected with siPORCN or siWLS, and after 24hrs, medium was changed and conditioned for 5 days. Total protein was extracted from medium as above for immunoblot.

Figure 3. WNT4 does not activate paracrine Wnt signaling.
(A-D) HT1080-PKO control cells were treated for 24 hours with conditioned media (CM, at 50%, 25% or 12.5% final volume supplemented with fresh medium) from either MM134 (A-B) or HT1080 cell lines (C-D). (A,C) Immunoblots of whole cell lysates from the treated HT1080-PKO cells were run and probed for DVL2, DVL3, total LRP6 and phosphorylated LRP6. (B,D) mRNA from the treated HT1080-PKO cells was extracted for qPCR for AXIN2 mRNA expression levels, vs RPLP0. Points represent mean of 2 technical replicates ±SD. (B,D) Cells treated with 62.5ng/mL rWNT3A as a positive control. Statistics obtained using ANOVA with Dunnett's multiple correction. (E) HT1080-PKO cells were transfected with a TOP-FLASH reporter plasmid, then co-cultured with either HT1080 or HT1080-PKO WNT overexpressing (OE) cells, with or without 10nM LGK974. 'Ctrl' represents TOP-FLASH transfected HT1080-PKO without co-culture. WNT signaling activity, as measured by luminescence, was performed using a dual luciferase assay. Statistics obtained using Student's unpaired t-test compared to the no co-culture control (ctrl). * represents $p < 0.005$. Points represent mean of 3 technical replicates ±SD. Results are representative of two independent experiments. (F-G) HT1080-PKO control cells were treated for 24 hours with recombinant WNT protein (at concentrations of 10ng/ml, 50ng/ml, 100ng/ml, 250ng/ml, or 500ng/ml). (F) Immunoblots of whole cell lysates were run and (G) mRNA extracted for qPCR were performed as above (A-D). Statistics obtained using ANOVA with Dunnett's multiple correction. Points represent mean of 2 technical replicates ±SD.

3T3-E1 Alkaline Phosphatase Activity

Figure 4. Secreted versus recombinant Wnt protein differentially activate paracrine signaling.
(A-C), MC3T3-E1 cells were plated into a 96-well plate and 24hrs later were treated with (A), rWNT3A (0, 18.75, 32.5, 75ng/ml) or rWNT4 (0, 125, 250, 500ng/ml), (B), rWNT3A (62.5ng/ml), rWNT4 (250ng/ml) or 50% conditioned media (CM) from HT1080 cell lines, or (C), 50% CM from HT1080 or MM134 cell lines that were treated with or without 10nM LGK974. Points represent a mean of 4 (A), 3 (B) and 6 (C) biological replicates ±SD. Estimated concentrations of secreted WNTs (sWNT) present in CM based on linear regression of rWNTs is shown in Supplemental Fig. 4. *, p<0.05 vs control, ANOVA with Dunnett's multiple correction.

Figure 5. WNT4 from various cell types lacks paracrine activity.
Western blots of whole cell lysate or conditioned media from WNT3A or WNT4 overexpressing (A) SUM44PE, (D) HCC1428, or (F) PEO1 cell lines were probed for either WNT3A or WNT4. (B), A western blot of conditioned media from SUM44PE WNT4 over expressing cells treated with either LGK974, siPORCN or siWLS. Statistics obtained using ANOVA with Dunnett's multiple correction. Alkaline phosphatase production induced by conditioned media from WNT3A or WNT4 overexpressing (C) SUM44PE, (E) HCC1428 or (G) PEO1 cell lines was measured. Points represent a mean of 4 biological replicates ±SD.

Figure 6. WNT4 and WNT3A have cell autonomous activity independent of PORCN.
(A), Whole cell lysates from HT1080 cell lines were harvested and immunoblotted for DVL2, DVL3, total LRP6 and phosphorylated LRP6. (B), mRNA from HT1080 cell lines was extracted and qPCR was performed to determine expression levels of AXIN2. Points represent a mean of 3 technical replicates ±SD. (C-D), HT1080 (WT or PORCN null) cell lines transfected with siControl, siWLS, or siWNT4. (C), Cells were live cell imaged for cell death (SyTOX green) and (D), total double-stranded DNA was measured at timecourse completion. Points represent a mean of 6 biological replicates ±SD. (E), HT1080-PKO cell lines were transfected as indicated. Whole cell lysates were harvested and immunoblotted for DVL3.

Figure 7. WNT4 activates signaling in an atypical manner independent of secretion.
WNT4 can be modified by PORCN during the process of secretion, but WNT4 can instead be secreted in a PORCN-independent, WLS-dependent manner. In either case, secreted WNT4 does not have paracrine activity in any context tested herein. Cell-autonomous WNT4 signaling can activate DVL proteins intracellularly, and likely signals via other pathways in ILC cells.

Supplemental Figure 1. PORCNi (WntC59) does not impact ILC growth.
MM134 and SUM44PE were plated and 24hrs later started treatment with either anti-estrogen fulvestrant (Fulv) or increasing concentrations of WntC59. At the timecourse completion of either 5 or 7 days, total ATP was measured using Promega Cell-Titer Glo. Points represent the mean of 6 biological replicates ± SD.

Supplemental Figure 2. Experimental Workflow for Assessing Wnt Secretion and Function.
Diagram of the workflow for experiments aimed at assessing the ability of Wnt proteins to be secreted and their functionality. Experimental endpoints are labeled with corresponding figures.

A

B

Supplemental Figure 3. WNT4 secretion is PORCN-independent, but secreted WNT4 is post-translationally modified.

(A), HT1080 cells were reverse transfected with siWNT4. 24hrs later media was changed and cells were treated with or without LGK974 (10nM). Conditioned media (CM) was harvested 5 days later and total protein was collected from immunoblotting (as described in the Methods and Materials) for WNT3A or WNT4. **(B),** Conditioned media was harvested from either HT1080:WT or MM134:Wnt4 over-expressing cells after treatment (3 or 5 days respectively) with increasing concentrations of tunicamycin (0.5uM, 1uM or 10uM). WNT4 presence was detected from untreated whole cell lysate and conditioned media.

A

3T3-E1

Treatment:　−　r3A　r4

DVL2

DVL3

tLRP6

pLRP6

Vinculin

B

HT1080
Conditioned Medium

C

MM134
Conditioned
Medium

D

Cell Line	Concentration (ng/ml)	
	sWNT3A	sWNT4
HT1080	-	-
HT1080 WNT3A	62	-
HT1080 WNT4	-	190
HT1080-PKO	-	42
HT1080-PKO WNT3A	-	55
HT1080-PKO WNT4	-	64
MM134	-	-
MM134 WNT3A	157	-
MM134 WNT4	-	1808

Supplemental Figure 4. Secreted Wnt proteins are at concentrations sufficient to activate paracrine signaling.
(A), 3T3-E1 cells were treated for 24hrs with 300ng/ml of either rWNT3A or rWNT4. Immunoblots of whole cell lysates were probed for DVL2, DVL3, total LRP6 and phosphorylated LRP6. (E-F), Serial dilutions of rWNTs (2.5, 5, 10, 20ng) were run on an immunoblot alongside CM collected from HT1080 or MM134 cell lines. Blots were probed for either WNT3A or WNT4, respectively. (G), Estimated concentrations of secreted WNTs (sWNT) present in CM based on linear regression of rWNTs. Volume of CM loaded per well varied from 25-70uL, normalized based on total protein concentration (see Materials and Methods).

Supplemental Figure 5. siWLS does not phenocopy siWNT4 and CM does not rescue siWNT4 in MM134.
(A-B), MM134 were reverse transfected with siRNA as indicated. **(C-D),** 24hrs after WNT4 knockdown, cells
were treated with either rWNT3A (62.5mg/ml), rWNT4 (250ng/ml) or conditioned media (CM). The cells were
live cell imaged for **(A,C),** proliferation (phase-contrast confluence) and death (SyTOX green incorporation).
(B,D), Total double stranded DNA was neasured from assays in **(A,C)** at timecourse completion. Point repre-
sent mean of six biological replicates ± SD.

A

Supplemental Figure 6. WNT overexpression does not induce autonomous phosphorylation of DVL2/3 in MM134.
Whole cell lysates from MM134 were harvested and immunoblotted for DVL2, DVL3, total and phosphorylated LRP6.

A

B

Wnt Source	Wnt OE	Mode	Receiver	Active?
HT1080	WNT4	Paracrine	HT1080:PKO	No
HT1080	WNT4	Paracrine	3T3-E1	No
HT1080:PKO	WNT4	Paracrine	HT1080:PKO	No
HT1080:PKO	WNT4	Paracrine	3T3-E1	No
MM134	WNT4	Paracrine	HT1080:PKO	No
MM134	WNT4	Paracrine	3T3-E1	No
HCC1428	WNT4	Paracrine	3T3-E1	No
PEO1	WNT4	Paracrine	3T3-E1	No
SUM44PE	WNT4	Paracrine	3T3-E1	No
Recombinant	rWNT4	Paracrine	HT1080:PKO	No
Recombinant	rWNT4	Paracrine	3T3-E1	Yes
HT1080	WNT4	Autonomous	(HT1080)	Yes
HT1080:PKO	WNT4	Autonomous	(HT1080:PKO)	Yes
MM134	WNT4	Autonomous	(MM134)	Yes

C

Wnt Source	Wnt OE	Mode	Receiver	Active?
HT1080	WNT3A	Paracrine	HT1080:PKO	Yes
HT1080	WNT3A	Paracrine	3T3-E1	Yes
HT1080:PKO	WNT3A	Paracrine	HT1080:PKO	No*
HT1080:PKO	WNT3A	Paracrine	3T3-E1	No*
MM134	WNT3A	Paracrine	HT1080:PKO	No
MM134	WNT3A	Paracrine	3T3-E1	Yes
HCC1428	WNT3A	Paracrine	3T3-E1	Yes
PEO1	WNT3A	Paracrine	3T3-E1	Yes
SUM44PE	WNT3A	Paracrine	3T3-E1	No
Recombinant	rWNT3A	Paracrine	HT1080:PKO	Yes
Recombinant	rWNT3A	Paracrine	3T3-E1	Yes
HT1080	WNT3A	Autonomous	(HT1080)	Yes
HT1080:PKO	WNT3A	Autonomous	(HT1080:PKO)	Yes
MM134	WNT3A	Autonomous	(MM134)	n/a

*No WNT3A secretion from HT1080-PKO

Supplemental Figure 7. Summary of paracrine Wnt activity per source and receiver context.
(A), WNT3A can be secreted independent of PORCN from ILC cells, but PORCN function is overall required for paracrine activity. However, paracrine activity of WNT3A was dependent on the Wnt source and the receiver cell context. (B-C), Summary of results herein for (B) WNT4 and (C) WNT3A.