1 **DATE: 2018/05/29**
2 **VERSION: 5**
3
4 **Title: Pan-cancer machine learning predictors of tissue of origin and molecular subtype**

6 **Authors**

Author	Symbol
William F. Flynn	1¶
Sandeep Namburi	1¶
Carolyn A. Paisie	1
Honey V. Reddi	1,2
Sheng Li	1,2*
R. Krishna Murthy Karuturi	1,2*
Joshy George	1,2*

Affiliations

1. The Jackson Laboratory for Genomic Medicine, 10 Discovery Drive, Farmington,

Connecticut, USA

2. The Jackson Laboratory Cancer Center, Bar Harbor, Maine, USA

¶ These authors contributed equally to this work.

* Corresponding Authors: sheng.li@jax.org, krishna.karuturi@jax.org, joshy.george@jax.org

ABSTRACT

Background: It is estimated by the American Cancer Society that approximately 5% of all

metastatic tumors have no defined primary tissue of origin and are classified as <u>c</u>ancers of

<u>unknown primary</u> origin (CUPs). The current standard of care for CUP patients depends on

immunohistochemistry (IHC) based approaches to identify the primary site. The addition of post-

mortem evaluation to IHC based tests helps to reveal the identity of the primary site for only

25% of the CUPs, emphasizing the acute need for better methods of determination of the site of

origin. CUP patients are therefore given generic chemotherapeutic agents resulting in poor

prognosis. When the tissue of origin is known, patients can be given site specific therapy with

significant improvement in clinical outcome. Similarly, identifying the primary origin of metastatic

cancer is of great importance for designing treatment.

Identification of the primary site of origin is an import first step but may not be sufficient

information for optimal treatment of the patient. Recent studies, primarily from The Cancer

Genome Atlas (TCGA) project, and others, have revealed molecular subtypes in several cancer

types with distinct clinical outcome. The molecular subtype captures the fundamental

mechanisms driving the cancer and provides information that is essential for the optimal

treatment of a cancer. Thus, along with primary site of origin, molecular subtype of a tumor is

emerging as a criterion for personalized medicine and patient entry into clinical trials.

However, there is no comprehensive toolset available for precise identification of tissue of origin

or molecular subtype for precision medicine and translational research.

Methods and Findings: We posited that metastatic tumors will harbor the gene expression

profiles of the primary tissue of origin of the cancer. Therefore, we decided to learn the

characteristics of the primary tumors using the large number of cancer genome profiles

available from the TCGA project. Our predictors were trained for 33 cancer types and for the 11

cancers where there are established molecular subtypes. We estimated the accuracy of several

machine learning models using cross-validation methods and external validation sets. The

extensive testing using independent test sets revealed that the predictors had a median

sensitivity and specificity of 97.2% and 99.9% respectively without losing classification of any

tumor. Subtype classifiers achieved median sensitivity of 87.7% and specificity of 94.5% via

cross validation and presented median sensitivity of 79.6% and specificity of 94.6% in two

external datasets of 1,999 total samples. Importantly, these external data shows that our

classifiers can robustly predict the cancer primary origin from microarray data, metastatic

cancer, and patient-derived xenograft (PDX) mouse models.

Conclusion: We have demonstrated the utility of gene expression profiles to solve the

important clinical challenge of identifying the primary site of origin and the molecular subtype of

cancers based on machine learning algorithms. We show, for the first time to our knowledge,

that our pan-cancer classifiers can predict multiple cancers' primary tissue of origin from

metastatic samples. The predictors will be made available as open source software, freely

available for academic non-commercial use.

KEYWORDS

Cancer, TCGA, RNA-seq, Classification, Subtypes, Transcriptome, Machine learning, Cell-of-

origin, Cancer-of-unknown-primary

1. INTRODUCTION

Precision cancer therapy requires the knowledge of tissue of origin and accurate subtyping of the cancer to identify an appropriate therapeutic regimen. However, according to the American Cancer Society, an estimated 2 to 5 percent of all cancer patients have metastatic tumors for which routine testing cannot locate the primary site and is therefore classified as a cancer of unknown primary (CUP). CUP patients have very poor prognosis, primarily because the course of treatment is generic and not tailored for a specific tumor type [1, 2]. In addition, lack of knowledge of the cause of the disease puts CUP patients under severe psychological distress that may lead to clinically significant depressive symptoms [3].

In a study of CUP patients that were predicted to have primary tumors originating in the colon, the median survival of patients increased in those that received site-specific chemotherapy as compared to those who received empirically-determined treatments [4, 5]. Furthermore, multiple studies have supported the use of molecular profiling to diagnose CUP and to determine specific treatment based upon the predicted tissue of origin leading to an improvement in overall survival [6-9]. Therefore, it is important to develop systematic methods to identify the primary site of origin of the disease.

Currently, immunohistochemistry (IHC) utilizing antibodies targeted to certain tumor-specific antigens is the main method for primary tumor identification in patients with CUP [2, 10]. However, there is not a single, specific marker that can be used to conclusively diagnose the primary tumor, leading to the use of multiple different IHC markers, and generating the possibility that different clinicians will arrive at different diagnoses of the primary tumor type [2, 11-14]. Recent advances in genomics has led to the development of potential new means for diagnosing these tumors; multiple studies have utilized gene expression profiles and other molecular markers to diagnose CUP [2]. One such method involved comparisons of gene

expression profiles between CUPs and a set of primary and metastatic tumors with known

origins to predict the tissue of origin for the CUP [2, 6, 15-20]. None of these tools were able to

identify the tissue of origin with high sensitivity and specificity for all the CUP samples. For

example, the EPICUP tool, which had the best performance to date provides high accuracy for

only 87% of the CUP samples.

Several studies, including the Cancer Genome Atlas (TCGA) and International Cancer Genome

Consortium (ICGC) studies, have shown that the cancers from the primary tissue can be

classified into molecular subtypes with distinct clinical outcome and therapeutic options [21-28].

The molecular subtype information can also have predictive power. For example, Bevacizumab,

a monoclonal antibody that block angiogenesis, is shown to benefit patients of mesenchymal

and proliferative subtypes in ovarian cancer and therefore may be used as a criterion for the

entry into a clinical trial [29]. In addition, molecular subtype information can guide the selection

of targeted therapies and to suggest new treatment strategies (e.g. the potential use of JAK2

inhibitors and PD-L1/2 antagonists for the treatment of EBV-positive gastric cancer) [27].

However, identification of molecular subtypes is clinically challenging and thus clinicians are

unable to utilize molecular subtype information to inform treatment decisions [30, 31] due to a

lack of tools and assays for pan-cancer subtyping despite the availability of genomic

technologies for clinical diagnostics.

To fill this important gap in the clinical and translational research setting, using expression data

available from the TCGA project, we developed Machine Learning based predictors that enable

accurate identification of tissue of origin and subtype of cancer (Figure 1). Our predictors were

trained for 33 cancer types and for the 11 cancers where there are established molecular

subtypes. The extensive testing using independent test sets revealed that the predictors had a

median sensitivity and specificity of 97.2% and 99.9% respectively without losing classification

of any tumor using in total 1,959 samples. Subtype classifiers achieved median sensitivity of

79.6% and specificity of 94.6% using two external validation sets consisting of 1,999 breast and

ovarian cancer samples in total. Our gene expression-based pan-cancer classifier can, for the

first time to our knowledge, robustly predict multiple cancers' primary tissue of origin from

metastatic samples using an independent validation dataset (specificity: 99.3%, sensitivity:

82.1%). Compared to other pan-cancer classifiers based on somatic mutations, our classifier is

not limited to only cancer types with high mutation burden and has much greater potential for

clinical diagnosis and therapeutic design.

2. METHODS

2.1 Expression datasets

2.1.1 Learning set: TCGA expression data

RSEM [32] normalized mRNA expression matrices were downloaded for each of the 33 unique

cancer cohorts (listed in Table 1) available from the Broad Institute GDAC Firehose (run

2016_01_28) [33]. Individual cohort expression matrices were converted to Biobase

ExpressionSet objects [34] for standardization and then combined as an ExpressionSet stored

in Apache Feather format (version 0.4.0) to allow downstream analysis in both R and Python

(https://github.com/wesm/feather). The raw expression matrix consisted 11,330 samples and

20,531 genes, which was reduced to 10,446 samples and 9,642 genes after filtering (detailed

below).

136
137 **Table 1. 33 GDAC Cancer Cohorts**

Cohort Abbr.	Cases	Disease Name
ACC	92	Adrenocortical carcinoma
BLCA	412	Bladder urothelial carcinoma
BRCA	1,098	Breast invasive carcinoma
CESC	307	Cervical and endocervical cancers
CHOL	51	Cholangiocarcinoma
COAD	460	Colon adenocarcinoma
DLBC	58	Lymphoid Neoplasm Diffuse Large B-cell Lymphoma

ESCA	185	Esophageal carcinoma
GBM	613	Glioblastoma multiforme
HNSC	528	Head and Neck squamous cell carcinoma
KICH	113	Kidney Chromophobe
KIRC	537	Kidney renal clear cell carcinoma
KIRP	323	Kidney renal papillary cell carcinoma
LAML	200	Acute Myeloid Leukemia
LGG	516	Brain Lower Grade Glioma
LIHC	377	Liver hepatocellular carcinoma
LUAD	585	Lung adenocarcinoma
LUSC	504	Lung squamous cell carcinoma
MESO	87	Mesothelioma
OV	602	Ovarian serous cystadenocarcinoma
PAAD	185	Pancreatic adenocarcinoma
PCPG	179	Pheochromocytoma and Paraganglioma
PRAD	499	Prostate adenocarcinoma
READ	171	Rectum adenocarcinoma
SARC	261	Sarcoma
SKCM	470	Skin Cutaneous Melanoma
STAD	443	Stomach adenocarcinoma
TGCT	150	Testicular Germ Cell Tumors
THCA	503	Thyroid carcinoma
THYM	124	Thymoma
UCEC	560	Uterine Corpus Endometrial Carcinoma
UCS	57	Uterine Carcinosarcoma
UVM	80	Uveal Melanoma

2.1.2 External validation data

External cancer gene expression datasets, i.e. those used exclusively for testing the

performance of models, were obtained from the publicly available Gene Expression Omnibus

(GEO) and the patient-derived xenograft (PDX) mouse models generated at the Jackson

Laboratory. These datasets were not introduced during model fitting (hence external validation)

and were generated using both RNA-seq and microarray technologies.

The first dataset used to test the model accuracy was the microarray gene expression profile of

2,158 cancer samples from the expression project for oncology (expO, GSE2109) [35]. Of the

2,158 samples, we were able to identify relevant primary tumor types for 1,558 samples and

excluded LGG/GBM from classification due to too few samples, resulting in classification of

1,552 samples. The second dataset, GSE18549, contained expression profiles of 96 tumors

from their metastatic sites [36]. We used 88 of these tumors whose primary sites could be

identified and those with more than 1 sample per primary site. The third dataset used contains

the expression profile of 338 PDX RNA-seq samples generated at the Jackson Laboratory and

available through the Mouse Tumor Biology (MTB) gene expression portal

(http://tumor.informatics.jax.org) [37]. Of these 338 samples, 325 samples could be mapped to

one of the 33 TCGA primary cancer types and the 7 OV samples were excluded as they

originate from the same two patients. The distribution of the primary types in the external

datasets used for validation are shown in Table 2.

Table 2. Distribution of samples in the three external datasets used to validate the

primary classification model

Primary site	External dataset		
	GSE2109	GSE18549	PDX
BLCA	32	0	29
BRCA	354	14	41
COAD	312	35	68
DLBC	0	0	4
KIRC/KIRP/KIRH	281	6	8
LAML	0	0	13
LGG/GBM	6	0	5
LIHC	46	0	0
LUAD/LUSC	133	10	88
OV	279	14	7
PAAD	0	0	12
PRAD	83	9	0
READ	0	1	0
SARC	0	0	32
SKCM	0	0	18
THCA	32	0	0
N/A	600	7	13
Used	1,552	88	318
Total	2,158	96	338

For external validation of our subtype classifiers, we acquired two additional microarray

datasets. The first, accession number GSE9899, contains 215 ovarian cancer samples [15] and

the second, EGA study EGAS00000000083 (https://www.ebi.ac.uk/ega), contains 1,784 breast

cancer samples [38]. Both datasets comprise 4 molecular subtypes each: mesenchymal,

immunoreactive, differentiated, and proliferative for the ovarian set; and basal-like, HER2-

enriched, luminal A, and luminal B for the breast set.

2.1.3 Normalization, filtering, and preprocessing

All expression data was log2-transformed, and only genes with (a) maximum log2 expression

greater than 8 and (b) variance in log2 expression greater than 1 were retained. After filtering,

the genes in each dataset was scaled to zero mean expression and unit variance. This scaling

allows expression to be measured in terms of standard deviations and affords platform-

independent use of subsequently trained models.

2.1.4 Molecular subtype label curation

Molecular subtype information was downloaded from cBioPortal [39, 40] for 3,367 samples from

the following primary cancers: glioblastoma multiforme (GBM), stomach adenocarcinoma

(STAD), breast (BRCA), ovarian (OV), prostate (PRAD), and lung squamous cell cancers

(LUSC). Further annotations were curated from the following supplemental data files: lower

grade glioma (LGG) [41], head and neck squamous cell carcinoma (HNSC) [22], uterine corpus

endometrial carcinoma (UCEC) [42], cutaneous melanoma (SKCM) [43], papillary (KIRP) [44]

and clear cell (KIRC) [45] renal cancers, and lung adenocarcinoma (LUAD) [27]. R (version 3+)

scripts were written to extract relevant information (e.g. sample id, specific subtype) from

downloaded data and supplemental files. These scripts are available in the public project

GitHub repository as described under Code availability.

2.1.5 Pan-organ group labels

A recently published study performed integrated clustering on the multiomic data from the

roughly 10,000 The Cancer Genome Atlas (TCGA) samples of 33 types of cancer. The authors

identified multi-cancer groups, which tend to span whole organs or related organ groups [46].

We use these pan-organ groups assignments to present our classification results in a whole- or

multi-organ context and for logical grouping of primary cancer types in figures. These

classifications are reproduced here: central nervous system (GBM LGG), core gastrointestinal

(ESCA, STAD, COAD, READ), developmental gastrointestinal (LIHC, PAAD, CHOL), endocrine

(THCA and ACC), gynecologic (OV, UCEC CESC BRCA), head and neck (HNSC), hematologic

and lymphatic malignancies (LAML, DLBC, THYM), melanocytic (SKCM and UVM), neural-

crest-derived tissues (PCPG), soft tissue (SARC and UCS), thoracic (LUAD, LUSC, MESO),

urologic (BLCA, PRAD, TGCT, KIRC, KICH, KIRP).

2.2 Machine Learning Algorithms for Cancer Classification

We evaluated several popular machine learning algorithms to develop CUP classification and

subtype identification: DLDA, KNN, SVM and Random Forest. We used R-packages

sparsediscrim (version 0.2.4)*, base::knn, e1071* (version 1.6-8)*,* and *randomForest* (version 4.6-

14), respectively for the training, testing; and *caret* (version 6.0-79) for tool development.

Unless otherwise specified, default parameters were chosen for model construction.

2.2.1 Diagonal Linear Discriminant Analysis (DLDA)

The DLDA classifier belongs to the family of Naive Bayes classifiers, where the distribution of

each class is assumed to be a multivariate normal and to share a common covariance matrix.

The DLDA classifier is a modification to LDA, where the off-diagonal elements of the pooled

sample covariance matrix are set to zero [47]. DLDA was used as classifier in several genomic

based cancer classification tasks [48, 49].

2.2.2 k-Nearest Neighbor (KNN) classifier

A KNN classifier offers the simplest classifier training strategy, also referred to as 'lazy

classifier', and has been successfully applied in the classification of cancer and non-cancer

related classification tasks [50-52]. A KNN classifier uses the training set as reference vectors

and, for every sample in the test set, the k nearest (in Euclidean distance) training set vectors

are found. The classification is decided by majority vote of the k-nearest neighbors' class. Note

that, if multiple neighbor vectors are found with identical distances, all such neighbor vectors are

included in the voting pool, which can lead to k being exceeded in these cases [53].

2.2.3 Support Vector Machine (SVM)

The SVM algorithm builds a predictive model by constructing a representation of the training

samples as points in higher dimensional space and builds a linear model (separating linear

boundary) in that space such that the mapped samples of the different categories are separated

by a gap that is as wide as possible. New examples are then mapped into that same space and

predicted to belong to a category based on which side of the separating boundary they fall. For

multiclass classification among N classes, N*(N-1)/2 binary SVM classifiers are constructed and

trained in a one-versus-one manner; ultimate class predictions come from voting amongst the

ensemble of binary classifiers. The implementation used for our classifiers employed a

Gaussian kernel. SVMs were successfully used for classification of samples in variety of studies

[54-56].

2.2.4 Random Forest

The random forest algorithm employs a collection of decision trees constructed from

bootstrapped input data and classification is done by majority voting among the ensemble of

trees [57]. Single decision trees are prone to overfitting; multiple trees constructed from

randomly sampled copies of the input data allows the consensus classification to be robust and

extensible to new samples. Each of our random forest models constructed 1000 trees, each

tree constructed from randomly sampled input with replacement, and each decision tree node

uses 31 randomly selected features to partition the tree.

2.3 Model training and external validation

The schema for classifier design for tumor type classification and subtype classification are

depicted in Figure 1.

2.3.1 Design of Tumor type classifier

All models were trained using the same feature selection and cross-validation schedule. Each

model was then trained using a 3-fold cross validation procedure as follows. The expression set

is partitioned into 3 random subsamples and for each partition: (1) the selected partition is used

as the testing set and the remaining 9 are combined into a training set; (2) the 100 most

differentially expressed genes in each class (cancer type) are selected, measured by log-fold-

change of differential expression between in-class and out-of-class samples ($p < 0.001$); (3) the

model is trained using the selected features; (4) predictions for the selected partition is

recorded. The cross-validation procedure yields an estimate of the model performance with the

selected parameters. The final model is then constructed using the entire set of samples and

1,971 unique genes selected via the procedure in (2) above.

2.3.2 Molecular subtype classification

For each of the 11 primary cancer types with established molecular subtypes, a model is

constructed as described above using the scaled, log2-transformed expression of the sample

corresponding to the selected primary type as input. For each cancer type, similar to cancer

type classification, features are selected by computing the differential expression (p<0.001) in

each subtype in comparison with the other subtypes of the same cancer type.

2.3.3 Comparison metrics

Each classification algorithm was compared using per class and overall positive predictive

value, sensitivity, and specificity. Additional metrics such as per-class balance accuracy, and

F1 score are included in the supplementary tables. Per class metrics are computed using a

one-versus-all scheme.

2.4 Visualization

An interactive web application was constructed using the Python Dash framework (version

0.21.0). The application shows the TCGA data embedded in three dimensions using UMAP

(umap-learn, version 0.2.1, [58]) and t-SNE (MulticoreTSNE, version 0.1, [59, 60]). Data points

are color coded by tumor type (with cancer and match normal samples), cancer type, and

molecular subtype, which can be controlled interactively through the interface.

The web application and the associated code are freely available for non-commercial, academic

use at https://pccportal.jax.org (pan-cancer classification portal) and the source code is

available as described in the Code availability subsection.

2.5 Code availability

All code to download, process, train, and validate these data and models is

available in the following GitHub repository:

. All results and the

majority of figures can be easily reproduced by cloning and running make.

3. RESULTS

3.1 Precise classification of primary cancer types across platforms

The classification of the 33 primary cancer types from the TCGA cohort (9,642 samples) by

random forest is presented in Figure 2A. Classification yields a median sensitivity and

specificity of 97.2% and 99.9% (n=33), respectively (Figure 2C). The major misclassifications

are primarily within organ systems (Figure 2B). Indeed, when primary types are grouped by

pan-organ groups [46], the median sensitivity increases to 98.5% with a substantial

improvement in the minimum sensitivity to 86.0% (Figure 2B,D). It is important to note that

every sample was classified by our model; no samples were excluded from classification, either

by a sample quality metric or through lack of consensus during label assignment. The most

frequent misclassifications occur between nearby locations in the gastrointestinal tract; rectal

adenocarcinoma (READ) is completely misclassified as colon adenocarcinoma (COAD), and

esophageal carcinoma (ESCA) is often misclassified as stomach adenocarcinoma

(STAD). COAD and READ are so similar that they are typically considered as a single primary

type, colorectal carcinoma (CRC) [23]. Misclassification between ESCA and STAD is expected,

as a certain class of ESCAs (esophageal adenocarcinomas) present at the interface of the

esophagus and stomach [61]. Also of note is the misclassification of uterine carcinosarcoma

(UCS) as uterine corpus endometrial carcinoma (UCEC). Histologically, USC presents features

of both UCEC and sarcoma (SARC) [62].

To understand these misclassifications, the expression profiles of every training sample was

embedded into a two-dimensional latent space using UMAP (see Methods) and colored by

primary tumor type, shown in Figure 3. Several anatomical and histological structures readily

emerge from the embedding. Some cancers are observed to form disparate, well-separated

clusters by organ system, such as brain (GBM-LGG), liver and gallbladder (LIHC, CHOL), and

kidneys (KIRC, KIRP, KIRH), while other cancers are grouped by histological features, such as

the melanomas (SKCM, UVM) and squamous cell cancers (BLCA, CECS, HNSC, LUSC, and

some ESCA) forming distinct clusters. The core gastrointestinal tract cancers cluster tightly, with

COAD and READ embedded into an inseparable mass which is adjoined by STAD and some

ESCA samples. ESCA samples clearly segregate into two populations, consistent with both

esophageal adenocarcinoma (clustered with STAD) and squamous cell carcinoma (clustered

with LUSC, HNSC, etc.) being classified under ESCA (Zheng 2013). Similarly, the known

similarities between USC, UCEC, and SARC clearly emerges, with the embedding of USC

forming a bridge between UCEC and SARC clusters; we also observe two distinct clusters of

SARC samples, one most similar to USC and the other most similar to UCEC. As this

embedding is heavily dependent on the input samples and number thereof, it may be that some

misclassifications are unavoidable without a larger cohort of samples.

3.1.1 Cancer type classifier performed well on external expression data

To further validate the primary type classifier, we classified 1,552 samples across 9 primary

tumor types (Figure 4A) profiled using microarrays from the Expression Project for Oncology

(expO, GSE2109 [35]). Due to the age of the dataset, primary cancer types corresponding to

the brain (LGG, GBM), lung (LUAD, LUSC), and kidney (KIRC, KIRP, KIRH) were aggregated to

match the respective primary tumor annotations of the dataset. Further, the genes used for

classification were reduced to 1,788 genes from the initially selected genes of 1,971 in order to

match those found in the external dataset, and the model was retrained on the training set with

only these 1,788 features. This external validation set not only tests the classifier performance

independent of batch variation, but also its independence of the platform and robustness to

feature loss which are critical for the application of the classifiers in clinical and translational

research.

Classification by primary cancer types, shown in Figure 4B, yields specificity of 99.3% and

median sensitivity of 78.1% (n=9) (Figure 4C), with misclassifications largely within organ

systems. For example, misclassification arises between gastrointestinal cancers STAD, COAD,

and LIHC, and ovarian serous cystadenocarcinoma (OV) is misclassified as cancers with similar

histology or anatomical location. When the classification is reorganized by pan-organ group, as

shown in Figure 4D, median sensitivity increases to 86.0% (n=6) with the misclassification only

between core and developmental gastrointestinal cancers (Figure 4E).

3.1.2 Classifier can identify cancer of unknown primary

Identification of the primary cancer (tissue) of origin from a metastatic sample is a significant

clinical challenge. As metastases are expected to retain the transcriptional signature of primary

tumor of origin, we hypothesize that our classifier can identify the primary tumor type from

metastatic samples. We examined the performance of our classifiers using an external

validating dataset from metastatic tumors.

Primary type classification of expression profiles of 88 metastatic samples across 6 known

primary types is shown in Figure 5A,B [36]. The median specificity is 99.3%, and the median

sensitivity is 82.1%. The most common misclassifications were, again, between COAD and

STAD, the vast majority involving metastatic tumors in the liver. Within-organ-system

classification (Figure 5C) shows the minimum sensitivity substantially improves from 20.0% to

69.0%, where predictions of these gastrointestinal cancers are combined (Figure 5D). Further

examination revealed that the most common metastases in the data, 30/88 (34%) samples, are

to the liver or lung from the colon, illustrated in Figure 5E. Of the 88 tumors, 52 metastases

(59.1%) are classified to the correct primary tumor type, 72 metastases (81.8%) are classified to

the correct primary organ system, 7 metastases (7.9%) are classified as the tumor from the

respective metastasized sites, and 9 metastases (10.1%) are classified incorrectly (Figure 5F)

i.e. neither as tumor of primary site nor as the metastasized site.

We further validate our classifiers independently using an external dataset from patient-derived

xenograft (PDX) models of cancer. PDX models of cancer are a great resource to evaluate

therapeutic regimens but can also be used as a tool to study metastatic cancer, as illustrated in

Figure 6A. Primary tumor is resected from human patients and tumor fragments are implanted

into a cohort of immunodeficient mice [63]. After a growth period, the mouse-grown tumor is

resected and implanted into a new generation of mice. This process can be repeated several

times.

We performed the primary tumor type classification of 318 PDX-derived mouse-grown tumors

(samples were taken from the second generation of mice) spanning 11 cancer types (Figure 6B-

C). Classification of primary cancer types yields a median specificity of 99.5% (n=11), and

median sensitivity of 76% (n=11) (Figure 6D). When classified by pan-organ system, the median

sensitivity increases to 83% (Figure 6E,F). Despite not being present in the set of primary

cancers, several tumors including COAD and PAAD are classified as STAD, possibly due to the

close proximity of anatomic positions.

These three external validations of our model overwhelmingly support the hypothesis that

metastatic and xenograft tumors retain the molecular signature of the primary

tumor. Application of such models in the clinic will allow for more effective treatments of CUPs.

3.2 Subtype specific classification accurately identifies molecular subtypes

Molecular subtypes have been defined for 11 cancer types: BRCA, HNSC, KIRC, KIRP, LGG,

LUAD, LUSC, OV, PRAD, SKCM, and STAD. Each of these primary types has two to four

molecular subtypes. For example, breast cancers are frequently subtyped into Basal-like, Her2-

enriched, Luminal A and Luminal B. This subtyping has prognostic power and can be used as

predictive marker for therapeutic approaches [64].

Eleven models were constructed, one model for each primary tumor type, into its molecular

subtypes, as illustrated schematically in Figure 1. The positive predictive value, sensitivity,

specificity, and number of samples per subtype are shown in Figure 7A-D. The best performing

subtype classifiers, LGG, LUAD, PRAD, have median sensitivity above 90%, with PRAD

yielding nearly perfect classification.

3.2.1 Subtype classifiers are accurate on external data of different platforms

To further validate the cancer subtype classifiers, we classified samples from two external

datasets: ovarian cancer [15] and breast cancer [38] annotated with molecular subtypes, with

215 and 1,784 samples respectively. The ovarian cancer subtype classifier (Figure 7E,F)

reached a median specificity of 94.9% (the best performance is for mesenchymal: 99.2%) and a

median sensitivity of 88.4% (the best performance is proliferative: 97.2%). The breast cancer

subtype classifier, shown in Figure 7G,H, presented a median specificity of 95% (the best

performance is for basal-like: 99.9%) and a median sensitivity of 72.4% (the best performance is

for luminal-A subtype: 95%). Notably, the basal-like molecular subtype of breast cancer is a

particularly aggressive subtype and patients relapse rapidly. The accurate classification of this

subtype is important for precise treatment—recently, a diabetes drug has been shown as a

potential therapy for basal-like breast cancer patients [65]. The two external validation datasets

from 1,999 patients with breast or ovarian cancers demonstrate that our subtype classifier can

distinguish clinically favorable subtypes from those associated with poor prognosis.

Together with the identification of primary origin for metastatic tumors, followed by subtype

classification together allows for informed therapeutic decision making for clinicians, thereby

improving treatment outcomes for CUP patients.

4. DISCUSSION

It is widely appreciated that cancer is a disease at the scale of the entire genome, but it remains

difficult to effectively translate this complexity into clinical utilities. Two important pieces of

information that are relevant in clinical care and translational research are knowledge of tissue

of origin, or CUP, and subtype of the cancer. Identifying tissue of origin of CUPs and molecular

subtype is critical for personalized medicine, where the treatment is tailored to the molecular

profile of individual tumor [66]. Because of the lack of primary site information, CUP patients

receive palliative chemotherapy that lacks the precision of modern targeted cancer medicine

and results in no clear benefit in survival [67, 68]. Various approved targeted therapies for

cancer by the Food and Drug Administration (FDA) include signal transduction inhibitors, gene

expression modulators, apoptosis inducers, angiogenesis inhibitors, immunotherapies. The

targeted therapies have been approved for the treatment of over 28 types of cancer. As CUP

may retain molecular signatures of its primary site, CUP patients with primary or metastatic

tumor might benefit from established therapeutic regimens appropriate for cancers of that

tissue; therefore, identifying the primary site is key to choosing effective therapeutic options.

Another major challenge in clinical cancer research is accurately classifying cancers into

appropriate homogeneous subtypes to improve prognosis and treatment [5]. Analyses of The

Cancer Genome Atlas (TCGA) and the International Cancer Genome Consortium (ICGC) have

established that a cancer at any primary site can be further classified into molecular subtypes

with potentially distinct clinical outcome and therapeutic options [21-28, 62]. For example, the

EBV-positive subtype of gastric cancer is associated with overexpression of JAK2, PD-L1 and

PD-L2 genes, suggesting that PD-L1/2 antagonists and JAK2 inhibitors are potential therapeutic

options for these tumors [64]. Similar clinical and therapeutic relevance of subtype information

has been demonstrated in multiple cancers [21, 23-25, 27, 28]. Increasingly, tumor molecular

subtype is being considered as an eligibility criterion for the entry into clinical trials [29].

However, for many cancers, the molecular subtype information is not available for use in clinical

practice because of the difficulty in identifying the subtype of a given tumor [30, 31]. Besides

being an important factor in clinical decision making, the knowledge of the molecular subtype

will be helpful in translational research. For example, cancer avatar trials use PDX models to

test panels of drugs to determine the best regime for personalized human therapy. Knowledge

of subtype of the tumor may narrow down the choice of treatment regimens to test which

increases efficiency of the cancer avatar trials.

We developed classifiers of high sensitivity and specificity for prediction of tissue of origin of 33

cancers and molecular subtyping of 11 cancers using gene expression data from the TCGA. We

show, for the first time to our knowledge, our pan-cancer classifiers can predict multiple cancers'

primary of origin from metastatic samples. Compared to the other classifiers based on somatic

mutations [69], our classifier is not limited to only cancer types with high mutation burden and

has much greater potential for clinical diagnosis and therapeutic design. Further, the classifiers

trained on the TCGA RNA-seq data generalize to different cohorts of primary and metastatic

samples profiled using microarray sequencing technologies. Such external validation qualifies

our classifiers to be robust across technology platforms, batches and sample processing

protocols. A combination of primary tissue of origin and subtype classification from metastases

will serve as important tools for clinicians in effectively treating CUPs.

The classification tools discriminated all cancers from each other well, except among the gastro-

intestinal cancers. However, the classification by cancer group could be achieved with very high

sensitivity and specificity. Different cancers among gastrointestinal cancers are less

distinguishable due to their anatomic proximity and molecular similarity, as shown in Figure 3.

To circumvent this problem, in our future work, we will adopt a hierarchical classification of

tumors: (1) granular classification by cancer group (2) finer classification by cancer type in each

group. In addition, we can include features from copy number, mutation and methylation data to

augment our feature set for both accuracy as well as robustness for technology and batch

variations. Molecular subtyping can also benefit from adding features from heterogeneous data

as several subtypes were identified to exhibit genomic features that span whole spectrum of

omics data. For example, the CIN subtype in gastric cancer is known to exhibit large structural

variations which may not be captured accurately by expression data. Thus, our future work will

encompass comprehensive multi-omic data to identify tissue of origin and molecular subtyping.

Though the overall performance of the classifiers designed using DLDA, SVM and KNN is not

as good as Random Forest on this data, their performance is on par with or better than Random

Forest based classifiers on certain tumor type and subtype classification. However, the

classification predicted by Random Forest classifiers is easier to interpret.

As we continue to enhance the classifier, we do recognize that the clinical utility of the classifier

is dependent on its ability to classify FFPE (formalin-fixed paraffin-embedded) samples, which is

the standard specimen type used for molecular profiling of cancers in clinical diagnostics in

addition to fresh-frozen samples. We have previously shown that the classifiers designed can

work with FFPE samples if they are profiled using nanoString arrays [70]. Therefore, it is

feasible to generalize our classifiers to work on FFPE samples for clinical applications. In

summary, we have demonstrated the utility of gene expression profiles to solve the important

clinical challenge of identifying the primary site of origin and the molecular subtype of cancers

based on machine learning algorithms. These predictors will be made available as open source

software, freely available for academic non-commercial use.

In an effort make these tools available to as wide an audience as possible, we offer our models

and results in two publicly available forms: a web-based portal and a software package which

can be used to apply these tools to other datasets and to reproduce the results presented

here. The web-based portal provides interactive visualizations showing the expression profiles

of the TCGA cancer samples and the classification results of our classifiers. These

visualizations allow for the exploration of relationships between cancer types in the context of

pan-cancer expression profiles.

SUPPORTING INFORMATION

Table S1. Contingency table and all performance metrics – random forest

Table S2. Contingency table and all performance metrics – SVM

Table S3. Contingency table and all performance metrics – kNN

Table S4. Contingency table and all performance metrics – DLDA

Table S5. Performance metrics for external datasets – random forest

ACKNOWLEDGEMENTS

Research reported in this publication was partially supported by the National Cancer Institute of

the National Institutes of Health under Award Number P30CA034196. The content is solely the

responsibility of the authors and does not necessarily represent the official views of the National

Institutes of Health. This study makes use of data generated by the Molecular Taxonomy of

Breast Cancer International Consortium. Funding for the project was provided by Cancer

Research UK and the British Columbia Cancer Agency Branch.

AUTHOR CONTRIBUTIONS

R.K.M.K. and J.G. designed research. S.L., R.K.M.K. and J.G. guided the project. C.A.P. and

J.G. performed data acquisition. W.F.F. and J.G. wrote software and performed analysis. H.R.

provided input on clinical oncology and contributed to the interpretation of the results. S.N.

provided computational support. W.F.F., S.N., C.A.P., H.R., S.L., R.K.M.K., J.G. wrote the

manuscript.

FIGURE LEGENDS

Figure 1. Platform-independent learning and validation from TCGA transcriptomes.

(A) Schematic showing the learning procedure used to train machine learning models from

labeled TCGA transcriptomes spanning 33 cancer types and 11 molecular subtypes. Models

were trained and evaluated using k-fold cross validation on normalized and standard scaled

expression profiles. For each fold, N features were selected from each class (see Methods) and

pooled, which were used to train the classification model. Using this schema, we constructed

one primary type and eleven molecular subtype classifiers for each type of model: random

forest (RF), support vector machine (SVM), k-nearest neighbor classifier (kNN), and diagonal

linear discriminant analysis (DLDA). (B) Classification performance was evaluated via the

cross-validation on the learning set and external validation utilizing five datasets; using two of

these datasets, we challenged the classifier to infer primary tumor types from transcriptomes of

metastatic or passaged patient-derived xenograft samples.

Figure 2. Precise classification of tumors by primary type and organ system.

(A-B) Random forest classification of primary cancer types and grouped by pan-organ system.

Text in contingency table cell $c_{j,i}$ shows tumor of class i classified as class j. Grayscale shading

of table cells is proportional to the number of samples of each tumor type, represented as bars

above the table. Color shading along the main diagonal shows pan-organ groups. Positive

predictive value (precision) for each prediction class are shown to the right of the table. (C-D)

Sensitivity and specificity for each classification in (A) and (B), respectively.

Figure 3. Unsupervised embedding of expression profiles reveals relationships among

primary cancer types.

Expression profiles from all samples were embedded into two dimensions using uniform

manifold approximation and projection (UMAP) [58] and colored by primary cancer type. For

each cancer, labels are placed near the centroid of the expression profile in the UMAP latent

space. Anatomical and histological relationships are emergent and add context to the most

common misclassifications in Figure 2. The following groups of cancers are highlighted with

green, blue, and purple ellipses, respectively: COAD, READ, STAD; BLCA, CESC, ESCA,

HNSC, LUSC; OV, SARC, UCEC, UCS.

Figure 4. External validation of primary type classifier using microarray data.

1,552 microarray expression profiles (GSE2109) from 9 cancer types or related cancer types

(LUSC/LUAD and KIRC/KIRP/KICH) were classified to further validate the primary type

classifier (A). Markers in (A) are scaled by the number of samples in each class. Classification

was evaluated by primary cancer type (B) and pan-organ group (C). (D-E) Sensitivity and

specificity for each classification in (B) and (C), respectively.

Figure 5. Classifier infers primary cancer of origin from metastatic tumor samples.

(A-B) 88 expression profiles of metastatic tumors (GSE18549) from primary cancers of origin

spanning 6 organs were classified by primary cancer type and primary pan-organ group. (C-D)

Sensitivity and specificity for each classification in (A) and (B), respectively. (E) 30/88 (34%) of

samples are liver or lung metastases from the colon. (F) The majority of misclassifications of

primary type are within pan-organ system; of the remainder, 7 misclassifications identify the

metastatic tumor whereas 9 are true misclassifications.

Figure 6. Classifier infers primary cancer of origin from passaged patient-derived

xenografts.

(A) 295 expression profiles of resected passaged patient-derived xenograft (PDX) tumors from

primary cancers spanning 11 organs were classified by primary cancer type (samples generated

at the Jackson Laboratory, available via MTB [37]). PDX tumor samples were taken for

sequencing from the second generation of mice. Classification of primary cancer identification

was evaluated by primary cancer type (B) and pan-organ group (C). (D-E) Sensitivity and

specificity for each classification in (B) and (C), respectively.

Figure 7. Cross- and external validation of molecular subtype classifiers.

A classifier of molecular subtypes was constructed for each of 11 primary cancer types,

spanning 38 molecular subtypes. (A) Per-class positive predictive value, (B) specificity, and (C)

sensitivity of molecular subtype classifications evaluated through cross-validation (Figure 1).

(D) Number of training samples for each molecular subtype. To further validate these subtype

classifiers, breast (E) and ovarian (F) subtype classifiers were used to predict the respective

molecular subtypes in two external datasets (GSE9899 and EGAS00000000083, respectively).

(G-H) Sensitivity and specificity for each classification in (E) and (F), respectively.

REFERENCES

1.	Pavlidis N, Khaled H, Gaafar R. A mini review on cancer of unknown primary site: A clinical puzzle for the oncologists. J Adv Res. 2015;6(3):375-82. Epub 2015/08/11. doi: 10.1016/j.jare.2014.11.007. PubMed PMID: 26257935; PubMed Central PMCID: PMC4522587.

2.	Moran S, Martinez-Cardus A, Boussios S, Esteller M. Precision medicine based on epigenomics: the paradigm of carcinoma of unknown primary. Nat Rev Clin Oncol. 2017;14(11):682-94. Epub 2017/07/05. doi: 10.1038/nrclinonc.2017.97. PubMed PMID: 28675165.

3.	Hyphantis T, Papadimitriou I, Petrakis D, Fountzilas G, Repana D, Assimakopoulos K, et al. Psychiatric manifestations, personality traits and health-related quality of life in cancer of unknown primary site. Psychooncology. 2013;22(9):2009-15. Epub 2013/01/30. doi: 10.1002/pon.3244. PubMed PMID: 23359412.

4.	Hainsworth JD, Schnabel CA, Erlander MG, Haines DW, 3rd, Greco FA. A retrospective study of treatment outcomes in patients with carcinoma of unknown primary site and a colorectal cancer molecular profile. Clin Colorectal Cancer. 2012;11(2):112-8. Epub 2011/10/18. doi: 10.1016/j.clcc.2011.08.001. PubMed PMID: 22000811.

5.	Varadhachary GR, Talantov D, Raber MN, Meng C, Hess KR, Jatkoe T, et al. Molecular profiling of carcinoma of unknown primary and correlation with clinical evaluation. J Clin Oncol. 2008;26(27):4442-8. Epub 2008/09/20. doi: 10.1200/JCO.2007.14.4378. PubMed PMID: 18802157.

6.	Yoon HH, Foster NR, Meyers JP, Steen PD, Visscher DW, Pillai R, et al. Gene expression profiling identifies responsive patients with cancer of unknown primary treated with carboplatin, paclitaxel, and everolimus: NCCTG N0871 (alliance). Ann Oncol. 2016;27(2):339-44. Epub 2015/11/19. doi: 10.1093/annonc/mdv543. PubMed PMID: 26578722; PubMed Central PMCID: PMC4907341.

7.	Varadhachary GR, Spector Y, Abbruzzese JL, Rosenwald S, Wang H, Aharonov R, et al. Prospective gene signature study using microRNA to identify the tissue of origin in patients with carcinoma of unknown primary. Clin Cancer Res. 2011;17(12):4063-70. Epub 2011/05/03. doi: 10.1158/1078-0432.CCR-10-2599. PubMed PMID: 21531815.

8.	Hainsworth JD, Rubin MS, Spigel DR, Boccia RV, Raby S, Quinn R, et al. Molecular gene expression profiling to predict the tissue of origin and direct site-specific therapy in patients with carcinoma of unknown primary site: a prospective trial of the Sarah Cannon research institute. J Clin Oncol. 2013;31(2):217-23. Epub 2012/10/04. doi: 10.1200/JCO.2012.43.3755. PubMed PMID: 23032625.

9.	Varadhachary GR, Karanth S, Qiao W, Carlson HR, Raber MN, Hainsworth JD, et al. Carcinoma of unknown primary with gastrointestinal profile: immunohistochemistry and survival data for this favorable subset. Int J Clin Oncol. 2014;19(3):479-84. Epub 2013/07/03. doi: 10.1007/s10147-013-0583-0. PubMed PMID: 23813044.

10.	Green AC. Cancer of unknown primary: does the key lie in molecular diagnostics? Cytopathology. 2015;26(1):61-3. Epub 2015/02/17. doi: 10.1111/cyt.12235. PubMed PMID: 25683360.

11.	Kandalaft PL, Gown AM. Practical Applications in Immunohistochemistry: Carcinomas of Unknown Primary Site. Arch Pathol Lab Med. 2016;140(6):508-23. Epub 2015/10/13. doi: 10.5858/arpa.2015-0173-CP. PubMed PMID: 26457625.

12.	Conner JR, Hornick JL. Metastatic carcinoma of unknown primary: diagnostic approach using immunohistochemistry. Adv Anat Pathol. 2015;22(3):149-67. Epub 2015/04/07. doi: 10.1097/PAP.0000000000000069. PubMed PMID: 25844674.

13.	Rubin BP, Skarin AT, Pisick E, Rizk M, Salgia R. Use of cytokeratins 7 and 20 in determining the origin of metastatic carcinoma of unknown primary, with special emphasis on lung cancer. Eur J Cancer Prev. 2001;10(1):77-82. Epub 2001/03/27. PubMed PMID: 11263595.

14.   Gunia S, Koch S, May M. Is CDX2 immunostaining useful for delineating anorectal from
penile/vulvar squamous cancer in the setting of squamous cell carcinoma with clinically unknown
primary site presenting with histologically confirmed inguinal lymph node metastasis? J Clin
Pathol. 2013;66(2):109-12. Epub 2012/10/30. doi: 10.1136/jclinpath-2012-201138. PubMed
PMID: 23105122.
15.   Tothill RW, Kowalczyk A, Rischin D, Bousioutas A, Haviv I, van Laar RK, et al. An
expression-based site of origin diagnostic method designed for clinical application to cancer of
unknown origin. Cancer Res. 2005;65(10):4031-40. Epub 2005/05/19. doi: 10.1158/0008-
5472.CAN-04-3617. PubMed PMID: 15899792.
16.   Horlings HM, van Laar RK, Kerst JM, Helgason HH, Wesseling J, van der Hoeven JJ, et
al. Gene expression profiling to identify the histogenetic origin of metastatic adenocarcinomas of
unknown primary. J Clin Oncol. 2008;26(27):4435-41. Epub 2008/09/20. doi:
10.1200/JCO.2007.14.6969. PubMed PMID: 18802156.
17.   Varadhachary GR, Raber MN, Matamoros A, Abbruzzese JL. Carcinoma of unknown
primary with a colon-cancer profile-changing paradigm and emerging definitions. Lancet Oncol.
2008;9(6):596-9. Epub 2008/05/31. doi: 10.1016/S1470-2045(08)70151-7. PubMed PMID:
18510991.
18.   van Laar RK, Ma XJ, de Jong D, Wehkamp D, Floore AN, Warmoes MO, et al.
Implementation of a novel microarray-based diagnostic test for cancer of unknown primary. Int J
Cancer. 2009;125(6):1390-7. Epub 2009/06/19. doi: 10.1002/ijc.24504. PubMed PMID:
19536816.
19.   Handorf CR, Kulkarni A, Grenert JP, Weiss LM, Rogers WM, Kim OS, et al. A multicenter
study directly comparing the diagnostic accuracy of gene expression profiling and
immunohistochemistry for primary site identification in metastatic tumors. Am J Surg Pathol.
2013;37(7):1067-75. Epub 2013/05/08. doi: 10.1097/PAS.0b013e31828309c4. PubMed PMID:
23648464; PubMed Central PMCID: PMC5266589.
20.   Greco FA, Lennington WJ, Spigel DR, Hainsworth JD. Molecular profiling diagnosis in
unknown primary cancer: accuracy and ability to complement standard pathology. J Natl Cancer
Inst. 2013;105(11):782-90. Epub 2013/05/04. doi: 10.1093/jnci/djt099. PubMed PMID: 23641043.
21.   Cancer Genome Atlas N. Comprehensive molecular portraits of human breast tumours.
Nature. 2012;490(7418):61-70. Epub 2012/09/25. doi: 10.1038/nature11412. PubMed PMID:
23000897; PubMed Central PMCID: PMC3465532.
22.   Cancer Genome Atlas N. Comprehensive genomic characterization of head and neck
squamous cell carcinomas. Nature. 2015;517(7536):576-82. Epub 2015/01/30. doi:
10.1038/nature14129. PubMed PMID: 25631445; PubMed Central PMCID: PMC4311405.
23.   Cancer Genome Atlas N. Comprehensive molecular characterization of human colon and
rectal cancer. Nature. 2012;487(7407):330-7. Epub 2012/07/20. doi: 10.1038/nature11252.
PubMed PMID: 22810696; PubMed Central PMCID: PMC3401966.
24.   Cancer Genome Atlas Research N. Comprehensive genomic characterization defines
human glioblastoma genes and core pathways. Nature. 2008;455(7216):1061-8. Epub
2008/09/06. doi: 10.1038/nature07385. PubMed PMID: 18772890; PubMed Central PMCID:
PMC2671642.
25.   Cancer Genome Atlas Research N. Integrated genomic analyses of ovarian carcinoma.
Nature. 2011;474(7353):609-15. Epub 2011/07/02. doi: 10.1038/nature10166. PubMed PMID:
21720365; PubMed Central PMCID: PMC3163504.
26.   Cancer Genome Atlas Research N. Comprehensive genomic characterization of
squamous cell lung cancers. Nature. 2012;489(7417):519-25. Epub 2012/09/11. doi:
10.1038/nature11404. PubMed PMID: 22960745; PubMed Central PMCID: PMC3466113.
27.   Cancer Genome Atlas Research N. Comprehensive molecular profiling of lung
adenocarcinoma. Nature. 2014;511(7511):543-50. Epub 2014/08/01. doi: 10.1038/nature13385.
PubMed PMID: 25079552; PubMed Central PMCID: PMC4231481.

28.   Cancer Genome Atlas Research N. The Molecular Taxonomy of Primary Prostate Cancer.
Cell. 2015;163(4):1011-25. Epub 2015/11/07. doi: 10.1016/j.cell.2015.10.025. PubMed PMID:
26544944; PubMed Central PMCID: PMC4695400.
29.   Kommoss S, Winterhoff B, Oberg AL, Konecny GE, Wang C, Riska SM, et al.
Bevacizumab May Differentially Improve Ovarian Cancer Outcome in Patients with Proliferative
and Mesenchymal Molecular Subtypes. Clin Cancer Res. 2017;23(14):3794-801. Epub
2017/02/06. doi: 10.1158/1078-0432.CCR-16-2196. PubMed PMID: 28159814; PubMed Central
PMCID: PMC5661884.
30.   Prat A, Fan C, Fernandez A, Hoadley KA, Martinello R, Vidal M, et al. Response and
survival of breast cancer intrinsic subtypes following multi-agent neoadjuvant chemotherapy.
BMC Med. 2015;13:303. Epub 2015/12/20. doi: 10.1186/s12916-015-0540-z. PubMed PMID:
26684470; PubMed Central PMCID: PMC4683815.
31.   Prat A, Pineda E, Adamo B, Galvan P, Fernandez A, Gaba L, et al. Clinical implications
of the intrinsic molecular subtypes of breast cancer. Breast. 2015;24 Suppl 2:S26-35. Epub
2015/08/09. doi: 10.1016/j.breast.2015.07.008. PubMed PMID: 26253814.
32.   Li B, Dewey CN. RSEM: accurate transcript quantification from RNA-Seq data with or
without a reference genome. BMC Bioinformatics. 2011;12(1). doi: 10.1186/1471-2105-12-323.
33.   Center BITGDA. Analysis-ready standardized TCGA data from Broad GDAC Firehose
2016_01_28 run. Broad Institute of MIT and Harvard: Broad Institute of MIT and Harvard; 2016.
34.   Huber W, Carey VJ, Gentleman R, Anders S, Carlson M, Carvalho BS, et al. Orchestrating
high-throughput genomic analysis with Bioconductor. Nat Methods. 2015;12(2):115-21. Epub
2015/01/31. doi: 10.1038/nmeth.3252. PubMed PMID: 25633503; PubMed Central PMCID:
PMC4509590.
35.   Singh R, Maganti RJ, Jabba SV, Wang M, Deng G, Heath JD, et al. Microarray-based
comparison of three amplification methods for nanogram amounts of total RNA. American Journal
of Physiology-Cell Physiology. 2005;288(5):C1179-C89. doi: 10.1152/ajpcell.00258.2004.
36.   Hsu SD, Kim MK, Foye A, Silvestri A, Lyerly HK, Morse M, et al. Use of gene expression
signatures to identify origin of primary and therapeutic strategies for patients with advanced solid
tumors.   Journal   of   Clinical   Oncology.   2010;28(15_suppl):10504-.   doi:
10.1200/jco.2010.28.15_suppl.10504.
37.   Krupke DM, Begley DA, Sundberg JP, Bult CJ, Eppig JT. The Mouse Tumor Biology
database. Nat Rev Cancer. 2008;8(6):459-65. Epub 2008/04/25. doi: 10.1038/nrc2390. PubMed
PMID: 18432250; PubMed Central PMCID: PMC2574871.
38.   Curtis C, Shah SP, Chin SF, Turashvili G, Rueda OM, Dunning MJ, et al. The genomic
and transcriptomic architecture of 2,000 breast tumours reveals novel subgroups. Nature.
2012;486(7403):346-52. Epub 2012/04/24. doi: 10.1038/nature10983. PubMed PMID: 22522925;
PubMed Central PMCID: PMC3440846.
39.   Cerami E, Gao J, Dogrusoz U, Gross BE, Sumer SO, Aksoy BA, et al. The cBio cancer
genomics portal: an open platform for exploring multidimensional cancer genomics data. Cancer
Discov. 2012;2(5):401-4. Epub 2012/05/17. doi: 10.1158/2159-8290.CD-12-0095. PubMed PMID:
22588877; PubMed Central PMCID: PMC3956037.
40.   Gao J, Aksoy BA, Dogrusoz U, Dresdner G, Gross B, Sumer SO, et al. Integrative analysis
of complex cancer genomics and clinical profiles using the cBioPortal. Sci Signal.
2013;6(269):pl1. Epub 2013/04/04. doi: 10.1126/scisignal.2004088. PubMed PMID: 23550210;
PubMed Central PMCID: PMC4160307.
41.   Ceccarelli M, Barthel FP, Malta TM, Sabedot TS, Salama SR, Murray BA, et al. Molecular
Profiling Reveals Biologically Discrete Subsets and Pathways of Progression in Diffuse Glioma.
Cell. 2016;164(3):550-63. Epub 2016/01/30. doi: 10.1016/j.cell.2015.12.028. PubMed PMID:
26824661; PubMed Central PMCID: PMC4754110.

42. Getz G, Gabriel SB, Cibulskis K, Lander E, Sivachenko A, Sougnez C, et al. Integrated
genomic characterization of endometrial carcinoma. Nature. 2013;497(7447):67-73. doi:
10.1038/nature12113.
43. Akbani R, Akdemir Kadir C, Aksoy BA, Albert M, Ally A, Amin Samirkumar B, et al.
Genomic Classification of Cutaneous Melanoma. Cell. 2015;161(7):1681-96. doi:
10.1016/j.cell.2015.05.044.
44. Comprehensive Molecular Characterization of Papillary Renal-Cell Carcinoma. New
England Journal of Medicine. 2016;374(2):135-45. doi: 10.1056/NEJMoa1505917.
45. Comprehensive molecular characterization of clear cell renal cell carcinoma. Nature.
2013;499(7456):43-9. doi: 10.1038/nature12222.
46. Hoadley KA, Yau C, Hinoue T, Wolf DM, Lazar AJ, Drill E, et al. Cell-of-Origin Patterns
Dominate the Molecular Classification of 10,000 Tumors from 33 Types of Cancer. Cell.
2018;173(2):291-304 e6. Epub 2018/04/07. doi: 10.1016/j.cell.2018.03.022. PubMed PMID:
29625048; PubMed Central PMCID: PMC5957518.
47. Dudoit S, Fridlyand J, Speed TP. Comparison of Discrimination Methods for the
Classification of Tumors Using Gene Expression Data. Journal of the American Statistical
Association. 2002;97(457):77-87. doi: 10.1198/016214502753479248.
48. Tabchy A, Valero V, Vidaurre T, Lluch A, Gomez H, Martin M, et al. Evaluation of a 30-
gene paclitaxel, fluorouracil, doxorubicin, and cyclophosphamide chemotherapy response
predictor in a multicenter randomized trial in breast cancer. Clin Cancer Res. 2010;16(21):5351-
61. Epub 2010/09/11. doi: 10.1158/1078-0432.CCR-10-1265. PubMed PMID: 20829329;
PubMed Central PMCID: PMC4181852.
49. Kurokawa C, Iankov ID, Anderson SK, Aderca I, Leontovich AA, Maurer MJ, et al.
Constitutive Interferon Pathway Activation in Tumors as an Efficacy Determinant Following
Oncolytic Virotherapy. J Natl Cancer Inst. 2018. Epub 2018/05/23. doi: 10.1093/jnci/djy033.
PubMed PMID: 29788332.
50. Khondoker M, Dobson R, Skirrow C, Simmons A, Stahl D. A comparison of machine
learning methods for classification using simulation with multiple real data examples from mental
health studies. Stat Methods Med Res. 2016;25(5):1804-23. Epub 2013/09/21. doi:
10.1177/0962280213502437. PubMed PMID: 24047600; PubMed Central PMCID:
PMC5081132.
51. Li Y, Kang K, Krahn JM, Croutwater N, Lee K, Umbach DM, et al. A comprehensive
genomic pan-cancer classification using The Cancer Genome Atlas gene expression data. BMC
Genomics. 2017;18(1):508. Epub 2017/07/05. doi: 10.1186/s12864-017-3906-0. PubMed PMID:
28673244; PubMed Central PMCID: PMC5496318.
52. Li Z, Mao Y, Li H, Yu G, Wan H, Li B. Differentiating brain metastases from different
pathological types of lung cancers using texture analysis of T1 postcontrast MR. Magn Reson
Med. 2016;76(5):1410-9. Epub 2015/12/02. doi: 10.1002/mrm.26029. PubMed PMID: 26621795.
53. Venables WN, Ripley BD. Modern Applied Statistics with S2002.
54. Wang S, Cai Y. Identification of the functional alteration signatures across different cancer
types with support vector machine and feature analysis. Biochim Biophys Acta. 2018;1864(6 Pt
B):2218-27. Epub 2017/12/27. doi: 10.1016/j.bbadis.2017.12.026. PubMed PMID: 29277326.
55. Galvez JM, Castillo D, Herrera LJ, San Roman B, Valenzuela O, Ortuno FM, et al.
Multiclass classification for skin cancer profiling based on the integration of heterogeneous gene
expression series. PLoS One. 2018;13(5):e0196836. Epub 2018/05/12. doi:
10.1371/journal.pone.0196836. PubMed PMID: 29750795; PubMed Central PMCID:
PMC5947894.
56. Zhi J, Sun J, Wang Z, Ding W. Support vector machine classifier for prediction of the
metastasis of colorectal cancer. Int J Mol Med. 2018;41(3):1419-26. Epub 2018/01/13. doi:
10.3892/ijmm.2018.3359. PubMed PMID: 29328363; PubMed Central PMCID: PMC5819940.

57.   Breiman L. Consistency for a simple model of random forests. Univ. California, Berkeley,
CA: 2004   Contract No.: Technical Report 670.
58.   McInnes L, Healy J. UMAP: Uniform Manifold Approximation and Projection for Dimension
Reduction. ArXiv e-prints [Internet]. 2018.
59.   van der Maaten L. Accelerating t-SNE using Tree-Based Algorithms. Journal of Machine
Learning Research. 2014;15:3221-45.
60.   Ulyanov D. Multicore-TSNE. GitHub; 2016.
61.   Zhang Y. Epidemiology of esophageal cancer. World J Gastroenterol. 2013;19(34):5598-
606. Epub 2013/09/17. doi: 10.3748/wjg.v19.i34.5598. PubMed PMID: 24039351; PubMed
Central PMCID: PMC3769895.
62.   Cherniack AD, Shen H, Walter V, Stewart C, Murray BA, Bowlby R, et al. Integrated
Molecular Characterization of Uterine Carcinosarcoma. Cancer Cell. 2017;31(3):411-23. Epub
2017/03/16. doi: 10.1016/j.ccell.2017.02.010. PubMed PMID: 28292439; PubMed Central
PMCID: PMC5599133.
63.   Hidalgo M, Amant F, Biankin AV, Budinska E, Byrne AT, Caldas C, et al. Patient-derived
xenograft models: an emerging platform for translational cancer research. Cancer Discov.
2014;4(9):998-1013. Epub 2014/09/04. doi: 10.1158/2159-8290.CD-14-0001. PubMed PMID:
25185190; PubMed Central PMCID: PMC4167608.
64.   Bass AJ, Thorsson V, Shmulevich I, Reynolds SM, Miller M, Bernard B, et al.
Comprehensive molecular characterization of gastric adenocarcinoma. Nature.
2014;513(7517):202-9. doi: 10.1038/nature13480.
65.   Wu X, Li X, Fu Q, Cao Q, Chen X, Wang M, et al. AKR1B1 promotes basal-like breast
cancer progression by a positive feedback loop that activates the EMT program. The Journal of
Experimental Medicine. 2017;214(4):1065-79. doi: 10.1084/jem.20160903.
66.   Varadhachary GR, Abbruzzese JL, Lenzi R. Diagnostic strategies for unknown primary
cancer. Cancer. 2004;100(9):1776-85. Epub 2004/04/28. doi: 10.1002/cncr.20202. PubMed
PMID: 15112256.
67.   Fizazi K, Greco FA, Pavlidis N, Daugaard G, Oien K, Pentheroudakis G, et al. Cancers of
unknown primary site: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up.
Ann Oncol. 2015;26 Suppl 5:v133-8. Epub 2015/09/01. doi: 10.1093/annonc/mdv305. PubMed
PMID: 26314775.
68.   Lee J, Hahn S, Kim DW, Kim J, Kang SN, Rha SY, et al. Evaluation of survival benefits
by platinums and taxanes for an unfavourable subset of carcinoma of unknown primary: a
systematic review and meta-analysis. Br J Cancer. 2013;108(1):39-48. Epub 2012/11/24. doi:
10.1038/bjc.2012.516. PubMed PMID: 23175147; PubMed Central PMCID: PMC3553519.
69.   Chen Y, Sun J, Huang L-C, Xu H, Zhao Z. Classification of Cancer Primary Sites Using
Machine Learning and Somatic Mutations. BioMed Research International. 2015;2015:1-9. doi:
10.1155/2015/491502.
70.   Leong HS, Galletta L, Etemadmoghadam D, George J, Australian Ovarian Cancer S,
Kobel M, et al. Efficient molecular subtype classification of high-grade serous ovarian cancer. J
Pathol. 2015;236(3):272-7. Epub 2015/03/27. doi: 10.1002/path.4536. PubMed PMID: 25810134.