# Co-outbreak of ST37 and a novel ST3006 *Klebsiella pneumoniae* from multi-site infection in a neonatal intensive care unit, Fuzhou, China

Dongjie Chen[1,2], Xinlan Hu[2], Falin Chen[2], Hongru Li[3], Daxuan Wang[3], Xiaoqin Li[3], Changsheng Wu[2], Ning Li[2], Shaolian Wu[2], Zhen Li[2], Liqing Chen[2], Yusheng Chen[1,3]

1 Shengli Clinical Medical College of Fujian Medical University, Fuzhou, 350001, China

2 Clinical Microbiology Laboratory, Fujian Provincial Hospital, Fuzhou 350001, China

3 Department of Respiratory Medicine and Critical Care Medicine, Fujian Provincial Hospital, Fuzhou 350001, China

Corresponding author:

Yusheng Chen

No.134 East Street, Fuzhou 350001, China

+86 13600895421

chenyusheng8201@163.com

**Summary**

**Background:** The spread of carbapenem resistance among *Klebsiella pneumoniae* (*K.*

*pneumoniae*) is a major public health problem, particularly in neonatal intensive care

units (NICUs).

**Aim:** To describe the nosocomial co-outbreak of ST37 and a new sequence type

ST3006 *K. pneumoniae*, causing catheter-related bloodstream infections in NICU.

**Methods:** Fifteen strains of *K. pneumoniae* were isolated from seven neonates during

June 3–28, 2013, in a tertiary hospital neonatal intensive care unit in Fujian, China.

Antimicrobial susceptibility was determined by the Vitek 2 system and micro-broth

dilution method. Multi-locus sequence typing (MLST) and pulsed-field gel

electrophoresis (PFGE) were used to analyse the genetic relatedness of isolates.

Genome sequencing and gene function analyses were performed for investigating

pathogenicity and drug resistance and screening genomic islands.

**Findings:** Two *K. pneumoniae* clones were identified from seven neonates with

multi-site infection, one ST37 strain and one new sequence type ST3006.

Antimicrobial susceptibility testing revealed that the ST37 strain exhibited multi-drug

resistance and carbapenem resistance. The ST3006 strain was only resistant to

ampicillin-sulbactam, cefazolin, ceftriaxone. MLST and PFGE showed that 15 strains

were divided into three groups, with a high level of homology. Gene sequencing and

analysis indicated that KPN1343 harboured 12 resistance genes, 15 genomic islands

and 205 reduced virulence genes. KPN1344 harboured four resistance genes, 19

genomic islands and 209 reduced virulence genes.

**Conclusion:** Co-outbreak of *K. pneumoniae* involved two clones, ST36 and ST3006,

causing multi-site infection. Genome sequencing and analysis is an effective method

for studying bacterial resistance genes and their functions.

**Introduction**

Klebsiella pneumonia, particularly multidrug resistant *K. pneumoniae*, is a common and important pathogen in hospitals, posing a serious and urgent threat to public health (1, 2). Neonates in the neonatal intensive care units often have underlying conditions, such as prematurity, or the presence of indwelling catheters, a history of antibiotic treatment and parenteral nutrition, which are risk factors for infection (3). In addition, relaxed vigilance by doctors and nurses regarding nosocomial infection promotes nosocomial infection outbreaks, with considerable impact on neonatal treatment and prognosis, prolonged hospital stays, increased hospital costs and mortality.

Nosocomial outbreaks in the neonatal intensive care unit are frequently reported (4-7); however, it is rare to isolate the same clonal pathogen from different sites in the same patient. This study identified 15 *K. pneumoniae* strains isolated from sputum specimens, blood specimens and umbilical vein catheter tips in seven neonates that belonged to the two cloned strains ST37 and ST3006, which produced carbapenemase and extended-spectrum beta-lactamases, respectively.

**Methods**

**Patients and bacterial strains**

Patient characteristics were obtained from electronic medical records. Bacterial strains were isolated from seven neonates and stored in a refrigerator (SANYO, Japan) at -70°C. Strains were identified using the Vitek 2 system (BioMérieux, France). The ethics approval number is K2018-01-001, and we have obtained the written consents from the patients' parents.

**Multi-locus sequence typing**

Bacterial DNA was extracted using the Bacteria Genomic DNA Kit (CWBIO, China). Multi-locus sequence typing (MLST) for *K. pneumoniae* was performed following the methods described previously (8). The allelic profiles and sequence types (STs) were Determined using online databases (https://pubmlst.org/bigsdb?db=pubmlst_mlst_s

eqdef). The novel allele profiles were sent to klebsiellaMLST@pasteur.fr for

confirmation.

**Pulsed-field gel electrophoresis**

Pulsed-field gel electrophoresis (PFGE) of XbaI-digested genomic DNA samples of *K.*

*pneumoniae* was performed with a CHEF MAPPER XA apparatus (Bio-Rad, USA)

according to the literature (9). Electrophoresis was performed for 24 h at 14°C with

pulse time ranging from 5 to 35 s at 6 V/cm. PFGE profiles were analysed and

compared using the Gel Doc XR+ system, version 2.0 (Bio-Rad, USA).

**Antimicrobial susceptibility testing**

Antibiotic susceptibility testing was performed by the Vitek 2 system (BioMérieux,

France), including ampicillin sulbactam, cefazolin, ceftriaxone, cefotetan, ceftazidime,

cefepime, gentamicin, tobramycin, amikacin, levofloxacin, ciprofloxacin, aztreonam,

imipenem, ertapenem, piperacillin-tazobactam, trimethoprim-sulfamethoxazole. The

MICs of tigecycline and polymyxin B was determined by the micro-broth dilution

method. All antibiotics were interpreted according to the Clinical and Laboratory

Standards Institute (CLSI) document M100-S23 (10), except tigecycline and

polymyxin B. For tigecycline and polymyxin B, the European Committee on

Antimicrobial Susceptibility Testing (EUCAST) breakpoint (11) was used. *E. coli*

ATCC 25922 and *P. aeruginosa* ATCC27853 were used as quality control.

**Genome sequencing and gene analysis**

Whole-genome sequencing was performed on the Illumina HiSeq PE150 platform

(Novogene Bioinformatics Technology Co., Ltd., Beijing, China). The Island

Path-DIOMB program was used to predict the genomic islands. For these pathogenic

bacteria, we used the Pathogen–Host Interactions Database (PHI) and the

Comprehensive Antibiotic Resistance Database (CARD) to perform the pathogenicity

and drug resistance analyses.

**Results**

**Clinical characteristics of patients**

Patient clinical characteristics were determined from the retrospective study of the electronic medical records and are presented in Table I. Subjects included five premature infants and two infants with hyperbilirubinaemia. All of the five premature infants were also diagnosed as low birth weight infants (two were very low birth weight infants) and with neonatal respiratory distress syndrome (NRDS). Three infants were delivered by caesarean section, two infants were delivered by midwives, one infant was delivered by forceps and only one infant was a normal delivery. The average age of the five premature infants was approximately 30 weeks, with an average body weight of 1.37 kg. Patient 1 was treated with piperacillin/tazobactam and meropenem, while the remaining patients were treated with piperacillin/tazobactam. Patient 1, Patient 2, Patient 3, Patient 4, Patient 6 and Patient 7, had a lower than normal Apgar score (10-10-10). The clinical condition of Patient 4 (the second of the twins) was poor but was discharged upon request of the family, while the remaining patients were treated and discharged.

**Characteristics and Antibiotic Susceptibility of *K. pneumoniae***

Twenty-one *K. pneumoniae* strains were identified by the VIETK 2 system from different specimens, including sputum, blood, umbilical vein catheter tip. Only 15 *K. pneumoniae* strains were collected and frozen in a -70°C refrigerator. In Patient 2, Patient 3, Patient 4 and Patient 6, we simultaneously isolated two *K. pneumoniae strains* from blood and/or umbilical vein catheter tips and ST typing and drug resistance were consistent. (ST37, a novel ST typing ST3006: gapA 69, infB 19, mdh 90, pgi 20, phoE 125, rpoB 18, novel allele tonB 406). Antibiotic susceptibility testing showed that the ST37 strain was resistant to ampicillin-sulbactam, cefazolin, ceftriaxone, cefotetan, ceftazidime, cefepime, gentamicin, aztreonam, imipenem, ertapenem, piperacillin-tazobactam. The ST3006 strain was only resistant to ampicillin-sulbactam, cefazolin, ceftriaxone. (Table II, Table III). After investigation and nosocomial infection control are carried out by the infection department, no

further bacteria were isolated from the patients, except for a strain ST1224 from

Patient 3 on June 28, 2013.

**Antibiotic Resistance Genes**

The CARD was used to search for the names of resistance-related genes. KPN1343

was found to have 12 antibiotic resistant genes: beta-lactam resistance (OXA-33,

TEM-1, SHV-11), aminoglycoside resistance [AAC (6')-IId, AAC (3)-IIa, AAC

(6')-Ib-cr], phenicol resistance (catB3), rifamycin resistance (arr-3), sulfonamide

resistance (sul1), efflux pump complex or subunit conferring antibiotic resistance

(oqxB, oqxA, CRP), streptogramin resistance (catB3). KPN1344 was found to have

four antibiotic resistant genes: beta-lactam resistance (TEM-1, CTX-M-3), efflux

pump complex or subunit conferring antibiotic resistance (vgaC, CRP). We chose the

resistance gene with a best-identities rate≥0.99 (Table IV).

**PFGE**

PFGE homology analysis showed that 15 *K. pneumoniae* strains were divided into

three clusters: ClusterA:1344,1346,1348,1351,1352,1354,1355; ClusterB:1362 and

ClusterC:1342,1343,1345,1347,1349,1350, 1353. MLST typing also divided 15 *K.*

*pneumoniae* strains into three groups. (Fig. 1)

**Pathogen-Host Interactions and Genomic Islands**

Using the BLAST software, the amino acid sequences of KPN1343 and KPN1344

were compared using the PHI database. PHI Phenotype classification showed that the

number of reduced virulence genes matched the database mostly, 205 and 209,

respectively (Fig. 2). Using Island Path-DIOMB to predict genomic islands,

KPN1343 has 15 genomic islands and KPN1344 has 19 genomic islands, the length

and direction of the genes are shown in Fig. 3 (length of <15 kb only is shown).

**Discussion**

*K. pneumoniae* is an important nosocomial pathogen that can cause pneumonia,

urinary tract infection, digestive tract infection, bloodstream infection, liver abscess

and meningitis (12-17). Infection outbreaks have been frequently reported in neonatal

intensive care units. Extended spectrum beta-lactamase-producing and

carbapenemase-producing *K. pneumoniae* can cause large outbreaks with significant

morbidity and mortality effects (18). Underlying conditions, including premature

delivery, low birth weight and neonatal respiratory distress syndrome, are risk factors

for neonatal infection. Inadequate medical equipment, environmental disinfection and

hand hygiene as well as understaffing have been reported to be important factors for

infection outbreak (19, 20). In our study, infection control practitioners took samples

from incubators, air, the hands of doctors and nurses, objects and patient skin for

bacterial culture. Only one *K. pneumoniae* strain was isolated from the inner surface

of an incubator. However, this strain was not stored for further research.

Screening and monitoring of extended spectrum beta-lactamase-producing and

carbapenemase-producing *K. pneumoniae* in hospitals is crucial (21). However, the

current molecular biological techniques, such as PCR and fluorescence quantitative

PCR, have limitations and cannot be used extensively to screen for drug resistance

genes. Genome sequencing maybe an effective method for screening resistant genes,

although the technique entails high costs and is time-consuming, limiting it

application. Homology analysis methods, such as ERIC-PCR, PFGE and MLST are

complex and also time-consuming, such that timely detection of outbreaks is not

feasible (22-24). Newer technologies need to be developed for monitoring the rapid

outbreak of nosocomial infections.

Most previous reports on *K. pneumoniae* outbreaks involved single-site

infections, diagnosed from respiratory specimens (25, 26), blood specimens (27-29),

or urine specimens (30-32). This study showed that the pathogens were

simultaneously isolated from respiratory specimens, blood specimens and umbilical

vein catheter tips in the same patient. However, isolates of kpn1340, kpn1341,

kpn1356, kpn1357, kpn1358, kpn1359 were not collected and could not be further

analysed. Antibiotic susceptibility reports revealed that in all cases antibiotic

susceptibility was consistent with strain ST37 or ST3006 and we suspect that these

may belong to the same clones.

A number of limitations to our research should be noted. Specifically, we were

not aware of a nosocomial infection outbreak; therefore, some strains of *K. pneumoniae* were not collected. It is important to enhance awareness of infection outbreak and strengthen bacterial preservation. Furthermore, we should strengthen communication and cooperation with infection control practitioners to screen and block nosocomial transmission at an earlier stage. Hospitals should implement different strategies to combat outbreaks, such as strict hand-washing by staff, frequent equipment change and extensive cleaning of paediatric wards.

**Conclusions**

In summary, we reported two clones of *K. pneumoniae* ST37 and ST3006 outbreak in a neonatal intensive care unit. The clones were isolated from multi-location, sputum specimens, blood and umbilical vein catheter tips. Whole-genome sequencing would be a useful method for the detection of drug resistance genes and the analysis gene function. Active and effective infection control measures are indispensable for preventing and controlling nosocomial infection outbreaks.

**Acknowledgement**

We thank the team of curators of the Institut Pasteur MLST and whole-genome MLST databases for curating the data and making them publicly available at http://bigsdb.pasteur.fr

**Ethics approval and consent to participate**

The ethics approval number is K2018-01-001，and we have obtained the written consents from the patients' parents.

**Conflict of interest statement**

The authors declare that they have no conflict of interest.

**Funding sources**

This work was supported by the Startup Fund for scientific research, Fujian Medical University（grant no. 2016QH116), the Youth Scientific Research Project of Fujian Provincial Health and Family Planning Commission (grant no. 2017-1-9), the Scientific Innovation Project of Fujian Provincial Health and Family Planning Commission (grant no. 2016-CX-3), the National Major Science and Technology Project for the Control and Prevention of Major Infectious Diseases of China (grant no.2017ZX10103004).

**References**

[1]   Huang W, Wang G, Sebra R, Zhuge J, Yin C, Aguero-Rosenfeld ME, Schuetz AN, Dimitrova N, Fallon JT. Emergence and Evolution of Multidrug-Resistant Klebsiella pneumoniae with both blaKPC and blaCTX-M Integrated in the Chromosome. Antimicrob Agents Chemother. 2017. 61(7).

[2]   Navon-Venezia S, Kondratyeva K, Carattoli A. Klebsiella pneumoniae: a major worldwide source and shuttle for antibiotic resistance. FEMS Microbiol Rev. 2017. 41(3): 252-275.

[3]   Nour I, Eldegla HE, Nasef N, Shouman B, Abdel-Hady H, Shabaan AE. Risk factors and clinical outcomes for carbapenem-resistant Gram-negative late-onset sepsis in a neonatal intensive care unit. J Hosp Infect. 2017. 97(1): 52-58.

[4]   Ruiz E, Rojo-Bezares B, Sáenz Y, Olarte I, Esteban I, Rocha-Gracia R, Zarazaga M, Torres C. Outbreak caused by a multi-resistant Klebsiella pneumoniae strain of new sequence type ST341 carrying new genetic environments of aac(6')-Ib-cr and qnrS1 genes in a neonatal intensive care unit in Spain. Int J Med Microbiol. 2010. 300(7): 464-9.

[5]   Zhou J, Li G, Ma X, Yang Q, Yi J. Outbreak of colonization by carbapenemase-producing Klebsiella pneumoniae in a neonatal intensive care unit: Investigation, control measures and assessment. Am J Infect Control. 2015. 43(10): 1122-4.

[6]   Karaaslan A, Soysal A, Altinkanat GG, Kepenekli KE, Söyletir G, Bakir M. Molecular characterization and risk factors for carbapenem-resistant Gram-negative bacilli colonization in children: emergence of NDM-producing Acinetobacter baumannii in a newborn intensive care unit in Turkey. J Hosp Infect. 2016. 92(1): 67-72.

[7]   Nakamura K, Kaneko M, Abe Y, Yamamoto N, Mori H, Yoshida A, Ohashi K, Miura S, Yang

TT, Momoi N, Kanemitsu K. Outbreak of extended-spectrum β-lactamase-producing

Escherichia coli transmitted through breast milk sharing in a neonatal intensive care unit. J

Hosp Infect. 2016. 92(1): 42-6.

[8]  Diancourt L, Passet V, Verhoef J, Grimont PA, Brisse S. Multilocus sequence typing of

Klebsiella pneumoniae nosocomial isolates. J Clin Microbiol. 2005. 43(8): 4178-82.

[9]  Zhan L, Wang S, Guo Y, Jin Y, Duan J, Hao Z, Lv J, Qi X, Hu L, Chen L, Kreiswirth BN,

Zhang R, Pan J, Wang L, Yu F. Outbreak by Hypermucoviscous Klebsiella pneumoniaeST11

Isolates with Carbapenem Resistance in a Tertiary Hospital in China. Front Cell Infect

Microbiol. 2017. 7: 182.

[10] Mendes RE, Sader HS, Flamm RK, Jones RN. Activity of oritavancin tested against

uncommonly isolated Gram-positive pathogens responsible for documented infections in

hospitals worldwide. J Antimicrob Chemother. 2014. 69(6): 1579-81.

[11] Neuert S, Nair S, Day MR, Doumith M, Ashton PM, Mellor KC, Jenkins C, Hopkins KL,

Woodford N, de Pinna E, Godbole G, Dallman TJ. Prediction of Phenotypic Antimicrobial

Resistance Profiles from Whole Genome Sequences of Non-typhoidal Salmonella enterica.

Front Microbiol. 2018. 9: 592.

[12] Mills G, Dumigan A, Kidd T, Hobley L, Bengoechea JA. Identification and Characterization

of Two Klebsiella pneumoniae lpxL Lipid A Late Acyltransferases and Their Role in

Virulence. Infect Immun. 2017. 85(9).

[13] Wolfensberger A, Meier AH, Kuster SP, Mehra T, Meier MT, Sax H. Should International

Classification of Diseases codes be used to survey hospital-acquired pneumonia. J Hosp

Infect. 2018. 99(1): 81-84.

[14] Giannella M, Trecarichi EM, Giacobbe DR, De Rosa FG, Bassetti M, Bartoloni A, Bartoletti

M, Losito AR, Del BV, Corcione S, Tedeschi S, Raffaelli F, Saffioti C, Spanu T, Rossolini

GM, Marchese A, Ambretti S, Cauda R, Viscoli C, Lewis RE, Viale P, Tumbarello M. Effect

of combination therapy containing a high-dose carbapenem on mortality in patients with

carbapenem-resistant Klebsiella pneumoniae bloodstream infection. Int J Antimicrob Agents.

2018. 51(2): 244-248.

[15] Tan TY, Ong M, Cheng Y, LSY N. Hypermucoviscosity, rmpA, and aerobactin are associated

with community-acquired Klebsiella pneumoniae bacteremic isolates causing liver abscess in

Singapore. J Microbiol Immunol Infect. 2017.

[16] Ku YH, Chuang YC, Chen CC, Lee MF, Yang YC, Tang HJ, Yu WL. Klebsiella pneumoniae

Isolates from Meningitis: Epidemiology, Virulence and Antibiotic Resistance. Sci Rep. 2017.

7(1): 6634.

[17] De Rosa FG, Corcione S, Cavallo R, Di PG, Bassetti M. Critical issues for Klebsiella

pneumoniae KPC-carbapenemase producing K. pneumoniae infections: a critical agenda.

Future Microbiol. 2015. 10(2): 283-94.

[18] Arena F, Giani T, Becucci E, Conte V, Zanelli G, D'Andrea MM, Buonocore G, Bagnoli F,

Zanchi A, Montagnani F, Rossolini GM. Large oligoclonal outbreak due to Klebsiella

pneumoniae ST14 and ST26 producing the FOX-7 AmpC β-lactamase in a neonatal intensive

care unit. J Clin Microbiol. 2013. 51(12): 4067-72.

[19] Stapleton PJ, Murphy M, McCallion N, Brennan M, Cunney R, Drew RJ. Outbreaks of

extended spectrum beta-lactamase-producing Enterobacteriaceae in neonatal intensive care

units: a systematic review. Arch Dis Child Fetal Neonatal Ed. 2016. 101(1): F72-8.

[20] Jørgensen SB, Bojer MS, Boll EJ, Martin Y, Helmersen K, Skogstad M, Struve C.

Heat-resistant, extended-spectrum β-lactamase-producing Klebsiella pneumoniae in

endoscope-mediated outbreak. J Hosp Infect. 2016. 93(1): 57-62.

[21] Yun HC, Tully CC, Mende K, Castillo M, Murray CK. A single-center, six-year evaluation of

the role of pulsed-field gel electrophoresis in suspected burn center outbreaks. Burns. 2016.

42(6): 1323-30.

[22] Ostria-Hernandez ML, Juárez-de lRKC, Arzate-Barbosa P, Lara-Hernández A, Sakai F, Ibarra

JA, Castro-Escarpulli G, Vidal JE. Nosocomial, Multidrug-Resistant Klebsiella pneumoniae

Strains Isolated from Mexico City Produce Robust Biofilms on Abiotic Surfaces but Not on

Human Lung Cells. Microb Drug Resist. 2018. 24(4): 422-433.

[23] Diago-Navarro E, Chen L, Passet V, Burack S, Ulacia-Hernando A, Kodiyanplakkal RP, Levi

MH, Brisse S, Kreiswirth BN, Fries BC. Carbapenem-resistant Klebsiella pneumoniae

exhibit variability in capsular polysaccharide and capsule associated virulence traits. J Infect

Dis. 2014. 210(5): 803-13.

[24] Bosch T, SPM L, MHA H, Wever PC, Schneeberger PM, NHM R, ACAP L, JAJW K,

Schoffelen A, Notermans D, Witteveen S, Bathoorn E, Schouls LM. An outbreak of NDM-1

producing Klebsiella pneumoniae in a Dutch hospital with interspecies transfer of the resistance plasmid and unexpected occurrence in unrelated healthcare centers. J Clin Microbiol. 2017.

[25] Yu F, Ying Q, Chen C, Li T, Ding B, Liu Y, Lu Y, Qin Z, Parsons C, Salgado C, Qu D, Pan J, Wang L. Outbreak of pulmonary infection caused by Klebsiella pneumoniae isolates harbouring blaIMP-4 and blaDHA-1 in a neonatal intensive care unit in China. J Med Microbiol. 2012. 61(Pt 7): 984-9.

[26] Johnson J, Quach C. Outbreaks in the neonatal ICU: a review of the literature. Curr Opin Infect Dis. 2017. 30(4): 395-403.

[27] Richards C, Alonso-Echanove J, Caicedo Y, Jarvis WR. Klebsiella pneumoniae bloodstream infections among neonates in a high-risk nursery in Cali, Colombia. Infect Control Hosp Epidemiol. 2004. 25(3): 221-5.

[28] Matsumura Y, Tanaka M, Yamamoto M, Nagao M, Machida K, Ito Y, Takakura S, Ogawa K, Yoshizawa A, Fujimoto Y, Okamoto S, Uemoto S, Ichiyama S. High prevalence of carbapenem resistance among plasmid-mediated AmpC β-lactamase-producing Klebsiella pneumoniae during outbreaks in liver transplantation units. Int J Antimicrob Agents. 2015. 45(1): 33-40.

[29] Giani T, Arena F, Vaggelli G, Conte V, Chiarelli A, De Angelis L H, Fornaini R, Grazzini M, Niccolini F, Pecile P, Rossolini GM. Large Nosocomial Outbreak of Colistin-Resistant, Carbapenemase-Producing Klebsiella pneumoniae Traced to Clonal Expansion of an mgrB Deletion Mutant. J Clin Microbiol. 2015. 53(10): 3341-4.

[30] Kanerva M, Skogberg K, Ryynänen K, Pahkamäki A, Jalava J, Ollgren J, Tarkka E, Lyytikäinen O. Coincidental detection of the first outbreak of carbapenemase-producing Klebsiella pneumoniae colonisation in a primary care hospital, Finland, 2013. Euro Surveill. 2015. 20(26).

[31] Seara N, Oteo J, Carrillo R, Pérez-Blanco V, Mingorance J, Gómez-Gil R, Herruzo R, Pérez-Vázquez M, Astray J, García-Rodríguez J, Ruiz-Velasco LM, Campos J, de Burgos C, Ruiz-Carrascoso G. Interhospital spread of NDM-7-producing Klebsiella pneumoniae belonging to ST437 in Spain. Int J Antimicrob Agents. 2015. 46(2): 169-73.

[32] de Galassus A G, Cizeau F, Agathine A, Domrane C, Ducellier D, Fihman V, Decousser JW.

350　Contribution and limits of clinical specimens for the screening of intestinal

351　multi-drug-resistant bacteria in view of laboratory automation. J Hosp Infect. 2017. 97(1):

352　59-63.

356    Table I. Clinical characteristics of patients

| Patients | Clinical diagnosis | Mode of delivery | Age (w) | Sex | Body weight (kg) | Antibiotics | Apgar score | Outcome |
|---|---|---|---|---|---|---|---|---|
| P1 | Premature infant Low birth weight infant NRDS | Caesarean delivery | 33 | F | 1.55 | TZP, MEM | 6-8-8 | Treated |
| P2 | Premature infant Low birth weight infant NRDS | Normal delivery | 31 | F | 1.60 | TZP | 8-8-8 | Treated |
| P3 (first of twins) | Premature infant Very low birth weight infant NRDS | Midwifery delivery | 29 | F | 1.45 | TZP | 8-9-9 | Treated |
| P4 (second of twins) | Premature infant Very low birth weight infant NRDS | Midwifery delivery | 29 | F | 1.26 | TZP | 8-9-9 | Automatic discharge |
| P5 | Neonatal hyperbilirubinemia | Forceps delivery | 40 | M | 2.76 | TZP | 10-10-10 | Treated |
| P6 | Neonatal hyperbilirubinemia | Caesarean delivery | 40 | F | 3.83 | TZP | 10-10-10 | Treated |
| P7 | Premature infant Very low birth weight infant NRDS | Caesarean delivery | 30 | M | 1.00 | TZP | 6-8-8 | Treated |

P: patient; NRDS: neonatal respiratory distress syndrome; W: weeks; F: female; M: male; kg: kilogram; TZP: piperacillin/tazobactam. MEM: meropenem.

360 Table II. The characteristics of *K. pneumoniae*

| Patient | Date of detection | Isolate No. | Isolate site | Isolates | STs | Resistance phenotype |
|---|---|---|---|---|---|---|
| P1 | 2013/6/3 AM 10:03 | kpn1340 | Blood | kpn | ND | ND |
| | | kpn1341 | UVCP | kpn | ND | ND |
| | | kpn1342 | Sputum | kpn | 37 | Carbapenemases |
| P2 | 2013/6/13 AM 07:20 | kpn1343 | Blood | kpn | 37 | Carbapenemases |
| | | kpn1344 | Blood | kpn | 3006 | ESBLs |
| | | Negative | UVCP | Negative | ND | ND |
| | | kpn1356 | Sputum | kpn | ND | ND |
| P3 (first of twins) | 2013/6/14 AM 09:00 | kpn1345 | Blood | kpn | 37 | Carbapenemases |
| | | kpn1346 | Blood | kpn | 3006 | ESBLs |
| | | kpn1347 | UVCP | kpn | 37 | Carbapenemases |
| | | kpn1357 | Sputum | kpn | ND | ND |
| | 2013/6/28 | kpn1362 | Sputum | kpn | 1224 | Carbapenemases |
| P4 (second of twins) | 2013/6/14 AM10:40 | kpn1348 | Blood | kpn | 3006 | ESBLs |
| | | kpn1349 | Blood | kpn | 37 | Carbapenemases |
| | | kpn1350 | UVCP | kpn | 37 | Carbapenemases |
| | | kpn1351 | UVCP | kpn | 3006 | ESBLs |
| | | kpn1358 | Sputum | kpn | ND | ND |
| P5 | 2013/6/15 AM10:56 | kpn1352 | Blood | kpn | 3006 | ESBLs |
| | | Negative | Sputum | Negative | ND | ND |
| P6 | 2013/6/15 PM14:46 | kpn1353 | Blood | kpn | 37 | Carbapenemases |
| | | kpn1354 | Blood | kpn | 3006 | ESBLs |
| | | Negative | Sputum | Negative | ND | ND |
| P7 | 2013/6/15 AM10:30 | kpn1355 | UVCP | kpn | 3006 | ESBLs |
| | 2013/6/18 AM11:08 | kpn1359 | Sputum | kpn | ND | ND |

361 P: patient; UVCP: umbilical vein catheter tip; kpn: *Klebsiella pneumonia*; N: negative; ND: not done;

362 AM: ante meridiem; PM: post meridiem

Table III. Antibiotic susceptibility of *K. pneumoniae* Isolates ST37 and ST3006

| Antibiotics | MIC (ug/mL) S/I/R | | | |
| --- | --- | --- | --- | --- |
| | *K. pneumoniae* ST37(n=7) | | *K. pneumoniae* ST3006(n=7) | |
| Ampicillin-sulbactam | ≥32/ | R | ≥32 | R |
| Cefazolin | ≥64/ | R | ≥64 | R |
| Ceftriaxone | ≥64 | R | ≥64 | R |
| Cefotetan | ≥64 | R | ≤4 | S |
| Ceftazidime | ≥64 | R | ≤1 | S |
| Cefepime | ≥64 | R | ≤1 | S |
| Gentamicin | ≥16 | R | ≤1 | S |
| Tobramycin | =8 | I | ≤1 | S |
| Amikacin | ≤2 | S | ≤2 | S |
| Levofloxacin | ≤0.25 | S | ≤0.25 | S |
| Ciprofloxacin | ≤0.25 | S | ≤0.25 | S |
| Aztreonam | ≥64 | R | =64 | R |
| Imipenem | =4 | R | ≤1 | S |
| Ertapenem | ≥8 | R | ≤0.5 | S |
| Tigecycline* | ≤0.25 | S | ≤0.25 | S |
| Polymyxin B* | ≤0.25 | S | ≤0.25 | S |
| Piperacillin–Tazobactam | =64 | R | ≤4 | S |
| Trimethoprim–Sulfamethoxazole | ≤20 | S | ≤20 | S |

MIC: minimal inhibitory concentration; S: sensitive; I: intermediary; R: resistance; *: tigecycline and polymyxin B antibiotic susceptibility were determined by micro-broth dilution method

367 Table IV. Antibiotic resistance ontology annotates for antibiotic resistance genes

| ORF_ID | Best Hit ARO | Best Identities | ARO | ARO_category |
|---|---|---|---|---|
| KPN1343_GM005273 | OXA-33 | 1 | ARO:3001781 | antibiotic inactivation enzyme, determinant of beta-lactam resistance |
| KPN1343_GM005323 | AAC (6')-IId | 1 | ARO:3002548 | antibiotic inactivation enzyme, determinant of aminoglycoside resistance |
| KPN1343_GM005284 | AAC (3)-IIa | 1 | ARO:3002533 | antibiotic inactivation enzyme, determinant of aminoglycoside resistance |
| KPN1343_GM005274 | AAC (6')-Ib-cr | 1 | ARO:3002547 | antibiotic inactivation enzyme, determinant of aminoglycoside resistance |
| KPN1343_GM005278 | TEM-1 | 1 | ARO:3000873 | antibiotic inactivation enzyme, determinant of beta-lactam resistance |
| KPN1343_GM002829 | SHV-11 | 1 | ARO:3001070 | antibiotic inactivation enzyme, determinant of beta-lactam resistance |
| KPN1343_GM005272 | catB3 | 1 | ARO:3002676 | antibiotic inactivation enzyme, determinant of phenicol resistance |
| KPN1343_GM005271 | arr-3 | 1 | ARO:3002848 | antibiotic inactivation enzyme, determinant of rifamycin resistance |
| KPN1343_GM005269 | sul1 | 1 | ARO:3000410 | antibiotic target replacement protein, determinant of sulfonamide resistance |
| KPN1343_GM004245 | oqxB | 1 | ARO:3003923 | efflux pump complex or subunit conferring antibiotic resistance |
| KPN1343_GM004244 | oqxA | 1 | ARO:3003922 | efflux pump complex or subunit conferring antibiotic resistance |
| KPN1343_GM005322 | catB3 | 0.99 | ARO:3002670 | antibiotic inactivation enzyme, determinant of streptogramin resistance |
| KPN1343_GM000019 | CRP | 0.99 | ARO:3000518 | efflux pump complex or subunit conferring antibiotic resistance |
| KPN1344_GM005191 | TEM-1 | 1 | ARO:3000873 | antibiotic inactivation enzyme, determinant of beta-lactam resistance |
| KPN1344_GM005193 | CTX-M-3 | 1 | ARO:3001866 | antibiotic inactivation enzyme, determinant of beta-lactam resistance |
| KPN1344_GM000088 | vgaC | 1 | ARO:3002831 | efflux pump complex or subunit conferring antibiotic resistance |
| KPN1344_GM003641 | CRP | 0.99 | ARO:3000518 | efflux pump complex or subunit conferring antibiotic resistance |

368 ORF: open reading frame, ID: identity, ARO: antibiotic resistance ontology

371　　Fig. 1

376    Fig. 2

378    Fig. 3

**Figure Legends:**

Fig. 1 Pulsed-field gel electrophoresis (PFGE) of *K. pneumoniae*.

MLST: multi-locus sequence typing; ST: sequence type.

15 *K. pneumoniae* strains were divided into three clusters: Clusters A: 1344, 1346, 1348, 1351, 1352, 1354, 1355; Cluster B: 1362; Cluster C: 1342, 1343, 1345, 1347, 1349,1350, 1353. MLST typing also divided 15 *K. pneumoniae* strains into three groups: ST37, ST1224, ST3006.

Fig. 2 KPN1343 and KPN1344 PHI Phenotype classification.

PHI: pathogen–host interactions database. PHI Phenotype classification showed that the number of reduced virulence genes matched the database mostly, 205 and 209, respectively.

Fig. 3 KPN1343 and KPN1344 genomic islands.

KPN1343 has 15 genomic islands and KPN1344 has 19 genomic islands, the length and direction of the genes are shown in Fig. 3 (length of <15 kb only is shown).