CHARACTERIZATION OF INFLUENZA B VIRUS VARIANTS WITH REDUCED
NEURAMINIDASE INHIBITOR SUSCEPTIBILITY

Farrukee R[1,2], Zarebski AE[3], McCaw JM[3,4,5, 6] , Bloom JD[7], Reading PC[1,2], #Hurt AC[1,2, *]

[1] WHO Collaborating Centre for Reference and Research on Influenza, at the Peter
Doherty Institute for Infection and Immunity, Melbourne, Victoria, 3000, Australia

[2] Department of Microbiology and Immunology, The University of Melbourne, at the
Peter Doherty Institute for Infection and Immunity, Melbourne, Victoria, 3000, Australia

[3] School of Mathematics and Statistics, The University of Melbourne, Melbourne,
Australia

[4] Centre for Epidemiology and Biostatistics, Melbourne School of Population and Global
Health, The University of Melbourne, Melbourne, Australia

[5] Victorian Infectious Diseases Reference Laboratory Epidemiology Unit, at the Peter
Doherty Institute for Infection and Immunity, Melbourne, Victoria, 3000, Australia

[6] Infection and Immunity theme, Murdoch Children's Research Institute, Royal
Children's Hospital, Melbourne, Australia

[7] Division of Basic Sciences and Computational Biology Program, Fred Hutchinson
Cancer Research Center, Seattle, WA, USA.

.

#Address correspondence to Aeron C Hurt ,aeron.hurt@influenzacentre.org

*WHO Collaborating Centre for Reference and Research on Influenza, Peter Doherty
Institute, 792 Elizabeth Street, Melbourne, VIC 3000, Australia. Tel: +613-93429314;
Fax: +613-93429393

Running Title: Characterization of influenza B virus variants

Abstract

Treatment options for influenza B virus infections are limited to neuraminidase inhibitors (NAIs) which block the neuraminidase (NA) glycoprotein on the virion surface. The development of NAI resistance would therefore result in a loss of antiviral treatment options for influenza B infections. This study characterized two contemporary influenza B viruses with known resistance-conferring NA amino acid substitutions, D197N and H273Y, detected during routine surveillance. The D197N and H273Y variants were characterized *in vitro* by assessing NA enzyme activity and affinity, as well as replication in cell culture compared to NAI-sensitive wild-type viruses. *In vivo* studies were also performed in ferrets to assess the replication and transmissibility of each variant. Mathematical models were used to analyse within-host and between-host fitness of variants relative to wild-type viruses. The data revealed that the H273Y variant had similar NA enzyme function relative to its wild-type but had slightly reduced replication and transmission efficiency *in vivo*. The D197N variant had impaired NA enzyme function but there was no evidence of reduction in replication or transmission efficiency in ferrets. Our data suggest that the influenza B variant with H273Y NA substitution had a more notable reduction in fitness compared to wild-type viruses than the influenza B variant with the D197N NA substitution. Although a D197N variant is yet to become widespread, it is the most commonly detected NAI-resistant influenza B virus in surveillance studies. Our results highlight the need to carefully monitor circulating viruses for the spread of influenza B viruses with the D197N NA substitution.

Introduction

Influenza A and B viruses co-circulate in the human population and cause yearly

epidemics worldwide. Even though influenza B viruses generally cause a milder disease

than influenza A viruses, in certain seasons they can contribute to a substantial burden

of disease and up to 25% of influenza-related mortality (1, 2). Vaccines are the main

control measure for influenza disease prevention, but a recent analysis of data between

2004-2014 revealed that the effectiveness of the vaccine against influenza B infections

was only 54% (3).

The neuraminidase inhibitors (NAIs) (zanamivir, oseltamivir, laninamivir and peramivir)

are a class of antiviral drugs that target the conserved amino acid residues of the viral

neuraminidase (NA) active site and competitively inhibit enzyme function (4, 5).

However, mutations in the NA enzyme active site or adjacent framework residues can

abrogate NAI interaction with the NA and reduce the susceptibility of viruses to one or

more of these drugs (6). Antiviral options for the treatment of influenza B viruses are

limited to NAIs as the older class of influenza antivirals, the adamantanes, are only

effective against influenza A viruses (6). Development of NAI resistance in influenza B

viruses is therefore a public health concern as it would remove the only available

treatment option.

In vitro studies consistently demonstrate that compared to influenza A viruses, influenza

B viruses exhibit reduced sensitivity to oseltamivir, suggesting that the drug may already

have reduced effectiveness against influenza B viruses (6-10). The clinical relevance of

this has not been fully elucidated, but in 7 out of 9 clinical studies it was shown that

oseltamivir treatment resolved symptoms faster in influenza A patients than in influenza

B patients (11). Considering this, it is possible that NA mutations that only moderately

alter the oseltamivir susceptibility of influenza B viruses may have a significant impact

on the clinical effectiveness of the drug.

A number of different NA substitutions at conserved amino acid positions (e.g. E117,

D197, I221 and H273) have previously been described to confer reduced inhibition by

the NAIs *in vitro* (8, 12-21), but the impact of these substitutions on enzyme function,

virus replication or transmissibility, has only been assessed in a limited number of

studies (14, 22, 23). The fitness of influenza B viruses with either the H273Y or D197N

NA substitution is of particular interest as a number of viruses with either substitution

have been recently found in community settings in patients who did not receive NAI

treatment (8, 9, 17, 18, 24). Two reports have identified household transmission of

influenza B viruses with the D197N NA substitution (18, 25) and more recently, a global

surveillance report identified a cluster of six influenza B viruses with the D197N NA

substitution in Japan in early 2014, further suggesting potential community transmission

of the variant virus (18). There has also been examples of suspected transmission of

influenza B viruses with the H273Y NA substitution (9). The H273Y NA substitution in

influenza B viruses occurs at the equivalent residue to that of the H275Y NA substitution

in A(H1N1) viruses, which was present in the oseltamivir resistant A(H1N1) viruses that

spread globally in 2008/9 (26, 27).

The effect of H273Y NA substitutions in influenza B viruses has been previously studied

using reverse genetics (rg) in the B/Yamanashi/166/98 virus background (15, 22, 23).

To date, few studies have reported the effect of the H273Y or the D197N NA

substitution in contemporary viruses, which is important because it has been shown that

the fitness consequences of resistance-conferring mutations can vary due to the genetic

background of the NA (28, 29). Although experiments using reverse genetics can be

useful in defining the impact of a single mutation on viral fitness; they don't evaluate the

effect of compensatory mutations that may play an important role in the fitness of that

virus. Our aim was to characterize two 'naturally' occurring influenza B variant viruses

containing either the H273Y or D197N NA substitutions which had been detected during

routine surveillance in patients not being treated with NAIs, compared to closely

matched wild-type viruses by assessing their enzyme function, *in vitro* replication and *in*

vivo replication and transmission.

Results

NAI susceptibility, NA activity, surface expression and substrate affinity

The effect of the D197N and H273Y substitutions on NA enzyme function were

assessed using four different assays. The MUT-Y273 variant had a 3-fold increase in

oseltamivir IC_{50} and an 85-fold increase in peramivir IC_{50} compared to WT-H273, but the

IC_{50} for zanamivir and laninamivir were not significantly different (Table I). The MUT-

Y273 virus had comparable K_m (substrate affinity) to that of the WT-H273 virus (Table I).

Similarly, the relative NA surface expression and enzyme activity of the MUT-Y273 virus

compared to the WT-H273 virus was 115 ± 13.4% (mean ± SEM) and 119 ± 23.1%

respectively, neither of which was significantly different (Figure I).

The MUT-N197 virus had a 2- 3-fold reduction in susceptibility to all four NAIs compared

to that of the WT-D197 virus (Table I). The substrate affinity (K_m) was 3-fold lower for

the MUT-N197 virus compared to the WT-D197 virus (Km values are inversely

proportional to substrate affinity) but this difference was not statistically significant

(Table I). The cell-surface expression of the NA enzyme of the Mut-N197 variant was

123 ± 10.9% relative to the WT-D197 virus but the NA enzyme activity of the Mut-N197

variant was only 14.2 ± 3.0%, which was significantly lower (p<0.01) than the WT-D197

virus (Figure I).

In vitro replication kinetics

Multi-step growth curves were performed in MDCK and MDCK-SIAT (SIAT) cells to

assess the *in vitro* replication kinetics of the WT and MUT viruses. The MUT-Y273

variant showed significantly reduced viral titres compared to the WT-H273 virus at 24

and 36 hours post-infection in MDCK cells and at 36 hours post-infection in SIAT cells

(Figure II a, b). However, at 48 hours post-infection there were no significant

differences in viral titres between the MUT-Y273 and WT-H273 viruses in either cell

line.

The MUT-N197 variant demonstrated no reduction in replication compared to the WT

virus in MDCK cells and actually outgrew the WT-D197 virus in SIAT cells at 48 hours

post-infection where it reached a significantly higher titre of 6.61 ± 0.11 vs 5.5 ± 0.2

Log_{10} $TCID_{50}$/ml (p<0.01) (Figure II c, d).

Replication and transmission of pure viral populations in ferrets

To gain insight regarding virus replication *in vivo*, we determined titres of infectious virus

in nasal wash samples taken at various times after intranasal infection of ferrets. Figure

III a & c summarizes the results for animals infected with either the WT-H273 or the

Mut-Y273 virus. In both donor groups (infected by direct intranasal inoculation),

infectious virus was detectable until day 8 post-infection and peak viral titres were not

significantly different. There was also no significant difference in the area under the

curve (AUC) of $TCID_{50}$ plots over the 10 day period (Figure III a, c). Body temperature

and weight were also assessed daily after infection and no significant differences were

noted between donor animals infected with WT-H273 or Mut-Y273, although there was

a trend towards greater weight loss in donors infected with WT-H273 (Figure S2).

We examined virus replication in nasal washes from naïve recipient ferrets who were

co-housed with experimentally infected donors to assess the transmissibility of WT-

H273 and Mut-Y273 viruses. All four recipients co-housed with WT-H273-infected

donors shed infectious virus but only two out of four recipients co-housed with MUT-

Y273-infected donors shed detectable infectious virus (Figure III a, c). There was a

delay in transmission of the MUT-Y273 virus compared to the WT-273 virus, as peak

virus titres in WT-H273-infected recipients was detected on day 4 post-exposure while

the two recipients infected with the MUT-Y273 virus reached peak viral titres on day 7

post-exposure (Figure III a, c).

Next, we examined virus replication following experimental infection of animals with WT-

D197 and Mut-N197 viruses. The kinetics of viral shedding was similar between donor

ferrets experimentally infected with either WT-D197 or MUT-N197 viruses (Figure III b,

d). The AUC of the viral replication plots and the peak viral titres between the donor

groups were not significantly different. Infectious viral titres were detectable in nasal

washes of at least one ferret until day 6 post-infection and donors did not experience

any significant changes in either body weight or temperature (Figure S2).

The WT-D197 virus transmitted successfully to all four recipients, while the MUT-N197

virus transmitted to three out of the four recipients. However, length of viral shedding

and the peak viral titres were not significantly different between recipients infected with

either the WT-D197 virus or the MUT-N197 virus (Figure III b, d); there was also no

transmission delay observed as with MUT-Y273. Similar to the donor ferrets, the

recipient ferrets infected with either WT-D197 or MUT-N197 virus experienced no

significant changes in either body weight or temperature.

Pyrosequencing analysis of relevant NA substitutions in nasal wash samples from all

infected animals confirmed that neither MUT-N197 nor the MUT-Y273 viruses reverted

to wild-type genotype following replication within the airways of donor ferrets, or

following transmission to recipient animals.

Relative fitness between viral pairs based on 'competitive mixtures' analyses

To gain further insight into the fitness differential between each variant virus and their

corresponding wild-type viruses, a competitive mixtures experiment was performed in

ferrets. Mathematical models for the progression of the infection and its transmission

were fitted to measurements collected during the experiment. The parameter estimates

from this fitting process provide information about the relative fitness of the variant for

replication within-host and its ability to transmit between hosts.

These inferences are summarized in Table II and visualizations are shown in Figure IV.

We define the relative within-host fitness as the ratio of the basic reproduction numbers

of the variant and its wild-type counterpart. . For the MUT-Y273 virus the Ro ratio is

estimated to be 0.84 with a 2-sided 95% CI of (0.70, 1.02), with our analysis indicating

that there is a 95% chance that this ratio is less than 1. So while the MUT-Y273 variant

is almost certainly less fit than its wild-type counterpart, the difference is very small.

There was little evidence that the mutation affected transmission fitness, with the

mutant's ability to transmit reduced by a factor of 0.90 (0.60, 1.35).

For MUT-D197, we find little evidence for a difference in within-host fitness, with an

estimated ratio of 0.94 (0.81, 1.10) and a 75% chance that the variant is less fit than the

wild-type. Interestingly, the transmission fitness of the MUT-N197 virus was higher by a

factor of 2.78 (1.56, 5.59) over the wild-type. However, it is important to note that this

estimate is driven heavily by two recipient ferrets where a large increase in the

population of MUT-N197 was observed following transmission from donors infected with

a 50:50 ratio of WT-D197: MUT-N197 viruses. Although the statistical model is

designed to account for the variation among ferrets, we are careful to not draw too

strong a conclusion on this result given the limited number of animals in the study.

Discussion

This study aimed to characterize two contemporary influenza B variants with either the

D197N or H273Y NA substitution. The data showed that the MUT-N197 variant had

significantly reduced NA enzyme activity but the replication and transmission efficiency

of this virus in an *in vitro* or *in vivo* model was not notably reduced compared to a

closely matched wild type virus. Further to this, mathematical analysis from competitive

mixtures experiments, found little evidence of reduced relative within-host fitness and no

evidence of reduced between-host transmission of the MUT-N197 virus. A previous

study using an influenza B virus (B/Rochester/02/2001) with the D197N NA substitution

also showed that when ferrets were infected with a mixture of wild type and variant

virus, a mixture was still maintained 5-days post infection (30).

It is interesting that the significantly reduced NA enzyme activity of the MUT-N197 virus

did not predict the outcome of the replication or transmission fitness of this variant in the

ferret model. This may be because the enzyme-based assay is inherently limited in its

ability to capture the complex interplay between the different viral proteins during

replication and transmission in a biological setting. This kind of discrepancy between *in*

vitro and *in vivo* data has been described previously (31), and highlights the need for

caution when assessing viral fitness using only *in vitro* parameters.

In this study, the MUT-Y273 virus had equivalent NA activity and expression to the WT-H273 virus, but delayed growth in cell culture. Previous experiments using a rg-H273Y virus showed that relative to the rg-WT, the rg-H273Y virus had significantly higher NA activity and surface expression, and superior replication kinetics in a competitive mixtures experiment in normal human bronchial epithelial cells (14, 22, 23). The discrepancy in these fitness outcomes between our 2015 H273Y virus and a 1998 rg-H273Y virus may be due to differences in viral background (16 amino acid differences). The MUT-Y273 virus had similar replicative fitness in donor ferrets following intranasal infection viruses and delayed transmission between co-housed ferrets compared to WT-H273 virus when tested in a pure population. Mathematical analysis showed evidence of slight reduction in the within-host fitness of the variant in a competitive mixture with the wild-type but no evidence of reduction in between-host transmission.

There is suggestive evidence that transmission efficiency of the MUT-Y273 variant when tested as a pure population had a lower fitness that the wild-type (transmission to 2/4 vs 4/4 ferrets), compared to when the variant was assessed in a competitive mixtures experiment where it had similar fitness to the WT-H273 virus. It is possible that a variant virus with reduced NA enzyme function may benefit from replicating in the presence of a fully functional wild-type virus rather than as a pure population, because the NA enzyme from the wild-type virus could compensate for the loss of activity of the variant. This phenomenon has been previously shown with viruses differing by one amino acid at residue 151 of the NA, where mixed viral populations grew better in cell culture than either variant alone (32).

The viruses assessed in our study were studied in comparison to closely matched

circulating wild-type viruses which had identical HA, NA (except for the resistance

mutation of interest) and MP sequences; however a limitation of this comparison is that

some amino acid differences did exist between the WT and MUT viruses in the internal

genes (Figure S1) and it is possible that one or more of the substitutions in the internal

genes may have contributed to the fitness differences observed between the WT and

MUT viruses. Nevertheless these variant viruses represent real circulating strains and

therefore have greater relevance than performing experiments using reverse genetics

using laboratory reference strains in lab as has been done in the past with B/Yamanashi

viruses. Further to none of the changes (or equivalent changes in influenza A viruses)

have been reported in the literature to alter virus fitness, replication or transmission.

It is also important to note that our study contributes to the very limited number of

animal studies that have reported influenza B virus transmission to date. One previous

study demonstrated successful influenza B transmission between guinea pigs in contact

and non-contact models (33), while only two studies (with mixed results) have

previously reported on influenza B transmission in the ferret mode l(34, 35). Of these

two, one study reported successful aerosol and contact transmission of a mouse-

adapted influenza B virus, but could not demonstrate transmission by either model

using a non-mouse adapted isolate (34). The second study using a modified influenza B

virus lacking the NB protein showed that one out of the four recipient ferrets were

successfully infected by aerosol transmission (35). Our results show successful contact

transmission of both the influenza B wild-type viruses tested (4 out of 4 ferrets) and

transmission of influenza B variants in the ferret model with circulating human influenza

viruses.

Conclusion

The results from our experiments suggest that of two influenza B viruses with reduced

NAI susceptibility, the viral fitness of a recent influenza B virus with the H273Y NA

substitution was reduced compared to an equivalent wild-type virus, but an influenza B

virus with a D197N NA substitution showed little reduction in reproduction or

transmission in the ferret model. Although the frequency of influenza B viruses with the

D197N NA substitution circulating in the community is still less than 1%, it remained the

most commonly reported NAI resistant strains in recent year. Future studies to

determine if the MUT-D197 virus has reduced capacity to transmit via aerosol

transmission will be important.

In the future it is possible that the NA variants examined in our study may gain

additional 'permissive' mutations that improve their fitness, as was observed with the

H275Y NA substitution in seasonal H1N1 viruses (28). This may be more likely for

viruses with the D197N NA substitution, which appear to have a smaller fitness deficit

than viruses with the H273Y NA substitution. This knowledge, alongside the fact that

clinical reports suggests that oseltamivir is less effective against influenza B and

therefore minor changes in the viral NA may further reduce oseltamivir effectiveness,

emphasizes the importance of continued monitoring of influenza B viruses for NA

substitutions in the future.

Materials and Methods

Virus selection and stocks

A B/Yamagata/16/88-like Influenza B viruses with the D197N NA substitution

designated B/Singapore/GP702/2015 (MUT-N197), was isolated from a 8-year old

female patient not receiving NAI therapy and submitted to the WHO Collaborating

Centre for Reference and Research on Influenza, Melbourne, Australia by the Ministry

of Health, Singapore. Another B/Yamagata/16/88-like virus with the H273Y NA

substitution designated B/Perth/136/2015 (MUT-Y273) was isolated from a 61-year old

female patient not receiving NAI therapy and submitted during routine surveillance.

Wild-type viruses B/Minnesota/23/2015 (WT-D197) and B/England/598/2014 (WT-

H273) were kindly provided by the Centers for Disease Control and Prevention (CDC),

Atlanta, United States and the Crick Institute, London, England, respectively, and were

specifically chosen as wild-type pairs for MUT-N197 and MUT-Y273 respectively, due to

their close genetic homology to the two variant viruses. The WT-D197: MUT-N197 pair

had identical gene sequences except for the desired D197N mutation the NA, one

amino acid difference in the NS gene and one amino acid difference in the PA gene

Similarly the WT-H273: MUT-Y273 virus had two amino acid changes in the NP gene

one change in the PB1 gene and one change in the PA gene (see Figure S1 for details).

Viruses were propagated in Madin-Darby canine kidney (MDCK) cells and infectivity

titres were determined by calculating the 50% tissue culture infectious dose ($TCID_{50}$/mL)

as previously described(36).

In vitro NA inhibition and NA kinetics assay

To determine the IC_{50} (50% inhibitory concentration) of the influenza B viruses for each

NAI, a fluorometric MUNANA (2-(4-methylumbelliferyl)-a-d-N-acetylneuraminic acid)-

based assay was performed as previously described(37). NAI compounds were

purchased from Carbosynth, UK. The Michaelis-Menten constant K_m (substrate affinity

was also determined by measuring the rate of reaction at different MUNANA

concentrations, as previously described(38), but using a MUNANA concentration range

from 0-800 µM. The IC_{50} and enzyme kinetics assays were determined by three

independent experiments.

Neuraminidase surface expression and activity assay

Measurement of cell-surface NA expression and activity was performed by transfecting

HEK-293T (293T) cells with an expression plasmid bearing the NA gene of interest with

a C-terminal V5 epitope tag(28, 39, 40).

At 20 hours post-transfection, the NA enzyme activity on the cell surface was measured

using a modified MUNANA-based assay as previously described(39). The enzyme

expression in these cells was also measured by staining with an Anti-V5 Alexaflour 647

Antibody (Thermofisher, Australia) and measuring staining intensity using flow

cytometry(39). The NA activity results were normalized for total surface expression and

results for each variant were normalized against fluorescent staining intensity/activity of

corresponding wild-type viruses. Two independent experiments were done to assess

NA expression and activity, where each virus was tested in triplicates.

In vitro replication kinetics

A multi-step replication experiment was carried out in MDCK and MDCK-SIAT1(SIAT)

cell lines, which were grown to confluence in 6-well plates and infected at a MOI of 0.01

$TCID_{50}$/cell as described previously[36]. Infectivity titres in the samples were then

determined by $TCID_{50}$ assay.

Ethics statement

Ferrets (weight range 600–1500 g) were housed in the Bioresources Facility at the

Peter Doherty Institute for Infection. Experiments were conducted with approval from

the University of Melbourne Biochemistry & Molecular Biology, Dental Science,

Medicine, Microbiology & Immunology, and Surgery Animal Ethics Committee, in

accordance with the NHMRC Australian code of practice for the care and use of animals

for scientific purposes (project license number 1313040).

Ferret experiments

Male and female ferrets aged 6-12 months old were used in experiments. All animals

received food and water ad libitum. All test animals were confirmed to be seronegative

to circulating human influenza strains using a hemagglutination inhibition assay prior to

the experiments. The replication and contact transmission efficiency of the wild-type and

variant viruses in the ferret model was tested by experimental infection of ferrets with a

'pure' population of wild-type (WT) or mutant (MUT) viruses as described below. In

independent experiments, ferrets were infected with 'mixtures' of wild-type and mutant

viruses to determine the relative fitness of each in 'competitive mixtures'

experiments(36, 40, 41). Viral stocks were standardized to 1×10^5 TCID$_{50}$/ml in

Dulbecco's Phosphate Buffered Saline (Sigma, Australia) for ferret inoculation. To make

viral mixtures for WT-H273: MUT-Y273 and WT-D197: MUT-N197, the standardized

stocks were volumetrically mixed at 80:20, 50:50 and 20:80 ratios.

On day 0, donor ferrets were anaesthetized by intramuscular injection with 20 mg/ml

Ilium Xylazil-20 (Troy-laboratories, Australia) and intranasally inoculated with 0.5 mL of

(i) pure populations of WT (n=4) and MUT (n=4) viruses, or (ii) mixtures at ratios of

50:50 (n=4), 80:20 (n=2) and 20:80 (n=2) of WT and MUT viruses. The ferrets were

then housed in high-efficiency particulate air (HEPA) filtered cages. After 24 hours (Day

1 p.i.), one naïve recipient ferret was co-housed with one donor ferret to allow for

contact transmission. The donor and recipients were co-housed for the entire length of

the experiment (11 days). Nasal washes were collected from donors and recipients

each day throughout the experiment and temperature and weight was monitored daily,

as described previously (40). All ferrets were euthanized on Day 11. Infectious viral load

in ferret nasal washes were quantified by TCID$_{50}$ assay.

380   *RNA extraction and quantitative analysis of viral RNA in ferret nasal washes*

The viral RNA from ferret nasal wash was extracted and quantified using real-time PCR. The real-time primers were provided by the CDC (Atlanta, USA), and capable of detecting the influenza B NS gene. Internal controls with known RNA copy numbers of B/England/598/2014 NS transcripts were generated in-house and included for each assay. This was used to correlate cycle threshold (Ct) with RNA copy number and quantitate viral RNA in nasal wash samples.

Molecular analysis of ferret nasal washes

The proportion of WT and MUT viruses in each nasal wash sample was determined by pyrosequencing[40] using primer pairs specifically designed for PCR amplification and sequencing analysis (Table S3). To validate the performance of the assay, pure populations and known mixtures for each WT:MUT pair were serially diluted and tested across three separate assays. Based on these validation tests, only samples with greater than 4.3 Log_{10} NS gene copies/µL were analysed for mixture estimation.

Quantitative assessment of differences in virus replication and transmission fitness

The fitness of a virus depends on its ability replicate within a host and to transmit between hosts. We used a mechanistic model of viral replication and competition to assess the relative within-host replicative fitness of the variant compared to the wild-type virus[40, 42].

As previously described, our model expanded on the standard Target Cell - Infectious Cell – Virus (TIV) model of viral dynamics[42]. The viral population is stratified by strain

and accounts for infectious and non-infectious viral material by simultaneously fitting to

$TCID_{50}$, real-time PCR and pyrosequencing data(43).

From the model fit, we estimate the ratio of the basic reproduction number, R_0, of the

two viral strains in the competitive mixture, where R_0 describes the expected number of

secondary infections produced by a single infected cell in a population of susceptible

cells(44).

To assess the relative ability of the variant to transmit between hosts we fitted a

transmission model as described in McCaw et al. to pyrosequencing data(41). We

utilized the pyrosequencing measurements of the proportion of mutant virus in the

donors (on the inferred day of infection) and the recipients. We extended on our

previous work to account for pyrosequencing measurement errors in producing our

estimates. The transmission model has a single parameter describing the relative

transmission fitness of the two strains where a value greater than 1 indicates enhanced

transmissibility of the mutant.

Statistical analysis

Viral titres between different groups from the *in vitro* and *in vivo* experiments were

analysed by a two-way ANOVA with Bonferroni's post-hoc analysis using GraphPad

Prism 5.0. All other comparisons such as Area under the Curve (AUC), peak viral titres,

NA activity and K_m values were made using non-parametric Mann Whitney's U test.

Parameters for the mathematical models used to determine relative finesses (both

within and between hosts) were estimated from the data in a Bayesian framework using

the R interface to STAN (45, 46) (see Text S4).

Acknowledgements

The Melbourne WHO Collaborating Centre for Reference and Research on Influenza is

Supported by the Australian Government Department of Health

We gratefully acknowledge GISRS laboratories for providing influenza viruses to the

Melbourne WHO Collaborating Centre and Thomas Cumming, Celeste Tai, Leonard

Izzard, Ding Yuan Oh, Rebecca Jayne Bower and Charlene Plascenia for assistance

with animal experiments.

441 **TABLE I: Effect of neuraminidase mutation on IC_{50} value and enzyme kinetics**

	IC_{50} (mean ± SD)[a]	Kinetic (mean ± SD)[a]

Virus	Zanamivir	Oseltamivir	Laninamivir	Peramivir	K_m (µM)
WT-H273	2.2±1.3	36.1±5.4	2.6±2.0	1.0±0.05	11.5±1.5
Mut-Y273	1.4±0.5 (1)	96.7±10.8 (3)	1.63±0.4 (1)	80.9 ±6.8 (85)*	12.3±0.8 (1)
WT-D197	1.6±0.2	35.1±3.3	2.8±1.3	1.0±0.1	13.8±1.3
Mut-N197	4.5±0.7 (3)	66.2±7.5 (2)	5.0±1.2 (2)	3.0±0.7 (3)	30.3±3.3 (3)

443   (n) Fold difference from corresponding wild type values.

444   * MUT-Y273 has significantly higher IC_{50} for peramivir than WT-H273 (Mann-Whitney's

445   U Test, $p < 0.05$).

446   [a] The results summarise the mean± SD of three independent experiments

457　**TABLE II: Relative Within-Host and Transmission fitness between viral pairs**

458　**tested in 'competitive mixtures' during ferret infection**

		Relative within-host fitness	Relative Transmission fitness
Mutant	**Wild Type**	**R_0 ratio (95% CI)[a]**	**Relative fitness value (95% CI)[b]**
Mut-Y273	WT-H273	0.84 (0.70,1.02)	0.90 (0.60, 1.35)
Mut-N197	WT-D197	0.94 (0.81,1.10)	2.78 (1.56, 5.53)

460　[a]If the R_0 ratio <1 it indicates fitness advantage of wild-type over mutant, and if R_0

461　ratio>1 the mutant is fitter than wild-type

462　[b]Relative transmission fitness value<1 indicates fitness advantage of wild type over

463　mutant and fitness value>1 indicate mutant is fitter than wild-type

Figure Legends:

Figure I: The mean NA surface expression and activity of influenza B variants relative to the corresponding WT. HEK-293T cells were transfected with plasmids containing the NA gene encoding for WT and variant proteins. At 20 hours post-transfection cells were analysed for NA activity using a MUNANA-based assay, and NA expression using a FACs-based assay. The activity and expression data of the variants were normalized relative to their corresponding WT and expressed as the mean±SEM. Data are derived from two independent experiments, each containing triplicate samples and compared between relevant groups using the non-parametric Mann-Whitney's U-Test.

Figure II: Multi-cycle replication of influenza B WT and MUT viruses in MDCK and SIAT cells. Cells were infected at a MOI of 0.01 and supernatants collected and assayed for infectious virus at 2, 24, 36 and 48 hr post-infection. Data show the mean (± SEM) of triplicate samples for Mut-Y273 vs WT-H273 (a, b) and Mut-N197 vs WT-D197(c, d). A two-way ANOVA analysis with Bonferroni's post-test analysis was done to compare mean viral titres at each time point between WT and MUT viruses. * $p < 0.05$, **$p < 0.01$, ***$p < 0.001$

Figure III: Viral shedding in nasal wash samples from donor and recipient ferrets infected with WT or MUT influenza B viruses. Donor ferrets (n= 4 per group) were experimentally infected with 10^5 $TCID_{50}$ of 'pure' WT or MUT viruses. Naïve recipients were co-housed with each donor one day after experimental infection of donors. Daily nasal wash samples were collected from both donors and recipients over a period of 11 days and titres of infectious virus in nasal wash samples were determined by $TCID_{50}$ assay. A-D summarizes the $TCID_{50}$ results with the bar graphs showing the mean (± SEM) titres at each day for each group. At each time point, mean viral titres were compared between donors infected with either WT or MUT virus by two-way ANOVA with Bonferroni's post-test analysis. Similarly, analysis was done to compare titres between recipients infected with either WT or MUT virus. Significant differences in mean viral titres were observed between WT-H273 and Mut-Y273 infected recipients on day 3 and 4 post-exposure and between WT-D197 and Mut-N197 infected donors on day 1 post-inoculation. Area under the curve (AUC) of $TCID_{50}$ graphs were also calculated and compared by Mann-Whitney's U test; no significant differences were detected between donor animals infected with WT and corresponding MUT viruses. * $p<0.05$, **$p<0.01$, ***$p<0.001$.

Figure IV: Summary of changes in mutant virus proportions in nasal washes within-host and during transmission in 'competitive mixtures' ferret experiments.

To assess within-host and transmission fitness of mutant viruses relative to their corresponding wild-types, ferrets were experimentally co-infected with viral mixtures of WT:MUT at different ratios (50:50, 80:20 and 20:80) and co-housed with naïve recipients. The proportion of WT vs MUT was determined in the collected nasal wash samples by pyrosequencing and data were analysed only from animals where the proportion of WT and MUT could be accurately determined. A-B plots individual points that correspond to the change in the proportion of mutant virus in one ferret between the first day of infection and last day of infection (y-axis). This allows for a visual representation of within-host fitness. C-D illustrate individual points that corresponds to a transmission event between two ferrets. In these plots the abscissa is the proportion of MUT virus in donor ferrets the day preceding confirmed transmission and the ordinate shows the proportion of MUT virus in recipient ferrets within the first 24 hours of confirmed infection. The solid grey line is the representative curve of the shape parameter, calculated by fitting our data to a model described in more detail in the methods and supplementary. The 95% confidence interval is represented by the shaded region. If our fitted model lies above the symmetry line (y=x solid line) then the mutant virus is fitter than the wild-type virus, whereas if the fitted model lies below the symmetry line the converse is true.

Reference

1. Paul Glezen W, Schmier JK, Kuehn CM, Ryan KJ, Oxford J. 2013. The burden of influenza B: a structured literature review. Am J Public Health 103:17.
2. Gutierrez-Pizarraya A, Perez-Romero P, Alvarez R, Aydillo TA, Osorio-Gomez G, Milara-Ibanez C, Sanchez M, Pachon J, Cordero E. 2012. Unexpected severity of cases of influenza B infection in patients that required hospitalization during the first postpandemic wave. J Infect 65:423-30.
3. Belongia EA, Simpson MD, King JP, Sundaram ME, Kelley NS, Osterholm MT, McLean HQ. 2016. Variable influenza vaccine effectiveness by subtype: a systematic review and meta-analysis of test-negative design studies. Lancet Infect Dis 16:942-51.
4. Itzstein Mv, Wen-Yang W, Kok GB, Pegg MS, Dyason JC, Jin B, Van Phan T, Smythe ML, White HF, Oliver SW, Colman PM, Varghese JN, Ryan MD, Woods JM, Bethell RC, Hotham VJ, Cameron JM, Penn CR. 1993. Rational design of potent sialidase-based inhibitors of influenza virus replication. Nature:418.
5. von Itzstein M. 2007. The war against influenza: discovery and development of sialidase inhibitors. Nat Rev Drug Discov 6:967-74.
6. Gubareva LV. 2004. Molecular mechanisms of influenza virus resistance to neuraminidase inhibitors. Virus Res 103:199-203.
7. McKimm-Breschkin J, Trivedi T, Hampson A, Hay A, Klimov A, Tashiro M, Hayden F, Zambon M. 2003. Neuraminidase sequence analysis and susceptibilities of influenza virus clinical isolates to zanamivir and oseltamivir. Antimicrob Agents Chemother 47:2264-72.
8. Sheu TG, Deyde VM, Okomo-Adhiambo M, Garten RJ, Xu X, Bright RA, Butler EN, Wallis TR, Klimov AI, Gubareva LV. 2008. Surveillance for neuraminidase inhibitor resistance among human influenza A and B viruses circulating worldwide from 2004 to 2008. Antimicrob Agents Chemother 52:3284-92.
9. Meijer A, Rebelo-de-Andrade H, Correia V, Besselaar T, Drager-Dayal R, Fry A, Gregory V, Gubareva L, Kageyama T, Lackenby A, Lo J, Odagiri T, Pereyaslov D, Siqueira MM, Takashita E, Tashiro M, Wang D, Wong S, Zhang W, Daniels RS, Hurt AC. 2014. Global update on the susceptibility of human influenza viruses to neuraminidase inhibitors, 2012-2013. Antiviral Res 110:31-41.
10. Escuret V, Frobert E, Bouscambert-Duchamp M, Sabatier M, Grog I, Valette M, Lina B, Morfin F, Ferraris O. 2008. Detection of human influenza A (H1N1) and B strains with reduced sensitivity to neuraminidase inhibitors. J Clin Virol 41:25-8.
11. Farrukee R, Mosse J, Hurt AC. 2013. Review of the clinical effectiveness of the neuraminidase inhibitors against influenza B viruses. Expert Rev Anti Infect Ther 11:1135-45.
12. Garg S, Moore Z, Lee N, McKenna J, Bishop A, Fleischauer A, Springs CB, Nguyen HT, Sheu TG, Sleeman K, Finelli L, Gubareva L, Fry AM. 2013. A cluster of patients infected with I221V influenza b virus variants with reduced oseltamivir susceptibility--North Carolina and South Carolina, 2010-2011. J Infect Dis 207:966-73.
13. Carr S, Ilyushina NA, Franks J, Adderson EE, Caniza M, Govorkova EA, Webster RG. 2011. Oseltamivir-resistant influenza A and B viruses pre- and postantiviral therapy in children and young adults with cancer. Pediatr Infect Dis J 30:284-8.
14. Pascua PNQ, Marathe BM, Burnham AJ, Vogel P, Webby RJ, Webster RG, Govorkova EA. 2016. Competitive Fitness of Influenza B Viruses Possessing E119A and H274Y Neuraminidase Inhibitor Resistance–Associated Substitutions in Ferrets. PLoS ONE 11:1-19.
15. Burnham AJ, Baranovich T, Govorkova EA. 2013. Neuraminidase inhibitors for influenza B virus infection: efficacy and resistance. Antiviral research 100:10.1016/j.antiviral.2013.08.023.

16.   Okomo-Adhiambo M, Sleeman K, Ballenger K, Nguyen HT, Mishin VP, Sheu TG, Smagala J, Li Y,
Klimov AI, Gubareva LV. 2010. Neuraminidase inhibitor susceptibility testing in human influenza
viruses: a laboratory surveillance perspective. Viruses 2:2269-89.
17.   Takashita E, Meijer A, Lackenby A, Gubareva L, Rebelo-de-Andrade H, Besselaar T, Fry A, Gregory
V, Leang SK, Huang W, Lo J, Pereyaslov D, Siqueira MM, Wang D, Mak GC, Zhang W, Daniels RS,
Hurt AC, Tashiro M. 2015. Global update on the susceptibility of human influenza viruses to
neuraminidase inhibitors, 2013-2014. Antiviral Res 117:27-38.
18.   Hurt AC, Besselaar TG, Daniels RS, Ermetal B, Fry A, Gubareva L, Huang W, Lackenby A, Lee RT,
Lo J, Maurer-Stroh S, Nguyen HT, Pereyaslov D, Rebelo-de-Andrade H, Siqueira MM, Takashita E,
Tashiro M, Tilmanis D, Wang D, Zhang W, Meijer A. 2016. Global update on the susceptibility of
human influenza viruses to neuraminidase inhibitors, 2014-2015. Antiviral Res 132:178-85.
19.   Colacino JM, Laver WG, Air GM. 1997. Selection of influenza A and B viruses for resistance to 4-
guanidino-Neu5Ac2en in cell culture. J Infect Dis 176:S66-8.
20.   Cheam AL, Barr IG, Hampson AW, Mosse J, Hurt AC. 2004. In vitro generation and
characterisation of an influenza B variant with reduced sensitivity to neuraminidase inhibitors.
Antiviral Res 63:177-81.
21.   Barnett JM, Cadman A, Burrell FM, Madar SH, Lewis AP, Tisdale M, Bethell R. 1999. In vitro
selection and characterisation of influenza B/Beijing/1/87 isolates with altered susceptibility to
zanamivir. Virology 265:286-95.
22.   Burnham AJ, Armstrong J, Lowen AC, Webster RG, Govorkova EA. 2015. Competitive fitness of
influenza B viruses with neuraminidase inhibitor-resistant substitutions in a coinfection model of
the human airway epithelium. J Virol 89:4575-87.
23.   Burnham AJ, Baranovich T, Marathe BM, Armstrong J, Webster RG, Govorkova EA. 2014. Fitness
costs for Influenza B viruses carrying neuraminidase inhibitor-resistant substitutions:
underscoring the importance of E119A and H274Y. Antimicrob Agents Chemother 58:2718-30.
24.   Okomo-Adhiambo M, Sleeman K, Lysen C, Nguyen HT, Xu X, Li Y, Klimov AI, Gubareva LV. 2013.
Neuraminidase inhibitor susceptibility surveillance of influenza viruses circulating worldwide
during the 2011 Southern Hemisphere season. Influenza Other Respir Viruses 7:645-58.
25.   Hatakeyama S, Sugaya N, Ito M, Yamazaki M, Ichikawa M, Kimura K, Kiso M, Shimizu H,
Kawakami C, Koike K, Mitamura K, Kawaoka Y. 2007. Emergence of influenza B viruses with
reduced sensitivity to neuraminidase inhibitors. Jama 297:1435-42.
26.   Moscona A. 2009. Global Transmission of Oseltamivir-Resistant Influenza. New England Journal
of Medicine 360:953-956.
27.   Kramarz P, Monnet D, Nicoll A, Yilmaz C, Ciancio B. 2009. USE OF OSELTAMIVIR IN 12 EUROPEAN
COUNTRIES BETWEEN 2002 AND 2007 – LACK OF ASSOCIATION WITH THE APPEARANCE OF
OSELTAMIVIR-RESISTANT INFLUENZA A(H1N1) VIRUSES. Eurosurveillance 14.
28.   Bloom JD, Gong LI, Baltimore D. 2010. Permissive Secondary Mutations Enable the Evolution of
Influenza Oseltamivir Resistance. Science 328:1272-1275.
29.   Baz M, Abed Y, Simon P, Hamelin ME, Boivin G. 2010. Effect of the neuraminidase mutation
H274Y conferring resistance to oseltamivir on the replicative capacity and virulence of old and
recent human influenza A(H1N1) viruses. J Infect Dis 201:740-5.
30.   Mishin VP, Hayden FG, Gubareva LV. 2005. Susceptibilities of antiviral-resistant influenza viruses
to novel neuraminidase inhibitors. Antimicrob Agents Chemother 49:4515-20.
31.   Abed Y, Baz M, Boivin G. 2006. Impact of neuraminidase mutations conferring influenza
resistance to neuraminidase inhibitors in the N1 and N2 genetic backgrounds. Antivir Ther
11:971-6.
32.   Xue KS, Hooper KA, Ollodart AR, Dingens AS, Bloom JD. 2016. Cooperation between distinct viral
variants promotes growth of H3N2 influenza in cell culture. eLife 5:e13974.

33.   Pica N, Chou Y-Y, Bouvier NM, Palese P. 2012. Transmission of influenza B viruses in the guinea
631   pig. Journal of Virology 86:4279-4287.
632   34.   Kim EH, Park SJ, Kwon HI, Kim SM, Kim YI, Song MS, Choi EJ, Pascua PN, Choi YK. 2015. Mouse
633   adaptation of influenza B virus increases replication in the upper respiratory tract and results in
634   droplet transmissibility in ferrets. Sci Rep 5.
635   35.   Elderfield RA, Koutsakos M, Frise R, Bradley K, Ashcroft J, Miah S, Lackenby A, Barclay WS. 2016.
636   NB protein does not affect influenza B virus replication in vitro and is not required for replication
637   in or transmission between ferrets. J Gen Virol 97:593-601.
638   36.   Hurt AC, Nor'e SS, McCaw JM, Fryer HR, Mosse J, McLean AR, Barr IG. 2010. Assessing the Viral
639   Fitness of Oseltamivir-Resistant Influenza Viruses in Ferrets, Using a Competitive-Mixtures
640   Model. Journal of Virology 84:9427-9438.
641   37.   Hurt AC, Barr IG, Hartel G, Hampson AW. 2004. Susceptibility of human influenza viruses from
642   Australasia and South East Asia to the neuraminidase inhibitors zanamivir and oseltamivir.
643   Antiviral Res 62:37-45.
644   38.   Marathe BM, Leveque V, Klumpp K, Webster RG, Govorkova EA. 2013. Determination of
645   neuraminidase kinetic constants using whole influenza virus preparations and correction for
646   spectroscopic interference by a fluorogenic substrate. PLoS ONE 8.
647   39.   Bloom JD, Nayak JS, Baltimore D. 2011. A Computational-Experimental Approach Identifies
648   Mutations That Enhance Surface Expression of an Oseltamivir-Resistant Influenza
649   Neuraminidase. PLoS ONE 6:e22201.
650   40.   Butler J, Hooper KA, Petrie S, Lee R, Maurer-Stroh S, Reh L, Guarnaccia T, Baas C, Xue L, Vitesnik
651   S, Leang SK, McVernon J, Kelso A, Barr IG, McCaw JM, Bloom JD, Hurt AC. 2014. Estimating the
652   fitness advantage conferred by permissive neuraminidase mutations in recent oseltamivir-
653   resistant A(H1N1)pdm09 influenza viruses. PLoS Pathog 10.
654   41.   McCaw JM, Arinaminpathy N, Hurt AC, McVernon J, McLean AR. 2011. A Mathematical
655   Framework for Estimating Pathogen Transmission Fitness and Inoculum Size Using Data from a
656   Competitive Mixtures Animal Model. PLoS Computational Biology 7:e1002026.
657   42.   Petrie SM, Butler J, Barr IG, McVernon J, Hurt AC, McCaw JM. 2015. Quantifying relative within-
658   host replication fitness in influenza virus competition experiments. J Theor Biol 382:259-71.
659   43.   Petrie SM, Guarnaccia T, Laurie KL, Hurt AC, McVernon J, McCaw JM. 2013. Reducing
660   uncertainty in within-host parameter estimates of influenza infection by measuring both
661   infectious and total viral load. PLoS One 8:e64098.
662   44.   Baccam P, Beauchemin C, Macken CA, Hayden FG, Perelson AS. 2006. Kinetics of influenza A
663   virus infection in humans. J Virol 80:7590-9.
664   45.   Team SD. 2017. Rstan: The R interface to Stan. http://mc-stan.org. Accessed
665   46.   Carpenter B, Gelman A, Hoffman MD, Lee D, Goodrich B, Betancourt M, Brubaker M, Guo J, Li P,
666   Riddell A. 2017. Stan: A Probabilistic Programming Language. 2017 76:32.

FIGURE I

Figure I: The mean NA surface expression and activity of influenza B variants relative to the
corresponding WT. HEK-293T cells were transfected with plasmids containing the NA gene encoding for
WT and variant proteins. At 20 hours post-transfection cells were analysed for NA activity using a
MUNANA-based assay, and NA expression using a FACs-based assay. The activity and expression data
of the variants were normalized relative to their corresponding WT and expressed as the mean±SEM.
Data are derived from two independent experiments, each containing triplicate samples and compared
between relevant groups using the non-parametric Mann-Whitney's U-Test.

 FIGURE II

Figure II: Multi-cycle replication of influenza B WT and MUT viruses in MDCK and SIAT cells. Cells were infected at a MOI of 0.01 and supernatants collected and assayed for infectious virus at 2, 24, 36 and 48 hr post-infection. Data show the mean (± SEM) of triplicate samples for Mut-Y273 vs WT-H273 (a, b) and Mut-N197 vs WT-D197(c, d). A two-way ANOVA analysis with Bonferroni's post-test analysis was done to compare mean viral titres at each time point between WT and MUT viruses. * $p < 0.05$, **$p < 0.01$, ***$p < 0.001$

FIGURE III

---- Limit of detection

Figure III: Viral shedding in nasal wash samples from donor and recipient ferrets infected with WT or MUT influenza B viruses. Donor ferrets (n= 4 per group) were experimentally infected with 10^5 $TCID_{50}$ of 'pure' WT or MUT viruses. Naïve recipients were co-housed with each donor one day after experimental infection of donors. Daily nasal wash samples were collected from both donors and recipients over a period of 11 days and titres of infectious virus in nasal wash samples were determined by $TCID_{50}$ assay. A-D summarizes the $TCID_{50}$ results with the bar graphs showing the mean (± SEM) titres at each day for each group. At each time point, mean viral titres were compared between donors infected with either WT or MUT virus by two-way ANOVA with Bonferroni's post-test analysis. Similarly, analysis was done to compare titres between recipients infected with either WT or MUT virus. Significant differences in mean viral titres were observed between WT-H273 and Mut-Y273 infected recipients on day 3 and 4 post-exposure and between WT-D197 and Mut-N197 infected donors on day 1 post-inoculation. Area under the curve (AUC) of $TCID_{50}$ graphs were also calculated and compared by Mann-Whitney's U test; no significant differences were detected between donor animals infected with WT and corresponding MUT viruses. * $p<0.05$, **$p<0.01$, ***$p<0.001$

Figure IV: Summary of changes in mutant virus proportions in nasal washes within-host and during transmission in 'competitive mixtures' ferret experiments. To assess within-host and transmission fitness of mutant viruses relative to their corresponding wild-types, ferrets were experimentally co-infected with viral mixtures of WT:MUT at different ratios (50:50, 80:20 and 20:80) and co-housed with naïve recipients. The proportion of WT vs MUT was determined in the collected nasal wash samples by pyrosequencing and data were analysed only from animals where the proportion of WT and MUT could be accurately determined. A-B plots individual points that correspond to the change in the proportion of mutant virus in one ferret between the first day of infection and last day of infection (y-axis).This allows for a visual representation of within-host fitness. C-D illustrate individual points that corresponds to a transmission event between two ferrets. In these plots the abscissa is the proportion of MUT virus in donor ferrets the day preceding confirmed transmission and the ordinate shows the proportion of MUT virus in recipient ferrets within the first 24 hours of confirmed infection. The solid grey line is the representative curve of the shape parameter, calculated by fitting our data to a model described in more detail in the methods and supplementary. The 95% confidence interval is represented by the shaded region. If our fitted model lies above the symmetry line (y=x solid line) then the mutant virus is fitter than the wild-type virus, whereas if the fitted model lies below the symmetry line the converse is true.