Classification:

PHYSICAL SCIENCES: Biophysics and Computational Biology

Title:

Cellular heterogeneity in pressure and growth emerges from tissue topology and geometry

Yuchen Long[1,*], Ibrahim Cheddadi[2,4], Vincent Mirabet[1], Gabriella Mosca[3], Mathilde Dumond[1,5], Jan Traas[1], Christophe Godin[1,2], Arezki Boudaoud[1,*].

1. Laboratoire Reproduction et Développement des Plantes, Université de Lyon, ENS de Lyon, UCB Lyon 1, CNRS, INRA, INRIA, F-69342, Lyon, France.

2. INRIA/CIRAD/INRA Project-team Virtual Plants, France.

3. Department of Plant and Microbial Biology, University of Zürich, Zollikerstrasse 107, CH-8008 Zürich, Switzerland.

4. Current address: Univ. Grenoble Alpes, CNRS, Grenoble INP, TIMC-IMAG, 38000 Grenoble, France.

5. Current address: Department of Biosystems Science and Engineering, ETH Zurich, Mattenstrasse 26, 4058, Basel, Switzerland.

*. Correspondence and requests for materials should be addressed to yuchen.long@ens-lyon.fr or arezki.boudaoud@ens-lyon.fr

The authors declare no competing financial interests.

Abstract

Cell-to-cell heterogeneity prevails in many biological systems, although its origin and function are often unclear. Cell hydrostatic pressure, alias turgor pressure, is an essential property in tissue physiology and morphogenesis, and its spatial variations are often overlooked. Here, using micro-indentation by atomic force microscope, we find that hydrostatic pressure is heterogeneous between adjacent cells in the epidermis of Arabidopsis shoot apical meristem, a population of stem cells from which originate all aerial organs of the plant. Combining experiments and physical modelling of cell wall mechanics and water movement, we propose that cell size and tissue topology contribute to the patterning of heterogeneities in pressure and growth, partially reminiscent of a liquid foam. Pressure correlates either positively or negatively with cellular growth rate depending on conditions, suggesting that the meristem is at a tipping point between limitation of growth by cell wall extensibility and by water conductivity. Together, our results shed light on spatial variations of pressure and growth rate, with potential roles in tissue homeostasis.

Key words

Cellular heterogeneity, hydrostatic pressure, tissue topology, growth mechanics, tissue hydraulics, atomic force microscopy, biophysical modelling.

Significance statement

Within a living tissue or an organ, cells grow at markedly different rates. Nevertheless, the causes and consequences of such heterogeneity in growth are poorly understood. We therefore examined cell internal pressure and cell growth in a population of stem cells in the model plant Arabidopsis, by combining optical microscopy and biophysical measurements. We found that pressure and growth in a given cell were both correlated with the number of its neighbours, i.e. with local topology. We retrieved experimental observations by using a mathematical model accounting for force propagation and water movement in the tissue. Our work suggests that water availability may be partially limiting for growth and puts forward topology as a source of spatial variations in cell growth.

Introduction

Cell-to-cell fluctuations are observed in many biological phenomena like gene expression, signalling, cell size regulation and growth (1–8). Notably, heterogeneity in cell size and growth rate often prevails and impacts tissue patterning and macroscopic growth robustness (1, 2). Cell volume change is driven by osmosis (9–11) and the resulting intracellular hydrostatic pressure, and is restrained by peripheral constraints (plasma membrane, cytoskeletal cortex, extracellular matrix, or cell wall) in plant (12), animal (13), tumorous (14) and microbial cells (15). Despite numerous studies (15–17), the spatial variation of hydrostatic pressure in multicellular tissues, and its relation with cell-to-cell growth variability, remain elusive.

Due to high difference between internal and external osmotic potential, cells with rigid cell walls – like in plants, bacteria and fungi – accumulate hydrostatic pressure, alias turgor pressure, often greater than atmospheric pressure (Fig. S1) (12). Animal cells also accumulate hydrostatic pressure, especially when compacted or contracting (13, 14), though to a lesser extent than walled cells. In plants, turgor pressure is believed to positively correlate with growth rate, as depicted in the Lockhart-Ortega equation (Fig. S1) (17). However, observations suggest that growth rate and pressure level are not always associated (16), making it difficult to assess and understand the link between growth regulation and cellular pressure build-up. Here, we aimed at addressing this issue in a model plant tissue. We quantified turgor pressure and growth rate in a multicellular tissue and we use a physical model coupling tissue growth mechanics with water movement to account for our observations.

Results

Atomic force microscopy reveals turgor pressure heterogeneity in Arabidopsis shoot apical meristem

Recent advances in atomic force microscopy (AFM) enabled non-invasive turgor pressure retrieval utilizing indentation force-displacement and surface topography in living plant cells (Fig. 1A) (18). Accordingly, we used a pressurized thin-shell model to deduce the turgor pressure value from the AFM-measured force-displacement curve, which is influenced by turgor pressure, cell wall mechanical properties, and cell 3D geometry (Fig. S2; see

Supplementary Discussion) (19). We carefully evaluated the contributions of these parameters

to the indentation results, and excluded many possible biases in deducing turgor from our AFM

approaches (Fig. S2; see Supplementary Discussion). Additionally, influence of tissue context

on indentation readout (20) was rigorously tested and ruled out by numerical simulations of

indentations on realistic 3D meshes accounting for pressurized epidermal cells (Fig. S8; see

Supplementary Discussion). Taken together, we concluded that our AFM measurements are

sensitive to cell-to-cell turgor pressure variation, and semi-quantitative in regards to the

absolute value of pressure. Accordingly, in the following, we use 'turgor pressure' to qualify

the AFM-determined values.

We applied AFM-based pressure deduction to the epidermis of the *Arabidopsis thaliana* shoot

apical meristem (SAM), a system featuring substantial growth heterogeneity (1). We included

untreated wild-type SAMs and a conceptually simpler model of chemically treated SAM that

resembles a "foam of cells" (co-treated with naphthylphthalamic acid (NPA), a polar auxin

transport inhibitor that induces pin-formed SAMs, and oryzalin, a microtubule-depolymerizing

drug that blocks cell division but permits continuous, isotropic growth; hereafter referred to as

"oryzalin-treated SAMs") (21). Walls between cells are often curved in growing oryzalin-

treated SAMs, like liquid films in a foam, suggesting that neighbouring cells have different

turgor pressure (21) in a growing tissue (Fig. S3). Cell-specific AFM measurements on nine

SAMs revealed that turgor pressure is markedly heterogeneous in oryzalin-treated SAMs

epidermis (Fig. 1E and Fig. S2). Oryzalin-treated SAMs had average turgor pressures spanning

the range 0.65 to 1.72 MPa (1.21 ± 0.34 MPa, mean ± standard error of mean; Fig. S4). These

values are comparable to the 2.0 MPa osmotic pressure estimated by incipient plasmolysis (Fig.

S5). The AFM-determined turgor also decreased with increasing osmolarity of external medium

(Fig. S5), further supporting this AFM-based approach.

Cell-specific measurements revealed that cellular pressure anticorrelates with the number of

epidermal cell-neighbours: cells with fewer neighbours have higher turgor pressure (Fig. S6).

We focused on cell-to-cell variations of turgor and we removed variations between SAMs by

normalizing cellular pressure to the average pressure per SAM; normalized pressure decreases

with neighbour number, being 1.3-fold greater in four- than in eight-neighboured cells ($n = 202$

cells, Pearson correlation coefficient R = -0.31, $p < 10^{-5}$) (Fig. 1F), consistent with previous

prediction (21).

Untreated SAMs exhibit more constant turgor, as determined by AFM, at 2.62 ± 0.03 MPa (Fig.

S4), with the same magnitude as the estimated osmotic pressure at 1.2 MPa (Fig. S5) and the

preliminary values obtained previously (22) (see Supplementary Discussion). Consistently, we

also detected similar, though slightly weaker, turgor–neighbour-number anticorrelation in

untreated SAMs (7 SAMs, n = 503 cells, R = -0.16, $p = 10^{-4}$) (Fig. 1G-J and Fig. S6, S7),

confirming that non-random turgor pressure heterogeneity establish in tissues with static

topology (no neighbour number change) or dynamic topology (neighbour numbers change due

to division).

A mechanical-hydraulic model predicts that pressure heterogeneity emerges from tissue

topology

Earlier tissue models (21, 23) treated intracellular pressure as an input or required differential

osmotic pressure. Nevertheless, the anticorrelation that we find between pressure and neighbour

number is reminiscent of liquid foams (24). We therefore tested whether pressure heterogeneity

could emerge from the classical plant water relations and wall mechanics (25) (Fig. 2A). We

use a 2D vertex model, with topological distribution similar to SAM (Fig. S9), that actually

expands the Lockhart-Ortega equation of visco-elasto-plastic growth of single cell (Fig. S1)

(17) to cell walls in a multicellular tissue with more realistic 2D polygonal geometry. Each cell

wall has a thickness w and behaves as an elastic material of modulus E when wall tension

induced by turgor pressure P is low enough, more specifically when the elastic strain (ε,

reversible deformation) is lower than a threshold ε^Y. When tension in the wall exceeds the

corresponding threshold ($E\varepsilon^Y$), the cell wall undergoes irreversible expansion, akin to visco-

plastic flow, with an extensibility (effective viscosity) Φ^w (Fig. 2A). Water flux from

extracellular space is proportional to $L^a \Delta\Psi$, where $\Delta\Psi = \Delta\Pi - P$ is the cross-membrane water

potential difference (external pressure is assumed to be zero; for the sake of simplicity we set

$\Delta\Pi$ constant) and L^a is the conductivity. Intercellular water redistribution via plant

plasmodesmata (26), animal gap junctions, or cytoplasmic bridges (27, 28) is driven by

intercellular differences in turgor pressure P, with a conductivity L^s. We did not prescribe turgor

pressure, instead letting it emerge from local mechanical and hydraulic interplays (see

Supplementary Materials and Methods for detailed model description).

Whereas cell wall extensibility can be more limiting for growth than water conductivity in fast

elongating organs (29), we found that most aquaporin (channel protein allowing rapid water

flux (30)) isoforms had significantly lower expression in inflorescence than in other fast

growing tissues like stem and root (31) (with a reduction up to ~80%). Therefore we considered

limitation of growth by cell wall extensibility and by water conductivity on equal footing. We

first imposed slower water flux than cell wall extension, and found that, with homogenous

effective parameters, we could recover the turgor–neighbour-number anticorrelation in our

model, without or with cell divisions to mimic oryzalin-treated or untreated SAM (no division

5 simulations, $n = 1219$ cells, $R = -0.87$, $p < 10^{-100}$; dividing 8 simulations, $n = 3240$ cells, $R =$

-0.58, $p < 10^{-100}$) (Fig. 2B-I). This implies that local hydrostatic pressure heterogeneity does not

require differential cellular osmotic pressure (21) in a growing tissue, and predicts a topological

origin of pressure variability, similar to liquid foams where pressure heterogeneity is due to fast

liquid flow and limited gas diffusion rate between bubbles (24). Indeed, water flow in a tissue

parallels gas flow between cells in a foam and mechanics of cell walls replaces the rheology of

the liquid films.

In order to test the robustness of this result, we explored the parameter space of the vertex

model. Analytical exploration in a two-cell system (25) had showed that systems dynamics is

controlled by two main dimensionless parameters, α^s, which compares intercellular water

conductivity to total (intercellular and transmembrane) conductivity, and α^a, which compares

the limitation of growth by transmembrane permeability to the combined limitation of growth

by cell wall extensibility and transmembrane permeability (expression of these dimensionless

parameters are given in the Methods). Accordingly, our first results, obtained with $\alpha^a = 0.1$ and

$\alpha^s = 0.1$, denoted water flux-limited growth. We uplifted this limit, consistently with

observations in many cell types (29), by increasing α^a from 0.1 to 0.5 and 0.9, where water

conductivity equals or is relatively one magnitude higher than wall extensibility (32), and still

recovered the turgor–neighbour-number anticorrelation (Fig. 2J), demonstrating that strong flux

limitation is not required for pressure heterogeneity. Next, we relaxed the intercellular flux

limitation, by increasing α^s from 0.1 to 0.5 and 0.9, and showed that the anticorrelation is

maintained even when intercellular flux is much stronger than osmotic water uptake (Fig. 2J).

Shifting cell size checkpoint at division (by either -25% or +50%), which effectively changes

average cell surface-to-volume ratio that antagonistically weighs the contribution of flux and

wall limitations, also yields steady turgor–neighbour-number anticorrelation (Fig. 2J).

Altogether, the parameter space exploration indicates that topology-related pressure

heterogeneity is a robust outcome of the model.

In SAM surface, cell neighbour number and size are coupled (Fig. S9). Consistently, cell-

specific turgor pressure anticorrelates with normalized cell area (Fig. S9). To uncouple cell size

and topology, we modified the initial state of our simulation to have four-neighboured cells

bigger than eight-neighboured cells, and found that higher turgor pressure accumulated in four-

neighboured cells, not in the smaller cells (Fig. S9), confirming that turgor pressure

heterogeneity emerges from tissue topology rather than cell size differences. This can be

explained by local topology, which determines cell wall angles and the subsequent tension

distribution at each three-cell junction (Fig. 2K-N): in our model, wall stress and strain above

the growth threshold are relaxed at a rate limited by wall extensibility and hydraulic

conductivity. Therefore, stress and strain remain close to their values at growth threshold and

are almost homogeneous in oryzalin-like simulations (Fig. 2C), and the sum of wall tension at

each three-cell junction (vertex) mostly depends on the angles between walls. The vertex

between three hexagonal cells with 120° internal angles has a sum of tension at zero (Fig. 2K,

L). Fewer-neighboured cells have sharper internal angles, so the sum of tension at vertex is

greater towards the cell interior, creating additional inward compression and prompting higher

pressure build-up at equilibrium (Fig. 2M, N). In dividing simulations, as in the untreated case,

new walls do not bear stress right after division because they form from cell interior. These new

walls are gradually strained due to mesh growth but do not yield (expand) to release stress

before reaching the threshold (Fig. 2G). Consequently, cell division keeps the wall stress from

homogenizing and dampens the turgor–neighbour-number anticorrelation (Fig. 2I). Notably,

although tissue topology is sufficient to explain pressure heterogeneity, we do not exclude a

role of other biological parameters and/or sources of noise in pressure variations, as suggested

by weaker correlations in experiments (Fig. 1F, J) than in simulations. Nonetheless, since tissue

topology is highly conserved in many biological systems (33), we propose that hydrostatic

pressure heterogeneity may emerge in compact tissues with polygonal cells (34) and non-

instantaneous water movement, due to the adjustment to reconcile local mechanical and

hydraulic conditions.

Heterogeneity in growth rate is patterned according to the balance between wall extensibility and tissue conductivity

Next, we monitored areal growth rate of SAM epidermal cells by time-lapse confocal

microscopy. As observed previously, untreated SAMs exhibited slower growth in the central

zone (CZ), where stem cells reside, than in the surrounding peripheral zone (PZ) (35) (Fig. S7).

In addition, cellular growth rate anticorrelates with neighbour number (11 SAMs, $n = 1491$

cells, $R = -0.15$, $p = 10^{-8}$; Fig. 3E) and cell size (Fig. 3F; $R = -0.33$, $p = 10^{-38}$), supporting

previous reports that smaller cells in SAM grow faster (36, 37), and suggests that higher turgor

pressure in fewer-neighboured cells associates with faster growth. In oryzalin-treated SAMs,

however, the fewer-neighboured and small cells grew slower (14 SAMs, $n = 1160$ cells; Fig.

3K, neighbour number $R = 0.20$, $p = 10^{-11}$; Fig. 3L, cell size $R = 0.16$, $p = 10^{-8}$). This suggests

that higher turgor pressure associates with either faster or slower growth depending on

conditions (Fig. S9). Although seemingly a small shift, this negative-to-positive slope change

of local growth heterogeneity captures a strong qualitative inversion of growth behaviour (Fig.

3D, J), where relative cell size increases exponentially in opposite fashions (Fig. S10).

Accordingly, smaller cells expand more than larger cells in untreated SAMs, which may

contribute to cell size homeostasis, a phenomenon that would be absent in oryzalin-treated

SAMs.

We further explored the growth trend in our vertex model. Based on the parameter exploration

on dividing mesh aimed to reproduce untreated SAM behaviour, we found both positive and

negative correlations between growth rate and cell neighbour number (Fig. S11). This indicates

that growth trend is sensitive to parameters that alter the balance between water flux and wall

expansion. Conservatively, we chose equal contribution by flux and wall in our model ($\alpha^a =$

0.5), which required scaling up theoretical cell size (by +50%) to maintain growth trend as in

untreated SAM: Figures 4A-C show that the fewer-neighboured and smaller cells grow faster

(10 simulations, n = 5971; G vs N, R = -0.25, p = 10^{-84}; G vs A, R = -0.96, $p < 10^{-100}$).

We then attempted to reproduce oryzalin-treated behaviour, guided by the experimental

observations that, besides stopping cell division, oryzalin treatment also yields higher osmotic

pressure (1.6-fold) and drastically thicker cell walls (4-fold) (Fig. S5, S12). We found that both

stalling division and increasing osmotic pressure failed to invert the growth trend in the model,

while doubling and quadrupling wall thickness inverted the correlation of growth rate to

neighbour number and cell size (Fig. 4D-F, G; S13), like in oryzalin-treated SAM. Combining

higher osmotic pressure and thicker wall revealed that quadrupling wall thickness can robustly

trigger growth trend inversion (Fig. 4 and Fig. S13).

Effectively, changing osmotic pressure and wall thickness also alters the wall–flux limitation

balance of the system (Fig. 4G). Higher osmotic pressure induces faster water influx (reduced

limitation by flux). The extra volume strains the walls to accumulate stress farther beyond the

threshold, which is relaxed by wall yielding (growth) with extensibility as the rate limit.

Meanwhile, wall thickening reduces wall stress and strain towards the threshold for expansion,

effectively reducing the mechanical drive of growth and increasing the weight of water

permeability in limiting growth, similar to liquid foam. We do not exclude other possible

parameter changes triggered by oryzalin treatment, like water conductivity and wall synthesis

rate, that also contribute to the wall–flux limitation balance. Nevertheless, implementing the

observed cell wall thickening in the model is sufficient to explain the observed growth rate

inversion from untreated to oryzalin-treated scenario.

Conclusions

In this study, we analysed the intercellular heterogeneity of turgor pressure and its effect on cell

growth rate in the epidermis of Arabidopsis SAM, utilizing both indentation experiments and

mechanical modelling. We propose that tissue mechanics and hydraulics act in parallel with the

established genetic regulations in SAM cell growth. We demonstrate that cellular hydrostatic

pressure is innately heterogeneous in Arabidopsis SAM epidermis; however, it is not a proxy

of local growth rate variability, contrary to expectations. Instead, our analysis suggests that

turgor pressure and growth rate are combinatorial outputs of local mechanics and hydraulics. Each of these parameters can be controlled by genetic and biochemical inputs, and small uniform changes in these biological inputs can enable drastic shifts of cell size distribution (Fig. S14), unlike liquid foams in which cell size distribution always enlarge over time. Our simulation and experiment data suggest that Arabidopsis SAM is likely close to the tipping point between the limits of wall synthesis and water influx. Such inversion in growth trend is not irrelevant to normal development: for example, a growth variability switch is observed during sepal development, where clones of smaller and often fewer-neighboured cells grow faster in young sepals then slower in older sepals, effectively switching from homogenizing to amplifying cell size variability (38). The emergent heterogeneity of local growth and hydrostatic pressure likely underlies morphogenesis in other compact tissues with polygonal cells.

Materials and Methods

Plant materials, treatments and growth conditions

Arabidopsis thaliana GFP-LTi6b (ecotype WS-4) reporter line and ecotype Col-0 was used(39). Untreated inflorescence meristems were obtained from soil-grown plants, first in short-day (8 h light 20°C / 16 h dark 19°C cycle) for 3 to 4 weeks then transferred to long-day (16 h light 20°C / 8 h dark 19°C cycle) for 1 to 2 weeks to synchronize bolting. Oryzalin-treated inflorescence meristems were obtained from plants grown on custom-made Arabidopsis medium (40) (Duchefa) supplemented with 1% agar-agar (Merck) and 10 μM N-1-naphthylphthalamic acid (NPA, Sigma-Aldrich/Merck) for 3 weeks. Pin-formed inflorescence meristems from NPA medium were immersed in 10 μg/mL oryzalin (Sigma-Aldrich/Merck) twice (3 h duration, 24 h interval) (21). For mechanical measurements and time-lapse confocal imaging, meristems were mounted on Arabidopsis apex culture medium (ACM) (40) with 2% agarose and 0.1% plant preservation mixture (PPM, Plant Cell Technology) to prevent contamination, and cultivated in long-day condition.

Atomic force microscopy and force curve analysis

AFM measurement and analysis were performed as in Beauzamy et al., 2015 (18) with

278 modifications. For detail, see Supplementary Materials and Methods.

Time-lapse confocal microscopy and image processing

Untreated (dissected) and oryzalin-treated meristems were mounted and grown on ACM with

0.8% agarose and 0.1% PPM for live imaging. Confocal stacks were taken on an LSM 700

confocal microscope (Carl Zeiss) operated with the ZEN 2010 software (version 6.0, Carl

Zeiss), using a W N-Achroplan 40x/0.75 M27 water immersion objective, and on a TCS SP8

confocal microscope (Leica) operated with the Leica Application Suite X software (version 3.5,

Leica), using a Fluotar VISIR 25x/0.95 water immersion lens. GFP was excited at 488 nm and

emission detected between 415 – 735 nm. Stacks have resolution of 1028×1028 pixels, with

resolution ranging between 3.2 to 4.4 pixels/μm; Z steps were between 0.5 and 0.85μm.

For hyperosmotic treatments, meristems were mounted in Petri-dishes on Patafix (UHU), then

submerged in liquid ACM containing 0.1% PPM. Samples were first imaged in liquid ACM

(plus 0.1% PPM), then submersion medium was changed to ACM plus desired concentration

of NaCl (plus 0.1% PPM) by first rinsing with 3~5 mL target solution, then soaked in target

solution for 5 minutes before imaging. Because of the reduced signal in hyperosmotic solutions,

possibly due to the altered refractive index, stronger gain was used to reach comparable signal

intensity.

Confocal image stacks were analysed with MorphoGraphX (MGX version 1.0) (41). For details,

see Supplementary Materials and Methods.

Statistical analysis

Data were processed using Excel 2000 (Microsoft). All Tukey box plots depict the first, second

(median) and third quartiles of data distribution, with whiskers marking the lowest/highest data

within 1.5 interquartile ranges (IQR) of the lower/upper quartiles. Tukey's outliers are depicted

as small circles outside the whiskers. Values like turgor pressure, cell area and growth rate were

normalized to the average per meristem. After normalization, every cell was considered as one

biological sample, and all linear regressions and Pearson correlations were performed on whole

datasets. For simulations, cells on the edge of the mesh were not analysed due to border effect.

Extremely rare polygon classes (i.e. triangle and nonagon) were not shown on the box plots in

the main figures but were included in linear regression and Pearson correlation tests and were

plotted in Extended data Figure 6. Kolmogorov–Smirnov test was used to distinguish cell

neighbour number distribution (significance level $\alpha = 0.01$).

Modelling and FEM simulation

We build a vertex-based model of plant tissues at cellular level that couples osmosis-driven

hydraulic fluxes between cells and from apoplast with a fixed water potential, and cell wall

mechanics which resists and grows under tension, based on the modelling framework from

Cheddadi et al. 2019 (25). Realistic meshes for FEM indentation simulations were generated

from confocal stacks, while inflation and indentation simulations were performed with the same

procedure adopted from Mosca et al., 2017 (20). For details, see Supplementary Materials and

Methods.

Acknowledgements

We thank P. Bolland and A. Lacroix for plant care, G. Cloarec for help with TEM, L. Beauzamy

for AFM training, C. Mollier for help in estimating confocal technical errors, A. Kiss for help

with anisotropic filtering parameterization, G. Cerrutti for help in implementing the division

algorithm and in segmentation and mesh generation for the indentation simulations, and R.

Smith for help in segmentation and mesh generation for the indentation simulations. We

acknowledge the contribution of the PLATIM facility of SFR Biosciences (UMS3444/CNRS,

US8/Inserm, ENS de Lyon, UCBL) for AFM and confocal microscopy, and of Centre

Technologique des Microstructures (UCBL, Lyon) for electron microscopy. This work was

supported by a fellowship from Institut Universitaire de France and an ERC Starting Grant

"PhyMorph" to A.B. (ERC-2012-StG-307387), an EMBO Long-term Fellowship to Y.L.

(EMBO ALTF 168-2015), and an Agropolis Foundation grant (MecaFruit3D) to I.C. and C.G.

Author contributions

This study was initiated by A.B. Y.L. and A.B. designed the experiments. Y.L. executed AFM

and confocal microscopy, acquired experimental data except EM, and analysed experimental

and simulation data. J.T. performed electron microscopy. V.M and M.D. wrote scripts to

facilitate experimental data analysis. I.C. and C.G. designed physical model of tissue growth.

I.C. implemented the model, ran simulations, optimized model parameters and analysed

simulation data. G.M. designed, ran, and analysed indentation simulations. Y.L. and A.B.

contributed to the design of simulations. Y.L., I.C., V.M., G.M., C.G. and A.B. contributed to

data interpretation. Y.L. and A.B. wrote the manuscript with inputs from the other authors.

Data availability

339 All materials, scripts and datasets generated and analysed during the current study are available

340 from the corresponding authors upon reasonable request.

Figure 1. Turgor pressure heterogeneity in SAM is associated with variation of cell topology.

(**a** and **b**) Schematic representation of AFM nanoindentation for turgor pressure measurement. *r*, probe tip radius; *k*, cantilever stiffness; *F*, indentation force; *Z*, indentation depth; *P*, turgor pressure; *σ*, cell wall tension. (**c** to **j**) Cell-specific turgor pressure in oryzalin-treated (c to f) and untreated SAM (g to j). (c and g) Top-view surface projections of SAM, oryzalin-treated (c) or untreated (g), with plasma membrane GFP signal; scale bars represent 20 μm. (d and h) Cell topology determined on segmented cell contour images, around SAM centre; numbers indicate neighbour number in epidermis. (e and i) AFM-determined cell-specific turgor pressure heat maps. (f and j) Box plots of cellular turgor pressure *P* normalized per SAM against cell topology *N* (f, oryzalin-treated 9 SAMs, *n* = 202 cells; j, untreated 7 SAMs, *n* = 503 cells). Circles are Tukey's outliers; lowercase letters indicate statistically different populations (Student's *t*-test, *p* < 0.05); red lines indicate linear regressions, with Pearson correlation coefficient *R* and corresponding *p*-value.

Figure 2. Turgor pressure heterogeneity emerges from cell topology.
(**a**) Schematic representations of model components, including apoplasmic and intercellular water fluxes (left), mechanical equilibrium at three-cell junctions (middle) and the visco-elasto-plastic cell wall rheology. P, cell-specific turgor pressure; σ, cell wall tension; Φ^w, wall extensibility; ϵ^Y, wall strain threshold; E, wall Young's modulus. (**b** to **i**) Turgor pressure in non-dividing (oryzalin-treated, b to e) and dividing (untreated) simulations (f to i), scale bars are 5 unit length. (b and f) Cell topology; (c and g) cell wall strain relative to strain threshold; (d and h) cellular turgor pressure heat maps; (e and i) Box plots of normalized cellular turgor pressure P against cell topology N (d, non-dividing 5 simulations, $n = 1219$ cells; g, dividing 8 simulations, $n = 3240$ cells). Cells on the mesh edge were not analysed due to border effect. Circles are Tukey's outliers; lowercase letters indicate statistically different populations (Student's t-test, $p < 0.05$); red lines indicate linear regressions, with Pearson correlation coefficient R and corresponding p-value. (**j**) Model parameter exploration. Colours indicate Person correlation R, with perfect anticorrelation as blue ($R = -1$), perfect correlation in red ($R = 1$), and insignificant correlation ($p > 0.05$) in black. A, normalized cell area; N, cell neighbour number; P, normalized turgor pressure; α^a, dimensionless parameter for flux-wall limit balance; α^s, dimensionless parameter for apoplasmic-symplasmic limit balance; s, cell size at division (**k** to **n**) Schematic explanation of topology-derived turgor pressure heterogeneity. (k and l) Three-cell junctions in a tissue of hexagonal cells are at mechanical

equilibrium with equal tensions and equal wall-wall angles. (m and n) Fewer-neighboured cells have sharper wall-wall angles, a tension that is roughly constant per wall effectively results in mechanical compression due to unequal projected tension distribution (red dash-line arrow) that is balanced by higher turgor pressure build-up (big blue arrow).

Figure 3. Cellular growth rate bifurcates between conditions.
(**a** to **f**) Relative growth rate per hour G of untreated SAM cells between 12-hour interval. (**g** to **l**) Cellular growth rate per hour of oryzalin-treated SAM between 12-hour interval (48 and 60 hours post treatment). (a, b, g and h) Surface projections of untreated or oryzalin-treated SAM at initial time point (a and g) and 12 hours later (b and h); scale bars are 20 μm unless otherwise noted. (c and i) Heat maps of areal relative growth rate per hour. (d and j) Example 4 and 8-neighbored cells during 24-hour growth, with areal normalization at initial time point. Cell contour and relative size (blue for 4-neighbored, red for 8-neighbored) depict the diverging growth trends. Scale bars are as indicated. (e, f, k and l) Box plots of relative growth rate per hour G against cell topology N (e and k) and dot plots of relative growth rate per hour G against normalized cell area A (f and l) (e and f, untreated 11 SAMs, $n = 1491$ cells;

k and l, oryzalin-treated 14 SAMs, $n = 1160$ cells). Note that Tukey's outliers are not plotted, but are included for statistical analyses. Lowercase letters indicate statistically different populations (Student's t-test, $p < 0.05$); red lines indicate linear regressions, with Pearson correlation coefficient R and corresponding p-value.

Figure 4. Model effective parameters that enable recapitulating the untreated and oryzalin-treated growth trends.

(**a** to **c**) Relative growth rate per hour G in simulations with cell division, which corresponds to $\alpha^a = 0.5$ and division size $s +50\%$, and mimics untreated SAM. Cells that just divided (dark blue with 0 growth) are not included in analysis. (**d** to **f**) Growth simulations with no cell division, $\alpha^a = 0.5$, and quadruple wall thickness to mimic oryzalin treatment. (a, d) Heat maps of areal relative growth rate per hour. Scale bars are 10 unit lengths. (b, e) Box plots of relative growth rate per hour G against cell topology N. Lowercase letters indicate statistically different populations (Student's t-test, $p < 0.05$); red lines indicate linear regressions, with Pearson correlation coefficient R and corresponding p-value. (c, f) Dot plots

of relative growth rate per hour G against normalized cell area A. Numbers of repeats see Methods. (**g**) Model exploration to fit oryzalin-treated case. Colours indicate Person correlation R, with perfect anticorrelation as blue ($R = -1$), perfect correlation in red ($R = 1$), and insignificant correlation ($p > 0.05$) in black. Triangles indicate parameter's influence to the flux-wall limit balance. * indicates the same parameter choices ($\alpha^a = 0.5$ and division size s +50%), with Unt indicating dividing, untreated-like simulations and nd for non-dividing; $\Delta\Pi$, osmotic drive; w, wall thickness; A, normalized cell area; N, cell neighbour number; P, normalized turgor pressure; G, relative growth rate; Φ^w, wall extensibility; L^a, cross-membrane water conductivity; L^s, cell-to-cell symplasmic conductivity; size, cell size at division. (**h**) Summary of growth trend inversion according to the flux-wall limits shown in (g).

References

1. Hong L, et al. (2018) Heterogeneity and Robustness in Plant Morphogenesis: From Cells to Organs. *Annu Rev Plant Biol* 69(1):469–495.

2. Kamimoto K, et al. (2016) Heterogeneity and stochastic growth regulation of biliary epithelial cells dictate dynamic epithelial tissue remodeling. *eLife* 5. doi:10.7554/eLife.15034.

3. Ietswaart R, Rosa S, Wu Z, Dean C, Howard M (2017) Cell-Size-Dependent Transcription of FLC and Its Antisense Long Non-coding RNA COOLAIR Explain Cell-to-Cell Expression Variation. *Cell Syst* 4(6):622-635.e9.

4. Long Y, et al. (2017) In vivo FRET-FLIM reveals cell-type-specific protein interactions in Arabidopsis roots. *Nature* 548(7665):97–102.

5. Chubb JR (2017) Symmetry breaking in development and stochastic gene expression. *Wiley Interdiscip Rev Dev Biol* 6(6):e284.

6. Donati G, Watt FM (2015) Stem Cell Heterogeneity and Plasticity in Epithelia. *Cell Stem Cell* 16(5):465–476.

7. Gerdes MJ, et al. (2014) Emerging Understanding of Multiscale Tumor Heterogeneity. *Front Oncol* 4. doi:10.3389/fonc.2014.00366.

8. Eldar A, Elowitz MB (2010) Functional roles for noise in genetic circuits. *Nature* 467(7312):167–173.

9. Watanabe K, Umeda T, Niwa K, Naguro I, Ichijo H (2018) A PP6-ASK3 Module Coordinates the Bidirectional Cell Volume Regulation under Osmotic Stress. *Cell Rep* 22(11):2809–2817.

10. Xie K, Yang Y, Jiang H (2018) Controlling Cellular Volume via Mechanical and Physical Properties of Substrate. *Biophys J* 114(3):675–687.

11. Guo M, et al. (2017) Cell volume change through water efflux impacts cell stiffness and stem cell fate. *Proc Natl Acad Sci* 114(41):E8618–E8627.

12. Dumais J, Forterre Y (2012) "Vegetable Dynamicks": The Role of Water in Plant Movements. *Annu Rev Fluid Mech* 44(1):453–478.

13. Stewart MP, et al. (2011) Hydrostatic pressure and the actomyosin cortex drive mitotic cell rounding. *Nature* 469(7329):226–230.

14. Montel F, et al. (2011) Stress clamp experiments on multicellular tumor spheroids. *Phys Rev Lett* 107(18):188102.

15. Rojas ER, Huang KC (2017) Regulation of microbial growth by turgor pressure. *Curr Opin Microbiol* 42:62–70.

376   16.   Kroeger JH, Zerzour R, Geitmann A (2011) Regulator or driving force? The role of turgor
377   pressure in oscillatory plant cell growth. *PloS One* 6(4):e18549.

378   17.   Ortega JK (1985) Augmented growth equation for cell wall expansion. *Plant Physiol*
379   79(1):318–320.

380   18.   Beauzamy L, Derr J, Boudaoud A (2015) Quantifying Hydrostatic Pressure in Plant Cells by
381   Using Indentation with an Atomic Force Microscope. *Biophys J* 108(10):2448–2456.

382   19.   Vella D, Ajdari A, Vaziri A, Boudaoud A (2012) The indentation of pressurized elastic shells:
383   from polymeric capsules to yeast cells. *J R Soc Interface* 9(68):448–455.

384   20.   Mosca G, Sapala A, Strauss S, Routier-Kierzkowska A-L, Smith RS (2017) On the micro-
385   indentation of plant cells in a tissue context. *Phys Biol* 14(1):015003.

386   21.   Corson F, et al. (2009) Turning a plant tissue into a living cell froth through isotropic growth.
387   *Proc Natl Acad Sci U S A* 106(21):8453–8458.

388   22.   Beauzamy L, Louveaux M, Hamant O, Boudaoud A (2015) Mechanically, the Shoot Apical
389   Meristem of Arabidopsis Behaves like a Shell Inflated by a Pressure of About 1 MPa. *Front*
390   *Plant Sci* 6. doi:10.3389/fpls.2015.01038.

391   23.   Ishihara S, et al. (2013) Comparative study of non-invasive force and stress inference
392   methods in tissue. *Eur Phys J E Soft Matter* 36(4):9859.

393   24.   Weaire DL, Hutzler S (2001) *The Physics of Foams* (Clarendon Press).

394   25.   Cheddadi I, Génard M, Bertin N, Godin C (2019) Coupling water fluxes with cell wall
395   mechanics in a multicellular model of plant development. *PLOS Comput Biol* 15(6):e1007121.

396   26.   Lucas WJ, Ham B-K, Kim J-Y (2009) Plasmodesmata - bridging the gap between neighboring
397   plant cells. *Trends Cell Biol* 19(10):495–503.

398   27.   Kumar NM, Gilula NB (1996) The Gap Junction Communication Channel. *Cell* 84(3):381–388.

399   28.   McLean PF, Cooley L (2013) Protein Equilibration through Somatic Ring Canals in Drosophila.
400   *Science* 340(6139). doi:10.1126/science.1234887.

401   29.   Cosgrove DJ (1993) Wall extensibility: its nature, measurement and relationship to plant
402   cell growth. *New Phytol* 124(1):1–23.

403   30.   Maurel C, et al. (2015) Aquaporins in Plants. *Physiol Rev* 95(4):1321–1358.

404   31.   Klepikova AV, Kasianov AS, Gerasimov ES, Logacheva MD, Penin AA (2016) A high resolution
405   map of the Arabidopsis thaliana developmental transcriptome based on RNA-seq profiling.
406   *Plant J Cell Mol Biol* 88(6):1058–1070.

407   32.   Ortega JKE (2019) Dimensionless Numbers to Analyze Expansive Growth Processes. *Plants*

408   8(1):17.

409   33.   Gibson WT, Gibson MC (2009) Chapter 4 Cell Topology, Geometry, and Morphogenesis in
410   Proliferating Epithelia. *Current Topics in Developmental Biology*, Current Topics in
411   Developmental Biology. (Academic Press), pp 87–114.

412   34.   Péret B, et al. (2012) Auxin regulates aquaporin function to facilitate lateral root emergence.
413   *Nat Cell Biol* 14(10):991–998.

414   35.   Kwiatkowska D (2004) Surface growth at the reproductive shoot apex of Arabidopsis thaliana
415   pin-formed 1 and wild type. *J Exp Bot* 55(399):1021–1032.

416   36.   Serrano-Mislata A, Schiessl K, Sablowski R (2015) Active Control of Cell Size Generates Spatial
417   Detail during Plant Organogenesis. *Curr Biol* 25(22):2991–2996.

418   37.   Willis L, et al. (2016) Cell size and growth regulation in the Arabidopsis thaliana apical stem
419   cell niche. *Proc Natl Acad Sci U S A* 113(51):E8238–E8246.

420   38.   Tsugawa S, et al. (2017) Clones of cells switch from reduction to enhancement of size
421   variability in Arabidopsis sepals. *Development* 144(23):4398–4405.

422   39.   Cutler SR, Ehrhardt DW, Griffitts JS, Somerville CR (2000) Random GFP：：cDNA fusions enable
423   visualization of subcellular structures in cells of Arabidopsis at a high frequency. *Proc Natl*
424   *Acad Sci* 97(7):3718–3723.

425   40.   Stanislas T, Hamant O, Traas J (2017) Chapter 11 - In-vivo analysis of morphogenesis in plants.
426   *Methods in Cell Biology*, Cell Polarity and Morphogenesis., ed Lecuit T (Academic Press), pp
427   203–223.

428   41.   Reuille PB de, et al. (2015) MorphoGraphX: A platform for quantifying morphogenesis in 4D.
429   *eLife* 4:e05864.