Title:

Cellular heterogeneity in pressure and growth emerges from tissue

topology and geometry

Yuchen Long[1,*], Ibrahim Cheddadi[2,4], Vincent Mirabet[1], Gabriella Mosca[3], Mathilde Dumond[1,5],

Jan Traas[1], Christophe Godin[1,2], Arezki Boudaoud[1,*].

1. Laboratoire Reproduction et Développement des Plantes, Université de Lyon, ENS de

Lyon, UCB Lyon 1, CNRS, INRA, INRIA, F-69342, Lyon, France.

2. INRIA/CIRAD/INRA Project-team Virtual Plants, France.

3. Department of Plant and Microbial Biology, University of Zürich, Zollikerstrasse 107, CH-

8008 Zürich, Switzerland.

4. Current address: Univ. Grenoble Alpes, CNRS, Grenoble INP, TIMC-IMAG, 38000

Grenoble, France.

5. Current address: Department of Biosystems Science and Engineering, ETH Zurich,

Mattenstrasse 26, 4058, Basel, Switzerland.

*. Correspondence and requests for materials should be addressed to yuchen.long@ens-

lyon.fr or arezki.boudaoud@ens-lyon.fr

The authors declare no competing financial interests.

Summary

Cell-to-cell heterogeneity prevails in many biological systems, although its origin and function are often unclear. Cell hydrostatic pressure, alias turgor pressure, is essential in physiology and morphogenesis, and its spatial variations are often overlooked. Here, based on a mathematical model describing cell mechanics and water movement in a plant tissue, we predict that cell pressure anticorrelates with cell neighbour number. Using atomic force microscopy, we confirm this prediction in the Arabidopsis shoot apical meristem, a population of stem cells that generate all plant aerial organs. Pressure is predicted to correlate either positively or negatively with cellular growth rate depending on osmotic drive, cell wall extensibility, and hydraulic conductivity. The meristem exhibits one of these two regimes depending on conditions, suggesting that, in this tissue, water conductivity is non-negligible in growth control. Our results illustrate links between local topology, cell mechanical state and cell growth, with potential roles in tissue homeostasis.

Key words

Cellular heterogeneity, hydrostatic pressure, tissue topology, growth mechanics, tissue hydraulics, atomic force microscopy, biophysical modelling.

Highlights

- A mechano-hydraulic model of tissue growth predicts heterogeneity in cell pressure and in cell growth according to local tissue topology.
- Indentation-based measurement of cell pressure in the shoot apical meristem of *Arabidopsis thaliana*, together with a realistic mechanical model of tissue indentation, reveals pressure variations according to cell size and cell neighbour number.
- Cell growth in the Arabidopsis shoot apical meristem either correlates or anti-correlates with cell neighbour number, which can be retrieved by the model using experimentally-based parameters.

Introduction

Cell-to-cell fluctuations are observed in many biological processes like gene expression, signalling, cell size regulation and growth [1–8]. Notably, heterogeneity in cell size and growth rate often prevails and may impact tissue patterning and macroscopic growth robustness [1,2]. Cell volume change is driven by osmosis [9–11] and the resulting intracellular hydrostatic pressure, and is restrained by peripheral constraints — plasma membrane, cytoskeletal cortex, extracellular matrix, or cell wall — in plant cells [12], animal cells [13] including tumorous [14], and microbial cells [15] (Fig. 1a).

Due to high difference between internal and external osmotic potential, cells with rigid cell walls – like in plants, bacteria and fungi – accumulate hydrostatic pressure, alias turgor pressure, often greater than atmospheric pressure (Fig. S1) [12]. Animal cells also accumulate hydrostatic pressure, especially when compacted or contracting [13,14], though to a lesser extent than walled cells. Whereas it is increasingly realized that pressure regulation is crucial for the general physiology, growth and signalling in animal [9,11,13] and plant cells [16–19], pressure remains poorly characterized in multicellular contexts.

In plants, turgor pressure drives cells expansion, which is classically modelled as visco-elasto-plastic process: irreversible expansion occurs when the pressure exceeds a threshold value and the cell deforms reversibly when pressure is smaller, as depicted in the Lockhart-Ortega equation (Fig. 1b) [20]:

$$\frac{1}{V}\frac{\mathrm{d}V}{\mathrm{d}t} = \varphi(P - P^Y)_+ + \frac{1}{E^V}\frac{\mathrm{d}P}{\mathrm{d}t}$$

where V is cell volume, P is turgor pressure, P^Y is the pressure threshold for irreversible expansion, φ is global wall extensibility (akin to inverse of viscosity) and E^V is the cell volumetric modulus, while the $_+$ index indicates that irreversible expansion occurs only if $P > P^Y$. For single cell, turgor pressure positively correlates with total growth rate (Fig. 1b). Other experimental observations in single-cell systems suggest that growth rate and pressure level are not always associated [15,21], making it difficult to assess and understand the link between growth regulation and cellular pressure build-up.

In multicellular context, like the plant shoot apex and sepal epidermis, neighbouring cells grow at notably different rates [1]. This prompts the question, according to the Lockhart-Ortega equation, of whether turgor pressure also varies between neighbouring plant cells. Intuitively, pressure difference should be equalized by plasmodesmata, symplasmic bridges connecting most plant cells [22]. This is supported by the association of plasmodesmata closing (symplasmic isolation) and pressure build-up in specialized cells, such as guard cells or cotton fibres [23–25]. However, pressure gradient was also observed in plant tissues with symplasmic continuity [26], and predicted to be crucial for perception of water availability by roots [27]. In addition, computational models of tissue mechanics suggest that neighbouring cells need to have different pressure to recapitulate tissue arrangement and mechanical status in chemical-treated Arabidopsis epidermis [28] and in Drosophila epithelia [29], although such spatial variation is yet to be demonstrated, and its relation with cell-to-cell growth variability remains elusive.

Here, we explore this issue in a model plant tissue, the epidermis of the *Arabidopsis thaliana* shoot apical meristem (SAM), by combining computational modelling, dimensional analysis, and experimental observations. Our physical model that couples tissue growth mechanics with water movement accounts for our experimental observations of turgor pressure and growth rate variability in different conditions. We show that cell pressure heterogeneity emerges from local tissue topology to influence growth variability. Pressure level, however, is not a direct predictor of growth rate, and dimensional analysis of the model reveals that growth behaviour depends on the extent of the osmotic drive and on the balance between wall extensibility and hydraulic conductivity. Together, we propose a link between cell topology and cell hydro-mechanical status that may be involved in tissue homeostasis.

Results

A mechanical-hydraulic model predicts pressure heterogeneity emerging from tissue topology

Earlier tissue models [28,29] retrieved intracellular pressure from static tissue geometry or required differential osmotic pressure, whereas a recent model by Cheddadi et al. proposed that

both pressure and growth emerge from the coupling between cell wall mechanics and classical

plant hydraulics [30] in a tissue with hexagonal topology, i.e. with every cell having six

neighbours. We therefore tested, based on this model, the consequences of unequal neighbour

numbers on the mechanical status of the tissue, and notably on pressure (Fig. 1). We expanded

on the 2D vertex model to cell walls in a multicellular tissue with more realistic 2D polygonal

geometry. Generalizing the Lockhart-Ortega equation of visco-elasto-plastic 1D growth of

single cell (Fig. 1b) [20], each cell wall has a thickness w and behaves as an elastic material of

modulus E when wall tension induced by turgor pressure P is low enough, more specifically

when the elastic strain ε (reversible deformation) is lower than a threshold ε^Y. When tension in

the wall exceeds the corresponding threshold $E\varepsilon^Y$, the cell wall undergoes irreversible expansion,

akin to visco-plastic flow, with an extensibility (effective viscosity) Φ^w (Fig. 1c). Water flux

from extracellular space is proportional to $L^a \Delta \Psi$, where $\Delta \Psi = \Delta \Pi - P$ is the cross-membrane

water potential difference (external pressure is assumed to be zero; for parsimony we set $\Delta \Pi$

constant) and L^a is the conductivity of the cell membrane. Intercellular water redistribution via

plant plasmodesmata [31], animal gap junctions, or cytoplasmic bridges [32,33] is driven by

intercellular differences in turgor pressure P, with a conductivity L^s. We did not prescribe turgor

pressure, instead letting it emerge from local mechanical and hydraulic interplays (see STAR

Methods for detailed model description).

To prescribe cell divisions, we utilized the recently introduced Willis-Refahi rule that derives

from experimental data in the SAM: cells divide according to their size and to the size increment

since the previous division [34]. The simulations recreate distributions of neighbour number

(topological distributions) that are similar to those observed in the SAM (Fig. 1e).

We set the model parameters based on classical measurements from Boyer [35] and Cosgrove

[36] (Table 1). We found turgor pressure to be heterogeneous, with a clear anticorrelation with

topology: cells with less neighbours have relatively higher pressure (Fig 1f, 3 simulations, cell

number $n = 1535$, Pearson correlation coefficient $R = -0.57$, $p < 10^{-100}$).

Initial parameter	Value	Initial parameter	Value
Unit length	10 μm	Apoplasmic	
Cell height h	10 μm	conductivity L^a	3.4×10^{-10} m MPa^{-1} s^{-1}
Cell wall thickness w	1 μm	Symplasmic	
Elastic modulus E	33.8 MPa	conductivity L^s	3.4×10^{-10} m MPa^{-1} s^{-1}
Wall extensibility Φ^w	2.4×10^{-5} MPa^{-1} s^{-1}		
Strain threshold ε^Y	0.1	Osmotic pressure $\Delta\Pi$	0.7 MPa

Perturbation	Modified parameter	Perturbation	Modified parameter
$\alpha^s = 0.1$	$L^s = 3.8 \times 10^{-11}$ m MPa^{-1} s^{-1}	$\theta = 10/3$	$\Delta\Pi = 1$ MPa
$\alpha^s = 0.9$	$L^s = 3.1 \times 10^{-9}$ m MPa^{-1} s^{-1}	$\theta = 20/3$	$\Delta\Pi = 2$ MPa
$\alpha^a = 0.1$	$\Phi^w = 2.2 \times 10^{-4}$ MPa^{-1} s^{-1}	$\theta = 10$	$\Delta\Pi = 3$ MPa
$\alpha^a = 0.9$	$\Phi^w = 2.7 \times 10^{-6}$ MPa^{-1} s^{-1}	$\theta = 40/3$	$\Delta\Pi = 4$ MPa

Table 2. Parameters used for the untreated model. (Upper) The initial parameters correspond to $\alpha^s = 0.5$, $\alpha^a = 0.5$, $\theta = 7/3$ in Figure 1f, which are based on measurements by Boyer (1985) [35] for α^a and θ, while α^s data is not available thus is conservatively ascribed an intermediate value of 0.5. (Lower) Non-dimensional parameter exploration around the reference values (initial parameters, except , $\theta = 20/3$) and the corresponding modified parameters. Note that $\alpha^a = 0.9$ corresponds to measurements by Cosgrove (1985) [36], compiled by Ortega (2019) [37].

In order to test the robustness of this result, we explored the parameter space of the vertex model. Analytical exploration in a two-cell system [30] had showed that systems dynamics is mostly controlled by three dimensionless parameters. The first, α^s, compares intercellular water conductivity to total (intercellular and transmembrane) conductivity,

$$\alpha^s = \frac{A^s L^s}{A^s L^s + A^a L^a}$$

with A^a the average surface of a cell, A^s the average surface of a cell in contact with neighbouring cells, L^a the conductivity (per unit surface) of the plasma membrane, and L^s the conductivity (per unit surface) due to plasmodesmata. The second, α^a, compares the limitation of growth by transmembrane conductivity to the combined limitation of growth by cell wall extensibility and transmembrane conductivity,

143
$$\alpha^a = \frac{L^a}{L^a + \Phi^w \dfrac{h^2}{4w}}$$

144 where h and w are cell height and cell wall thickness. Both α^s and α^a are bound between

145 0 and 1. The third parameter, θ, assesses the osmotic drive by comparing the cross-membrane

146 osmotic pressure difference, $\Delta\Pi$, and a representative threshold pressure for growth to occur,

147 P^Y,

148
$$\theta = \frac{\Delta\Pi}{P^Y}$$

Contrary to the Lockhart-Ortega model that was formulated at cell scale, our model accounts

for specific cell wall geometry and mechanical properties. Accordingly, we express the

threshold in terms of a yield strain ε^Y and the threshold pressure depends on cell geometry and

topology. We found empirically [30] that half the threshold pressure of a single hexagonal cell

provides a good order of magnitude for the threshold pressure in the multicellular model and

hence use

155
$$P^Y = \frac{w}{2R \cos(\pi/N)} E \varepsilon^Y$$

where R is a representative cell size and the number of cell neighbours is $N = 6$ (Fig. S1).

The tissue globally grows if θ is greater than 1. Our first results, obtained with $\alpha^s = 1/2$,

$\alpha^a = 1/2$, and $\theta = 7/3$, corresponded to a balance between mechanical and hydraulic

limitations to growth (Fig. 1f). We explored the parameter space by considering 4 values of θ

reaching up to 40/3 (Fig. 1i, S2). As will be clarified below (see last subsection), we considered

$\theta = 20/3$ as a reference value (Fig. 1g). We then decreased and increased either α^s or α^a

down to 0.1 or up to 0.9, respectively (Fig. 1i, S2). Note that the values 0.5 and 0.9 of α^a span

available measurements of extensibility and conductivity [35,36] (Table 1). Next, we tested the

effect of cell divisions by arresting them (Fig. 1h). In all cases, we recovered the turgor–

neighbour-number anticorrelation (Fig. 1f-i, S2), demonstrating that topology-related pressure

heterogeneity is a robust behaviour of the model.

In simulations, like in SAM surface, cell neighbour number and size are coupled (Fig. 1f-i, S2).

Consistently, cell-specific turgor pressure anticorrelates with normalized cell area (Fig. 1f-I,

S2). To uncouple cell size and topology, we modified the initial state of our simulation to have

four-neighboured cells bigger than eight-neighboured cells, and found that higher turgor

pressure accumulated in four-neighboured cells, not in the smaller cells (Fig. 1j), showing that

turgor pressure heterogeneity emerges from tissue topology rather than cell size differences.

This can be explained by local topology, which determines cell wall angles and the subsequent

tension distribution at each tricellular junction (Fig. 1k, l): in our model, wall stress and strain

above the growth threshold are relaxed at a rate limited by wall extensibility and hydraulic

conductivity. Therefore, stress and strain are relatively homogenised in non-dividing

simulations (Fig. 1h), and the sum of wall tension at each tricellular junction (vertex) mostly

depends on the angles between walls. The vertex between three hexagonal cells with 120°

internal angles has a sum of tension at zero (Fig. 1k). Fewer-neighboured cells have sharper

internal angles, so the sum of tension at vertex is greater towards the cell interior, creating

additional inward compression and prompting higher pressure build-up at equilibrium (Fig. 1l).

In dividing simulations, new walls do not bear stress right after division because they form from

cell interior. These new walls are gradually strained due to mesh growth but do not yield

(expand) to release stress before reaching the threshold (Fig. 1f, g). Consequently, cell division

keeps the wall stress from homogenizing and dampens the turgor–neighbour-number

anticorrelation, compared to non-dividing simulations (Fig. 1g).

Altogether, our results imply that local hydrostatic pressure heterogeneity does not require

differential cellular osmotic pressure [28] in a growing tissue, and predict a topological origin

of pressure variability.

190 Atomic force microscopy reveals heterogeneous turgor pressure in Arabidopsis shoot
191 apical meristem

To test predictions, we built upon recent advances in atomic force microscopy (AFM) that

enabled non-invasive turgor pressure retrieval utilizing indentation force-displacement and

surface topography in living plant cells (Fig. 2) [38–42]. We used a pressurized thin-shell model

to deduce the turgor pressure value from the AFM-measured force-displacement curves, which

are influenced by turgor pressure, cell wall mechanical properties, cell 3D geometry [38], and

may reflect mechanical properties at different sub- to supra-cellular scales according to

indentation depths (Fig. 2b-e). We applied AFM measurements to the Arabidopsis SAM

epidermis, a system featuring substantial growth heterogeneity [1]. We included untreated soil-

grown SAMs (Fig. 3) and a conceptually simpler model SAM co-treated with

naphthylphthalamic acid (NPA), a polar auxin transport inhibitor that induces pin-formed

SAMs, and oryzalin, a microtubule-depolymerizing drug that blocks cell division but permits

continuous, isotropic growth; hereafter referred to as "oryzalin-treated SAMs" (Fig. 4) [28].

Walls between cells are often curved in growing oryzalin-treated SAMs, suggesting that

neighbouring cells have different turgor pressure [28] in a growing tissue.

Figure 2 illustrates the AFM measurement pipeline, where we determined SAM surface

topography with AFM (Fig. 2f) and performed indentations near the cell centre to have near-

perpendicular indentation and minimize any bias due to surface slope (Fig 2g). Physical values

required to deduce pressure using the published formula [39,40], except cell wall thickness

separately measured by electron microscopy (EM), are simultaneously determined by AFM

scan and indentation. Specifically, both mean curvature κ_M and Gaussian curvature κ_G of outer

cell walls directly contribute to the capacity to sustain turgor pressure [38,39,43], and were

determined from AFM scan topography (Fig. 2f). As previously suggested in tomato SAM [44],

outer periclinal wall thickness is not very variable between cells or within cells in Arabidopsis

SAM, with untreated meristem $t_u = 179 \pm 7$ nm (mean ± standard error of mean, SEM) and

oryzalin $t_o = 742 \pm 29$ nm (Fig. S3). Based on previous work [45], we used the indentation

depth range 0 to 150 nm to determine apparent Young's modulus of cell wall, E (Fig. 2b, e). We

determined indentation stiffness, k, using the depth range 0.3 to 0.5 μm for untreated and 1.1 to

1.5 μm for oryzalin-treated meristems (Fig. 2c, e); these ranges were chosen so that depth is

greater than wall thickness (to minimize cell wall contributions to k) and the deformed region

remains smaller than cell size (to minimize the contributions of neighbouring cells to k, Fig. 2d,

e), so as to be in the validity range of the pressurized shell model [40]. Note that these values

of depth for untreated SAMs differ from the larger values (1 to 2 μm) used in preliminary

experiments [41], which were interpreted using supracellular curvature (unlike here) and were

chosen to reveal supracellular pressure (averaged over many cells and possibly over cell layers)

for comparison with the large-scale pressure obtained using indentation with a large flat tip (100 μm diameter). To verify the validity of the depth ranges used here, we tested influence of tissue context on indentation readout [46] by numerical simulations of indentations on realistic 3D meshes accounting for pressurized epidermal cells (Fig. 2m-o''); in conditions corresponding to experiments, the deformation of neighbouring cells was negligible, demonstrating that the depth range enables cell-level measurements.

Cell-specific AFM indentations on seven untreated and nine oryzalin-treated SAMs revealed that surface wall curvature, Young's moduli, cell apparent stiffness and the deduced turgor pressure are all markedly heterogeneous across SAM epidermis (Fig. 3, 4). We computed intracellular variability of quantities measured at subcellular scale (which excludes curvature, measured at cell scale), and found that apparent Young's modulus E and stiffness k show significantly higher variability than pressure P (as quantified by intracellular coefficients of variation, Fig. 3i, 4i), which agrees with previous work [40], and supports deduced P being a cellular property. Additionally, we found that E shows relatively comparable subcellular and intercellular variability, as previously observed [45,47], whereas P show significantly bigger heterogeneity between than within cells (Fig. 3j, 4j). Deduced pressure, P, is relatively insensitive to variability in thickness as described in [40], and change in thickness by $1\times$ standard deviation (SD) only alters the coefficient of variance (CV) of P by 3% and 9% for untreated and oryzalin-treated samples respectfully, both significantly smaller than the intercellular P variability (untreated 21% and oryzalin 15%). All these indicate that P deduction is cell-specific, and that variability in cell wall mechanics does not account for deduced pressure heterogeneity.

Osmotic treatments reveal large values of osmotic pressure in the shoot apical meristem

Based on AFM, we find that deduced turgor pressure is heterogeneous, with averages values per meristem of 2.61 ± 0.03 MPa and 1.21 ± 0.34 MPa (mean $\pm$ SEM) in untreated and oryzalin-treated meristems, respectively (Fig. S3). These values appear high with respect to the classical 0.3 to 1 MPa range found in most other plant tissues, though values of up to 5 MPa were measured in guard cells of other species using the pressure probe [48].

As neither the classic pressure probe nor the pico gauge [49] can be applied to cells as small as

in the shoot apical meristem, we assessed quantitatively the values of turgor pressure P, by

using an incipient plasmolysis assay to determine the osmotic pressure of SAM, and found that

untreated meristems had a rather invariable osmolarity of about 0.5 Osm, slightly higher than

values reported in tomato SAM [50], while oryzalin samples showed a wider variability of 0.6

to 1.0 Osm (Fig. S4). The corresponding values of osmotic pressure, 1.2 MPa for untreated and

2.0 MPa for oryzalin treated, are comparable to the values of AFM-deduced turgor (2.6 MPa

and 1.2 MPa, respectively) (Fig. S3, S4). The quantitative agreement is less good in the case of

untreated meristems, possibly due to tissue tension in the epidermis which could artificially

increase the deduced pressure due to contributions from inner layers. We note however that

tissue tension would smooth out intercellular differences and would thus not contribute to the

cell-to-cell heterogeneity of pressure. An alternative explanation would be that solute

penetration differs between untreated and treated SAMs. Altogether, the values of turgor found

with AFM are in semi-quantitative agreement with the values of osmotic pressure deduced from

incipient plasmolysis.

Cell pressure in shoot apical meristem anticorrelates with local topology and size

Next, we tested model predictions and found that, indeed, AFM-determined cellular pressure

anticorrelates with the number of epidermal cell-neighbours N, or local topology (7 SAMs, n =

503 cells, R = -0.16, p = 10^{-4}, Fig. S5). With the linear relationship between cell area and

neighbour number (Fig. 3d) due to fundamental geometrical constraints in compact tissues [51],

we also identified strong anticorrelation between pressure and cell size A (R = -0.19, p = 10^{-5}).

We focused on cell-to-cell turgor variations and we removed variations between SAMs by

normalizing cellular pressure to the average pressure per SAM; given that untreated SAMs have

unvaried average turgor values (Fig. S3), the anticorrelation is conserved for pressure-versus-

topology (R = -0.16, p = 10^{-4}) and versus-size (R = -0.23, p = 10^{-7}) (Fig. 3h-h'').

Encased in rigid cell walls, plant cells seldom exchange neighbours, and the main source of

topological change is via division, where dividing cells tend to lose neighbours and cells

adjacent to the division plain tend to gain neighbours [52]. We therefore considered oryzalin-

treated SAM, in which cell divisions are arrested while growth is continuous, and recovered

similar anticorrelation for pressure against neighbour number (9 SAMs, n = 202 cells, R = -

0.19, p = 0.008) and against cell area (R = -0.48, p = 10^{-12}) (Fig. S5). Normalization further

enhanced the pressure-versus-topology anticorrelation (R = -0.31, p < 10^{-5}) while preserving

the pressure-size trend (R = -0.33, p < 10^{-5}) (Fig. 4h-h''), indicating that the pressure-topology

and pressure-size relationships are robust system-level trends, regardless of the absolute values.

We then examined whether such trends may be caused by trends in cell wall thickness, t, or

modulus, E, or by trends in stiffness, k. There is no correlation between t and cell size (R = -

0.10, p = 0.65, Fig. S3). So t only introduces small unbiased noise, and it cannot account for

the P heterogeneity. We found that E and k showed no significant correlation to cell neighbour

number N in untreated SAMs (Fig. 3e-f'), indicating that they introduce no systemic bias to P

heterogeneity. In oryzalin-treated SAMs, however, both E and k showed weak positive

correlation to N (Fig. 4e-f'), which is opposite to the P vs N anticorrelation (Fig. 4h'). This is

interesting, because the higher P in N = 4 cells cannot be explained by lower E and k ("softer"

cell). This indicates that, although feedback from shape on E may occur in oryzalin-treated

meristems, P heterogeneity is not a direct consequence of feedbacks from wall tension or cell

geometry/topology on wall stiffness.

Moreover, we found that turgor pressure heterogeneity may be removed when sample is

osmotically challenged: the same SAM shows heterogeneous pressure when turgid and

homogeneous pressure when at intermediate turgidity (Fig. S4). This shows that the AFM

approach is not technically biased by tissue topology and/or cell size.

Finally, we used cell side wall convexity as a proxy for differences in turgor, because cells with

higher pressure would be expected to bulge out into cells with lower pressure (Fig. 4q-q'') [53].

We constructed a weighed convexity index (wCI) to quantify convexity (Fig. 3o-q, 4o, p), and

found that convexity significantly anticorrelates with number of neighbours, in agreement with

qualitative observations in oryzalin-treated meristems [28], and in agreement with the pressure

trends.

Altogether, our data indicate that non-random turgor pressure heterogeneity establishes in

tissues with static topology (no neighbour number change) or dynamic topology (neighbour

numbers change due to division). Although tissue topology is sufficient to explain pressure heterogeneity, we do not exclude a role of other biological parameters and/or sources of noise in pressure variations, as suggested by weaker correlations in experiments (Fig. 3h', 4h') than in simulations (Fig. 1g, h).

Realistic mechanical models of tissue indentation are inconsistent with homogeneous pressure

Whereas the aforementioned pressure deduction is based on a model utilizing local cell shape [39], Mosca et al. [46] proposed that cell packing also contribute to the indentation stiffness. We therefore implemented realistic indentation using a thin-shell indentation finite element method (FEM) model following Mosca et al. [46]. We constructed two realistic templates (untreated- and treated-like) from confocal images; the surface was meshed and the outermost cell layer was constructed to represent the epidermis, with uniform values of cell wall thickness taken from our EM-based measurements and same cell dimensions as in confocal images; all cells were inflated by turgor pressure (Fig. 2m). We used the null hypothesis that all cells were inflated by uniform turgor pressure (2 MPa, rounded from experimental values), and performed indentations on the exact corresponding cells indented experimentally from these two templates, excluding cells at periphery of the template to avoid boundary effects. We chose indentation near the cell centre to have near-perpendicular indentation, like in experiments. We recovered values of apparent stiffness k that have the same magnitude as experimental values: untreated-like $k_{\mathrm{FEM}} = 14.9 \pm 0.1$ N/m (mean $\pm$ SEM, compared to $k_{\mathrm{AFM}} = 12.9 \pm 0.2$ N/m), treated-like $k_{\mathrm{FEM}} = 38.2 \pm 1.0$ N/m (compared to $k_{\mathrm{AFM}} = 14.5 \pm 0.6$ N/m). However, we found that the k variability differs from experiments, notably in untreated SAMs (Fig. 3k, 4k). Closer inspection revealed that cell surface mean curvature κ_{M}, which is directly linked to P, also showed strong discrepancy between simulation and experiment (Fig. 3l, 4l). Accordingly, the null hypothesis of uniform turgor is not consistent with experimental data. We corrected such discrepancy based on the theory of thin shells (apparent stiffness mainly depends on these two parameters following $k \sim \pi P \kappa_{\mathrm{M}}^{-1}$, see [39]) by rescaling κ_{M} and P according to their experimental variability, and recovered the heterogeneous distribution of k determined by AFM (Fig. 3m, 4m). Conversely, correction of k and κ_{M} also recovered the experimentally deduced P

distribution. This indicates that the discrepancy between FEM and AFM indentations can be

fully explained by the heterogeneity in pressure (and subsequently curvature), and the

contribution of cell packing to the measured variability is negligible with our depths of

indentation.

Two types of correlations between cell growth and local topology or size

Next, we monitored areal growth rate of SAM epidermal cells by time-lapse confocal

microscopy. As observed previously, untreated SAMs exhibited slower growth in the centre,

where stem cells reside, than the surrounding cell [54] (Fig. S5). In addition, cellular growth

rate anticorrelates with neighbour number (11 SAMs, $n = 1491$ cells, $R = -0.15$, $p = 10^{-8}$; Fig.

5e) and cell size (Fig. 5f; $R = -0.33$, $p = 10^{-38}$), supporting previous reports that smaller cells in

SAM grow faster [34,55], and suggesting that higher turgor pressure in fewer-neighboured cells

associates with faster growth. In oryzalin-treated SAMs, however, the fewer-neighboured and

small cells grew slower (14 SAMs, $n = 1160$ cells; Fig. 5k, neighbour number $R = 0.20$, $p = 10^{-11}$;

Fig. 5l, cell size $R = 0.16$, $p = 10^{-8}$). This suggests that higher turgor pressure associates with

either faster or slower growth depending on conditions. Although seemingly a small shift, this

negative-to-positive slope change of local growth heterogeneity captures a strong qualitative

inversion of growth behaviour (Fig. 5d, j). Accordingly, smaller cells expand more than larger

cells in untreated SAMs, which may contribute to cell size homeostasis, a phenomenon that

would be absent in oryzalin-treated SAMs.

Heterogeneity in growth rate is patterned according to the balance between wall
extensibility, tissue conductivity and osmotic drive

We further explored the growth trend in our vertex model. Based on the parameter exploration

on dividing mesh aimed to reproduce untreated SAM behaviour, we found both positive and

negative correlations between growth rate and cell neighbour number (Fig. S2). This indicates

that growth trend is sensitive to parameters that alter the balance between water flux and wall

expansion (governed by the non-dimensional parameter α^a) and to the ratio between osmotic

pressure and yield pressure ($\theta = \Delta\Pi / P^Y$). This can be rationalized by examining the relative

growth rate G of a cell according to the Lockhart model (see [30]).

$$368 \qquad G = \alpha^a \Phi^w \frac{h}{2w} (\Delta\Pi - P^Y)$$

Both the prefactor $\alpha^a \Phi^w h / 2w$ and the yield pressure P^Y decrease with cell size R and

neighbour number N, while $\Delta\Pi$ is constant. Consequently two regimes are expected: when the

osmotic drive θ is smaller than a threshold θ^T, G is dominated by the variations of P^Y,

therefore G increases with cell size and with neighbour number, which corresponds to the trend

in oryzalin-treated meristems; when $\theta > \theta^T$, the variations of P^Y are negligible, so G follows

the prefactor and decreases with cell size, which corresponds to the trend in untreated meristems.

These two regimes occur whatever the value of α^a, and the threshold value θ^T increases with

increasing α^a. In a tissue context, many global parameter shifts can invert growth trend

through changes in these dimensionless parameters (Fig. 6k, l).

Overall, the model retrieves untreated SAM trends if transmembrane conductivity partially

limits growth (α^a not too large), symplasmic conductivity is not on par with transmembrane

conductivity (α^s not too large) or if the osmotic drive is sufficient (the ratio of osmotic to yield

pressure is large enough). Conservatively, we chose equal contribution by flux and wall in our

model ($\alpha^a = 0.5$), and increased osmotic pressure to 2 MPa, a value comparable to the

experimental measurements: Figures 6a-c show that the fewer-neighboured and smaller cells

grow faster (3 simulations, $n = 1496$; G vs N, $R = -0.10$, $p = 10^{-4}$; G vs A, $R = -0.76$, $p < 10^{-100}$).

We then attempted to reproduce oryzalin-treated behaviour, guided by the experimental

observations that, besides stopping cell division, oryzalin treatment also yields higher osmotic

pressure (1.6-fold) and drastically thicker cell walls (4-fold) (Fig. S3, S4). We found that both

stalling division and increasing osmotic pressure failed to invert the growth trend in the model,

as expected, while doubling and quadrupling wall thickness inverted the correlation of growth

rate to neighbour number and cell size (Fig. 6d-j; S6), like in oryzalin-treated SAM. Combining

higher osmotic pressure and thicker wall revealed that quadrupling wall thickness can robustly

trigger growth trend inversion (Fig. 6j, S6).

Effectively, changing osmotic pressure and wall thickness alter the wall–flux limitation balance

of the system (Fig. 6j-l). Higher osmotic pressure induces faster water influx (reduced limitation

by hydraulics). The extra volume strains the walls to accumulate stress farther beyond the

threshold, which is relaxed by wall yielding (growth) with extensibility as the rate limit.

Meanwhile, wall thickening reduces wall stress and strain towards the threshold for expansion,

effectively reducing the mechanical drive of growth and increasing the weight of water

permeability in limiting growth. We do not exclude other possible parameter changes triggered

by oryzalin treatment, like water conductivity and wall synthesis rate, that also contribute to the

wall–flux limitation balance. Nevertheless, implementing the observed cell wall thickening in

the model is sufficient to explain the observed growth rate inversion from untreated to oryzalin-

treated scenario.

DISCUSSION

In this study, we modelled the growth of a plant tissue by coupling tissue mechanics and tissue

hydraulics. This generalizes previous models focusing only on mechanics [56–58]. In this

model, both cell growth and turgor pressure emerge from mechanics and hydraulics. Each of

these parameters can be controlled by genetic and biochemical inputs, and small uniform

changes in these biological inputs can enable drastic shifts of system behaviour and the final

cell size distribution (Fig. 6m). We predicted that a broad distribution of neighbour number

leads to heterogeneity in cell growth and in turgor pressure, even when hydraulics has a minor

contribution to the control of growth. We verified this prediction in the context of the shoot

apical meristem (SAM) of *Arabidopsis thaliana*. It remains to be seen whether our results apply

to other plant tissues, to animals or whether this is specific to the SAM.

In order to test our predictions and deduce pressure in the SAM, we combined a recently

developed indentation-based approach [40] with FEM-based realistic mechanical models of

indentation [46]. We found values of turgor in the range 1-3 MPa range, higher than the range

0.2-1 MPa generally measured in plant tissues [18], though values of up to 5 MPa were

measured in guard cells [48]. For instance, the Arabidopsis root epidermis has a turgor of about

0.4 MPa, as measured with the pressure probe [59]; the Arabidopsis leaf epidermis has a turgor

of about 1-2 MPa, as deduced form indentation and mechanical modelling [60]. Accordingly,

we speculate that the SAM function might require relatively high turgor. Another specificity of

the SAM could be a relatively low transmembrane conductivity, as most aquaporin (channel

protein allowing rapid transmembrane water flux [61]) isoforms had significantly lower

expression in inflorescence than in other fast growing tissues like stem and root [62] (with a

reduction up to ~80%). Nevertheless, a wide range of water conductivity can produce pressure

heterogeneity, including very high conductivity (i.e. $\alpha^a = 0.9$ [36]), and the agreement between

model predictions and experimental measurements suggest that tissue hydraulics has a

(possibly small) contribution to limiting growth in the SAM. Similarly, altered expression of

the aquaporin PIP2;1 delays the emergence of lateral roots [63], also hinting to a developmental

role for hydraulic conductivity. Altogether, we propose that tissue mechanics and hydraulics act

in parallel with the established genetic regulations in SAM cell growth.

We found that both cell growth and turgor pressure are heterogeneous in the SAM. It has already

been reported that smaller cells [55] or the smaller of two sister cells [34] grow faster in the

SAM of untreated and NPA-treated plants, respectively. We found the same trend in untreated

SAM, with an inversion in oryzalin-treated SAM. Such inversion is not irrelevant to normal

development: for example, a growth variability switch is observed during sepal development,

where clones of smaller and often fewer-neighboured cells grow faster in young sepals then

slower in older sepals, effectively shifting from homogenizing to amplifying cell size variability

[64]. In untreated SAM, small cells grow faster than big cells, possibly contributing to tissue

homeostasis. Finally, irrespective of the conditions, we find that turgor pressure is smaller in

big cells in experiments, which might contribute to reducing mechanical stress in the cell wall

in these cells [65]; this would act in parallel with the mechanism proposed in the context of leaf

epidermal cells, based on cells adopting puzzle shapes that limit cell wall stress [65].

Our results point towards a link from cell topology (number of neighbours) to cell mechanical

status. This might also be relevant to animal epithelia [29], though this appears unexplored

experimentally. Feedbacks from cell mechanics to cell topology are more established: cell

division and thus number of neighbours can be oriented by mechanical stress in animals and in

plants [66–69]. Since tissue topology is highly conserved in many biological systems [70], we

propose that pressure heterogeneity may emerge in compact tissues with polygonal cells [63]

and non-instantaneous water movement, due to the adjustment to reconcile local mechanical

and hydraulic conditions.

Finally, we note that heterogeneous patterns may not always be stochastic [71]. The emergent

heterogeneity of local growth and hydrostatic pressure is coupled with the characteristic yet

dynamic tissue topology [51,52] (Fig. 6m), all based on stringent rules and likely underlies

morphogenesis in compact tissues. With the discovery of many cell-size-dependent transcripts

[3,72], our model proposes another source for non-random variability in a tissue.

STAR Methods

Plant materials, treatments and growth conditions

Arabidopsis thaliana GFP-LTi6b (ecotype WS-4) reporter line and ecotype Col-0 was used[73].

Untreated inflorescence meristems were obtained from soil-grown plants, first in short-day (8

h light 20°C / 16 h dark 19°C cycle) for 3 to 4 weeks then transferred to long-day (16 h light

20°C / 8 h dark 19°C cycle) for 1 to 2 weeks to synchronize bolting. Oryzalin-treated

inflorescence meristems were obtained from plants grown on custom-made Arabidopsis

medium [74] (Duchefa) supplemented with 1% agar-agar (Merck) and 10 µM N-1-

naphthylphthalamic acid (NPA, Sigma-Aldrich/Merck) for 3 weeks. Pin-formed inflorescence

meristems from NPA medium were immersed in 10 µg/mL oryzalin (Sigma-Aldrich/Merck)

twice (3 h duration, 24 h interval) [28]. For mechanical measurements and time-lapse confocal

imaging, meristems were mounted on Arabidopsis apex culture medium (ACM) [74] with 2%

agarose and 0.1% plant preservation mixture (PPM, Plant Cell Technology) to prevent

contamination, and cultivated in long-day condition.

Atomic force microscopy

Untreated meristems (dissected, with most late stage-2 floral primordia removed to prevent

blocking of the cantilever) and oryzalin-treated meristems were mounted on ACM (2% agarose,

0.1% PPM) the night before. Drops of 2% low melting agarose (Duchefa) were applied around

the lower parts of meristems for mechanical stabilization. For oryzalin-treated meristems, 72 h

post-treatment meristems were measured.

AFM indentations were performed as in Beauzamy et al., 2015 [40]. Specifically, a BioScope

Catalyst model AFM (Bruker) operated with the NanoScope software (version 9.1, Bruker),

under a MacroFluo optical epifluorescence macroscope (Leica), was used. All measurements

were done with customized 0.8 μm diameter spherical probes mounted on silicon cantilevers

of 42 N/m spring constant (SD-Sphere-NCH-S-10, Nanosensors). Cantilever deflection

sensitivity was calibrated against a clean sapphire wafer submerged in water before each session.

Meristems were submerged in water during AFM measurements. PeakForce QNM mode was

used to record sample surface topography and cell contours (aided by the stiffness difference

between periclinal and anticlinal cell walls on DMT modulus maps) in overlapping square tiles

of 30×30 to 50×50 μm^2 (128×128 pixels). Force curves were obtained by the point-and-shoot

mode of the NanoScope software, with at least 3 locations chosen near the barycentre of each

cell, and 3 consecutive indentations per location, making at least 9 force curves per cell.

Approximately 10 μN maximum force was applied during each indentation, corresponding to

approximately 1 μm indentation depth.

For hyperosmotic treatments, oryzalin-treated meristems were mounted in Petri-dishes on

Patafix (UHU), then the gap between Patafix and sample base was quickly sealed with bio-

compatible glue Reprorubber-Thin Pour (Flexbar) for stabilization. After the glue solidified

(less than 2 minutes), samples were submerged in liquid ACM containing 0.1% PPM. Samples

were first measured in liquid ACM (plus 0.1% PPM), then submersion medium was changed

to ACM plus desired concentration of NaCl (plus 0.1% PPM) by first rinsing with 3~5 mL

target solution, then soaked in target solution for 5 minutes before AFM measurements. Each

new measurement per solution change took around 30 minutes.

Force curve analysis

Turgor pressure was determined as previously reported [40]. Specifically, cell wall elastic

modulus (Hertzian model, 1~10% maximal indentation force) and cell apparent stiffness (linear,

75~99% maximal indentation force) were retrieved from each force curve by the NanoScope

Analysis software (version 1.5, Bruker). Quality of force curves were checked empirically and by the fit coefficient of determination $r^2 > 0.99$. Cells with only low quality force curves were not analysed. Cell surface curvatures (mean and Gaussian) were estimated from AFM topographic images, with the curvature radii fitted to the long and short axes of each cell. Turgor pressure was further deduced from each force curve (100 iterations) with the electron-microscopy-determined cell wall thickness 190 nm for untreated and 740 nm for oryzalin-treated SAMs, and cell-specific turgor pressure is retrieved by averaging all turgor deductions per cell.

For cell registration, confocal stacks of each meristem were obtained prior to AFM measurements by an LSM 700 confocal (Carl Zeiss). Surface projection of *GFP-LTi6b* signal was generated by the software MerryProj [75], then rescaled and rotated (affine transformation) to overlay the AFM image tiles. The resulting surface projection image was used to generate cell contour image of the whole meristemic surface using morphological segmentation plugin [76] for the software ImageJ (https://fiji.sc/), while the relative positions of each AFM indentation location is then registered onto the cell contour image, along with cellular geometrical and topological analyses, using the NanoIndentation plugin (version alpha) for ImageJ [77].

Since each meristem had different turgor pressure range, cellular turgor pressure was normalized to the average of each meristem for comparing cell-to-cell turgor pressure heterogeneity without meristem-specific effects.

Electron microscopy

For serial block-face imaging SEM (SBF-SEM), plants were grown in vitro on medium containing the auxin transport inhibitor NPA (Naphtalene Phtalamic Acid) to generate stems with naked meristems and were locally treated for 48h with the microtubule depolymerizing drug oryzalin (Sigma) in lanolin at a concentration of 2 μg/μl [78]. These plantlets were subsequently taken off the inhibitor and left to regenerate for 48h on normal Arabidopsis medium. Meristems with young organ primordia were fixed in 0.5% glutaraldehyde (in demineralized water), from 25% Sigma stock in Microscopy Facility lab. The plantlets were

left at room temp for 2h in an eppendorf and rinsed 1x in water before post fixation and de-

hydrating and embedding in Spurr's epoxy as described in [79]. The samples were then

sectioned and viewed in a Zeiss Merlin SEM [79].

For standard transmission electron microscopy fixed meristems of soil grown plants were

embedded in Spurr's resin and sectioned before viewing in a Jeol 2100F (at the Centre

Technologique des Microstructures, UCBL, Lyon).

Time-lapse confocal microscopy

Untreated (dissected) and oryzalin-treated meristems were mounted and grown on ACM with

0.8% agarose and 0.1% PPM for live imaging. Confocal stacks were taken on an LSM 700

confocal microscope (Carl Zeiss) operated with the ZEN 2010 software (version 6.0, Carl

Zeiss), using a W N-Achroplan 40x/0.75 M27 water immersion objective, and on a TCS SP8

confocal microscope (Leica) operated with the Leica Application Suite X software (version 3.5,

Leica), using a Fluotar VISIR 25x/0.95 water immersion lens. GFP was excited at 488 nm and

emission detected between 415 – 735 nm. Stacks have resolution of 1028×1028 pixels, with

resolution ranging between 3.2 to 4.4 pixels/μm; Z steps were between 0.5 and 0.85μm.

For hyperosmotic treatments, meristems were mounted in Petri-dishes on Patafix (UHU), then

submerged in liquid ACM containing 0.1% PPM. Samples were first imaged in liquid ACM

(plus 0.1% PPM), then submersion medium was changed to ACM plus desired concentration

of NaCl (plus 0.1% PPM) by first rinsing with 3~5 mL target solution, then soaked in target

solution for 5 minutes before imaging. Because of the reduced signal in hyperosmotic solutions,

possibly due to the altered refractive index, stronger gain was used to reach comparable signal

intensity.

Image processing and geometric analysis

3D shell mesh and surface projection of untreated meristems were generated from confocal

stacks using the level set method (LSM) addon [80] for the software MorphoGraphX (MGX

version 1.0) [81]. For oryzalin-treated meristems, 2D surface projections were generated by

MerryProj [75] and imported into MGX for further processing. Projected images were

segmented using watershed method after manual seeding, and cell lineage between time points

was manually assigned in the meristem proper. To limit Z distortion and biases due to change

in inclination of the surface, which may affect analysis accuracy, only cells within 20° of

inclination angle from the highest position of the SAM were included. A custom-made Python

script was used to trace cell lineage between multiple time points and determine cell topology

based on the anticlinal wall number exported from MGX. Areal relative exponential growth

rate per hour was calculated as:

568
$$G = \frac{\ln(A_t/A_0)}{\Delta t}$$

where Δt is time interval in hours, A_0 is original cellular area at time t_0, and A_t is final area at

time $t_0 + \Delta t$. Cells undergone topological changes (i.e. divided cells and cells adjacent to new

division planes) during the acquisition were not included in the growth analyses. To analyse

variation, cell-specific growth rate was further co-aligning by the median per SAM then

stretching the distribution to the average first and third quartile positions to each data point.

To enhance the confocal images in hyperosmotic solution, anisotropic diffusion filter [82] was

re-implemented and applied to the raw images with the following parameters (specifications

see http://cbp-domu-forge.ens-lyon.fr/redmine/projects/anifilters/wiki): $K = 0.3$, $\sigma = 5$, $\gamma = 0.9$,

$D = 10$, 50 iterations.

For figure panels, brightness and contract of confocal images were linearly enhanced for better

visual. To synchronize panel shape and size, black background with no relevant information

was cropped from or added to the edge of the panels.

Finite element method (FEM) indentation simulation

Inflation and indentation simulations were performed with the same procedure adopted from

Mosca et al., 2017 [46]. For each meristem, a surface projection of the L1 anticlinal walls

obtained from confocal microscopy was segmented and extruded to the average anticlinal wall

height as observed in the template (5 μm for untreated and 10μm for oryzalin-treated) with the

CellMaker plugin for MorphoGraphX [81]. This generates a multicellular template made of

triangular membrane elements. The extruded meshes keep the average overall organ curvature.

In order to give the extruded cells a more realistic rest curvature (in the unpressurized state),

both meshes were pre-inflated with a small pressure (0.05 MPa for untreated and 0.15 MPa

oryzalin-treated SAM) and saved as the rest configuration. Both meshes were refined around

the indentation points to increase the accuracy of the indentation simulations. The untreated

mesh used in the indentation simulation and curvature analysis has an average edge size of 0.1

μm near the indentation points and increases to 0.5 μm far away from those points. The

oryzalin-treated mesh has instead an average edge size of 0.2 μm and 1 μm near and far from

the indentation the indentation areas, respectively.

The cell wall was modelled as a Saint-Venant Kirchhoff material represented by membrane

elements with zero transversal stress and mathematically prescribed thickness [46]. The two

templates were assigned the following material properties:

- isotropic material, 200 MPa Young's modulus, 0.3 Poisson ratio;

- cell wall thickness of 0.19 μm and 0.74 μm for the untreated and oryzalin-treated meristem,

following experimental data.

The templates were inflated from the rest configuration with a turgor pressure of 2 MPa inside

each cell for both meshes, until the forces equilibrium was reached (sum of the forces squared).

With the chosen parameters, the total cell height after inflation is similar to the experimental

observation from the confocal stacks for the referring templates (6.8 μm for untreated and 14.5

μm for oryzalin-treated). Afterwards the bottom of the template (bottom anticlinal cell walls)

was blocked in all space degrees of freedom to simulate the presence of the supporting inner

tissue during the indentation process.

The indentation is modelled as in [46] and is performed in the global z-direction as given by

the confocal images. The indentation process is performed on each cell independently, where

the specific indentation point is chosen to be close to the uppermost anticlinal wall location.

This reproduces the AFM indentation modality. As in analysis of AFM experiments, the

stiffness of the untreated cell was computed as the slope of the indentation curve at 0.5 μm

indentation depth, with the reaction forces on the indenter between 0.3 and 0.7 μm depth used

for slope computation. For the oryzalin-treated, given the thicker walls, the indentations were

deeper and the stiffness was computed around 1 μm indentation depth (reaction forces between

0.8 and 1.2 μm).

For both meshes, a refinement analysis was performed to verify the accuracy of the simulation

results. The results are reported in Supplementary Table 1. We considered the variation of the

coarse oryzalin mesh so small to justify using it for the AFM and curvature comparison analysis,

while we preferred using the refined mesh for the untreated to reduce the error due to mesh

resolution.

Mesh	coarse size	refined size	Relative MD	sd of relative MD
Untreated	0.2 μm	0.1 μm	3%	2.4%
Oryzalin-treated	0.4 μm	0.2 μm	2%	1.3%

Supplementary Table 1. FEM refinement analysis. Coarse mesh edge size near
indentation; refined mesh average edge size near indentation; relative MD, relative mean
absolute difference; sd, standard deviation.

Statistical analysis

Data were processed using Excel 2000 (Microsoft). All Tukey box plots depict the first, second

(median) and third quartiles of data distribution, with whiskers marking the lowest/highest data

within 1.5 interquartile ranges (IQR) of the lower/upper quartiles. Tukey's outliers are depicted

as small circles outside the whiskers. Values like turgor pressure, cell area and growth rate were

normalized to the average per meristem. After normalization, every cell was considered as one

biological sample, and all linear regressions and Pearson correlations were performed on whole

datasets. For simulations, cells on the edge of the mesh were not analysed due to border effect.

Extremely rare polygon classes (i.e. triangle and nonagon) were not shown on the box plots in

the main figures but were included in linear regression and Pearson correlation tests and were

plotted in Figure S5. Kolmogorov–Smirnov test was used to distinguish cell neighbour number

distribution (significance level $\alpha = 0.05$).

Mechanical-hydraulic modelling

Summary. We build a vertex-based model of plant tissues at cellular level that couples osmosis-

driven hydraulic fluxes between cells and from apoplast with a fixed water potential, and cell

wall mechanics which resists and grows under tension. Turgor and growth rate heterogeneities

emerge from this coupling and from the heterogeneities in cells sizes and topology (number of

neighbours).

We consider a collection of N polygonal cells $i = 1, \ldots, N$ that form a mesh; this mesh evolves

with the appearance of new cells because of cell division. The walls between cells are

discretized into one or several segments. Given the topology, the mesh is fully characterized by

the position of the vertices. The walls are given a height h and a thickness w.

Cell wall rheology. The cell walls are modelled as a visco-elasto-plastic material, which would

be equivalent to the Ortega model [20] in the case of an elongating cell. Let σ_k be the stress of

a wall segment k; the constitutive law writes $\sigma_k = E_k \varepsilon_k^e$ where E_k is the elastic modulus and

ε_k^e is the elastic deformation of the wall. Let l_k be the length of segment k, the rate of change

of ε_k^e is given by:

$$\frac{\mathrm{d}\varepsilon_k^e}{\mathrm{d}t} + \Phi_k^w E_k \max\left(0, \varepsilon_k^e - \varepsilon_k^Y\right) = \frac{1}{l_k}\frac{\mathrm{d}l_k}{\mathrm{d}t}$$

654   where Φ_k^w is the extensibility and ε_k^Y is the yield deformation of segment k. Equivalently, we

655   could define a yield stress.

656   *Mechanical equilibrium.* Let P_i be the turgor pressure in each cell i. The tissue being at every

657   moment in a quasi-static equilibrium, pressure forces on wall edges and elastic forces within

658   walls balance exactly at each vertex v:

$$\frac{1}{2}\sum_{k\in f(v)} \Delta_k P A_k \boldsymbol{n}_k + \sum_{k\in f(v)} E_k \varepsilon_k^e a_k \boldsymbol{e}_{k,v} = 0$$

Where $f(v)$ is the set of walls adjacent to junction v, and $\Delta_k P = P_{k_1} - P_{k_2}$ is the pressure jump

across wall face k, with $k_1 < k_2$ as indices of the cells separated by face k, $A_k = h l_k$ is the area

of the face k on which pressure is exerted, $\boldsymbol{n}_k$ is the normal vector to face k, oriented from cell

k_1 to cell k_2, and $a_k = h w$ is the cross-section of the face, on which the elastic stress is exerted;

finally, $\boldsymbol{e}_{k,v}$ is the unit vector in the direction of face k, oriented from junction v to the other end

of face k. In the case of a single cylindrical cell for which growth is restricted to its principal

direction, the model is equivalent to the Lockhart-Ortega model.

Fluxes. For each cell i, the apoplasmic pathway is represented as a flux U_i^a (in volume per

time unit) from the apoplast of constant water potential Ψ^a through a perfectly semi-permeable

membrane: $U_i^a = A_i L_i^a (\Delta\Pi - P_i)$, where A_i is the area of each cell in contact with the apoplast,

L_i^a is the corresponding water conductivity, $\Delta\Pi = \pi_i + \Psi^a$ is assumed constant, and π_i is the

osmotic pressure of cell i.

The symplasmic pathway corresponds to flows that occur through plasmodesmata, channels

between cells that convey both water and solutes. The symplasmic flows thus only depend on

turgor pressure difference. Let L_{ij} be the symplasmic water conductivity corresponding to the

interface between two neighbour cells i and j, and A_{ij} their contact area, both assumed

symmetric: $L_{ij} = L_{ji}$ and $A_{ij} = A_{ji}$. The symplasmic flux U_{ji}^s (in volume per time unit) from

cell j to i is defined by:

678
$$U_{ji}^s = A_{ij} L_{ij}^s (P_j - P_i)$$

679   Finally, the total water flux for cell i is the sum of the apoplasmic flux U_i^a and the symplasmic

680   fluxes U_{ji}^s with all its neighbors, so that its volume variation can be expressed as:

681
$$\frac{dV_i}{dt} = A_i L_i^a (\Delta\Pi - P_i) + \sum_{j \in n(i)} A_{ij} L_{ij}^s (P_j - P_i)$$

where $n(i)$ is the set of neighbours of cell i.

Cell division. We implemented the Willis-Refahi rule [34], in which the division volume is

given by $V_0 = f V_b + \mu_b (2 - f + Z)$, where $f = 0.5$, V_b is the volume at birth, $\mu_b = 3.31$

is the mean birth volume and Z is a Gaussian noise with zero mean and $\left(4\sigma_d^2 - f^2 \sigma_b^2\right)^{1/2}$

standard deviation, with $\sigma_b = 0.2$ and $\sigma_d = 0.1$.

Numerical resolution. In the Lockhart-Ortega model, the compatibility between wall elongation

and cell volume increase is automatically enforced through the geometrical constraint of

unidirectional growth that leads to equal relative growth rate of the cell and strain rate of the

walls. In our multicellular model, this equality is no longer true. Instead, the lengths $l(X)$ of the

edges and the volumes $V(X)$ of the cells are expressed as functions of the positions X of the

vertices; then, given an initial position X of the vertices and elastic deformation ε^e of the edges,

the equations of wall rheology, mechanical equilibrium, and water fluxes form a closed set of

equations with respect to the unknowns X, P, and ε^e that allow to predict their evolution.

To give an idea of the mathematical complexity of the problem, one may consider the following

example: in a connected tissue, if one cell is stretched and forced to increase its volume, an

equal volume of water has to enter the cell, either from the apoplastic compartment or the

neighbour cells. In the latter case, pressure should drop in the neighbour cells, which should

attract water from their own neighbours, and this could propagate to further cells depending on

the geometry of the tissue and the effective parameters. Volume and therefore positions of the

vertices could be also affected. Finally, one can see that the interaction between hydraulics and

mechanics implies long range interactions where pressure plays a key role.

We developed an original algorithm and implemented it in an in-house code, where at each time

step, the mechanical equilibrium is resolved under constraints on the cell volume (from the

water fluxes), and constraints on the cells edges (from the rheological law of the walls). This

was implemented in Python, using the open source libraries NumPy, SciPy, and the Topomesh

class from the OpenAlea project

(http://openalea.gforge.inria.fr/doc/vplants/container/doc/html/container/openalea_container_t

opomesh_ref.html). This algorithm is described in more detail in a separate publication [30].

The computations were run on a computer with a 3.6GHz Intel Xeon E5 processor, 64 GB of

RAM, and running Linux Debian Stretch. The typical computing time was one week for each

computation.

Parameterization of the model. The reference values of the parameters were chosen based on

the literature or on our experiments (main text Table 1), except for α^s, for which no data is

available and was conservatively ascribed an intermediate value of 0.5. Model behaviour was

explored by changing non-dimensional parameter values as explained in the main text.

Procedure for the computations. We first run in parallel three computations with cell division

to around 300 cells. To mimic untreated case, simulations continue until around 600 cells. To

mimic the oryzalin treatment, the current states of the "untreated" computations at around 300

cells are used as initial conditions for the oryzalin case: division is stopped, and we run

computations either with the same effective parameters, or with some parameters modified so

that the behaviour of the oryzalin treated meristems is recovered (see supplementary table 2);

the computations are run until the total volume has been multiplied by three from this initial

state. Typical runtime for one parameter set is one week.

Dividing simulations	SIM#	n	Non-dividing simulations	SIM#	n
Initial (Fig. 1f) $\alpha^s = 0.5, \alpha^a = 0.5, \theta = 7/3$	3	1535	No division (Fig. 1h, 6d)	3	759
Reference (Fig. 1g, 6a) $\alpha^s = 0.5, \alpha^a = 0.5, \theta = 20/3$	3	1496	$\Delta\Pi \times 2$	3	759
$\theta = 10/3$	3	1515	$w \times 2$	3	759
$\theta = 10$	3	1519	$w \times 4$ (Fig. 6g)	3	759
$\theta = 40/3$	3	1530	$\Delta\Pi \times 2 , w \times 2$	3	759
$\alpha^s = 0.1$	3	1506	$\Delta\Pi \times 2 , w \times 4$	3	759
$\alpha^s = 0.9$	3	1525			
$\alpha^a = 0.1$	3	1507			
$\alpha^a = 0.9$	3	1519			

Supplementary table 2. Parameter modification and simulation repeats. Parameters are

systemically changed, with the reference parameter set $\alpha^s = 0.5$, $\alpha^a = 0.5$, $\theta = 20/3$ for

exploration in untreated simulations and for mimicking oryzalin in non-dividing simulations.

Annotations are as described above. SIM#, number of simulations; n, total cell number at last

time point without boundary cells.

Acknowledgements

We thank P. Bolland and A. Lacroix for plant care, G. Cloarec for help with TEM, L. Beauzamy

for AFM training, C. Mollier for help in estimating confocal technical errors, A. Kiss for help

with anisotropic filtering parameterization, G. Cerutti for help in implementing the division

algorithm and in segmentation, R. Smith for help in segmentation and mesh generation for the

indentation simulations, and V. Battu, F. Zhao and C. Galvan-Ampudia for providing plant

materials. We acknowledge the contribution of the PLATIM facility of SFR Biosciences

(UMS3444/CNRS, US8/Inserm, ENS de Lyon, UCBL) for AFM and confocal microscopy, and

of Centre Technologique des Microstructures (UCBL, Lyon) for electron microscopy. This

work was supported by a fellowship from Institut Universitaire de France and an ERC Starting

Grant "PhyMorph" to A.B. (ERC-2012-StG-307387), an EMBO Long-term Fellowship to Y.L.

(EMBO ALTF 168-2015), and an Agropolis Foundation grant (MecaFruit3D) to I.C. and C.G.

Author contributions

This study was initiated by A.B. Y.L. and A.B. designed the experiments. Y.L. executed AFM

and confocal microscopy, acquired experimental data except EM, and analysed experimental

and simulation data. J.T. performed electron microscopy. V.M and M.D. wrote scripts to

facilitate experimental data analysis. I.C. and C.G. designed physical model of tissue growth.

I.C. implemented the model, ran simulations, optimized model parameters and analysed

simulation data. G.M. designed, ran, and analysed indentation simulations. Y.L. and A.B.

contributed to the design of simulations. Y.L., I.C., V.M., G.M., C.G. and A.B. contributed to

data interpretation. Y.L. and A.B. wrote the manuscript with inputs from the other authors.

Data availability

All materials, scripts and datasets generated and analysed during the current study are available

from the corresponding authors upon reasonable request.

Figure 1. Turgor pressure heterogeneity emerges from cell topology in the mechano-hydraulic model.

(**a**) In a plant cell, the turgor pressure P is contained by the cell wall tensile stress σ. (**b**) A schematic representation of the Lockhart-Ortega equation, where 1D cell length L elongation is a combination of reversible stretch ϵL (elasticity, ϵ is elastic strain) and cell wall yield ΔL at longer timescale (viscosity) if P is higher than a threshold Y (effective plasticity, $\Delta L = \Phi t L (P - Y)$, Φ is wall extensibility, t is time). (**c**) Schematic representations of model components, including cell geometry with height h and radius R (left), apoplasmic/transmembrane and symplasmic/intercellular water fluxes (middle left), mechanical equilibrium at tricellular junctions (middle right) and the visco-elasto-plastic cell wall rheology (right). P_i, cell-specific turgor pressure; σ, cell wall tension; Φ^w, wall extensibility; ε^Y, wall strain threshold; E, wall Young's modulus. (**d**) Simulation snapshots: in "dividing" simulations, 16 initial cells grow and divide until about 600 cells; in "non-dividing" simulations, divisions are stopped when cell number reaches about 300 cells and growth continues until they triple in size. Colour indicates cell neighbour number as in (e). (**e**) Similar distributions of cell neighbour number in the experimentally observed (Exp.) shoot apical meristem and in simulations (Sim.) by Kolmogorov–Smirnov test (confidence level $\alpha = 0.05$, $D_{n,m} < D_\alpha$), error bars are standard deviations. *, Student's t-test $p < 0.05$; **, $p < 0.01$. (**f-h**) Cell wall strain, turgor pressure, neighbour number, and area in dividing (f, g) and non-dividing (h) simulations; the dimensionless parameter for flux-wall balance α^a, apoplasmic-symplasmic balance α^s and the osmotic drive are as indicated. Left: cell wall elastic strain ε normalized by yield strain ε^Y; middle left: cell turgor pressure P normalized by average pressure; middle: boxplots of normalized cellular turgor pressure P against cell topology N (**f**, $n = 1535$ cells; **g**, $n = 1496$ cells; **h**, $n = 759$ cells); middle right: normalized pressure against normalized area; right: normalized area against neighbour number. Cells on the mesh edge were not analysed due to border effect. Circles are Tukey's outliers; lowercase letters indicate statistically different populations (Student's t-test, $p < 0.05$); red lines indicate linear regressions, with Pearson correlation coefficient R and corresponding p-value. In (**d, f-h**) scale bars are 5 unit length. (**i**) Model parameter exploration. Colours indicate Person correlation coefficient R, with perfect anticorrelation as blue ($R = -1$), perfect correlation in red ($R = 1$); all correlations were statistically significant ($p < 0.05$). A, normalized cell area; N, cell neighbour number; P, normalized turgor pressure; α^a, dimensionless parameter for flux-wall balance; α^s, dimensionless parameter for apoplasmic-symplasmic balance; θ, dimensionless osmotic drive. (**j**) Simulations initiated with 4-edged bigger than 8-edged cells, shown at two simulation times, with area and pressure shown in top and bottom, respectively; 4-edged cells have higher pressure although bigger. (**k to l**) Schematic explanation of topology-derived turgor pressure heterogeneity. (**k**) Three-cell junctions in a tissue of hexagonal cells are at mechanical equilibrium with equal tensions and equal wall-wall angles. (**l**) Fewer-neighboured cells have sharper wall-wall angles, a tension that is roughly constant per wall effectively results in mechanical compression due to unequal projected tension distribution (red dash-line arrow) that is balanced by higher turgor pressure build-up (big blue arrow).

Figure 2. The experimental pipeline for turgor pressure deduction.

(**a**) Schematic representation of AFM nanoindentation for turgor pressure measurement. r, probe tip radius; k, cantilever stiffness. (**b** to **e**) Illustration for force curve interpretation at different indentation depth. (b) When indentation depth Z_1 is smaller than wall thickness t, the force-indentation curve is sensitive to cell wall property, (c) deeper-than-wall indentation $Z_2 > t$ is also sensitive to turgor pressure P. (d) Even deeper indentation Z_3 deforms surrounding cells and is also sensitive to tissue context. Dotted line marks the shell position before indentation, which is used to determine surface mean and Gaussian curvature, κ_M and κ_G, by AFM topographic scan. (e) Three regimes of the force-indentation curve are used to fit for cell wall Young's modulus E, apparent stiffness at cell-scale k_s and tissue scale k_d. F denotes indentation force. (**f** to **l**) The AFM-confocal pipeline of measurement and deduction on an example untreated SAM. (f) Derjaguin-Muller-Toporov (DMT) modulus map, highlighting cell contours, is projected on the surface topography of an AFM scan. Smaller scan region is chosen to bypass the SAM surface unevenness. (g) Multiple indentations (marked by crosshair) are performed near the barycentre of each cell. (h and i) Confocal stack and its surface projection of the same SAM with plasma membrane GFP signal. (j) Tiled AFM scans are overlaid and stitched on the confocal surface projection image, red square marking the same tile from (f and g). Individual indentation positions are registered on a global coordinate and assigned to segmented cells. (k) Force curves are analysed, and the cellular average of physical values are mapped. (l) Turgor pressure is deduced per force curve and averaged per cell. Stiffness from very deep indentation is not used for cell-specific deduction. (**m** to **o''**) Virtual indentation on realistic 3D mesh. (m) The epidermal layer is meshed based on confocal image. Cells are pressurized with a uniform 2 MPa turgor pressure. (n and n') A cell before (red) and being indented (magenta). (o to o'') Longitudinal section of the indented mesh; black, before indentation; magenta, being indented. Cell junctions

(rectangles) are magnified to highlight the neighbour cell deformation (marked by asterisks) by very deep indentation. (**p** and **q**) AFM-determined surface profiles of untreated (p) and oryzalin-treated SAM (q). Scale bars represent 20 μm if not specified. Note that z axes in (f, p and q) are stretched to emphasize SAM surface unevenness.

Figure 3. Turgor pressure heterogeneity in untreated SAM.

(**a**) Top-view surface projections of untreated SAM with plasma membrane GFP signal; scale bars represent 20 μm. (**b**) Map of epidermal cell neighbour number. (**c**) Distribution of neighbour number on 7 untreated SAM surfaces, error bars are standard errors of means. (**d**) Normalized cell area A and cell neighbour number N are linearly correlated. (**e** to **h"**) AFM-determined cellular values and their association with neighbour number N and cell area A, all plotted values are normalized per SAM (7 SAMs, $n = 503$ cells). (e to e") Cell wall Young's moduli E, (f to f") apparent stiffness k at intermediate indentation depth, (g to g") cell surface mean curvature κ_M, (h to h") deduced turgor pressure P. (**i** and **j**) Coefficient of variance (CV) of AFM-determined values in untreated meristems. (i) At subcellular scale (intra), P is much less variable than E and k. (j) P, unlike E and k, are more heterogeneous between cells (inter) than between measurements within single cells (intra). (**o** to **q**) Weighed convex index (wCI) anticorrelates with cell neighbour number (p) and normalized cell area (q). In box plots, circles are Tukey's outliers; lowercase letters indicate statistically different populations (Student's t-test, $p < 0.05$); red lines indicate linear regressions, with Pearson correlation coefficient R and corresponding p-value.

Figure 4. Turgor pressure heterogeneity in oryzalin-treated SAM.
(**c**) Top-view surface projections of oryzalin SAM with plasma membrane GFP signal; scale bars represent 20 μm. (**b**) Map of epidermal cell neighbour number. (**c**) Distribution of neighbour number on 9 untreated SAM surfaces, error bars are standard errors of means. (**d**) Normalized cell area A and cell neighbour number N are linearly correlated. (**e** to **h"**) AFM-determined cellular values and their association with neighbour number N and cell area A, all plotted values are normalized per SAM (9 SAMs, n = 202 cells). (e to e") Cell wall

Young's moduli E, (f to f") apparent stiffness k at intermediate indentation depth, (g to g") cell surface mean curvature κ_M, (h to h") deduced turgor pressure P. (**i** and **j**) Coefficient of variance (CV) of AFM-determined values in untreated meristems. (i) At subcellular scale (intra), P and k are much less variable than E. (j) P and k, unlike E, are more heterogeneous between cells (inter) than between measurements within single cells (intra). (**o** to **q"**) Weighed convex index (wCI) anticorrelates with cell neighbour number (o) and normalized cell area (p). (q to q") Cell walls curve towards cell with low pressure, as signified by the enhanced curvature towards the ablated cell with released pressure. In box plots, circles are Tukey's outliers; lowercase letters indicate statistically different populations (Student's t-test, $p < 0.05$); red lines indicate linear regressions, with Pearson correlation coefficient R and corresponding p-value.

Figure 5. Cellular growth rate bifurcates between conditions.
(**a** to **f**) Relative growth rate per day G of untreated SAM cells. (**g** to **l**) Cellular growth rate per day of oryzalin-treated SAM between 12-hour interval (48 and 60 hours post treatment). (a, b, g and h) Surface projections of untreated or oryzalin-treated SAM at initial time point

(a and g) and 12 hours later (b and h); scale bars are 20 μm unless otherwise noted. (c and i) Heat maps of areal relative growth rate per day. (d and j) Example 4 and 8-neighbored cells during 24-hour growth, with areal normalization at initial time point. Cell contour and relative size (blue for 4-neighbored, red for 8-neighbored) depict the diverging growth trends. Scale bars are as indicated. (e, f, k and l) Box plots of relative growth rate per day G against cell topology N (e and k) and dot plots of relative growth rate per day G against normalized cell area A (f and l) (e and f, untreated 11 SAMs, $n = 1491$ cells; k and l, oryzalin-treated 14 SAMs, $n = 1160$ cells). Note that Tukey's outliers are plotted in Figure S5 and all data are included for statistical analyses. Lowercase letters indicate statistically different populations (Student's t-test, $p < 0.05$); red lines indicate linear regressions, with Pearson correlation coefficient R and corresponding p-value.

Figure 6. The model recapitulates the untreated and oryzalin-treated growth trends.
(**a** to **i**) Relative growth rate G normalized by average growth rate, neighbour number N, and
normalized cell area A, with reference values of dimensionless parameters ($\alpha^a = 0.5$, $\alpha^s = 0.5$,

θ = 20/3): dividing simulations (a-c), non-dividing simulations (d-f), and non-dividing simulations with quadruple wall thickness w to mimic oryzalin treatment (g-i). (a, d, g) Heat maps of normalized areal relative growth rate. (b, e, h) Box plots of normalized relative growth rate G against cell topology N. Lowercase letters indicate statistically different populations (Student's t-test, $p < 0.05$); red lines indicate linear regressions, with Pearson correlation coefficient R and corresponding p-value. (c, f, i) Dot plots of normalized relative growth rate G against normalized cell area A. For numbers of repeats see STAR Methods. (**j**) Model exploration to fit oryzalin-treated case. Colours indicate Person correlation R, with perfect anticorrelation as blue (R = -1), perfect correlation in red (R = 1), and insignificant correlation ($p > 0.05$) in black. A, normalized cell area; N, cell neighbour number; P, normalized turgor pressure; G, normalized relative growth rate; $\Delta\Pi$, transmembrane osmotic pressure difference; w, wall thickness. (**k, l**) Influence of dimensionless (k) and dimensional parameters (l) on growth trends. Triangles indicate parameter's influence to the mechanical–hydraulic balance. α^a, dimensionless parameter for flux-wall balance; α^s, dimensionless parameter for symplasmic-apoplasmic balance; θ, dimensionless osmotic drive. w, wall thickness; E, cell wall modulus; ε^Y, strain threshold; L^a, cross-membrane water conductivity; L^s, cell-to-cell symplasmic conductivity; Φ^w, wall extensibility; $\Delta\Pi$, transmembrane osmotic pressure difference; R, representative cell size. (**m**) Summary of growth trend inversion according to the mechanical–hydraulic control shown in (j). The wall and flux limits correspond to untreated and oryzalin-treated SAMs, respectively.

Figure S1. Osmotic pressure and hydrostatic pressure.

(**a**) Osmosis occurs when solutions of different osmotic pressure Π, due to different solute concentration (depicted by the density of white dots), are separated by a semipermeable membrane. Difference of osmotic potential Ψ_Π (depicted by horizontal black lines, always negative in solutions) across the membrane dictates that solvent flows from high to low

potential compartment (low to high solute concentration). In a U-shaped tube setup (right), osmosis may stop before the two compartments reach the same concentration, as the extra volume (in fact height, Δh) in the higher-concentration compartment exerts a hydrostatic pressure P due to gravity that pushes the solvent back. P increases until reaching the same value as $\Delta\Pi$ at equilibrium where the total water potential Ψ_W is equal on both sides, and the net solvent flux $I = 0$ (indicated by the opposite blue arrows). M, solute molar concentration; i, van t'Hoff index of solute that disassociates; R, ideal gas constant; T, absolute temperature. (**b**) Osmosis in a plant cell, where gravity is neglected for its small size. Because of the rigid cell wall that restricts the cell volume, hydrostatic pressure P, alias turgor pressure, builds up alongside cell wall tension σ to counterbalance the difference of osmotic potential Ψ_Π, until $P = \Delta\Pi$. (**c**) Geometrical extension of Lockhart model for a single polygonal cell. σ, cell wall stress; P, turgor pressure; R, cell radius; N, polygon class; v, vertex; $\boldsymbol{n_1}$ and $\boldsymbol{n_2}$; normal vector; $\boldsymbol{e_1}$ and $\boldsymbol{e_2}$, tangent vector. (**d**) Half of the threshold pressure of a polygonal cells base on $P^Y = \frac{w}{2R\cos(\pi/N)} E\varepsilon^Y$. For growing cells, turgor pressure P falls between the transmembrane osmotic pressure $\Delta\Pi$ and pressure threshold P^Y.

Figure S2. Simulations for vertex model parameter exploration.

Model parameters were systemically changed. α_a, dimensionless parameter determining the weight of the limitation of growth by transmembrane permeability to the combined limitation of growth by cell wall extensibility and transmembrane permeability; α_s, dimensionless

parameter determining the weight of intercellular water conductivity over total (intercellular and transmembrane) conductivity; *s*, cell size scale at division relative to the initial setting. *P*, normalized turgor pressure, *A*, normalized cell area; *G*, relative growth rate per hour. Polygons indicate cell neighbour number. Red lines indicate linear regressions, with Pearson correlation coefficient *R* and corresponding *p*-value.

Figure S3. Cell wall thickness and individual SAM measurements of turgor pressure and growth rate.

(**a** and **b**) TEM and serial block-face imaging SEM images of untreated (a) and oryzalin-treated meristem (b). (**c**) Boxplots of surface wall thickness of untreated (Unt) and oryzalin-treated (Ory) meristems. (**d**) Surface wall thickness does not correlate to cell width in oryzalin-treated meristems. (**e-h**) Oryzalin-treated SAMs showed highly variable turgor pressure and growth rate per SAM, while untreated samples show little variation; *n* is cell number measured per meristem. Circles are Tukey's outliers; lowercase letters indicate statistically different populations (Student's *t*-test, *p* < 0.05). (g) The first five untreated meristems measured by AFM were of ecotype WS-4, while the last two are of Col-0.

Figure S4. Hyperosmotic treatments on SAMs.

(**a** and **b**) Confocal images of untreated (a) and oryzalin-treated (b) meristems submerged in liquid apex culture medium (ACM) before and after the supplementation of NaCl. The

increased osmolality resulted in blurred images (each upper-right subpanel), possibly due to altered refraction limit, which required anisotropic filtering to enhance the trajectory of cell contours. Out of 8 untreated meristems, half showed incipient plasmolsysis at 200 mM NaCl (~0.4 Osm) and the other half at 300 mM (~0.6 Osm). For 7 oryzalin-treated meristems, incipient plasmolysis occurred among a broader osmolarity (2 meristems at 300 mM, 3 meristems at 400 mM, 2 meristems at 500 mM). Arrows points to noticeable membrane detachment and rounding, indicating plasmolysis. (**c**) Osmolarity calibration by osmometer measurements ($r^2 > 0.99$). (**d**) Moderate hyperosmotic treatment decreases AFM-determined turgor pressure, and reduces turgor heterogeneity plotted against cell neighbour number (linear regression slope a, Pearson correlation coefficient R and corresponding p-value).

Figure S5. Pressure and growth rate before and after normalization, and regional measurements.

(**a** to **j**) Normalized and raw turgor pressure P and growth rate G of untreated meristems plotted against cell neighbour number N and cell area A. (**k** to **t**) Normalized and raw turgor pressure P and growth rate G of oryzalin-treated meristems plotted against cell neighbour number and cell area. (a, f, k, p) Normalized P and co-aligned G against N plots complete

with outliers and the rare 3- and 9-neighboured cells, same data as in Figure 3 and 4. (b, g, l, q) Same normalized data with only the major polygonal classes (N = 5, 6, 7), showing that the correlations do not depend on the rare polygonal classes at the ends of distributions. (c, h, m, r) Same normalized data were binned for cells with 5 or fewer neighbours (≤5), cells with 6 neighbours and cells with 7 or more neighbours (≥7), showing the same trends as unbinned data. (d, i, n, s) Raw data of P and G before normalization / co-alignment against neighbour number; (e, j, o, t) raw data of P and G before normalization / co-alignment against unnormalized cell area, all showing the same trend as normalized data. Red lines are linear regression line, with Pearson correlation coefficient R and corresponding p-value. Lower case letters denote statistically significant groups (Student's t-test $p < 0.05$). (**u**) Frequency of N-neighboured cells in central zone (CZ) and peripheral zone (PZ) shows no significant difference (Kolmogorov–Smirnov test, confidence level α = 0.05, $D_{n,m} < D_{\alpha}$). (**v, w**) Normalized turgor pressure and co-aligned growth rate in subregions of SAM; n is cell number, error bars indicate standard error of mean, lower case letters denote statistically significant groups (Student's t-test $p < 0.05$).

Figure S6. Simulations to recapitulate oryzalin-treated SAM.
Based on the reference parameter set (α^s = 0.5, α^a = 0.5, θ = 20/3), which represents the untreated-like behaviour with cell divisions (Dividing), oryzalin-treated-like behaviour was further fitted by stopping cell division (No div.) and subsequently changing osmotic pressure $\Delta\Pi$ and cell wall thickness w, based on experimental observations. P, normalized turgor pressure, A, normalized cell area; G, normalized relative growth rate. Polygons indicate cell neighbour number. Red lines indicate linear regressions, with Pearson correlation coefficient R and corresponding p-value.

758   **References**

759   1.   Hong, L.; Dumond, M.; Zhu, M.; Tsugawa, S.; Li, C.-B.; Boudaoud, A.; Hamant, O.; Roeder,
760   A.H.K. Heterogeneity and Robustness in Plant Morphogenesis: From Cells to Organs. *Annu.*
761   *Rev. Plant Biol.* **2018**, *69*, 469–495.

762   2.   Kamimoto, K.; Kaneko, K.; Kok, C.Y.-Y.; Okada, H.; Miyajima, A.; Itoh, T. Heterogeneity and
763   stochastic growth regulation of biliary epithelial cells dictate dynamic epithelial tissue
764   remodeling. *eLife* **2016**, *5*.

765   3.   Ietswaart, R.; Rosa, S.; Wu, Z.; Dean, C.; Howard, M. Cell-Size-Dependent Transcription of FLC
766   and Its Antisense Long Non-coding RNA COOLAIR Explain Cell-to-Cell Expression Variation.
767   *Cell Syst.* **2017**, *4*, 622-635.e9.

768   4.   Long, Y.; Stahl, Y.; Weidtkamp-Peters, S.; Postma, M.; Zhou, W.; Goedhart, J.; Sánchez-Pérez,
769   M.-I.; Gadella, T.W.J.; Simon, R.; Scheres, B.; et al. In vivo FRET-FLIM reveals cell-type-specific
770   protein interactions in Arabidopsis roots. *Nature* **2017**, *548*, 97–102.

771   5.   Chubb, J.R. Symmetry breaking in development and stochastic gene expression. *Wiley*
772   *Interdiscip. Rev. Dev. Biol.* **2017**, *6*, e284.

773   6.   Donati, G.; Watt, F.M. Stem Cell Heterogeneity and Plasticity in Epithelia. *Cell Stem Cell* **2015**,
774   *16*, 465–476.

775   7.   Gerdes, M.J.; Sood, A.; Sevinsky, C.; Pris, A.D.; Zavodszky, M.I.; Ginty, F. Emerging
776   Understanding of Multiscale Tumor Heterogeneity. *Front. Oncol.* **2014**, *4*.

777   8.   Eldar, A.; Elowitz, M.B. Functional roles for noise in genetic circuits. *Nature* **2010**, *467*, 167–
778   173.

779   9.   Watanabe, K.; Umeda, T.; Niwa, K.; Naguro, I.; Ichijo, H. A PP6-ASK3 Module Coordinates the
780   Bidirectional Cell Volume Regulation under Osmotic Stress. *Cell Rep.* **2018**, *22*, 2809–2817.

781   10.   Xie, K.; Yang, Y.; Jiang, H. Controlling Cellular Volume via Mechanical and Physical Properties
782   of Substrate. *Biophys. J.* **2018**, *114*, 675–687.

783   11.   Guo, M.; Pegoraro, A.F.; Mao, A.; Zhou, E.H.; Arany, P.R.; Han, Y.; Burnette, D.T.; Jensen, M.H.;
784   Kasza, K.E.; Moore, J.R.; et al. Cell volume change through water efflux impacts cell stiffness
785   and stem cell fate. *Proc. Natl. Acad. Sci.* **2017**, *114*, E8618–E8627.

786   12.   Dumais, J.; Forterre, Y. "Vegetable Dynamicks": The Role of Water in Plant Movements. *Annu.*
787   *Rev. Fluid Mech.* **2012**, *44*, 453–478.

788   13.   Stewart, M.P.; Helenius, J.; Toyoda, Y.; Ramanathan, S.P.; Muller, D.J.; Hyman, A.A. Hydrostatic
789   pressure and the actomyosin cortex drive mitotic cell rounding. *Nature* **2011**, *469*, 226–230.

790   14.   Montel, F.; Delarue, M.; Elgeti, J.; Malaquin, L.; Basan, M.; Risler, T.; Cabane, B.; Vignjevic, D.;
791   Prost, J.; Cappello, G.; et al. Stress clamp experiments on multicellular tumor spheroids. *Phys.*
792   *Rev. Lett.* **2011**, *107*, 188102.

793   15.   Rojas, E.R.; Huang, K.C. Regulation of microbial growth by turgor pressure. *Curr. Opin.*
794   *Microbiol.* **2017**, *42*, 62–70.

795   16.   Zerzour, R.; Kroeger, J.; Geitmann, A. Polar growth in pollen tubes is associated with spatially
796   confined dynamic changes in cell mechanical properties. *Dev. Biol.* **2009**, *334*, 437–446.

797   17.   Lopez, R.; Badel, E.; Peraudeau, S.; Leblanc-Fournier, N.; Beaujard, F.; Julien, J.-L.; Cochard,
798   H.; Moulia, B. Tree shoot bending generates hydraulic pressure pulses: a new long-distance
799   signal? *J. Exp. Bot.* **2014**, *65*, 1997–2008.

18.   Beauzamy, L.; Nakayama, N.; Boudaoud, A. Flowers under pressure: ins and outs of turgor

regulation in development. *Ann. Bot.* **2014**, mcu187.

19.   Feng, W.; Lindner, H.; Robbins, N.E.; Dinneny, J.R. Growing Out of Stress: The Role of Cell- and
Organ-Scale Growth Control in Plant Water-Stress Responses. *Plant Cell* **2016**, *28*, 1769–1782.

20.   Ortega, J.K. Augmented growth equation for cell wall expansion. *Plant Physiol.* **1985**, *79*, 318–
320.

21.   Kroeger, J.H.; Zerzour, R.; Geitmann, A. Regulator or driving force? The role of turgor pressure
in oscillatory plant cell growth. *PloS One* **2011**, *6*, e18549.

22.   Sager, R.E.; Lee, J.-Y. Plasmodesmata at a glance. *J. Cell Sci.* **2018**, *131*, jcs209346.

23.   Willmer, C.M.; Sexton, R. Stomata and plasmodesmata. *Protoplasma* **1979**, *100*, 113–124.

24.   Wille, A.C.; Lucas, W.J. Ultrastructural and histochemical studies on guard cells. *Planta* **1984**,
160, 129–142.

25.   Ruan, Y.L.; Llewellyn, D.J.; Furbank, R.T. The control of single-celled cotton fiber elongation
by developmentally reversible gating of plasmodesmata and coordinated expression of
sucrose and K+ transporters and expansin. *Plant Cell* **2001**, *13*, 47–60.

26.   Rygol, J.; Pritchard, J.; Zhu, J.J.; Tomos, A.D.; Zimmermann, U. Transpiration Induces Radial
Turgor Pressure Gradients in Wheat and Maize Roots. *Plant Physiol.* **1993**, *103*, 493–500.

27.   Robbins, N.E.; Dinneny, J.R. Growth is required for perception of water availability to pattern
root branches in plants. *Proc. Natl. Acad. Sci.* **2018**, *115*, E822–E831.

28.   Corson, F.; Hamant, O.; Bohn, S.; Traas, J.; Boudaoud, A.; Couder, Y. Turning a plant tissue into
a living cell froth through isotropic growth. *Proc. Natl. Acad. Sci. U. S. A.* **2009**, *106*, 8453–
8458.

29.   Ishihara, S.; Sugimura, K.; Cox, S.J.; Bonnet, I.; Bellaïche, Y.; Graner, F. Comparative study of
non-invasive force and stress inference methods in tissue. *Eur. Phys. J. E Soft Matter* **2013**,
36, 9859.

30.   Cheddadi, I.; Génard, M.; Bertin, N.; Godin, C. Coupling water fluxes with cell wall mechanics
in a multicellular model of plant development. *PLOS Comput. Biol.* **2019**, *15*, e1007121.

31.   Lucas, W.J.; Ham, B.-K.; Kim, J.-Y. Plasmodesmata - bridging the gap between neighboring
plant cells. *Trends Cell Biol.* **2009**, *19*, 495–503.

32.   Kumar, N.M.; Gilula, N.B. The Gap Junction Communication Channel. *Cell* **1996**, *84*, 381–388.

33.   McLean, P.F.; Cooley, L. Protein Equilibration through Somatic Ring Canals in Drosophila.
Science **2013**, *340*.

34.   Willis, L.; Refahi, Y.; Wightman, R.; Landrein, B.; Teles, J.; Huang, K.C.; Meyerowitz, E.M.;
Jönsson, H. Cell size and growth regulation in the Arabidopsis thaliana apical stem cell niche.
Proc. Natl. Acad. Sci. U. S. A. **2016**, *113*, E8238–E8246.

35.   Boyer, J.S.; Cavalieri, A.J.; Schulze, E.-D. Control of the rate of cell enlargement: Excision, wall
relaxation, and growth-induced water potentials. *Planta* **1985**, *163*, 527–543.

36.   Cosgrove, D.J. Cell Wall Yield Properties of Growing Tissue: Evaluation by in Vivo Stress
Relaxation. *Plant Physiol.* **1985**, *78*, 347–356.

37.   Ortega, J.K.E. Dimensionless Numbers to Analyze Expansive Growth Processes. *Plants* **2019**,
8, 17.

38.   Vella, D.; Ajdari, A.; Vaziri, A.; Boudaoud, A. The indentation of pressurized elastic shells: from
polymeric capsules to yeast cells. *J. R. Soc. Interface* **2012**, *9*, 448–455.

39.   Vella, D.; Ajdari, A.; Vaziri, A.; Boudaoud, A. Indentation of Ellipsoidal and Cylindrical Elastic
Shells. *Phys. Rev. Lett.* **2012**, *109*, 144302.

40.   Beauzamy, L.; Derr, J.; Boudaoud, A. Quantifying Hydrostatic Pressure in Plant Cells by Using
Indentation with an Atomic Force Microscope. *Biophys. J.* **2015**, *108*, 2448–2456.

41.   Beauzamy, L.; Louveaux, M.; Hamant, O.; Boudaoud, A. Mechanically, the Shoot Apical
Meristem of Arabidopsis Behaves like a Shell Inflated by a Pressure of About 1 MPa. *Front.*
Plant Sci. **2015**, *6*.

42.   Malgat, R.; Faure, F.; Boudaoud, A. A Mechanical Model to Interpret Cell-Scale Indentation
Experiments on Plant Tissues in Terms of Cell Wall Elasticity and Turgor Pressure. *Front. Plant*
Sci. **2016**, *7*.

43.   Lazarus, A.; Florijn, H.C.B.; Reis, P.M. Geometry-induced rigidity in nonspherical pressurized
elastic shells. *Phys. Rev. Lett.* **2012**, *109*, 144301.

44.   Kierzkowski, D.; Nakayama, N.; Routier-Kierzkowska, A.-L.; Weber, A.; Bayer, E.; Schorderet,
M.; Reinhardt, D.; Kuhlemeier, C.; Smith, R.S. Elastic domains regulate growth and
organogenesis in the plant shoot apical meristem. *Science* **2012**, *335*, 1096–1099.

45.   Milani, P.; Gholamirad, M.; Traas, J.; Arnéodo, A.; Boudaoud, A.; Argoul, F.; Hamant, O. In vivo
analysis of local wall stiffness at the shoot apical meristem in Arabidopsis using atomic force
microscopy. *Plant J. Cell Mol. Biol.* **2011**, *67*, 1116–1123.

46.   Mosca, G.; Sapala, A.; Strauss, S.; Routier-Kierzkowska, A.-L.; Smith, R.S. On the micro-
indentation of plant cells in a tissue context. *Phys. Biol.* **2017**, *14*, 015003.

47.   Hong, L.; Dumond, M.; Tsugawa, S.; Sapala, A.; Routier-Kierzkowska, A.-L.; Zhou, Y.; Chen, C.;
Kiss, A.; Zhu, M.; Hamant, O.; et al. Variable Cell Growth Yields Reproducible Organ
Development through Spatiotemporal Averaging. *Dev. Cell* **2016**, *38*, 15–32.

48.   Yi, H.; Chen, Y.; Wang, J.Z.; Puri, V.M.; Anderson, C.T. The stomatal flexoskeleton: how the
biomechanics of guard cell walls animate an elastic pressure vessel. *J. Exp. Bot.* **2019**, *70*,
3561–3572.

49.   Knoblauch, J.; Mullendore, D.L.; Jensen, K.H.; Knoblauch, M. Pico gauges for minimally
invasive intracellular hydrostatic pressure measurements. *Plant Physiol.* **2014**,
pp.114.245746.

50.   Nakayama, N.; Smith, R.S.; Mandel, T.; Robinson, S.; Kimura, S.; Boudaoud, A.; Kuhlemeier, C.
Mechanical Regulation of Auxin-Mediated Growth. *Curr. Biol.* **2012**, *22*, 1468–1476.

51.   Lewis, F.T. The correlation between cell division and the shapes and sizes of prismatic cells in
the epidermis of cucumis. *Anat. Rec.* **1928**, *38*, 341–376.

52.   Gibson, W.T.; Veldhuis, J.H.; Rubinstein, B.; Cartwright, H.N.; Perrimon, N.; Brodland, G.W.;
Nagpal, R.; Gibson, M.C. Control of the mitotic cleavage plane by local epithelial topology.
Cell **2011**, *144*, 427–438.

53.   Hamant, O.; Heisler, M.G.; Jönsson, H.; Krupinski, P.; Uyttewaal, M.; Bokov, P.; Corson, F.;
Sahlin, P.; Boudaoud, A.; Meyerowitz, E.M.; et al. Developmental Patterning by Mechanical
Signals in Arabidopsis. *Science* **2008**, *322*, 1650–1655.

54.   Kwiatkowska, D. Surface growth at the reproductive shoot apex of Arabidopsis thaliana pin-
formed 1 and wild type. *J. Exp. Bot.* **2004**, *55*, 1021–1032.

55.   Serrano-Mislata, A.; Schiessl, K.; Sablowski, R. Active Control of Cell Size Generates Spatial
Detail during Plant Organogenesis. *Curr. Biol.* **2015**, *25*, 2991–2996.

56.   Ali, O.; Mirabet, V.; Godin, C.; Traas, J. Physical models of plant development. *Annu. Rev. Cell*
Dev. Biol. **2014**, *30*, 59–78.

57.   Dumond, M.; Boudaoud, A. Physical Models of Plant Morphogenesis. In *Mathematical*

Modelling in Plant Biology; Morris, R.J., Ed.; Springer International Publishing: Cham, 2018;
pp. 1–14 ISBN 978-3-319-99070-5.

58. Alt, S.; Ganguly, P.; Salbreux, G. Vertex models: from cell mechanics to tissue morphogenesis.
Philos. Trans. R. Soc. B Biol. Sci. **2017**, *372*.

59. Dyson, R.J.; Vizcay-Barrena, G.; Band, L.R.; Fernandes, A.N.; French, A.P.; Fozard, J.A.;
Hodgman, T.C.; Kenobi, K.; Pridmore, T.P.; Stout, M.; et al. Mechanical modelling quantifies
the functional importance of outer tissue layers during root elongation and bending. *New*
Phytol. **2014**, *202*, 1212–1222.

60. Forouzesh, E.; Goel, A.; Mackenzie, S.A.; Turner, J.A. In vivo extraction of Arabidopsis cell
turgor pressure using nanoindentation in conjunction with finite element modeling. *Plant J.*
2013, *73*, 509–520.

61. Maurel, C.; Boursiac, Y.; Luu, D.-T.; Santoni, V.; Shahzad, Z.; Verdoucq, L. Aquaporins in Plants.
Physiol. Rev. **2015**, *95*, 1321–1358.

62. Klepikova, A.V.; Kasianov, A.S.; Gerasimov, E.S.; Logacheva, M.D.; Penin, A.A. A high resolution
map of the Arabidopsis thaliana developmental transcriptome based on RNA-seq profiling.
Plant J. Cell Mol. Biol. **2016**, *88*, 1058–1070.

63. Péret, B.; Li, G.; Zhao, J.; Band, L.R.; Voß, U.; Postaire, O.; Luu, D.-T.; Da Ines, O.; Casimiro, I.;
Lucas, M.; et al. Auxin regulates aquaporin function to facilitate lateral root emergence. *Nat.*
Cell Biol. **2012**, *14*, 991–998.

64. Tsugawa, S.; Hervieux, N.; Kierzkowski, D.; Routier-Kierzkowska, A.-L.; Sapala, A.; Hamant, O.;
Smith, R.S.; Roeder, A.H.K.; Boudaoud, A.; Li, C.-B. Clones of cells switch from reduction to
enhancement of size variability in Arabidopsis sepals. *Development* **2017**, *144*, 4398–4405.

65. Sapala, A.; Runions, A.; Routier-Kierzkowska, A.-L.; Das Gupta, M.; Hong, L.; Hofhuis, H.;
Verger, S.; Mosca, G.; Li, C.-B.; Hay, A.; et al. Why plants make puzzle cells, and how their
shape emerges. *eLife* **2018**, *7*.

66. LeGoff, L.; Rouault, H.; Lecuit, T. A global pattern of mechanical stress polarizes cell divisions
and cell shape in the growing Drosophila wing disc. *Development* **2013**, *140*, 4051–4059.

67. Campinho, P.; Behrndt, M.; Ranft, J.; Risler, T.; Minc, N.; Heisenberg, C.-P. Tension-oriented
cell divisions limit anisotropic tissue tension in epithelial spreading during zebrafish epiboly.
Nat. Cell Biol. **2013**, *15*, 1405–1414.

68. Mao, Y.; Tournier, A.L.; Hoppe, A.; Kester, L.; Thompson, B.J.; Tapon, N. Differential
proliferation rates generate patterns of mechanical tension that orient tissue growth. *EMBO*
J. **2013**, *32*, 2790–2803.

69. Louveaux, M.; Julien, J.-D.; Mirabet, V.; Boudaoud, A.; Hamant, O. Cell division plane
orientation based on tensile stress in Arabidopsis thaliana. *Proc. Natl. Acad. Sci.* **2016**, *113*,
E4294–E4303.

70. Gibson, W.T.; Gibson, M.C. Chapter 4 Cell Topology, Geometry, and Morphogenesis in
Proliferating Epithelia. In *Current Topics in Developmental Biology*; Current Topics in
Developmental Biology; Academic Press, 2009; Vol. 89, pp. 87–114.

71. Long, Y.; Boudaoud, A. Emergence of robust patterns from local rules during plant
development. *Curr. Opin. Plant Biol.* **2018**, *47*, 127–137.

72. Vargas-Garcia, C.A.; Ghusinga, K.R.; Singh, A. Cell size control and gene expression
homeostasis in single-cells. *Curr. Opin. Syst. Biol.* **2018**, *8*, 109–116.

73. Cutler, S.R.; Ehrhardt, D.W.; Griffitts, J.S.; Somerville, C.R. Random GFP∷cDNA fusions enable

visualization of subcellular structures in cells of Arabidopsis at a high frequency. *Proc. Natl.*
Acad. Sci. **2000**, *97*, 3718–3723.

74. Stanislas, T.; Hamant, O.; Traas, J. Chapter 11 - In-vivo analysis of morphogenesis in plants. In
Methods in Cell Biology; Lecuit, T., Ed.; Cell Polarity and Morphogenesis; Academic Press,
2017; Vol. 139, pp. 203–223.

75. Reuille, P.B. de; Bohn-Courseau, I.; Godin, C.; Traas, J. A protocol to analyse cellular dynamics
during plant development. *Plant J.* **2005**, *44*, 1045–1053.

76. Legland, D.; Arganda-Carreras, I.; Andrey, P. MorphoLibJ: integrated library and plugins for
mathematical morphology with ImageJ. *Bioinformatics* **2016**, *32*, 3532–3534.

77. Mirabet, V.; Dubrulle, N.; Rambaud, L.; Beauzamy, L.; Dumond, M.; Long, Y.; Milani, P.;
Boudaoud, A. NanoIndentation, an ImageJ Plugin for the Quantification of Cell Mechanics.
Methods Mol. Biol. **in press**.

78. Sassi, M.; Ali, O.; Boudon, F.; Cloarec, G.; Abad, U.; Cellier, C.; Chen, X.; Gilles, B.; Milani, P.;
Friml, J.; et al. An auxin-mediated shift toward growth isotropy promotes organ formation at
the shoot meristem in Arabidopsis. *Curr. Biol. CB* **2014**, *24*, 2335–2342.

79. Kremer, A.; Lippens, S.; Bartunkova, S.; Asselbergh, B.; Blanpain, C.; Fendrych, M.; Goossens,
A.; Holt, M.; Janssens, S.; Krols, M.; et al. Developing 3D SEM in a broad biological context. *J.*
Microsc. **2015**, *259*, 80–96.

80. Kiss, A.; Moreau, T.; Mirabet, V.; Calugaru, C.I.; Boudaoud, A.; Das, P. Segmentation of 3D
images of plant tissues at multiple scales using the level set method. *Plant Methods* **2017**,
13, 114.

81. Reuille, P.B. de; Routier-Kierzkowska, A.-L.; Kierzkowski, D.; Bassel, G.W.; Schüpbach, T.;
Tauriello, G.; Bajpai, N.; Strauss, S.; Weber, A.; Kiss, A.; et al. MorphoGraphX: A platform for
quantifying morphogenesis in 4D. *eLife* **2015**, *4*, e05864.

82. Schmidt, T.; Pasternak, T.; Liu, K.; Blein, T.; Aubry-Hivet, D.; Dovzhenko, A.; Duerr, J.; Teale, W.;
Ditengou, F.A.; Burkhardt, H.; et al. The iRoCS Toolbox – 3D analysis of the plant root apical
meristem at cellular resolution. *Plant J.* **2014**, *77*, 806–814.