Journal:

Title: **The inhibition of automatic imitation: a meta-analysis and synthesis of fMRI studies**

Kohinoor M. Darda and Richard Ramsey

Wales Institute for Cognitive Neuroscience, School of Psychology, Bangor University,

Bangor, Gwynedd, Wales, LL57 2AS, United Kingdom

Corresponding author: r.ramsey@bangor.ac.uk

Abstract

Humans copy other people without their conscious awareness, a behaviour known as automatic imitation. Although automatic imitation forms a key part of daily social interactions, we do not copy other people indiscriminately. Instead, we control imitative tendencies by prioritising some actions and inhibiting others. To date, neuroimaging studies investigating the control of automatic imitation have produced inconsistent findings. Some studies suggest that imitation control relies on a domain-specific neural circuit related to social cognition (the theory-of-mind network). In contrast, other studies show engagement of a domain-general neural circuit that is engaged during a diverse range of cognitive control tasks (the multiple demand network). Given the inconsistency of prior findings, in the current paper we avoided problems associated with interpreting individual studies by performing a meta-analysis. To do so, we used a multi-level kernel density analysis to quantitatively identify consistent patterns of activation across functional magnetic resonance imaging studies investigating the control of imitation. Our results show clear and consistent evidence across studies that the control of automatic imitation is guided by brain regions in the multiple demand network including dorsolateral frontoparietal cortex. In contrast, there was only limited evidence that regions in the theory of mind network were engaged. Indeed, medial prefrontal cortex showed no consistent engagement and right temporoparietal junction engagement may reflect spatial rather than imitative control. As such, the current meta-analysis reinforces the role of domain-general control mechanisms and questions the role of domain-specific processes in regulating imitative tendencies. Consequently, neurocognitive models of imitation need updating to place more emphasis on domain-general control mechanisms, as well as to consider more complex organisational structures of control, which may involve contributions from multiple cognitive systems.

Introduction

In day-to-day social interactions, humans copy or are copied, even without their conscious awareness. This involuntary tendency to imitate others' gestures, speech patterns, and postures, is known as automatic imitation (Heyes, 2011). Imitative behaviours are present in almost all animal taxa (Iacoboni, 1999) and have been considered an important evolutionary inheritance mechanism (Boyd & Richerson, 1995; Whiten & Van Schaik, 2007). In many species, for example, a variety of behavioural phenotypes are found to spread and transmit across generations with the help of imitation (Shea, 2009). Given the influence of imitation on the development of social species, researchers across a broad spectrum of the life sciences have started to investigate the psychological and biological mechanisms that underpin imitation. For example, over the last 20 years, researchers have used functional magnetic resonance imaging (fMRI) in order to better understand the neural underpinnings of the control of automatic imitative tendencies. However, these studies have provided mixed findings regarding the contributions of domain-general or domain-specific neural networks in imitation control. The current paper, therefore, meta-analyses fMRI studies to date on the control of automatic imitation in order to provide a combined quantitative estimate of the extant evidence of many individual studies (Lipsey & Wilson, 2001).

In the last two decades, automatic imitation has been widely studied with an attempt to interconnect different disciplines like cognitive science, social psychology, evolutionary biology, and cognitive neuroscience (Prinz & Meltzoff, 2002; Chartrand & Bargh, 1999; Byrne & Russon, 1998). This convergence across multiple disciplines has allowed for a range of perspectives on imitation to emerge in which theory and empirical data can strengthen each other. In social psychology, automatic imitation has been studied in naturalistic social interactions (Chartrand & Lakin, 2013). Evidence

primarily from this field suggests that automatic imitation functions as a "social glue" as it strengthens feelings of affiliation, and co-operative behaviour between interacting partners (Chartrand & Lakin, 2013). Consequent imitation behaviour is also moderated by other variables including, but not limited to, personality variables, self-construal, goal to affiliate or disaffiliate, cultural and social contexts, as well as the similarity, familiarity, and status of the person being imitated (Chartrand & Lakin, 2013; Caspers et al., 2010; Duffy & Chartrand, 2015; Figure 1).

Even though automatic imitation seems to be an important evolutionary behaviour that facilitates social situations, we do not always copy others' behaviours. In many situations, imitation can be maladaptive, and it is essential to circumvent the tendency to automatically imitate (Cross & Iacoboni, 2014; Cross, Torrisi, Losin, & Iacoboni, 2013; van Schie, van Waterschoot, & Bekkering, 2008; Newman- Norlund, van Schie, van Zuijlen, & Bekkering, 2007). This need to regulate imitative tendencies indicates the existence of a selection mechanism that inhibits unwanted actions, and prioritises alternatives (Brass et al., 2009). Thus, imitation control can be divided into at least two component processes – action representation and action selection. We observe an interaction partner and their actions (representation), and then select the action that needs to be executed (selection).

In cognitive psychology and neuroscience, researchers have generally used computer-based reaction-time (RT) measures of the inhibition of automatic imitation (Brass et al., 2000; Stürmer et al., 2000). One of the most commonly used tasks in this field consists in making finger movements while simultaneously observing a compatible or incompatible finger movement (Brass et al., 2000). For example, participants may be asked to make a finger movement in response to an imperative cue i.e. they are

instructed to lift their index finger when they see a number '1' on screen, and their middle finger when they see a number '2.' Simultaneously, participants also observe a task-irrelevant index or middle finger movement, which is compatible or incompatible with their own response. Other variants of this task include hand opening and closing movements instead of finger movements (Press et al., 2005; Wang et al., 2011) or pre-specifying the participant's response before the imperative cue (i.e. participants are asked to always lift their index finger when they see a finger movement; Brass, Bekkering & Prinz, 2001; Heyes et al., 2005). In these variants as well, participants observe a hand or finger movement which is compatible or incompatible with their own response. Irrespective of the task used, greater cognitive resources are required when inhibiting movements incompatible to one's own responses, thus leading to greater RTs (Heyes, 2011; Brass & Heyes, 2005). The difference between the incompatible and compatible conditions (referred to as the general compatibility effect) is said to be a measure of imitation control (Heyes et al., 2005; Heyes, 2011).

To date, a number of neuroimaging studies have investigated the neural mechanisms of imitation control using RT paradigms. However, the evidence demonstrating the extent to which RT measures of imitation control rely on domain-general or domain-specific neural networks is mixed. Domain-specific processes operate on particular types of stimuli or aspects of cognition, while domain-general processes operate across a range of stimuli and tasks (Barett, 2012; Spunt & Adolphs, 2017). One of the prevailing theories of automatic imitation proposes that imitation control relies on a domain-specific neural circuit related to social cognition (Brass et al., 2009). This "specialist" theory has gained traction with evidence from patient and neuroimaging data pointing to the engagement of two key candidate regions – the

anterior medial prefrontal cortex (mPFC) and the right temporoparietal junction (rTPJ) (Brass & Heyes, 2005; Brass et al., 2009). For example, mPFC and rTPJ have been engaged in human brain imaging investigations of imitation inhibition (Brass et al., 2001; 2005; 2009; Spengler et al., 2009; Wang et al., 2011). Brass and colleagues further proposed a dissociation of roles for the mPFC and rTPJ during imitation control - the rTPJ distinguishes between self- and other-generated actions, and the mPFC enforces the self-generated action when faced with conflict from an action representation generated by another agent (Brass et al., 2009). In addition, patients with frontal lobe lesions show disrupted imitation inhibition behaviour (Brass et al., 2003; Spengler et al., 2010) and an increased tendency to automatically imitate even when they are clearly instructed to not do so (Lhermitte et al., 1986). More evidence for the involvement of rTPJ comes from neuro-stimulation studies: inhibiting the activity in the rTPJ by transcranial magnetic stimulation (TMS) interfered with imitative responses impairing imitation inhibition (Hogeveen et al., 2014; Sowden & Catmur, 2015). Irrespective of the method used, it is worth noting that, to date, there have only been a small number of studies implicating mPFC and rTPJ in the control of imitation. Moreover, these studies have used relatively small sample sizes between 10 and 25 participants and there have been few, if any, direct replications. Therefore, the sum total of evidence for mPFC and rTPJ engagement during imitation control is suggestive rather than compelling.

Along with imitative control, neuroimaging findings suggest mPFC and rTPJ are also engaged in a variety of socio-cognitive tasks that are associated with theory of mind, including distinguishing between self from other, perspective taking, as well as attributing beliefs, desires and attitudes to others (ToM; Gallagher et al., 2000; Amodio & Frith, 2006; Ruby & Decety, 2001; Aichhorn et al., 2006; Decety et al., 2002;

Santiesteban et al., 2012; Brass et al., 2009; Spengler et al., 2010). Based on these findings, self-other control processes have thus been proposed as a candidate mechanism for a range of socio-cognitive functions. For example, it is important to inhibit one's own perspective or mental state and enhance that of the other when empathising with others, taking their perspective, or engaging a successful theory-of-mind (de Guzman et al., 2016; Sowden & Shah, 2014). Further, atypical self-other control has been linked to disorders characterised by social dysfunction including autism and schizophrenia (Cook and Bird , 2012; Ferri et al., 2012). Overall, this evidence suggests that in imitation control, it is crucial to inhibit the representation of the other's action, and enforce your own, and this mechanism is guided by a domain-specific neural circuit unique to social cognition (Brass et al., 2009).

In contrast to this "specialist" view of imitation control, however, "generalist" theories of imitation suggest that the inhibition of automatic imitation does not differ from any other pre-potent tendencies or general cognitive functions (Heyes, 2011; Cooper et al., 2013). Multiple cognitive control tasks like the Flanker, Stroop, and Simon tasks, which require the inhibition of automatic overlearned response tendencies, have been found to engage a domain-general control network identified in the dorsolateral fronto-parietal cortices (Aron et al., 2014; Bunge et al., 2002; Hazeltine et al., 2007; Nee et al., 2007; Wager et al., 2005). This network is also called the multiple demand (MD) network as it is engaged across a diversity of mental operations (Duncan et al., 2010). Across studies that investigate imitation inhibition, some have found engagement of the mPFC and rTPJ (Brass et al., 2001; 2005; 2009; Spengler et al., 2009), whereas others show engagement of the MD network (Bien, Roebroeck, Goebel, & Sack, 2009; Crescentini, Mengotti, Grecucci, & Rumiati, 2011; Cross & Iacoboni, 2013; Mengotti,

Corradi-Dell'Acqua, & Rumiati, 2012; Marsh et al., 2016). However, most previous fMRI studies have been limited by low statistical power and small sample sizes. More recently, a multi-experiment study using larger sample sizes (N=28, N=50) and a functional region of interest (fROI) approach that bolsters statistical power and functional sensitivity has shown that imitation control engages only the MD network, and not mPFC or rTPJ (Darda, Butler & Ramsey, 2018). Indeed, even with an *a priori* power analysis ensuring 80% power to detect medium effect sizes, Darda and colleagues (2018) did not even find a directional trend to suggest that the ToM network was directly engaged during imitation control.

As mentioned before, imitation control can be divided into at least two component processes – action representation and action selection. The above review of literature suggests two possible neural mechanisms as being key to action selection during imitation control. On one hand, during imitation control, the neural representation generated by the observed person's action is inhibited, and the self-generated action is selected and enforced and this selection mechanism engages a domain-specific neural network i.e. the mPFC and rTPJ. On the other hand, the selection mechanism may be guided by a domain-general neural network i.e. the MD network. In both possible mechanisms, the input is the same i.e. the observed person and action may engage domain-specific socio-perceptual neural circuits. However, the difference lies in the selection or control mechanism that underlies the inhibition of automatic imitative tendencies which finally leads to consequent behaviour (see graphical representation in Figure 1).

The question of interest for the current meta-analysis, therefore, lies at the selection stage of imitation control with the evidence to date for engagement of domain-

specific and domain-general neural networks being inconsistent. Even though the most statistically powerful fMRI study to date only shows the engagement of the MD network (Darda et al., 2018), the interpretation of individual studies remains limited in scope for several reasons. First, many single studies are likely to be underpowered leading to missed or spurious results (Button et al., 2013). Second, empirical work involves design choices that strongly influence results, making it harder to generalise effects across analysis pipelines and differing experimental procedures (Carp, 2012). Given the inconsistency of prior findings and the absence of a quantitative synthesis of evidence, taking a meta-analytical approach to further investigate the neural basis of imitation has many benefits. As such, by means of a meta-analysis, the current paper enables the detection of consistent patterns of activation across studies.

In order to quantify the consistency and specificity of regional activation for imitation control across studies, we performed a multi-level kernel density analysis (MKDA; see Methods and Materials for details). We included all fMRI studies (N=11) investigating imitation control using the RT measure of imitation inhibition (see Table 1). Our primary measure aimed to quantify the consistency of region engagement across studies with particular focus on the engagement of the ToM network and the MD network. Given the prior mixed findings across studies, this meta-analysis aimed to quantify the extent to which ToM, MD or both neural networks may be engaged when during the inhibition of automatic imitation.

We also ran two more exploratory analyses, which were based on a small subset of the total studies. The most common measure of imitation inhibition, the general compatibility effect, also includes a spatial component (Heyes, 2011). In order to measure imitative compatibility more specifically, therefore, imitative and spatial

effects need to be dissociated (Gowen et al., 2016; Boyer et al., 2012; Catmur & Heyes, 2011). However, only a few fMRI studies have measured the imitative compatibility effect independent of the spatial component (Darda et al., 2018; Marsh et al., 2016; Cross et al., 2013). This makes it difficult to interpret the roles of the ToM (mPFC and rTPJ) and MD networks in imitation control – their engagement could reflect both social (imitative) and/or non-social (spatial) control. Indeed, the rTPJ has been previously associated with orienting to both social and non-social stimuli (Corbetta et al., 2008; Thiel et al., 2004). Thus, given that only a few studies have dissociated between imitative (N=3) and spatial compatibility (N=4) effects, we also ran two further exploratory MKDAs in order to quantify consistency of patterns across studies for both imitative and spatial compatibility effects.

Methods and Materials

Literature search and data collection

In the current paper, we follow recent guidelines put forward for meta-analysing neuroimaging studies (Muller et al., 2018). FMRI studies exploring the inhibition of automatic imitative tendencies were searched for on the online database PubMed, as well as the article search engine Google Scholar. Combinations of keywords including 'imitation inhibition,' 'fMRI,' 'imitation,' 'automatic imitation,' and 'imitation control' were used to identify relevant literature (prior to January 2018). A total of 14 studies were found. We rejected studies if the primary method of investigation was not fMRI, if the study did not report results in stereotactic coordinate space (either Montreal neurological Institute (MNI) or Talaraich coordinates) (N=1; Bien et al., 2009), if reported results were based on region-of-interest (ROI) analyses, and the study did not report whole-brain analysis coordinates either in the main article or in

supplementary materials (or we could not obtain them from the authors) (N=1; Brass et al., 2009), and if the study involved children or atypical populations (and the coordinates for controls were not reported separately) (N=1; Spengler, Bird, & Brass, 2010).

A wide variety of contrasts are used in studies that investigate the inhibition of automatic imitation. However, in order to minimise heterogeneity, studies that used a paradigm that was not based on or was not conceptually similar to the Brass et al. (2000) measure of the inhibition of automatic imitation were also excluded. Thus, 11 studies with a total of 276 participants were included in the meta-analysis (see Table 1).

Even though our main analysis was on the general compatibility effect, we also ran two separate meta-analyses for spatial and imitative compatibility. Table 2 summarises the contrasts used in the current meta-analysis for general, spatial, and imitative compatibility effects. A total of 12 contrasts across 11 studies with 131 foci were used for general compatibility, 4 contrasts across 4 studies with 42 foci were used for spatial compatibility, and a total of 3 contrasts across 3 studies with 20 foci were used for imitative compatibility.

Data analysis

All analyses in the current paper were performed in MatlabR2015b (Mathworks, Naticks, MA) using the MKDA toolbox developed by Wager et al., 2007; http://wagerlab.colorado.edu). MKDA is an analysis technique that uses a random effects model to assess convergence across studies. This allows for assessing convergence across studies as opposed to between individual foci (as implemented in classical meta-analysis techniques that use fixed effects analyses). Thus, results are not biased by a small number of individual studies. Further, each contrast is weighted by

the sample size and study quality (i.e. whether the study used fixed or random effects model; Wager et al., 2007; Kober and Wager, 2010).

MKDA was performed on all three compatibility types separately. Before performing the analyses, we extracted the following information from each study and included it in our database: authors, year of publication, sample size, task contrasts, fixed or random effects model, and MNI or Talaraich co-ordinates. Co-ordinates reported in Talairach space were converted to MNI stereotactic space using Lancaster transformation (tal2icbm transform; Lancaster et al., 2007). Peak coordinates from each contrast map were then convolved with a 10mm spherical kernel in order to create a contrast indicator map (CIM). The resulting voxels within 10 mm of the peak were deemed "significant" and given a value of one; other voxels were given a value of zero which indicated no significant effect. A density map was then created by averaging the indicator maps, weighted by sample size, and whether the study used a fixed or random effects model. More specifically, as recommended by Wager and colleagues (Wager, Lindquist, & Kaplan, 2007), this density map was weighted by the square root of the sample size of the study, and then multiplied by an adjustment factor of 1 for random effects analysis, and .75 for a fixed effects analysis.

Each voxel of the density map was given a density statistic P. P stands for the proportion of contrasts included in the analysis that show activity within 10mm of the voxel. A Monte Carlo simulation (with 5000 iterations) was then carried out in order to identify voxels that had a P-statistic that was higher than the frequency predicted by chance. This was tested against the null hypothesis that activated regions in the resulting pairwise contrast maps (from the 5000 iterations) were randomly distributed across the brain. To test for the significance of the cluster size, a similar procedure was used. This allowed for the identification of a threshold for cluster size at which a specific

number of voxels needed to be activated contiguously so that the cluster could be deemed significant.

In order to maximise sensitivity in testing our hypotheses, we report results using two thresholding techniques. One thresholding technique is based on height and the other is based on cluster size. For the weighted P-statistic (height-based threshold), the family wise error (FWE) corrected threshold is the proportion of studies which yielded activity within 10 mm of a voxel that showed a higher P-statistic than the maximum P-statistic across 95% of the Monte Carlo maps. For the cluster size threshold, the FWE corrected threshold is the contiguous voxels observed at two different thresholds (p<.001 and p<.01) whose cluster size is more than the extent of clusters found across 95% of the Monte Carlo maps. We use two cluster-based thresholds in order to also detect regions that show a lower response in magnitude over a larger cluster size both at more stringent (p<.001) and less stringent (p<.01) thresholds. Voxels that exceed the height-based threshold in our analysis appear on the resulting maps in Figure 2 in yellow, and those that exceed the cluster extent-based threshold appear in orange (p<.001) and red (p<.01).

In the resulting table (Table 3), peak activation foci that pass the height-based threshold are reported. If activations do not pass the height-based threshold, foci of the cluster-extent-based thresholding are reported. The number of voxels in each cluster that survived height-based and/or extent-based thresholding is also reported. Resulting coordinates were localised using the SPM Anatomy Toolbox (Eickhoff et al., 2005).

Results

For general compatibility effect, across 12 contrasts from 11 studies, consistent activation was found in right MFG and right supramarginal gyrus extending to the temporo-parietal junction (see Table 3; Figure 2A). These clusters survived both height-

based and the more stringent extent-based thresholding (p<.001). Activation was also found in the right IPL and right SFG, which survived both height and the less stringent extent-based thresholding (p<.01). Activation in the left insula and left IPL survived the more stringent extent-based threshold (p<.001), but not the height-based threshold.

We ran two further MKDAs separately for spatial and imitative compatibility. For spatial compatibility, across 4 contrasts from 4 studies, we found consistent activation that withstood the height-based thresholding in the left IPL and the right SFG (see Table 3; Figure 2B). No regions showed cluster-based thresholding. For imitative compatibility, across 3 contrasts from 3 studies, we found consistent activation in the left IPL that survived height-based (see Table 3; Figure 2C). Activation was also found in the right IPL, which withstood height-based as well as the less stringent extent-based thresholding (p<.01).

These density maps showing regions that withstood both height and/or cluster-extent thresholding for each compatibility type were then overlaid with the ToM and MD network masks separately. The ToM network mask consisted of four parcels including the dorsal, medial, and ventral prefrontal cortex (DMPFC, MMPFC, VMPFC), and the right temporo-parietal junction (rTPJ), which have previously been implicated in mentalising or theory-of-mind (Dufour et al., 2013). For the MD network mask, 16 parcels were used which included areas in bilateral superior and inferior parietal lobules (SPL, IPL), inferior parietal sulcus (IPS), inferior and middle frontal gyrus (IFG, MFG), precentral gyrus (PrecG), insula (Ins), and the supplementary motor area (SMA) (available at: https://evlab.mit.edu/funcloc/download-parcels). Overlay of the density maps with the ToM and MD network masks allowed for identification of overlap between regions that were consistently activated in the MKDA and the ToM and MD networks (Figure 3). For all compatibility types (general, imitative and spatial), all

regions that passed height or extent-based thresholding overlapped with regions in the MD network (Figure 3A). Additionally, one cluster, which showed consistent activation for general compatibility, also overlapped with the right TPJ in the ToM network (Figure 3B). There was no overlap with the mPFC node of the ToM network for any compatibility type.

In order to break down the role of the right TPJ in general compatibility, we performed a further, more exploratory analysis. We compared peak coordinates from prior studies with a right TPJ mask, which has been previously implicated in theory-of-mind (Dufour et al., 2013). To do so, the ToM network mask for rTPJ was overlaid with the contrast indicator maps of all studies used for imitative (N=3) and spatial (N=4) compatibility. The contrast indicator maps include 10mm spherical kernels around peak coordinates of each contrast. This allows coordinates from prior imitative and spatial compatibility contrasts to be displayed without any thresholding restrictions and overlaid with the rTPJ node of the ToM network. Figure 4 shows overlap between contrast indicator maps for general compatibility and spatial compatibility with the right TPJ node of the ToM network mask. By contrast, there is no overlap between contrast indicator maps for imitative compatibility and the same right TPJ mask.

Discussion

In the current paper, a meta-analysis of fMRI studies was performed to quantify the consistency and specificity of regional activation during the inhibition of automatic imitation. Our findings largely supported the "generalist" rather than the "specialist" view of action selection when controlling the urge to imitate. Indeed, we found consistent engagement of the dorsolateral frontoparietal cortices when observing an action that conflicted with a current motor intention, which overlapped considerably

with regions associated with the MD network. In addition, we found limited evidence for the engagement of brain regions associated with ToM, such as mPFC and rTPJ, during imitation control. Indeed, mPFC showed no consistent engagement across studies and rTPJ showed engagement, but it is currently unclear if this engagement reflects spatial or social control. Thus, our results provide clear support for the engagement of a domain-general neural network during imitation control and only limited evidence for the engagement of a domain-specific neural network that is tied to social cognition.

Studies investigating the neural correlates of imitation control have to date shown mixed evidence for the engagement of domain-general and domain-specific neural networks in imitation inhibition. While some studies have found engagement of the mPFC and rTPJ (Brass et al., 2001; 2005; 2009; Spengler et al., 2009), others show engagement of the MD network (Bien, Roebroeck, Goebel, & Sack, 2009; Marsh et al., 2016; Darda et al., 2018). The current MKDA demonstrated that brain regions in the multiple demand network are reliably and consistently engaged across studies that investigate imitation inhibition using the general compatibility effect. Brain regions in the MD network are also engaged for imitative (bilateral IPL) and spatial compatibility effects (left IPL, right SFG). Thus, our findings suggest that brain regions that are engaged across a range of cognitive control tasks are also reliably engaged when controlling the automatic tendency to imitate others, as measured by general and imitative compatibility effects.

In contrast, evidence supporting the engagement of a domain-specific neural circuit that is central to social cognition and includes mPFC and rTPJ is less consistent. Brass et al. (2009) proposed that the rTPJ was involved in distinguishing between self- and other-generated actions, whereas the mPFC was engaged when enforcing the

correct action. However, the current MKDA did not find any evidence of anterior mPFC

engagement for either general, spatial, or imitative compatibility effects. An absence of

mPFC engagement for imitation control across studies is thus inconsistent with the

hypothesis that a specific neural system related to social cognition is also engaged

during the inhibition of automatic imitation (Brass et al., 2009).

In contrast to mPFC, rTPJ was consistently engaged by the general compatibility

effect. The role of rTPJ in imitation control, however, is difficult to interpret given its

functional heterogeneity and involvement in both social and non-social processes

(Corbetta et al., 2008; Krall et al., 2015; 2016; Lee & McCarthy, 2016; Schuwerk et al.,

2017). Indeed, in the largest and most sensitive fMRI study of imitation inhibition to

date, Darda and colleagues (2018) showed no engagement of rTPJ for the imitative

compatibility effect, but engagement of rTPJ for general and spatial compatibility

effects. Thus, in the current meta-analysis, in order to break down the role of rTPJ in

imitation control as measured by the general compatibility effect, we overlapped rTPJ

with unthresholded spatial and imitative compatibility effect contrast indicator maps

separately. rTPJ did not overlap with the imitative compatibility effect map, but there

was a partial overlap for the spatial compatibility effect. Current findings from fMRI

studies, therefore, provide limited evidence for the univariate engagement of rTPJ in the

control of imitative tendencies. In contrast, current fMRI findings provide more

evidence that rTPJ is involved in resolving spatial conflict, which is in keeping with

patient work (Vallar & Perani, 1987; Valler et al., 1993), as well as evidence using

spatial cueing tasks (Thiel et al., 2004; Corbetta et al., 2008).

fMRI is only one form of measurement, however, and it is important to consider

how these findings mesh with results from other neuroscience techniques. For instance,

neurostimulation studies have implicated rTPJ in imitation control (Santiesteban et al.,

2015; Bardi et al., 2017). Using repetitive transcranial magnetic stimulation (TMS), dampening of activity in the rTPJ interfered with imitative, but not spatial responses (Hogeveen et al., 2014; Sowden & Catmur, 2015), whereas excitatory stimulation of the rTPJ by anodal transcranial direct current stimulation (tDCS) caused increased performance on the imitation task (Santiesteban et al., 2012). Further, in patients with lesions in the temporoparietal junction area, imitation inhibition deficits have been found to correlate with deficits in visual and cognitive perspective taking tasks, further supporting the role of rTPJ in imitation control (Spengler et al., 2010). Thus, there seems to be a discrepancy between neurostimulation and patient studies, and results from the current meta-analysis of fMRI studies. The evidence from neurostimulation and patient studies for the engagement of rTPJ in imitation control is, however, limited to a few studies with small sample sizes. Therefore, for more confirmatory evidence, future investigations with pre-registered and adequately powered replications are essential (Munafo et al., 2017; Zwaan et al., 2018; Nelson et al., 2018). In addition, it is also possible that the role of rTPJ in imitation control cannot be captured by univariate measurements and a more complex neural organisation is at play during imitation control.

The lack of consistent activation of mPFC in the current meta-analysis and a difficulty in interpreting the role of rTPJ have implications for "specialist" theories of imitation. "Specialist" theories suggest that based on a dedicated neural circuit for social cognition, self-other control is crucial for the regulation of imitation, empathy, autism, and theory-of-mind (Brass et al., 2009; de Guzman et al., 2016; Sowden & Shah, 2016). However, more recent behavioural evidence suggests that imitation may not vary as a function of autistic-like traits or empathy, thus questioning the reliance of imitation inhibition on a distinctly social mechanism (Butler et al., 2015; Cracco et al., 2018;

Genschow et al., 2017). Instead of a distinctly social mechanism, imitation control may involve domain-general cognitive control mechanisms, which are also engaged during the control of other non-social pre-potent response tendencies (Heyes, 2011; Cooper et al., 2012). Indeed, the dual-route model of automatic imitation proposed by Heyes (2011) can explain the control of automatic imitative tendencies without assuming a reliance on a self-other distinction. The model suggests that like other stimulus-response compatibility tasks, imitation control is mediated by long-term stimulus-response associations which are a product of learning. In line with this, the computational model put forth by Cooper et al. (2012) further substantiates this notion by demonstrating that spatial and imitative compatibility effects depend on similar cognitive processes, and any behavioural differences are accounted for by different sets of input nodes for spatial and imitative effects in a general dual-route framework (but see Berthental & Scheutz (2013) for a critique of this model).

Even though it is possible that imitation and spatial compatibility rely on a partly shared set of cognitive processes, this does not address the question of whether these processes also rely on similar or distinct neurobiological mechanisms. The current meta-analysis suggests that the selection mechanism in imitation inhibition is guided by a domain-general multiple demand system, which is also engaged during the inhibition of other non-social external influences. However, a lack of engagement of mPFC (and possibly rTPJ) in imitation control does not imply that they do not also play a regulatory role in imitation control. For example, mPFC has been demonstrated to exert a top-down influence during modulation of imitation via direct gaze (Wang et al., 2011). In addition, rTPJ showed a higher response when an interaction partner was believed to be human and looked human compared to when these animacy cues were absent (Klapper et al., 2014). These findings suggest that mPFC and rTPJ may play a

regulatory role in imitation control and may be functionally connected to other networks without being directly engaged (Burnett & Blakemore, 2009). The current findings suggest that future work should postulate and test more complex models of imitation control, which extend beyond the operations of the theory of mind network.

In a similar manner, other socio-perceptual circuits, which extend beyond the MD network, may also be involved when inhibiting automatic imitative tendencies. That is, we may use a domain-general selection mechanism to select between both social and non-social stimuli, but other domain-specific person and action perception systems are likely to be selectively engaged during action representation component of imitation (Caspers et al., 2010; Downing & Peelen, 2011; Kanwisher, 2010). For example, as mentioned in the introduction (also see Figure 1), domain-specific action observation and person perception networks may functionally interact with the MD network during imitation control. It could be these links between domain-specific and domain-general neural architectures that distinguish between social and non-social processes. Thus, the engagement of domain specific and domain general neural networks in imitation control may be more complicated that what current models of imitation suggest.

In conclusion, the current meta-analysis provides evidence that the selection mechanism when inhibiting automatic imitative tendencies is guided by the regions of the domain-general multiple demand network rather than a domain-specific system related to social cognition. Our meta-analysis questions the role of mPFC and right TPJ in imitation control, and suggests that current neurocognitive models of imitation control need further revision in order to account for the more complex nature of functional interplay between domain-general and domain-specific systems.

References

Aichhorn, M., Perner, J., Kronbichler, M., Staffen, W., & Ladurner, G. (2006). Do visual perspective tasks need theory of mind?. Neuroimage, 30(3), 1059-1068.

Amodio DM, Frith CD. 2006. Meeting of minds: the medial frontal cortex and social cognition. Nat Rev Neurosci. 7:268-277.

Aron AR, Robbins TW, Poldrack RA. 2014. Inhibition and the right inferior frontal cortex: one decade on. Trends Cogn Sci. 18:177-185.

Bardi, L., Gheza, D., & Brass, M. 2017. TPJ-M1 interaction in the control of shared representations: new insights from tDCS and TMS combined. NeuroImage, 146:734-740.

Bargh, J. A., & Chartrand, T. L. (1999). The unbearable automaticity of being. American psychologist, 54(7), 462.

Barrett, H. C. (2012). A hierarchical model of the evolution of human brain specializations. Proceedings of the National Academy of Sciences, 109(Supplement 1), 10733-10740. doi: 10.1073/pnas.1201898109

Bertenthal, B. I., & Scheutz, M. (2013). In praise of a model but not its conclusions: Commentary on Cooper, Catmur, and Heyes (2012). Cognitive science, 37(4), 631-641.

Bien N, Roebroeck A, Goebel R, Sack AT. 2009. The brain's intention to imitate: the neurobiology of intentional versus automatic imitation. Cereb Cortex. 19:2338-51.

Boyd, R., & Richerson, P. J. (1996, January). Why culture is common, but cultural evolution is rare. In Proceedings-British Academy (Vol. 88, pp. 77-94). OXFORD UNIVERSITY PRESS INC.

Boyer TW, Longo MR, Bertenthal BI. 2012. Is automatic imitation a specialized form of stimulus–response compatibility? Dissociating imitative and spatial compatibilities. Acta Psychologica. 139:440-8.

Brass M, Bekkering H, Wohlschlager A, Prinz W. 2000. Compatibility between observed and executed finger movements: comparing symbolic, spatial, and imitative cues. Brain and Cognition, 44: 124-143.

Brass M, Zysset S, von Cramon DY. 2001. The inhibition of imitative response tendencies. NeuroImage, 14:1416-1423.

Brass M, Derrfuss J, Matthes-von Cramon G, von Cramon DY. 2003. Imitative response tendencies in patients with frontal brain lesions. Neuropsychology, 17:265-271.

Brass M, Derrfuss J, von Cramon DY. 2005. The inhibition of imitative and overlearned responses: a functional double dissociation. Neuropsychologia, 43:89-98.

Brass, M., & Heyes, C. 2005. Imitation: is cognitive neuroscience solving the correspondence problem? Trends Cogn Sci, 9:489-495.

Brass M, Ruby P, Spengler S. 2009. Inhibition of imitative behaviour and social cognition. Philos Trans R Soc Lond B Biol Sci. 364:2359-67.

Bunge SA, Hazeltine E. Scanlon MD, Rosen AC, Gabrieli JDE. 2002. Dissociable contributions of prefrontal and parietal cortices to response selection. NeuroImage. 17:1562-1571.

Button, K. S., Ioannidis, J. P., Mokrysz, C., Nosek, B. A., Flint, J., Robinson, E. S., & Munafò, M. R. (2013). Power failure: why small sample size undermines the reliability of neuroscience. Nature Reviews Neuroscience, 14(5), 365.

Byrne, R. W., & Russon, A. E. (1998). Learning by imitation: A hierarchical approach. Behavioral and brain sciences, 21(5), 667-684.

Carp, J. (2012). The secret lives of experiments: methods reporting in the fMRI literature. Neuroimage, 63(1), 289-300.

Caspers, S., Zilles, K., Laird, A. R., & Eickhoff, S. B. (2010). ALE meta-analysis of action observation and imitation in the human brain. NeuroImage, 50(3), 1148-1167.

Catmur C, Heyes C. 2011. Time course analyses confirm independence of imitative and spatial compatibility. Journal of experimental psychology. Human perception and performance. 37:409-21.

Chartrand, T. L., & Lakin, J. L. (2013). The Antecedents and Consequences of Human Behavioral Mimicry. Annual Review of Psychology, 64(1), 285-308. doi: doi:10.1146/annurev-psych-113011-143754

Cook, J. L., & Bird, G. (2012). Atypical social modulation of imitation in autism spectrum conditions. Journal of autism and developmental disorders, 42(6), 1045-1051.

Cooper, R. P., Catmur, C., & Heyes, C. 2012. Are automatic imitation and spatial compatibility mediated by different processes? Cogn Sci. 37:605–30.

Corbetta, M., Patel, G., & Shulman, G. L. (2008). The Reorienting System of the Human Brain: From Environment to Theory of Mind. Neuron, 58(3), 306-324.

Cracco E, Bardi L, Desmet C, Rigoni D, Radkova I, Deschrijver E, Genschow O, De Coster
L, Brass M. Automatic Imitation: A Meta-Analysis. (2018). Psychological Bulletin.

Crescentini C, Mengotti P, Grecucci A, Rumiati RI. 2011. The effect of observed biological
and non-biological movements on action imitation: an fMRI study. Brain
research. 1420:80-92.

Cross KA, Iacoboni M. 2013. Optimised neural coding? Control mechanisms in large
cortical networks implemented by connectivity changes. Human Brain Mapping.
34:213-25.

Cross, K. A., Torrisi, S., Losin, E. A. R., & Iacoboni, M. 2013. Controlling automatic
imitative tendencies: interactions between mirror neuron and cognitive control
systems. Neuroimage, 83:493-504.

Cross, K. A., & Iacoboni, M. 2014. To imitate or not: Avoiding imitation involves
preparatory inhibition of motor resonance. Neuroimage, 91:228-236.

Darda, K. M., Butler, E. E., & Ramsey, R. (2018). Functional Specificity and Sex
Differences in the Neural Circuits Supporting the Inhibition of Automatic
Imitation. Journal of cognitive neuroscience, 30(6), 914-933.

Decety, J., Chaminade, T., Grezes, J., & Meltzoff, A. N. (2002). A PET exploration of the
neural mechanisms involved in reciprocal imitation. Neuroimage, 15(1), 265-
272.

de Guzman M, Bird G, Banissy MJ, Catmur C. 2016. Self–other control processes in social
cognition: from imitation to empathy. Phil. Trans. R. Soc.
B. 371(1686):20150079.

Downing, P. E., & Peelen, M. V. (2011). The role of occipitotemporal body-selective
regions in person perception. Cognitive Neuroscience, 2(3-4), 186-203.

Duffy, K. A., & Chartrand, T. L. (2015). Mimicry: causes and consequences. Current
Opinion in Behavioral Sciences, 3, 112-116.

Dufour N, Redcay E, Young L, Mavros PL, Moran JM, Triantafyllou C, Gabrieli JD, Saxe R.
2013. Similar brain activation during false belief tasks in a large sample of adults
with and without autism. PloS One. 8:e75468.

Duncan J. 2010. The multiple-demand (MD) system of the primate brain: mental
programs for intelligent behaviour. Trends Cogn Sci. 14:172-9.

Ferri, F., Frassinetti, F., Mastrangelo, F., Salone, A., Ferro, F. M., and Gallese, V. (2012). Bodily self and schizophrenia: the loss of implicit self-body knowledge. Conscious. Cogn. 21, 1365–1374. doi: 10.1016/j.concog.2012.05.001

Gallagher, H. L., Happé, F., Brunswick, N., Fletcher, P. C., Frith, U., & Frith, C. D. (2000). Reading the mind in cartoons and stories: an fMRI study of 'theory of mind'in verbal and nonverbal tasks. Neuropsychologia, 38(1), 11-21.

Genschow, O., van Den Bossche, S., Cracco, E., Bardi, L., Rigoni, D., & Brass, M. 2017. Mimicry and automatic imitation are not correlated. PloS one, 12(9), e0183784.

Gowen E, Bolton E, Poliakoff E. 2016. Believe it or not: Moving non-biological stimuli believed to have human origin can be represented as human movement. Cognition. 146:431-8.

Hazeltine E, Poldrack R, Gabrieli JDE. 2007. Neural activation during response competition. J Cogn Neurosci. 12:118-129.

Heyes, C., Bird, G., Johnson, H., & Haggard, P. (2005). Experience modulates automatic imitation. Cognitive Brain Research, 22(2), 233-240.

Heyes, C. (2011). Automatic imitation. Psychological bulletin, 137(3), 463.

Hogeveen J, Obhi SS, Banissy MJ, Santiesteban I, Press C, Catmur C, Bird G. 2014. Task-dependent and distinct roles of the temporoparietal junction and inferior frontal cortex in the control of imitation. Social Cognitive and Affective Neuroscience, 10:1003-1009.

Iacoboni, M., Woods, R. P., Brass, M., Bekkering, H., Mazziotta, J. C., & Rizzolatti, G. (1999). Cortical mechanisms of human imitation. science, 286(5449), 2526-2528.

Kanwisher N. 2010. Functional specificity in the human brain: a window into the functional architecture of the mind. Proc Natl Acad Sci. 107:11163-70.

Kober, H., & Wager, T. D. (2010). Meta-analysis of neuroimaging data. Wiley Interdisciplinary Reviews: Cognitive Science, 1(2), 293-300.

Krall, S. C., Rottschy, C., Oberwelland, E., Bzdok, D., Fox, P. T., Eickhoff, S. B., ... & Konrad, K. (2015). The role of the right temporoparietal junction in attention and social interaction as revealed by ALE meta-analysis. Brain Structure and Function, 220(2), 587-604.

Lancaster, J. L., Tordesillas-Gutiérrez, D., Martinez, M., Salinas, F., Evans, A., Zilles, K., ... & Fox, P. T. (2007). Bias between MNI and Talairach coordinates analyzed using the ICBM-152 brain template. Human brain mapping, 28(11), 1194-1205.

Lee, S. M., & McCarthy, G. (2014). Functional heterogeneity and convergence in the right temporoparietal junction. Cerebral Cortex, 26(3), 1108-1116.

Lhermitte, F., Pillon, B., & Serdaru, M. (1986). Human autonomy and the frontal lobes. Part I: Imitation and utilization behavior: a neuropsychological study of 75 patients. Annals of neurology, 19(4), 326-334.

Lipsey, M. W., & Wilson, D. B. (2001). Practical meta-analysis. Sage Publications, Inc.

Marsh LE, Bird G, Catmur C. 2016. The imitation game: Effects of social cues on 'imitation' are domain-general in nature. NeuroImage. 139:368-75.

Mengotti P, Corradi-Dell'Acqua C, Rumiati RI. 2012. Imitation components in the human brain: an fMRI study. Neuroimage. 59:1622-30.

Munafò, M. R., Nosek, B. A., Bishop, D. V. M., Button, K. S., Chambers, C. D., Percie du Sert, N., . . . Ioannidis, J. P. A. (2017). A manifesto for reproducible science. Nature Human Behaviour, 1, 0021. doi: 10.1038/s41562-016-0021

Nelson, L. D., Simmons, J., & Simonsohn, U. (2018). Psychology's Renaissance. Annu Rev Psychol, 69, 511-534. doi: 10.1146/annurev-psych-122216-011836.

Nee DE, Wager TD, Jonides J. 2007. Interference resolution: Insights from a meta-analysis of neuroimaging tasks. Cogn Affect Behav Neurosci. 7:1-17.

Newman-Norlund RD, van Schie HT, van Zuijlen AM, Bekkering H. 2007. The mirror neuron system is more active during complementary compared with imitative action. Nat Neurosci. 10:817-819.

Press, C., Bird, G., Flach, R., & Heyes, C. (2005). Robotic movement elicits automatic imitation. Cognitive Brain Research, 25(3), 632-640.

Prinz, W., & Meltzoff, A. N. (2002). An introduction to the imitative mind and brain. The imitative mind: Development, evolution and brain bases, 1-15.

Ruby, P., & Decety, J. (2001). Effect of subjective perspective taking during simulation of action: a PET investigation of agency. Nature neuroscience, 4(5), 546.

Santiesteban I, Banissy MJ, Catmur C, Bird G. 2012. Enhancing Social Ability by Stimulating Right Temporoparietal Junction. Curr Biol. 22: 2274-2277.

Santiesteban, I., Banissy, M. J., Catmur, C., & Bird, G. 2015. Functional lateralization of temporoparietal junction–imitation inhibition, visual perspective-taking and theory of mind. European Journal of Neuroscience, 42(8), 2527-2533.

Schuwerk T, Schurz M, Müller F, Rupprecht R, Sommer M. 2017. The rTPJ's overarching cognitive function in networks for attention and theory of mind. Social cognitive and affective neuroscience. 12(1):157-168.

Shea, N. (2009). Imitation as an inheritance system. Philosophical Transactions of the Royal Society B: Biological Sciences, 364(1528), 2429-2443.

Spengler S, von Cramon DY, Brass M. 2009. Control of shared representations relies on key processes in mental state reasoning. Human Brain Mapping. 30:3704-3718.

Spengler S, von Cramon DY, Brass M. 2010. Resisting motor mimicry: Control of imitation involves processes central to social cognition in patients with frontal and temporo-parietal lesions. Social Neuroscience, 5: 401-416.

Spunt, R. P., & Adolphs, R. (2017). A new look at domain specificity: insights from social neuroscience. Nat Rev Neurosci, 18(9), 559-567. doi: 10.1038/nrn.2017.76

Stürmer, B., Aschersleben, G., & Prinz, W. (2000). Correspondence effects with manual gestures and postures: a study of imitation. Journal of Experimental Psychology: Human Perception and Performance, 26(6), 1746.

Thiel, C. M., Zilles, K., & Fink, G. R. (2004). Cerebral correlates of alerting, orienting and reorienting of visuospatial attention: an event-related fMRI study. Neuroimage, 21(1), 318-328.

Vallar, G. (1993). The anatomical basis of spatial hemineglect in humans. In Unilateral neglect (pp. 37-69). Psychology Press.

Vallar, G., & Perani, D. (1987). The anatomy of spatial neglect in humans. In Advances in psychology (Vol. 45, pp. 235-258). North-Holland.

van Schie HT, van Waterschoot BM, Bekkering H. 2008. Understanding action beyond imitation: reversed compatibility effects of action observation in imitation and joint action. J Exp Psychol Hum Percept Perform. 34:1493.

Wager TD, Sylvester CYC, Lacey SC, Nee DE, Franklin, M, Jonides J. 2005. Common and unique components of response inhibition revealed by fMRI. NeuroImage. 27:323-340.

Wager, T. D., Lindquist, M., & Kaplan, L. (2007). Meta-analysis of functional neuroimaging data: current and future directions. Social cognitive and affective neuroscience, 2(2), 150-158.

Wang Y, Ramsey R, Hamilton AF de C. 2011. The control of mimicry by eye contact is mediated by medial prefrontal cortex. J Neurosci. 31:12001e12010.

Whiten, A., & Van Schaik, C. P. (2007). The evolution of animal 'cultures' and social intelligence. Philosophical Transactions of the Royal Society B: Biological Sciences, 362(1480), 603-620.

Zwaan, R. A., Etz, A., Lucas, R. E., & Donnellan, M. B. (2017). Making Replication Mainstream. Behav Brain Sci, 1-50. doi: 10.1017/S0140525X17001972

Table 1. Data extracted from the studies included in the meta-analysis.

Authors	Year	Sample	Contrasts			Fixed/Random Effects Model	MNI or Talaraich coordinates
			GC	SC	IC		
Brass et al.	2001	10	x			Fixed	Talaraich
Brass et al.	2005	20	x			Fixed	Talaraich
Spengler et al.	2009	20	x			Random	Talaraich
Crescentini et al.	2011	19	x			Random	MNI
Wang et al.	2011	20	x			Random	MNI
Mengotti et al.	2012	22	x	x		Random	MNI
Cross & Iacoboni	2013	24	x	x		Random	MNI
Cross et al.	2013	20	x		x		
Klapper et al.	2014	19	x			Random	MNI
Marsh et al.	2016	24	x	x	x	Random	MNI
Darda et al. (Exp1)	2018	28	x			Random	MNI
Darda et al. (Exp2)	2018	50	x	x	x	Random	MNI
TOTAL = 11		276					

Table 2. Contrasts used in the meta-analysis for general, spatial, and imitative compatibility.

Authors	Year	General Compatibility	Spatial Compatibility	Imitative Compatibility
Brass et al.	2001	General Incompatible > General Compatible		
Brass et al.	2005	General Incompatible > General Compatible		
Spengler et al.	2009	General Incompatible > General Compatible		
Crescentini et al.	2011	General Incompatible > General Compatible		
Wang et al.	2011	General Incompatible > General Compatible*		
Mengotti et al.	2012	Non-specular > Specular^	Spatially Incompatible > Spatially Compatible	
Cross & Iacoboni	2013	General Incompatible > General Compatible	Spatially Incompatible > Spatially Compatible	
Cross et al.	2013	General Incompatible > General Compatible		General Compatibility > Spatial Compatibility
Klapper et al.	2014	General Incompatible > General Compatible*		
Marsh et al.	2016	General Incompatible > General Compatible	Spatial Incompatible > Spatially Compatible	Imitatively Incompatible > Imitatively Compatible
Darda et al. (Exp1)	2018	General Incompatible > General Compatible		
Darda et al. (Exp2)	2018	General Incompatible > General Compatible	Spatial Incompatible > Spatial Compatible	Imitative Incompatible > Imitatively Compatible
No. of studies		11	4	3
No. of contrasts		12	4	3
No. of foci		131	42	20

Table 2 shows the contrasts used in the current meta-analysis, and the number of contrasts, foci, and studies for each compatibility type (general, spatial, and imitative). *Collapsed across conditions; for Wang et al., 2011: collapsed across direct and averted gaze, for Klapper et al., 2014: collapsed across belief (motion-capture, computer animation) and form (human, non-human).
^Non-specular > Specular i.e. {(spatially incompatible and imitatively compatible) + (imitatively incompatible and spatially compatible) > general compatible)}]

Table 3. Areas consistently activated for general compatibility, spatial compatibility, and imitative compatibility.

GENERAL COMPATIBILITY					
Region	**MNI**			Maximum P	No. of Voxels	Threshold
	x	**y**	**z**			
Right supramarginal gyrus/ temporo-parietal junction	56	-34	36	0.48	3	**^
Right IPL/supramarginal gyrus	52	-30	38	0.40	5	*^
Right middle frontal gyrus	36	0	54	0.49	22	**^
Right superior frontal gyrus	26	-2	64	0.41	46	*^
Left insula lobe	-36	14	0	0.35	453	**
Left IPL	-44	-36	42	0.39	339	**
SPATIAL COMPATIBILITY						
Left IPL	-36	-40	48	0.78	8	^
Right superior frontal gyrus	24	-4	58	0.78	190	^
	24	-6	54		56	^
	24	-4	60		134	^
IMITATIVE COMPATIBILITY						
Left supramarginal gyrus/ IPL	-48	-28	34	0.73	11	^
Right supramarginal gyrus/ IPL	48	-26	44	0.73	18	*^
	52	-30	42		7	
	46	-26	46		11	

Table 3 shows areas consistently activated for general compatibility, spatial compatibility, and imitative compatibility. Maximum P stands for the maximum proportion of studies exhibiting the effect at the peak density weighted by sample size. MNI = Montreal Neurological Institute (MNI) standard stereotaxic space coordinates. The voxel size is $2 \times 2 \times 2$ mm^3.

*Clusters withstanding p<.01 cluster extent-based threshold.
**Clusters withstanding p<.001 cluster extent-based threshold.
^Clusters withstanding the height-based threshold.

Figure 1. A Theory Map of Automatic Imitation

¹ Detecting the presence of other humans (e.g., Kanwisher, 2010)
² Detection of dynamic and goal-directed behaviour (e.g., Caspers et al., 2010)
³ Domain General control systems are associated with the multiple demand network and by definition operate across multiple tasks / domains (e.g., Duncan, 2010)
⁴ Domain Specific control systems are associated with the theory of mind network and are functionally tuned towards social information processing (Brass et al., 2009)
⁵ Unintentional copying of others actions in social contexts (see Chartrand & Lakin, 2013)

Figure 1. A Theory Map of Automatic Imitation A Theory Map of Automatic Imitation.

For more details on the development of Theory Mapping, see Gray (2017);

www.theorymaps.org.

Figure 2. Consistency of brain activation from the MKDA Analyses.

Figure 2. Consistency of brain activation from the MKDA Analyses.

Brain areas that are consistently engaged for general compatibility (A), spatial compatibility (B), and imitative compatibility (C). Voxels that exceed the height-based threshold (p<.05, FDR corrected) in our analysis appear in yellow, and those that exceed the cluster extent-based threshold appear in orange (p<.001) and red (p<.01).

Figure 3. Overlay of the MKDA maps with the ToM and MD network masks.

Overlay of the density maps with the ToM and MD network masks allowed for identification of overlap between regions that were consistently activated in the MKDA and the ToM and MD networks. For all compatibility types (general, imitative and spatial), all regions that passed height or extent-based thresholding overlapped with regions in the MD network (A). Additionally, one cluster, which showed consistent activation for general compatibility, also overlapped with the right TPJ in the ToM network (B). There was no overlap with the mPFC node of the ToM network for any compatibility type.

Figure 4. Overlay of Contrast Indicator Maps with rTPJ.

Figure 4. Overlay of Contrast Indicator Maps with rTPJ.

The ToM network mask for rTPJ overlaid with the contrast indicator maps of all studies used for general (N=11; A) imitative (N=3; B) and spatial (N=4; C) compatibility. There was overlap between contrast indicator maps for general compatibility and spatial compatibility with right TPJ. By contrast, there is no overlap between contrast indicator maps for imitative compatibility and the same right TPJ mask.