Transmembrane Protein 135 (TMEM135) is a Liver X Receptor Target Gene that Mediates an

Auxiliary Peroxisome Matrix Protein Import Pathway.

Benjamin J. Renquist[1], Thushara W. Madanayake[1], Susma Ghimire[1], Caroline E. Geisler[1], Yafei

Xu[1], Randy L. Bogan[1]

[1]School of Animal and Comparative Biomedical Sciences, College of Agriculture and Life

Sciences, University of Arizona, Tucson, Arizona, USA

Running Title: TMEM135 is an Inducible Mediator of Peroxisome Function

#Address correspondence to:

Randy L. Bogan, PhD, MCR

School of Animal and Comparative Biomedical Sciences

University of Arizona

PO Box 210090

Tucson, AZ 85721-0090

Phone: 520-621-1487

Fax: 520-621-9435

Email: boganr@email.arizona.edu

Summary

The liver x receptors (LXRs) are key regulators of systemic lipid metabolism. We identified transmembrane protein 135 (TMEM135), a peroxisomal protein with an unknown function, as a novel LXR target gene. The LXRs directly induce *TMEM135* transcription in humans via an LXR response element in the 5' untranslated region, but do not increase *Tmem135* in murine cells. Functionally, knockdown of TMEM135 *in vitro* and *in vivo* results in many features typical of peroxisomal disorders such as steatosis and reduced peroxisomal β-oxidation. Mechanistically, proteomic and Western blot analyses indicated that TMEM135 mediates import of peroxisome matrix proteins necessary for β-oxidation and bile acid synthesis. These findings indicate that TMEM135 is an LXR-inducible regulator of peroxisome catabolic and anabolic processes by mediating an auxiliary matrix protein import pathway, and thus may represent a novel therapeutic target for disorders associated with peroxisome dysfunction.

Introduction

The liver x receptors (LXR) α (NR1H3) and β (NR1H2) belong to the nuclear receptor superfamily and are key regulators of cholesterol and fatty acid metabolism (1). Natural LXR ligands include oxysterols (2) that increase during cholesterol loading and activate LXR-mediated transcription. The LXRs heterodimerize with the retinoic acid receptor (RXR) and bind to LXR response elements (LXRE) within the promoters of target genes (2). The LXREs belong to the direct repeat 4 (DR4) class that consist of two hexanucleotide half-sites separated by a 4-nucleotide spacer (3,4). The protein products of known LXR target genes affect systemic lipid metabolism. Well known examples include increasing cholesterol efflux via ATP-binding cassette (ABC) sub-family A, member 1 (ABCA1) (5), and sub-family G, member 1 (ABCG1) (6); limiting cholesterol uptake via induction of myosin regulatory light chain interacting protein (MYLIP) that targets the low-density lipoprotein receptor (LDLR) for ubiquitin-mediated degradation (7); and increased lipogenesis via induction of sterol regulatory element binding protein 1c (SREBP1c, encoded by sterol regulatory element binding transcription factor 1 gene or *SREBF1*) (8,9). Also,

the *NR1H3* isoform of the LXRs is itself an LXR target gene (10,11), leading to auto-amplification

of LXR actions in certain species and tissues. Although many LXR target genes have been

discovered, additional target genes may remain unidentified.

We have previously investigated LXR actions within the steroidogenic corpus luteum of

the ovary (12-14). In an attempt to uncover novel LXR target genes, rhesus macaque luteal cells

were treated with vehicle or 1 μM T0901317 (T09, synthetic LXR agonist) for 24 hours, mRNA

were isolated and used for microarray analysis of gene expression on the Affymetrix® Rhesus

Macaque genome chip (unpublished data). Genes that were differentially expressed (>2-fold

change in T09 vs vehicle, p< 0.05 with Benjamini and Hochberg correction for false discovery

rate) were identified. Several known LXR target genes were found to be differentially expressed

including *ABCA1*, *ABCG1*, *NR1H3*, *SREBF1*, and *MYLIP*. One gene that was differentially

expressed and has not been previously determined to be an LXR target gene was transmembrane

protein 135 (*TMEM135*). Therefore, the objective of the current study is to determine if *TMEM135*

is an LXR target gene and to identify its physiologic role in lipid metabolism.

Results

<u>LXR Agonist Increases Expression of TMEM135 in Human Hepatocyte and Macrophage Cell</u>

<u>Lines</u>

Macrophage and hepatocyte cell lines were selected for initial studies on LXR regulation

of *TMEM135* transcription as these cell types play a pivotal role in LXR-mediated reverse

cholesterol transport from peripheral tissues to the liver (1). The LXR ligand T09 increased the

mRNA expression of *TMEM135* in both HepG2 (Figure 1A) and Hep3B (Figure 1B) hepatocyte

cell lines, with an approximately 2-fold maximum increase in HepG2 cells and a 3-fold increase

in Hep3B cells. In monocyte-derived macrophages (THP-1), T09 also induced *TMEM135* mRNA

expression up to 4-fold (Figure 1C). The protein synthesis inhibitor cycloheximide was used to

determine whether the effect of T09 on *TMEM135* in THP-1 cells requires synthesis of new

proteins. Cycloheximide did not significantly affect basal or T09-stimulated *TMEM135* mRNA

expression (Figure 1D), indicating that T09 induces *TMEM135* via a direct transcriptional

mechanism not involving the synthesis of intermediary proteins. As TMEM135 has previously

been reported to be localized to mitochondria-enriched fractions (15,16), the effect of T09 on

TMEM135 protein expression in mitochondria-enriched fractions from HepG2 cells was

determined. Relative concentrations of TMEM135 were significantly ($p < 0.05$) increased

approximately 1.8-fold following a 24-hour T09 treatment (Figure 1E and 1F).

A LXRE in the 5' UTR Mediates LXR Agonist Induction of TMEM135

A 4 kb region spanning from -3319 to +682 bp (relative to the transcriptional start site) of

the human *TMEM135* promoter was analyzed with MatInspector (17) for potential LXRE binding

sites. This analysis yielded three potential LXREs (Figure S1), and these were arbitrarily

designated LXRE1, LXRE2 and LXRE3 in order from most distal to most proximal to the

translation start site. Electrophoretic mobility shift assays (EMSA) were used to determine if

LXR/RXR heterodimers bind these LXREs. As shown in Figure 2A, NR1H3 and RXRA nuclear

receptors together, but not individually, caused a shift in mobility of the fluorescent LXRE1 probe

indicating that NR1H3/RXRA heterodimers bind the LXRE1 sequence in the *TMEM135* promoter.

A 200-fold molar excess of unlabeled wild type LXRE1, LXRE2, and LXRE3 oligonucleotides

eliminated the appearance of the shifted fluorescent band while the same molar excess of mutant

LXRE1, LXRE2, and LXRE3 oligonucleotides (Figure S1B) did not prevent NR1H3/RXRA binding

to the fluorescent probe; which validated the specificity of NR1H3/RXRA heterodimer binding to

all 3 LXREs in the *TMEM135* promoter (Figure 2A). Similar results were obtained for

NR1H2/RXRA heterodimers (Figure 2B), indicating that both LXR isoforms specifically bind to all

3 LXRE sites in the *TMEM135* promoter. The LXRE1 and LXRE2 sites have identical

hexanucleotide half-sites, whereas the LXRE3 site is unique (Figure S1B). To indicate if these

sequence variations may cause differences in relative binding affinity, a fluorescent LXRE3 probe

was incubated with NR1H3/RXRA heterodimers and increasing concentrations (from 0.25 to 10-

fold molar excess) of either LXRE1 or LXRE3 unlabeled competitor DNA. Oligonucleotides with

the LXRE3 sequence more effectively inhibited binding to the fluorescent probe than

oligonucleotides with the LXRE1 sequence (Figure 2C). As a complementary approach, because

the LXRE1 and LXRE3 sites were fluorescently labelled with spectrally distinct fluorophores, an

EMSA was performed where a mixture (1 nM each) of fluorescent LXRE1 and LXRE3 probes

were incubated with increasing concentrations of NR1H3/RXRA heterodimers. This approach

resulted in a higher percentage of LXRE3 than LXRE1 bound across a range of NR1H3/RXRA

heterodimer concentrations (Figure 2D). Collectively, this indicates that LXRE3 has a higher

binding affinity for NR1H3/RXR heterodimers than LXRE1.

Luciferase reporter assays were used to study transcription initiation from the *TMEM135*

promoter. Transfection of HepG2 cells with wild type *TMEM135* promoter/pGL 4.17 resulted in a

large increase in luciferase expression that was responsive to T09 as compared to the empty pGL

4.17 vector (Figure 2E). Furthermore, luciferase activity was further amplified by co-transfection

of plasmids that constitutively express NR1H3 and RXRA (Figure 2E). Next, we determined the

requirement of each individual LXRE to LXR agonist-induced transcription from the *TMEM135*

promoter. Promoters containing all combinations of wild type and mutant LXREs (Figure S1B)

were generated, and T09-induced luciferase activity was determined. LXR-agonist induced

luciferase activity was significantly ($p < 0.05$) higher in cells transfected with the wild type

TMEM135 promoter as compared to the *TMEM135* promoter with all 3 LXREs mutated (Figure

2F). Furthermore, constructs containing the wild type LXRE3 site and mutated LXRE1 and/or

LXRE2 sites were not significantly different from the wild type promoter, while all constructs that

contained a mutated LXRE3 were not significantly different from the promoter that had all three

LXREs mutated (Figure 2F). This indicates that even though all 3 LXREs bind LXR/RXR

heterodimers, the LXRE3 site alone mediates LXR agonist-induced transcription from the

TMEM135 promoter.

We next determined whether *Tmem135* mRNA expression was induced by T09 in mice.

The murine LXRE3 site showed relatively little homology with the human sequence as a total of

3 nucleotides within both hexanucleotide half-sites were unique in mice (Figure S2A). In mouse

hepatocyte (BNL 1NG A.2) and macrophage (RAW 264.7) cell lines, T09 did not significantly

induce *Tmem135* mRNA expression (Figure S2B) whereas it did increase the known LXR target

gene *Abca1* (Figure S2C). These data indicate that *Tmem135* is not an LXR target gene in mice.

<u>*TMEM135* is a Direct Target Gene of the LXRs</u>

To directly determine whether the LXRs are needed for T09-induced *TMEM135*

transcription, siRNA-mediated knockdown of the LXRs was performed in HepG2 cells. The

NR1H2 siRNA caused an approximately 75% decrease ($p < 0.05$) in *NR1H2* mRNA expression

compared to the control and *NR1H3* siRNA groups (Figure 3A). The *NR1H3* siRNA resulted in an

approximately 70-80% decrease ($p < 0.05$) in *NR1H3* mRNA expression compared to the control

siRNA depending on the presence or absence of T09 (Figure 3B). The *NR1H2* siRNA by itself

also caused a significant reduction in T09-stimulated *NR1H3* mRNA expression and tended to

slightly improve *NR1H3* knockdown when co-transfected with the *NR1H3* siRNA (Figure 3B).

Decreased *NR1H3* mRNA expression is expected to occur following NR1H2 knockdown because

NR1H3 itself is an LXR target gene (6,11). Consistent with *NR1H3* being an LXR target gene,

T09 induced a significant increase in *NR1H3* in the control siRNA group, while T09 was not as

effective at inducing *NR1H3* in the *NR1H2* siRNA group (Figure 3B).

The known LXR target gene *ABCA1* was significantly increased by T09 in the control

siRNA group, and the effect of T09 was completely abolished by *NR1H2/NR1H3* siRNA co-

transfection (Figure 3C). Also, *NR1H3* knockdown alone significantly inhibited the T09-induced

increase in *ABCA1* (Figure 3C). Results for *TMEM135* were very similar to *ABCA1* with T09

causing a significant increase in the control siRNA group, and the effect of T09 was completely

blocked by *NR1H2/NR1H3* siRNA co-transfection (Figure 3D). Furthermore, *NR1H3* knockdown

itself significantly inhibited the T09-induced increase in *TMEM135* as compared to the control

siRNA (Figure 3D).

The LXRE3 site of the human *TMEM135* promoter lies downstream of the putative

transcriptional start site, but before the translation initiation site (Figure S1). To determine whether

LXR-mediated transcription of *TMEM135* results in an mRNA with a truncated 5' UTR, primers

and a probe were designed to amplify and detect a 76 base pair region encompassing the LXRE3

site in the 5' UTR. There was a significant ($p < 0.05$) reduction in mRNA expression of full-length

5' UTR transcripts caused by T09 in the control siRNA group, while this effect was blocked by

NR1H2/NR1H3 siRNA co-transfection (Figure 3E). Furthermore, *NR1H2/NR1H3* co-knockdown

resulted in a significant ($p < 0.05$) increase in basal mRNA expression of full-length 5' UTR

transcripts compared to all other groups (Figure 3E). ChIP was used to determine if the reduction

in full-length 5' UTR transcripts is associated with increased NR1H2 and/or NR1H3 binding to the

LXRE3 site on the *TMEM135* gene. Monoclonal antibodies specific for each LXR isoform were

used for ChIP, and QPCR of the LXRE3 region on chromatin purified via ChIP indicated that T09

treatment caused a significant increase in binding of NR1H3 to LXRE3 in the *TMEM135* gene

(Figure 3F). This indicates that LXR binding to LXRE3 increases mRNA expression of *TMEM135*

via a transcript with a truncated 5' UTR relative to the putative mRNA sequence.

<u>TMEM135 Mediates Fatty Acid Metabolism and Proliferation in HepG2 Cells</u>

To begin unraveling the biologic function of TMEM135, a series of knockdown experiments

were performed. Transfection of HepG2 cells with siRNA against *TMEM135* decreased its mRNA

expression by 70-90%. As TMEM135 has previously been implicated in fat metabolism (15), and

the LXRs are known to induce lipogenesis (1), we first determined its effect on triglycerides and

mRNA expression of fatty acid oxidation and lipogenesis-associated genes. Knockdown of

TMEM135 significantly increased basal triglyceride accumulation in HepG2 cells (Figure 4A).

Furthermore, T09 significantly increased intracellular triglyceride concentrations, while TMEM135

knockdown further increased triglyceride accumulation in the presence of T09 (Figure 4A). An

increase in triglyceride accumulation could result from a decrease in fatty acid oxidation and/or

an increase in lipogenesis. There were no significant effects of TMEM135 knockdown on mRNA

expression of the key regulator of fatty acid β-oxidation, peroxisome proliferator activated receptor

alpha (*PPARA*), while it tended to increase the PPARA target gene carnitine palmitoyltransferase

1A (*CPT1A*) (18) (Figure 4B). It is known that the LXRs induce lipogenesis via induction of

SREBP1c (*SREBF1* gene) (8,9). As expected, T09 caused a significant increase in *SREBF1* and

the SREBP1c/LXR target gene fatty acid synthase (*FASN*) (Figure 4B), consistent with the T09-

induced increase in triglyceride accumulation (Figure 4A). However, TMEM135 knockdown

significantly suppressed both basal and T09-induced *SREBF1* mRNA expression (Figure 4B).

This indicates that the increase in triglyceride accumulation in HepG2 cells due to TMEM135

knockdown is likely not due to enhanced lipogenesis. Thus, an inhibition of fatty acid oxidation

seems a more likely explanation for the increase in triglyceride accumulation. Further supporting

the possibility that fatty acid oxidation is impaired, TMEM135 knockdown reduced ATP

concentrations when HepG2 cells were incubated in glucose-free medium (Figure 4C).

During these experiments it appeared that TMEM135 knockdown also inhibited replication

of HepG2 cells. Because enhanced fatty acid β-oxidation is a hallmark of hepatocellular

carcinoma (HCC) (19), HepG2 cells are an HCC-derived cell line, and our previous experiments

indicated a key role for TMEM135 in fatty acid metabolism; we determined whether TMEM135

regulated proliferation of HepG2 cells. Knockdown of *TMEM135* significantly ($p < 0.05$) reduced

viable HepG2 cell numbers at 48 and 72 hours post-transfection as compared to cells transfected

with the control siRNA (Figure 4D). Cell cycle analysis indicated that *TMEM135* knockdown

significantly ($p < 0.05$) increased the percentage of HepG2 cells in the G0/G1 stage, with a

corresponding significant reduction in the percentage of cells in the S phase (Figure 4E). The

increase in G0/G1 arrest was associated with alterations in mRNA expression of tumor

suppressor and cell cycle genes. There were significant increases in cyclin dependent kinase

inhibitor 2A (*CDKN2A*) and tumor protein p53 (*TP53*) (Figure 4F), which restrict passage through

the G1/S checkpoint (20). There were also significant increases in cyclin dependent kinase 2

(*CDK2*) and cyclin E1 (*CCNE1*) (Figure 4F), which are key mediators of passage through the

G1/S checkpoint, but no change in cyclin A2 (*CCNA2*) (Figure 4F) which is increased in the S phase (21). Collectively, these data indicate that TMEM135 knockdown reduces HepG2 proliferation by restricting passage through the G1/S checkpoint.

Liver-Selective TMEM135 Knockdown Reduces Peroxisomal β-Oxidation

To determine the physiologic function of TMEM135, a siRNA knockdown experiment was performed in male C57BL/6 mice. Mice received either a non-targeting control or *Tmem135* siRNA and were sacrificed 4 days later in either the *ad libitum* fed state or after a 12 hour fast. Because we determined that *Tmem135* is not an LXR target gene in mice (Figure S2B), we used fasting to induce hepatic fat accumulation. Tail-vein delivery of the *Tmem135* siRNA caused an approximately 60% mRNA knockdown in the liver while no knockdown was observed in other tissues including skeletal muscle, adipose, and heart (Figure 5A). Knockdown of TMEM135 in the liver was further confirmed by Western blot and proteomic analysis of mitochondria/peroxisome-enriched fractions (Figure 5B). Fed mice injected with the *Tmem135* siRNA gained significantly less weight during the 4-day siRNA treatment, while the loss in weight due to fasting was similar for both siRNAs (Figure 5C). There was no significant effect of the *Tmem135* siRNA on basal or fasting-induced hepatic triglycerides (Figure S3A), while there was a tendency for increased hepatic non-esterified fatty acid (NEFA) concentrations in TMEM135 knockdown mice in the fed state (Figure S3B). There were no significant effects of TMEM135 knockdown on liver ATP or glycogen concentrations (Figure S3C-D). Also, there were no significant effects of TMEM135 on serum lipids (total cholesterol, HDL cholesterol, triglycerides), although there was a trend for reduced serum triglycerides in fed mice receiving the control siRNA that was not observed in TMEM135 knockdown mice (Figure S4A). There were no significant effects of TMEM135 knockdown on serum NEFA, glucose, or insulin concentrations (Figure S4B-D). The mRNA expression of key genes involved in fatty acid β-oxidation were determined in the liver. As expected, fasting significantly induced mRNA expression of *Ppara*, *Cpt1a*, acyl-CoA dehydrogenase medium chain (*Acadm*), uncoupling protein 2 (*Ucp2*), and sirtuin 3 (*Sirt3*) (Figure

5D). Similarly, knockdown of TMEM135 significantly increased *Acadm* and *Sirt3* in fed animals,

while *Ucp2* increased with fasting in control but not TMEM135 knockdown animals (Figure 5D).

As expected with increased β-oxidative flux, fasting significantly increased NADH in animals

receiving the control siRNA, which was prevented by TMEM135 knockdown (Figure 5E) indicating

an impairment in β-oxidation in these mice. Reduced hepatic NADH during fasting could be due

to reduced β-oxidation in peroxisomes and/or mitochondria. However, there was a reciprocal

increase in hepatic ketone concentrations in fasted mice with TMEM135 knockdown (Figure 5F).

Because ketogenesis occurs exclusively in mitochondria and is dependent on fatty acid oxidation

(22), this indicates that mitochondrial fatty acid β-oxidation was not impaired. Thus, knockdown

mice have impaired fatty acid β-oxidation in peroxisomes but not mitochondria.

TMEM135 Mediates Import of Peroxisome Matrix Proteins Involved in Fatty Acid β-Oxidation and

Bile Acid Synthesis

TMEM135 is a peroxisomal protein with high homology to the Tim17 family that mediate

translocation of proteins across mitochondrial membranes (23). In addition to peroxisomes,

TMEM135 has been reported to be localized to mitochondria (15,16). Therefore, we hypothesized

that TMEM135 mediates protein import into peroxisomes and/or mitochondria. A

mitochondria/peroxisome-enriched fraction was prepared from frozen mouse livers. This fraction

was validated to contain both mitochondria and peroxisomes as indicated by presence of the

mitochondria marker cytochrome c oxidase subunit 4I1 (COX4I1) and the peroxisome marker

ATP binding cassette subfamily D member 3 (ABCD3, also known as PMP70), while COX4I1 and

ABCD3 were undetectable in the cytosolic supernatant (Figure 6A). Furthermore, the cytosolic

protein tubulin beta class I (TUBB) segregated to the cytosolic fraction and was not detected in

the mitochondria/peroxisome-enriched fraction. Proteomic analysis of a subset of fed mice was

used to provide an unbiased estimate of differential protein abundance in the

mitochondria/peroxisome-enriched fractions. A total of 23 proteins were significantly less

abundant (Fisher's Exact Test, p< 0.05) in the mitochondria/peroxisome-enriched fraction from

TMEM135 knockdown livers (Figure 6B). Of particular interest was the finding that these 23

proteins included nearly all of the matrix proteins known to be necessary for bile acid synthesis

and peroxisomal β-oxidation of very-long chain fatty acids, branched chain fatty acids, and

dicarboxylic acids (24,25) (Figure 6 C-D). Four proteins were selected for further analysis via

Western blot: acetyl-CoA acyltransferase 1 (ACAA1), acyl-CoA oxidase 1 (ACOX1), sterol carrier

protein 2 (SCP2), and catalase (CAT). All these proteins were enriched in the

mitochondria/peroxisome fraction, but except for SCP2 (which may be due to low detection

sensitivity) each protein was also clearly detectable in the cytosolic fraction (Figure 6E). This

contrasts with the peroxisome membrane protein ABCD3 which appeared to be exclusively

localized to the mitochondria/peroxisome fraction (Figure 6A). Collectively, detection of these

proteins in the cytoplasm indicates that their import into peroxisomes may be rate-limiting.

We next used Western blot to quantitatively determine import of ACAA1, ACOX1, SCP2,

and CAT into peroxisomes. Whole cell lysates (using TUBB as loading control) and

mitochondria/peroxisome-enriched fractions (using COX4I1 and ABCD3 as loading controls for

mitochondria and peroxisome content, respectively) were analyzed (Figure 7A). The signal for

each protein was normalized to the respective loading control, and the ratio of peroxisome to

whole cell lysate was used to determine import. There were no significant differences in

peroxisome content of the mitochondria/peroxisome-enriched fraction with TMEM135 knockdown

as indicated by the ABCD3:COX4I1 ratio, indicating that differences detected by proteomic

analysis are not simply due to effects on peroxisome proliferation (Figure 7B). When using ABCD3

to correct for variations in peroxisome content of the mitochondria/peroxisome-enriched fraction,

there were no significant differences between siRNAs for peroxisome import of ACOX1 and CAT

in either fed or fasted animals (Figure 7C). However, relative ACAA1 import was significantly

reduced in TMEM135 knockdown mice when fasted (Figure 7C). Furthermore, SCP2 import was

significantly lower in TMEM135 knockdown mice in both the fed and fasted state (Figure 7C).

Thus, TMEM135 mediates the import of ACAA1 and SCP2 into peroxisomes.

Discussion

Peroxisomes have key catabolic functions including α and β-oxidation of fatty acids, and anabolic activities including the synthesis of bile acids and ether phospholipids (24,25). There are 2 main classes of peroxisome disorders: 1) peroxisome biogenesis disorders, and 2) single peroxisomal enzyme deficiencies (24,25). Outcomes from these disorders range from infant death to mild degenerative neurosensory problems (24,25). Hepatic pathologies are one of the principal features associated with peroxisome disorders, which are often caused by a failure to import matrix enzymes into peroxisomes (24). In the current study we determined that TMEM135, a protein predicted to be associated with peroxisomes based on homology but with a previously unknown function, is an LXR-inducible protein that mediates the import of proteins into peroxisomes.

Many effects of hepatocyte-specific TMEM135 knockdown in the current study are consistent with typical features of peroxisomal disorders. Knockdown of TMEM135 causes an accumulation of fatty acids both *in vitro* and *in vivo*. In HepG2 cells, TMEM135 knockdown increased triglyceride concentrations despite a simultaneous significant reduction in mRNA expression of *SREBF1*, the key regulator of lipogenesis (Figure 4A-B). This resembles liver-specific peroxisome deficiency in mice (peroxisomal biogenesis factor 5 or *Pex5* conditional knockout), which results in hepatic steatosis despite significantly reduced mRNA expression of *Srebf1* (26). *In vivo* there was a tendency for TMEM135 knockdown to increase hepatic NEFA concentrations in fed mice (Figure S3B), which was coupled with a general increase in mRNA expression of *PPARA* and fasting-induced genes in the fed state such as *Acadm* (18) and *Sirt3* (27) (Figure 5D). This is consistent with several mouse models of peroxisome deficiency or peroxisomal β-oxidation defects, which are characterized by steatosis and consequently induction of PPARA target genes due to an increase in endogenous PPARA agonists (24). When β-oxidation (both peroxisomal and mitochondrial) was stimulated by fasting, TMEM135 knockdown mice had significantly lower NADH concentrations (Figure 5E), indicating an impairment in β-

oxidation. Meanwhile, hepatic ketones were significantly increased in knockdown mice when

fasted (Figure 5F). Because ketogenesis occurs exclusively in mitochondria, the increase in

ketones may have resulted from increased availability of fatty acids to the mitochondria due to a

decrease in competition with peroxisomes (28), or due to the increased NAD+/NADH ratio (22).

Collectively, this indicates that an impairment in fatty acid β-oxidation occurs at the level of

peroxisomes and not mitochondria. One final similarity between TMEM135 knockdown and

peroxisomal disorders is that the significant reduction in body weight gain of knockdown mice in

the fed state (Figure 5C) is consistent with liver-specific peroxisome deficiency in mice that results

in a stunting of body growth (26) and severe weight loss when pair-fed with control mice (29).

The TMEM135-mediated import of proteins into peroxisomes is a novel pathway. The

classical mechanism involves peroxisomal targeting signals (PTS), which are short amino acid

sequences at the N or C-terminus that interact with various members of the peroxin family of

proteins to facilitate import into the peroxisome matrix (30). Among the proteins whose import was

quantified by Western blot in the current study ACOX1 and SCP2 contain a PTS1 sequence on

their C-terminus that is recognized by PEX5, and ACAA1 contains a PTS2 sequence at its N-

terminus that interacts with PEX7 (peroxisomal biogenesis factor 7) (30,31). Therefore, TMEM135

is likely not obligatory for their import, but rather serves an auxiliary role. It is not known whether

TMEM135-mediated protein import occurs via a mechanism that involves, or occurs

independently of, peroxins. Because TMEM135 is a transmembrane protein with TIM17 homology

(23), a peroxin-independent mechanism seems most likely. Regardless, TMEM135-mediated

protein import may be necessary under conditions of high peroxisome activity when peroxin-

dependent import becomes saturated.

An auxiliary role of TMEM135 in peroxisome protein import makes it a potential therapeutic

target. One potential pharmacologic therapy comes from our findings in the current study that

TMEM135 is an LXR target gene in humans. The function of TMEM135 fits well with the known

roles of the LXRs in cholesterol metabolism. The LXRs are master regulators of reverse

cholesterol transport (RCT), a process whereby excess cholesterol is removed from peripheral

tissues and transported to the liver for elimination in bile acids (1). There has been much interest

in the development of LXR agonists to treat human diseases including atherosclerosis,

Alzheimer's disease, and other metabolic disorders (1). Because RCT results in cholesterol being

converted to bile acids in the liver and the final steps in bile acid synthesis occur in peroxisomes

(24), an LXR-induced increase in TMEM135 would help stimulate peroxisomal synthesis of bile

acids during periods of cholesterol loading. The LXRs are also known to increase hepatic

lipogenesis via their induction of SREBP1c (8,9), which is a master regulator of genes necessary

for fatty acid synthesis. In fact, increased hepatic lipogenesis has been the primary obstacle in

the development of LXR agonists as therapeutic agents (1). Elevated TMEM135 expression may

increase peroxisomal β-oxidation of fatty acids, and thus limit fat accumulation when lipogenesis

is elevated due to LXR activation. It should be noted that our data indicate that *TMEM135* is an

LXR target gene in humans, but in mice *Tmem135* is either not induced by the LXRs or is a low-

affinity target gene (Figure S2). Thus, LXR agonist-induced hepatic fat accumulation may be less

pronounced in humans than in mice due to species differences in LXR regulation of *TMEM135*

transcription.

Elevated TMEM135 activity may protect against cholestasis and fatty liver, but may also

contribute to hepatic pathologies. For example, metabolomics studies have reported that elevated

serum bile acids and urinary bile salts represent the core metabolic phenotype in the transition

from healthy liver to fatty liver and HCC (19). Additionally, elevated fatty acid β-oxidation and

reduced mitochondrial respiration are part of the metabolic remodeling that occurs in liver disease

(19). These metabolic changes that occur during HCC pathogenesis are consistent with excessive

peroxisome activity as peroxisomes play a key role in bile acid synthesis, they contribute to fatty

acid β-oxidation, and can also indirectly alter mitochondrial respiration (24). In the current study

TMEM135 knockdown reduced proliferation of the HepG2 HCC cell line by increasing G0/G1

arrest. There is additional evidence in the literature, albeit limited, of an oncogenic role for

TMEM135. It was among genes identified by transcriptomic analysis as being significantly

reduced in expression when proliferation of the HCC cell lines HepG2 and Huh7 were inhibited

by knockdown of the COP9 signalosome subunit 5 (32). Also, a genomic rearrangement that

results in a TMEM135-containing fusion transcript has been linked with aggressive prostate

cancer (33), and Cas9-mediated insertion of a suicide gene into cells expressing this same fusion

transcript reduces tumor burden and mortality in mouse xenografts (34). Thus, TMEM135 may

represent a novel target in the treatment of certain cancers, as well as in metabolic diseases

associated with fatty liver.

There have only been two previous studies that address the biological function of

TMEM135 (15,16). The first indication of its function came from a mouse model of mitochondrial

acyl-CoA dehydrogenase very long chain (ACADVL) deficiency (15). Under conditions of

increased demand for β-oxidation such as during fasting and cold stress, ACADVL-deficient mice

have reduced survival due to cardiac dysfunction, which resembles clinical features of human

ACADVL deficiency (15). It was discovered that TMEM135 was elevated more than four-fold in

the hearts of ACADVL-deficient mice that survived birth (15). Furthermore, cardiac TMEM135

expression was increased by both fasting and cold stresses in wild-type mice (15). The ACADVL

enzyme catalyzes the first step in mitochondrial fatty acid β-oxidation using fatty acids ranging in

length from 14 up to 24 carbons as substrates (35). It is generally believed that mitochondria

preferentially oxidize short and medium chain fatty acids (<C12), both mitochondria and

peroxisomes oxidize long chain fatty acids (C14-C20), while peroxisomes exclusively oxidize

very-long chain fatty acids (>C20) (28). Peroxisomal β-oxidation produces acetyl-CoA and

medium chain fatty acids that can be transported to the mitochondria for further oxidation and

ATP generation (28,36). Because ACADVL deficiency restricts the mitochondria from directly

oxidizing long and very-long chain fatty acids, an upregulation of TMEM135 would promote

survival by increasing peroxisomal β-oxidation capacity and the processing of long and very-long

chain fatty acids to medium chain fatty acids that are subsequently oxidized in the mitochondria.

Furthermore, the first step in peroxisomal β-oxidation generates hydrogen peroxide that is

subsequently metabolized by CAT into oxygen and water with the release of heat, which is

another mechanism whereby TMEM135 could aid cold stress. The second mechanistic study into

TMEM135 function reported that a mutation in *Tmem135* was responsible for accelerated aging

of the retina in a mouse model of age-related macular degeneration (16). In this study alterations

in mitochondrial function associated with TMEM135 were found (16). As will be subsequently

discussed, it is not clear whether TMEM135 acts directly within mitochondria, or if its effects on

mitochondria are indirect via altered peroxisome function. Nevertheless, it is becoming

increasingly apparent that peroxisomal function declines with aging (28), which is consistent with

a *Tmem135* mutation causing accelerated aging in the retina (16).

There is some controversy regarding the subcellular localization of TMEM135. Neither the

murine or human TMEM135 proteins contain a PEX19 (peroxisomal biogenesis factor 19)

sequence that mediates insertion of some peroxisome membrane proteins (30) as determined by

PeroxisomeDB 2.0 (37). There are also no PTS1 or PTS2 motifs in TMEM135 (37). Thus, there

is no indication that TMEM135 incorporation into peroxisome membranes involves interaction with

peroxins. One potential mechanism of TMEM135 incorporation into peroxisome membranes

involves its translation in the endoplasmic reticulum (ER), and incorporation into peroxisomes via

an ER-dependent mechanism (30). It has previously been reported that TMEM135 specifically

localizes to peroxisomes but not mitochondria as determined using a GFP fusion protein (38),

while other studies have reported it to be localized to mitochondria (15,16). However, in studies

reporting TMEM135 localization to mitochondria, non-mitochondrial localization was also

observed in "dot-like" (15) or a "vesicular expression pattern in the cytoplasm" (16), which could

be descriptions of peroxisomes. In Lee *et. al.* (16), TMEM135 was found in punctate structures

on the surface of mitochondria. It is possible that these punctate structures are peroxisomes

associated with mitochondria. Indeed, it has become clear that peroxisomes and mitochondria

can physically associate with each other via the formation of tethering complexes, which may help

in functions such as metabolite transfer between organelles (39). Also, peroxisomes and

mitochondria share proteins involved in their division (39), and peroxisome-specific induction of

reactive oxygen species production causes increased mitochondrial fragmentation (40). There is

some evidence that TMEM135 may regulate mitochondrial division as a *Tmem135* mutation was

associated with over-fused mitochondrial networks whereas TMEM135 overexpression caused

an increase in mitochondrial fission (16). Overall, it seems plausible that TMEM135 is not

localized in mitochondria but rather in peroxisomes that can associate with mitochondria, and its

effects on mitochondria may be indirectly mediated via altered peroxisome activity.

The critical role of peroxisomes in human health is illustrated by the spectrum of

peroxisome disorders; but in addition to these inborn errors of metabolism it is becoming

increasingly apparent that peroxisomes play key roles in other metabolic and age-related

diseases including cancer, neurodegenerative disorders, and diabetes (28). This may be due to

direct catabolic and anabolic actions of peroxisomes, as well as interactions between

peroxisomes and the ER or mitochondria (28,39). Our data indicate that TMEM135 regulates

peroxisome activity by mediating an auxiliary, and inducible, matrix protein import pathway.

Furthermore, *TMEM135* is an LXR target gene in humans, and is likely to be regulated by other

metabolic signaling pathways. These findings implicate TMEM135 as a potential therapeutic

target in the treatment of diseases associated with peroxisomal dysfunction.

Acknowledgements

Mass spectrometry and proteomics data were acquired by the Arizona Proteomics

Consortium supported by NIEHS grant ES06694 to the SWEHSC, NIH/NCI grant CA023074 to

the UA Cancer Center, and by the BIO5 Institute of the University of Arizona. The authors would

like to thank Dr. Jon Hennebold at the Oregon National Primate Research Center of Oregon

Health & Science University for assistance with the unpublished microarray experiment in

macaque luteal cells that provided preliminary data for the current study.

Funding Statement: This research was supported by the Eunice Kennedy Shriver National

Institute of Child Health & Human Development, National Institutes of Health, Award Number R00

HD067678 to R.L.B. The content is solely the responsibility of the authors and does not

necessarily represent the official views of the National Institutes of Health.

References

1. Hong, C., and Tontonoz, P. (2014) Liver X receptors in lipid metabolism: opportunities for
drug discovery. *Nat Rev Drug Discov* **13**, 433-444
2. Peet, D. J., Janowski, B. A., and Mangelsdorf, D. J. (1998) The LXRs: a new class of
oxysterol receptors. *Curr Opin Genet Dev* **8**, 571-575
3. Teboul, M., Enmark, E., Li, Q., Wikstrom, A. C., Pelto-Huikko, M., and Gustafsson, J. A.
(1995) OR-1, a member of the nuclear receptor superfamily that interacts with the 9-cis-
retinoic acid receptor. *Proc Natl Acad Sci U S A* **92**, 2096-2100
4. Willy, P. J., Umesono, K., Ong, E. S., Evans, R. M., Heyman, R. A., and Mangelsdorf, D.
J. (1995) LXR, a nuclear receptor that defines a distinct retinoid response pathway. *Genes
Dev* **9**, 1033-1045
5. Costet, P., Luo, Y., Wang, N., and Tall, A. R. (2000) Sterol-dependent transactivation of
the ABC1 promoter by the liver X receptor/retinoid X receptor. *J Biol Chem* **275**, 28240-
28245
6. Kennedy, M. A., Venkateswaran, A., Tarr, P. T., Xenarios, I., Kudoh, J., Shimizu, N., and
Edwards, P. A. (2001) Characterization of the human ABCG1 gene: liver X receptor
activates an internal promoter that produces a novel transcript encoding an alternative
form of the protein. *J Biol Chem* **276**, 39438-39447
7. Zelcer, N., Hong, C., Boyadjian, R., and Tontonoz, P. (2009) LXR regulates cholesterol
uptake through Idol-dependent ubiquitination of the LDL receptor. *Science* **325**, 100-104
8. Repa, J. J., Liang, G., Ou, J., Bashmakov, Y., Lobaccaro, J. M., Shimomura, I., Shan, B.,
Brown, M. S., Goldstein, J. L., and Mangelsdorf, D. J. (2000) Regulation of mouse sterol
regulatory element-binding protein-1c gene (SREBP-1c) by oxysterol receptors, LXRalpha
and LXRbeta. *Genes Dev* **14**, 2819-2830
9. Yoshikawa, T., Shimano, H., Amemiya-Kudo, M., Yahagi, N., Hasty, A. H., Matsuzaka, T.,
Okazaki, H., Tamura, Y., Iizuka, Y., Ohashi, K., Osuga, J., Harada, K., Gotoda, T., Kimura,
S., Ishibashi, S., and Yamada, N. (2001) Identification of liver X receptor-retinoid X
receptor as an activator of the sterol regulatory element-binding protein 1c gene promoter.
Mol Cell Biol **21**, 2991-3000
10. Laffitte, B. A., Joseph, S. B., Walczak, R., Pei, L., Wilpitz, D. C., Collins, J. L., and
Tontonoz, P. (2001) Autoregulation of the human liver X receptor alpha promoter. *Mol Cell
Biol* **21**, 7558-7568
11. Whitney, K. D., Watson, M. A., Goodwin, B., Galardi, C. M., Maglich, J. M., Wilson, J. G.,
Willson, T. M., Collins, J. L., and Kliewer, S. A. (2001) Liver X receptor (LXR) regulation
of the LXRalpha gene in human macrophages. *J Biol Chem* **276**, 43509-43515
12. Bogan, R. L., Debarber, A. E., and Hennebold, J. D. (2012) Liver x receptor modulation of
gene expression leading to proluteolytic effects in primate luteal cells. *Biol Reprod* **86**, 89
13. Bogan, R. L., and Hennebold, J. D. (2010) The reverse cholesterol transport system as a
potential mediator of luteolysis in the primate corpus luteum. *Reproduction* **139**, 163-176

14.   Seto, N. L., and Bogan, R. L. (2015) Decreased cholesterol uptake and increased liver x
484   receptor-mediated cholesterol efflux pathways during prostaglandin F2 alpha-induced and
485   spontaneous luteolysis in sheep. *Biol Reprod* **92**, 128

486   15.   Exil, V. J., Silva Avila, D., Benedetto, A., Exil, E. A., Adams, M. R., Au, C., and Aschner,
487   M. (2010) Stressed-induced TMEM135 protein is part of a conserved genetic network
488   involved in fat storage and longevity regulation in Caenorhabditis elegans. *PLoS One* **5**,
489   e14228

490   16.   Lee, W. H., Higuchi, H., Ikeda, S., Macke, E. L., Takimoto, T., Pattnaik, B. R., Liu, C., Chu,
491   L. F., Siepka, S. M., Krentz, K. J., Rubinstein, C. D., Kalejta, R. F., Thomson, J. A., Mullins,
492   R. F., Takahashi, J. S., Pinto, L. H., and Ikeda, A. (2016) Mouse Tmem135 mutation
493   reveals a mechanism involving mitochondrial dynamics that leads to age-dependent
494   retinal pathologies. *Elife* **5**

495   17.   Cartharius, K., Frech, K., Grote, K., Klocke, B., Haltmeier, M., Klingenhoff, A., Frisch, M.,
496   Bayerlein, M., and Werner, T. (2005) MatInspector and beyond: promoter analysis based
497   on transcription factor binding sites. *Bioinformatics* **21**, 2933-2942

498   18.   Kersten, S., and Stienstra, R. (2017) The role and regulation of the peroxisome proliferator
499   activated receptor alpha in human liver. *Biochimie* **136**, 75-84

500   19.   Beyoglu, D., and Idle, J. R. (2013) The metabolomic window into hepatobiliary disease. *J*
501   *Hepatol* **59**, 842-858

502   20.   LaPak, K. M., and Burd, C. E. (2014) The molecular balancing act of p16(INK4a) in cancer
503   and aging. *Mol Cancer Res* **12**, 167-183

504   21.   Harper, J. W., and Adams, P. D. (2001) Cyclin-dependent kinases. *Chem Rev* **101**, 2511-
505   2526

506   22.   Newman, J. C., and Verdin, E. (2017) beta-Hydroxybutyrate: A Signaling Metabolite. *Annu*
507   *Rev Nutr* **37**, 51-76

508   23.   Zarsky, V., and Dolezal, P. (2016) Evolution of the Tim17 protein family. *Biol Direct* **11**, 54

509   24.   Baes, M., and Van Veldhoven, P. P. (2016) Hepatic dysfunction in peroxisomal disorders.
510   *Biochim Biophys Acta* **1863**, 956-970

511   25.   Waterham, H. R., Ferdinandusse, S., and Wanders, R. J. (2016) Human disorders of
512   peroxisome metabolism and biogenesis. *Biochim Biophys Acta* **1863**, 922-933

513   26.   Peeters, A., Swinnen, J. V., Van Veldhoven, P. P., and Baes, M. (2011) Hepatosteatosis
514   in peroxisome deficient liver despite increased beta-oxidation capacity and impaired
515   lipogenesis. *Biochimie* **93**, 1828-1838

516   27.   Hirschey, M. D., Shimazu, T., Goetzman, E., Jing, E., Schwer, B., Lombard, D. B., Grueter,
517   C. A., Harris, C., Biddinger, S., Ilkayeva, O. R., Stevens, R. D., Li, Y., Saha, A. K.,
518   Ruderman, N. B., Bain, J. R., Newgard, C. B., Farese, R. V., Jr., Alt, F. W., Kahn, C. R.,
519   and Verdin, E. (2010) SIRT3 regulates mitochondrial fatty-acid oxidation by reversible
520   enzyme deacetylation. *Nature* **464**, 121-125

521   28.   Cipolla, C. M., and Lodhi, I. J. (2017) Peroxisomal Dysfunction in Age-Related Diseases.
522   *Trends Endocrinol Metab* **28**, 297-308

523   29.   Peeters, A., Fraisl, P., van den Berg, S., Ver Loren van Themaat, E., Van Kampen, A.,
524   Rider, M. H., Takemori, H., van Dijk, K. W., Van Veldhoven, P. P., Carmeliet, P., and Baes,
525   M. (2011) Carbohydrate metabolism is perturbed in peroxisome-deficient hepatocytes due
526   to mitochondrial dysfunction, AMP-activated protein kinase (AMPK) activation, and
527   peroxisome proliferator-activated receptor gamma coactivator 1alpha (PGC-1alpha)
528   suppression. *J Biol Chem* **286**, 42162-42179

529   30.   Haimovich, G., Cohen-Zontag, O., and Gerst, J. E. (2016) A role for mRNA trafficking and
530   localized translation in peroxisome biogenesis and function? *Biochim Biophys Acta* **1863**,
531   911-921

532   31.   Mizuno, Y., Ninomiya, Y., Nakachi, Y., Iseki, M., Iwasa, H., Akita, M., Tsukui, T.,
533   Shimozawa, N., Ito, C., Toshimori, K., Nishimukai, M., Hara, H., Maeba, R., Okazaki, T.,

Alodaib, A. N., Al Amoudi, M., Jacob, M., Alkuraya, F. S., Horai, Y., Watanabe, M., Motegi,
H., Wakana, S., Noda, T., Kurochkin, I. V., Mizuno, Y., Schonbach, C., and Okazaki, Y.
(2013) Tysnd1 deficiency in mice interferes with the peroxisomal localization of PTS2
enzymes, causing lipid metabolic abnormalities and male infertility. *PLoS Genet* **9**,
e1003286

32.   Lee, Y. H., Judge, A. D., Seo, D., Kitade, M., Gomez-Quiroz, L. E., Ishikawa, T., Andersen,
J. B., Kim, B. K., Marquardt, J. U., Raggi, C., Avital, I., Conner, E. A., MacLachlan, I.,
Factor, V. M., and Thorgeirsson, S. S. (2011) Molecular targeting of CSN5 in human
hepatocellular carcinoma: a mechanism of therapeutic response. *Oncogene* **30**, 4175-
4184

33.   Yu, Y. P., Ding, Y., Chen, Z., Liu, S., Michalopoulos, A., Chen, R., Gulzar, Z. G., Yang, B.,
Cieply, K. M., Luvison, A., Ren, B. G., Brooks, J. D., Jarrard, D., Nelson, J. B.,
Michalopoulos, G. K., Tseng, G. C., and Luo, J. H. (2014) Novel fusion transcripts
associate with progressive prostate cancer. *Am J Pathol* **184**, 2840-2849

34.   Chen, Z. H., Yu, Y. P., Zuo, Z. H., Nelson, J. B., Michalopoulos, G. K., Monga, S., Liu, S.,
Tseng, G., and Luo, J. H. (2017) Targeting genomic rearrangements in tumor cells through
Cas9-mediated insertion of a suicide gene. *Nat Biotechnol* **35**, 543-550

35.   McAndrew, R. P., Wang, Y., Mohsen, A. W., He, M., Vockley, J., and Kim, J. J. (2008)
Structural basis for substrate fatty acyl chain specificity: crystal structure of human very-
long-chain acyl-CoA dehydrogenase. *J Biol Chem* **283**, 9435-9443

36.   Wanders, R. J., Waterham, H. R., and Ferdinandusse, S. (2015) Metabolic Interplay
between Peroxisomes and Other Subcellular Organelles Including Mitochondria and the
Endoplasmic Reticulum. *Front Cell Dev Biol* **3**, 83

37.   Schluter, A., Real-Chicharro, A., Gabaldon, T., Sanchez-Jimenez, F., and Pujol, A. (2010)
PeroxisomeDB 2.0: an integrative view of the global peroxisomal metabolome. *Nucleic
Acids Res* **38**, D800-805

38.   Wiese, S., Gronemeyer, T., Ofman, R., Kunze, M., Grou, C. P., Almeida, J. A., Eisenacher,
M., Stephan, C., Hayen, H., Schollenberger, L., Korosec, T., Waterham, H. R., Schliebs,
W., Erdmann, R., Berger, J., Meyer, H. E., Just, W., Azevedo, J. E., Wanders, R. J., and
Warscheid, B. (2007) Proteomics characterization of mouse kidney peroxisomes by
tandem mass spectrometry and protein correlation profiling. *Mol Cell Proteomics* **6**, 2045-
2057

39.   Schrader, M., Godinho, L. F., Costello, J. L., and Islinger, M. (2015) The different facets
of organelle interplay-an overview of organelle interactions. *Front Cell Dev Biol* **3**, 56

40.   Ivashchenko, O., Van Veldhoven, P. P., Brees, C., Ho, Y. S., Terlecky, S. R., and Fransen,
M. (2011) Intraperoxisomal redox balance in mammalian cells: oxidative stress and
interorganellar cross-talk. *Mol Biol Cell* **22**, 1440-1451

41.   Stolowich, N. J., Petrescu, A. D., Huang, H., Martin, G. G., Scott, A. I., and Schroeder, F.
(2002) Sterol carrier protein-2: structure reveals function. *Cell Mol Life Sci* **59**, 193-212

42.   Schwartz, D. M., and Wolins, N. E. (2007) A simple and rapid method to assay
triacylglycerol in cells and tissues. *J Lipid Res* **48**, 2514-2520

43.   Jensen, J. T., Addis, I. B., Hennebold, J. D., and Bogan, R. L. (2017) Ovarian Lipid
Metabolism Modulates Circulating Lipids in Premenopausal Women. *J Clin Endocrinol
Metab* **102**, 3138-3145

44.   Geisler, C. E., Hepler, C., Higgins, M. R., and Renquist, B. J. (2016) Hepatic adaptations
to maintain metabolic homeostasis in response to fasting and refeeding in mice. *Nutr
Metab (Lond)* **13**, 62

45.   Salway, J. G. (1969) The simultaneous determination of acetoacetate and glucose in
capillary blood. *Clin Chim Acta* **25**, 109-116

46.   Walker, P. G. (1954) A colorimetric method for the estimation of acetoacetate. *Biochem J*
58, 699-704

 Figure Legends

Figure 1: LXR agonist induction of TMEM135 expression in human hepatocyte (HepG2,

Hep3B) and monocyte-derived macrophage (THP-1) cell lines via a direct transcriptional

mechanism. Panel A contains the results from treatment of HepG2 cells in the presence or

absence of the synthetic LXR agonist T09 for 6, 12, 24 and 48 hours; while panel B is the effect

of a 24-hour T09 treatment on *TMEM135* mRNA expression in Hep3B cells (n = 4). Panel C is a

time course of T09 treatment in THP-1 cells (n = 4). For panels A-C, asterisks denote significant

(p< 0.05) difference between DMSO and T09 at the indicated timepoint. Panel D is the effect of

the protein synthesis inhibitor cycloheximide on *TMEM135* mRNA expression in THP-1 cells.

Cells were treated for 24 hours in a 2 × 2 factorial with T09 and cycloheximide (n = 5). Columns

with different letters are significantly different (p< 0.05). Panel E is Western blot analysis of

TMEM135 protein in the mitochondria-enriched fraction of HepG2 cells treated with or without

T09 for 24 hours (n = 4). The approximate molecular weights of the protein ladder are indicated,

as well as the TMEM135 and COX4I1 (housekeeping control) bands. Panel F is results of

densitometry analysis of the blot shown in Panel E, asterisk denotes significant difference (p<

0.05). For all panels, error bars indicate ± one standard error of the mean (SEM).

Figure 2: LXR/RXR heterodimers bind all three potential LXREs in the human *TMEM135*

promoter with LXRE3 mediating LXR-induced *TMEM135* transcription. Panel A is an EMSA

using LXRE1 as the fluorescent probe. The nuclear receptor(s) used in the reaction are indicated

beneath the image, with the unlabeled competitor DNA (200-fold molar excess) shown above the

image. Panel B substitutes NR1H2 for NR1H3 in the binding reactions. Panel C utilizes

fluorescent LXRE3 as the probe. The image has been cropped to only show the bound and free

fluorescent LXRE3 probe. The nuclear receptor(s) used in the reaction are indicated beneath the

image. Increasing amounts (0.25, 0.5, 1, 2, and 10-fold molar excess per competitor) of unlabeled

LXRE1 or LXRE3 were included in some reactions as indicated above the images. In Panel D, an

EMSA was performed using a mixture of LXRE1 and LXRE3 probes that were labeled with

spectrally-distinct fluorescent dyes, as well as increasing quantities of LXR/RXR proteins. The

percent of each probe bound in each reaction was determined by densitometry and plotted. Panel

E displays luciferase activity (arbitrary units normalized to β-galactosidase) derived from cells

transfected with the empty vector or the *TMEM135* promoter-containing construct in the presence

and absence of T09. The effect of increased expression of NR1H3 and RXRA is also shown (n =

4). Panel F contains the fold-increase in luciferase activity induced by T09 from the wild type

TMEM135 promoter, as well as *TMEM135* promoters containing all possible combinations of

mutant LXRE sites (n = 4). All transfections in Panel F included *NR1H3* and *RXRA* co-transfection

to increase basal expression of these nuclear receptors. An X indicates point mutations were

introduced into the corresponding LXRE (see Figure S1), Luc = luciferase. Error bars indicate one

SEM, columns without common letters are significantly ($p < 0.05$) different.

Figure 3: The LXRs are obligatory for LXR agonist-induced *TMEM135* mRNA expression,

and cause transcription of *TMEM135* with an abbreviated 5' UTR. Panels A, B, C, and D show

the effects of siRNA transfection in HepG2 cells on T09-induced mRNA expression of the LXR

isoforms *NR1H2* and *NR1H3*, *ABCA1*, and *TMEM135*; respectively. Panel E shows the effect of

LXR knockdown in HepG2 cells on T09-induced expression of *TMEM135* mRNA transcripts

containing the full 5' UTR. Primers and probe used for QPCR in panel E amplify a 76 bp region

within the 5' UTR. Panel F is results from ChIP analysis. For all panels, error bars indicate one

SEM, columns without common letters are significantly ($p < 0.05$) different (n = 4).

Figure 4: TMEM135 regulates fatty acid metabolism and proliferation of HepG2 cells. Panel

A displays the effect of TMEM135 knockdown in the presence and absence of the lipogenic LXR

agonist T09 on intracellular triglyceride accumulation in HepG2 cells. Panel B is mRNA

expression in HepG2 cells following the same treatments as Panel A for *TMEM135*, genes

involved in fatty acid oxidation (*PPARA*, *CPT1A*), and genes involved in fatty acid synthesis

(*SREBF1*, *FASN*). For panels A and B, columns without a common letter are significantly ($p <

0.05$) different (n = 4). Panel C is the effect of TMEM135 knockdown on ATP content (normalized

to total protein content) in HepG2 cells after a 4-hour incubation in glucose-free media. The chart

is plotted with the control siRNA set to 1, and the asterisk denotes significant difference (n = 4).

Panel D is the effect of TMEM135 knockdown on proliferation of HepG2 cells. Asterisks denote

significant difference at the corresponding timepoint (n = 4). Panel E is the effect of TMEM135

knockdown on cell cycle progression in HepG2 cells, and panel F is the effect on mRNA

expression of cell cycle-associated genes. For panels E-F, asterisks denote significant differences

due to TMEM135 knockdown (n = 4).

Figure 5: Liver-specific knockdown of TMEM135 *in vivo* inhibits peroxisomal β-oxidation

in male C57BL/6 mice. Panel A is mRNA expression of *Tmem135* in the listed tissues isolated

from fed mice (n = 5). Panel B displays TMEM135 protein expression in the

mitochondria/peroxisome-enriched fraction from livers. The right half of the image was incubated

with antibody pre-adsorbed with the immunizing peptide. The sample contained in each lane is

indicated beneath the images: MW = molecular weight marker, Con = control siRNA, KD =

Knockdown, *Tmem135* siRNA. The chart in panel B displays the results of densitometry analysis

of all 3 TMEM135 bands in the mitochondria/peroxisome-enriched fraction of mice liver (fed and

fasted animals pooled, n = 10 mice), as well as spectral counts from one peptide identified via

proteomic analysis that was mapped to TMEM135 (n = 3, fed animals only). Panel C displays the

change in body weight of animals from the time of siRNA injection until sacrifice 4 days later (n =

5). In Panels A-C, asterisks denote significant difference due to *Tmem135* siRNA. Panel D

displays hepatic mRNA expression of *Tmem135* and genes involved in fatty acid oxidation, panel

E displays hepatic NAD concentrations, and Panel F is hepatic and serum ketone concentrations

(BHB = β-hydroxy butyrate, total ketones = BHB + acetoacetate). For panels D-F, asterisks denote

significant difference (p< 0.05) due to *Tmem135* siRNA within feeding status, and # indicates

significant difference due to fasting in animals receiving control siRNA (n = 5).

Figure 6: Proteomic identification of a potential role for TMEM135 in import of peroxisome

proteins. Panel A displays the relative purity of the mitochondria/peroxisome-enriched fraction

used for proteomic analysis as determined by Western blot analysis of controls for mitochondria

(COX4I1), peroxisome (ABCD3), and cytosolic (TUBB) proteins. For each image, the lanes from

left to right are the molecular weight marker (MW), whole cell lysate (WC), cytosolic fraction

(Cyto), and the mitochondria/peroxisome-enriched fraction (M/P). Panel B are proteins

significantly decreased (Fisher's Exact Test, $p < 0.05$) in the mitochondria/peroxisome-enriched

fraction from TMEM135 knockdown livers, as well as the respective mean ± SEM of their spectral

counts (n = 3). Panel C illustrates peroxisomal β-oxidation of very-long chain fatty acids (VLCFA)

and dicarboxylic acids (DCA). VLCFA and DCA are transported into peroxisomes via ABCDs

(ABCD members 1-3). The enzymes catalyzing each step in the reaction are shown in boxes, and

enzymes that were significantly less abundant in the mitochondria/peroxisome-enriched fraction

from TMEM135 knockdown mice (Panel B) are shaded. Panel D illustrates peroxisomal bile acid

synthesis and β-oxidation of branched chain fatty acids (BCFA). Bile acid intermediates (BAI) and

BCFA are transported into peroxisomes via the ABCD proteins. Enzymes catalyzing each step

are shown in boxes, and shading indicates they were significantly less abundant following

TMEM135 knockdown (Panel B). CA = cholic acid, CDCA = chenodeoxycholic acid, BrAcyl-CoA

= branched chain acyl-CoA. Panel E is subcellular distribution of ACAA1, ACOX1, SCP2, and

CAT. ACOX1 migrates in 2 separate bands in Western blot, with the smaller band being

proteolytically processed ACOX1 (31). Furthermore, SCP2 (also known as SCPx) is a 58 kDa

protein that contains a 45 kDa 3-ketoacyl CoA thiolase domain and a 13 kDa sterol-carrier domain

(41), which are separated by proteolytic cleavage (31). The antibody used in the current study

recognizes the full-length 58 kDa protein (indicated by arrow) and the 45 kDa 3-ketoacyl CoA

thiolase domain, but not the 13 kDa sterol-carrier domain. The identity of the other bands in the

SCP2 blot is not known.

Figure 7: Effect of TMEM135 on import of ACAA1, ACOX1, SCP2, and CAT into

peroxisomes. Panel A contains images from Western blot analysis of whole cell and

mitochondria/peroxisome-enriched fractions (Mito/Perox-Enriched Fraction) in livers of fed and

fasted mice. Loading controls were TUBB for whole cell lysates, COX4I1 for mitochondria, and

ABCD3 for peroxisomes. The siRNA treatment is indicated beneath the lanes, Con = control

siRNA, KD = Knockdown, *Tmem135* siRNA. For SCP2, the band used for densitometry analysis

is indicated by asterisks. Panels B-C display the results of densitometry analysis of the blots

shown in Panel A. In panel B, differences between groups in peroxisome content of the

mitochondria/peroxisome-enriched fraction was determined by the ratio of ABCD3 to COX4I1. In

panel C, peroxisome import was determined by first normalizing each protein in the

mitochondria/peroxisome-enriched fraction to ABCD3 to account for differences in peroxisome

content, and the ratio relative to the whole cell lysate (normalized to TUBB) was used to

determine peroxisome import. Asterisks denote significant difference ($p < 0.05$) within feeding

status due to *Tmem135* siRNA, and # indicates significant difference due to fasting in mice

receiving the control siRNA (n = 5).

STAR Methods

CONTACT FOR REAGENT AND RESOURCE SHARING

Further information and requests for resources and reagents should be directed to and

will be fulfilled by the Lead Contact, Randy L. Bogan (boganr@email.arizona.edu). Transfer of

resources is subject to the acceptance of a Material Transfer Agreement as required by policy at

the University of Arizona. Resources are transferred and distributed by several means according

to the needs of the end-user. For other academic and non-profit researchers, the resources are

typically transferred under standard Material Transfer Agreements. For requests by for-profit

organizations, the resources are made available under standard non-exclusive license.

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Cell Lines

All cell lines were obtained from American Type Culture Collection (ATCC) and were

cultured at 37°C, 5% CO_2 in air, in a humidified incubator. Human cells (all derived from males)

included the hepatocyte lines HepG2 (ATCC® HB-8065™) and Hep 3B2.1-7 [Hep3B] (ATCC® HB-

8064™), and the monocyte line THP-1 (ATCC® TIB-202™). Murine cells included the hepatocyte

line (embryonic-derived) BNL 1NG A.2 (ATCC® TIB-76™) and the macrophage line (male) RAW

264.7 (ATCC® TIB-71™). Growth media for all cell lines except THP-1 was DMEM/F12 (Sigma

Aldrich, Inc.) supplemented with Pen/Strep (100 units/ml penicillin and 100 µg/ml streptomycin)

and 10% fetal bovine serum (FBS). THP-1 cells were incubated in RPMI media (Sigma)

supplemented with 2-mercaptoethanol (0.05 mM), Pen/Strep, and 10% FBS. THP-1 cells were

differentiated into macrophages by treatment with 12-myristate-13-acetate (PMA, 100 ng/ml) for

5 days prior to experimentation.

<u>Mice</u>

All procedures involving mice were approved by the University of Arizona IACUC. Male

C57BL/6 mice, 12 weeks of age, were used. Mice were group housed until initiation of

experimental procedures, at which point they were switched to individual caging.

METHOD DETAILS

<u>Cell Treatments</u>

T0901317 (Cayman Chemical, Inc.) was dissolved in DMSO and used at a 1 µM

concentration unless indicated otherwise. Cycloheximide (Fisher Scientific, Inc.) was dissolved in

DMSO and used at a final concentration of 50 µg/ml. The final DMSO concentration was held

constant in all groups at less than 0.2% (*v:v*). For siRNA knockdown experiments, Silencer® select

pre-designed siRNAs against human *NR1H2*, *NR1H3*, *TMEM135*, and a negative control siRNA

were purchased and transfected into HepG2 cells using Lipofectamine® RNAiMAX (Life

Technologies, Inc.) or TransIT-siQuest® reagent (Mirus Bio, LLC.) according to manufacturer

instructions. All experiments were repeated 4-5 times on different days or with different passages

of cells.

<u>HepG2 Mitochondria Isolation</u>

Cells were harvested and washed three times in 1 ml cell resuspension buffer (1 mM Tris-

HCl, pH 7.4, 130 mM NaCl, 5 mM KCl, 7.5 mM $MgCl_2$). Cells were then resuspended in 600 µl

homogenization buffer (10 mM Tris-HCl, pH 7.4, 10 mM KCl, 0.15 mM $MgCl_2$, plus protease and

phosphatase inhibitor cocktail), incubated on ice for 10 min, and homogenized for 30 strokes in a

glass dounce homogenizer with a tight-fitting pestle. The homogenate was mixed with 100 µl of

2M sucrose and centrifuged at 1,200 × g for 5 min at 4° C, and the supernatant transferred to a

new tube. This low speed centrifugation step was repeated two more times. Mitochondria were

pelleted by centrifugation at 7,000 × g for 10 min at 4° C and the supernatant removed. The pellet

was washed once with 300 µl of mitochondria wash buffer (10 mM Tris-HCl, pH 7.4, 0.15 mM

$MgCl_2$, 250 mM sucrose), and centrifuged at 9,500 × g for 5 min. Mitochondria were solubilized in

RIPA buffer containing protease and phosphatase inhibitors, and protein concentration was

determined by BCA assay (Fisher Scientific) following manufacturer recommendations.

Electrophoretic Mobility-Shift Analysis (EMSA)

The DNA sequences for human *NR1H2*, *NR1H3*, and RXR α (*RXRA*) were synthesized

by Life Technologies and cloned into the pTarget™ mammalian expression vector (Promega

Corporation) using XhoI and KpnI restriction enzyme sites. Inserts were confirmed by DNA

sequence analysis. Recombinant NR1H2, NR1H3, and RXRA proteins were produced using the

TNT® Quick Coupled Transcription/Translation System (Promega). Fluorescent probes were

synthesized by Integrated DNA Technologies and corresponded to the LXRE1 sequence with

IRDye 700 attached to the 5' end, and probes corresponding to the LXRE3 sequence were

labelled with IRDye 800. Additionally, unlabeled DNA oligos corresponding to LXREs 1-3, as well

as mutated LXREs 1-3, were synthesized (Figure S1B). Single stranded DNA oligos were

annealed by mixing sense and antisense oligos in a 1:1 molar ratio, heating to 100°C for 5

minutes, and slowly cooling to room temperature. The EMSA binding reactions were carried out

for 30 min at room temperature in 10 mM Tris, 50 mM KCl, pH 7.5; 3.5 mM DTT, 0.25% Tween-

20, 1 µg Poly (dI.dC), 5 nM fluorescent probe, and 1.5 µl of TNT lysate per protein (non-induced

TNT lysate substituted in control reactions). Unlabeled competitor DNA was added prior to the

addition of fluorescent probe at a 200-fold molar excess unless indicated otherwise. Following the

incubation period, EMSA reactions were resolved on a 6% TBE gel (Life Technologies). Imaging

and densitometry were performed with a Li-COR Odyssey CLx and Image Studio version 3.1

software, respectively.

Reporter Assays

The region from -2662 to -1 bp (relative to translation start site) of the human *TMEM135*

promoter was synthesized by Life Technologies. Additionally, the same region of the human

TMEM135 promoter was synthesized except that point mutations were introduced into each of

the three LXREs (Figure S1B). The normal and mutant *TMEM135* promoters were cloned into the

pGL4.17 vector (Promega) using SacI and XhoI restriction enzyme sites, which expresses firefly

luciferase from a promoter cloned into the vector. A unique PacI site was identified in the region

between LXRE1 and LXRE2, and a unique EcoRI site was identified in the region between LXRE2

and LXRE3. These restriction sites were used to generate six more *TMEM135* promoter/pGL4.17

constructs so that all possible combinations of wild type and mutated LXREs were obtained. All

constructs were verified using DNA sequence analysis. These reporter vectors were transfected

into HepG2 cells along with a β-galactosidase control plasmid (Promega) to normalize

transfection efficiency. In some transfections, *NR1H3*/pTarget and *RXRA*/pTarget constructs

were co-transfected. Vectors were transfected in a ratio of 60:20:10:10 (luciferase:β-

galactosidase:NR1H3:RXRA) using Lipofectamine® 3000 according to manufacturer

recommendations, with no-insert plasmids substituted in control transfections to keep DNA

concentrations constant. Each vector combination was transfected in triplicate to HepG2 cells in

96 well plates, and cells were incubated with DNA/Lipofectamine complexes for 24 hours at which

point media were changed and cells were treated for an additional 48 hours with DMSO or T09.

Cell lysates were prepared using mammalian protein extraction reagent (Fisher), and the lysate

was fractionated to quantify luciferase and β-galactosidase activity using luminescent detection

kits (Fisher). The ratio of luciferase to β-galactosidase activity was calculated, and the fold-change

in T09 vs DMSO treated cells was determined for each transfection.

<u>Chromatin Immunoprecipitation (ChIP) Analysis</u>

HepG2 cells were fixed for 10 min at room temperature with 1% formaldehyde. The

reaction was quenched by adding glycine to a 125 mM final concentration and incubating 5 min

at room temperature. The cells were trypsinized, harvested, and washed twice with PBS. Nuclei

were isolated by incubating cell pellets in 500 µl of a hypotonic buffer for 15 min on ice (20 mM

Tris-HCL, pH 7.4, 10 mM NaCl, 3 mM MgCL$_2$, with protease and phosphatase inhibitor cocktail

from Promega). Triton X-100 detergent was added to a 0.5% *v:v* final concentration to lyse the

cells followed by centrifugation at 1200 × g for 10 min to pellet nuclei. The supernatant was

removed, and the nuclei pellet washed once with 500 µl hypotonic buffer. The pellet was

resuspended in 500 µl of RIPA buffer with protease and phosphatase inhibitors, and nuclei were

sonicated in an ice-water bath using a Branson 250 digital sonifier programmed to cycle at a 10%

amplitude with 5 seconds on followed by 20 seconds off, and a 3-minute total sonication time that

was determined in preliminary time-course studies to produce chromatin of optimal length

(approximately 200-1000 base pairs). The sonicated lysates were centrifuged at 8,000 × g for 1

minute and the supernatant recovered. An aliquot of cleared lysate was treated with 25 µg RNAse

A and 50 µg proteinase K in a buffer containing 500 mM NaCl and heated at 65°C for 1.5 hours

to reverse cross-links. The DNA concentration in this aliquot was determined using a fluorometric

assay (Life Technologies) following manufacturer recommendations. The quantity of DNA used

in each ChIP was held constant within replicates (8 to 10 µg), and 10% of the input amount was

saved. Chromatin was diluted to a 500 µl final volume in binding/wash buffer (50 mM Tris, 150

mM NaCl, 1% Triton X-100) containing protease and phosphatase inhibitors, and 3 µg of antibody

was added per reaction. Tubes were incubated on an end-over-end rotator at 4°C overnight.

Antibodies used for ChIP (Abcam) were mouse monoclonal IgG2a including: 1) anti NR1H3 (clone

PPZ0412), 2) anti NR1H2 (clone K8917), and 3) non-specific control IgG2a (clone MOPC-173).

The following day 25 µl of Protein G-conjugated Dynabeads (Life Technologies) were added and

incubated for 1 hour at 4°C. The beads were recovered using a magnet and washed three times

at 4°C for 5 minutes each wash with 1 ml of binding/wash buffer. The beads were then washed

once in final wash buffer (50mM Tris, 500 mM NaCl, 1% Triton X-100). Elutions were performed

with 130 µl elution buffer (100mM $NaHCO_3$, 500mM NaCl, 1% SDS) containing 25 µg RNAse A

and 50 µg proteinase K, and ChIP reactions and input controls were incubated on a thermal mixer

at 65°C for 1.5 hours. The supernatant was recovered, and DNA purified using PCR purification

columns (Life Technologies) following manufacturer recommendations. Purified DNA was

analyzed by QPCR in a duplex reaction with primers and a Taqman probe specific for the region

spanning LXRE3 on the *TMEM135* gene, and a non-specific locus was simultaneously quantified

in the same reaction as a control for non-specific DNA carryover (Table S1). Serial dilutions of

purified sonicated DNA were included to verify linear amplification and PCR efficiency, and Ct

values from ChIP reactions and input controls were used to calculate the percentage of DNA

recovered relative to the input. Data were normalized by calculating the ratio of the TMEM135

LXRE3 to the non-specific chromatin locus.

<u>Semi-Quantitative Real-Time PCR</u>

Cells or tissue were homogenized in Trizol reagent for isolation of mRNA and were further

purified over RNeasy columns according to manufacturer recommendations (Life Technologies).

Concentrations of RNA were quantified by spectrophotometry, and 0.2-1 µg RNA was treated with

DNAse I and reverse transcribed into cDNA using the High Capacity cDNA Reverse Transcription

Kit (Life Technologies). Following reverse transcription, cDNA were diluted to 10 ng/µl (based on

original RNA concentration) and used for semi-quantitative real-time PCR analysis (QPCR).

Primer and probe sequences used for QPCR are listed in Table S1. The QPCR assays were

performed on a StepOne Plus Real-Time PCR system (Life Technologies) using TaqMan® MGB

probe or SYBR green detection. Relative mRNA abundance was determined by extrapolation of

threshold (Ct) values from a standard curve of serial cDNA dilutions and normalized to the

housekeeping gene mitochondrial ribosomal protein S10 (*MRPS10*).

<u>Cell Proliferation Assay</u>

HepG2 cells were plated at 15,000 cells per well in 96-well plates and transfected with

siRNAs immediately. At the appropriate post-transfection time, viable cells were quantified using

a CyQuant® Direct Cell Proliferation assay (Life Technologies) according to manufacturer

recommendations, and fluorescence (excitation 495 nm, emission 527 nm) determined using a

Synergy H1 plate reader (BioTek Instruments, Inc.) in area scanning mode to account for uneven

cell distribution. The number of cells per well was estimated by extrapolation from a standard

curve of increasing cell numbers that was generated at the time of plating.

<u>Flow Cytometry Analysis of Cell Cycle</u>

Freshly harvested HepG2 cell pellets were fixed by slow addition of 1 ml ice-cold 70%

ethanol while vortexing and incubated at -20°C overnight. Fixed cells were harvested by

centrifugation at 850 × g for 5 min and resuspended in 1 ml DNA staining buffer (100 µg/ml RNAse

A, 50 µg/ml propidium iodide, in PBS). Cells were incubated for 30 min at 37°C, placed on ice,

and analyzed within one hour. A Becton Dickinson FACSCANTO II flow cytometer equipped with

a 488 nm argon ion laser and a 585/42 bandpass filter was used for detection of bound propidium

iodide. List mode data files consisting of 10,000 events were acquired and the percentage of cells

in G0/G1, S, and G2/M stages was determined with CellQuest PRO software (BD Biosciences).

<u>HepG2 ATP Quantification</u>

At 48 hours post-siRNA transfection, HepG2 cells were switched to serum-free, glucose-

free, sodium pyruvate-free DMEM media and incubated at 37°C for 4 hours. Cellular ATP

concentrations were determined using a homogenous luminescent ATP detection kit (Abcam

catalog ab113849). Additional HepG2 cells treated in parallel to those used for ATP determination

were lysed in RIPA buffer and a BCA protein assay (Fisher) was performed, and ATP content

was normalized to protein concentrations.

<u>Triglyceride Quantification</u>

HepG2 cells or frozen mouse liver were sonicated in PBS containing protease and

phosphatase inhibitors (Promega) in an ice-water bath using a Branson 250 digital sonifier

programmed to cycle at a 10% amplitude with 5 seconds on followed by 20 seconds off, and a 30

second total sonication time. Triglycerides were extracted using an organic solvent procedure

optimized for triglycerides (42). Lysates were diluted 1:1 with deionized water, and 400 µl of

diluted lysate was combined in a glass tube with 1.8 ml of a 4:1 isopropanol:hexane mixture.

Samples were vortexed for 1 minute and incubated at room temperature for 30 min. Next, 500 µl

of a 1:1 hexane:diethyl ether solution was added, the samples were vortexed as before, and

incubated at room temperature for 1 minute. One ml of deionized water was added, samples were

vortexed as before, and incubated at room temperature for 20 minutes to allow phase separation.

The organic phase was transferred to a new glass tube and dried under a gentle stream of

nitrogen in a water bath at 30°C. Dried extracts were dissolved in 200 µl assay buffer (50 mM

potassium phosphate, pH 7.0, 0.01% Triton X-100) by 5 total cycles of vortexing for 1 minute and

incubating at room temperature for 3 minutes. Extracts (50 µl/well) were transferred to a clear 96

well plate, and 250 µl of triglyceride detection reagent (Pointe Scientific, Inc., catalog T7532) was

added. Triglyceride standards (Pointe Scientific) diluted in assay buffer were also run with each

assay. Plates were incubated at 37°C for 30 minutes with vigorous shaking, and absorbance at

500 nm was determined with a Synergy H1 plate reader. The protein concentration in sonicated

lysates was determined using a BCA protein assay, and triglyceride content was normalized to

protein concentrations.

<u>In Vivo siRNA Delivery and Tissue Harvest</u>

Ambion® In Vivo pre-designed siRNA (Life Technologies) against *Tmem135* (siRNA ID

s91285) or *in vivo* negative control #1 siRNA (catalog 4459405) were complexed with

Invivofectamine® 3.0 reagent (Life Technologies) according to manufacturer recommendations.

The siRNA complexes were delivered via a 200 µl tail-vein injection at a final siRNA concentration

of 1 mg/kg body weight. Mice were weighed prior to siRNA injection and again before

euthanization. Mice were euthanized 4 days after siRNA injection at 4 hours after lights on, with

or without a preceding 12 hour fast. Serum, liver, heart, adipose, and skeletal muscle were

collected and snap frozen. Tissues were powdered while frozen and stored at -80° C until used.

<u>Serum Analytes</u>

899    Total cholesterol, HDL cholesterol, and triglyceride concentrations in serum were

determined as we have described previously (43). Briefly, the HDL fraction of serum was isolated

by mixing serum 1:1 with 15% polyethylene glycol, centrifuging at 2,000 × g for 10 minutes at

room temperature, and harvesting the HDL-containing supernatant. The cholesterol

concentrations in whole serum and the HDL fraction were determined by a fluorometric enzyme

assay consisting of cholesterol esterase (0.3 units/ml), cholesterol oxidase (1 unit/ml), horse

radish peroxidase (1 unit/ml), and Amplex UltraRed® (50 µM, Life Technologies) in assay buffer

(0.1 M potassium phosphate, pH 7.0, 50 mM NaCl, 5 mM cholic acid, 0.1% Triton X-100). The

cholesterol assay was incubated at 37°C for 30 minutes, and fluorescence measured in a Synergy

H1 plate reader at 555 nm excitation and 585 nm emission wavelengths. Triglyceride

concentrations in whole serum were determined using the same assay described earlier for

HepG2 and mouse liver samples. Commercial enzymatic assays were used to quantify serum

concentrations of β-hydroxybutyrate (BHB) (Cayman Chemical, catalog 700190), non-esterified

fatty acids (NEFA) (Catachem Inc., catalog C514-0A), glucose (Pointe Scientific, Inc., catalog

G7521), and a commercial ELISA was used to quantify serum insulin (Mercodia AB, catalog 10-

1247) according to manufacturer recommendations.

<u>Liver Analytes</u>

916    Commercial kits were used for the extraction and quantification of liver NAD+ and NADH

(Abcam Inc., catalog ab176723), and glycogen (BioAssay Systems, catalog E2GN-100),

according to manufacturer recommendations. Concentrations of ATP and NEFAs in liver were

determined as we have described elsewhere (44). To quantify ketones (BHB and acetoacetate),

powdered liver was sonicated in PBS, acidified with 10% (*w:v*) trichloroacetic acid, centrifuged to

remove proteins, and the supernatant was recovered and neutralized. Concentrations of BHB in

the extract were determined using the same enzymatic assay described for serum.

Concentrations of acetoacetate were determined using a modified assay based on methods

described previously (45,46). Briefly, samples and standards were divided with one portion

subjected to boiling (100° C for 5 minutes) to degrade acetoacetate, with the remaining portion

kept on ice. A three-fold volume of freshly prepared diazo reagent (46) was then added to all

samples and incubated at room temperature in the dark for 30 minutes. The reaction was stopped

by adding a 20% volume of 2N NaOH, and absorbance was measured at 480 nm. The

absorbance from the boiled sample was subtracted from the respective non-boiled sample, and

acetoacetate concentrations were calculated from the resultant standard curve.

Liver Subcellular Fractionation

Powdered liver (approximately 50 mg) was homogenized for 40 strokes in a glass dounce

homogenizer with a tight-fitting pestle using 600 µl of cold fractionation buffer (10 mM Tris-HCl,

pH 7.4, 1 mM EGTA, 200 mM sucrose, protease and phosphatase inhibitor cocktail) on ice. The

homogenate was transferred to a new tube, and the dounce homogenizer was washed with an

additional 600 µl of fractionation buffer and transferred to the same tube. The homogenate was

centrifuged at 600 × g for 10 minutes at 4°C to pellet unbroken cells and nuclei, and the

supernatant transferred to a new tube. This low speed spin was repeated once more.

Mitochondria and peroxisomes were pelleted by centrifugation at 7,000 × g for 10 minutes, and

the supernatant (cytosolic fraction) was transferred to a new tube. The pellet was washed with

300 µl fractionation buffer and pelleted as before. The pellet was resuspended in 500 µl of

fractionation buffer and a 30 µl aliquot was pelleted and solubilized in RIPA buffer for a BCA

protein assay. Following protein assay mitochondria/peroxisome fractions were partitioned into

30 µg (protein content) aliquots followed by centrifugation and removal of the supernatant, and

the pellets were frozen at -80°C.

<u>Proteomic Analyses</u>

The mitochondria/peroxisome-enriched fraction from a subset of fed control and

TMEM135 knockdown mice was solubilized in RIPA buffer, proteins were precipitated with

acetone, and proteins were sent to the Arizona Proteomics Consortium for mass spectrometry

identification of proteins. Proteins were digested with trypsin and equal quantities of protein (500

ng) were loaded. Samples were analyzed on a Thermo Q Exactive Plus Orbitrap mass

spectrometer. Peptide identification in the resultant tandem mass spectra was performed using

Sequest version 1.3.0.339 scanning against the mouse proteome database

(Mouse_unitprotkb_proteome_2016_0720_cont.fasta). Data analysis was performed using

Scaffold version 4.8.1, with a peptide identification threshold of 95.0% and a minimum protein

identification threshold of 99.9% with 2 unique peptides.

<u>Western Blot</u>

All primary antibodies were obtained from Abcam unless otherwise indicated. Catalog

numbers and final concentrations or dilutions of each antibody were: rabbit polyclonal anti-ACAA1

(catalog ab154091, 1 µg/ml), rabbit monoclonal anti-ACOX1 (catalog ab184032, 1:2500 dilution),

rabbit monoclonal anti-CAT (catalog ab209211, 1:2000 dilution), rabbit monoclonal anti-SCP2

(catalog ab140126, 1:2500 dilution), rabbit polyclonal anti-TMEM135 (catalog ab167096, 1

µg/ml), mouse monoclonal anti-COX4I1 (catalog 14744, 0.25 µg/ml), mouse monoclonal anti-

TUBB (Sigma Aldrich catalog T8328, 0.5 µg/ml), and rabbit polyclonal anti-PMP70 (catalog

ab3421, 0.5 µg/ml).

Proteins (12.5 µg/well) were resolved on 4-12% Bis-Tris gels (Life Technologies) and

subsequently transferred to nitrocellulose membranes. The membranes were blocked with 5%

non-fat dry milk (NFDM) (*w:v*) in TBS with 0.1% (*v:v*) Tween (TBST) for 1 hr at room temp. All

primary antibodies were diluted in TBST with 1% NFDM and incubated with the membranes on a

rocking platform at 4° C overnight. The membranes were washed 4x with TBST, and IRDye®

680RD or 800CW-conjugated secondary antibodies (LI-COR Biosciences, Inc.) were diluted

1:4000 in TBST + 1% NFDM and incubated with membranes for 1 hour at room temp. The

membranes were washed as before. Imaging and densitometry were performed with a Li-COR

Odyssey CLx and Image Studio version 3.1 software, respectively.

QUANTIFICATION AND STATISTICAL ANALYSIS

Statistical analyses were performed using Stata Version 14 and all differences were

considered significant at $p < 0.05$. Information on the sample size, mean and error estimates, and

indicators of statistical significance are presented in the figures and their corresponding legends.

For immortalized cell lines, the total sample size (n) corresponds to cells on a different passage

number and/or experiments performed on different days. Time course data in immortalized cell

lines were analyzed by mixed effects regression analysis to account for repeated measures.

Treatments were coded 0 or 1, and hours post-treatment and treatment × time interaction were

included as fixed effects in the regression model with experimental replicate as the random effect.

Differences between treatments at each timepoint were determined from the treatment × time

interaction coefficient. For all other comparisons involving more than 2 groups, mixed effects

regression analysis was performed using treatment as the fixed effect and experimental replicate

as the random effect to account for repeated measures. Pairwise differences between treatments

were then determined using the Bonferroni multiple comparison method to control for the type 1

error rate. For comparisons involving only 2 groups, a paired t-test was performed.

For all mice data, the sample size refers to the number of mice. Regression analysis was

used for mice outcomes except where indicated otherwise. Samples were coded 0 or 1 for fasting,

TMEM135 knockdown fed mice, and TMEM135 knockdown fasted mice. The regression model

utilized fasting, TMEM135 knockdown in the fed state, and TMEM135 knockdown in the fasted

state, as predictors. The fasting coefficient indicated the difference due to fasting in control mice

only, while the TMEM135 knockdown coefficients indicated the difference between control mice

and TMEM135 knockdown mice within the fed state or fasted state. Differences were considered

significant at $p < 0.05$, with p values ≥ 0.05 and < 0.10 considered trends. Only fed mice were

998 used in proteomic analysis. Because the proteomic data does not follow a normal distribution due

999 to some proteins not being detected in all samples, significant differences in proteomic data were

1000 determined from weighted spectra using the non-parametric Fisher's Exact test ($p < 0.05$)

1001 contained within the Scaffold version 4.8.1 software. The list of proteins that were significantly

1002 reduced in knockdown livers was further trimmed by hand to exclude proteins that were not

1003 detected in all the samples from mice that received the control siRNA.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

A

COX4I1 ABCD3 TUBB

(MW: 75, 50, 37, 25, 20, 15, 10)

MW WC Cyto M/P

B

| | Weighted Spectral Counts ± SEM | |
Protein Symbol	Control siRNA	Tmem135 siRNA
CPS1	110 ± 16.3	84.7 ± 34.1
CAT	125 ± 7.6	76 ± 14
SCP2	128 ± 13.1	69.7 ± 35.3
HSD17B4	66 ± 8.5	46.7 ± 9.5
CES1	61.3 ± 11.8	43.3 ± 7.2
NUDT7	28 ± 8.2	17 ± 17
ACOX1	27 ± 6.7	13 ± 10.5
PC	24.3 ± 3.8	15.7 ± 4.7
ACOX2	22 ± 4.4	13.3 ± 11.9
CES3	17.7 ± 3.7	10.7 ± 6.1
GLUD1	21 ± 2.3	13 ± 3.1
DHRS4	23.3 ± 3.4	11.3 ± 6.4
HACL1	20.7 ± 3.7	9.7 ± 5.7
COMT	17.7 ± 5.8	8.3 ± 8.3
MTTP	15 ± 4.7	7.7 ± 4.3
PSAP	14 ± 4.5	5.3 ± 3.2
ACAA1	35 ± 6.5	9.3 ± 9.3
ALDH3A2	11 ± 2.9	5.3 ± 3.5
ALDH6A1	11 ± 2.9	5.7 ± 3.5
AMACR	10 ± 2.6	4 ± 4
SUCLA2	9 ± 1.7	3.3 ± 2.4
ACADL	5 ± 0.6	1.7 ± 0.9
SLCO1A1	3.7 ± 1.2	1 ± 1

C

VLCFA, DCA

ABCDs ABCDs

ACOX1
↓
EHHADH
↓
ACAA1
↓
Acyl/DCA-CoA(n-2)
+
Acetyl-CoA

D

BAI, BCFA

ABCDs ABCDs

AMACR
↓
ACOX2
↓
HSD17B4
↓
SCP2

CA/CDCA-CoA
+
Propionyl-CoA

BrAcyl-CoA(n-2)
+
Acetyl-CoA

E

ACAA1 ACOX1 SCP2 CAT

(MW: 75, 50, 37, 25, 20, 15, 10)

MW WC Cyto M/P

Figure 6

Figure 7