TMEM135 is an LXR-Inducible Regulator of Peroxisomal Metabolism

Benjamin J. Renquist[1], Thushara W. Madanayake[1], Susma Ghimire[1], Caroline E. Geisler[1], Yafei

Xu[1], Randy L. Bogan*[1]

[1]School of Animal & Comparative Biomedical Sciences, College of Agriculture & Life Sciences,

University of Arizona, Tucson, Arizona, 85721, USA

*Address correspondence to:

Randy L. Bogan, PhD, MCR

School of Animal and Comparative Biomedical Sciences

University of Arizona

PO Box 210090

Tucson, AZ 85721-0090

Phone: 520-621-1487

Fax: 520-621-9435

Email: boganr@email.arizona.edu

Summary

The liver x receptors (LXRs) are key regulators of systemic lipid metabolism. We identified transmembrane protein 135 (TMEM135), a peroxisomal protein with an unknown function, as an LXR target gene. The LXRs induce *TMEM135* transcription in humans via an LXR response element located downstream of the putative transcriptional start site, but do not increase *Tmem135* in murine cells. Functionally, knockdown of TMEM135 in hepatocytes *in vitro* and *in vivo* results in fatty acid accumulation due to reduced peroxisomal β-oxidation, which was most pronounced with elevated fatty acid loading. Mechanistically, proteomic and Western blot analyses indicated that TMEM135 mediates import of peroxisome matrix proteins necessary for β-oxidation and bile acid synthesis. These findings indicate that the LXRs regulate peroxisomal metabolism via transcriptional control of *TMEM135*, and TMEM135 mediates an auxiliary matrix protein import pathway and thus may serve as a therapeutic target for metabolic and aging disorders associated with peroxisome dysfunction.

Keywords: peroxisome, liver x receptors, nuclear receptors, lipid metabolism, β-oxidation, liver metabolism, steatosis, fatty acid metabolism, cholesterol, bile acid synthesis

Introduction

The liver x receptors (LXR) α (NR1H3) and β (NR1H2) belong to the nuclear receptor superfamily and are key regulators of cholesterol and fatty acid metabolism (Hong and Tontonoz, 2014). Oxysterols are natural LXR ligands (Peet et al., 1998) that increase during cholesterol loading. The LXRs heterodimerize with the retinoic acid receptor (RXR) and bind to LXR response elements (LXRE) within the promoters of target genes (Peet et al., 1998). The LXREs belong to the direct repeat 4 (DR4) class that consist of two hexanucleotide half-sites separated by a 4-nucleotide spacer (Teboul et al., 1995; Willy et al., 1995). The protein products of known LXR target genes affect systemic lipid metabolism. Well known examples include increasing cholesterol efflux via ATP-binding cassette (ABC) sub-family A, member 1 (ABCA1) (Costet et al., 2000), and sub-family G, member 1 (ABCG1) (Kennedy et al., 2001); limiting cholesterol

uptake via induction of myosin regulatory light chain interacting protein (MYLIP) that targets the low-density lipoprotein receptor (LDLR) for ubiquitin-mediated degradation (Zelcer et al., 2009); and increased lipogenesis via induction of sterol regulatory element binding protein 1c (SREBP1c, encoded by sterol regulatory element binding transcription factor 1 gene or *SREBF1*) (Repa et al., 2000; Yoshikawa et al., 2001). Also, the *NR1H3* isoform of the LXRs is itself an LXR target gene (Laffitte et al., 2001; Whitney et al., 2001), leading to auto-amplification of LXR actions in certain species and tissues. Although many LXR target genes have been discovered, additional target genes may remain unidentified.

We have previously investigated LXR actions within the steroidogenic corpus luteum of the ovary (Seto and Bogan, 2015; Xu et al., 2018a; Xu et al., 2018b). To uncover novel LXR target genes, rhesus macaque luteal cells were treated with vehicle or 1 µM T0901317 (T09, synthetic LXR agonist) for 24 hours, mRNA were isolated and used for microarray analysis of gene expression on the Affymetrix® Rhesus Macaque genome chip (unpublished data). Genes that were differentially expressed (>2-fold change in T09 vs vehicle, $p < 0.05$ with Benjamini and Hochberg correction for false discovery rate) were identified. Several known LXR target genes were differentially expressed including *ABCA1*, *ABCG1*, *NR1H3*, *SREBF1*, and *MYLIP*. One gene that was differentially expressed and not previously determined to be an LXR target gene was transmembrane protein 135 (*TMEM135*). Therefore, the objective of the current study is to determine if *TMEM135* is an LXR target gene and to identify its physiologic role in lipid metabolism.

Results

LXR Agonist Increases Expression of TMEM135 in Human Hepatocyte and Macrophage Cell Lines

Macrophage and hepatocyte cell lines were selected for initial studies on LXR regulation of *TMEM135* transcription as these cell types play a pivotal role in LXR-mediated reverse cholesterol transport from peripheral tissues to the liver (Hong and Tontonoz, 2014). The LXR

ligand T09 increased the mRNA expression of *TMEM135* in both HepG2 (Figure 1A) and Hep3B

(Figure 1B) hepatocyte cell lines, with an approximately 2-fold maximum increase in HepG2 cells

and a 3-fold increase in Hep3B cells. In monocyte-derived macrophages (THP-1), T09 also

induced *TMEM135* mRNA expression up to 4-fold (Figure 1C). The protein synthesis inhibitor

cycloheximide was used to determine whether the effect of T09 on *TMEM135* in THP-1 cells

requires synthesis of new proteins. Cycloheximide did not significantly affect basal or T09-

stimulated *TMEM135* mRNA expression (Figure 1D), indicating that T09 induces *TMEM135* via

a direct transcriptional mechanism not involving the synthesis of intermediary proteins. As

TMEM135 has previously been reported to be localized to mitochondria (Exil et al., 2010; Lee et

al., 2016), the effect of T09 on TMEM135 protein expression in mitochondria/peroxisome-

enriched fractions from HepG2 cells was determined. Relative concentrations of TMEM135 were

significantly ($p < 0.05$) increased approximately 1.8-fold following a 24-hour T09 treatment (Figure

1E and 1F).

A LXRE Downstream of the Putative Transcription Start Site Mediates LXR Agonist Induction of

TMEM135

A 4 kb region spanning from -3319 to +682 bp (relative to the transcriptional start site) of

the human *TMEM135* promoter was analyzed with MatInspector (Cartharius et al., 2005) for

potential LXRE binding sites. This analysis yielded three potential LXREs (Figure S1), and these

were arbitrarily designated LXRE1, LXRE2 and LXRE3 in order from most distal to most proximal

to the translation start site. Electrophoretic mobility shift assays (EMSA) were used to determine

if LXR/RXR heterodimers bind these LXREs. As shown in Figure 2A, NR1H3 and RXRA nuclear

receptors together, but not individually, caused a shift in mobility of the fluorescent LXRE1 probe

indicating that NR1H3/RXRA heterodimers bind the LXRE1 sequence in the *TMEM135* promoter.

A 200-fold molar excess of unlabeled LXRE1, LXRE2, and LXRE3 oligonucleotides eliminated

the appearance of the shifted fluorescent band while the same molar excess of mutant LXRE1,

LXRE2, and LXRE3 oligonucleotides (Figure S1B) did not prevent NR1H3/RXRA binding to the

fluorescent probe; which validated the specificity of NR1H3/RXRA heterodimer binding to all 3

LXREs in the *TMEM135* promoter (Figure 2A). Similar results were obtained for NR1H2/RXRA

heterodimers (Figure 2B), indicating that both LXR isoforms specifically bind to all 3 LXRE sites

in the *TMEM135* promoter. The LXRE1 and LXRE2 sites have identical hexanucleotide half-sites,

whereas the LXRE3 site is unique (Figure S1B). To indicate if these sequence variations may

cause differences in relative binding affinity, a fluorescent LXRE3 probe was incubated with

NR1H3/RXRA heterodimers and increasing concentrations (from 0.25 to 10-fold molar excess) of

either LXRE1 or LXRE3 unlabeled competitor DNA. Oligonucleotides with the LXRE3 sequence

more effectively inhibited binding to the fluorescent probe than oligonucleotides with the LXRE1

sequence (Figure 2C). As a complementary approach, because the LXRE1 and LXRE3 sites

were fluorescently labelled with spectrally distinct fluorophores, an EMSA was performed where

a mixture of fluorescent LXRE1 and LXRE3 probes were incubated with increasing concentrations

of NR1H3/RXRA heterodimers. This approach resulted in a higher percentage of LXRE3 than

LXRE1 bound across a range of NR1H3/RXRA heterodimer concentrations (Figure 2D).

Collectively, this indicates that LXRE3 has a higher binding affinity for NR1H3/RXR heterodimers

than LXRE1.

Luciferase reporter assays were used to study transcription initiation from the *TMEM135*

promoter. Transfection of HepG2 cells with wild type *TMEM135* promoter/pGL 4.17 resulted in a

large increase in luciferase expression that was responsive to T09 as compared to the empty pGL

4.17 vector (Figure 2E). Furthermore, luciferase activity was further amplified by co-transfection

of plasmids that constitutively express NR1H3 and RXRA (Figure 2E). Next, we determined the

requirement of each individual LXRE to LXR agonist-induced transcription from the *TMEM135*

promoter. Promoters containing all combinations of wild type and mutant LXREs (Figure S1B)

were generated, and T09-induced luciferase activity was determined. LXR-agonist induced

luciferase activity was significantly ($p < 0.05$) higher in cells transfected with the wild type

TMEM135 promoter as compared to the *TMEM135* promoter with all 3 LXREs mutated (Figure

2F). Furthermore, constructs containing the wild type LXRE3 site and mutated LXRE1 and/or

LXRE2 sites were not significantly different from the wild type promoter, while all constructs that

contained a mutated LXRE3 were not significantly different from the promoter that had all three

LXREs mutated (Figure 2F). This indicates that even though all 3 LXREs bind LXR/RXR

heterodimers, the LXRE3 site alone mediates LXR agonist-induced transcription from the

TMEM135 promoter.

We next determined whether *Tmem135* mRNA expression was induced by T09 in mice.

The murine LXRE3 site showed relatively little homology with the human sequence as a total of

3 nucleotides within both hexanucleotide half-sites were unique in mice (Figure S2A). In mouse

hepatocyte (BNL 1NG A.2) and macrophage (RAW 264.7) cell lines, T09 did not significantly

induce *Tmem135* mRNA expression (Figure S2B) whereas it did increase the known LXR target

gene *Abca1* (Figure S2C). These data indicate that *Tmem135* is not an LXR target gene in mice.

<u>*TMEM135* is a Direct Target Gene of the LXRs</u>

To directly determine whether the LXRs are needed for T09-induced *TMEM135*

transcription, siRNA-mediated knockdown of the LXRs was performed in HepG2 cells. The

NR1H2 siRNA caused an approximately 75% decrease ($p < 0.05$) in *NR1H2* mRNA expression

compared to the control and *NR1H3* siRNA groups (Figure 3A). The *NR1H3* siRNA resulted in an

approximately 70-80% decrease ($p < 0.05$) in *NR1H3* mRNA expression compared to the control

siRNA depending on the presence or absence of T09 (Figure 3B). The *NR1H2* siRNA by itself

also caused a significant reduction in T09-stimulated *NR1H3* mRNA expression and tended to

slightly improve *NR1H3* knockdown when co-transfected with the *NR1H3* siRNA (Figure 3B).

Decreased *NR1H3* mRNA expression is expected to occur following NR1H2 knockdown because

NR1H3 itself is an LXR target gene (Kennedy et al., 2001; Whitney et al., 2001). Consistent with

NR1H3 being an LXR target gene, T09 induced a significant increase in *NR1H3* in the control

siRNA group, while T09 was not as effective at inducing *NR1H3* in the *NR1H2* siRNA group

(Figure 3B).

The known LXR target gene *ABCA1* was significantly increased by T09 in the control siRNA group, and the effect of T09 was completely abolished by *NR1H2/NR1H3* siRNA co-transfection (Figure 3C). Also, *NR1H3* knockdown alone significantly inhibited the T09-induced increase in *ABCA1* (Figure 3C). Results for *TMEM135* were very similar to *ABCA1* with T09 causing a significant increase in the control siRNA group, and the effect of T09 was completely blocked by *NR1H2/NR1H3* siRNA co-transfection (Figure 3D). Furthermore, *NR1H3* knockdown itself significantly inhibited the T09-induced increase in *TMEM135* as compared to the control siRNA (Figure 3D).

The LXRE3 site of the human *TMEM135* promoter lies downstream of the putative transcriptional start site, but before the translation initiation site (Figure S1). To determine whether LXR-mediated transcription of *TMEM135* results in an mRNA with a truncated 5' UTR, primers and a probe were designed to amplify and detect a 76 base pair region encompassing the LXRE3 site in the 5' UTR. There was a significant ($p < 0.05$) reduction in mRNA expression of full-length 5' UTR transcripts caused by T09 in the control siRNA group, while this effect was blocked by *NR1H2/NR1H3* siRNA co-transfection (Figure 3E). Furthermore, *NR1H2/NR1H3* co-knockdown resulted in a significant ($p < 0.05$) increase in basal mRNA expression of full-length 5' UTR transcripts compared to all other groups (Figure 3E). ChIP was used to determine if the reduction in full-length 5' UTR transcripts is associated with increased NR1H2 and/or NR1H3 binding to the LXRE3 site on the *TMEM135* gene. Monoclonal antibodies specific for each LXR isoform were used for ChIP, and QPCR of the LXRE3 region on chromatin purified via ChIP indicated that T09 treatment caused a significant increase in binding of NR1H3 to LXRE3 in the *TMEM135* gene (Figure 3F). This indicates that LXR binding to LXRE3 increases mRNA expression of *TMEM135* via a transcript with a truncated 5' UTR relative to the putative mRNA sequence.

TMEM135 Mediates Fatty Acid Metabolism and Proliferation in HepG2 Cells

To begin unraveling the biologic function of TMEM135, a series of knockdown experiments were performed. Transfection of HepG2 cells with siRNA against *TMEM135* decreased its mRNA

expression by 70-90%. As TMEM135 has previously been implicated in fat metabolism (Exil et

al., 2010), and the LXRs are known to induce lipogenesis (Hong and Tontonoz, 2014), we first

determined its effect on triglycerides and mRNA expression of fatty acid oxidation and

lipogenesis-associated genes. Knockdown of TMEM135 significantly increased basal triglyceride

accumulation in HepG2 cells (Figure 4A). Furthermore, T09 significantly increased intracellular

triglyceride concentrations, while TMEM135 knockdown further increased triglyceride

accumulation in the presence of T09 (Figure 4A). An increase in triglyceride accumulation could

result from a decrease in fatty acid oxidation and/or an increase in lipogenesis. There were no

significant effects of TMEM135 knockdown on mRNA expression of the key regulator of fatty acid

β-oxidation, peroxisome proliferator activated receptor alpha (*PPARA*), while it tended to increase

the PPARA target gene carnitine palmitoyltransferase 1A (*CPT1A*) (Kersten and Stienstra, 2017)

(Figure 4B). It is known that the LXRs induce lipogenesis via induction of SREBP1c (*SREBF1*

gene) (Repa et al., 2000; Yoshikawa et al., 2001). As expected, T09 caused a significant increase

in *SREBF1* and the SREBP1c/LXR target gene fatty acid synthase (*FASN*) (Figure 4B), consistent

with the T09-induced increase in triglyceride accumulation (Figure 4A). However, TMEM135

knockdown significantly suppressed both basal and T09-induced *SREBF1* mRNA expression

(Figure 4B). This indicates that the increase in triglyceride accumulation in HepG2 cells occurred

despite an apparent reduction in lipogenesis. Thus, an inhibition of fatty acid oxidation seems a

more likely explanation for the increase in triglycerides. The tendency for an increase in *CPT1A*

mRNA caused by TMEM135 knockdown is not evidence against a decrease in fatty acid oxidation

because reduced β-oxidation would result in fatty acid accumulation, and because fatty acids are

the endogenous ligands for PPARA (Kersten and Stienstra, 2017), this would also cause

increased *CPT1A* mRNA expression. Further supporting an impairment in fatty acid β-oxidation,

TMEM135 knockdown significantly reduced ATP concentrations when HepG2 cells were

incubated in glucose-free medium (Figure 4C).

During these experiments it appeared that TMEM135 knockdown also inhibited replication of HepG2 cells. Because enhanced β-oxidation is a hallmark of hepatocellular carcinoma (HCC) (Beyoglu and Idle, 2013) and our previous experiments indicated a key role for TMEM135 in fatty acid metabolism, we determined whether TMEM135 regulated proliferation of HepG2 cells. Knockdown of *TMEM135* significantly ($p < 0.05$) reduced viable HepG2 cell numbers at 48 and 72 hours post-transfection as compared to cells transfected with the control siRNA (Figure 4D). Cell cycle analysis indicated that *TMEM135* knockdown significantly ($p < 0.05$) increased the percentage of HepG2 cells in the G0/G1 stage, with a corresponding significant reduction in the percentage of cells in the S phase (Figure 4E). The increase in G0/G1 arrest was associated with alterations in mRNA expression of tumor suppressor and cell cycle genes. There were significant increases in cyclin dependent kinase inhibitor 2A (*CDKN2A*) and tumor protein p53 (*TP53*) (Figure 4F), which restrict passage through the G1/S checkpoint (LaPak and Burd, 2014). There were also significant increases in cyclin dependent kinase 2 (*CDK2*) and cyclin E1 (*CCNE1*) (Figure 4F), which increase prior to passage through the G1/S checkpoint, but no change in cyclin A2 (*CCNA2*) (Figure 4F) which is increased in the S phase (Harper and Adams, 2001). Collectively, these data indicate that TMEM135 knockdown reduces HepG2 proliferation by restricting passage through the G1/S checkpoint.

Liver-Selective TMEM135 Knockdown Reduces Peroxisomal β-Oxidation

To determine the physiologic function of TMEM135, a siRNA knockdown experiment was performed in male C57BL/6 mice. Mice received either a non-targeting control or *Tmem135* siRNA and were sacrificed 4 days later in either the *ad libitum* fed state or after a 12-hour fast. Because we determined that *Tmem135* is not an LXR target gene in mice (Figure S2), we used fasting to induce hepatic fat accumulation. The *Tmem135* siRNA caused an approximately 60% mRNA knockdown in the liver while no knockdown was observed in other tissues including skeletal muscle, adipose, and heart (Figure 5A). Knockdown of TMEM135 in the liver was further confirmed by Western blot and proteomic analysis of mitochondria/peroxisome-enriched fractions

(Figure 5B). Fed mice injected with the *Tmem135* siRNA gained significantly less weight during

the 4-day treatment, while the loss in weight from fasting was similar for both siRNAs (Figure 5C).

There was no significant effect of the *Tmem135* siRNA on basal or fasting-induced hepatic

triglycerides (Figure S3A), while there was a tendency for increased hepatic non-esterified fatty

acid (NEFA) concentrations in TMEM135 knockdown mice in the fed state (Figure S3B). There

were no significant effects of TMEM135 knockdown on liver ATP or glycogen concentrations

(Figure S3C-D). Also, there were no significant effects of TMEM135 on serum lipids (total

cholesterol, HDL cholesterol, triglycerides), although there was a trend for fasting to reduce serum

triglycerides in control siRNA mice that was not observed in TMEM135 knockdown mice (Figure

S4A). There were no significant effects of TMEM135 knockdown on serum NEFA, glucose,

insulin, or β-hydroxybutyrate concentrations (Figure S4B-E). The mRNA expression of key genes

involved in fatty acid β-oxidation were determined in the liver. As expected, fasting significantly

induced mRNA expression of *Ppara*, *Cpt1a*, acyl-CoA dehydrogenase medium chain (*Acadm*),

uncoupling protein 2 (*Ucp2*), and sirtuin 3 (*Sirt3*) (Figure 5D). Similarly, knockdown of TMEM135

significantly increased *Acadm* and *Sirt3* in fed animals, while *Ucp2* was significantly reduced by

TMEM135 knockdown in fasted animals (Figure 5D). As expected with increased β-oxidative flux,

fasting significantly increased NADH in animals receiving the control siRNA, while TMEM135

knockdown significantly reduced NADH (Figure 5E) indicating an impairment in β-oxidation in

fasted mice. Reduced hepatic NADH during fasting could be due to reduced β-oxidation in

peroxisomes and/or mitochondria. However, there was a reciprocal significant increase in hepatic

ketone concentrations in fasted mice with TMEM135 knockdown (Figure 5F). Because

ketogenesis occurs exclusively in mitochondria and is intricately linked with fatty acid oxidation

(Newman and Verdin, 2017), this indicates that mitochondrial fatty acid β-oxidation was not

impaired. Thus, knockdown mice have impaired fatty acid β-oxidation in peroxisomes but not

mitochondria.

TMEM135 Knockdown Reduces Import of Peroxisome Matrix Enzymes

TMEM135 is a peroxisomal protein with high homology to the Tim17 family that mediate

translocation of proteins across mitochondrial membranes (Zarsky and Dolezal, 2016). In addition

to peroxisomes, TMEM135 has been reported to be localized to mitochondria (Exil et al., 2010;

Lee et al., 2016). Therefore, we hypothesized that TMEM135 mediates protein import into

peroxisomes and/or mitochondria. A mitochondria/peroxisome-enriched fraction was prepared

from frozen mouse livers. This fraction was validated to contain both mitochondria and

peroxisomes as indicated by presence of the mitochondria marker cytochrome c oxidase subunit

4I1 (COX4I1) and the peroxisome marker ATP binding cassette subfamily D member 3 (ABCD3,

also known as PMP70), while COX4I1 and ABCD3 were not detected in the cytosolic fraction

(Figure 6A). Furthermore, the cytosolic protein tubulin beta class I (TUBB) segregated to the

cytosolic fraction and was not detected in the mitochondria/peroxisome-enriched fraction.

Proteomic analysis of a subset of fed mice was used to provide an unbiased estimate of

differential protein abundance in the mitochondria/peroxisome-enriched fractions. A total of 23

proteins were significantly less abundant (Fisher's Exact Test, $p < 0.05$) in the

mitochondria/peroxisome-enriched fraction from TMEM135 knockdown livers (Figure 6B).

Interestingly, these 23 proteins included nearly all the matrix proteins known to be necessary for

peroxisomal bile acid synthesis and β-oxidation of very-long chain fatty acids, branched chain

fatty acids, and dicarboxylic acids (Baes and Van Veldhoven, 2016; Waterham et al., 2016)

(Figure 6 C-D). Analysis of ABCD3 expression by proteomic and Western blot analysis

demonstrated that total peroxisome content was not significantly altered by TMEM135 knockdown

(Figure 6E), indicating that the lower abundance of peroxisomal proteins in the proteomic data

was not simply due to reduced peroxisome content. Four peroxisomal proteins were selected for

further analysis via Western blot: acetyl-CoA acyltransferase 1 (ACAA1), acyl-CoA oxidase 1

(ACOX1), sterol carrier protein 2 (SCP2), and catalase (CAT). All these proteins were enriched in

the mitochondria/peroxisome fraction, but except for SCP2 (which had low detection sensitivity)

each protein was also clearly detectable in the cytosolic fraction (Figure 6F). This contrasts with

the peroxisome membrane protein ABCD3 which appeared to be exclusively localized to the

mitochondria/peroxisome fraction (Figure 6A). Collectively, detection of these proteins in the

cytoplasm indicates that their import into peroxisomes may be rate-limiting.

We next used Western blot to quantitatively determine import of ACAA1, ACOX1, SCP2,

and CAT into peroxisomes. Whole cell lysates (using TUBB as loading control) and

mitochondria/peroxisome-enriched fractions (using ABCD3 as loading control for peroxisomes)

were analyzed (Figure 7A). The signal for each protein was normalized to the respective loading

control, and the ratio of peroxisome to whole cell lysate was used to determine import. Using this

method, there were no significant differences between siRNAs for peroxisome import of ACOX1

and CAT in either fed or fasted animals (Figure 7B). However, relative ACAA1 import was

significantly reduced in TMEM135 knockdown mice when fasted (Figure 7B). Furthermore, SCP2

import was significantly lower in TMEM135 knockdown mice in both the fed and fasted state

(Figure 7B). Collectively, Western blot confirmed that TMEM135 mediates the import of ACAA1

and SCP2 into peroxisomes, and thus may mediate the import of additional proteins identified via

proteomics.

TMEM135 Knockdown Inhibits Fasting-Induced β-Oxidation

We next determined the effect of TMEM135 on markers of peroxisome function. In fasted

mice, TMEM135 knockdown resulted in a significant increase in total fatty acids and linoleic acid,

with trends ($0.05 < p < 0.1$) for increases in several other fatty acids (Figure 7C-D). In general,

mitochondria preferentially oxidize short and medium chain fatty acids (<C12), both mitochondria

and peroxisomes oxidize long chain fatty acids (C14-C18), while peroxisomes exclusively oxidize

very-long chain fatty acids (>C20) (Cipolla and Lodhi, 2017). While mitochondria are believed to

be the principal site of long chain fatty acid oxidation, peroxisomes also directly oxidize long chain

fatty acids in a cooperative manner with mitochondria (Noland et al., 2007). The peroxisomal

contribution to long chain fatty acid oxidation becomes quantitatively greater during physiologic

states of increased fatty acid load such as fasting, diabetes, and high fat diets (Baes and Van

Veldhoven, 2016). The significant increase in linoleic and total fatty acids is consistent with an

impairment in β-oxidation in fasted TMEM135 knockdown mice, and when considering hepatic

NADH and ketone concentrations (Figure 5E-F), indicates that the impairment occurred in

peroxisomes. Also, linoleic acid is an essential fatty acid obtained via dietary sources (Saini and

Keum, 2018), which indicates that TMEM135 is not altering fatty acid synthesis or desaturation.

With the exception of 22:6 that is derived from 18:3 (Saini and Keum, 2018), fasting itself did not

increase any of the very-long chain fatty acids that are exclusively oxidized by peroxisomes and

may explain why TMEM135 knockdown did not affect very-long chain fatty acid concentrations in

fasted mice. Interestingly, in fed mice TMEM135 knockdown significantly decreased 22:0 and

tended to decrease 24:0, indicating a slight increase in peroxisomal β-oxidation in the fed state.

Regarding bile acids, hepatic concentrations of cholic acid were not significantly altered by

TMEM135 knockdown (Figure S5). Collectively, these data indicate that TMEM135 may not be

obligatory to basal peroxisome function, but under physiologic conditions of elevated hepatic fatty

acid flux, TMEM135 may play a key role in enhancing peroxisomal function to assist the liver in

meeting the increased demand for β-oxidation.

Discussion

Peroxisomes have key catabolic functions including α and β-oxidation of fatty acids, and

anabolic activities including the synthesis of bile acids and ether phospholipids (Baes and Van

Veldhoven, 2016; Waterham et al., 2016). There are 2 main classes of peroxisome disorders: 1)

peroxisome biogenesis disorders, and 2) single peroxisomal enzyme deficiencies (Baes and Van

Veldhoven, 2016; Waterham et al., 2016). Outcomes from these disorders range from infant death

to mild degenerative neurosensory problems (Baes and Van Veldhoven, 2016; Waterham et al.,

2016). Hepatic pathologies are one of the principal features associated with peroxisome

disorders, which are often caused by a failure to import matrix enzymes into peroxisomes (Baes

and Van Veldhoven, 2016). In the current study we determined that TMEM135 is an LXR-

inducible protein that regulates peroxisome function by mediating an auxiliary import pathway for

matrix enzymes to enter peroxisomes.

Multiple lines of evidence support TMEM135 as being a regulator of peroxisome function.

When hepatic β-oxidation (both peroxisomal and mitochondrial) was stimulated by fasting,

TMEM135 knockdown mice had significantly lower NADH (Figure 5E) and significantly higher

total fatty acid and linoleic acid concentrations (Figure 7C-D), indicating an impairment in β-

oxidation. Meanwhile, hepatic ketones were significantly increased in knockdown mice when

fasted (Figure 5F), and ketogenesis occurs exclusively in mitochondria and is intricately linked to

β-oxidation (Newman and Verdin, 2017). Collectively, these findings indicate that β-oxidation was

impaired in peroxisomes, but not in mitochondria. The increase in ketones may have resulted

from the increased availability of fatty acids to the mitochondria and/or due to the increased

NAD+/NADH ratio, both of which stimulate ketogenesis (Newman and Verdin, 2017). Additionally,

many effects of hepatocyte-specific TMEM135 knockdown in the current study are consistent with

typical features of peroxisome dysfunction. Liver-specific peroxisome deficiency in mice

(peroxisomal biogenesis factor 5 or *Pex5* conditional knockout) results in hepatic fat accumulation

despite significantly reduced mRNA expression of *Srebf1* and increased PPARA target gene

mRNA expression, and a stunting of body growth (Peeters et al., 2011). In our study, knockdown

of TMEM135 caused fatty acid accumulation both *in vitro* and *in vivo.* In HepG2 cells, increased

triglycerides occurred despite a simultaneous significant reduction in mRNA expression of

SREBF1 (Figure 4A-B), and some PPARA target genes were increased by TMEM135 knockdown

in fed mice (Figure 5D). Also, TMEM135 knockdown caused a significant reduction in body weight

gain of knockdown mice in the fed state (Figure 5C). Collectively, these data indicate that

TMEM135 mediates peroxisome function.

Mechanistically, our data indicate that TMEM135 modulates peroxisome function by

regulating the import of matrix enzymes into peroxisomes. This differs from the classical

mechanism involving peroxisomal targeting signals (PTS), which are short amino acid sequences

at the N or C-terminus that interact with various members of the peroxin family of proteins to

facilitate import into the peroxisome matrix (Haimovich et al., 2016). Among the proteins whose

import was quantified by Western blot in the current study ACOX1 and SCP2 contain a PTS1

sequence on their C-terminus that is recognized by PEX5, and ACAA1 contains a PTS2 sequence

at its N-terminus that interacts with PEX7 (peroxisomal biogenesis factor 7) (Haimovich et al.,

2016; Mizuno et al., 2013). Therefore, TMEM135 is likely not obligatory for their import, but rather

serves an auxiliary role. It is not known whether TMEM135-mediated protein import occurs via a

mechanism that involves, or occurs independently of, peroxins. Because TMEM135 is a

transmembrane protein with TIM17 homology (Zarsky and Dolezal, 2016), a peroxin-independent

mechanism seems most likely. Regardless, TMEM135-mediated protein import may be

necessary under conditions of high peroxisome activity when peroxin-dependent import becomes

saturated, which is consistent with our finding that fasting-induced hepatic β-oxidation is

dependent on TMEM135.

An auxiliary role of TMEM135 in peroxisome protein import makes it a potential therapeutic

target, and because it is also an LXR target gene in humans, pharmacologic modulation of the

LXRs may be useful for altering peroxisome metabolism. The LXRs are master regulators of

reverse cholesterol transport (RCT), a process whereby excess cholesterol is removed from

peripheral tissues and transported to the liver for elimination in bile acids (Hong and Tontonoz,

2014). There has been much interest in the development of LXR agonists to treat human diseases

including atherosclerosis, Alzheimer's disease, and other metabolic disorders (Hong and

Tontonoz, 2014). However, the therapeutic potential of LXR agonists has been limited because

they also increase hepatic lipogenesis via induction of SREBP1c (Repa et al., 2000; Yoshikawa

et al., 2001). Our data indicate that the LXRs can stimulate peroxisomal β-oxidation by increasing

transcription of *TMEM135*, which consequently may limit steatosis during LXR-induced

lipogenesis. It should be noted that our data indicate that *TMEM135* is an LXR target gene in

humans, but in mice *Tmem135* is either not induced by the LXRs or is a low-affinity target gene

(Figure S2). Thus, LXR agonist-induced steatosis may be less pronounced in human compared

to mice hepatocytes due to species differences in LXR regulation of *TMEM135* transcription.

Previous evidence of a link between the LXRs and peroxisomal β-oxidation is limited and

conflicting. One group reported increased hepatic peroxisomal β-oxidation in mice treated with an

LXR agonist through an undefined mechanism involving increased oxidative gene expression

(Beyer et al., 2004; Hu et al., 2005). However, another report indicates that the LXRs suppress

peroxisomal β-oxidation by inhibiting expression of the peroxisomal membrane transporter

ABCD2 (Gondcaille et al., 2014). It would be interesting to determine how LXR-mediated

suppression of ABCD2 and induction of TMEM135 influences overall peroxisome function.

Elevated TMEM135 may protect against lipid loading but may also contribute to hepatic

pathologies. For example, metabolomics studies have reported that elevated serum bile acids

and urinary bile salts represent the core metabolic phenotype in the transition from healthy liver

to nonalcoholic fatty liver disease, nonalcoholic steatohepatitis, cirrhosis, and HCC (Beyoglu and

Idle, 2013). Additionally, elevated β-oxidation and reduced mitochondrial respiration are part of

the metabolic remodeling that occurs in chronic liver disease (Beyoglu and Idle, 2013). These

metabolic changes are consistent with high peroxisome activity as peroxisomes play a key role in

bile acid synthesis, they contribute to fatty acid β-oxidation, and can also indirectly alter

mitochondrial respiration (Baes and Van Veldhoven, 2016). In the current study TMEM135

knockdown reduced proliferation of the HepG2 cell line by increasing G0/G1 arrest. Also, a

metabolomics study found that linoleic acid is significantly reduced in human HCC tumors

compared to nontumor liver tissue from the same subjects (Beyoglu et al., 2013), and in our data

linoleic acid was significantly increased by TMEM135 knockdown *in vivo* (Figure 7C), consistent

with a role for TMEM135 in pathogenic β-oxidation. There is additional evidence in the literature

of an oncogenic role for TMEM135. It was among genes identified by transcriptomic analysis as

being significantly reduced in expression when proliferation of HepG2 and Huh7 cells were

inhibited by knockdown of the COP9 signalosome subunit 5 (Lee et al., 2011). Also, a genomic

rearrangement that results in a TMEM135-containing fusion transcript has been linked with

aggressive prostate cancer (Yu et al., 2014), and Cas9-mediated insertion of a suicide gene into

cells expressing this same fusion transcript reduces tumor burden and mortality in mouse

xenografts (Chen et al., 2017). Thus, TMEM135 may represent a novel target in the treatment of

certain cancers and chronic liver disease.

Previous studies to determine the biological function of TMEM135 are limited. The first

indication of its function came from a mouse model of mitochondrial acyl-CoA dehydrogenase

very long chain (ACADVL) deficiency (Exil et al., 2010). Under conditions of increased demand

for β-oxidation such as fasting and cold stress, ACADVL-deficient mice have reduced survival

due to cardiac dysfunction (Exil et al., 2010). It was discovered that TMEM135 was elevated more

than four-fold in the hearts of ACADVL-deficient mice that survived birth (Exil et al., 2010). The

ACADVL enzyme catalyzes the first step in mitochondrial β-oxidation using fatty acids larger than

14 carbons as substrates (McAndrew et al., 2008). Because ACADVL deficiency restricts the

mitochondria from directly oxidizing long chain fatty acids, data from the current study indicate

that the increase in TMEM135 likely promoted survival by increasing peroxisomal β-oxidation and

the processing of long and very-long chain fatty acids to medium chain fatty acids that were

subsequently oxidized in the mitochondria in an ACADVL-independent manner. Another study

into TMEM135 function reported that a mutation in *Tmem135* was responsible for accelerated

aging of the retina in a mouse model of age-related macular degeneration (Lee et al., 2016).

Neurodegenerative disorders are commonly associated with peroxisome dysfunction (Cipolla and

Lodhi, 2017), which is consistent with a *Tmem135* mutation causing accelerated aging in the

retina. Both of these previous studies on TMEM135 reported that it is localized to mitochondria

(Exil et al., 2010; Lee et al., 2016), whereas our data indicates it functions in peroxisomes. In

support of a peroxisomal origin of action, another previous study clearly demonstrated that

TMEM135 (formerly known as PMP52) is localized to peroxisomes and not mitochondria (Wiese

et al., 2007). Because peroxisomes and mitochondria can physically associate via the formation

of tethering complexes (Schrader et al., 2015), it seems likely that reports of TMEM135

association with mitochondria (Exil et al., 2010; Lee et al., 2016) were due to peroxisomes that

were tethered to mitochondria, and TMEM135 effects on mitochondria are indirectly mediated via

altered peroxisome function.

The critical role of peroxisomes in human health is illustrated by the spectrum of

peroxisome disorders; but in addition to these inborn errors of metabolism it is becoming

increasingly apparent that peroxisomes play key roles in other metabolic and age-related

diseases including cancer, neurodegenerative disorders, and diabetes (Cipolla and Lodhi, 2017).

This may be due to direct catabolic and anabolic actions of peroxisomes, as well as interactions

between peroxisomes and other organelles including mitochondria (Cipolla and Lodhi, 2017;

Schrader et al., 2015). Our data indicate that TMEM135 regulates peroxisome function by

mediating an auxiliary matrix protein import pathway. Furthermore, the effects of TMEM135 are

inducible as it is an LXR target gene in humans, and it is likely to be regulated by other metabolic

signaling pathways. In support of this, PPARA is the master regulator of β-oxidation in the liver

and in primary human hepatocytes *TMEM135* mRNA expression is consistently induced by

PPARA agonists (Kersten and Stienstra, 2017) and its promoter is bound by PPARA as

determined using ChIP-seq (McMullen et al., 2014), indicating it is also a PPARA target gene.

These findings implicate TMEM135 as a potential therapeutic target in the treatment of metabolic

and age-related diseases that are associated with peroxisome dysfunction.

Acknowledgements

Mass spectrometry and proteomics data were acquired by the Arizona Proteomics

Consortium at the BIO5 Institute of the University of Arizona. The authors would like to thank Dr.

Jon Hennebold at the Oregon National Primate Research Center of Oregon Health & Science

University for assistance with the unpublished microarray experiment in macaque luteal cells that

provided preliminary data for the current study. The authors would also like to thank Dr. André-

Denis Wright for arranging Departmental support to ensure completion of this project. This

research was supported by the Eunice Kennedy Shriver National Institute of Child Health &

Human Development, National Institutes of Health, Award Number R00 HD067678 to R.L.B. The

content is solely the responsibility of the authors and does not necessarily represent the official

views of the National Institutes of Health. This work was also supported by the Arizona Biomedical

Research Commission Early Stage Investigator Award No. ADHS14-082986 to B.J.R.

Author Contributions

Conceptualization, R.L.B. and B.J.R.; Methodology R.L.B. and B.J.R.; Investigation,

R.L.B., B.J.R., T.W.M., S.G., C.E.G., and Y.X.; Resources, R.L.B. and B.J.R.; Writing – Original

Draft, R.L.B.; Writing – Review & Editing, R.L.B. and B.J.R.; Funding Acquisition, R.L.B. and

B.J.R.

Declaration of Interests

The authors declare no competing interests.

References

Baes, M., and Van Veldhoven, P.P. (2016). Hepatic dysfunction in peroxisomal disorders. Biochim
Biophys Acta *1863*, 956-970.
Beyer, T.P., Schmidt, R.J., Foxworthy, P., Zhang, Y., Dai, J., Bensch, W.R., Kauffman, R.F., Gao,
H., Ryan, T.P., Jiang, X.C., et al. (2004). Coadministration of a liver X receptor agonist
and a peroxisome proliferator activator receptor-alpha agonist in Mice: effects of nuclear
receptor interplay on high-density lipoprotein and triglyceride metabolism in vivo. J
Pharmacol Exp Ther *309*, 861-868.
Beyoglu, D., and Idle, J.R. (2013). The metabolomic window into hepatobiliary disease. J Hepatol
59, 842-858.
Beyoglu, D., Imbeaud, S., Maurhofer, O., Bioulac-Sage, P., Zucman-Rossi, J., Dufour, J.F., and
Idle, J.R. (2013). Tissue metabolomics of hepatocellular carcinoma: tumor energy
metabolism and the role of transcriptomic classification. Hepatology *58*, 229-238.
Cartharius, K., Frech, K., Grote, K., Klocke, B., Haltmeier, M., Klingenhoff, A., Frisch, M.,
Bayerlein, M., and Werner, T. (2005). MatInspector and beyond: promoter analysis based
on transcription factor binding sites. Bioinformatics *21*, 2933-2942.
Chen, Z.H., Yu, Y.P., Zuo, Z.H., Nelson, J.B., Michalopoulos, G.K., Monga, S., Liu, S., Tseng, G.,
and Luo, J.H. (2017). Targeting genomic rearrangements in tumor cells through Cas9-
mediated insertion of a suicide gene. Nat Biotechnol *35*, 543-550.
Cipolla, C.M., and Lodhi, I.J. (2017). Peroxisomal Dysfunction in Age-Related Diseases. Trends
Endocrinol Metab *28*, 297-308.
Costet, P., Luo, Y., Wang, N., and Tall, A.R. (2000). Sterol-dependent transactivation of the ABC1
promoter by the liver X receptor/retinoid X receptor. J Biol Chem *275*, 28240-28245.
Exil, V.J., Silva Avila, D., Benedetto, A., Exil, E.A., Adams, M.R., Au, C., and Aschner, M. (2010).
Stressed-induced TMEM135 protein is part of a conserved genetic network involved in fat
storage and longevity regulation in Caenorhabditis elegans. PLoS One *5*, e14228.

Geisler, C.E., Hepler, C., Higgins, M.R., and Renquist, B.J. (2016). Hepatic adaptations to
maintain metabolic homeostasis in response to fasting and refeeding in mice. Nutr Metab
(Lond) *13*, 62.
Gondcaille, C., Genin, E.C., Lopez, T.E., Dias, A.M., Geillon, F., Andreoletti, P., Cherkaoui-Malki,
M., Nury, T., Lizard, G., Weinhofer, I., et al. (2014). LXR antagonists induce ABCD2
expression. Biochim Biophys Acta *1841*, 259-266.
Haimovich, G., Cohen-Zontag, O., and Gerst, J.E. (2016). A role for mRNA trafficking and
localized translation in peroxisome biogenesis and function? Biochim Biophys Acta *1863*,
911-921.
Harper, J.W., and Adams, P.D. (2001). Cyclin-dependent kinases. Chem Rev *101*, 2511-2526.
Hong, C., and Tontonoz, P. (2014). Liver X receptors in lipid metabolism: opportunities for drug
discovery. Nat Rev Drug Discov *13*, 433-444.
Hu, T., Foxworthy, P., Siesky, A., Ficorilli, J.V., Gao, H., Li, S., Christe, M., Ryan, T., Cao, G.,
Eacho, P., et al. (2005). Hepatic peroxisomal fatty acid beta-oxidation is regulated by liver
X receptor alpha. Endocrinology *146*, 5380-5387.
Jensen, J.T., Addis, I.B., Hennebold, J.D., and Bogan, R.L. (2017). Ovarian Lipid Metabolism
Modulates Circulating Lipids in Premenopausal Women. J Clin Endocrinol Metab *102*,
3138-3145.
Kennedy, M.A., Venkateswaran, A., Tarr, P.T., Xenarios, I., Kudoh, J., Shimizu, N., and Edwards,
P.A. (2001). Characterization of the human ABCG1 gene: liver X receptor activates an
internal promoter that produces a novel transcript encoding an alternative form of the
protein. J Biol Chem *276*, 39438-39447.
Kersten, S., and Stienstra, R. (2017). The role and regulation of the peroxisome proliferator
activated receptor alpha in human liver. Biochimie *136*, 75-84.
Laffitte, B.A., Joseph, S.B., Walczak, R., Pei, L., Wilpitz, D.C., Collins, J.L., and Tontonoz, P.
(2001). Autoregulation of the human liver X receptor alpha promoter. Mol Cell Biol *21*,
7558-7568.
LaPak, K.M., and Burd, C.E. (2014). The molecular balancing act of p16(INK4a) in cancer and
aging. Mol Cancer Res *12*, 167-183.
Lee, W.H., Higuchi, H., Ikeda, S., Macke, E.L., Takimoto, T., Pattnaik, B.R., Liu, C., Chu, L.F.,
Siepka, S.M., Krentz, K.J., et al. (2016). Mouse Tmem135 mutation reveals a mechanism
involving mitochondrial dynamics that leads to age-dependent retinal pathologies. Elife *5*.
Lee, Y.H., Judge, A.D., Seo, D., Kitade, M., Gomez-Quiroz, L.E., Ishikawa, T., Andersen, J.B.,
Kim, B.K., Marquardt, J.U., Raggi, C., et al. (2011). Molecular targeting of CSN5 in human
hepatocellular carcinoma: a mechanism of therapeutic response. Oncogene *30*, 4175-
4184.
Masood, A., Stark, K.D., and Salem, N., Jr. (2005). A simplified and efficient method for the
analysis of fatty acid methyl esters suitable for large clinical studies. J Lipid Res *46*, 2299-
2305.
McAndrew, R.P., Wang, Y., Mohsen, A.W., He, M., Vockley, J., and Kim, J.J. (2008). Structural
basis for substrate fatty acyl chain specificity: crystal structure of human very-long-chain
acyl-CoA dehydrogenase. J Biol Chem *283*, 9435-9443.
McMullen, P.D., Bhattacharya, S., Woods, C.G., Sun, B., Yarborough, K., Ross, S.M., Miller, M.E.,
McBride, M.T., LeCluyse, E.L., Clewell, R.A., et al. (2014). A map of the PPARalpha
transcription regulatory network for primary human hepatocytes. Chem Biol Interact *209*,
14-24.
Mizuno, Y., Ninomiya, Y., Nakachi, Y., Iseki, M., Iwasa, H., Akita, M., Tsukui, T., Shimozawa, N.,
Ito, C., Toshimori, K., et al. (2013). Tysnd1 deficiency in mice interferes with the
peroxisomal localization of PTS2 enzymes, causing lipid metabolic abnormalities and
male infertility. PLoS Genet *9*, e1003286.

Newman, J.C., and Verdin, E. (2017). beta-Hydroxybutyrate: A Signaling Metabolite. Annu Rev
Nutr *37*, 51-76.
Noland, R.C., Woodlief, T.L., Whitfield, B.R., Manning, S.M., Evans, J.R., Dudek, R.W., Lust,
R.M., and Cortright, R.N. (2007). Peroxisomal-mitochondrial oxidation in a rodent model
of obesity-associated insulin resistance. Am J Physiol Endocrinol Metab *293*, E986-
E1001.
Peet, D.J., Janowski, B.A., and Mangelsdorf, D.J. (1998). The LXRs: a new class of oxysterol
receptors. Curr Opin Genet Dev *8*, 571-575.
Peeters, A., Swinnen, J.V., Van Veldhoven, P.P., and Baes, M. (2011). Hepatosteatosis in
peroxisome deficient liver despite increased beta-oxidation capacity and impaired
lipogenesis. Biochimie *93*, 1828-1838.
Repa, J.J., Liang, G., Ou, J., Bashmakov, Y., Lobaccaro, J.M., Shimomura, I., Shan, B., Brown,
M.S., Goldstein, J.L., and Mangelsdorf, D.J. (2000). Regulation of mouse sterol regulatory
element-binding protein-1c gene (SREBP-1c) by oxysterol receptors, LXRalpha and
LXRbeta. Genes Dev *14*, 2819-2830.
Saini, R.K., and Keum, Y.S. (2018). Omega-3 and omega-6 polyunsaturated fatty acids: Dietary
sources, metabolism, and significance - A review. Life Sci *203*, 255-267.
Salway, J.G. (1969). The simultaneous determination of acetoacetate and glucose in capillary
blood. Clin Chim Acta *25*, 109-116.
Schrader, M., Godinho, L.F., Costello, J.L., and Islinger, M. (2015). The different facets of
organelle interplay-an overview of organelle interactions. Front Cell Dev Biol *3*, 56.
Schwartz, D.M., and Wolins, N.E. (2007). A simple and rapid method to assay triacylglycerol in
cells and tissues. J Lipid Res *48*, 2514-2520.
Seto, N.L., and Bogan, R.L. (2015). Decreased cholesterol uptake and increased liver x receptor-
mediated cholesterol efflux pathways during prostaglandin F2 alpha-induced and
spontaneous luteolysis in sheep. Biol Reprod *92*, 128.
Stolowich, N.J., Petrescu, A.D., Huang, H., Martin, G.G., Scott, A.I., and Schroeder, F. (2002).
Sterol carrier protein-2: structure reveals function. Cell Mol Life Sci *59*, 193-212.
Teboul, M., Enmark, E., Li, Q., Wikstrom, A.C., Pelto-Huikko, M., and Gustafsson, J.A. (1995).
OR-1, a member of the nuclear receptor superfamily that interacts with the 9-cis-retinoic
acid receptor. Proc Natl Acad Sci U S A *92*, 2096-2100.
Walker, P.G. (1954). A colorimetric method for the estimation of acetoacetate. Biochem J *58*, 699-
704.
Waterham, H.R., Ferdinandusse, S., and Wanders, R.J. (2016). Human disorders of peroxisome
metabolism and biogenesis. Biochim Biophys Acta *1863*, 922-933.
Whitney, K.D., Watson, M.A., Goodwin, B., Galardi, C.M., Maglich, J.M., Wilson, J.G., Willson,
T.M., Collins, J.L., and Kliewer, S.A. (2001). Liver X receptor (LXR) regulation of the
LXRalpha gene in human macrophages. J Biol Chem *276*, 43509-43515.
Wiese, S., Gronemeyer, T., Ofman, R., Kunze, M., Grou, C.P., Almeida, J.A., Eisenacher, M.,
Stephan, C., Hayen, H., Schollenberger, L., et al. (2007). Proteomics characterization of
mouse kidney peroxisomes by tandem mass spectrometry and protein correlation
profiling. Mol Cell Proteomics *6*, 2045-2057.
Willy, P.J., Umesono, K., Ong, E.S., Evans, R.M., Heyman, R.A., and Mangelsdorf, D.J. (1995).
LXR, a nuclear receptor that defines a distinct retinoid response pathway. Genes Dev *9*,
1033-1045.
Xu, Y., Hernandez-Ledezma, J.J., Hutchison, S.M., and Bogan, R.L. (2018a). The liver X
receptors and sterol regulatory element binding proteins alter progesterone secretion and
are regulated by human chorionic gonadotropin in human luteinized granulosa cells. Mol
Cell Endocrinol *473*, 124-135.

Xu, Y., Hutchison, S.M., Hernandez-Ledezma, J.J., and Bogan, R.L. (2018b). Increased 27-
hydroxycholesterol production during luteolysis may mediate the progressive decline in
progesterone secretion. Mol Hum Reprod *24*, 2-13.
Yoshikawa, T., Shimano, H., Amemiya-Kudo, M., Yahagi, N., Hasty, A.H., Matsuzaka, T.,
Okazaki, H., Tamura, Y., Iizuka, Y., Ohashi, K., et al. (2001). Identification of liver X
receptor-retinoid X receptor as an activator of the sterol regulatory element-binding protein
1c gene promoter. Mol Cell Biol *21*, 2991-3000.
Yu, Y.P., Ding, Y., Chen, Z., Liu, S., Michalopoulos, A., Chen, R., Gulzar, Z.G., Yang, B., Cieply,
K.M., Luvison, A., et al. (2014). Novel fusion transcripts associate with progressive
prostate cancer. Am J Pathol *184*, 2840-2849.
Zarsky, V., and Dolezal, P. (2016). Evolution of the Tim17 protein family. Biol Direct *11*, 54.
Zelcer, N., Hong, C., Boyadjian, R., and Tontonoz, P. (2009). LXR regulates cholesterol uptake
through Idol-dependent ubiquitination of the LDL receptor. Science *325*, 100-104.

Figure Legends

Figure 1: A synthetic LXR agonist induces TMEM135 expression in human hepatocyte

(HepG2, Hep3B) and monocyte-derived macrophage (THP-1) cell lines via a direct

transcriptional mechanism. Panel A contains the results from treatment of HepG2 cells in the

presence or absence of the synthetic LXR agonist T09 for 6, 12, 24 and 48 hours; while panel B

is the effect of a 24-hour T09 treatment on *TMEM135* mRNA expression in Hep3B cells (n = 4).

Panel C is a time course of T09 treatment in THP-1 cells (n = 4). For panels A-C, asterisks denote

significant (p< 0.05) difference between DMSO and T09 at the indicated timepoint. Panel D is the

effect of the protein synthesis inhibitor cycloheximide on *TMEM135* mRNA expression in THP-1

cells. Cells were treated for 24 hours in a 2 × 2 factorial with T09 and cycloheximide (n = 5).

Columns with different letters are significantly different (p< 0.05). Panel E is Western blot analysis

of TMEM135 protein in the mitochondria/peroxisome-enriched fraction of HepG2 cells treated with

or without T09 for 24 hours (n = 4). The approximate molecular weights of the protein ladder are

indicated, as well as the TMEM135 and COX4I1 (housekeeping control) bands. Panel F is results

of densitometry analysis of the blot shown in Panel E, asterisk denotes significant difference (p<

0.05). For all panels, error bars indicate ± one standard error of the mean (SEM).

Figure 2: The LXRs bind all three potential LXREs in the human *TMEM135* promoter with

LXRE3 mediating LXR-induced *TMEM135* transcription. Panel A is an EMSA using LXRE1 as

the fluorescent probe. The nuclear receptor(s) used in the reaction are indicated beneath the

image, with the unlabeled competitor DNA (200-fold molar excess) shown above the image. Panel

B substitutes NR1H2 for NR1H3 in the binding reactions. Panel C utilizes fluorescent LXRE3 as

the probe. The image has been cropped to only show the bound and free fluorescent LXRE3

probe. The nuclear receptor(s) used in the reaction are indicated beneath the image. Increasing

amounts (0.25, 0.5, 1, 2, and 10-fold molar excess per competitor) of unlabeled LXRE1 or LXRE3

were included in some reactions as indicated above the images. In Panel D, an EMSA was

performed using a mixture of LXRE1 and LXRE3 probes that were labeled with spectrally-distinct

fluorescent dyes, as well as increasing quantities of LXR/RXR proteins. The percent of each probe

bound in each reaction was determined by densitometry and plotted. Panel E displays luciferase

activity (arbitrary units normalized to β-galactosidase) derived from cells transfected with the

empty vector or the *TMEM135* promoter-containing construct in the presence and absence of

T09. The effect of increased expression of NR1H3 and RXRA is also shown (n = 4). Panel F

contains the fold-increase in luciferase activity induced by T09 from the wild type *TMEM135*

promoter, as well as *TMEM135* promoters containing all possible combinations of mutant LXRE

sites (n = 4). All transfections in Panel F included *NR1H3* and *RXRA* co-transfection to increase

basal expression of these nuclear receptors. An X indicates point mutations were introduced into

the corresponding LXRE (see Figure S1), Luc = luciferase. Error bars indicate one SEM, columns

without common letters are significantly (p< 0.05) different. See also Figures S1 and S2.

Figure 3: The LXRs are obligatory for LXR agonist-induced *TMEM135* mRNA expression,

and cause transcription of *TMEM135* with an abbreviated 5' UTR. Panels A, B, C, and D show

the effects of siRNA transfection in HepG2 cells on T09-induced mRNA expression of the LXR

isoforms *NR1H2* and *NR1H3*, *ABCA1*, and *TMEM135*; respectively. Panel E shows the effect of

LXR knockdown in HepG2 cells on T09-induced expression of *TMEM135* mRNA transcripts

containing the full 5' UTR. Primers and probe used for QPCR in panel E amplify a 76 bp region

within the 5' UTR that encompasses the LXR binding site in genomic DNA. Panel F is results from

ChIP analysis. For all panels, error bars indicate one SEM, columns without common letters are

significantly (p< 0.05) different (n = 4).

Figure 4: TMEM135 regulates fatty acid metabolism and proliferation of HepG2 cells. Panel

A displays the effect of TMEM135 knockdown in the presence and absence of the lipogenic LXR

agonist T09 on intracellular triglyceride accumulation in HepG2 cells. Panel B is mRNA

expression in HepG2 cells for *TMEM135*, genes involved in fatty acid oxidation (*PPARA*, *CPT1A*),

and genes involved in fatty acid synthesis (*SREBF1*, *FASN*). For panels A and B, columns without

a common letter are significantly (p< 0.05) different (n = 4). Panel C is the effect of TMEM135

knockdown on ATP content (normalized to total protein content) in HepG2 cells after a 4-hour

incubation in glucose-free, sodium pyruvate-free media. The chart is plotted with the control

siRNA set to 1, and the asterisk denotes a significant difference (n = 4). Panel D is the effect of

TMEM135 knockdown on proliferation of HepG2 cells. Asterisks denote significant difference at

the corresponding timepoint (n = 4). Panel E is the effect of TMEM135 knockdown on cell cycle

progression in HepG2 cells, and panel F is the effect on mRNA expression of cell cycle-associated

genes. For panels E-F, asterisks denote significant differences due to TMEM135 knockdown (n

= 4).

Figure 5: Liver-specific knockdown of TMEM135 inhibits peroxisomal β-oxidation. Panel A

is mRNA expression of *Tmem135* in the listed tissues isolated from fed mice (n = 5). Panel B

displays TMEM135 protein expression in the mitochondria/peroxisome-enriched fraction from

livers. The right half of the image was incubated with antibody pre-absorbed with the immunizing

peptide. The sample contained in each lane is indicated beneath the images: MW = molecular

weight marker, Con = control siRNA, KD = Knockdown, *Tmem135* siRNA. The chart in panel B

displays the results of densitometry analysis of all 3 TMEM135 bands in the

mitochondria/peroxisome-enriched fraction of mice liver (fed and fasted animals pooled, n = 10

mice), as well as spectral counts from one peptide identified via proteomic analysis that was

mapped to TMEM135 (n = 3, fed animals only). In Panels A-B, asterisks denote significant

difference due to *Tmem135* siRNA. Panel C displays the change in body weight of animals from

the time of siRNA injection until sacrifice 4 days later (n = 5). Panel D displays hepatic mRNA

expression of *Tmem135* and genes involved in fatty acid oxidation, panel E displays hepatic NAD

concentrations, and Panel F is hepatic ketone concentrations (total ketones = β-hydroxybutyrate

+ acetoacetate). For panels C-F, asterisks denote significant difference ($p < 0.05$) due to

Tmem135 siRNA within feeding status, and # indicates significant difference due to fasting in

animals receiving control siRNA (n = 5). See also Figures S3 and S4.

Figure 6: TMEM135 mediates import of peroxisome matrix enzymes. Panel A displays the

relative purity of the mitochondria/peroxisome-enriched fraction used for proteomic analysis as

determined by Western blot analysis of controls for mitochondria (COX4I1), peroxisome (ABCD3),

and cytosolic (TUBB) proteins. For each image, the lanes from left to right are the molecular

weight marker (MW), whole cell lysate (WC), cytosolic fraction (Cyto), and the

mitochondria/peroxisome-enriched fraction (M/P). Panel B are proteins significantly decreased

(Fisher's Exact Test, $p < 0.05$) in the mitochondria/peroxisome-enriched fraction from TMEM135

knockdown livers, as well as the respective mean ± SEM of their spectral counts (n = 3, fed mice

only). Panel C illustrates peroxisomal β-oxidation of very-long chain fatty acids (VLCFA) and

dicarboxylic acids (DCA). VLCFA and DCA are transported into peroxisomes via ABCDs (ABCD

members 1-3). Panel D illustrates peroxisomal bile acid synthesis and β-oxidation of branched

chain fatty acids (BCFA). Bile acid intermediates (BAI) and BCFA are transported into

peroxisomes via the ABCD proteins. CA = cholic acid, CDCA = chenodeoxycholic acid, BrAcyl-

CoA = branched chain acyl-CoA. For panels C-D, the enzymes catalyzing each step are shown

in boxes, and enzymes that were significantly less abundant in the mitochondria/peroxisome-

enriched fraction from TMEM135 knockdown mice (Panel B) are shaded. Panel E displays relative

peroxisome content in the mitochondria/peroxisome-enriched fractions as determined by

proteomic and Western blot analysis. For proteomics, only fed mice were analyzed (n = 3). For

Western blot, the ABCD3:COX4I1 ratio in the mitochondria/peroxisome-enriched fraction is

shown (n = 5). Panel F is subcellular distribution of ACAA1, ACOX1, SCP2, and CAT. ACOX1

migrates in 2 separate bands in Western blot, with the smaller band being proteolytically

processed ACOX1 (Mizuno et al., 2013). Furthermore, SCP2 (also known as SCPx) is a 58 kDa

protein that contains a 45 kDa 3-ketoacyl CoA thiolase domain and a 13 kDa sterol-carrier domain

(Stolowich et al., 2002), which are separated by proteolytic cleavage (Mizuno et al., 2013). The

antibody used in the current study recognizes the full-length 58 kDa protein (indicated by arrow)

and the 45 kDa 3-ketoacyl CoA thiolase domain, but not the 13 kDa sterol-carrier domain. The

identity of the other bands in the SCP2 blot is not known.

Figure 7: Reduced peroxisomal import of matrix enzymes impairs β-oxidation. Panel A

contains images from Western blot analysis of whole cell and mitochondria/peroxisome-enriched

fractions (Mito/Perox-Enriched Fraction) in livers of fed and fasted mice. Loading controls were

TUBB for whole cell lysates, COX4I1 for mitochondria, and ABCD3 for peroxisomes. The siRNA

treatment is indicated beneath the lanes, Con = control siRNA, KD = Knockdown, *Tmem135*

siRNA. For SCP2, the band used for densitometry analysis is indicated by asterisks. Panel B

displays the results of densitometry analysis of the blots shown in Panel A. Peroxisome import

was determined by first normalizing each protein in the mitochondria/peroxisome-enriched

fraction to ABCD3 to account for differences in peroxisome content, and the ratio relative to the

whole cell lysate (normalized to TUBB) was used to determine peroxisome import. Panel C plots

concentrations of C18 fatty acids quantified by GC. In panels B-C, asterisks denote significant

difference ($p < 0.05$) within feeding status due to *Tmem135* siRNA, and # indicates significant

difference due to fasting in mice receiving the control siRNA (n = 5). Panel D displays the mean

± SEM for all fatty acids (µg fatty acid per mg protein) quantified by GC. In the P column, asterisks

denote significant difference ($p < 0.05$) and ^ denotes trends ($0.05 < p < 0.1$) within feeding status

due to *Tmem135* siRNA. In the fasted mice control siRNA column, # indicates significant ($p < 0.05$)

difference due to fasting in control siRNA mice. See also Figure S5.

STAR Methods

CONTACT FOR REAGENT AND RESOURCE SHARING

Further information and requests for resources and reagents should be directed to and will be fulfilled by the Lead Contact, Randy L. Bogan (boganr@email.arizona.edu). Transfer of resources is subject to the acceptance of a Material Transfer Agreement as required by policy at the University of Arizona. Resources are transferred and distributed by several means according to the needs of the end-user. For other academic and non-profit researchers, the resources are typically transferred under standard Material Transfer Agreements. For requests by for-profit organizations, the resources are made available under standard non-exclusive license.

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Cell Lines

All cell lines were obtained from American Type Culture Collection (ATCC) and were cultured at 37°C, 5% CO_2 in air, in a humidified incubator. Human cells (all derived from males) included the hepatocyte lines HepG2 (ATCC® HB-8065™) and Hep 3B2.1-7 [Hep3B] (ATCC® HB-8064™), and the monocyte line THP-1 (ATCC® TIB-202™). Murine cells included the hepatocyte line (embryonic-derived) BNL 1NG A.2 (ATCC® TIB-76™) and the macrophage line (male) RAW 264.7 (ATCC® TIB-71™). Growth media for all cell lines except THP-1 was DMEM/F12 (Sigma Aldrich, Inc.) supplemented with Pen/Strep (100 units/ml penicillin and 100 μg/ml streptomycin) and 10% fetal bovine serum (FBS). THP-1 cells were incubated in RPMI media (Sigma) supplemented with 2-mercaptoethanol (0.05 mM), Pen/Strep, and 10% FBS. THP-1 cells were differentiated into macrophages by treatment with 12-myristate-13-acetate (PMA, 100 ng/ml) for 5 days prior to experimentation.

Mice

All procedures involving mice were approved by the University of Arizona IACUC. Male C57BL/6 mice, 12 weeks of age, were used. Mice were group housed until initiation of experimental procedures, at which point they were switched to individual caging.

METHOD DETAILS

 Cell Treatments

T0901317 (Cayman Chemical, Inc.) was dissolved in DMSO and used at a 1 μM

concentration unless indicated otherwise. Cycloheximide (ACROS Organics, Inc.) was dissolved

in DMSO and used at a final concentration of 50 μg/ml. The final DMSO concentration was held

constant in all groups at less than 0.2% (*v:v*). For siRNA knockdown experiments, Silencer® select

pre-designed siRNAs (Life Technologies, Inc.) against human *NR1H2* (s14684), *NR1H3*

(s19568), *TMEM135* (s35201), and a negative control siRNA (catalog 4390843) were purchased

and transfected into HepG2 cells using Lipofectamine® RNAiMAX (Life Technologies) or TransIT-

siQuest® reagent (Mirus Bio, LLC.) according to manufacturer instructions. All experiments were

repeated 4-5 times on different days or with different passages of cells.

HepG2 Mitochondria/Peroxisome-Enriched Fraction Isolation

Cells were harvested and washed three times in 1 ml cell resuspension buffer (1 mM Tris-

HCl, pH 7.4, 130 mM NaCl, 5 mM KCl, 7.5 mM $MgCl_2$). Cells were then resuspended in 600 μl

homogenization buffer (10 mM Tris-HCl, pH 7.4, 10 mM KCl, 0.15 mM $MgCl_2$, plus protease and

phosphatase inhibitor cocktail), incubated on ice for 10 min, and homogenized for 30 strokes in a

glass dounce homogenizer with a tight-fitting pestle. The homogenate was mixed with 100 μl of

2M sucrose and centrifuged at 1,200 × g for 5 min at 4° C, and the supernatant transferred to a

new tube. This low speed centrifugation step was repeated two more times. Mitochondria and

peroxisomes were pelleted by centrifugation at 7,000 × g for 10 min at 4° C and the supernatant

removed. The pellet was washed once with 300 μl of wash buffer (10 mM Tris-HCl, pH 7.4, 0.15

mM $MgCl_2$, 250 mM sucrose), and centrifuged at 9,500 × g for 5 min. Pellets were solubilized in

RIPA buffer containing protease and phosphatase inhibitors, and protein concentration was

determined by BCA assay (Fisher Scientific) following manufacturer recommendations.

Electrophoretic Mobility-Shift Analysis (EMSA)

The DNA sequences for human *NR1H2*, *NR1H3*, and RXR α (*RXRA*) were synthesized

by Life Technologies and cloned into the pTarget™ mammalian expression vector (Promega

Corporation) using XhoI and KpnI restriction enzyme sites. Inserts were confirmed by DNA

sequence analysis. Recombinant NR1H2, NR1H3, and RXRA proteins were produced using the

TNT® Quick Coupled Transcription/Translation System (Promega). Fluorescent probes were

synthesized by Integrated DNA Technologies and corresponded to the LXRE1 sequence with

IRDye 700 attached to the 5' end, and probes corresponding to the LXRE3 sequence were

labelled with IRDye 800. Additionally, unlabeled DNA oligos corresponding to LXREs 1-3, as well

as mutated LXREs 1-3, were synthesized (Figure S1B). Single stranded DNA oligos were

annealed by mixing sense and antisense oligos in a 1:1 molar ratio, heating to 100°C for 5

minutes, and slowly cooling to room temperature. The EMSA binding reactions were carried out

for 30 min at room temperature in 10 mM Tris, 50 mM KCl, pH 7.5; 3.5 mM DTT, 0.25% Tween-

20, 1 μg Poly (dI.dC), 5 nM fluorescent probe (1 nM each for LXRE1/LXRE3 competition

experiment), and 1.5 μl of TNT lysate per protein (non-induced TNT lysate substituted in control

reactions). Unlabeled competitor DNA was added prior to the addition of fluorescent probe at a

200-fold molar excess unless indicated otherwise. Following the incubation period, EMSA

reactions were resolved on a 6% TBE gel (Life Technologies). Imaging and densitometry were

performed with a Li-COR Odyssey CLx and Image Studio version 3.1 software, respectively.

Reporter Assays

The region from -2662 to -1 bp (relative to translation start site) of the human *TMEM135*

promoter was synthesized by Life Technologies. Additionally, the same region of the human

TMEM135 promoter was synthesized except that point mutations were introduced into each of

the three LXREs (Figure S1B). The normal and mutant *TMEM135* promoters were cloned into the

pGL4.17 vector (Promega) using SacI and XhoI restriction enzyme sites, which expresses firefly

luciferase from a promoter cloned into the vector. A unique PacI site was identified in the region

between LXRE1 and LXRE2, and a unique EcoRI site was identified in the region between LXRE2

and LXRE3. These restriction sites were used to generate six more *TMEM135* promoter/pGL4.17

constructs so that all possible combinations of wild type and mutated LXREs were obtained. All

constructs were verified using DNA sequence analysis. These reporter vectors were transfected

into HepG2 cells along with a β-galactosidase control plasmid (Promega) to normalize

transfection efficiency. In some transfections, *NR1H3*/pTarget and *RXRA*/pTarget constructs

were co-transfected. Vectors were transfected in a ratio of 60:20:10:10 (luciferase:β-

galactosidase:NR1H3:RXRA) using Lipofectamine$^®$ 3000 according to manufacturer

recommendations, with no-insert plasmids substituted in control transfections to keep DNA

concentrations constant. Each vector combination was transfected in triplicate to HepG2 cells in

96 well plates, and cells were incubated with DNA/Lipofectamine complexes for 24 hours at which

point media were changed and cells were treated for an additional 48 hours with DMSO or T09.

Cell lysates were prepared using mammalian protein extraction reagent (Fisher), and the lysate

was fractionated to quantify luciferase and β-galactosidase activity using luminescent detection

kits (Fisher). The ratio of luciferase to β-galactosidase activity was calculated, and the fold-change

in T09 vs DMSO treated cells was determined for each transfection.

<u>Chromatin Immunoprecipitation (ChIP) Analysis</u>

HepG2 cells were fixed for 10 min at room temperature with 1% formaldehyde. The

reaction was quenched by adding glycine to a 125 mM final concentration and incubating 5 min

at room temperature. The cells were trypsinized, harvested, and washed twice with PBS. Nuclei

were isolated by incubating cell pellets in 500 µl of a hypotonic buffer for 15 min on ice (20 mM

Tris-HCL, pH 7.4, 10 mM NaCl, 3 mM MgCL$_2$, with protease and phosphatase inhibitor cocktail

from Promega). Triton X-100 detergent was added to a 0.5% *v:v* final concentration to lyse the

cells followed by centrifugation at 1200 × g for 10 min to pellet nuclei. The supernatant was

removed, and the nuclei pellet washed once with 500 µl hypotonic buffer. The pellet was

resuspended in 500 µl of RIPA buffer with protease and phosphatase inhibitors, and nuclei were

sonicated in an ice-water bath using a Branson 250 digital sonifier programmed to cycle at a 10%

amplitude with 5 seconds on followed by 20 seconds off, and a 3-minute total sonication time that

was determined in preliminary time-course studies to produce chromatin of optimal length

(approximately 200-1000 base pairs). The sonicated lysates were centrifuged at 8,000 × g for 1

minute and the supernatant recovered. An aliquot of cleared lysate was treated with 25 µg RNAse

A and 50 µg proteinase K in a buffer containing 500 mM NaCl and heated at 65°C for 1.5 hours

to reverse cross-links. The DNA concentration in this aliquot was determined using a fluorometric

assay (Life Technologies) following manufacturer recommendations. The quantity of DNA used

in each ChIP was held constant within replicates (8 to 10 µg), and 10% of the input amount was

saved. Chromatin was diluted to a 500 µl final volume in binding/wash buffer (50 mM Tris, 150

mM NaCl, 1% Triton X-100) containing protease and phosphatase inhibitors, and 3 µg of antibody

was added per reaction. Tubes were incubated on an end-over-end rotator at 4°C overnight.

Antibodies used for ChIP were mouse monoclonal IgG2a including: 1) anti-NR1H3 (clone

PPZ0412, R&D Systems catalog PP-PPZ0412-00), 2) anti NR1H2 (clone K8917, R&D Systems

catalog PP-K8917-00), and 3) non-specific control IgG2a (clone MOPC-173, Abcam catalog

ab18413). The following day 25 µl of Protein G-conjugated Dynabeads (Life Technologies) were

added and incubated for 1 hour at 4°C. The beads were recovered using a magnet and washed

three times at 4°C for 5 minutes each wash with 1 ml of binding/wash buffer. The beads were

then washed once in final wash buffer (50mM Tris, 500 mM NaCl, 1% Triton X-100). Elutions

were performed with 130 µl elution buffer (100mM NaHCO$_3$, 500mM NaCl, 1% SDS) containing

25 µg RNAse A and 50 µg proteinase K, and ChIP reactions and input controls were incubated

on a thermal mixer at 65°C for 1.5 hours. The supernatant was recovered, and DNA purified using

PCR purification columns (Life Technologies) following manufacturer recommendations. Purified

DNA was analyzed by QPCR in a duplex reaction with primers and a Taqman probe specific for

the region spanning LXRE3 on the *TMEM135* gene, and a non-specific locus was simultaneously

quantified in the same reaction as a control for non-specific DNA carryover (Table S1). Serial

dilutions of purified sonicated DNA were included to verify linear amplification and PCR efficiency,

and Ct values from ChIP reactions and input controls were used to calculate the percentage of

DNA recovered relative to the input. Data were normalized by calculating the ratio of the

TMEM135 LXRE3 to the non-specific chromatin locus.

<u>Semi-Quantitative Real-Time PCR</u>

Cells or tissue were homogenized in Trizol reagent for isolation of mRNA and were further

purified over RNeasy columns according to manufacturer recommendations (Life Technologies).

Concentrations of RNA were quantified by spectrophotometry, and 0.2-1 μg RNA was treated with

DNAse I and reverse transcribed into cDNA using the High Capacity cDNA Reverse Transcription

Kit (Life Technologies). Following reverse transcription, cDNA were diluted to 10 ng/μl (based on

original RNA concentration) and used for semi-quantitative real-time PCR analysis (QPCR).

Primer and probe sequences used for QPCR are listed in Table S1. The QPCR assays were

performed on a StepOne Plus Real-Time PCR system (Life Technologies) using TaqMan® MGB

probe or SYBR green detection. Relative mRNA abundance was determined by extrapolation of

threshold (Ct) values from a standard curve of serial cDNA dilutions and normalized to the

housekeeping gene mitochondrial ribosomal protein S10 (*MRPS10*).

<u>Cell Proliferation Assay</u>

HepG2 cells were plated at 15,000 cells per well in 96-well plates and transfected with

siRNAs immediately. At the appropriate post-transfection time, viable cells were quantified using

a CyQuant® Direct Cell Proliferation assay (Life Technologies) according to manufacturer

recommendations, and fluorescence (excitation 495 nm, emission 527 nm) determined using a

Synergy H1 plate reader (BioTek Instruments, Inc.) in area scanning mode to account for uneven

cell distribution. The number of cells per well was estimated by extrapolation from a standard

curve of increasing cell numbers that was generated at the time of plating.

<u>Flow Cytometry Analysis of Cell Cycle</u>

Freshly harvested HepG2 cell pellets were fixed by slow addition of 1 ml ice-cold 70%

ethanol while vortexing and incubated at -20°C overnight. Fixed cells were harvested by

centrifugation at 850 × g for 5 min and resuspended in 1 ml DNA staining buffer (100 μg/ml RNAse

A, 50 µg/ml propidium iodide, in PBS). Cells were incubated for 30 min at 37°C, placed on ice,

and analyzed within one hour. A Becton Dickinson FACSCANTO II flow cytometer equipped with

a 488 nm argon ion laser and a 585/42 bandpass filter was used for detection of bound propidium

iodide. List mode data files consisting of 10,000 events were acquired and the percentage of cells

in G0/G1, S, and G2/M stages was determined with CellQuest PRO software (BD Biosciences).

<u>HepG2 ATP Quantification</u>

At 48 hours post-siRNA transfection, HepG2 cells were switched to serum-free, glucose-

free, sodium pyruvate-free DMEM media and incubated at 37°C for 4 hours. Cellular ATP

concentrations were determined using a homogenous luminescent ATP detection kit (Abcam

catalog ab113849). Additional HepG2 cells treated in parallel to those used for ATP determination

were lysed in RIPA buffer and a BCA protein assay (Fisher) was performed, and ATP content

was normalized to protein concentrations.

<u>Triglyceride Quantification</u>

HepG2 cells or frozen mouse liver were sonicated in PBS containing protease and

phosphatase inhibitors (Promega) in an ice-water bath using a Branson 250 digital sonifier

programmed to cycle at a 10% amplitude with 5 seconds on followed by 20 seconds off, and a 30

second total sonication time. Triglycerides were extracted using an organic solvent procedure

optimized for triglycerides (Schwartz and Wolins, 2007). Lysates were diluted 1:1 with deionized

water, and 400 µl of diluted lysate was combined in a glass tube with 1.8 ml of a 4:1

isopropanol:hexane mixture. Samples were vortexed for 1 minute and incubated at room

temperature for 30 min. Next, 500 µl of a 1:1 hexane:diethyl ether solution was added, the

samples were vortexed as before, and incubated at room temperature for 1 minute. One ml of

deionized water was added, samples were vortexed as before, and incubated at room

temperature for 20 minutes to allow phase separation. The organic phase was transferred to a

new glass tube and dried under a gentle stream of nitrogen in a water bath at 30°C. Dried extracts

were dissolved in 200 µl assay buffer (50 mM potassium phosphate, pH 7.0, 0.01% Triton X-100)

by 5 total cycles of vortexing for 1 minute and incubating at room temperature for 3 minutes.

Extracts (50 µl/well) were transferred to a clear 96 well plate, and 250 µl of triglyceride detection

reagent (Pointe Scientific, Inc., catalog T7532) was added. Triglyceride standards (Pointe

Scientific) diluted in assay buffer were also run with each assay. Plates were incubated at 37°C

for 30 minutes with vigorous shaking, and absorbance at 500 nm was determined with a Synergy

H1 plate reader. The protein concentration in sonicated lysates was determined using a BCA

protein assay, and triglyceride content was normalized to protein content.

<u>In Vivo siRNA Delivery and Tissue Harvest</u>

Ambion® In Vivo pre-designed siRNA (Life Technologies) against *Tmem135* (s91285) or

in vivo negative control #1 siRNA (catalog 4459405) were complexed with Invivofectamine® 3.0

reagent (Life Technologies) according to manufacturer recommendations. The siRNA complexes

were delivered via a 200 µl tail-vein injection at a final siRNA concentration of 1 mg/kg body

weight. Mice were weighed prior to siRNA injection and again before euthanization. Mice were

euthanized 4 days after siRNA injection at 4 hours after lights on, with or without a preceding 12

hour fast. Serum, liver, heart, adipose, and skeletal muscle were collected and snap frozen.

Tissues were powdered while frozen and stored at -80° C until used.

<u>Serum Analytes</u>

Total cholesterol, HDL cholesterol, and triglyceride concentrations in serum were

determined as we have described previously (Jensen et al., 2017). Briefly, the HDL fraction of

serum was isolated by mixing serum 1:1 with 15% polyethylene glycol, centrifuging at 2,000 × g

for 10 minutes at room temperature, and harvesting the HDL-containing supernatant. The

cholesterol concentrations in whole serum and the HDL fraction were determined by a

fluorometric enzyme assay consisting of cholesterol esterase (0.3 units/ml), cholesterol oxidase

(1 unit/ml), horse radish peroxidase (1 unit/ml), and Amplex UltraRed® (50 µM, Life Technologies)

in assay buffer (0.1 M potassium phosphate, pH 7.0, 50 mM NaCl, 5 mM cholic acid, 0.1% Triton

X-100). The cholesterol assay was incubated at 37°C for 30 minutes, and fluorescence measured

in a Synergy H1 plate reader at 555 nm excitation and 585 nm emission wavelengths. Triglyceride

concentrations in whole serum were determined using the same assay described earlier for

HepG2 and mouse liver samples. Commercial enzymatic assays were used to quantify serum

concentrations of β-hydroxybutyrate (BHB) (Cayman Chemical, catalog 700190), non-esterified

fatty acids (NEFA) (Catachem Inc., catalog C514-0A), glucose (Pointe Scientific, Inc., catalog

G7521), and a commercial ELISA was used to quantify serum insulin (Mercodia AB, catalog 10-

1247) according to manufacturer recommendations.

<u>Liver Analytes</u>

Commercial kits were used for the extraction and quantification of liver NAD+ and NADH

(Abcam Inc., catalog ab176723), and glycogen (BioAssay Systems, catalog E2GN), according to

manufacturer recommendations. Concentrations of ATP and NEFAs in liver were determined as

we have described elsewhere (Geisler et al., 2016). Briefly, for ATP determination frozen

powdered liver was homogenized in ATP releasing agent (Sigma Chemical Co., catalog FLSAR).

ATP was quantified with a luminescent ATP Determination Kit (Life Technologies, catalog

A22066) and normalized to tissue weight. For NEFAs, powdered liver was sonicated in PBS as

described earlier, and 100 μl lysate was mixed with 1 ml of 100% ethanol. Samples were vortexed

for 10 minutes, centrifuged at 16,000 × g for 5 minutes, and the supernatant recovered. A

commercial colorimetric assay for NEFAs (Catachem Inc., catalog C514-0A) was used and

standards were diluted in ethanol, and NEFA concentrations were expressed on an equal protein

basis.

To quantify ketones (BHB and acetoacetate), powdered liver was sonicated in PBS,

acidified with 10% (*w:v*) trichloroacetic acid, centrifuged to remove proteins, and the supernatant

was recovered and neutralized. Concentrations of BHB in the extract were determined using the

same enzymatic assay described for serum. Concentrations of acetoacetate were determined

using a modified assay based on methods described previously (Salway, 1969; Walker, 1954).

Briefly, samples and standards were divided with one portion subjected to boiling (100° C for 5

minutes) to degrade acetoacetate, with the remaining portion kept on ice. A three-fold volume of

freshly prepared diazo reagent (Walker, 1954) was then added to all samples and incubated at

room temperature in the dark for 30 minutes. The reaction was stopped by adding a 20% volume

of 2N NaOH, and absorbance was measured at 480 nm. The absorbance from the boiled sample

was subtracted from the respective non-boiled sample, and acetoacetate concentrations were

calculated from the resultant standard curve. Concentrations of cholic acid in liver were quantified

using a competitive enzyme immunoassay according to manufacturer recommendations (Cell

Biolabs, Inc., catalog MET-5007).

Liver Subcellular Fractionation

Powdered liver (approximately 50 mg) was homogenized for 40 strokes in a glass dounce

homogenizer with a tight-fitting pestle using 600 µl of cold fractionation buffer (10 mM Tris-HCl,

pH 7.4, 1 mM EGTA, 200 mM sucrose, protease and phosphatase inhibitor cocktail) on ice. The

homogenate was transferred to a new tube, and the dounce homogenizer was washed with an

additional 600 µl of fractionation buffer and transferred to the same tube. The homogenate was

centrifuged at 600 × g for 10 minutes at 4°C to pellet unbroken cells and nuclei, and the

supernatant transferred to a new tube. This low speed spin was repeated once more.

Mitochondria and peroxisomes were pelleted by centrifugation at 7,000 × g for 10 minutes, and

the supernatant (cytosolic fraction) was transferred to a new tube. The pellet was washed with

300 µl fractionation buffer and pelleted as before. The pellet was resuspended in 500 µl of

fractionation buffer and a 30 µl aliquot was pelleted and solubilized in RIPA buffer for a BCA

protein assay. Following protein assay mitochondria/peroxisome fractions were partitioned into

30 µg (protein content) aliquots followed by centrifugation and removal of the supernatant, and

the pellets were frozen at -80°C.

Proteomic Analyses

The mitochondria/peroxisome-enriched fraction from a subset of fed control and

TMEM135 knockdown mice was solubilized in RIPA buffer, proteins were precipitated with

acetone, and proteins were sent to the Arizona Proteomics Consortium for mass spectrometry

identification of proteins. Proteins were digested with trypsin and equal quantities of protein (500

ng) were loaded. Samples were analyzed on a Thermo Q Exactive Plus Orbitrap mass

spectrometer. Peptide identification in the resultant tandem mass spectra was performed using

Proteome Discoverer Software version 1.3.0.339 scanning with the SEQUEST algorithm against

the mouse proteome database (Mouse_unitprotkb_proteome_2016_0720_cont.fasta). Data

analysis was performed using Scaffold version 4.8.1, with a peptide identification threshold of

95.0% and a minimum protein identification threshold of 99.9% and 2 unique peptides.

<u>Western Blot</u>

All primary antibodies were obtained from Abcam unless otherwise indicated. Catalog

numbers and final concentrations or dilutions of each antibody were: rabbit polyclonal anti-ACAA1

(catalog ab154091, 1 μg/ml), rabbit monoclonal anti-ACOX1 (catalog ab184032, 1:2500 dilution),

rabbit monoclonal anti-CAT (catalog ab209211, 1:2000 dilution), rabbit monoclonal anti-SCP2

(catalog ab140126, 1:2500 dilution), rabbit polyclonal anti-TMEM135 (catalog ab167096, 1

μg/ml), rabbit monoclonal anti-COX4I1 (Li-COR Inc., catalog 926-42214, 1:1000 dilution), mouse

monoclonal anti-COX4I1 (catalog 14744, 0.25 μg/ml), mouse monoclonal anti-TUBB (Sigma

Aldrich catalog T8328, 0.5 μg/ml), and rabbit polyclonal anti-PMP70 (catalog ab3421, 0.5 μg/ml).

Proteins (12.5 μg/well) were resolved on 4-12% Bis-Tris gels (Life Technologies) and

subsequently transferred to nitrocellulose membranes. The membranes were blocked with 5%

non-fat dry milk (NFDM) (*w:v*) in TBS with 0.1% (*v:v*) Tween (TBST) for 1 hour at room temp. All

primary antibodies were diluted in TBST with 1% NFDM and incubated with the membranes on a

rocking platform at 4° C overnight. The membranes were washed 4x with TBST, and IRDye®

680RD or 800CW-conjugated secondary antibodies (LI-COR Biosciences, Inc.) were diluted

1:4000 in TBST + 1% NFDM and incubated with membranes for 1 hour at room temp. The

membranes were washed as before. Imaging and densitometry were performed with a Li-COR

Odyssey CLx and Image Studio version 3.1 software, respectively.

Gas Chromatography (GC)

1019

Lysates were prepared by sonicating frozen powdered liver in PBS as described earlier.

Insoluble material was pelleted by centrifugation at 600 × g for 5 minutes. A BCA protein assay

was performed on the cleared lysate. A one-step transesterification reaction was performed

(Masood et al., 2005). Briefly, 1.7 ml of acidic methanol (1.6 ml methanol and 100 µl acetyl

chloride) was added to borosilicate glass tubes with Teflon-lined caps. Butylated hydroxytoluene

(BHT, 50 µg/ml) was added to methanol to prevent fatty acid oxidation. Tridecanoic acid (13:0,

Cayman Chemical) was used as an internal standard because preliminary studies determined

that it was not present in liver lysates. Tridecanoic acid was diluted in methanol/BHT to 100 µg/ml,

and 100 µl was added to all samples and standards. Authentic standards for the following fatty

acids were obtained from Cayman Chemical: 12:0, 14:0, 16:0, 16:1, 18:0, 18:1 cis(n9), 18:2

cis(n6), 20:3 cis(n6), 20:4 (n6), 22:0, 22:6 (n3), 24:0, 24:1, and 26:0. Pooled mixtures containing

increasing concentrations of these fatty acids were prepared and dissolved in chloroform. 100 µl

of authentic standards were added to standard tubes, and 100 µl of chloroform was added to

unknown tubes. Next, 100 µl PBS was added to standard tubes, and 800 µg (protein content) of

lysate diluted to a 100 µl volume with PBS was added to unknown tubes. The tubes were tightly

capped and vortexed for 1 minute, and then incubated in a heating block at 100°C for one hour.

The tubes were cooled to room temperature, 750 µl hexane was added and the tubes were

vortexed for 30 seconds, and the upper organic layer was transferred to a GC sample vial. The

hexane extraction was repeated once more. Hexane extracts were evaporated under a gentle

stream of nitrogen to dryness and dissolved in 50 µl hexane.

GC was performed on an Agilent Technologies 6890N GC equipped with a Varian CP-Sil

88 column for fatty acid methyl ester analysis (100 m × 0.25 mm inner diameter × 0.2 µm film

thickness) and a flame ionization detector. Inlet temperature was 250° C, and 1 µl of each

standard or unknown was manually injected at a split ratio of 7.5:1. Helium was used as carrier

gas at a 1.0 ml/minute constant flow. The oven was programmed for an initial temperature of 80°

C with a 4° C/min ramp to 220° C, a 5-minute hold, then a 4° C/min ramp to 240° C followed by a

10-minute hold. The detector was set at 270°C using air (450 ml/minute) and hydrogen (40

ml/minute), and nitrogen was used as make-up gas (10 ml/minute constant flow). The sampling

frequency was 20 Hz.

OpenLAB CDS ChemStation Edition software version C.01.06 (Agilent Technologies) was

used to analyze GC data. Retention times of authentic standards were matched with

corresponding peaks in unknown samples. For each fatty acid, the peak area for standards and

unknowns was divided by the internal standard peak area. The normalized peak area from

increasing concentrations of authentic standards was plotted, and the R^2 for the resultant standard

curves were > 0.995 in all cases. All fatty acids except 12:0 and 26:0 were identified in the

unknowns, and fatty acid concentrations were determined by extrapolating the normalized peak

area from the respective standard curve. Concentrations were expressed on an equal protein

basis.

QUANTIFICATION AND STATISTICAL ANALYSIS

Statistical analyses were performed using Stata Version 14 and all differences were

considered significant at $p < 0.05$. Information on the sample size, mean and error estimates, and

indicators of statistical significance are presented in the figures and their corresponding legends.

For immortalized cell lines, the total sample size (n) corresponds to cells on a different passage

number and/or experiments performed on different days. Time course data in immortalized cell

lines were analyzed by mixed effects regression analysis to account for repeated measures.

Treatments were coded 0 or 1, and hours post-treatment and treatment × time interaction were

included as fixed effects in the regression model with experimental replicate as the random effect.

Differences between treatments at each timepoint were determined from the treatment × time

interaction coefficient. For all other comparisons involving more than 2 groups, mixed effects

regression analysis was performed using treatment as the fixed effect and experimental replicate

as the random effect to account for repeated measures. Pairwise differences between treatments

were then determined using the Bonferroni multiple comparison method to control for the type 1

error rate. For comparisons involving only 2 groups, a paired t-test was performed.

For all mice data, the sample size refers to the number of mice. Regression analysis was

used for mice outcomes except where indicated otherwise. Samples were coded 0 or 1 for fasting,

TMEM135 knockdown fed mice, and TMEM135 knockdown fasted mice. The regression model

utilized fasting, TMEM135 knockdown in the fed state, and TMEM135 knockdown in the fasted

state, as predictors. The fasting coefficient indicated the difference due to fasting in control mice

only, while the TMEM135 knockdown coefficients indicated the difference between control mice

and TMEM135 knockdown mice within the fed state or fasted state. Differences were considered

significant at $p < 0.05$, with p values ≥ 0.05 and < 0.10 considered trends. Only fed mice were

used in proteomic analysis. Because the proteomic data does not follow a normal distribution due

to some proteins not being detected in all samples, significant differences in proteomic data were

determined from weighted spectra using the non-parametric Fisher's Exact test ($p < 0.05$)

contained within the Scaffold version 4.8.1 software. The list of proteins that were significantly

reduced in knockdown livers was further trimmed by hand to exclude proteins that were not

detected in all the samples from mice that received the control siRNA.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

B

| Protein Symbol | Weighted Spectral Counts ± SEM | |
	Control siRNA	Tmem135 siRNA
CPS1	110 ± 16.3	84.7 ± 34.1
CAT	125 ± 7.6	76 ± 14
SCP2	128 ± 13.1	69.7 ± 35.3
HSD17B4	66 ± 8.5	46.7 ± 9.5
CES1	61.3 ± 11.8	43.3 ± 7.2
NUDT7	28 ± 8.2	17 ± 17
ACOX1	27 ± 6.7	13 ± 10.5
PC	24.3 ± 3.8	15.7 ± 4.7
ACOX2	22 ± 4.4	13.3 ± 11.9
CES3	17.7 ± 3.7	10.7 ± 6.1
GLUD1	21 ± 2.3	13 ± 3.1
DHRS4	23.3 ± 3.4	11.3 ± 6.4
HACL1	20.7 ± 3.7	9.7 ± 5.7
COMT	17.7 ± 5.8	8.3 ± 8.3
MTTP	15 ± 4.7	7.7 ± 4.3
PSAP	14 ± 4.5	5.3 ± 3.2
ACAA1	35 ± 6.5	9.3 ± 9.3
ALDH3A2	11 ± 2.9	5.3 ± 3.5
ALDH6A1	11 ± 2.9	5.7 ± 3.5
AMACR	10 ± 2.6	4 ± 4
SUCLA2	9 ± 1.7	3.3 ± 2.4
ACADL	5 ± 0.6	1.7 ± 0.9
SLCO1A1	3.7 ± 1.2	1 ± 1

Figure 6

Figure 7

Fatty Acids						
	Fed Mice			**Fasted Mice**		
	Control siRNA	*Tmem135 siRNA*	*P*	*Control siRNA*	*Tmem135 siRNA*	*P*
14:0	0.32 ± 0.06	0.29 ± 0.03		0.73 ± 0.12 #	0.85 ± 0.18	
16:0	48.9 ± 3.6	45.7 ± 2.2		70.2 ± 4.4 #	83.1 ± 7.5	^
16:1	2.4 ± 0.5	2.2 ± 0.2		3.5 ± 0.7	5.3 ± 1.0	^
18:0	31.3 ± 1.3	29.4 ± 1.2		31.3 ± 1.6	33.2 ± 0.4	
18:1	22.9 ± 3.4	22.0 ± 1.9		41.1 ± 4.8 #	56.7 ± 9.9	^
18:2	27.1 ± 2.0	28.5 ± 1.9		58.1 ± 5.1 #	78.0 ± 10.0	*
20:3	2.2 ± 0.2	2.1 ± 0.2		1.1 ± 0.1 #	1.6 ± 0.1	^
20:4	21.0 ± 1.6	19.2 ± 1.4		20.8 ± 1.1	23.4 ± 0.7	
22:0	1.7 ± 0.1	1.4 ± 0.1	*	1.4 ± 0.1	1.6 ± 0.1	
22:6	13.0 ± 1.2	11.7 ± 1.0		17.1 ± 1.1 #	19.0 ± 0.9	
24:0	1.01 ± 0.03	0.83 ± 0.05	^	0.95 ± 0.08	1.01 ± 0.05	
24:1	0.87 ± 0.04	0.87 ± 0.05		1.0 ± 0.1	1.19 ± 0.06	^
Total	172.5 ± 12.9	164.1 ± 7.7		247.4 ± 14.2 #	304.8 ± 28.3	*