Disentangling the effects of genetic architecture, mutational bias and selection on evolutionary forecasting

Peter A. Lind [1,2], Eric Libby [1,3,4], Jenny Herzog[1] and Paul B. Rainey[1,5,6]

[1]New Zealand Institute for Advanced Study, Massey University at Albany, Auckland, 0745, New Zealand.

[2]Department of Molecular Biology, Umeå University, SE-901 87 Umeå, Sweden.

[3]Santa Fe Institute, Santa Fe, New Mexico, United States of America.

[4]Department of Mathematics, Umeå University, SE-901 87 Umeå, Sweden.

[5]Department of Microbial Population Biology, Max Planck Institute for Evolutionary Biology, Plön 24306, Germany.

[6]Ecole Supérieure de Physique et de Chimie Industrielles de la Ville de Paris (ESPCI Paris-Tech), CNRS UMR 8231, PSL Research University, 75231 Paris, France.

Correspondence and requests for materials should be addressed to

Peter A. Lind, Dept. Molecular Biology, Umeå University, SE-901 87 Umeå, Sweden.

email: peter.lind@umu.se

Keywords: Pseudomonas fluorescens, experimental evolution, genetic architecture, wrinkly spreader, c-di-GMP, evolutionary forecasting

Short running title: complexities of the genotype-to-phenotype map

Impact statement: Using a combination of genetics, experimental evolution and mathematical modelling this work defines information necessary to predict the outcome of short-term adaptive evolution

Abstract

Predicting evolutionary change poses numerous challenges. Here we take advantage

of the model bacterium *Pseudomonas fluorescens* in which the genotype-to-

phenotype map determining evolution of the adaptive "wrinkly spreader" (WS) type

is known. We present mathematical descriptions of three necessary regulatory

pathways and use these to predict both the routes that evolution follows and the

expected mutational targets. To test predictions, mutation rates and targets were

determined for each pathway. Unanticipated mutational hotspots caused data to

depart from predictions but the new data were readily incorporated into refined

models. A mismatch was observed between the spectra of WS-causing mutations

obtained with and without selection due to low fitness of previously undetected WS-

causing mutations. Our findings contribute toward the development of mechanistic

models for forecasting evolution, highlight current limitations, and draw attention to

challenges in predicting locus-specific mutational biases and fitness effects.

Introduction

Adaptation requires the realization of beneficial mutations. As self-evident as this maybe, predicting the occurrence of beneficial mutations and their trajectories to improved fitness is fraught with challenges (Lässig, et al. 2017). Nonetheless progress has been made for phenotypically diverse asexual populations subject to strong selection. Effective approaches have drawn upon densely sampled sequence data and equilibrium models of molecular evolution to predict amino acid preferences at specific loci (Luksza and Lassig 2014). Predictive strategies have also been developed based on selection inferred from the shape of coalescent trees (Neher, et al. 2014). In both instances the models are coarse-grained and sidestep specific molecular and mutational details.

There is reason to by-pass molecular details: mutation, being a stochastic process, means that for the most part details are likely to be idiosyncratic and unpredictable. But an increasing number of investigations give reason to think otherwise – that adaptive molecular evolution might follow rules (Pigliucci 2010; Stern 2013; Laland, et al. 2015). This is particularly apparent in studies of parallel molecular evolution (Colosimo, et al. 2005; Woods, et al. 2006; Ostrowski, et al. 2008; Flowers, et al. 2009; Meyer, et al. 2012; Tenaillon, et al. 2012; Zhen, et al. 2012; Herron and Doebeli 2013; Galen, et al. 2015; Bailey, et al. 2017; Kram, et al. 2017; Stoltzfus and McCandlish 2017).

A standard starting position for predicting adaptive evolution recognises the importance of population genetic parameters including mutation rate, generation time, population size, selection and more recently information on the distribution of beneficial fitness effects, but these factors alone leave unconsidered mechanistic details that arise from the genotype-to-phenotype map and from mutational biases. To what extent do these details matter?

Mutations arise randomly with respect to utility, but genetic architecture can influence the translation of mutation into phenotypic variation: the likelihood that a given mutation generates phenotypic effects depends on the genotype-to-phenotype map (Alberch 1991; Gompel and Prud'homme 2009; Stern and Orgogozo 2009; Rainey, et

al. 2017). The function of gene products and their regulatory interactions thus

provides information on likely mutational targets underpinning particular phenotypes.

This is evident when considering a hypothetical structural determinant subject to both

positive and negative regulation and whose over-expression generates a given

adaptive phenotype. Assuming a uniform distribution of mutational events, mutations

in the negative regulator (and not the positive activator) will be the primary cause of

the adaptive phenotype. This follows from the fact that loss-of-function mutations are

more common than gain-of-function mutations. Indeed, an emerging rule indicates

that phenotypes determined by genetic pathways that are themselves subject to

negative regulation are most likely to arise by loss-of-function mutations in negative

regulatory components (McDonald, et al. 2009; Tenaillon, et al. 2012; Lind, et al.

2015; Fraebel, et al. 2017).

Mutation is not equally likely at each nucleotide of a given genome (Lind and

Andersson 2008; Lynch 2010; Seier, et al. 2011; Foster, et al. 2015; Reijns, et al.

2015; Sankar, et al. 2016; Stoltzfus and McCandlish 2017). Numerous instances of

mutational bias have been reported. Prime examples are simple sequence repeats

such as homopolymeric nucleotide tracts or di-, tri- and tetrameric repeats that mutate

at high frequency via slipped strand mispairing (Levinson and Gutman 1987). These

readily identifiable sequences define contingency loci in obligate human pathogens

and commensals (Moxon, et al. 1994) and are widespread in eukaryotic genomes

(Tautz and Renz 1984). The behaviour of contingency loci can be further modulated

by defects in components of methyl-directed mismatch repair systems (Richardson

and Stojiljkovic 2001; Martin, et al. 2004; Hammerschmidt, et al. 2014; Heilbron, et

al. 2014).

Certain palindromic structures also lead to mutational bias (Viswanathan, et al. 2000;

Lovett 2004) and promote amplification events including that increase mutational

target size (Roth, et al. 1996; Kugelberg, et al. 2010; Reams and Roth 2015),

transition-transversion bias (Stoltzfus and McCandlish 2017) and elevated mutation

rates at CpG sites (Galen, et al. 2015) can also skew the distributions of mutational

effects. Further bias arises from the chromosomal neighbourhood of genes under

selection (Steinrueck and Guet 2017), the location of genes with regard to interactions

with DNA replication/transcription machineries (Sankar, et al. 2016), and

environmental factors that affect not only mutation rate but also the spectra of

mutational events (Krasovec, et al. 2017; Maharjan and Ferenci 2017; Shewaramani,

et al. 2017).

Beyond the genotype-to-phenotype map and mutational biases, predicting adaptive

evolution requires ability to know *a priori* the fitness effects of specific mutations. At

the present time there is much theoretical and empirical interest in the distribution of

fitness effects (DFE) (Eyre-Walker and Keightley 2007) — and particularly the DFE

of beneficial mutations (Orr 2005) — because of implications for predicting the rate

of adaption and likelihood of parallel evolution (de Visser and Krug 2014), but

knowledge of the shape of the distribution is insufficient to connect specific mutations

to their specific effects, or to their likelihood of occurrence. Such connections require

a means of knowing the connection between mutations and their environment-specific

fitness effects. This is tall order. A starting point is to understand the relationship

between all possible mutational routes to a particular phenotype and the set that are

realised by selection.

Here we take a bacterial system in which the genetic pathways underpinning

evolution of the adaptive "wrinkly spreader" (WS) type are known and use this to

explore the current limits on evolutionary forecasting. *Pseudomonas fluorescens*

SBW25 growing in static broth microcosms rapidly depletes available oxygen

establishing selective conditions that favour mutants able to colonise the air-liquid

interface. The most successful mutant-class encompasses the WS types (Ferguson, et

al. 2013; Lind, et al. 2017b). These types arise from mutational activation of

diguanylate cyclases (DGCs), cause over-production of the second messenger c-di-

GMP (Goymer, et al. 2006; McDonald, et al. 2009), over-production of an acetylated

cellulose polymer (Spiers, et al. 2002; Spiers, et al. 2003) and ultimately formation of

a self-supporting microbial mat (Figure 1A).

Figure 1. The *Pseudomonas fluorescens* SBW25 wrinkly spreader model. (A)

Selection for access to oxygen allows wrinkly spreader (WS) mutants to invade the

ancestral smooth (SM) population in static microcosms. WS mutants form a mat at

the air-liquid interface through increased expression of the main structural

component, cellulose, encoded by the *wss* operon. Expression of cellulose is

controlled by the second messenger c-di-GMP, which is produced by diguanylate

cyclases (DGCs). Mutations in the *wsp*, *aws* and *mws* operons that activate their

respective DGCs (WspR, AwsR, MwsR) are the primary mutational pathways to WS.

(**B**) When a reporter construct connecting expression of the *wss* operon to resistance

to kanamycin is used under shaken non-selective conditions, WS mutants can be

isolated without the biasing influence of natural selection. This allows estimation of

the mutation rate to WS and an unbiased spectrum of mutations defining the

mutational target. Fitness can then be assayed in competition with a common

reference strain and the influence of fitness, mutational target size and mutational

biases on the outcome of evolution can be evaluated.

McDonald *et al.* (McDonald, et al. 2009) showed that each time the tape of WS

evolution is re-run mutations generating the adaptive type arise in one of three DGC-

encoding pathways (Wsp, Aws, or Mws) (Figure 1A). Subsequent work revealed that

when these three pathways are eliminated from the ancestral type that evolution

proceeds along multiple new pathways (Lind, et al. 2015). Preferential usage of Wsp,

Aws and Mws pathways stems from the fact that they are subject to negative

regulation and thus, relative to pathways subject to positive regulation, or requiring

promoter-activating mutations, gene fusion events, or other rare mutations, present a

large mutational target.

Given repeatability of WS evolution, knowledge of the Wsp/Aws/Mws pathways,

plus genetic tools for mechanistic investigation — including capacity to obtain WS

mutants in the absence of selection — the WS system offers a rare opportunity to

explore the feasibility of developing bottom-up strategies for evolutionary

forecasting. Our findings show that mechanistic-level predictions are possible, but

also draw attention to challenges that stem from current inability to *a priori* predict

locus specific mutational biases and environment-specific fitness effects.

Results

Obtaining an unbiased measure of pathway-specific mutation rates to WS

Knowledge of the rate at which mutation generates WS types via each of the Wsp,

Aws and Mws pathways — unbiased by the effects of selection — provides a

benchmark against which the predictive power of null models can be appraised. To

achieve such measures we firstly constructed a set of genotypes containing just one of

the three focal pathways: PBR721 carries the Wsp pathway but is devoid of Aws and

Mws, PBR713 carries the Aws pathway but is devoid of Wsp and Mws, while

PBR712 harbours the Mws pathway but is devoid of Wsp and Aws. Into each of

these genotypes a promoterless kanamycin resistance gene was incorporated

immediately downstream of the promoter of the cellulose-encoding Wss operon and

fused to an otherwise unaffected Wss operon (Figure 1B).

In the ancestral SM genotype the cellulose promoter is inactive in shaken King's

Medium B (KB) broth and thus the strain is sensitive to kanamycin. When a WS-

causing mutation occurs the *wss* promoter becomes active resulting in a kanamycin-

resistant WS type (Fukami, et al. 2007; McDonald, et al. 2011). Individual growth of

this set of three genotypes in shaken KB, combined with plating to detect kanamycin-

resistant mutants, makes possible a fluctuation assay (Luria and Delbruck 1943; Hall,

et al. 2009) from which a direct measure of the rate at which WS mutants arise can be

obtained. Importantly, because WS types are maladapted in shaken broth culture, the

screen for kanamycin-resistant clones allows the pathway-specific mutation rate to

WS to be obtained without the biasing effects of selection for growth at the air-liquid

interface (Figure 1B). The results are shown in Figure 2.

The mutation rate was highest for the Aws pathway (6.5×10^{-9}); approximately

double that of Wsp (3.7×10^{-9}) and an order of magnitude higher than that of the Mws

pathway (0.74×10^{-9}) (Figure 2). The rate at which WS mutants arose from the

ancestral genotype in which the three pathways are intact (11.2×10^{-9}) was

approximately the sum of the rates for the three pathways (11.0×10^{-9}) confirming

that the Wsp, Aws and Mws pathways are the primary routes by which WS types

evolve (Lind, et al. 2015). That the Aws pathway has the greatest capacity to

generate WS is surprising given the smaller target size (three genes compared to

seven genes in the Wsp pathway).

Figure 2. Mutation rates to WS. Fluctuation tests were used to estimate the mutation

rate to WS for the three common mutational pathways to WS. Error bars represent

mean and 95% confidence intervals.

Modelling the genotype-to-phenotype map underpinning WS evolution

Much is known about the function and interactions among components of each of the

three focal pathways. This knowledge allows development of models that capture the

dynamic nature of each pathway and thus allow predictions as to the likelihood that

evolution will precede via each of the three pathways. An unresolved issue is the

extent to which these models match experimental findings. Following a brief

description of each pathway we describe the models.

The 8.4 kb Wsp pathway is a chemotaxis-like system (Goymer, et al. 2006; Guvener

and Harwood 2007; Romling, et al. 2013; Micali and Endres 2016) comprised of

seven genes with the first six genes (*wspA-wspF*) being transcribed as a single unit

and the last (*wspR* from its own promoter (Bantinaki, et al. 2007). WspA

(PFLU1219) is a methyl-accepting chemotaxis (MCP) protein that forms a complex

with the CheW-like scaffolding proteins WspB (PFLU1220) and WspD (PFLU1222).

WspA senses environmental stimuli and transmits the information via conformational

changes in the WspA/WspB/WspD complex to effect activity of WspE (PFLU1223),

a CheA/Y hybrid histidine kinase response regulator. WspE activates both the WspR

(PFLU1225) diguanylate cyclase (DGC) and the CheB-like WspF methylesterase

WspF (PFLU1224) following transference of an active phosphoryl group. The

activity of WspA is modulated by methylation: the constitutively active CheR-like

methyltransferase WspC (PFLU1221) transfers methyl groups to conserved glutamine

residues on WspA while when phosphorylated, WspF serves to remove these groups.

WS mutants are known to arise by mutations in the WspF negative regulator and also

in the WspE kinase (McDonald, et al. 2009). In vitro manipulations of WspR that

abolish repression of the DGC domain by the response regulator domain are known,

but have never been observed to occur in experimental populations (Goymer, et al.

2006).

The 2.3 kb *aws* operon contains three genes transcribed from a single promoter

(*awsXRO). Homologous genes in *Pseudomonas aeruginosa* (*yfiRNB,* PA1121-1119)

have been characterised in detail (Malone, et al. 2010; Malone, et al. 2012; Xu, et al.

2016). The outer membrane lipoprotein AwsO (PFLU5209) has an OmpA domain, a

signal peptide and binds to peptidoglycan. AwsO is thought to be the sensor whose

activity is modulated in response to envelope stress (Malone, et al. 2012). AwsO

sequesters the periplasmic protein AwsX (PFLU5211) at the outer membrane. AwsX

functions as a negative regulator of the DGC AwsR (PFLU5210) in the inner

membrane. Both increased binding of AwsX to AwsO or loss of negative regulation

by inactivation of the interaction between AwsX and AwsR can lead to WS

(McDonald, et al. 2009; Malone, et al. 2010; Malone, et al. 2012).

The 3.9 kb *mwsR* gene (PFLU5329, known as *morA* (PA4601) in *Pseudomonas*

aeruginosa)*,* encodes a predicted membrane protein with both a DGC domain that

produces c-di-GMP and a phosphodiesterase (PDE) domain that degrades c-di-GMP.

Little is known of the molecular details determining its function, but both catalytic

domains appear to be active (Phippen, et al. 2014). Deletion of the PDE domain

results in a WS phenotype with activity being dependent on a functional DGC domain

(McDonald, et al. 2009).

If the specific effect of changing each nucleotide (and sets of nucleotides) was known

then models for each pathway would not be required, but such knowledge does not

exist. We therefore take a simplifying approach in which attention focuses on the

interactions between components that correspond to reactions whose rate can either

increase, decrease, or remain unaffected, depending on mutations in the component

parts. Such mutations increase the reaction (enabling mutations), decrease the

reaction (disabling mutations) or leave them unaffected. The components and

interactions are shown in Figure 3. Figure 3 along with figure supplements 1 to 3

describe the molecular reactions and the associated differential equations governing

the dynamics of each pathway. An advantage of this simplifying approach is that

changes to a reaction may encompass mutations in more than a single component.

For example, mutations in either WspE or WspR may increase reaction r_5 of the Wsp

pathway (Figure 3A).

Figure 3. Modeling of WS. (A) Model of the Wsp pathway. R* represents the

activated form of WspR and increase of R* leads to a WS phenotype **(B)** Model for

the Aws pathway. RR represents the activated form of AwsR and increase of RR

leads to a WS phenotype. (**C**) Model of the Mws Pathway. D* represents the activated

form of the DGC domain. Our functional model places the PDE domain as a negative

regulator of DGC activity. Details of the molecular networks are found in Figure 3

figure supplement 1 for Wsp, Figure 3 figure supplement 2 for Aws and Figure 3

figure supplement 3 for Mws. In the simple null models all genetic components are

the same size, but information of mutational target size can readily be accommodated

in the model by changing the individual probabilities of disabling and enabling

mutations. The benefit of doing so might however be small if not combined with

detailed data on mutation rates (see Discussion).

Equipped with the set of mathematical descriptions it is possible to consider

combinations of enabling, disabling, and no effect changes to reaction rates and

determine the likelihood that a WS type is generated. For the Wsp system this

amounts to 3^6 or 729 combinations. An example of one set of the possible mutations

(m_i) in Wsp is 1, −1, 0, 0, 0, 0 (an increase in r_1, a decrease in r_2, but no change in r_3,

r_4, r_5, or r_6 (Figure 3A)).

Predicting the pathways that evolution follows and genetic targets

To determine whether mutations producing WS occur more often in Wsp compared to

Aws or Mws pathways, we adopt a Bayesian approach in which the probability that a

particular pathway is used is decomposed into two terms: the probability that a

particular set of mutations (m_i) occurs in Wsp (or Aws, or Mws) represented as P (m_i

$\in$ Wsp) and the probability that those mutations give rise to a wrinkly spreader

represented as P (WS |m_i $\in$ Wsp) (or Aws, or Mws).

$P(WS \cap m \cap Wsp) = \sum_i P(WS|m_i \in Wsp)P(m_i \in Wsp)$ (1)

To estimate P(m_i $\in$ Wsp) we assume fixed probabilities of enabling and disabling

mutations and compute the product. Thus, the probability of m_i = 1, −1, 0, 0, 0, 0 is

$p_e p_d (1 − p_e − p_d)^4$, where p_e is the probability of a mutation with an enabling effect

and p_d is the probability of a mutation with a disabling effect. Recognising the value

of accommodating the possibility of localised mutational bias we note that p_e and p_d

can be adjusted for the affected reactants. The second term, P (WS |$m_i \in$ Wsp),

requires knowing both how gene products interact and how these interactions result in

a phenotype. This information is estimated based on the pathway dynamics

represented in Figure 3 and Figure 3 – figure supplement 1 by repeated sampling from

the space of all possible reaction rates, initial concentrations, and magnitude of effects

(see Materials and Methods).

The results of simulations are shown in Figure 4. Figure 4A shows that the Wsp

pathway is predicted to be the target of mutation 1.3 - 2.1 times more often than the

Aws pathway while Figure 4B shows that the Mws pathway is predicted to be the

target of mutation 0.7 - 1.0 times less often that the Aws pathway. While these results

agree with the experimental data showing Mws to be least likely pathway to be

followed, the predictions are at odds with the mutation rate data showing WS types to

be twice as likely to arise from mutation in Aws, versus Wsp. The causes of this

discrepancy are described in the following section.

Figure 4. Modelling of the probability of using the Wsp, Aws and Mws pathways

(A) Probability of Wsp relative to Aws. Equal mutation rates for all components.

(B) Probability of Mws relative to Aws. Equal mutation rates for all components.

(C) Probability of Mws relative to Wsp. Equal mutation rates for all

components.**(D) Probability of Wsp relative to Aws with the mutational hotspot**

in AwsX included. Based on the mutational data a hot spot for Aws is included in the

models that increase the mutation rate by a factor five for both enabling and disabling

mutations. **(E) Probability of Mws relative to Aws with the mutational hotspot in**

AwsX included. The relative contributions of individual reactions rates are available

in Figure 4 - figure supplement 1 for Wsp and Figure 4 - figure supplement 1 for

Aws.

In addition to predicting the preferred mutational pathways to produce a WS, the

Bayesian approach also predicts genes likely to be affected by mutation (Figure 4

figure supplement 1 and 2). Predictions as to specific genetic targets come from

appraisal of the relative importance of each reaction in terms of generating a wrinkly

spreader (Figure 4 – supplement 1 and 2). While it is recognised that a majority of

WS mutations arise from mutations in negative regulators of DGC activity, such as

WspF and AwsX (McDonald, et al. 2009; Lind, et al. 2015), further predictions are

possible based on impacts of alterations in gene function on reaction rates. For

example, with reference to the Wsp pathway (Figure 4 – figure supplement 1), there

are two reactions (r_2 and r_6) that affect WspF function: r_2 describes the rate of removal

of methyl groups from the signalling complex and r_6 the rate at which WspF is

activated by transfer of active phosphoric groups from the WspE kinase. Loss-of-

function (disabling) mutations being much more common than gain-of-function

(enabling) mutations means that both WspF and WspE are likely targets. The null

model predicts that in the area of parameter space in which Wsp is most likely

compared to Aws, 45% of the time WS will be generated when the second reaction,

r_2, is altered (Figure 4 – figure supplement 1). The same is true for reaction r_6. Thus

the presence of a negative regulator is predicted to extend the mutational target size

well beyond the gene itself. This is also true for Aws, where r_3 is the main contributor

to the WS phenotype in the case where disabling change is more common than

enabling change. Here mutations are predicted not only in the negative regulator

AwsX, but also in the interacting region of the DGC AwsR (Figure 4 – figure

supplement 2).

Loss-of-function mutations in negative regulators and their interacting partners are

not the only predicted targets. For Wsp r_1, r_3, r_4, and r_5 are altered approximately 5%

of the time in the parameter region where disabling mutations are more common than

enabling mutations, but contribute more when the rate of enabling mutations is

increased (Figure 4 – figure supplement 1). Enabling mutations based on the model

are likely to be found in WspC increasing r_1, WspABD increasing r_3, WspABD/WspE

increasing r_4 and WspR increasing r_5 (Figure 2A). For Aws, enabling mutations are

predicted to increase reaction r_1 by mutations causing constitutive activation of AwsO,

r_2 increasing binding of AwsO and AwsX and r_4 increasing formation of the active

AwsR dimer (Figure 3B, Figure 4 – figure supplement 2).

In summary, high rates of WS mutations are predicted for *wspF, wspE, wspA, awsX*

and *awsR* with lower rates for *wspC, wspR* and *awsO*. Several of these predictions sit

in accord with previous experimental observations, however, notable are predictions

that evolution might also target *wspA* and *wspR*, two genes that have not previously

been identified as mutational causes of WS types (McDonald, et al. 2009).

Analysis of mutants reveals sources of mutational bias

There are several reasons why predictions from the models might be out of kilter with

experimental data on mutation rates. We firstly looked to the distribution of WS

generating mutations among the 109 mutants collected during the course of the

fluctuation assays. Of the 109 mutants, 105 harboured a mutation in *wsp* (46

mutants), *aws* (41 mutants) or *mws* (18 mutants) (Figure 5A, Figure 5 – source data

1). The remaining four had mutations in previously described rare pathways

(PFLU0085, PFLU0183), again confirming that these non-focal pathways produce

just a fraction of the total set of WS mutants (Lind, et al. 2015).

The distribution of mutations for each of the three pathways is indicative of bias. As

shown in Figure 5B, almost 29% of all WS-causing mutations (adjusted for

differences in mutation rates between the three pathways) were due to an identical 33

base pair in-frame deletion in *awsX* (Δt229-g261, ΔY77-Q87), while a further 13 %

were due to an identical mutation (79 a->c, T27P) in *awsR*. At least 41 different

mutations in Aws can lead to WS: if mutation rates were equal for these sites the

probability of observing 20 identical mutations would be extremely small. In fact 10

million random samplings from the observed distribution of mutations failed to

recover this bias. While the Wsp pathway also contains sites that were mutated more

than once (six positions were mutated twice, one site three times and one five times),

sources of mutational bias in Wsp were less evident than in Aws (Figure 5B).

The mathematical models presented above assumed no mutational bias thus the lack

of fit between mutation rate data and predictions from the models. Nonetheless,

changing specific reaction rates within the models readily incorporates such

knowledge. For example, the mutational hotspot in *awsX* affects reactions r_2 and r_3 in

the Aws differential equation system (Figure 3B, Figure 3 – figure supplement 2).

The effect of a five-fold change in the probability of enabling/disabling change in

these reactions leads to the prediction that the Aws pathway is more likely to generate

WS types than Wsp for most probability values (see Figure 4D). The only area of

parameter space in which evolution is more likely to utilise the Wsp pathway is for

rare mutations that have a high probability of enabling change ($>> 10^{-2}$). One

interesting consequence is that it changes the phase-space over which evolution of

WS via mutations in the Wsp pathway is more likely with respect to the Aws

pathway. In Figure 4A, evolution is most likely to proceed via the Wsp pathway when

the probability of disabling change is greater than the probability of enabling change.

In contrast, when the likelihood of producing WS types is affected by the mutational

hotspot in *awsX*, then evolution will proceed via Wsp only when the probability of

enabling change is greater than the probability of disabling change (Figure 4D and

Figure 4 figure supplement 2).

Analysis of mutants reveals mutational targets and effects

Wsp pathway: Mutations were identified in five genes of the seven-gene pathway all

of which were predicted by the null model (Figure 4 – figure supplement 1). The most

commonly mutated gene was *wspA* (PFLU1219), with ten of 15 mutations (Figure 5)

being amino acid substitutions (six unique) clustered in the region 352-420 at the stalk

of the signalling domain. This region has been implicated in trimer-of-dimer

formation for the WspA homologue in *Pseudomonas aeruginosa* (O'Connor, et al.

2012) which is critical for self-assembly and localization of Wsp clusters in the

membrane. It is possible that these mutations stabilize trimer of dimer formation,

change the subcellular location of the Wsp complex, or affect interaction with WspD

(putative interface 383-420 in WspA) (Griswold, et al. 2002) and thus affecting relay

of signal to WspE. These effects we interpreted as enabling mutations increasing r_3 in

Figure 2A. The four additional mutations were in frame deletions in a separate region

of the transducer domain (ΔT293 - E299, ΔA281-A308). Although it is possible that

these mutations could also affect trimer-of-dimer formation, there are predicted

methylation sites in the region (Rice and Dahlquist 1991) that regulate the activity of

the protein via methyltransferase WspC and methylesterase WspF. Given that

disabling mutations are more common than enabling mutations it is likely that these

mutations decrease r_2 in Figure 3A by disrupting the interaction with WspF. We also

identified a single mutation that fused the open reading frame of WspC, the

methyltransferase that positively regulates WspA activity, to WspD, resulting in a

chimeric protein (Figure 5, Figure 5 – source data). This mutation is likely to be a rare

enabling mutation that increases the activity of WspC (increasing r_1 in Figure 3A) by

physically tethering it to the WspABD complex thus allowing it to more effectively

counteract the negative regulator WspF. Alternatively, the tethering may physically

block the interaction with WspF (decrease of r_2 in Figure 3A).

The second most commonly mutated gene in the *wsp* operon was *wspE* (PFLU1223)

(Figure 5). Four amino acids were repeatedly mutated in the response regulatory

domain of WspE and all cluster closely in a structural homology model made with

Phyre2 (Kelley, et al. 2015). All mutated residues surround the active site of the

phosphorylated D682 and it is likely that they disrupt feedback regulation by

decreasing phosphorylation of the negative regulator WspF (decreasing r_6) rather than

increasing activation of WspR (r_5 in Figure 3A).

Figure 5. Mutational targets. (**A**) 105 independent mutations in the *wsp* (n=46), *aws* (n=41) and *mws* (n=18) operons were identified. Numbers of independent mutants are shown in brackets. Full details on the mutations are available in Figure 5 – source data 1. (**B**) Diversity of mutations with area proportional to mutation rate (Figure 2). Two mutations (AwsX ΔY77-Q87 and AwsR T27P) contribute 41% of all mutations to WS suggesting that these are mutational hot spots. The increased mutation rate can be incorporated into the null model by increasing the probability of change for r_2 and r_3 in the Aws model (Figure 3B).

Twelve mutations were detected in *wspF* (PFLU1224). These are distributed throughout the gene and include amino acid substitutions, in-frame deletions as well as a frame-shift and a stop codon (Figure 5). The pattern of mutations is consistent with both the role of WspF as a negative regulator of WspA activity and the well-

characterised effect of loss-of-function mutations in this gene (Bantinaki, et al. 2007;

McDonald, et al. 2009). The mutations are interpreted as decreasing r_2 in Figure 3A.

Five mutations were found in WspR (PFLU1225), the DGC output response regulator

that produces c-di-GMP and activates expression of cellulose (Figure 5). All

mutations were located in the linker region between the response regulator and DGC

domains. Mutations in this region are known to generate constitutively active *wspR*

alleles by relieving the requirement for phosphorylation (Goymer, et al. 2006). They

may additionally affect subcellular clustering of WspR (Huangyutitham, et al. 2013)

or shift the equilibrium between the dimeric form of WspR, with low basal activity,

towards a tetrameric activated form (De, et al. 2009). In our model these increase

reaction r_5.

Aws pathway: Mutations were identified in all three genes of the Aws pathway – all of

which were predicted by the null model. In the Aws pathway, mutations were most

commonly found in *awsX* (25 out of 41 mutations (Figure 5)). The above-mentioned

mutational hotspot produced in-frame deletions likely mediated by 6 bp direct repeats

(Figure 5 – source data 1). The deletions are consistent with a loss of function and a

decrease in r_3 (Figure 3B) that would leave the partially overlapping open reading

frame of the downstream gene (*awsR*) unaffected.

The DGC AwsR, was mutated in 14 cases with an apparent mutational hot spot at

T27P (9 mutants) in a predicted transmembrane helix (amino acids 19-41). The

remaining mutations were amino acid substitutions in the HAMP linker and in the

PAS-like periplasmic domain between the two transmembrane helices. These amino

acid substitutions are removed from the output DGC domain (Figure 5) and their

effects are difficult to interpret, but they could cause changes in dimerization

(Malone, et al. 2012) or the packing of HAMP domains, which could, in turn, alter

transmission of conformational changes in the periplasmic PAS-like domain to the

DGC domain causing constitutive activation (Parkinson 2010). Such effects would

increase r_4 in Figure 3B. Mutations in the N-terminal part of the protein are easier to

interpret based on the existing functional model (Malone, et al. 2012) and most likely

disrupt interactions with the periplasmic negative regulator AwsX resulting in a

decrease in r_3 in Figure 3B.

Two mutations were found in the outer membrane lipoprotein protein AwsO between

the signal peptide and the OmpA domain (Figure 5). Both mutations were glutamine

to proline substitutions (Q34P, Q40P), which together with a previously reported

G42V mutation (McDonald, et al. 2009) suggest that multiple changes in this small

region can cause a WS phenotype. This is also supported by data from *Pseudomonas*

aeruginosa in which mutations in nine different positions in this region lead to a small

colony variant phenotype similar to WS (Malone, et al. 2012). A functional model

based on the YfiBNR in *P. aeruginosa* (Malone, et al. 2012; Xu, et al. 2016), suggest

that AwsO sequesters AwsX at the outer membrane and that mutations in the N-

terminal part of the protein lead to constitutive activation and increased binding of

AwsX. This would correspond to an increase in r_2 in Figure 3B, which would relieve

negative regulation of AwsR.

Mws pathway: The MwsR pathway (comprising just a single gene) harbours

mutations in both DGC and phosphodiesterase (PDE) domains . Only mutations in

the C-terminal phosphodiesterase (PDE) domain were predicted (Figure 3C). Eleven

of 18 mutations were identical in-frame deletions (ΔR1024-E1026) in the PDE

domain, mediated by 8 bp direct repeats (Figure 5, Figure 5 – source data 1). It has

been shown previously that deletion of the entire PDE domain generates the WS

phenotype (McDonald, et al. 2009), suggesting a negative regulatory role that causes

a decrease of r_2 in the model in Figure 3C. One additional mutation was found in the

PDE domain (E1083K) located close to R1024 in a structural homology model made

with Phyre2 (Kelley, et al. 2015), but distant to the active site residues (E1059-

L1061). Previously reported mutations (A1018T, ins1089DV) (McDonald, et al.

2009) are also removed from the active site and cluster in the same region in a

structural homology model. This suggests that loss of phosphodiesterase activity may

not be the mechanism leading to the WS phenotype. This is also supported by the

high solvent accessibility of the mutated residues, which indicates that major stability-

disrupting mutations are unlikely and changes in interactions between domains or

dimerization are more probable. Thus, it is likely that the WS phenotype resulting

from a deletion in the PDE domain is caused by disruption of domain interactions or

dimerization rather than loss of phosphodiesterase activity.

The remaining mutations within *mwsR* are amino acid substitutions in the GGDEF

domain, close to the DGC active site (927-931) with the exception of a duplication of

I978-G985. While it is possible that these mutations directly increase the catalytic

activity of the DGC, increasing r_1 in Figure 3C, such enabling mutations are

considered to be rare. An alternative hypothesis is that these mutations either interfere

with c-di-GMP feedback regulation or produce larger conformational changes that

change inter-domain or inter-dimer interactions, similar to the mutations in the PDE

domain. Based on these data we reject the current model of Mws function, which

predicted mutations decreasing r_2 (Figure 3C) through mutations inactivating the PDE

domain. We instead suggest that the mutations are likely to disrupt the conformational

dynamics between the domains and could be seen either as activating mutations

causing constitutive activation or disabling mutations with much reduced mutational

target size that must specifically disrupt the interaction surface between the domains.

In both cases the previous model lead to an overestimation of the rate to WS for the

Mws pathway.

Fitness of WS types

We measured the fitness of representative WS types with mutations in each of the

mutated genes (*wspA*, *wspC/D*, *wspE*, *wspF*, *wspR*, *awsX*, *awsR*, *awsO*, *mwsR*) in 1:1

competitions against a reference WspF ΔT226-G275 deletion mutant marked with

GFP (Figure 6). This type of fitness data should be interpreted with caution because

the fitness of WS mutants have been shown to be frequency dependent and some WS

mutants are superior in early phase attachment as opposed to growth at the air-liquid

interface (Lind, et al. 2015). Nevertheless, these competition experiments provide an

estimate of fitness when several different WS mutants compete at the air-liquid

interface (a likely situation given a $\sim 10^{-8}$ mutation rate to WS and a final population

size of $>10^{10}$). The fitness data account for the over- or under-representation of some

WS mutants when grown under selection (McDonald, et al. 2009) compared to those

uncovered without selection (as reported here).

Figure 6. Fitness of different WS mutants. Competitive fitness against a WspF ΔT226-G275 reference strain was measured for representative mutations in the Wsp, Aws, Mws pathways. Pairwise competitions were performed in quadruplicates and error bars represent +/- one standard deviation.

The three *wspF* mutants, the *wspC-wspD* fusion, and the *wspE* mutants have similar fitness. In contrast, both *wspA* mutants are slightly less fit and both *wspR* mutants are severely impaired (Figure 6). This sits in accord with previous work in which mutations generating WS obtained with selection have been detected in *wspF* and *wspE*, but not *wspA* or *wspR* (Goymer, et al. 2006; McDonald, et al. 2009). All *awsXRO* mutants have similar low fitness compared to the *wspF* reference strain (Figure 6), which explains why under selection these are found at lower frequencies compared to mutations in the *wsp* pathway (McDonald, et al. 2009) despite a roughly two-fold higher mutation rate to WS.

Differences of mutational spectra with and without selection

A final question concerns the outcome of the original experimental evolution under selection (McDonald, et al. 2009) and whether it can be explained by our detailed measurement of mutation rates, mutational targets and fitness assays. As indicated above, there exist major differences in the spectrum of mutations isolated with and without selection (Figure 7). The most obvious difference is in the use of the Wsp pathway, which is most commonly used (15/24) under selection and yet produces WS

types at a lower rate than the Aws pathway. The explanation lies in the lower fitness

of Aws mutants (Figure 6). Similarly, fitness effects also explains differences in the

spectrum of *wsp* mutations, with no *wspA* mutations being found under selection

despite being the most commonly mutated gene without selection (15/46). The

previous failure to detect *wspR* mutants in a screen of 53 WS mutants (Goymer, et al.

2006) is similarly explained by low fitness of WS types arising via mutations in this

gene.

Figure 7. WS mutations isolated with and without selection. Fitness effects bias

the mutational spectrum observed under selection resulting in underrepresentation of

WspA and WspR compared to WspF and WspE. Similar fitness effects of different

Aws mutants lead to similar patterns regardless of selective conditions. Only within

operon comparisons are valid for this figure as the mutants isolated without selection

had double deletions of the other operons. Between operon mutation rates are

available in Figure 2.

Discussion

The issue of evolutionary predictability and the relative importance of stochastic

events compared to deterministic processes have a long history in evolutionary

biology (Darwin 1872; Simpson 1949; Jacob 1977; Gould 1989; Conway Morris

2003; Orgogozo 2015). Recent interest has been sparked by an increasing number of

observations that evolution, under certain circumstances, can be remarkably

repeatable (Colosimo, et al. 2005; Shindo, et al. 2005; Jost, et al. 2008; Barrick, et al.

2009; Lee and Marx 2012; Meyer, et al. 2012; Zhen, et al. 2012; Herron and Doebeli

2013), but whether these cases are representative for evolutionary processes in

general remains to be determined. A related question, with greater potential for

practical applications, is whether it is possible to forecast short-term evolutionary

events and if so, then the challenge is to stipulate the data necessary to make

successful predictions.

Our uniquely detailed knowledge of the WS experimental evolution system has

provided a rare opportunity to disentangle the contributions of selection, mutational

biases and genetic architecture to evolutionary outcomes, and consequently explore

the limits of evolutionary forecasting. A thorough understanding of the function of the

molecular species and their interactions allowed development of a null model by

defining the genotype-to-phenotype map, which successfully predicted mutational

targets and the relative likelihood that evolution followed each of the three principle

pathways. Importantly, genetic architecture is likely to be transferable between

different species, which stands to allow the formulation of general principles and

evolutionary rules (Lind, et al. 2015). Despite the simplicity of the mathematical null

models, which contain only general information about functional interactions, we

successfully predicted mutational targets including previously unknown mutations in

wspA. Without information about fitness and mutational biases, however, only order

of magnitude predictions of mutant frequencies can be made. Thus, it is possible to

predict that Wsp, which is subject to negative regulation will be more common than a

DGC that requires enabling mutations (Lind, et al. 2015), but not which of two

pathways (Wsp and Aws) with differently wired negative regulation is likely to be

dominant after selection. Insights from the null model combined with data on

mutational targets also allowed us to reject our functional model of Mws.

Direct measurement of the fitness effects of large number of mutations is difficult and

time-consuming and typically only possible for microbial species. Therefore future

success in predicting fitness effects of mutations rests on the ability to infer them

from other parameters, such as estimated effects of mutations on thermodynamic

stability or molecular networks, or from incorporation of information concerning

evolutionary conservation of amino acid residues. Alternatively, it might be possible

to extrapolate findings from a small number of mutations that are either directly

constructed and assayed in the laboratory or through fitness estimates of

polymorphisms data for natural populations. Recent work on the prediction of the

fitness effects of random mutations in several genes suggests that in many cases large

effect mutations can be predicted using methods based on evolutionary conservation

(Lind, et al. 2017a).

Interestingly WS mutations in the same gene often have similar fitness effects (Figure

6). Obviously no general conclusions can be drawn from these few cases, but if

mutations with similar functional effects, for example disruption of a particular

interaction, can be assumed to be equally fit, this would greatly reduce the number of

specific mutants that need to be experimentally assayed for each gene. Several studies

suggest that fitness distributions are often bimodal, with a significant proportion being

complete loss-of-function mutations, which could explain the similar fitness effects of

mutations in the same genes if they result in inactivation of a particular biochemical

reaction of interaction (Sanjuan 2010; Jacquier, et al. 2013; Sarkisyan, et al. 2016;

Lind, et al. 2017a). The extent to which fitness effects are transferable between strains

with different genetic backgrounds or closely related species remains to be more fully

investigated (Ungerer, et al. 2003; Pearson, et al. 2012; Wang, et al. 2014).

Estimates of genomic mutation rates are remarkably consistent across species (Drake

1991), and mutational biases as evident in the types of base substitutions, are well-

characterized for a large number of bacterial species (Sung, et al. 2012; Wei, et al.

2014; Farlow, et al. 2015; Foster, et al. 2015; Long, et al. 2015). It is also known that

molecular processes, such as transcription and replication, can introduce mutational

biases (Beletskii and Bhagwat 1996; Hudson, et al. 2003; Lind and Andersson 2008;

Reijns, et al. 2015; Zhao, et al. 2015) and mutational hotspots caused by

homopolymeric tracts and direct repeats can greatly increase local mutation rate

(Streisinger and Owen 1985; Seier, et al. 2011). However, the distribution of mutation

rates across a gene or operon remains largely unknown. Absence of this knowledge

currently hinders efforts to forecast adaptive evolution.

There are several cases of probable mutational hotspots in the spectrum of WS

mutants found in this study before the influence of selection. One specific deletion

(ΔY77-Q87) in *awsX* accounts for nearly half (20/41) of the mutations in the Aws

pathway. Thus, despite the existence of hundreds of possible mutations leading to WS

(this work and (McDonald, et al. 2009; McDonald, et al. 2011; Lind, et al. 2015)) one

single mutation accounts for more than one quarter of all WS mutations. While the six

base pair direct repeat flanking the deletion provides a convincing explanation for its

increased rate, it is not clear why this deletion would be ten times more common than

the ΔP34-A46 deletion in the same gene that is flanked by ten base pair repeats and

contains five base pairs identical to those from the ΔY77-Q87 deletion (Figure 5 –

source data 1). There are also instances where single base pair substitutions are

overrepresented: the AwsR T27P mutation is found in nine cases, while eight other

single pair substitutions in Aws were found only once. Consider further the fact that

WspE (a gene of ~2.3 kb), where changes to only four specific amino acids repeatedly

cause WS, and WspF (a gene of ~1 kb) where any mutation that disrupts function

results in WS (Figure 5) contribute equally to the evolution of WS types. Together,

these findings draw attention to the limited value of including mutational target size

alone as a parameter in predictive models.

It is evident from these findings and from related studies (Pollock and Larkin 2004)

that there is need for detailed experimental measurement of local mutation rates in

specific systems. Such investigations stand to contribute to understanding of the

causes of mutational bias and the extent to which biases might be conserved among

related or even unrelated organisms. If local nucleotide sequence is the major

determinant, an estimate of mutation rate will apply strictly to very closely related

species, but if the dynamics of molecular processes, such as transcription and

replication (Sankar, et al. 2016), are major influences then estimates might be

applicable to a wider range of species.

Evolutionary forecasting is likely to be most successful for biological systems where

there are experimental data on a large number of independent evolutionary events,

such as influenza, HIV and cancer (Kouyos, et al. 2012; Fraser, et al. 2014; Lawrence,

et al. 2014; Luksza and Lassig 2014; Neher, et al. 2014; Eirew, et al. 2015). Evolution

might appear idiosyncratic indicating that every specific system requires detailed

investigation, but our hope is that deeper knowledge of the distribution of fitness

effects and mutational biases will allow short term forecasts to be produced using

modelling without the need for large-scale experimental studies. A major boost to

further refinement of evolutionary forecasting is likely to come from combining

coarse and fine-grained approaches. Our demonstration that simple null models of

functional networks can produce highly relevant quantitative predictions is an

important step forward allowing predictions to be directly tested in other experimental

systems.

Materials and methods

Strains and media

The strains used in the study are all *Pseudomonas fluorescens* SBW25 (Silby, et al.

2009) or derivatives thereof. The reporter construct (pMSC), used for isolation of WS

mutants before selection, fused the P_{wss} promoter to a kanamycin resistance marker

(*nptII*) (Fukami, et al. 2007; McDonald, et al. 2011). *P. fluorescens* strains with

deletions of the *wsp* (PFLU1219-1225), *aws* (PFLU5209-5211) and *mws* (PFLU5329)

operons were previously constructed as described by McDonald *et al.* (McDonald, et

al. 2011). All experiments used King's medium B (KB) (King, et al. 1954), solidified

with 1.5% agar and incubation was at 28°C. All strains were stored in glycerol saline

at -80°C.

Fluctuation tests and isolation of WS mutants before selection

Strains with the pMSC reporter construct and either wild type genetic background or

double deletions of *aws*/*mws*, *wsp*/*mws* or *wsp*/*aws* were used to estimate mutation

rates to WS before selection. Overnight cultures were diluted to approximately 10^3

cfu/ml and 60 independent 110 ul cultures were grown for 16-19 h (OD600= 0.9-1.0)

with shaking (200 rpm) in 96-well plates before plating on KB plates with 30 mg/l

kanamycin. Viable counts were estimated by serial dilution and plating on KB agar.

One randomly chosen colony per independent culture with WS colony morphology

was restreaked once on KB agar. The assay was repeated at least four times for the

double deletion mutants and twice for the wild type strain. Mutations rates were

estimated using the Ma-Sandri-Sarkar Maximum Likelihood Estimator (Hall, et al.

2009) available at www.keshavsingh.org/protocols/FALCOR.html.

Sequencing

Mutations causing the WS phenotype were identified by Sanger sequencing of

candidate genes in the remaining common pathway to WS, for example the *wsp*

operon for the *aws*/*mws* deletion strain. In a few cases where no mutations were

identified in the previously established WS target genes, we used genome sequencing

(Illumina HiSeq, performed by Macrogen Korea).

Fitness assays

Competition assays were performed as previously described (Lind, et al. 2015) by

mixing the WS mutant 1:1 with a reference strain labelled with green fluorescent

protein and measuring the ratio of each strain before and after static growth for 24 h

using flow cytometry (BD FACS Canto). We used a WspF ΔT226-G275 deletion

mutant as the reference strain because WspF mutants are the most commonly found

WS type when grown under selective conditions (McDonald, et al. 2009) and the in

frame deletion of 50 amino acids most likely represents a complete loss-of-function

mutation with minimal polar effects on the downstream *wspR*. Selection coefficients

per generation were calculated as $s = [\ln(R(t)/R(0))]/[t]$, as previously described

(Dykhuizen 1990) where R is the ratio of alternative WS mutant to WspF ΔT226-

G275 GFP and t the number of generations determined using viable counts. Control

competition experiments with isogenic WspF ΔT226-G275 reference strains with and

without GFP were used to correct for the cost of the GFP marker. Control

competitions were also used to determine the cost of the double deletions and the

reporter construct relative to a wild type genetic background, for example an AwsX

ΔY77-Q87 mutant in *Δwsp*/*Δmws* background with pMSC was competed with a GFP

labeled AwsX ΔY77-Q87 mutant in wild type background. Competitions were

performed independently inoculated quadruplicates for each strain.

Homology models

Homology models of the structure of WspA, WspE, WspR, AwsR, AwsO and MwsR

were made using Phyre2 in intensive mode (http://www.sbg.bio.ic.ac.uk/phyre2)

(Kelley, et al. 2015).

Probability estimation in the mathematical models

The differential equation models describe the interactions between proteins in each of

the three WS pathways. In order to solve the differential equations, two pieces of

information are required: i) the initial concentrations of the molecular species and ii)

the reaction rates. Although this information is unavailable a random-sampling

approach was used to generate different random sets of initial concentrations and

reaction rates. Each random set was used to establish a baseline of potential WS

expression making it possible to evaluate whether a set of mutations results in a WS

type. Effectively, this approach allows sampling of the probability distribution P (WS

$|m_i \in$ Wsp) used in our Bayesian model.

We randomly sample 1,000 different sets of reaction rates and initial concentrations

from uniform priors: reaction rates were sampled randomly from a uniform

distribution on log space (i.e. $10^{U[-2,2]}$) and initial concentrations of reactants were

sampled from a uniform distribution U[0,10]. For each set, the appropriate differential

equation model was integrated and the steady state concentration of the compounds

that correspond to a wrinkly spreader (RR in Aws, R* in Wsp and D* for Mws)

computed. This served as a baseline for the non-WS phenotype that was used for

comparison to determine whether combinations of mutations result in increased WS

expression. After obtaining the baseline, we implemented particular combinations of

enabling/disabling mutations (a m_i). Ideally, a distribution linking enabling/disabling

mutations to a fold change in reaction rates would be used, but this information is

unavailable. In order to progress the effect sizes for enabling and disabling mutations

were sampled from $10^{U[0,2]}$ and $10^{U[-2,0]}$, respectively, and then multiplied by the

reaction rates. The differential equations were then solved for the same time that it

took the baseline to reach steady state. The final concentration of R* (Figure 3A), RR

(Figure 3B) and D* (Figure 3C) was then compared to the baseline and the number of

times out of 1,000 that the WS-inducing compound increased served as an estimate of

P (WS$|m_i \in$ Wsp). The probability distribution stabilized by 500 random samples

and additional sampling did not produce significant changes (data not shown).

The absence of empirical data on reaction rates, initial concentrations, and expected

mutation effect size meant using a random sampling approach requiring estimates for

parameter ranges. Parameter ranges were chosen to be broad enough to capture

differences spanning several orders of magnitudes while allowing numerical

computations for solving the differential equations. To assess the effect of these

ranges on the results, the sampling procedure was repeated for WSP for three

different parameter regimes i) an expanded range for initial concentrations [0-50], ii)

an expanded range for reaction rates $10^{[-3,3]}$, iii) a compressed range for mutational

effect size $10^{\wedge[-1,1]}$. This analysis shows that qualitative results are robust to these

changes (see Figure S1).

Source code for the mathematical modelling is deposited as supplemental material.

Acknowledgements

824 This work was supported by Marsden Fund Council from New Zealand Government

825 funding, administered by the Royal Society of New Zealand.

Competing interests

828 The authors declare no competing interests.

Author contributions

Peter A. Lind Conceptualization, Methodology, Investigation, Data Curation,

Writing—original draft, Writing—review and editing, Visualization

Eric Libby Conceptualization, Methodology, Software, Formal analysis,

Investigation, Writing—original draft, Writing—review and editing, Visualization

Jenny Herzog Methodology, Investigation, Data Curation

Paul B. Rainey Conceptualization, Methodology, Validation, Writing—original

draft, Writing—review and editing, Visualization, Supervision, Project

administration, Funding acquisition

References

Alberch P. 1991. From genes to phenotype: dynamical systems and evolvability.

Genetica 84:5-11.

Bailey SF, Blanquart F, Bataillon T, Kassen R. 2017. What drives parallel evolution?:

How population size and mutational variation contribute to repeated evolution.

Bioessays 39:1-9. http://dx.doi.org/10.1002/bies.201600176

Bantinaki E, Kassen R, Knight CG, Robinson Z, Spiers AJ, Rainey PB. 2007.

Adaptive divergence in experimental populations of Pseudomonas fluorescens. III.

Mutational origins of wrinkly spreader diversity. Genetics 176:441-453.

http://dx.doi.org/10.1534/genetics.106.069906

Barrick JE, Yu DS, Yoon SH, Jeong H, Oh TK, Schneider D, Lenski RE, Kim JF.

2009. Genome evolution and adaptation in a long-term experiment with

Escherichia coli. Nature 461:1243-1247. http://dx.doi.org/10.1038/nature08480

Beletskii A, Bhagwat AS. 1996. Transcription-induced mutations: increase in C to T

mutations in the nontranscribed strand during transcription in Escherichia coli.

Proceedings of the National Academy of Sciences of the United States of America

93:13919-13924.

Colosimo PF, Hosemann KE, Balabhadra S, Villarreal G, Jr., Dickson M, Grimwood

J, Schmutz J, Myers RM, Schluter D, Kingsley DM. 2005. Widespread parallel

evolution in sticklebacks by repeated fixation of Ectodysplasin alleles. Science

307:1928-1933. http://dx.doi.org/10.1126/science.1107239

Conway Morris S. 2003. Life's solution inevitable humans in a lonely universe. New

York u.a.: Cambridge Univ. Press.

Darwin C. 1872. The origin of species by means of natural selection, or, The

preservation of favoured races in the struggle for life. London: John Murray.

De N, Navarro MV, Raghavan RV, Sondermann H. 2009. Determinants for the

activation and autoinhibition of the diguanylate cyclase response regulator WspR.

J Mol Biol 393:619-633. http://dx.doi.org/10.1016/j.jmb.2009.08.030

de Visser JA, Krug J. 2014. Empirical fitness landscapes and the predictability of

evolution. Nature reviews. Genetics 15:480-490. http://dx.doi.org/10.1038/nrg3744

Drake JW. 1991. A constant rate of spontaneous mutation in DNA-based microbes.

Proceedings of the National Academy of Sciences of the United States of America

88:7160-7164.

Dykhuizen DE. 1990. Experimental studies of natural selection in bacteria. Annu.

Rev. Ecol. Syst. 21:373-398.

Eirew P, Steif A, Khattra J, Ha G, Yap D, Farahani H, Gelmon K, Chia S, Mar C,

Wan A, et al. 2015. Dynamics of genomic clones in breast cancer patient

xenografts at single-cell resolution. Nature 518:422-426.

http://dx.doi.org/10.1038/nature13952

Eyre-Walker A, Keightley PD. 2007. The distribution of fitness effects of new

mutations. Nature reviews. Genetics 8:610-618. http://dx.doi.org/10.1038/nrg2146

Farlow A, Long H, Arnoux S, Sung W, Doak TG, Nordborg M, Lynch M. 2015. The

Spontaneous Mutation Rate in the Fission Yeast Schizosaccharomyces pombe.

Genetics 201:737-744. http://dx.doi.org/10.1534/genetics.115.177329

Ferguson GC, Bertels F, Rainey PB. 2013. Adaptive Divergence in Experimental

Populations of Pseudomonas fluorescens. V. Insight into the Niche Specialist

"Fuzzy Spreader" Compels Revision of the Model Pseudomonas Radiation.

Genetics. http://dx.doi.org/10.1534/genetics.113.154948

Flowers JM, Hanzawa Y, Hall MC, Moore RC, Purugganan MD. 2009. Population

genomics of the Arabidopsis thaliana flowering time gene network. Molecular

biology and evolution 26:2475-2486. http://dx.doi.org/10.1093/molbev/msp161

Foster PL, Lee H, Popodi E, Townes JP, Tang H. 2015. Determinants of spontaneous

mutation in the bacterium Escherichia coli as revealed by whole-genome

sequencing. Proceedings of the National Academy of Sciences of the United States

of America 112:E5990-5999. http://dx.doi.org/10.1073/pnas.1512136112

Fraebel DT, Mickalide H, Schnitkey D, Merritt J, Kuhlman TE, Kuehn S. 2017.

Environment determines evolutionary trajectory in a constrained phenotypic space.

Elife 6. http://dx.doi.org/10.7554/eLife.24669

Fraser C, Lythgoe K, Leventhal GE, Shirreff G, Hollingsworth TD, Alizon S,

Bonhoeffer S. 2014. Virulence and pathogenesis of HIV-1 infection: an

evolutionary perspective. Science 343:1243727.

http://dx.doi.org/10.1126/science.1243727

Fukami T, Beaumont HJ, Zhang XX, Rainey PB. 2007. Immigration history controls

diversification in experimental adaptive radiation. Nature 446:436-439.

http://dx.doi.org/10.1038/nature05629

Galen SC, Natarajan C, Moriyama H, Weber RE, Fago A, Benham PM, Chavez AN,

Cheviron ZA, Storz JF, Witt CC. 2015. Contribution of a mutational hot spot to

hemoglobin adaptation in high-altitude Andean house wrens. Proceedings of the

National Academy of Sciences of the United States of America 112:13958-13963.

http://dx.doi.org/10.1073/pnas.1507300112

Gompel N, Prud'homme B. 2009. The causes of repeated genetic evolution. Dev Biol

332:36-47. http://dx.doi.org/10.1016/j.ydbio.2009.04.040

Gould SJ. 1989. Wonderful life : the Burgess Shale and the nature of history. New

York: W.W. Norton.

Goymer P, Kahn SG, Malone JG, Gehrig SM, Spiers AJ, Rainey PB. 2006. Adaptive

divergence in experimental populations of Pseudomonas fluorescens. II. Role of

the GGDEF regulator WspR in evolution and development of the wrinkly spreader

phenotype. Genetics 173:515-526. http://dx.doi.org/10.1534/genetics.106.055863

Griswold IJ, Zhou HJ, Matison M, Swanson RV, McIntosh LP, Simon MI, Dahlquist

FW. 2002. The solution structure and interactions of CheW from Thermotoga

maritima. Nature Structural Biology 9:121-125. http://dx.doi.org/DOI

10.1038/nsb753

Hall BM, Ma CX, Liang P, Singh KK. 2009. Fluctuation analysis CalculatOR: a web

tool for the determination of mutation rate using Luria-Delbruck fluctuation

analysis. Bioinformatics 25:1564-1565.

http://dx.doi.org/10.1093/bioinformatics/btp253

Hammerschmidt K, Rose CJ, Kerr B, Rainey PB. 2014. Life cycles, fitness

decoupling and the evolution of multicellularity. Nature 515:75-79.

http://dx.doi.org/10.1038/nature13884

Heilbron K, Toll-Riera M, Kojadinovic M, MacLean RC. 2014. Fitness is strongly

influenced by rare mutations of large effect in a microbial mutation accumulation

experiment. Genetics 197:981-990. http://dx.doi.org/10.1534/genetics.114.163147

Herron MD, Doebeli M. 2013. Parallel evolutionary dynamics of adaptive

diversification in Escherichia coli. PLoS biology 11:e1001490.

http://dx.doi.org/10.1371/journal.pbio.1001490

Huangyutitham V, Guvener ZT, Harwood CS. 2013. Subcellular clustering of the

phosphorylated WspR response regulator protein stimulates its diguanylate cyclase

activity. MBio 4:e00242-00213. http://dx.doi.org/10.1128/mBio.00242-13

Hudson RE, Bergthorsson U, Ochman H. 2003. Transcription increases multiple

spontaneous point mutations in Salmonella enterica. Nucleic Acids Res 31:4517-

4522.

Jacob F. 1977. Evolution and tinkering. Science 196:1161-1166.

Jacquier H, Birgy A, Le Nagard H, Mechulam Y, Schmitt E, Glodt J, Bercot B, Petit

E, Poulain J, Barnaud G, et al. 2013. Capturing the mutational landscape of the

beta-lactamase TEM-1. Proc Natl Acad Sci U S A 110:13067-13072.

http://dx.doi.org/10.1073/pnas.1215206110

Jost MC, Hillis DM, Lu Y, Kyle JW, Fozzard HA, Zakon HH. 2008. Toxin-resistant

sodium channels: parallel adaptive evolution across a complete gene family.

Molecular biology and evolution 25:1016-1024.

http://dx.doi.org/10.1093/molbev/msn025

Kelley LA, Mezulis S, Yates CM, Wass MN, Sternberg MJ. 2015. The Phyre2 web

portal for protein modeling, prediction and analysis. Nat Protoc 10:845-858.

http://dx.doi.org/10.1038/nprot.2015.053

King EO, Ward MK, Raney DE. 1954. Two simple media for the demonstration of

pyocyanin and fluorescin. The Journal of laboratory and clinical medicine 44:301-

307.

Kouyos RD, Leventhal GE, Hinkley T, Haddad M, Whitcomb JM, Petropoulos CJ,

Bonhoeffer S. 2012. Exploring the complexity of the HIV-1 fitness landscape.

PLoS genetics 8:e1002551. http://dx.doi.org/10.1371/journal.pgen.1002551

Kram KE, Geiger C, Ismail WM, Lee H, Tang H, Foster PL, Finkel SE. 2017.

Adaptation of Escherichia coli to Long-Term Serial Passage in Complex Medium:

Evidence of Parallel Evolution. mSystems 2.

http://dx.doi.org/10.1128/mSystems.00192-16

Krasovec R, Richards H, Gifford DR, Hatcher C, Faulkner KJ, Belavkin RV,

Channon A, Aston E, McBain AJ, Knight CG. 2017. Spontaneous mutation rate is

a plastic trait associated with population density across domains of life. PLoS

biology 15:e2002731. http://dx.doi.org/10.1371/journal.pbio.2002731

Kugelberg E, Kofoid E, Andersson DI, Lu Y, Mellor J, Roth FP, Roth JR. 2010. The

tandem inversion duplication in Salmonella enterica: selection drives unstable

precursors to final mutation types. Genetics 185:65-80.

http://dx.doi.org/10.1534/genetics.110.114074

Laland KN, Uller T, Feldman MW, Sterelny K, Muller GB, Moczek A, Jablonka E,

Odling-Smee J. 2015. The extended evolutionary synthesis: its structure,

assumptions and predictions. Proceedings. Biological sciences / The Royal Society

282:20151019. http://dx.doi.org/10.1098/rspb.2015.1019

Lässig M, Mustonen V, Walczak AM. 2017. Predicting evolution. Nature Ecology &

Evolution 1:0077. http://dx.doi.org/10.1038/s41559-017-0077

Lawrence MS, Stojanov P, Mermel CH, Robinson JT, Garraway LA, Golub TR,

Meyerson M, Gabriel SB, Lander ES, Getz G. 2014. Discovery and saturation

analysis of cancer genes across 21 tumour types. Nature 505:495-501.

http://dx.doi.org/10.1038/nature12912

Lee MC, Marx CJ. 2012. Repeated, selection-driven genome reduction of accessory

genes in experimental populations. PLoS genetics 8:e1002651.

http://dx.doi.org/10.1371/journal.pgen.1002651

Levinson G, Gutman GA. 1987. Slipped-strand mispairing: a major mechanism for

DNA sequence evolution. Molecular biology and evolution 4:203-221.

http://dx.doi.org/10.1093/oxfordjournals.molbev.a040442

Lind PA, Andersson DI. 2008. Whole-genome mutational biases in bacteria.

Proceedings of the National Academy of Sciences of the United States of America

105:17878-17883. http://dx.doi.org/10.1073/pnas.0804445105

Lind PA, Arvidsson L, Berg OG, Andersson DI. 2017a. Variation in Mutational

Robustness between Different Proteins and the Predictability of Fitness Effects.

Molecular biology and evolution 34:408-418.

http://dx.doi.org/10.1093/molbev/msw239

Lind PA, Farr AD, Rainey PB. 2017b. Evolutionary convergence in experimental

Pseudomonas populations. ISME J 11:589-600.

http://dx.doi.org/10.1038/ismej.2016.157

Lind PA, Farr AD, Rainey PB. 2015. Experimental evolution reveals hidden diversity

in evolutionary pathways. Elife 4. http://dx.doi.org/10.7554/eLife.07074

Long H, Kucukyildirim S, Sung W, Williams E, Lee H, Ackerman M, Doak TG,

Tang H, Lynch M. 2015. Background Mutational Features of the Radiation-

Resistant Bacterium Deinococcus radiodurans. Molecular biology and evolution

32:2383-2392. http://dx.doi.org/10.1093/molbev/msv119

Lovett ST. 2004. Encoded errors: mutations and rearrangements mediated by

misalignment at repetitive DNA sequences. Mol Microbiol 52:1243-1253.

http://dx.doi.org/10.1111/j.1365-2958.2004.04076.x

Luksza M, Lassig M. 2014. A predictive fitness model for influenza. Nature 507:57-

61. http://dx.doi.org/10.1038/nature13087

Luria SE, Delbruck M. 1943. Mutations of Bacteria from Virus Sensitivity to Virus

Resistance. Genetics 28:491-511.

Lynch M. 2010. Rate, molecular spectrum, and consequences of human mutation.

Proceedings of the National Academy of Sciences of the United States of America

107:961-968. http://dx.doi.org/10.1073/pnas.0912629107

Maharjan RP, Ferenci T. 2017. A shifting mutational landscape in 6 nutritional states:

Stress-induced mutagenesis as a series of distinct stress input-mutation output

relationships. PLoS biology 15:e2001477.

http://dx.doi.org/10.1371/journal.pbio.2001477

Malone JG, Jaeger T, Manfredi P, Dotsch A, Blanka A, Bos R, Cornelis GR, Haussler

S, Jenal U. 2012. The YfiBNR signal transduction mechanism reveals novel targets

for the evolution of persistent Pseudomonas aeruginosa in cystic fibrosis airways.

PLoS Pathog 8:e1002760. http://dx.doi.org/10.1371/journal.ppat.1002760

Malone JG, Jaeger T, Spangler C, Ritz D, Spang A, Arrieumerlou C, Kaever V,

Landmann R, Jenal U. 2010. YfiBNR mediates cyclic di-GMP dependent small

colony variant formation and persistence in Pseudomonas aeruginosa. PLoS

Pathog 6:e1000804. http://dx.doi.org/10.1371/journal.ppat.1000804

Martin P, Sun L, Hood DW, Moxon ER. 2004. Involvement of genes of genome

maintenance in the regulation of phase variation frequencies in Neisseria

meningitidis. Microbiology 150:3001-3012.

http://dx.doi.org/10.1099/mic.0.27182-0

McDonald MJ, Cooper TF, Beaumont HJE, Rainey PB. 2011. The distribution of

fitness effects of new beneficial mutations in Pseudomonas fluorescens. Biology

Letters 7:98-100. http://dx.doi.org/Doi 10.1098/Rsbl.2010.0547

McDonald MJ, Gehrig SM, Meintjes PL, Zhang XX, Rainey PB. 2009. Adaptive

divergence in experimental populations of Pseudomonas fluorescens. IV. Genetic

constraints guide evolutionary trajectories in a parallel adaptive radiation. Genetics

183:1041-1053. http://dx.doi.org/10.1534/genetics.109.107110

Meyer JR, Dobias DT, Weitz JS, Barrick JE, Quick RT, Lenski RE. 2012.

Repeatability and contingency in the evolution of a key innovation in phage

lambda. Science 335:428-432. http://dx.doi.org/10.1126/science.1214449

Moxon ER, Rainey PB, Nowak MA, Lenski RE. 1994. Adaptive evolution of highly

mutable loci in pathogenic bacteria. Current biology : CB 4:24-33.

Neher RA, Russell CA, Shraiman BI. 2014. Predicting evolution from the shape of

genealogical trees. Elife 3. http://dx.doi.org/10.7554/eLife.03568

Orgogozo V. 2015. Replaying the tape of life in the twenty-first century. Interface

Focus 5:20150057. http://dx.doi.org/10.1098/rsfs.2015.0057

Orr HA. 2005. The genetic theory of adaptation: a brief history. Nature reviews.

Genetics 6:119-127. http://dx.doi.org/10.1038/nrg1523

Ostrowski EA, Woods RJ, Lenski RE. 2008. The genetic basis of parallel and

divergent phenotypic responses in evolving populations of Escherichia coli.

Proceedings. Biological sciences / The Royal Society 275:277-284.

http://dx.doi.org/10.1098/rspb.2007.1244

Parkinson JS. 2010. Signaling mechanisms of HAMP domains in chemoreceptors and

sensor kinases. Annu Rev Microbiol 64:101-122.

http://dx.doi.org/10.1146/annurev.micro.112408.134215

Pearson VM, Miller CR, Rokyta DR. 2012. The consistency of beneficial fitness

effects of mutations across diverse genetic backgrounds. Plos One 7:e43864.

http://dx.doi.org/10.1371/journal.pone.0043864

Phippen CW, Mikolajek H, Schlaefli HG, Keevil CW, Webb JS, Tews I. 2014.

Formation and dimerization of the phosphodiesterase active site of the

Pseudomonas aeruginosa MorA, a bi-functional c-di-GMP regulator. FEBS Lett

588:4631-4636. http://dx.doi.org/10.1016/j.febslet.2014.11.002

Pigliucci M. 2010. Genotype-phenotype mapping and the end of the 'genes as

blueprint' metaphor. Philosophical transactions of the Royal Society of London.

Series B, Biological sciences 365:557-566.

http://dx.doi.org/10.1098/rstb.2009.0241

Pollock DD, Larkin JC. 2004. Estimating the degree of saturation in mutant screens.

Genetics 168:489-502. http://dx.doi.org/10.1534/genetics.103.024430

Rainey PB, Remigi P, Farr AD, Lind PA. 2017. Darwin was right: where now for

experimental evolution? Curr Opin Genet Dev 47:102-109.

http://dx.doi.org/10.1016/j.gde.2017.09.003

Reams AB, Roth JR. 2015. Mechanisms of gene duplication and amplification. Cold

Spring Harb Perspect Biol 7:a016592.

http://dx.doi.org/10.1101/cshperspect.a016592

Reijns MA, Kemp H, Ding J, de Proce SM, Jackson AP, Taylor MS. 2015. Lagging-

strand replication shapes the mutational landscape of the genome. Nature 518:502-

506. http://dx.doi.org/10.1038/nature14183

Rice MS, Dahlquist FW. 1991. Sites of deamidation and methylation in Tsr, a

bacterial chemotaxis sensory transducer. J Biol Chem 266:9746-9753.

Richardson AR, Stojiljkovic I. 2001. Mismatch repair and the regulation of phase

variation in Neisseria meningitidis. Mol Microbiol 40:645-655.

Roth JR, Benson N, Galitsky T, Haack K, Lawrence JG, Miesel L. 1996.

Rearrangements of the Bacterial Chromosome: Formation and

Applications. In: Neidhardt FC, editor. Escherichia coli and Salmonella: Cellular and

Molecular Biology. Washington, D.C.: 2256-2276. p. 2256-2276.

Sanjuan R. 2010. Mutational fitness effects in RNA and single-stranded DNA viruses:

common patterns revealed by site-directed mutagenesis studies. Philosophical

transactions of the Royal Society of London. Series B, Biological sciences

365:1975-1982. http://dx.doi.org/10.1098/rstb.2010.0063

Sankar TS, Wastuwidyaningtyas BD, Dong Y, Lewis SA, Wang JD. 2016. The nature

of mutations induced by replication-transcription collisions. Nature.

http://dx.doi.org/10.1038/nature18316

Sarkisyan KS, Bolotin DA, Meer MV, Usmanova DR, Mishin AS, Sharonov GV,

Ivankov DN, Bozhanova NG, Baranov MS, Soylemez O, et al. 2016. Local fitness

landscape of the green fluorescent protein. Nature 533:397-401.

http://dx.doi.org/10.1038/nature17995

Seier T, Padgett DR, Zilberberg G, Sutera VA, Jr., Toha N, Lovett ST. 2011. Insights

into mutagenesis using Escherichia coli chromosomal lacZ strains that enable

detection of a wide spectrum of mutational events. Genetics 188:247-262.

http://dx.doi.org/10.1534/genetics.111.127746

Shewaramani S, Finn TJ, Leahy SC, Kassen R, Rainey PB, Moon CD. 2017.

Anaerobically Grown Escherichia coli Has an Enhanced Mutation Rate and

Distinct Mutational Spectra. PLoS genetics 13:e1006570.

http://dx.doi.org/10.1371/journal.pgen.1006570

Shindo C, Aranzana MJ, Lister C, Baxter C, Nicholls C, Nordborg M, Dean C. 2005.

Role of FRIGIDA and FLOWERING LOCUS C in determining variation in

flowering time of Arabidopsis. Plant physiology 138:1163-1173.

http://dx.doi.org/10.1104/pp.105.061309

Silby MW, Cerdeno-Tarraga AM, Vernikos GS, Giddens SR, Jackson RW, Preston

GM, Zhang XX, Moon CD, Gehrig SM, Godfrey SA, et al. 2009. Genomic and

genetic analyses of diversity and plant interactions of Pseudomonas fluorescens.

Genome biology 10:R51. http://dx.doi.org/10.1186/gb-2009-10-5-r51

Simpson GG. 1949. The meaning of evolution : a study of the history of life and of its

significance for man. New Haven: Yale University Press.

Spiers AJ, Bohannon J, Gehrig SM, Rainey PB. 2003. Biofilm formation at the air-

liquid interface by the Pseudomonas fluorescens SBW25 wrinkly spreader requires

an acetylated form of cellulose. Molecular Microbiology 50:15-27.

Spiers AJ, Kahn SG, Bohannon J, Travisano M, Rainey PB. 2002. Adaptive

divergence in experimental populations of Pseudomonas fluorescens. I. Genetic

and phenotypic bases of wrinkly spreader fitness. Genetics 161:33-46.

Steinrueck M, Guet CC. 2017. Complex chromosomal neighborhood effects

determine the adaptive potential of a gene under selection. Elife 6.

http://dx.doi.org/10.7554/eLife.25100

Stern DL. 2013. The genetic causes of convergent evolution. Nature reviews.

Genetics 14:751-764. http://dx.doi.org/10.1038/nrg3483

Stern DL, Orgogozo V. 2009. Is genetic evolution predictable? Science 323:746-751.

http://dx.doi.org/10.1126/science.1158997

Stoltzfus A, McCandlish DM. 2017. Mutational Biases Influence Parallel Adaptation.

Molecular biology and evolution 34:2163-2172.

http://dx.doi.org/10.1093/molbev/msx180

Streisinger G, Owen J. 1985. Mechanisms of spontaneous and induced frameshift

mutation in bacteriophage T4. Genetics 109:633-659.

Sung W, Ackerman MS, Miller SF, Doak TG, Lynch M. 2012. Drift-barrier

hypothesis and mutation-rate evolution. Proceedings of the National Academy of

Sciences of the United States of America 109:18488-18492.

http://dx.doi.org/10.1073/pnas.1216223109

Tautz D, Renz M. 1984. Simple sequences are ubiquitous repetitive components of

eukaryotic genomes. Nucleic Acids Res 12:4127-4138.

Tenaillon O, Rodriguez-Verdugo A, Gaut RL, McDonald P, Bennett AF, Long AD,

Gaut BS. 2012. The molecular diversity of adaptive convergence. Science

335:457-461. http://dx.doi.org/10.1126/science.1212986

Ungerer MC, Linder CR, Rieseberg LH. 2003. Effects of genetic background on

response to selection in experimental populations of Arabidopsis thaliana. Genetics

163:277-286.

Viswanathan M, Lacirignola JJ, Hurley RL, Lovett ST. 2000. A novel mutational

hotspot in a natural quasipalindrome in Escherichia coli. J Mol Biol 302:553-564.

http://dx.doi.org/10.1006/jmbi.2000.4088

Wang AD, Sharp NP, Agrawal AF. 2014. Sensitivity of the distribution of mutational

fitness effects to environment, genetic background, and adaptedness: a case study

with Drosophila. Evolution 68:840-853. http://dx.doi.org/10.1111/evo.12309

Wei W, Ning LW, Ye YN, Li SJ, Zhou HQ, Huang J, Guo FB. 2014. SMAL: A

Resource of Spontaneous Mutation Accumulation Lines. Molecular biology and

evolution 31:1302-1308. http://dx.doi.org/10.1093/molbev/msu073

Woods R, Schneider D, Winkworth CL, Riley MA, Lenski RE. 2006. Tests of parallel

molecular evolution in a long-term experiment with Escherichia coli. Proceedings

of the National Academy of Sciences of the United States of America 103:9107-

9112. http://dx.doi.org/10.1073/pnas.0602917103

Xu M, Yang X, Yang XA, Zhou L, Liu TZ, Fan Z, Jiang T. 2016. Structural insights

into the regulatory mechanism of the Pseudomonas aeruginosa YfiBNR system.

Protein Cell 7:403-416. http://dx.doi.org/10.1007/s13238-016-0264-7

Zhao HL, Xia ZK, Zhang FZ, Ye YN, Guo FB. 2015. Multiple Factors Drive

Replicating Strand Composition Bias in Bacterial Genomes. Int J Mol Sci

16:23111-23126. http://dx.doi.org/10.3390/ijms160923111

Zhen Y, Aardema ML, Medina EM, Schumer M, Andolfatto P. 2012. Parallel

molecular evolution in an herbivore community. Science 337:1634-1637.

http://dx.doi.org/10.1126/science.1226630

1168

1169 **Figure supplements**

A

1. Methylated WspC activates the WspABD complex via methylation

2. Phosphorylated WspF inactivates a methylated WspABD complex by demethylation

3. A signal S can activate a methylated WspABD complex via an autophosphorylation reaction

4. A methylated and phosphorylated WspABD complex can activate WspE by phosphorylation

5. Phosphorylated WspE can phosphorylate WspR to an active form

6. Phosphorylated WspE can phosphorylate WspF

7. WspR is degraded or bound by a competitor at constant concentration (mathematically equivalent). The rate (0.01) is chosen to provide a steady, slow decay. Whithout this all WspR is converted to R*

B

1. $C\# + ABD \xrightarrow{r_1} ABD\#$

2. $F^* + ABD\# \xrightarrow{r_2} F + ABD$

3. $ABD + S \xrightarrow{r_3} ABD\#^*$

4. $ABD\#^* + E \xrightarrow{r_4} ABD\# + E^*$

5. $E^* + R \xrightarrow{r_5} R^* + E$

6. $E^* + F \xrightarrow{r_6} F^* + E$

7. $R \xrightarrow{.01} \varnothing$

C

$\frac{\partial [ABD]}{\partial t} = r_2 [F^*][ABD\#] - r_1 [ABD][C\#]$

$\frac{\partial [ABD\#]}{\partial t} = r_1 [ABD][C\#] - r_2 [F^*][ABD\#] - r_3 S[ABD\#] + r_4 [E][ABD\#^*]$

$\frac{\partial [ABD\#^*]}{\partial t} = r_3 S[ABD\#] - r_4 [E][ABD\#^*]$

$\frac{\partial [E^*]}{\partial t} = r_4 [E][ABD\#^*] - r_5 [E^*][R] - r_6 [E^*][F]$

$\frac{\partial [R^*]}{\partial t} = r_5 [R][E^*]$

$\frac{d[E]}{dt} = r_6 [E^*][F] - r_4 [E][ABD\#^*] + r_5 [R][E^*]$

$\frac{\partial [F]}{\partial t} = r_2 [F^*][ABD\#] - r_6 [E^*][F]$

$\frac{\partial [F^*]}{d\partial t} = r_6 [E^*][F] - r_2 [F^*][ABD\#]$

$\frac{\partial [R]}{\partial t} = -r_5 [R][E^*] - .01[R]$

Figure 3 - figure supplement 1. Wsp model (A) Description of functional interactions **(B)** Description of molecular reaction **(C)** Differential equations describing the dynamics of the Wsp pathway. The activity of WspA is modulated by methylation, where it is activated by the CheR-like methyltransferase WspC (PFLU1221). The CheB-like methylesterase WspF (PFLU1224) functions as a negative regulator. Modulation of WspR activity through changes in oligomerization state and clustering is not explicitly included in the model (De, et al. 2008; Huangyutitham, et al. 2013), but can be interpreted as changes in the rate of WspR activation.

A

1. A signal S binds to AwsO and activates it

2. Activated AwsO binds to AwsX to form a complex that sequesters AwsX

3. AwsX binds AwsR to form a complex that sequesters AwsX

4. AwsR dimerizes (active form)

B

1. $S + O \xrightarrow{r_1} O^\wedge$

2. $X + O^\wedge \xrightarrow{r_2} OX$

3. $X + R \xrightarrow{r_3} XR$

4. $R + R \xrightarrow{r_4} RR$

C

$\frac{\partial [X]}{\partial t} = -r_2 [X][O^\wedge] - r_3 [X][R]$

$\frac{\partial [XR]}{\partial t} = r_3 [X][R]$

$\frac{\partial [O]}{\partial t} = -r_1 S[O]$

$\frac{\partial [O^\wedge]}{\partial t} = r_1 S[O] - r_2 [X][O^\wedge]$

$\frac{\partial [OX]}{\partial t} = r_2 [X][O^\wedge]$

$\frac{\partial [R]}{\partial t} = -r_3 [X][R] - r_4 [R][R]$

$\frac{\partial [RR]}{\partial t} = r_4 [R][R]$

Figure 3 - figure supplement 2. Aws model (A) Description of functional interactions **(B)** Description of molecular reaction **(C)** Differential equations describing the dynamics of the Aws pathway. Release of AwsX mediated repression results in a conformational shift that in the model is represented as formation of an active dimer.

1187

A

1. A signal S binds to DGC domain and activates it

2. Activated DGC interacts with PDE to inactivate it

3. Activated DGC interacts with C and to form alternative complex DC

B

1. $S + D \xrightarrow{r_1} D^*$

2. $D^* + P \xrightarrow{r_2} DP$

3. $D^* + C \xrightarrow{r_3} DC$

C

$$\frac{\partial[D]}{\partial t} = -r_1[D]S$$

$$\frac{\partial[D^*]}{\partial t} = r_1[D]S - r_2[D^*][P] - r_3[D^*][C]$$

$$\frac{\partial[P]}{\partial t} = -r_2[D^*][P]$$

$$\frac{\partial[DP]}{\partial t} = r_2[D^*][P]$$

$$\frac{\partial[C]}{\partial t} = -r_3[D^*][C]$$

$$\frac{\partial[DC]}{\partial t} = r_3[D^*][C]$$

1188

1189 **Figure 3 - figure supplement 3. Mws model (A)** Description of functional

1190 interactions **(B)** Description of molecular reaction **(C)** Differential equations

1191 describing the dynamics of the Aws pathway

1192

1193

1194 **Figure 4 - figure supplement 1. Relative contributions of reaction rates for Wsp**

1195

1196

1197

Figure 4 - figure supplement 2. Relative contributions of reaction rates for Aws.

Effect of the mutational hotspot on the relative contribution of each reaction in Aws

to the probability of generating a wrinkly spreader **(A)** Null model. **(B)** Hot spot in

AwsX increase rate mutation rate five times. Because the mutational hotspot increases

the likelihood that AwsX's reactions will be altered, r_2 and r_3 have an increased

probability of enabling or disabling change. As a consequence, r_2 shows the biggest

change as it contributes up to 70% (as opposed to 35% without the mutational

hotspot) of the probability that Aws is used to generate a WS in the area where

enabling change is more likely than disabling change. Furthermore, in the other area

of parameter space where disabling change is more likely, r_3 contributes up to 90% of

the probability that Aws is used with or without the hotspot.

1209

1210 **Supplementary files**

1211 Figure S1

1212 Figure 5 – source data 1. Table of all WS mutations in Wsp, Aws and Mws.

1213 ODE model and code

Figure S1. **Parameter sensitivity analysis.** To assess the effect of the chosen parameter (**A**) ranges on our results, we redid our sampling procedure for WSP for three different parameter regimes: (**B**) an expanded range for initial concentrations [0-50], (**C**) an expanded range for reaction rates $10^{[-3,3]}$, (**D**) a compressed range for mutational effect size $10^{[-1,1]}$. We found that our qualitative results are robust to these changes**.**

Figure 5 - Source data - Wsp mutations

Genome position	Type	Change	Gene positio	Gene locus	Gene symbol	Effect	Comment	Proposed molecular effect
1353597 - 1353602	Deletion	del 6 bp	1328 - 1333	PFLU1219	wspA	In frame del G443 - M444)	No homology	
1353112 - 1353195	Deletion	del 84 bp	843 - 926	PFLU1219	wspA	In frame del A281-A308	6 bp GGCCAC homology	methylation site
1353112 - 1353195	Deletion	del 84 bp	843 - 926	PFLU1219	wspA	In frame del A281-A308	6 bp GGCCAC homology	methylation site
1353140 - 1353160	Deletion	del 21 bp	878 - 898	PFLU1219	wspA	In frame del T293 - E299	5 bp ACTGA homology	methylation site
1353140 - 1353160	Deletion	del 21 bp	878 - 898	PFLU1219	wspA	In frame del T293 - E299	5 bp ACTGA homology	methylation site
1353423	Transversion	A->C	1154	PFLU1219	wspA	N385T		trimer-of-dimer association
1353451	Transversion	A->C	1182	PFLU1219	wspA	E394D		trimer-of-dimer association
1353451	Transversion	A->C	1182	PFLU1219	wspA	E394D		trimer-of-dimer association
1353411	Transition	C->T	1142	PFLU1219	wspA	A381V		trimer-of-dimer association
1353411	Transition	C->T	1142	PFLU1219	wspA	A381V		trimer-of-dimer association
1353528	Transition	C->T	1259	PFLU1219	wspA	A420V		possible trimer-of-dimer association
1353528	Transition	C->T	1259	PFLU1219	wspA	A420V		possible trimer-of-dimer association
1353323	Transition	G->A	1054	PFLU1219	wspA	A352T		possible trimer-of-dimer association
1353323	Transition	G->A	1054	PFLU1219	wspA	A352T		possible trimer-of-dimer association
1353408	Transversion	T->G	1139	PFLU1219	wspA	V380G		trimer-of-dimer association
1355373 - 1355707	Deletion	del 309 bp	973 - 51	PFLU1221	wspC/D	WspC M1-A324 fused to WspD D18-S232	9 bp ACCCTGGCC homology	increase wspC activity
1358267	Transversion	A->C	1916	PFLU1223	wspE	D639A		disrupt phoshorylation site of wspF
1358267	Transversion	A->C	1916	PFLU1223	wspE	D639A		disrupt phoshorylation site of wspF
1358267	Transversion	A->C	1916	PFLU1223	wspE	D639A		disrupt phoshorylation site of wspF
1358267	Transversion	A->C	1916	PFLU1223	wspE	D639A		disrupt phoshorylation site of wspF
1358267	Transversion	A->C	1916	PFLU1223	wspE	D639A		disrupt phoshorylation site of wspF
1358551	Transition	A->G	2200	PFLU1223	wspE	K734E		disrupt phoshorylation site of wspF
1358553	Transversion	A->C	2202	PFLU1223	wspE	K734N		disrupt phoshorylation site of wspF
1358553	Transversion	A->C	2202	PFLU1223	wspE	K734N		disrupt phoshorylation site of wspF
1358553	Transversion	A->C	2202	PFLU1223	wspE	K734N		disrupt phoshorylation site of wspF
1358270	Transition	C->T	1919	PFLU1223	wspE	S640L		disrupt phoshorylation site of wspF
1358266	Transversion	G->T	1915	PFLU1223	wspE	D639Y		disrupt phoshorylation site of wspF
1358279	Transversion	T->A	1928	PFLU1223	wspE	V643E		disrupt phoshorylation site of wspF
1358279	Transversion	T->A	1928	PFLU1223	wspE	V643E		disrupt phoshorylation site of wspF
1358766 - 1358780	Deletion	del 15 bp	151-165	PFLU1224	wspF	In frame del L51- I55	10 bp GGACCTGATC homology	disrupt demethylase activity
1358766 - 1358780	Deletion	del 15 bp	151-165	PFLU1224	wspF	In frame del L51- I55	10 bp GGACCTGATC homology	disrupt demethylase activity
1358810 -1358935	Deletion	del 126 bp	195 - 320	PFLU1224	wspF	In frame del R66 -L107	No homology	disrupt demethylase activity
1359285 - 1359289	Deletion	del 5 bp	670 - 674	PFLU1224	wspF	Frame shift after L223	3 bp GGC homology	disrupt demethylase activity
1359292 - 1359441	Deletion	del 150 bp	677 - 826	PFLU1224	wspF	In frame del T226-G275	3 bp ACC homology	disrupt demethylase activity
1359505	Transition	A->G	890	PFLU1224	wspF	Q297R		disrupt demethylase activity

1358755	Transition	C->T	140	PFLU1224	wspF	P47L		disrupt demethylase activity
1359091	Transition	C->T	476	PFLU1224	wspF	S159L		disrupt demethylase activity
1359171	Transition	C->T	556	PFLU1224	wspF	H186Y		disrupt demethylase activity
1359538	Transition	C->T	923	PFLU1224	wspF	P308L		disrupt demethylase activity
1359423	Transversion	G->C	808	PFLU1224	wspF	G270R		disrupt demethylase activity
1359604	Transversion	T->A	989	PFLU1224	wspF	L331*		disrupt demethylase activity
1360156 - 1360167	Deletion	del 12 bp	481 - 492	PFLU1225	wspR	In frame del N161 - L164	No homology	disrupt requirement for phosphorylation
1360099	Transversion	C->A	424	PFLU1225	wspR	Q142K		disrupt requirement for phosphorylation
1360102	Transition	C->T	427	PFLU1225	wspR	R143W		disrupt requirement for phosphorylation
1360102	Transition	C->T	427	PFLU1225	wspR	R143W		disrupt requirement for phosphorylation
1360107	Transversion	T->G	432	PFLU1225	wspR	D144E		disrupt requirement for phosphorylation

Figure 5 - Source data - Aws mutations

Genome position	Type	Change	Gene position	Gene locus	Gene symbol	Effect	Comment	Proposed molecular effect
5705729	Transversion	A->C	101	PFLU5209	awsO	Q34P		change binding with awsX
5705711	Transversion	A->C	119	PFLU5209	awsO	Q40P		change binding with awsX
5706522	Transition	C->T	575	PFLU5210	awsR	A192V		HAMP linker
5706522	Transition	C->T	575	PFLU5210	awsR	A192V		HAMP linker
5706909	Transition	C->T	188	PFLU5210	awsR	A63V		interaction awsX
5706441	Transversion	A->C	656	PFLU5210	awsR	D219A		HAMP linker
5706937	Transition	C->T	160	PFLU5210	awsR	R54C		interaction awsX
5707018	Transversion	A->C	79	PFLU5210	awsR	T27P		disrupt TM helix
5707018	Transversion	A->C	79	PFLU5210	awsR	T27P		disrupt TM helix
5707018	Transversion	A->C	79	PFLU5210	awsR	T27P		disrupt TM helix
5707018	Transversion	A->C	79	PFLU5210	awsR	T27P		disrupt TM helix
5707018	Transversion	A->C	79	PFLU5210	awsR	T27P		disrupt TM helix
5707018	Transversion	A->C	79	PFLU5210	awsR	T27P		disrupt TM helix
5707018	Transversion	A->C	79	PFLU5210	awsR	T27P		disrupt TM helix
5707018	Transversion	A->C	79	PFLU5210	awsR	T27P		disrupt TM helix
5707018	Transversion	A->C	79	PFLU5210	awsR	T27P		disrupt TM helix
5707574	Transition	C->T	92	PFLU5211	awsX	A31V		disrupt awsX function
5707528-5707566	Deletion	del 39 bp	100-138	PFLU5211	awsX	del P34-A46	10 bp homology CGCCCAGGCG	disrupt awsX function
5707528-5707566	Deletion	del 39 bp	100-138	PFLU5211	awsX	del P34-A46	10 bp homology CGCCCAGGCG	disrupt awsX function
5707460	Transversion	T->G	206	PFLU5211	awsX	L69R		disrupt awsX function
5707405-5707437	Deletion	del 33 bp	229-261	PFLU5211	awsX	del Y77-Q87	6 bp homology ACCCAG	disrupt awsX function
5707405-5707437	Deletion	del 33 bp	229-261	PFLU5211	awsX	del Y77-Q87	6 bp homology ACCCAG	disrupt awsX function
5707405-5707437	Deletion	del 33 bp	229-261	PFLU5211	awsX	del Y77-Q87	6 bp homology ACCCAG	disrupt awsX function
5707405-5707437	Deletion	del 33 bp	229-261	PFLU5211	awsX	del Y77-Q87	6 bp homology ACCCAG	disrupt awsX function

Genome position	Type	Change	Gene position	Gene locus	Gene symbol	Effect	Comment	Proposed molecular effect
5707405-5707437	Deletion	del 33 bp	229-261	PFLU5211	awsX	del Y77-Q87	6 bp homology ACCCAG	disrupt awsX function
5707405-5707437	Deletion	del 33 bp	229-261	PFLU5211	awsX	del Y77-Q87	6 bp homology ACCCAG	disrupt awsX function
5707405-5707437	Deletion	del 33 bp	229-261	PFLU5211	awsX	del Y77-Q87	6 bp homology ACCCAG	disrupt awsX function
5707405-5707437	Deletion	del 33 bp	229-261	PFLU5211	awsX	del Y77-Q87	6 bp homology ACCCAG	disrupt awsX function
5707405-5707437	Deletion	del 33 bp	229-261	PFLU5211	awsX	del Y77-Q87	6 bp homology ACCCAG	disrupt awsX function
5707405-5707437	Deletion	del 33 bp	229-261	PFLU5211	awsX	del Y77-Q87	6 bp homology ACCCAG	disrupt awsX function
5707405-5707437	Deletion	del 33 bp	229-261	PFLU5211	awsX	del Y77-Q87	6 bp homology ACCCAG	disrupt awsX function
5707405-5707437	Deletion	del 33 bp	229-261	PFLU5211	awsX	del Y77-Q87	6 bp homology ACCCAG	disrupt awsX function
5707405-5707437	Deletion	del 33 bp	229-261	PFLU5211	awsX	del Y77-Q87	6 bp homology ACCCAG	disrupt awsX function
5707405-5707437	Deletion	del 33 bp	229-261	PFLU5211	awsX	del Y77-Q87	6 bp homology ACCCAG	disrupt awsX function
5707405-5707437	Deletion	del 33 bp	229-261	PFLU5211	awsX	del Y77-Q87	6 bp homology ACCCAG	disrupt awsX function
5707405-5707437	Deletion	del 33 bp	229-261	PFLU5211	awsX	del Y77-Q87	6 bp homology ACCCAG	disrupt awsX function
5707405-5707437	Deletion	del 33 bp	229-261	PFLU5211	awsX	del Y77-Q87	6 bp homology ACCCAG	disrupt awsX function
5707405-5707437	Deletion	del 33 bp	229-261	PFLU5211	awsX	del Y77-Q87	6 bp homology ACCCAG	disrupt awsX function
5707405-5707437	Deletion	del 33 bp	229-261	PFLU5211	awsX	del Y77-Q87	6 bp homology ACCCAG	disrupt awsX function
5707405-5707437	Deletion	del 33 bp	229-261	PFLU5211	awsX	del Y77-Q87	6 bp homology ACCCAG	disrupt awsX function
5707262	Transversion	T->G	404	PFLU5211	awsX	L135R		disrupt awsX function

Figure 5 - Source data - Mws mutations

Genome position	Type	Change	Gene position	Gene locus	Gene symbol	Effect	Comment	Proposed molecular effect
5857733-5857741	Deletion	del 9 bp	3071-3079	PFLU5329	mwsR	del R1024 - E1026	8 bp homology CCTGGAGC	Disrupt interaction DGC and EAL domain
5857733-5857741	Deletion	del 9 bp	3071-3079	PFLU5329	mwsR	del R1024 - E1026	8 bp homology CCTGGAGC	Disrupt interaction DGC and EAL domain
5857733-5857741	Deletion	del 9 bp	3071-3079	PFLU5329	mwsR	del R1024 - E1026	8 bp homology CCTGGAGC	Disrupt interaction DGC and EAL domain
5857733-5857741	Deletion	del 9 bp	3071-3079	PFLU5329	mwsR	del R1024 - E1026	8 bp homology CCTGGAGC	Disrupt interaction DGC and EAL domain
5857733-5857741	Deletion	del 9 bp	3071-3079	PFLU5329	mwsR	del R1024 - E1026	8 bp homology CCTGGAGC	Disrupt interaction DGC and EAL domain
5857733-5857741	Deletion	del 9 bp	3071-3079	PFLU5329	mwsR	del R1024 - E1026	8 bp homology CCTGGAGC	Disrupt interaction DGC and EAL domain
5857733-5857741	Deletion	del 9 bp	3071-3079	PFLU5329	mwsR	del R1024 - E1026	8 bp homology CCTGGAGC	Disrupt interaction DGC and EAL domain
5857733-5857741	Deletion	del 9 bp	3071-3079	PFLU5329	mwsR	del R1024 - E1026	8 bp homology CCTGGAGC	Disrupt interaction DGC and EAL domain
5857733-5857741	Deletion	del 9 bp	3071-3079	PFLU5329	mwsR	del R1024 - E1026	8 bp homology CCTGGAGC	Disrupt interaction DGC and EAL domain
5857733-5857741	Deletion	del 9 bp	3071-3079	PFLU5329	mwsR	del R1024 - E1026	8 bp homology CCTGGAGC	Disrupt interaction DGC and EAL domain
5857733-5857741	Deletion	del 9 bp	3071-3079	PFLU5329	mwsR	del R1024 - E1026	8 bp homology CCTGGAGC	Disrupt interaction DGC and EAL domain
A5857192C	Transversion	A->C	A2530C	PFLU5329	mwsR	I844L		
C5857598/ C5857599	Insertion	duplication 2	2937-2959	PFLU5329	mwsR	dup I978-G985		Disrupt interaction DGC and EAL domain
G5857440A	Transition	G->A	G2778A	PFLU5329	mwsR	M926I		Disrupt interaction M992
G5857440A	Transition	G->A	G2778A	PFLU5329	mwsR	M926I		Disrupt interaction M926
G5857440A	Transition	G->A	G2778A	PFLU5329	mwsR	M926I		Disrupt interaction M926
G5857638A	Transition	G->A	G2976A	PFLU5329	mwsR	M992I		Disrupt interaction M926
G5857909A	Transition	G->A	G3247A	PFLU5329	mwsR	E1083K		Disrupt interaction DGC and EAL domain

Differential equations for WSP, AWS, & MWS pathways

WSP

$$\frac{d[ABD]}{dt} = r_2 F^*[ABDm] - r_1[ABD][Cm]$$

$$\frac{d[ABDm]}{dt} = r_1[ABD][Cm] - r_2 F^*[ABDm] - r_3 S[ABDm] + r_4 E[ABDm^*]$$

$$\frac{d[ABDm^*]}{dt} = -r_4 E[ABDm^*] + r_3 S[ABDm]$$

$$\frac{dE^*}{dt} = r_4 E[ABDm^*] - r_5 E^* R - r_6 E^* F$$

$$\frac{dR^*}{dt} = r_5 R E^*$$

$$\frac{dE}{dt} = r_6 E^* F - r_4 E[ABDm^*] + r_5 R E^*$$

$$\frac{dF}{dt} = -r_6 E^* F + r_2 F^*[ABDm]$$

$$\frac{dF^*}{dt} = r_6 E^* F - r_2 F^*[ABDm]$$

$$\frac{dR}{dt} = -r_5 R E^* - .01R$$

AWS

$$\frac{dX}{dt} = -r_2 X O^* - r_3 X R$$

$$\frac{d[XR]}{dt} = r_3 X R$$

$$\frac{dO}{dt} = -r_1 S O$$

$$\frac{dO^*}{dt} = r_1 S O - r_2 X O^*$$

$$\frac{d[OX]}{dt} = r_2 O^* X$$

$$\frac{dR}{dt} = -r_3 X R - r4 R R$$

$$\frac{d[RR]}{dt} = r_4 R R$$

MWS

$$
\begin{aligned}
\frac{dG}{dt} &= -r_1 G S \\
\frac{dG^*}{dt} &= r_1 G S - r_2 G^* E - r_3 G^* C \\
\frac{dE}{dt} &= -r_2 G^* E \\
\frac{d[GE]}{dt} &= r_2 G^* E \\
\frac{dC}{dt} &= -r_3 G^* C \\
\frac{d[GC]}{dt} &= r3 G^* C
\end{aligned}
$$

Julia code for WSP, AWS, and MWS differential equations

The following three functions implement the differential equation model (ODE model) for the WSP, AWS, and MWS pathways.

```
using ODE,StatsBase,MAT,HDF5,JLD

# code for WSP differential equations
function lindodeWSP(t,y)
   # convert y to reactants for ease of reading
   ABD=y[1];
   ABDm=y[2];
   ABDmp=y[3];
   Ep=y[4];
   Rp=y[5];
   E=y[6];
   F=y[7];
   Fp=y[8];
   R=y[9];
   # pull reaction rates from rs variable
   r1=rs[1];r2=rs[2];r3=rs[3];r4=rs[4];r5=rs[5];r6=rs[6];
   # compute derivatives, i.e. y'
   yp=zeros(size(y));
   yp[1]=r2*Fp*ABDm-r1*ABD*Cm; # dABD/dt
   yp[2]=r1*ABD*Cm-r2*Fp*ABDm-r3*S*ABDm+r4*E*ABDmp; # dABDm/dt
   yp[3]=-r4*E*ABDmp + r3*S*ABDm ; # dABDmp/dt
   yp[4]=r4*E*ABDmp - r5*Ep*R -r6*Ep*F ; # dEp/dt
   yp[5]=r5*R*Ep ; # dRp/dt
   yp[6]=r6*Ep*F-r4*E*ABDmp+r5*R*Ep ; # dE/dt
   yp[7]=-r6*Ep*F +r2*Fp*ABDm ; # dF/dt
   yp[8]=r6*Ep*F -r2*Fp*ABDm ; # dFp/dt
   yp[9]=-r5*R*Ep-.01*R; # dR/dt
   return yp
end

# code for AWS differential equations
```

```
function lindodeAWS(t,y)
    # convert y to reactants for ease of reading
    X=y[1];
    XR=y[2];
    O=y[3];
    Op=y[4];
    OX=y[5];
    R=y[6];
    RR=y[7];
    # pull reaction rates from rs variable
    r1=rs[1];r2=rs[2];r3=rs[3];r4=rs[4];
    # compute derivatives, i.e. y'
    yp=zeros(size(y));
    yp[1]=-r2*X*Op-r3*X*R; # dX/dt
    yp[2]=r3*X*R; # dXR/dt
    yp[3]=-r1*S*O; # dO/dt
    yp[4]=r1*S*O-r2*X*Op; # dOp/dt
    yp[5]=r2*Op*X; # dOX/dt
    yp[6]=-r3*X*R-r4*R*R; # dR/dt
    yp[7]=r4*R*R; # dRR/dt
    return yp;
end

# code for MWS differential equations
function lindodeMWS(t,y)
    # convert y to reactants for ease of reading
    G=y[1];
    Gp=y[2];
    E=y[3];
    GE=y[4];
    C=y[5];
    GC=y[6];
    # pull reaction rates from rs variable
    r1=rs[1];r2=rs[2];r3=rs[3];
    # compute derivatives, i.e. y'
    yp=zeros(size(y));
    yp[1]=-r1*G*S; # dG/dt
    yp[2]=r1*G*S-r2*Gp*E-r3*Gp*C;# dGp/dt
    yp[3]= -r2*Gp*E;# dE/dt
    yp[4]= r2*Gp*E;# dGE/dt
    yp[5]=-r3*Gp*C;# dC/dt
    yp[6]=r3*Gp*C; # dGC/dt
    return yp;
end
```

Julia code for running differential equation solvers

We include the code for implementing our Bayesian sampling method. The colored sections correspond to statements that make it specific to WSP (blue), AWS (red), or MWS (green). It was run in julia version 0.4.3.

```
# Variables for reactions
numrxns=6; # number of reaction rates for WSP
```

```
numrxns=4; # number of reaction rates for AWS
numrxns=3; # number of reaction rates for MWS
rs=rand(numrxns); # establish variable scope, will be reaction rates later
rs_save=copy(rs); # establish variable scope, will be a saved version of reaction rates later
totnumruns=3.^length(rs); # all possible combinations for reaction rates (down,nothing, up)
v=zeros(length(rs)); # establish variable scope (used to alter reaction rates)
S=0;Cm=0; # initialize constants used in differential equations
indexWS=5; # reactant corresponding to WS in WSP
indexWS=7; # reactant corresponding to WS in AWS
indexWS=6; # reactant corresponding to WS in MWS

# Variables for running ODE solver
testnums=1000; # number of runs
yorig=zeros(testnums); # storage for baseline WS production
yout=0; # establish variable scope
tf=1.0; # establish variable scope
tftimes=1.0; # establish variable scope

numreactants=9; # number of reactants
numreactants=7; # number of reactants
numreactants=6; # number of reactants
init=10*rand(numreactants); # establish variable scope

# Variables for storing data
res=-1*ones(totnumruns,testnums); # storage for altered WS production
numfinished=1; # counter for runs completed

# Code for Bayesian sampling method
while numfinished<=testnums
    done=0;
    try
        println(numfinished) # keeps track of how many sims have been done
        # Sample concentrations and rates to establish a baseline amount of WS production
        rs=10.^(4*rand(numrxns)-2); # sample reaction rates from [.01,100]
        rs_save=copy(rs); # saved copy as a reference when altering later
        init=10*rand(numreactants); # sample initial concentrations for reactants from [0,10]
        S=10*rand(); # sample initial concentration for constant reactant of signal (S) from [0,10]
        Cm=10*rand(); # sample initial concentration for constant reactant Cm from [0,10]

        # ODE solver for baseline
        tf=1.0; # initial time for ode solver
        dst=100; # initial distance, used to determine solution converged
        tol=10^(-8.0); # tolerance for ODE solver
        while dst>tol
            tout, yout = ode45(lindodeWSP, init, [0.0 ,tf]);
            tout, yout = ode45(lindodeAWS, init, [0.0 ,tf]);
            tout, yout = ode45(lindodeMWS, init, [0.0 ,tf]);
            dst=sum((yout[end-1]-yout[end]).^2); # Euclidean distance in final step of solution
            yorig[numfinished]=yout[end][indexWS]; # reactant corresponding to WS
            tftimes=tout[end]; # final time
            tf*=2;
        end
        done=1; # successful completion of try loop
    end
    if done==1 # baseline WS production is established, now sample changes/mutations
        i0=1; # counter for completed changes
        cct=0; # counter for total number of attempts
```

```
while i0<=totnumruns;
    cct+=1;
    println([numfinished cct]) # report status for tracking progress
    # Alter reaction rates
    num=i0-1; # used for determining which rates change down/none/up
        for i1=length(rs)-1:-1:0;
                v[i1+1]=floor(num/(3^i1)); # v is num into base 3 number
                num=num-v[i1+1]*3^i1;
        end
    v+=1;
    facs=[10.^(-2*rand()), 1, 10.^(2*rand())]; # factors to alter rxn rates [.01,1] down, 1 none, [1,100] up
    for i1=1:length(rs)
        rs[i1]=facs[v[i1]]*rs_save[i1]; # alter reaction rates
    end
    try
        tout, yout = ode45(lindodeWSP, init, [0.0,tftimes]);
        tout, yout = ode45(lindodeAWS, init, [0.0,tftimes]);
        tout, yout = ode45(lindodeMWS, init, [0.0,tftimes]);
        if abs(tout[end]-tftimes)<.01 # ODE solver finished
                res[i0,numfinished]=yout[end][indexWS]; # store amount of WS produced
            i0+=1
        end
    end
    if cct>10000
        i0=2*totnumruns; # baseline and sampling occurred in space with poorly conditioned ODEs, try again
    end
    end
    if i0<2*totnumruns # successful
        numfinished+=1;
        # Save data to a file for checking in MATLAB
        file=matopen("pathway_results_temp.mat","w")
        write(file,"res",res); altered WS production
        write(file,"yorig",yorig); baseline WS production
        close(file);
    end
    end
end

# Save data to a file for processing in MATLAB
file=matopen("pathway_results_complete_WSP.mat","w")
file=matopen("pathway_results_complete_AWS.mat","w")
file=matopen("pathway_results_complete_MWS.mat","w")
write(file,"res",res); altered WS production
write(file,"yorig",yorig); baseline WS production
close(file);
```

MATLAB code for interpreting saved data WSP vs MWS

This code shows how the data from the julia code is analyzed and transformed into the contour plots shown in the paper.

```
% create record of how parameters change down, none, up for WSP
rs1=rand(1,6);
```

```matlab
totnumruns=3.^length(rs1);
paramsWSP=zeros(totnumruns,length(rs1));
v=zeros(size(rs1));
i0=1;
while i0<=totnumruns;
    num=i0-1;
    for i1=length(rs1)-1:-1:0;
            v(i1+1)=floor(num/(3^i1));
            num=num-v(i1+1)*3^i1;
    end
    v=v+1;
    paramsWSP(i0,:)=v;
    i0=i0+1;
end
% create record of how parameters change down, none, up for MWS
rs1=rand(1,3);
totnumruns=3.^length(rs1);
paramsMWS=zeros(totnumruns,length(rs1));
v=zeros(size(rs1));
i0=1;
while i0<=totnumruns;
    num=i0-1;
    for i1=length(rs1)-1:-1:0;
            v(i1+1)=floor(num/(3^i1));
            num=num-v(i1+1)*3^i1;
    end
    v=v+1;
    paramsMWS(i0,:)=v;
    i0=i0+1;
end

% Load data
load pathway_results_complete_MWS.mat
resMWS=res;
yorigMWS=yorig';
clear res yorig
load pathway_results_complete_WSP.mat
resWSP=res;
yorigWSP=yorig';

% Variables and data storage to compare likelihood of pathways
numsampWSP=size(resWSP,2); % in case want to use fewer samples
numsampMWS=size(resMWS,2); % in case want to use fewer samples

perange=10.^[-7:.5:-1]; % range for probability enabling mutations
pdrange=10.^[-7:.5:-1]; % range for probability disabling mutations
pemat=zeros(length(perange),length(pdrange)); % matrix for plotting data and reference
pdmat=zeros(size(pemat)); % matrix for plotting data and reference
psummatWSP=zeros(size(pemat)); % matrix for probability WSP used
psummatMWS=zeros(size(pemat)); % matrix for probability MWS used
tol=0;
% Code to compare likelihood of pathways
for i0=1:length(perange)
    for j0=1:length(pdrange)
        % Retrieve probabilities
        pe=perange(i0);
        pd=pdrange(j0);
        pemat(i0,j0)=pe;
```

```matlab
        pdmat(i0,j0)=pd;
        % WSP computation
        pmatforr=ones(6,1)*[pd 1-pe-pd pe];
        psum=0; % total sum of (prob of rxn changes) X (number of times WS produced)
        for i1=1:size(resWSP,1)
            probevent=1; % initialize, probability to get combination of down, none, up for rxns
            for j1=1:6;
                probevent=probevent*pmatforr(j1,paramsWSP(i1,j1)); % multiply by prob of each change
                    end
            psum=psum+probevent*sum(resWSP(i1,1:numsampWSP)>yorigWSP(1:numsampWSP)+tol)/numsampWSP;
        end
        psummatWSP(i0,j0)=psum;

        % MWS computation
        pmatforr=ones(3,1)*[pd 1-pe-pd pe];
        psum=0; % total sum of (prob of rxn changes) X (number of times WS produced)
        for i1=1:size(resMWS,1)
            probevent=1; % initialize, probability to get combination of down, none, up for rxns
            for j1=1:3;
                probevent=probevent*pmatforr(j1,paramsMWS(i1,j1)); % multiply by prob of each change
            end
            psum=psum+probevent*sum(resMWS(i1,1:numsampMWS)>yorigMWS(1:numsampMWS)+tol)/numsampMWS;
        end
        psummatMWS(i0,j0)=psum;
    end
end

% Plot data
close all
figure
contourf(pemat,pdmat,log2(psummatWSP./psummatMWS),400,'LineStyle','None')
set(gca,'xScale','log','yScale','log','TickLength',[.025 .025],'LineWidth',3);
set(gca,'FontSize',18,'xTick',10.^[-7:1:-1],'yTick',10.^[-7:1:-1]);
xlabel('Probability of enabling change','FontSize',24);
ylabel('Probability of disabling change','FontSize',24);
c=colorbar('FontSize',18);
c.Label.String='log_2 ratio probability WSP/MWS';
colormap jet;
axis square
eval(['print -f1 -depsc -r300 WSP_vs_MWS_contour.eps']);
```

MATLAB code for interpreting saved data WSP vs AWS

```matlab
% create record of how parameters change down, none, up for WSP
rs1=rand(1,6);
totnumruns=3.^length(rs1);
paramsWSP=zeros(totnumruns,length(rs1));
v=zeros(size(rs1));
i0=1;
while i0<=totnumruns;
    num=i0-1;
    for i1=length(rs1)-1:-1:0;
            v(i1+1)=floor(num/(3^i1));
```

```matlab
                num=num-v(i1+1)*3^i1;
        end
    v=v+1;
    paramsWSP(i0,:)=v;
    i0=i0+1;
end
% create record of how parameters change down, none, up for AWS
rs1=rand(1,4);
totnumruns=3.^length(rs1);
paramsAWS=zeros(totnumruns,length(rs1));
v=zeros(size(rs1));
i0=1;
while i0<=totnumruns;
    num=i0-1;
    for i1=length(rs1)-1:-1:0;
            v(i1+1)=floor(num/(3^i1));
            num=num-v(i1+1)*3^i1;
    end
    v=v+1;
    paramsAWS(i0,:)=v;
    i0=i0+1;
end

% Load data
load pathway_results_complete_AWS.mat
resAWS=res;
yorigAWS=yorig';
clear res yorig
load pathway_results_complete_WSP.mat
resWSP=res;
yorigWSP=yorig';

% Variables and data storage to compare likelihood of pathways
numsampWSP=size(resWSP,2); % in case want to use fewer samples
numsampAWS=size(resAWS,2); % in case want to use fewer samples

perange=10.^[-7:.5:-1]; % range for probability enabling mutations
pdrange=10.^[-7:.5:-1]; % range for probability disabling mutations
pemat=zeros(length(perange),length(pdrange)); % matrix for plotting data and reference
pdmat=zeros(size(pemat)); % matrix for plotting data and reference
psummatWSP=zeros(size(pemat)); % matrix for probability WSP used
psummatAWS=zeros(size(pemat)); % matrix for probability MWS used
tol=0;
fac=5; % factor increase of mutation because of hotspot

% Code to compare likelihood of pathways
for i0=1:length(perange)
    for j0=1:length(pdrange)
        % Retrieve probabilities
        pe=perange(i0);
        pd=pdrange(j0);
        pemat(i0,j0)=pe;
        pdmat(i0,j0)=pd;
        % WSP computation
        pmatforr=ones(6,1)*[pd 1-pe-pd pe];
        psum=0; % total sum of (prob of rxn changes) X (number of times WS produced)
        for i1=1:size(resWSP,1)
                probevent=1; % initialize, probability to get combination of down, none, up for rxns
```

```matlab
            for j1=1:6;
                probevent=probevent*pmatforr(j1,paramsWSP(i1,j1)); % multiply by prob of each change
            end
            psum=psum+probevent*sum(resWSP(i1,1:numsampWSP)>yorigWSP(1:numsampWSP)+tol)/numsampWSP;
        end
        psummatWSP(i0,j0)=psum;

        % AWS computation
        pmatforr=ones(4,1)*[pd 1-pe-pd pe];
        pmatforr(4,:)=[.5*pd 1-.5*pd-.5*pe .5*pe]; % because only reactant in dimerization
        pmatforr(3,:)=[fac*pd 1-fac*pd-fac*pe fac*pe]; % effect of hotspot
        pmatforr(2,:)=[fac*pd 1-fac*pd-fac*pe fac*pe]; % effect of hotspot
        psum=0; % total sum of (prob of rxn changes) X (number of times WS produced)
        for i1=1:size(resAWS,1)
            probevent=1; % initialize, probability to get combination of down, none, up for rxns
            for j1=1:4;
                probevent=probevent*pmatforr(j1,paramsAWS(i1,j1)); % multiply by prob of each change
            end
            psum=psum+probevent*sum(resAWS(i1,1:numsampAWS)>yorigAWS(1:numsampAWS)+tol)/numsampAWS;
        end
        psummatAWS(i0,j0)=psum;
    end
end

% Plot data
close all
figure
contourf(pemat,pdmat,log2(psummatWSP./psummatAWS),400,'LineStyle','None')
set(gca,'xScale','log','yScale','log','TickLength',[.025 .025],'LineWidth',3);
set(gca,'FontSize',18,'xTick',10.^[-7:1:-1],'yTick',10.^[-7:1:-1]);
xlabel('Probability of enabling change','FontSize',24);
ylabel('Probability of disabling change','FontSize',24);
c=colorbar('FontSize',18);
c.Label.String='log_2 ratio probability WSP/AWS';
colormap jet;
axis square
eval(['print -f1 -depsc -r300 WSP_vs_AWS_contour.eps']);
```

9