The ecology of the Chloroflexi in full-scale activated sludge wastewater treatment plants

Marta Nierychlo[1], Aleksandra Miłobędzka[2,3], Francesca Petriglieri[1], Bianca
McIlroy[1], Per Halkjær Nielsen[1], and Simon Jon McIlroy[1§]*

[1]Center for Microbial Communities, Department of Chemistry and Bioscience,
Aalborg University, Aalborg, Denmark
[2]Microbial Ecology and Environmental Biotechnology Department, Institute of
Botany, Faculty of Biology, University of Warsaw; Biological and Chemical
Research Centre, Żwirki i Wigury 101, Warsaw 02-089, Poland
[3]Department of Biology, Faculty of Building Services, Hydro and Environmental
Engineering, Warsaw University of Technology, 00-653 Warsaw, Poland

* Corresponding author: Simon Jon McIlroy, Center for Microbial Communities,
Department of Chemistry and Bioscience, Aalborg University, Fredrik Bajers Vej 7H,
DK-9220 Aalborg, Denmark; Tel.: +45 9940 3573; Fax: +45 9814 1808; Email:
s.mcilroy@uq.edu.au
§ Present address: Australian Centre for Ecogenomics, University of Queensland,
Australia

Abstract
Filamentous bacteria belonging to the phylum Chloroflexi have received considerable
attention in wastewater treatment systems for their suggested role in operational
problem of impaired sludge settleability known as bulking. Their consistently high
abundance in full-scale systems, even in the absence of bulking, indicates that they
make a substantial contribution to the nutrient transformations during wastewater
treatment. In this study, extensive 16S rRNA amplicon surveys of full-scale Danish
WWTPs were screened to identify the most numerically important Chloroflexi
genera. Fluorescence *in situ* hybridization probes were designed for their *in situ*
characterization. All abundant phylotypes of the phylum were identified as facultative
anaerobic chemoorganotrophs involved in fermentation of sugars. These groups were
all filamentous but differed in their morphology and spatial arrangement. '*Candidatus*
Villogracilis' was predominantly located within the activated sludge flocs, where they
possibly have structural importance, and their abundance was relatively stable.
Conversely, the abundance of '*Candidatus* Amarilinum' was highly dynamic, relative
to other genera, sometimes reaching abundances in excess of 30% of the biovolume,
suggesting their likely role in bulking episodes. This study gives an important insight
into the role of Chloroflexi in WWTPs, thus contributing to the broader goal of
understanding the ecology of these biotechnologically important systems.

Keywords
Activated sludge; Filamentous bulking; Chloroflexi; '*Candidatus* Amarilinum', FISH;
Microautoradiography

Introduction

Members of the phylum Chloroflexi constitute a substantial proportion of the activated sludge community in full-scale activated sludge wastewater treatment plants (WWTPs), where they reportedly constitute up to 30% of the biovolume and often make up the majority of filamentous bacteria present (Beer *et al.* 2006; Morgan-Sagastume, Nielsen and Nielsen 2008; Mielczarek *et al.* 2012). Filamentous bacteria are generally believed to have structural importance for activated sludge flocs with good settling properties. However, overgrowth of certain filamentous species is associated with open and diffuse flocs as well as interfloc bridging, leading to a sometimes severe operational problem known as bulking (Wanner, Kragelund and Nielsen 2010). Some Chloroflexi species have also been proposed to be involved in the stabilization of problematic foams at WWTPs (Kragelund *et al.* 2011) and membrane fouling in membrane bioreactors (MBRs) (Ziegler *et al.* 2016). In addition, the high abundance of the Chloroflexi in wastewater treatment systems, often in the absence of bulking problems, indicates that they make a proportional contribution to the observed nutrient transformations of these systems. As such, the study of this phylum has implications for plant operation and our general understanding of the ecology of wastewater treatment.

Relatively little is known about the ecology of the Chloroflexi in nutrient removal WWTPs. *In situ* studies reveal a preference for the uptake of sugars, and a high level of surface associated hydrolytic enzymes indicates their involvement in the breakdown of complex organics (Kragelund *et al.* 2007, 2011; Xia, Kong and Nielsen 2007). Due to poor phylogenetic annotation of the routinely applied taxonomic databases, studies of activated sludge community dynamics often consider members of the Chloroflexi phylum as a whole - an approach that ignores the likely phenotypic diversity among its members (McIlroy *et al.* 2015). An understanding of the ecology of the phylum and how it relates to system function requires the identification and *in situ* characterization of the abundant genus-level phylotypes present.

Historically, the identification of bulking filaments has relied on classification keys based on morphological characteristics, with most morphotypes identified by an Eikelboom number (Eikelboom 2000; Jenkins, Richard and Daigger 2004). More recent phylogenetic identification of these filaments has relied on 16S rRNA gene-based clone library analyses coupled with fluorescent *in situ* hybridization (FISH) of plants with severe filamentous bulking. Most of the antecedent morphotypes are possessed by members of the Chloroflexi and include: 1851 (Beer *et al.* 2002); 0092 (Speirs *et al.* 2009); 0803 (Kragelund *et al.* 2011; Speirs, Tucci and Seviour 2015); 0914 (Speirs *et al.* 2011); 0041/0675 (Speirs *et al.* 2017), and several others (Kragelund *et al.* 2009) - noting that organisms possessing the same filamentous morphotype are often unrelated (Seviour *et al.* 1997; Speirs, Tucci and Seviour 2015), and closely related organisms can also possess different morphotypes (Speirs *et al.* 2017). FISH probes available for known phylotypes reportedly cover many of the Chloroflexi present, but there is still a substantial portion of the phylum without genus-level probes (up to 90% by FISH in some plants) (Kragelund *et al.* 2011). The recent extensive MiDAS 16S rRNA gene amplicon sequencing-based survey,

covering >50 Danish full-scale WWTPs over a 10 year period, has given a
comprehensive overview of the abundant core members of the full-scale activated
sludge treatment plants (McIlroy *et al.* 2015). As such, we are now able to
systematically target the numerically important Chloroflexi phylotypes. The *in situ*
physiology of some of these groups, such as 'Candidatus Promineofilum' (the B45
group) (McIlroy *et al.* 2016) and P2CN44 (Kragelund *et al.* 2011), has been
determined, but several abundant phylotypes are novel and known only by their 16S
rRNA gene sequence (McIlroy *et al.* 2015).
The aim of this study is to determine the *in situ* physiology of selected
abundant novel Chloroflexi activated sludge phylotypes. The extensive MiDAS
survey of full-scale Danish nutrient removal activated sludge plants was used to
identify the most abundant genus-level taxa belonging to the phylum Chloroflexi.
FISH probes were designed for these phylotypes and applied in combination with
microautoradiography (MAR) and histochemical staining for their *in situ*
characterization.

Methods
Biomass sampling and fixation for FISH
Biomass samples from the aerobic stage of selected full-scale activated sludge
WWTPs with nutrient removal were fixed with 4% paraformaldehyde (PFA) for 3 h
at 4°C. After fixation, samples were washed 3 times in sterile filtered tap water, re-
suspended in 50% ethanol in 1 x PBS solution [v/v], and stored at -20 °C. For basic
operational information for WWTPs sampled in this study, see Mielczarek *et al.*,
(2013).

Phylogenetic analysis and FISH probe design
Phylogenetic analysis and FISH probe design were performed with the ARB software
package (Ludwig *et al.* 2004) with the MiDAS database (Release 2.1), which is a
version of the SILVA database (Release 123 NR99) (Quast *et al.* 2013) curated for
activated sludge and anaerobic digester sequences (McIlroy *et al.* 2015, 2017b).
Potential probes were assessed *in silico* with the mathFISH software (Yilmaz,
Parnerkar and Noguera 2011). The Ribosomal Database Project (RDP) PROBE
MATCH function (Cole *et al.* 2014) was used to identify non-target sequences with
indels (McIlroy *et al.* 2011). Probe validation and optimization were based on
generated formamide dissociation curves (Daims, Stoecker and Wagner 2005), where
average relative fluorescent intensities of at least 50 cells calculated with ImageJ
software (National Institutes of Health, Maryland, USA) were measured for varied
hybridization buffer formamide concentration in increments of 5% (v/v) over a range
of 0-65% (v/v) (data not shown). Details for probes designed in this study have been
deposited in the ProbeBase database (Greuter *et al.* 2016).

FISH
FISH was performed as detailed by Daims *et al.,* (2005) using the probes designed in
this study as well as CFX197 and CFX223, targeting 'Ca. Promineofilum' (Speirs *et*

al. 2009); CFX1223 (Björnsson *et al.* 2002), and GNSB941 (Gich, Garcia-Gil and
Overmann 2001), applied as a mix to target the phylum Chloroflexi; EUB-338-I,
EUB338-II, and EUB338- III (Amann *et al.* 1990; Daims *et al.* 1999), applied as a
mix (EUBmix) to cover all bacteria; NON-EUB as a negative control for
hybridization (Wallner, Amann and Beisker 1993). The hybridization conditions
applied for each probe are given in **Table 1** or as recommended in their original
publications. Quantitative FISH (qFISH) biovolume fractions of individual
Chloroflexi genera were calculated as a percentage area of the total biovolume,
hybridizing the EUBmix probes, that also hybridizes with the specific probe. qFISH
analyses were based on 25 fields of view taken at 630 x magnification using the
Daime image analysis software (Daims, Lücker and Wagner 2006). Microscopic
analysis was performed with an Axioskop epifluorescence microscope (Carl Zeiss,
Oberkochen, Germany), an LSM510 Meta laser scanning confocal microscope (Carl
Zeiss), and a white light laser confocal microscope (Leica TCS SP8 X).

Morphological classification
Microscopic observations of wet mount preparations and Gram and Neisser staining
were performed according to the methods of Eikelboom (2000). Morphotype
classification was carried out conforming to the described classification keys, based
on morphological features of the filaments: shape, length, and diameter of filamentous
bacteria, motility, presence of branching, attached growth of other bacteria to the
filaments, (visible or not visible) septa between adjoining cells, shape of cells,
presence of a sheath and sulphur granules (Eikelboom 2000).

Histochemical staining
Following FISH, polyphosphate inclusions were stained with 28 μM 4',6-diamidino-
2-phenylindole (DAPI) for 1 h at 4°C in the dark. After staining at such relatively
high DAPI concentrations, the polyphosphate granules appear bright yellow with
fluorescence microscopy (Streichan, Golecki and Schon 1990).
Polyhydroxyalkanoates (PHA) were stained with 1% [w/v] Nile Blue A for 10 mins at
55°C essentially as described previously (Ostle and Holt 1982). FISH images were
acquired prior to staining and the same fields of view relocated.

Microsphere adhesion to cells (MAC)
MAC was performed on a sample of fresh sludge to identify the hydrophobicity of
target cells, applying the method of Kragelund and colleagues (2005) using a
sonicated solution of 0.2 μm FluoSpheres fluorescent sulphate-modified microspheres
with excitation and emission properties (505/515 nm) (Life Technologies
Corporation, Eugene OR, USA).

Microautoradiography
Biomass was sampled from the aerobic stage of full-scale activated sludge WWTPs in
Aalborg West, Bjergmarken, Ringkøbing, and Odense North-West, Denmark. All
plants are designed for N removal and enhanced biological phosphorus removal
(EBPR) and have stable performance. For further details on the plants assessed in this

study, see Mielczarek *et al.*, (2013). Biomass samples were stored at 4°C and all incubations performed within 24 h from sampling. The MAR incubation protocol was based on the method of Nierychlo et al., (2015). Activated sludge was aerated for 40 min at room temperature prior to MAR incubation to reduce the residual substrates, oxygen, and NO_x present. Sludge was then diluted with filtered sludge water from the same plant to yield a biomass concentration of 1 mgSSmL^{-1} and transferred to 11 ml serum bottles. Radiolabeled substrates were added to yield a total radioactivity of 10 µCi mg^{-1} SS. The following substrates were used: [^{3}H]acetate, [^{3}H]galactose (Amersham Biosciences, UK), [^{3}H]glucose, [^{3}H]mannose, [^{14}C]pyruvate (Perkin-Elmer, Waltham MA, USA), [^{3}H]amino acid mix, [^{14}C]butyric acid, [^{3}H]fructose, [^{3}H]glycerol, [^{3}H]ethanol, [^{3}H]lactate, [^{3}H]NAG, [^{14}C]propionate (American Radiolabeled Chemicals Inc., Saint Louis MO, USA). The corresponding cold substrate was added to yield a total concentration of 2 mM organic substrate. To achieve anaerobic conditions, prior to substrate addition, oxygen was removed by repeated evacuation of the headspace and subsequent injection of O_2-free N_2. Anaerobic incubations with selected substrates were supplemented with 0.5 mM nitrite or 2 mM nitrate to assess their use as electron acceptors. The supernatant concentrations were monitored using Quantofix Nitrate/Nitrite strips (Macherey-Nagel, Düren, Germany) and readjusted to their initial concentrations anaerobically to prevent exhaustion. Samples were incubated for 3 h at room temperature (approx. 21°C) on a rotary shaker at 250 rpm. Incubations with [^{14}C]carbonate (American Radiolabeled Chemicals Inc., Saint Louis MO, USA) contained 20 µCi mg^{-1} SS of radiolabeled substrate. 1mM NH_4Cl was added to half of the incubations to investigate ammonia and nitrite (produced from the oxidation of added ammonia) oxidation activity. These samples were incubated aerobically (same conditions as above) for 5h, as suggested by Daims et al., (2001). A pasteurized biomass (heated to 70°C for 10 min) incubation was prepared as a negative control to assess possible silver grain formation due to chemography. Incubations were terminated by the addition of cold PFA to a final concentration of 4% [w/v]. Samples were fixed for 3 h at 4°C and subsequently washed 3 times with sterile filtered tap water. Aliquots of 30 µl of the biomass were gently homogenized between glass coverslips. Following FISH (see earlier), coverslips were coated with Ilford K5D emulsion (Polysciences, Inc., Warrington, PA, USA), exposed in the dark for periods of 10 days, and developed with Kodak D-19 developer.

Results

Distribution of Chloroflexi in Danish full-scale WWTPs

Phylogenetic tree based on 16S rRNA gene sequences shows all the abundant Chloroflexi groups found in Danish WWTPs with nutrient removal and their phylogenetic relationship (**Figure 1**). The Chloroflexi phylum is among the most abundant phyla in full-scale systems in Denmark, constituting on average 10.6% of the total reads across all plants (**Figure 2A**). The Chloroflexi classes Anaerolineae, Caldilineae, Ardenticatenia, and SJA-15 made up the majority of members of the

phylum present (**Figure 1B**). All the abundant genus-level phylotypes (**Figure 1C**)
are novel, having no available cultured representatives, and were initially given
provisional alphanumeric names in the MiDAS database (McIlroy *et al.* 2015) (not
shown here). Based on their characterization, as presented in this study and previous
publications, we propose new, previously unpublished, candidate names (Murray and
Stackebrandt 1995). These names have been incorporated into the MiDAS taxonomy
version 2.1 (McIlroy *et al.* 2015) and are used throughout this report. These include:
'*Candidatus* Sarcinathrix' (L. fem. n. *Sarcina*, a package, bundle; Gr. fem. n. *thrix*,
hair), '*Candidatus* Villogracilis' (L. m. n., *Villo*, tuft of hair; L. fem. adj. *gracilis*,
slim, slender, thin; formerly MiDAS taxon SBR1029 (release 1.21)), '*Candidatus*
Defluviifilum' (L. n. *defluvium*, sewage; L. neut. n. *filum*, a thread; formerly MiDAS
taxon P2CN44 (release 1.21)), and '*Candidatus* Amarilinum' (Gr. n. *amara*, trench,
conduit, channel; here, a sewage duct; L. fem. n. *linea*, line; formerly MiDAS taxon
C10_SB1A (release 1.21)). 'Kouleothrix spp.', possessing the 1851 bulking filament
morphotype, was present in low abundance with a median and mean of 0.04 and
0.4%, respectively. 'Ca. Defluviifilum', 'Ca. Promineofilum', 'Ca. Villogracilis', and
'Ca. Sarcinathrix' represent the four most abundant genera by median read abundance
(**Figure 1C**), collectively constituting on average 6.2% of the total reads across all
Danish plants assessed in this study. These phylotypes were relatively stably present
across the different WWTPs (**Figure S1**) and therefore represent core members of the
microbial community of these systems. As little is known regarding the physiology of
the latter two genera, they were selected for a detailed characterization in this study.
Relative to these phylotypes, the 'Ca. Amarilinum' showed a much more dynamic
distribution and periodically reached abundances in excess of 30% of the amplicon
reads, which would indicate a likely role in bulking episodes in Denmark. As such,
this genus, known only by its 16S rRNA gene sequences, was also selected for
characterization in this study.

Phylogeny and FISH probe design
'Ca. Villogracilis' are members of the Anaerolineaceae, which is currently the sole
family of the class Anaerolineae in the MiDAS and SILVA taxonomies. These
sequences fall within order SBR1031 in the Greengenes taxonomy (McDonald *et al.*
2012). The CFX763A and CFX763B probes were designed to cover separate sub-
groups (A and B) of the 'Ca. Villogracilis' (**Figure S2**) - collectively covering >60%
of the MiDAS database sequences classified to the genus. The target region is not
covered by the V1-3 region amplicon sequences, although both probes match the full-
length database sequences most closely related to the abundant OTU sequences (data
not shown). When applied to full-scale activated sludge biomass, both probes
hybridized thin filaments (0.3-0.4 μm wide and 15-50 μm long) that were often
observed in bundles and almost exclusively located within the flocs. The
CFX763AB_H1A and CFX763AB_H1B helper probes are recommended to give
optimal fluorescence signal for both probes. Amplicon sequencing of the V1-3 region
of the 16S rRNA cannot be used to confidently separate the A and B sub-groups, due

to high sequence similarity, but qFISH indicates that the former is the more
numerically important of the two.

The 'Ca. Amarilinum' genus falls within the novel MiDAS Chloroflexi class-
level-group SJA-15, together with the also abundant genus 'Ca. Sarcinathrix' (**Figure**
2). A probe to cover the entire 'Ca. Amarilinum' group was not identified, so the
CFX64 (**Table 1**) was designed for the abundant amplicon OTUs (OTU_3 and
OTU_4592, **Figure S3**) and the most closely related full-length database sequence
(AF513086). These sequences share >97% similarity. As the probe covers these
abundant OTUs, it should cover the majority of members of the genus in full-scale
activated sludge in Denmark. When applied to activated sludge, the probe hybridized
exclusively to filamentous bacteria (see later for a detailed description of their
morphology). A strong positive signal was obtained without addition of designed
unlabeled helper probes CFX64_H1 and CFX64_H2 (**Table 1**), which did not
noticeably improve fluorescent signal (data not shown). The 'Ca. Amarilinum'
filaments constituted up to 30% of the community biovolume in some full-scale
WWTPs in Denmark - confirming their high abundance with amplicon sequencing
(see **Table 2**).

FISH probes are already available to target the 'Ca. Sarcinathrix' (CFX67),
which was shown to possess the Eikelboom 0914 morphotype in nutrient removal
activated sludge systems in Australia (Speirs *et al.* 2011). Application of CFX67
probe did not give significant positive fluorescence to bacterial cells in Danish plants.
The probe misses several full-length sequences classified to the genus and does not
cover any of the abundant OTU sequences (>0.1% average read abundance in at least
1 plant) from the MiDAS full-scale survey. As such, two new probes were designed
to give better coverage of the clade. The designed CFX449 and CFX1151 individually
target 85% of the full-length database sequences. As such, they can be applied with
different fluorochromes to confirm specific coverage of the genus, or together as a
mix to give higher signal (**Figure S4**). Helper probes were not required for either
probe, but did give a more even signal over the filament, which was also reported by
Speirs *et al.* (2011) for the CFX67 probe. The few filaments positive for the CFX67
probe in the Danish WWTPs assessed in this study also hybridized the CFX449 and
CFX1151 probes (**Figure S4**). Quantitative FISH values with the CFX449 and
CFX1151 were similar to amplicon-sequencing based estimates (**Table 2**).

All three phylotypes studied gave positive hybridization signal with the
EUBmix probe set, which is commonly applied as a universal probe targeting bacteria
(EUB338, EUB338-II, and EUB338-III). This is of interest, given that many
Chloroflexi reportedly lack the target site for the EUBmix probe set and fail to
hybridize the probe *in situ* (Kragelund *et al.* 2007, 2011; Speirs *et al.* 2009). Most 16S
rRNA gene sequences of the '*Ca.* Amarilinum', '*Ca.* Sarcinathrix' and 'Kouleothrix'
contain the site for the EUB338 probe, the 'Ca. Villogracilis' sequences possess the
EUB338-III site, and all groups have been shown to hybridize the probe *in situ*. Most
members of the '*Ca.* Defluviifilum' have one mismatch, though *in silico* analysis with
the MathFISH software (Yilmaz, Parnerkar and Noguera 2011) predicts positive
binding (with a calculated melting formamide point ($[\Delta FA]_m$) of 50%), which is

confirmed *in situ* for filaments hybridizing the T0803-0654 probe designed to target
the group (Kragelund *et al.* 2011). Thus, of the abundant phylotypes, only the 'Ca.
Promineofilum' genus is not covered by the EUBmix probe set (Speirs *et al.* 2009).

Morphological description and classification
The morphological properties of the CFX64 positive filaments, representing the 'Ca.
Amarilinum' genus, were investigated in detail for association to a morphotype of the
well-known antecedent classification systems (Eikelboom 1975). Though not clearly
visible, the cells appeared to be rectangular with no visible septa, a trichrome
thickness of 1.0-2.2 μm, and a length in the 20-140 μm range. They were non-motile,
Gram stain negative, with no branching or attached growth. The whole filaments
stained blue/violet with the Neisser stain, with no visible volutin granules. Excess
polyphosphate stores were not observed with DAPI staining. Cells did not appear to
contain excess stores of polyhydroxyalkanoates (PHAs), with negative results with
Nile blue A staining and only small positive granules observed with the Sudan black
stain. From these observations, primarily based on the characteristic violet color of
the cells after Neisser staining, it is suggested that the morphology of the filament is
most consistent with the Eikelboom type 0092 morphotype (Eikelboom 2000). The
'Ca. Promineofilum' also reportedly has the 0092 morphotype (Speirs *et al.* 2009),
but there was no observed overlap between the CFX64 and the CFX197 probes
targeting the '*Ca.* Amarilimum' and 'Ca. Promineofilum' genera, respectively
(**Figure S5**). The 'Ca. Promineofilum' phylotype is also thinner in appearance, with a
trichome diameter of approx. 0.8 μm (Speirs *et al.* 2009). Very few of the fluorescent
sulphate modified microspheres attached to the CFX64 positive filaments (data not
shown), indicating that they do not have a hydrophobic surface and are likely not
involved in foam formation.
Morphological descriptions are already reported for members of the 'Ca.
Sarcinathrix' (Speirs *et al.* 2011) and 'Ca. Defluviifilum' (Kragelund *et al.* 2011;
Speirs *et al.* 2017). The surface hydrophobicity of 'Ca. Sarcinathrix' was assessed for
the first time here, where it was determined to be hydrophilic and therefore unlikely to
be involved in foam formation. Morphological classification of the 'Ca. Villogracilis'
was not successful due to their location within the floc, making interpretation of
staining analyses difficult.

In situ substrate uptake
The results for substrate uptake by probe-defined 'Ca. Villogracilis' sub-groups A and
B, 'Ca. Amarilinum', and 'Ca. Sarcinathrix' using MAR-FISH under various
conditions are shown in **Table S1,** and a summary of known *in situ* traits of abundant
Chloroflexi is given in **Table 3**. 'Ca. Villogracilis' sub-group A, 'Ca. Amarilinum',
and 'Ca. Sarcinathrix' only utilized sugars of the 13 substrates tested, consistent with
other characterized Chloroflexi genera. The phylotypes differed in the types of sugars
taken up, noting that variation was also observed within the 'Ca. Amarilinum' genus –
with some filaments strongly positive and others clearly negative for fructose uptake.
'Ca. Villogracilis' sub-group B filaments were negative for all substrates and

conditions. It may be that they have a relatively lower activity than the much more
abundant sub-group A filaments that is below the detection of MAR. Further analyses
are required to assess the reason for the observed lack of substrate uptake. All three
genera were able to take up substrates under anoxic conditions, suggesting
fermentative metabolisms. Anoxic uptake of sugars in presence of nitrate/nitrite was
also observed, but their potential for denitrification is unclear, given that uptake was
also observed without nitrate/nitrite addition. The same ambiguous results were
obtained for the 'Ca. Defluviifilum' (Kragelund *et al.* 2011). The ability for
nitrification was also assessed for these groups in this study, given that *Nitrolancetus*
hollandicus, a nitrite oxidizing member of the class Thermomicrobia of the
Chloroflexi, was isolated from activated sludge (Sorokin *et al.* 2012); albeit in a
different class to the abundant members of the Chloroflexi Danish full-scale systems.
None of the genera appeared to be behaving as nitrifiers, with no observed uptake of
labeled CO_2 in the presence of ammonia (**Table 3 and S1**), while a positive MAR
signal was noted for *Nitrosomonas* as well as *Nitrospira,* targeted by the probe
Cluster6a_192 (Adamczyk *et al.* 2003) and Ntsp662 (Daims *et al.* 2001), respectively.

Discussion
The results of this study of the distribution and *in situ* morphology and physiology of
individual abundant phylotypes of the phylum Chloroflexi give valuable insight into
their potential importance to the operation of WWTPs. The 'Ca. Promineofilum', 'Ca.
Defluviifilum', 'Ca. Villogracilis', and 'Ca. Sarcinathrix' appear to be consistently
abundant filamentous members of the full-scale WWTP community, where they
certainly make an important contribution to the bulk nutrient transformations.
Relative to other abundant Chloroflexi genera, the distribution of the 'Ca.
Amarilinum' is much more dynamic, reaching levels of >30% of the biovolume in
some plants. As such, it is more likely that the 'Ca. Amarilinum' are responsible for
acute bulking episodes in Danish WWTPs, while other abundant phylotypes are core
members of the community that are possibly important to floc structure and the
breakdown of organics (see later). While the 'Ca. Villogracilis' are filamentous, they
are almost exclusively found within the flocs and are unlikely to contribute to bulking
episodes. This observation explains why they have never been detected before and
highlights the value of large-scale full-scale surveys, such as MiDAS, for describing
the abundant members of wastewater treatment systems and the importance of *in situ*
analyses to evaluate the role of organisms in floc structure and settleability. In
addition to a potential contribution to bulking, members of the 'Ca. Promineofilum'
are also putatively involved in membrane fouling in MBR systems (Ziegler *et al.*
2016), and 'Ca. Defluviifilum' has a hydrophobic cell surface that has been
implicated in the stabilization of problematic foams on the surface of reactor tanks
(Kragelund *et al.* 2011).
In recent studies Speirs *et al.*, (2015; 2017) described filamentous Chloroflexi
phylotypes in Australian WWTPs possessing the bulking Eikelboom morphotypes
0803 and 0041/0675 that are commonly observed by light microscopy in Danish

plants (Mielczarek *et al.* 2012). These phylotypes classified within the Anaerlolineae
and Caldilinaea, respectively. However, none of these were found to be abundant in
the amplicon based survey of Danish systems (data not shown). Their FISH probe
defined 0675 phylotype falls within the MiDAS defined 'Ca. Defluviifilum' genus (v.
2.1), shown previously to possess the Eikelboom 0803 phylotype (Kragelund *et al.*
2011), which they suggest should be split into a separate genus based on the
divergence of its 16S rRNA gene sequence (only ca. 90% similar). Further
characterisation of members of the current 'Ca. Defluviifilum' clade, including
obtaining representative genomes, will help to resolve the phylogeny of these
organisms. Members of the 'Kouleothrix' genus, shown to possess the Eikelboom
1851 bulking morphotype (Beer *et al.* 2002; Kohno, Sei and Mori 2002), were also
relatively low in abundance in Danish plants (Kragelund *et al.* 2011) but are common
in other countries – such as Japan, where they have been suggested as a major
contributor to filamentous bulking episodes (Nittami *et al.* 2017). Global surveys will
help to establish how relevant the abundant Danish phylotypes characterised in this
paper are worldwide.
The abundant phylotypes in Danish WWTPs appear to be
organoheterotrophic, fermentative, facultative anaerobes. This is suggested from their
exclusive utilization of sugars and ability for anaerobic carbon uptake. Fermentative
pathways were also annotated in the sole genome from these genera – '*Ca.*
Promineofilum breve' (McIlroy *et al.* 2016). The physiology determined for the 'Ca.
Defluviifilum' and 'Ca. Villogracilis' is consistent with other members of their
respective classes (Caldilineae and Anaerolinea), which are mostly fermentative
organoheterotrophic filaments growing on sugars and/or amino acids (Sekiguchi *et al.*
2003; Yamada *et al.* 2006, 2007; Grégoire *et al.* 2011; Nunoura *et al.* 2013;
Podosokorskaya *et al.* 2013; Imachi *et al.* 2014; McIlroy *et al.* 2017a). 'Ca.
Villogracilis' is the first reported facultative anaerobic genus of the Anaerolineae,
with all isolates of other described genera being obligate anaerobes.
In addition to the likely fermentation of sugars by the abundant Chloroflexi,
metabolic diversity within the less abundant members of the phylum is evident – with
reported assimilation of short and long chain fatty acids, amino acids, and glycerol
(**Table 3**). Some members of the phylum also appear unable to take up substrates
anaerobically and may scavenge sugars released from aerobic breakdown of complex
organic matter. Uptake of *N*-aminoglucosamine (NAG), a component of
peptidoglycan and lipopolysaccharides, for some Chloroflexi, suggests a specific role
in the breakdown of cellular material (Kragelund *et al.* 2007, 2011; Miura, Watanabe
and Okabe 2007). Of the known Chloroflexi genera reportedly abundant in activated
sludge, only the 'Kouleothrix' genus is seemingly unable to take up substrates
anaerobically *in situ* (Kragelund *et al.* 2007), although isolates of the genus are
capable of anaerobic fermentative growth on sugars (Kohno, Sei and Mori 2002).
In this study, the most abundant members of the Chloroflexi in Danish nutrient
removal WWTPs were identified and their ecophysiology described. These
phylotypes appear to differ in their impact on plant operation - with suggested
importance in sludge settleability, foaming, and membrane fouling being associated

with different groups. All abundant members of the phylum likely ferment sugars, and
future research should aim to obtain representative genomes for each in order to carry
out more detailed comparison of their metabolic activities. Such an approach will
explain important questions regarding how these organisms coexist, and what
conditions determine their relative abundances. The FISH probes designed in this
study will allow more hypothesis-based *in situ* investigation of their physiologies,
based on genomic evidence. The taxonomic annotation and design of FISH probes for
the abundant Chloroflexi, in combination with the high throughput nature of 16S
rRNA gene amplicon sequencing, will also allow their routine observation and study.
Defining and naming these novel genus level taxa importantly provides the
foundation, upon which information on their morphology, distribution, and
physiology can be gathered for an in-depth understanding of their ecology and how
this might relate to operational parameters.

Funding
This work was supported by the Danish Council for Independent Research (grant no.
4093-00127A); Innovation Fund Denmark (EcoDesign); The Obel Family
Foundation; Danish wastewater treatment plants in MiDAS; and Aalborg University.

Acknowledgements
We thank K. Vilstrup for help to latin names.

References

Adamczyk J, Hesselsoe M, Iversen N *et al.* The Isotope Array, a New Tool That
Employs Substrate-Mediated Labeling of rRNA for Determination of
Microbial Community Structure and Function. *Appl Environ Microbiol*
2003;**69**:6875–87.

Amann RI, Binder BJ, Olson RJ *et al.* Combination of 16S rRNA-targeted
oligonucleotide probes with flow cytometry for analyzing mixed microbial
populations. *Appl Env Microbiol* 1990;**56**:1919–25.

Beer M, Seviour EM, Kong Y *et al.* Phylogeny of the filamentous bacterium
Eikelboom Type 1851, and design and application of a 16S rRNA targeted
oligonucleotide probe for its fluorescence in situ identification in activated
sludge. *FEMS Microbiol Lett* 2002;**207**:179–83.

Beer M, Stratton HM, Griffiths PC *et al.* Which are the polyphosphate accumulating
organisms in full-scale activated sludge enhanced biological phosphate
removal systems in Australia? *J Appl Microbiol* 2006;**100**:223–43.

Björnsson L, Hugenholtz P, Tyson GW *et al.* Filamentous Chloroflexi (green non-
sulfur bacteria) are abundant in wastewater treatment processes with
biological nutrient removal. *Microbiology* 2002;**148**:2309–18.

Cole JR, Wang Q, Fish JA *et al.* Ribosomal Database Project: data and tools for high
throughput rRNA analysis. *Nucleic Acids Res* 2014;**42**:D633-42.

Daims H, Brühl A, Amann R *et al.* The domain-specific probe EUB338 is insufficient
for the detection of all Bacteria: development and evaluation of a more
comprehensive probe set. *Syst Appl Microbiol* 1999;**22**:434–44.

Daims H, Lücker S, Wagner M. daime, a novel image analysis program for microbial
ecology and biofilm research. *Env Microbiol* 2006;**8**:200–13.

Daims H, Nielsen JL, Nielsen PH *et al.* In Situ Characterization of Nitrospira-Like
Nitrite-Oxidizing Bacteria Active in Wastewater Treatment Plants. *Appl*
Environ Microbiol 2001;**67**:5273–84.

Daims H, Stoecker K, Wagner M. Fluorescence in situ hybridization for the detection
of prokaryotes. In: Osborn AM, Smith CJ (eds.). *Molecular Microbial*
Ecology. New York: Taylor & Francis, 2005, 213–39.

Eikelboom D. *Process Control of Activated Sludge Plants by Microscopic*
Investigation. London: IWA Publishing, 2000.

Eikelboom DH. Filamentous organisms observed in activated sludge. *Water Res*
1975;**9**:365–88.

Gich F, Garcia-Gil J, Overmann J. Previously unknown and phylogenetically diverse
members of the green nonsulfur bacteria are indigenous to freshwater lakes.
Arch Microbiol 2001;**177**:1–10.

Grégoire P, Bohli M, Cayol J-L *et al.* Caldilinea tarbellica sp. nov., a filamentous,
thermophilic, anaerobic bacterium isolated from a deep hot aquifer in the
Aquitaine Basin. *Int J Syst Evol Microbiol* 2011;**61**:1436–41.

Greuter D, Loy A, Horn M *et al.* probeBase—an online resource for rRNA-targeted
oligonucleotide probes and primers: new features 2016. *Nucleic Acids Res*
2016;**44**:D586–9.

Imachi H, Sakai S, Lipp JS *et al.* Pelolinea submarina gen. nov., sp. nov., an
anaerobic, filamentous bacterium of the phylum Chloroflexi isolated from
subseafloor sediment. *Int J Syst Evol Microbiol* 2014;**64**:812–8.

Jenkins D, Richard MG, Daigger GT. *Manual on the Causes and Control of Activated*
Sludge Bulking, Foaming and Other Solids Separation Problems. 3 rd.
London, England: CRC Press LLC, 2004.

Kindaichi T, Nierychlo M, Kragelund C *et al.* High and stable substrate specificities
of microorganisms in enhanced biological phosphorus removal plants. *Environ*
Microbiol 2013;**15**:1821–31.

Kohno T, Sei K, Mori K. Characterization of type 1851 organism isolated from
activated sludge samples. *Water Sci Technol J Int Assoc Water Pollut Res*
2002;**46**:111–4.

Kragelund C, Levantesi C, Borger A *et al.* Identity, abundance and ecophysiology of
filamentous Chloroflexi species present in activated sludge treatment plants.
FEMS Microbiol Ecol 2007;**59**:671–82.

Kragelund C, Müller E, Schade M *et al.* Identification of filamentous bacteria by
FISH. In: Nielsen PH, Daims H, Lemmer H (eds.). *FISH Handbook for*
Biological Wastewater Treatment. London: IWA Publishing, 2009, 33–68.

Kragelund C, Nielsen JL, Thomsen TR *et al.* Ecophysiology of the filamentous
Alphaproteobacterium Meganema perideroedes in activated sludge. *FEMS*
Microbiol Ecol 2005;**54**:111–22.

Kragelund C, Thomsen TR, Mielczarek AT *et al.* Eikelboom's morphotype 0803 in
activated sludge belongs to the genus Caldilinea in the phylum Chloroflexi.
FEMS Microbiol Ecol 2011;**76**:451–62.

Ludwig W, Strunk O, Westram R *et al.* ARB: a software environment for sequence
data. *Nucleic Acids Res* 2004;**32**:1363–71.

McDonald D, Price MN, Goodrich J *et al.* An improved Greengenes taxonomy with
explicit ranks for ecological and evolutionary analyses of bacteria and archaea.
ISME J 2012;**6**:610–8.

McIlroy SJ, Karst SM, Nierychlo M *et al.* Genomic and in situ investigations of the
novel uncultured Chloroflexi associated with 0092 morphotype filamentous
bulking in activated sludge. *ISME J* 2016;**In press**.

McIlroy SJ, Kirkegaard RH, Dueholm MS *et al.* Culture-Independent Analyses
Reveal Novel Anaerolineaceae as Abundant Primary Fermenters in Anaerobic
Digesters Treating Waste Activated Sludge. *Front Microbiol* 2017a;**8**, DOI:
10.3389/fmicb.2017.01134.

McIlroy SJ, Kirkegaard RH, McIlroy B *et al.* MiDAS 2.0: an ecosystem-specific
taxonomy and online database for the organisms of wastewater treatment
systems expanded for anaerobic digester groups. *Database* 2017b;**2017**, DOI:
10.1093/database/bax016.

McIlroy SJ, Saunders AM, Albertsen M *et al.* MiDAS: the field guide to the microbes
of activated sludge. *Database* 2015;**2015**:1–8.

McIlroy SJ, Tillett D, Petrovski S *et al.* Non-target sites with single nucleotide
insertions or deletions are frequently found in 16S rRNA sequences and can
lead to false positives in fluorescence in situ hybridization (FISH). *Env*
Microbiol 2011;**13**:38–47.

Mielczarek AT, Kragelund C, Eriksen PS *et al.* Population dynamics of filamentous
bacteria in Danish wastewater treatment plants with nutrient removal. *Water*
Res 2012;**46**:3781–95.

Mielczarek AT, Nguyen HT, Nielsen JL *et al.* Population dynamics of bacteria
involved in enhanced biological phosphorus removal in Danish wastewater
treatment plants. *Water Res* 2013;**47**:1529–44.

Miura Y, Watanabe Y, Okabe S. Significance of Chloroflexi in performance of
submerged membrane bioreactors (MBR) treating municipal wastewater.
Environ Sci Technol 2007;**41**:7787–94.

Morgan-Sagastume F, Nielsen JL, Nielsen PH. Substrate-dependent denitrification of
abundant probe-defined denitrifying bacteria in activated sludge. *FEMS*
Microbiol Ecol 2008;**66**:447–61.

Murray RG, Stackebrandt E. Taxonomic note: implementation of the provisional
status Candidatus for incompletely described procaryotes. *Int J Syst Bacteriol*
1995;**45**:186–7.

Nierychlo M, Nielsen JL, Nielsen PH. Studies of the Ecophysiology of Single Cells in
Microbial Communities by (Quantitative) Microautoradiography and
Fluorescence In Situ Hybridization (MAR-FISH). In: McGenity TJ, Timmis
KN, Nogales Fernández B (eds.). *Hydrocarbon and Lipid Microbiology*
Protocols, Springer Protocols Handbooks. 1st ed. Berlin-Heidelberg:
Springer-Verlag, 2015.

Nittami T, Speirs LBM, Yamada T *et al.* Quantification of *Chloroflexi* Eikelboom
morphotype 1851 for prediction and control of bulking events in municipal
activated sludge plants in Japan. *Appl Microbiol Biotechnol* 2017;**101**:3861–9.

Nunoura T, Hirai M, Miyazaki M *et al.* Isolation and characterization of a
thermophilic, obligately anaerobic and heterotrophic marine Chloroflexi
bacterium from a Chloroflexi dominated microbial community associated with
a japanese shallow hydrothermal system, and proposal for Thermomarinilin.
Microbes Environ JSME 2013;**28**:228–35.

Ostle AG, Holt JG. Nile blue A as a fluorescent stain for poly-beta-hydroxybutyrate.
Appl Env Microbiol 1982;**44**:238–41.

Podosokorskaya OA, Bonch-Osmolovskaya EA, Novikov AA *et al.* Ornatilinea
apprima gen. nov., sp. nov., a cellulolytic representative of the class
Anaerolineae. *Int J Syst Evol Microbiol* 2013;**63**:86–92.

Quast C, Pruesse E, Yilmaz P *et al.* The SILVA ribosomal RNA gene database
project: improved data processing and web-based tools. *Nucleic Acids Res*
2013;**41**:D590–6.

Sekiguchi Y, Yamada T, Hanada S *et al.* Anaerolinea thermophila gen. nov., sp. nov.
and Caldilinea aerophila gen. nov., sp. nov., novel filamentous thermophiles
that represent a previously uncultured lineage of the domain Bacteria at the
subphylum level. *Int J Syst Evol Microbiol* 2003;**53**:1843–51.

Seviour EM, Blackall LL, Christensson C *et al.* The filamentous morphotype
Eikelboom Type 1863 is not asingle genetic entity. *J Appl Microbiol*
1997;**82**:411–21.

Sorokin DY, Lücker S, Vejmelkova D *et al.* Nitrification expanded: discovery,
physiology and genomics of a nitrite-oxidizing bacterium from the phylum
Chloroflexi. *ISME J* 2012;**6**:2245–56.

Speirs L, Nittami T, McIlroy S *et al.* Filamentous bacterium Eikelboom type 0092 in
activated sludge plants in Australia is a member of the phylum Chloroflexi.
Appl Env Microbiol 2009;**75**:2446–52.

Speirs LBM, Dyson ZA, Tucci J *et al.* Eikelboom filamentous morphotypes 0675 and
0041 embrace members of the Chloroflexi: resolving their phylogeny, and
design of fluorescence in situ hybridisation probes for their identification.
FEMS Microbiol Ecol 2017;**93**, DOI: 10.1093/femsec/fix115.

Speirs LBM, McIlroy SJ, Petrovski S *et al.* The activated sludge bulking filament
Eikelboom morphotype 0914 is a member of the Chloroflexi. *Env Microbiol*
Rep 2011;**3**:159–65.

Speirs LBM, Tucci J, Seviour RJ. The activated sludge bulking filament Eikelboom
morphotype 0803 embraces more than one member of the *Chloroflexi*. Stams
A (ed.). *FEMS Microbiol Ecol* 2015;**91**:fiv100.

Streichan M, Golecki JR, Schon G. Polyphosphate-accumulating bacteria from
sewage plants with different proceses for biological phosphorus removal.
FEMS Microbiol Lett 1990;**73**:113–24.

Wallner G, Amann R, Beisker W. Optimizing fluorescent in situ hybridization with
rRNA-targeted oligonucleotide probes for flow cytometric identification of
microorganisms. *Cytometry* 1993;**14**:136–43.

Wanner J, Kragelund C, Nielsen PH. Microbiology of bulking. In: Seviour RJ,
Nielsen PH (eds.). *Microbial Ecology of Activated Sludge*. London: IWA
Publishing, 2010, 191–214.

Xia Y, Kong Y, Nielsen PH. In situ detection of protein-hydrolysing microorganisms
in activated sludge. *FEMS Microbiol Ecol* 2007;**60**:156–65.

Yamada T, Imachi H, Ohashi A *et al.* Bellilinea caldifistulae gen. nov., sp. nov. and
Longilinea arvoryzae gen. nov., sp. nov., strictly anaerobic, filamentous
bacteria of the phylum Chloroflexi isolated from methanogenic propionate-
degrading consortia. *Int J Syst Evol Microbiol* 2007;**57**:2299–306.

Yamada T, Sekiguchi Y, Hanada S *et al.* Anaerolinea thermolimosa sp. nov.,
Levilinea saccharolytica gen. nov., sp. nov. and Leptolinea tardivitalis gen.
nov., sp. nov., novel filamentous anaerobes, and description of the new classes
Anaerolineae classis nov. and Caldilineae classis nov. in the . *Int J Syst Evol*
Microbiol 2006;**56**:1331–40.

Yilmaz LS, Parnerkar S, Noguera DR. mathFISH, a web tool that uses
thermodynamics-based mathematical models for in silico evaluation of
oligonucleotide probes for fluorescence in situ hybridization. *Appl Env*
Microbiol 2011;**77**:1118–22.

Ziegler AS, McIlroy SJ, Larsen P *et al.* Dynamics of the Fouling Layer Microbial
Community in a Membrane Bioreactor. *PloS One* 2016;**11**:e0158811.

Figures and Tables

Figure 1. Maximum-likelihood (PhyML) 16S rRNA gene phylogenetic tree of
abundant activated sludge phylotypes (bold typeface) and isolated members of the
phylum Chloroflexi. The alignment used for the tree applied a 20% conservational
filter to remove hypervariable positions, giving 1108 aligned positions. Phylogenetic
classification is indicated with black brackets and is based on the MiDAS database
(Release 2.1). Probe coverage of probes relevant to the current study is shown with
red coloured brackets. Associated activated sludge morphotypes of previously
described phylotypes are designated with blue coloured brackets. Bootstrap values
from 100 resamplings are indicated for branches with >50% (white dot), 50-70%
(gray), and >90% (black) support. Species of the phylum Cyanobacteria were used as
the outgroup. The scale bar represents substitutions per nucleotide base.

Figure 2. Distribution of Chloroflexi in 25 full-scale activated sludge WWTPs
sampled 2-4 times per year from 2006 to 2015. (a) 10 most abundant phyla in Danish
WWTPs. (b) 10 most abundant Chloroflexi classes in Danish WWTPs (c) 10 most
abundant Chloroflexi genera in Danish WWTPs. X-axis shows the relative read
abundance in percentage of total bacteria.

Figure 3. Composite FISH micrographs of novel Chloroflexi genera in full-scale
activated sludge. Specific probes (Cy3-label, red) target (a) '*Ca.* Amarilinum', (b)
'*Ca.* Villogracilis', and (c) '*Ca.* Sarcinathrix', and EUBmix probe (Cy5-label, blue)
targets most bacteria. Activated sludge was sampled from (a) Bjergmarken WWTP,
(b) Odense North East WWTP, and (c) Aalborg West WWTP. Target filaments
appear magenta, while all other cells appear blue. Scale bars represent 20 μm.

Figure 4. FISH and corresponding bright-field MAR micrographs showing sugar
uptake by three abundant Chloroflexi phylotypes ('*Ca.* Amarilinum' hybridized probe
CFX64; '*Ca.* Villogracilis' sub-group A hybridized probe CFX763A; '*Ca.*
Sarcinathrix' hybridized probe CFX1151). Activated sludge was sampled in Aalborg
West or Odense North West WWTPs. Target cells in FISH micrograph overlays
appear yellow: specific probe (red) + EUBmix (green); and non-target cells appear
green (EUBmix only). Black silver granules indicate positive MAR signal. Scale bar
represents 10 μm.

Table 1. Probes designed and optimized in this study.

Probe	*E. coli* pos.	Target group	Coverage[*]	Non-target hits	Sequence (5'-3')	[FA]%[**]
CFX64	**64-99**	*'Ca.* Amarilinum spp.'	**1/2**	**0**	**TCT ACC TAA GCA GAC CGT TC**	**30**
CFX64_H1[#]	41-63	Helper for CFX64	N/A	N/A	AAC TTG CAT GTG TTA AGC ACG CC	N/A
CFX64_H2[#]	100-116	Helper for CFX64	N/A	N/A	TCA CCC GTG CGC CAC TG	N/A
CFX763A	**763-783**	*'Ca.* Villogracilis spp.' sub-group A	**68/260**	**1**	**GTT TAC TAC CCT AGC TTT CGC**	**45**
CFX763A_C1	763-783	Competitor for CFX763A probe	N/A	N/A	GTT CAC TAC CCT AGC TTT CGC	N/A
CFX763A_C2	763-783	Competitor for CFX763A probe	N/A	N/A	GTT TAC TCC CCT AGC TTT CGC	N/A
CFX763A_C3	763-783	Competitor for CFX763A probe	N/A	N/A	GTT TGC TAC CCT AGC TTT CGC	N/A
CFX763A_C4[Ψ]	763-783	Competitor for CFX763A probe	N/A	N/A	GTT TAC TAC CCT AGC TGT CGC	N/A
CFX763AB_H1A	784-808	Helper for CFX763A, CFX763B	N/A	N/A	TAG GAT TAC CGG GGT CTC TAA TCC C	N/A
CFX763AB_H1B	784-808	Helper for CFX763A, CFX763B	N/A	N/A	TAG GAT TAC CSG GGT CTC TA ATC CC	N/A
CFX763B	**763-783**	*'Ca.* Villogracilis spp.' sub-group B	**91/260**	**0**	**GTT TAC TAC CCT AGC TGT CGC**	**45**
CFX763B_C1[§]	763-783	Competitor for CFX763B probe	N/A	N/A	GTT TAC TAC CCT AGC TTT CGC	N/A
CFX763mix	**763-783**	*'Ca.* Villogracilis spp.'	**159/260**	**1**	**GTT TAC TAC CCT AGC TKT CGC**	**45**
CFX449	**449-491**	*'Ca.* Sarcinathrix spp.'	**12/14**	**1**	**GGG ATA CCG TCC TTG TCT CT**	**50**
CFX449_C1	449-491	Competitor for CFX449	N/A	N/A	GGG GTA CCG TCC TTG TCT CT	N/A
CFX449_H1	492-516	Helper for CFX449 probe	N/A	N/A	ACG TAG TTA GCC GAG ACT TAT TCC T	N/A
CFX449_H2	422-448	Helper for CFX449 probe	N/A	N/A	TCT CCC AGA AAA GRR GTT TAC GAC CCG	N/A
CFX1151	**1151-1170**	*'Ca.* Sarcinathrix spp.'	**12/14**	**1**	**TTG ACT CCG GCA GTC CCA CT**	**50**
CFX1151_C1	1151-1170	Competitor for CFX1151 probe	N/A	N/A	TTG ACA CCG GCA GTC CCA CT	N/A
CFX1151_H1	1171-1189	Helper for CFX1151 probe	N/A	N/A	ATC CCC ACC TTC CTC CGG T	N/A
CFX1151_H2A	1102-1127	Helper for CFX1151 probe	N/A	N/A	TAA CTA GTA GBG AGG GTT GCG CTC GT	N/A
CFX1151_H2B	1102-1127	Helper for CFX1151 probe	N/A	N/A	TAA CTA GTA GCA GGG GTT GCG CTC GT	N/A
CFX67[⌘]	**67-89**	*'Ca.* Sarcinathrix spp.'	**6/14**	**0**	**TTC CGA AGA TYA GGT TCG**	**35**
CFX67 comp[⌘]	67-89	Competitor for CFX67 probe	N/A	N/A	TTC CGA AGA TCG GGT TCG	N/A
CFX67-H46[⌘]	46-66	Helper for CFX67 probe	N/A	N/A	TTC GAC TTG CAT GTG TTA RGC	N/A
CFX67-H95[⌘]	95-112	Helper for CFX67 probe	N/A	N/A	CCG TRC GCC ACT AAC CYT	N/A

* Coverage of groups as defined in MiDAS database (Release 2.1)(McIlroy *et al.* 2015). Values given as group hits/ group totals; NA = Not applicable; **Recommended optimal formamide concentration for use in FISH hybridizations. # These helper probes are not required for optimal fluorescent signal of their respective probe. Ψ Do not use this competitor if CFX763A is applied with CFX763B. § Do not use this competitor if CFX763B is applied with CFX763A. ⌘These probes are taken from Speirs *et al.* (2011).

Table 2. Abundance estimation: 16S rRNA amplicon sequencing and qFISH (percentage of total).

WWTP	Sample date	Abundance (%)	
		Sequencing*	qFISH
'Ca. Villogracilis'			
Odense North East	September 2015	9.0	sub-group A: 12 ± 4 sub-group B: 1 ± 1
Aalborg East	August 2015	6.0	sub-group A: 11 ± 3 sub-group B: > 1
Horsens	August 2006	6.0	sub-group A: 11 ± 2 sub-group B: > 1
Fredericia	October 2015	3.3	sub-group A: 6 ± 1 sub-group B: > 1
'Ca. Amarilinum'			
Aalborg West	May 2014	11.9	15.8
Bjergmarken	August 2013	33.7	30.2
Odense North West	August 2010	24.2	16.8
'Ca. Sarcinathrix'			
Viby	August 2008	2.3	2
Boeslum	August 2008	2.1	3
Bjermarken	February 2013	1.2	2
Ejby Mølle	May 2008	1.0	1
Aalborg West	February 2012	0.8	1

* Taken from the 16S rRNA gene amplicon sequencing MiDAS survey of Danish WWTPs (McIlroy *et al.* 2015).

Table 3. Summary of the known *in situ* physiology for Chloroflexi commonly found in activated sludge.

Property	'Ca. Amarilinum'	'Ca. Villogracilis'	'Ca. Sarcinathrix'	'Ca. Promineofilum'	'Ca. Defluviifilum'	'Kouleothrix'	Chloroflexi Summary
Substrate uptake							
Formate	ND	ND	ND	ND	−[6]	−[2]	−[2,6]
Acetate	−[1]	−[1]	−[1]	−[9]	−[6,8]	−[2]	+[2]/−[2,6,8,9]
Propionate	−[1]	−[1]	−[1]	−[9]	−[6,8]	−[2]	−[2,6,8,9]
Butyrate	−[1]	−[1]	−[1]	−[9]	−[6]	+[2]/−[2]	+[2]/−[2,6,9]
Lactate	−[1]	−[1]	−[1]	ND	ND	ND	
Pyruvate	−[1]	−[1]	−[1]	−[9]	−[6,8]	+[2]/−[2]	+[2]/−[2,6,8,9]
Oleic acid	ND	ND	ND	−[9]	+[8]/−[6,8]	−[2]	+[5]/−[2,6,8,9]
Glucose	+[1]	−[1]	+[1]/−[1]	+[9]	+[6,8]	+[2]	+[2,3,6,8,9]
Mannose	−[1]	−[1]	−[1]	ND	+[8]/−[6]	+[2]/−[2]	+[2,7]/−[2,6]
Galactose	+[1]/−[1]	−[1]	+[1]/−[1]	ND	+[8]/−[6]	−[2]	+[2,7]/−[2,6]
Fructose	+[1]/−[1]	+[1]/−[1]	+[1]	ND	ND	ND	ND
NAG	−[1]	−[1]	−[1]	−[9]	+[6]/−[6]	+[2]	+[2,3,6]/−[2,9]
Glycine	ND	ND	ND	ND	−[6,8]	−[2]	−[2,6,8]
Leucine	ND	ND	ND	ND	−[6,8]	+[2]/−[2]	+[2]/−[2,6,8]
Amino acid mix	−[1]	−[1]	−[1]	−[9]	ND	ND	−[9]
Thymidine	ND	ND	ND	ND	−[6]	ND	+[2]/−[6]
Ethanol	−[1]	−[1]	−[1]	−[9]	−[6,8]	−[2]	−[2,6,8,9]
Glycerol	−[1]	−[1]	−[1]	−[9]	−[8]	ND	−[8,9]
Chemoautotrophy	−[1]*	−[1]*	−[1]*	ND	−[6]	−[2]**	−[2,6]
Electron acceptor conditions							
Aerobic	+[1]	+[1]/−[1]	+[1]	+[9]	+[6,8]	+[2]	+[2,6,8,9]
Anoxic	+[1]	+[1]/−[1]	+[1]	+[9]	+[6]	−[2]	+[6,9]/−[2,6]
Anoxic + NO$_2^-$	+[1]	+[1]/−[1]	ND	ND	+[6]	−[2]	+[6]/−[2,4,6]
Anoxic + NO$_3^-$	+[1]	+[1]/−[1]	ND	ND	+[6]	−[2]	+[6]/−[2,6]
Storage polymers (staining)							
PHA	−[1]	ND[1]	−[7]	−[5]	−[6]	ND	−[5,6]
Polyphosphate	−[1]	ND[1]	−[7]	−[5]	+[6]/−[6]	ND	+[6]/−[5,6]
Hydrophobic surface (MAC)	−[1]	ND[1]	−[1]	−[9]	+[6]	−[2]	+[6]/−[2,9]

References: 1. This study; **2.** (Kragelund *et al.* 2007); **3.** (Miura, Watanabe and Okabe 2007); **4.** (Morgan-Sagastume, Nielsen and Nielsen 2008); **5.** (Speirs *et al.* 2009); **6.** (Kragelund *et al.* 2011); **7.** (Speirs *et al.* 2011); **8.** (Kindaichi *et al.* 2013); **9.** (McIlroy *et al.* 2016). * carbonate with and without ammonia; ** anaerobic with 2 mM thiosulfate present.

Figure 1. Maximum-likelihood (PhyML) 16S rRNA gene phylogenetic tree of abundant activated sludge phylotypes (bold typeface) and isolated members of the phylum Chloroflexi. The alignment used for the tree applied a 20% conservational filter to remove hypervariable positions, giving 1108 aligned positions. Phylogenetic classification is indicated with black brackets and is based on the MiDAS database (Release 2.1). Probe coverage of probes relevant to the current study is shown with red coloured brackets. Associated activated sludge morphotypes of previously described phylotypes are designated with blue coloured brackets. Bootstrap values from 100 resamplings are indicated for branches with >50% (white dot), 50-70% (gray), and >90% (black) support. Species of the phylum Cyanobacteria were used as the outgroup. The scale bar represents substitutions per nucleotide base.

Figure 2. Distribution of Chloroflexi in 25 full-scale activated sludge WWTPs sampled 2-4 times per year from 2006 to 2015. (a) 10 most abundant phyla in Danish WWTPs. (b) 10 most abundant Chloroflexi classes in Danish WWTPs (c) 10 most abundant Chloroflexi genera in Danish WWTPs. X-axis shows the relative read abundance in percentage of total bacteria.

Figure 3. Composite FISH micrographs of novel Chloroflexi genera in full-scale activated sludge. Specific probes (Cy3-label, red) target (a) '*Ca.* Amarilinum', (b) '*Ca.* Villogracilis', and (c) '*Ca.* Sarcinathrix', and EUBmix probe (Cy5-label, blue) targets most bacteria. Activated sludge was sampled from (a) Bjergmarken WWTP, (b) Odense North East WWTP, and (c) Aalborg West WWTP. Target filaments appear magenta, while all other cells appear blue. Scale bars represent 20 μm.

Figure 4. FISH and corresponding bright-field MAR micrographs showing sugar uptake by three abundant Chloroflexi phylotypes ('*Ca.* Amarilinum' hybridized probe CFX64; '*Ca.* Villogracilis' sub-group A hybridized probe CFX763A; '*Ca.* Sarcinathrix' hybridized probe CFX1151). Activated sludge was sampled in Aalborg West or Odense North West WWTPs. Target cells in FISH micrograph overlays appear yellow: specific probe (red) + EUBmix (green); and non-target cells appear green (EUBmix only). Black silver granules indicate positive MAR signal. Scale bar represents 10 μm.