TITLE:

Stimulus-driven brain rhythms within the alpha band: The attentional-modulation conundrum

ABBREVIATED TITLE: Reversed attentional modulation of alpha and alpha-range SSRs

AUTHORS:

Christian Keitel[1^], Anne Keitel[1], Christopher SY Benwell[1], Christoph Daube[1], Gregor Thut[1], & Joachim Gross[1,2]

AFFILIATIONS:

1 – Institute of Neuroscience and Psychology, University of Glasgow, 58 Hillhead Street, G12 8QB Glasgow, UK; 2 – Institut für Biomagnetismus und Biosignalanalyse, Westfälische Wilhelms-Universität, Malmedyweg 15, 48149 Münster, Germany

^ – corresponding author, christian.keitel@glasgow.ac.uk

KEYWORDS: alpha rhythm, neural rhythm, entrainment, phase synchronisation, visual perception, spatial attention, steady-state response (SSR), steady-state visual evoked potential (SSVEP)

ACKNOWLEDGMENTS: Funded by a Wellcome Trust Joint Investigator Grant awarded to GT and JG (#098433/#098434). Lucy Dewhurst and Jennifer McAllister assisted in data collection. The experimental stimulation was realized using Cogent Graphics developed by John Romaya at the Laboratory of Neurobiology, Wellcome Department of Imaging Neuroscience, University College London (UCL).

0 Table(s), 5 Figure(s), 1 Footnote(s)

ABSTRACT

Two largely independent research lines use rhythmic sensory stimulation to study visual processing. Despite the use of strikingly similar experimental paradigms, they differ crucially in their notion of the stimulus-driven periodic brain responses: One regards them mostly as synchronised (entrained) intrinsic brain rhythms; the other assumes they are predominantly evoked responses (classically termed steady-state responses, or SSRs) that add to the ongoing brain activity. This conceptual difference can produce contradictory predictions about, and interpretations of, experimental outcomes. The effect of spatial attention on brain rhythms in the alpha-band (8 – 13 Hz) constitutes one such instance: alpha-range SSRs have typically been found to increase in power when participants focus their spatial attention on laterally presented stimuli, in line with a gain control of the visual evoked response. In nearly identical experiments, retinotopic decreases in entrained alpha-band power have been reported, in line with the inhibitory function of intrinsic alpha. Here, we demonstrate that this apparent discrepancy results from a small but far-reaching difference in data analysis. By re-analysing EEG data recorded during bilateral rhythmic visual stimulation, we show that averaging across trials before versus after computing power spectra results in the typical patterns reported by SSR and entrainment studies, respectively, therefore reconciling the seemingly contradictory findings. The opposite sign of attention effects suggests that two separate neural processes might be at play simultaneously.

SIGNIFICANCE STATEMENT

The apparent "attentional modulation conundrum" regarding stimulus-driven ~10 Hz brain rhythms stems from two different data analysis approaches. Either approach can be justified depending on the research question. However, only in employing both to the same dataset we found that attention may rely on two distinct cortical mechanisms – retinotopic alpha suppression and increased temporal tracking – to influence the processing of dynamic visual stimulation.

INTRODUCTION

Cortical visual processing has long been studied using rhythmic sensory stimulation (Adrian and Matthews, 1934; Walter et al., 1946; Regan, 1966). This type of stimulation drives continuous brain responses termed steady-state responses (SSRs) that reflect the temporal periodicities in the stimulation precisely. SSRs allow tracking of individual stimuli in multi-element displays (Vialatte et al., 2010; Norcia et al., 2015). Further, they readily indicate cognitive biases of cortical visual processing, such as the selective allocation of attention (Morgan et al., 1996; Keitel et al., 2013; Stormer et al., 2014).

Although SSRs can be driven using a wide range of frequencies (Herrmann, 2001), stimulation at alpha band frequencies (8 – 13 Hz) has stirred particular interest. Alpha rhythms dominate brain activity in occipital visual cortices (Groppe et al., 2013; Keitel and Gross, 2016) and influence perception (Benwell et al., 2017a; Benwell et al., 2017b; Iemi et al., 2017; Samaha et al., 2017). Researchers have therefore used alpha-rhythmic visual stimulation in attempts to align the phase of – or *entrain* – intrinsic alpha rhythms and consequently provided evidence for visual alpha entrainment (Mathewson et al., 2012; Zauner et al., 2012; Spaak et al., 2014; Gulbinaite et al., 2017). These findings suggest that at least part of the SSR driven by alpha-band stimulation should be attributed to entrained alpha generators (Notbohm et al., 2016).

Some issues remain with such an account (Capilla et al., 2011; Keitel et al., 2014). For instance, experiments have consistently reported SSR power increases when probing effects of spatial selective attention on SSRs driven by lateralised hemifield stimuli (Müller et al., 1998), also when using alpha-band frequencies (Kim et al., 2007; Kashiwase et al., 2012; Keitel et al., 2013). However, a recent study that used a similar paradigm, but treated alpha-frequency SSRs as phase-entrained alpha rhythms, reported the opposite effect (Kizuk and Mathewson, 2017). Oscillatory brain activity showed attentional modulations characteristic of the intrinsic alpha rhythm during stimulation: Alpha power decreased over the hemisphere contralateral to the attended position, an effect known to be part of a retinotopic alpha power lateralisation during selective spatial attention (Worden et al., 2000; Kelly et al., 2006; Thut et al., 2006; Rihs et al., 2007; Capilla et al., 2012). Briefly put, studies analysing SSRs show a power *increase,* whereas studies analysing "entrained alpha" show a power *decrease.* Both neural responses originate from visual cortices contralateral to the hemifield position of the driving stimuli (Keitel et al., 2013; Spaak et al., 2014). Assuming a single underlying neural process, reversed attention effects therefore seemingly contradict each other.

To reconcile these disparate findings, we turned to the respective data analysis approaches. Results in support of alpha entrainment differed in a small but potentially crucial detail from studies showing

the opposite effect, namely how exactly responses to the periodic stimulation were quantified. Effects consistent with SSR modulation resulted from spectral decompositions on trial-averaged EEG waveforms that yield power spectra with distinct peaks at the stimulation frequencies. This approach tunes the resulting power estimate to the part of the neural response that is sufficiently time-locked to the stimulation. Effects consistent with alpha entrainment instead resulted from averages of single-trial spectral transforms, thus emphasising intrinsic non-phase-locked activity. Hereby, narrow-band SSRs can be masked by a strong alpha peak (Covic et al., 2017).

We compared both approaches in a re-analysis of previously reported EEG data (Keitel et al., 2017b). Participants viewed two lateralised stimuli, both flickering at alpha band frequencies (10 and 12 Hz). They were cued to focus on one of the two and perform a target detection task at the attended position. We quantified spectral power estimates according to both approaches described above from the same EEG data. Should the outcome depend on the approach taken, we expected to find the typical alpha power lateralisation (contralateral < ipsilateral) when averaging single-trial spectral decompositions. Conversely, when applying the spectral decomposition to trial-averaged EEG, we expected SSR power modulation in the opposite direction (contralateral > ipsilateral).

Figure 1 Stimulus schematics and trial time course. (**a**) shows the exemplary time course of one trial with a cue displayed for 0.5 sec (here: Attend Right), followed by the bilateral visual stimulation for 3.5 sec. Left (L) stimulus contrast fluctuated with a rate of 10 Hz and Right (R) stimulus contrast at 12 Hz. Targets that participants were instructed to respond to were slightly altered versions of the stimuli (see inset) that were displayed occasionally for 0.3 sec. (**b**) Rhythmic visual stimulation was achieved by a frame-by-frame adjustment of global stimulus contrast (through local luminance changes) as exemplified here in one representative cycle.

METHODS

Participants

For the present report, we re-analysed EEG data of 17 volunteers recorded in an earlier study (Keitel et al., 2017a). Participants (13 women; median age = 22 yrs, range = 19 – 32 yrs) declared normal or corrected-to-normal vision and no history of neurological diseases or injury. All procedures were approved by the ethics committee of the College of Science & Engineering at the University of Glasgow (application no. 300140020) and adhered to the guidelines for the treatment of human subjects in the Declaration of Helsinki. Volunteers received monetary compensation of £6/h. They gave informed written consent before participating in the experiment. Note that we excluded five additional datasets on grounds reported in the original study (four showed excessive eye movements, one underperformed in the task).

Stimulation

Participants viewed experimental stimuli on a computer screen (refresh rate = 100 frames per sec) at a distance of 0.8 m that displayed a grey background (luminance = 6.5 cd/m^2). Small concentric circles in the centre of the screen served as a fixation point (*Figure 1*). Two blurry checkerboard patches (horizontal/vertical diameter = 4° of visual angle) were positioned at an eccentricity of 4.4° from central fixation, one each in the lower left and lower right visual quadrants. Both patches changed contrast rhythmically during trials: Stimulus contrast against the background was modulated by varying patch peak luminance between 7.5 cd/m^2 (minimum) and 29.1 cd/m^2 (maximum).

On each screen refresh, peak luminance changed incrementally to approach temporally smooth contrast modulations as opposed to a simple on-off flicker (Andersen and Muller, 2015). Further details of the stimulation can be found in Keitel et al. (2017a). The contrast modulation followed a 10-Hz periodicity for the left and a 12-Hz periodicity for the right stimulus. Note that the experiment featured further conditions displaying quasi-rhythmic contrast modulations in different frequency bands. Corresponding results can be found in the original report and will not be considered in the present analysis.

Procedure and Task

Participants performed the experiment in an acoustically dampened and electromagnetically shielded chamber. In total, they were presented with 576 experimental trials, subdivided into 8 blocks with durations of ~5 min each. Between blocks, participants took self-paced breaks. Prior to the experiment, participants practiced the behavioural task (see below) for at least one block. After each block they received feedback regarding their accuracy and response speed. The experiment was

comprised of 8 conditions (= 72 trials each) resulting from a manipulation of the two factors attended position (left vs. right patch) and stimulation frequency (one rhythmic and three quasi-rhythmic conditions) in a fully balanced design. Trials of different conditions were presented in pseudo-random order. As stated above, the present study focussed on the two conditions featuring fully rhythmic stimuli. Corresponding trials (*N* = 144) were thus selected a posteriori from the full design.

Single trials began with cueing participants to attend to the left or right stimulus for 0.5 sec, followed by presentation of the dynamically contrast-modulating patches for 3.5 sec (*Figure* 1). After patch offset, an idle period of 0.7 sec allowed participants to blink before the next trial started.

To control whether participants maintained a focus of spatial attention, they were instructed to respond to occasional brief "flashes" (0.3 sec) of the cued stimulus (= targets) while ignoring similar events in the other stimulus (= distracters). Targets and distracters occurred in one third of all trials and up to 2 times in one trial with a minimum interval of 0.8 sec between subsequent onsets. Detection was reported as speeded responses to flashes (recorded as space bar presses on a standard key board).

Behavioural data recording and analyses

Flash detections were considered a 'hit' when a response occurred from 0.2 to 1 sec after target onset. Delays between target onsets and responses were considered reaction times (RT). Statistical comparisons of mean accuracies (proportion of correct responses to the total number of targets and distracters) and median RTs between experimental conditions were conducted and reported in (2017a). In the present study, we did not consider the behavioural data further. Note that the original statistical analysis found that task performance in Attend-Left and Attend-Right conditions was comparable.

Electrophysiological data recording

EEG was recorded from 128 scalp electrodes and digitally sampled at a rate of 512 Hz using a BioSemi ActiveTwo system (BioSemi, Amsterdam, Netherlands). Scalp electrodes were mounted in an elastic cap and positioned according to an extended 10-20-system (Oostenveld and Praamstra, 2001). Lateral eye movements were monitored with a bipolar outer canthus montage (horizontal electro-oculogram). Vertical eye movements and blinks were monitored with a bipolar montage of electrodes positioned below and above the right eye (vertical electro-oculogram).

Electrophysiological data pre-processing

From continuous data, we extracted epochs of 5 s, starting 1 s before patch onset using the MATLAB toolbox EEGLAB (Delorme and Makeig, 2004). In further pre-processing, we excluded epochs that corresponded to trials containing transient targets and distracters (24 per condition) as well as epochs with horizontal and vertical eye movements exceeding 20 μV (~ 2° of visual angle) or containing blinks. For treating additional artefacts, such as single noisy electrodes, we applied the 'fully automated statistical thresholding for EEG artefact rejection' (FASTER; Nolan et al., 2010). This procedure corrected or discarded epochs with residual artefacts based on statistical parameters of the data. Artefact correction employed a spherical-spline-based channel interpolation. Epochs with more than 12 artefact-contaminated electrodes were excluded from analysis.

From 48 available epochs per condition, we discarded a median of 14/15 epochs (Attend Left/ -Right conditions) per participant with a between-subject range of 6/8 to 28/31 epochs. Within-subject variation of number of epochs per condition remained small with a median difference of 3 trials (maximum difference = 9 for one participant).

Subsequent analyses were carried out in Fieldtrip (Oostenveld et al., 2011) in combination with custom-written routines. We extracted segments of 3 s starting 0.5 s after patch onset from pre-processed artefact-free epochs (5 s). Data prior to stimulation onset (1 s), only serving to identify eye movements shortly before and during cue presentation, were omitted. To attenuate the influence of stimulus-onset evoked activity on EEG spectral decomposition, the initial 0.5 s of stimulation were excluded. Lastly, because stimulation ceased after 3.5 s, we also discarded the final 0.5 s of original epochs.

Electrophysiological data analyses – spectral decomposition

Artefact-free 3-sec Epochs were converted to scalp current densities (SCDs), a reference-free measure of brain electrical activity (Ferree, 2006; Kayser and Tenke, 2015), by means of the spherical spline method (Perrin et al., 1987) as implemented in Fieldtrip (function *ft_scalpcurrentdensity*, method 'spline', lambda = 10^{-4}). Detrended (i.e. mean and linear trend removed) SCD time series were then Tukey-tapered and subjected to Fourier transforms while employing zero-padding in order to achieve a frequency-resolution of 0.25 Hz. Crucially, from resulting complex Fourier spectra we calculated two sets of aggregate power spectra with slightly different approaches. First, we calculated power spectra as the average of squared absolute values of complex Fourier spectra (Z) as follows:

$$onPOW(f) = \frac{1}{n} \sum_{i=1}^{n} |Z_i(f)|^2 \qquad [1]$$

where *onPOW* is the classical power estimate for ongoing (intrinsic) oscillatory activity for frequency *f* and *n* is the number of trials. Secondly, we additionally calculated the squared absolute value of the averaged complex Fourier spectra according to:

$$evoPOW(f) = \left| \frac{1}{n} \sum_{i=1}^{n} Z_i(f) \right|^2 \qquad [2]$$

The formula yields *evoPOW*, or evoked power, an estimate that is identical with the frequency-tagging standard approach of averaging per-trial EEG time series before spectral decomposition. This step is usually performed to retain only the truly phase-locked response to the stimulus. Note that both formulas only differ in the order in which weighted sums and absolute values are computed. Also note that formula [2] is highly similar to the calculation of inter-trial phase coherence (ITC), a popular measure of phase locking (Cohen, 2014; Gross, 2014; van Diepen and Mazaheri, 2018). ITC calculation additionally includes a trial-by-trial amplitude normalisation. To complement our analysis we thus quantified ITC according to:

$$ITC(f) = \left| \frac{1}{n} \sum_{i=1}^{n} \frac{Z_i(f)}{|Z_i(f)|} \right| \qquad [3]$$

For further analyses, power spectra were normalised by converting them to decibel scale, i.e. taking the decadic logarithm, then multiplying by 10 (hereafter termed log power spectra). ITC was converted to ITCz to reduce the bias introduced by differences in trial numbers between conditions (Bonnefond and Jensen, 2012; Samaha et al., 2015).

Figure 2 EEG spectral decomposition. (**a**) Power spectra collapsed across conditions and all electrode positions below the sagittal midline for single subjects (light grey lines) and group averages (strong black line). Note the characteristic alpha peaks in the frequency range of 8 – 13 Hz. Inset scalp malp shows topographical distribution of alpha power on a dB scale based on scalp current denisities. (**b**) Same as in a but for ‚evoked' power. Distinct peaks are visible at stimulation frequencies 10 & 12 Hz (dashed vertical orange lines across plots). Inset scalp malps shows topographical distribution of SSR power at 10 & 12 on a dB scale. Note the difference in scale between ongoing power in a and evoked power b. (**c**) Same as in a but for inter-trial phase-locking (ITCz). Inset scalp malps shows topographical distribution of SSR ITCz at 10 & 12.

Alpha power – attentional modulation and lateralisation

Spectra of ongoing power (*onPOW*), pooled over both experimental conditions and all electrodes, showed a prominent peak in the alpha frequency range (*Figure 2*). Therefore, we used mean log ongoing power across the range of 8 – 13 Hz to assess intrinsic alpha power modulations by attention. Analysing Attend-Right and Attend-Left conditions separately, yielded two alpha power topographies for each participant. These were compared by means of cluster-based permutation statistics (Maris and Oostenveld, 2007) using *N* = 5000 random permutations. We clustered data across channel neighbourhoods with an average size of 7.9 channels that were determined by triangulated sensor proximity (function *ft_prepare_neighbours*, method 'triangulation'). The resulting probabilities (*P*-values) were corrected for two-sided testing. Subtracting left-lateralised (Attend-Left conditions) from right-lateralised (Attend-Right) alpha power topographies, we found a right-hemispheric positive and a left-hemispheric negative cluster of electrodes that was due to the retinotopic effects of spatial attention on alpha power lateralisation (*Figure 3*), similar to an earlier re-analysis of the other conditions of this experiment (Keitel et al., 2018).

Finally, we tested the difference between Attend-Left and Attend-Right conditions, i.e. attention effects for left- and right-hemispheric clusters separately. To this end we submitted alpha power differences (contralateral hemifield attended minus ignored) to Bayesian one-sample t-tests against zero (Rouder et al., 2009). Attention effects were further compared against each other by means of a Bayesian paired-samples t-test as implemented in JASP (JASP-Team, 2018) with a Cauchy prior scaled to r = 0.5, putting more emphasis on smaller effects (Rouder et al., 2012; Schonbrodt and Wagenmakers, 2017).

This procedure allowed us to quantify the evidence in favour of the null vs the alternative hypotheses (H_0 vs H_1). For each test, the corresponding Bayes factor (called BF_{10}) showed evidence for H_1 (compared to H_0) if it exceeded a value of 3, and no evidence for H_1 if $BF_{10} < 1$, with the intervening range 1 – 3 termed 'anecdotal evidence' by convention (Wagenmakers et al., 2011). Inversing BF_{10}, to

yield a quantity termed BF_{01}, served to quantify evidence in favour of H_0 on the same scale. For BF_{10} and BF_{01} values < 1 were taken as inconclusive evidence for either hypothesis. Note that for the sake of brevity we report errors in BF estimates only when exceeding 0.001%.

SSR power – attentional modulation

Spectra of evoked power, pooled over both experimental conditions and all electrodes, revealed periodic responses to the two stimuli at the respective stimulation frequencies, 10 and 12 Hz (*Figure 2*). Therefore, we assessed attention effects for these two spectral SSR representations. Two separate cluster-based permutation tests, one for each stimulation frequency, contrasted evoked power topographies between attended and ignored (= other stimulus attended) conditions. Two-sided tests were performed with the same parameters as for alpha power (see above).

Again, we found one electrode cluster carrying systematic attention effects per frequency. As for alpha, SSR power from these two clusters were subjected to separate Bayesian one-sample t-tests against zero (one-sided, attended > ignored) and compared against each other by means of a Bayesian paired-sample t-test (two-sided).

SSR inter-trial phase coherence – attentional modulation

We also evaluated a pure measure of neural phase-locking to the stimulation, SSR inter-trial phase coherence (ITC), because evoked power can be regarded as a hybrid measure depending on both the amplitude of the underlying rhythmic response and the consistency of its phase across trials. ITC indicates only the latter as SSRs are set to unit amplitude prior to summing across trials (see formula 3). ITC spectra, pooled over both experimental conditions and all electrodes, showed distinct neural phase-locking at the respective driving frequencies, 10 and 12 Hz (*Figure 2*). Cluster-based permutation testing confirmed topographic regions that showed systemic gain effects in ITC. Subsequently, the same Bayesian inference was applied to data from these clusters as for SSR power.

Correlation of alpha and SSR attention effects

As a consequence of our counter-intuitive finding that SSR attention effects seemed strongest over occipital regions ipsi-lateral to the driving stimulus (see Results section *SSR power & inter-trial phase locking – attentional modulation* below) we explored a posteriori whether these effects could be explained by ipsilateral increases in alpha power during focussed attention. We correlated attention effects on alpha and SSR power using Bayesian inference (rank correlation coefficient Kendall's tau-b or τ_b, beta-prior = 0.75) to test for a positive linear relationship. More specifically, we correlated the left-hemispheric alpha power suppression (Ignored minus Attended) with the 10-Hz SSR (evoked) power attention effect (Attended minus Ignored) and the right-hemispheric alpha power suppression

with the 12-Hz SSR power attention effect. We opted for these combinations because the corresponding effects overlapped topographically (see Results). Along with the correlation coefficient ρ, we report its 95%-Credible Interval (95%-CrI).

We also probed the linear relationship between alpha power and SSR ITC attention effects. Because ITC gains were not clearly lateralised we collapsed gain effects (Attended minus Ignored) across both stimulation frequencies and correlated these with a hemisphere-collapsed alpha suppression index. This index was quantified as the halved sum of left and right-hemispheric suppression effects as retrieved from significant clusters in the topographical analysis of alpha power differences (Attend Left minus Attend Right), shown in *Figure 3*. Again, we expected a positive correlation here if alpha power suppression influenced phase-locking to visual stimulation. For means of comparison, we repeated this analysis with attention effects on SSR power collapsed across frequencies.

Figure 3 Allocation of spatial attention produces retinotopic alpha power modulation. The scalp map (top, center) depicts alpha power lateralisation (Attend Left – Attend right conditions) on a dB scale. Black dots indicate left- and right-hemispheric electrode-clusters that showed a consistent difference in group statistics (two-tailed cluster-based permutation tests). Left and right spectra illustrate alpha power differences in respective clusters when the contralateral hemifield was attended (Att) versus ignored (Ign). The bottom grey inset depicts the distribution of individual alpha power suppression effects (Ignored minus Attended) within left (L) and right (R) hemisphere clusters in the 8 − 13 Hz band. Boxplots indicate interquartile ranges (boxes) and medians (coloured vertical intersectors). Dots below show individual effects (1 dot = 1 participant).

RESULTS

Ongoing alpha power – attentional modulation and lateralisation

The power of the ongoing alpha rhythm lateralised with the allocation of spatial attention to left and right stimuli. A topographic map of the differences in alpha power between Attend-Left and Attend-Right conditions shows significant left- and right-hemispheric electrode clusters (*Figure 3*). These

clusters signify retinotopic alpha power modulation when participants attended to left vs right stimulus positions (right cluster: t_{sum} = -21.454, P = 0.026; left cluster: t_{sum} = 81.264, P = 0.002). The differences are further illustrated in power spectra pooled over electrodes of each cluster (*Figure 3*). As predicted, alpha power at each cluster was lower when participants attended to the contralateral stimulus. Bayesian inference confirmed the alpha power attention effect for the right (M = 0.806 dB, SEM = 0.216; BF_{10} = 21.17) and left cluster (M = 0.790 dB, SEM = 0.133; BF_{10} = 906.36). Both effects were of comparable magnitude (BF_{01} = 4.009 ± 0.007).

SSR power & inter-trial phase locking – attentional modulation

Crucially, we found the opposite pattern when looking at SSRs, i.e. the exact same data but with a slightly different focus on oscillatory brain activity that was time-locked to the stimulation (compare formulas 1 and 2): SSRs showed increased power when the respective driving stimulus was attended versus ignored (*Figure 4*). The power of neural responses evoked by our stimuli (SSRs) was at least one order of magnitude smaller than ongoing alpha power on average (difference > 10dB, i.e. between 10 – 100 times). Nevertheless, SSRs could be clearly identified as distinct peaks in (evoked) power and ITC spectra. Consistent with the retinotopic projection to early visual cortices, topographical distributions of both measures showed a focal maxima contra-lateral to the respective stimulus positions that were attended (*Figure 2*). Counter-intuitively though, maximum attention effects on SSR power did not coincide topographically with sites that showed maximum SSR power overall (compare scalp maps in *Figure 2 & 4*). Also, due to their rather ipsilateral scalp distributions (with respect to the attended location), SSR attention effects did not match topographies of attention-related decreases in ongoing alpha power (compare scalp maps in *Figures 3 & 4*). The 10-Hz SSR driven by the left-hemifield stimulus showed a left-hemispheric power increase when attended (t_{sum} = 15.837, P = 0.059). Similarly, attention increased the power of the 12-Hz SSR driven by the right-hemifield stimulus in a right-hemispheric cluster (t_{sum} = 53.282, P < 0.001). Bayesian inference confirmed the attention effect on 10-Hz (M = 3.727 dB, SEM = 0.919; BF_{10} = 37.05) and 12-Hz SSR power (M = 4.473 dB, SEM = 0.841; BF_{10} = 329.75) averaged within clusters. Both effects were of comparable magnitude (BF_{01} = 3.443 ± 0.005).

SSR phase-locking (quantified as ITCz) also increased with attention to the respective stimulus. In contrast to evoked power, topographical representations of these effects showed greater overlap with the sites that showed maximum phase-locking in general (*Figure 4*). For both frequencies, ITCz increased in central occipital clusters (10 Hz: t_{sum} = 41.351, P = 0.004; 12 Hz: t_{sum} = 31.116, P = 0.012). Again, Bayesian inference confirmed the attention effect on 10-Hz (M = 1.386 au, SEM = 0.297; BF_{10} = 105.71, one-sided) and 12-Hz ITCz (M = 1.824 au, SEM = 0.451; BF_{10} = 36.11, one-sided).

Evidence for a greater attention effect on 12-Hz than on 10-Hz ITC remained inconclusive (BF_{10} = 0.473).

Figure 4 Attention effects on SSR evoked power (evoPow) and SSR inter-trial phase coherence. (**a**) SSR evoked power spectra show systematic power differences at the presentation frequency (10 Hz) of the left stimulus when it was attended (dark red) versus ignored (orange). The inset scalp map illustrates the topographical situation of attention effects. Power spectra were averaged across electrodes (black dots in scalp maps) that showed consistent attention effects in group statistics (two-tailed cluster-based permutation tests) for Attended and Ignored conditions separately. (**b**) Same as in a but for the 12-Hz stimulus presented in the right visual hemifield. (**c,d**) Same as in a,b but using ITCz as a measure of SSR inter-trial phase coherence.

Correlation of alpha and SSR attention effects

Lastly, we tested whether the SSR attention gain effects were mere reflections of the topographically coinciding ipsilateral ongoing alpha power increase during focussed attention that co-occurred with the contralateral ongoing alpha-power decrease (*Figure 3*). Speaking against this account, Bayesian inference provided moderate evidence against the expected positive correlations between the left-hemispheric alpha attention effect and the 10-Hz SSR attention effect (τ_b = -0.221, 95%-CrI = [0.002 0.269]; BF_{01} = 5.811) and between the right-hemispheric alpha attention effect and the 12-Hz SSR attention effect (τ_b = -0.088, 95%-CrI = [0.004 0.315]; BF_{01} = 3.904). These relationships are further illustrated by corresponding linear fits in *Figure 5*.

Following this analysis, we further explored the relationship between spatially non-overlapping decreases in alpha-power contralateral to the attended position and the ipsilateral SSR power gain

effects. For the lack of a specific hypothesis about the sign of the correlation in this case, we quantified the evidence for any relationship (two-sided test). The results remained inconclusive for a correlation between the left-hemispheric alpha attention effect and the right-hemispheric 12-Hz SSR attention effect ($\tau_b = 0.235$, 95%-CrI = [-0.110 0.487]; $BF_{01} = 1.280$) and between the right-hemispheric alpha attention effect and the left-hemispheric 10-Hz SSR attention effect ($\tau_b = 0.103$, 95%-CrI = [-0.218 0.383]; $BF_{01} = 2.400$).

Figure 5 Relationships between attention effects on alpha power and SSRs. (**a**) Individual 10-Hz (left stimulus) SSR evoked power gain (Attended minus Ignored; z-scored, y-axis) as a function of alpha suppression (Ignored minus Attended; z-scored, x axis) in overlapping left-hemispheric parieto-occipital electrode clusters. Grey dots represent participants. Coloured lines depict a straight line fit and its confidence interval (dashed lines). Goodness of fit of the linear model provided as R^2 along with corresponding P-Value. As confirmed by additional tests, both attention effects do not show a positive linear relationship that would be expected if the ipsilateral SSR power gain effect was a consequence of the ipsilateral alpha suppression. (**b**) Same as in a, but for the 12 Hz SSR driven by the right stimulus in overlapping right-hemispheric parieto-occipital electrode clusters. (**c,d**) Similar to a but for attention-related gain effects on SSR ITCz (z-scored, y-axis) in c and gain effects on SSR evoked power, both collapsed across electrode clusters showing 10- and 12-Hz SSR attention effects. Alpha suppression was collapsed across left- and right-hemispheric electrode clusters (see Figure 3).

Finally, we repeated this analysis for attention effects on inter-trial phase coherence (ITC). Because SSR ITC attention effects did not show a clear topographical lateralisation (*Figure 4*), they were collapsed across driving frequencies (10 & 12 Hz). Again, findings were inconclusive when looking into the correlation between these aggregate SSR ITC gain effects and a hemisphere-collapsed alpha

suppression index (τ_b = -0.059, 95%-CrI = [-0.349 0.251]; BF_{01} = 2.653). Correlating collapsed attention effects of SSR evoked power with the same pooled alpha suppression index yielded identical results regarding the rank correlation (also see linear fits in *Figure 5*).

DISCUSSION

Following a recent surge of interest, researchers have attempted to causally drive (entrain) intrinsic alpha rhythms by means of rhythmic visual stimulation with alpha-band frequencies. In support of this notion, stimulus-driven brain oscillations showed modulations by spatial attention that closely resembled the functional characteristics of intrinsic alpha (Kizuk and Mathewson, 2017). However, using similar stimulation paradigms, earlier "frequency-tagging" studies have found opposite gain effects on so-called steady-state responses (SSRs) within the alpha frequency range (Kim et al., 2007; Kashiwase et al., 2012; Keitel et al., 2013). In the present re-analyses of previously published data (Keitel et al., 2017a), we tested whether these conflicting results can be reconciled by assuming that both effects occur simultaneously. More specifically, we assumed that the direction of the effect depended on the particular analysis approach taken to quantify brain responses to the stimuli in the spectral domain.

We found that applying two slightly different approaches to the same dataset can indeed produce reversed outcomes regarding the effects of spatial attention. An approach typically used to evaluate ongoing oscillatory brain activity (averaging single-trial power spectra; see Methods, Formula 1) yielded a modulation of alpha power in line with the known effects of retinotopic alpha lateralisation during focussed spatial attention. In contrast, tuning in on stimulus-locked rhythmic neural activity (by calculating the power spectrum of the trial-averaged waveform; see Methods, Formulas 2 & 3) revealed a reversed pattern.

Analysis approach determines sign of attentional modulation

When focussing on the spectral representation of EEG ongoing power, we observed the prototypical broad peak in the alpha frequency range (8 – 13 Hz; *Figure 2*). Moreover, alpha showed functional modulations in line with numerous earlier reports: Alpha power decreased over the hemisphere contralateral to the attended stimulus position, an effect that has been related to a functional disinhibition of cortical areas representing task-relevant regions of the visual field (Worden et al., 2000; Kelly et al., 2006; Thut et al., 2006). Concurrently, alpha power increased over the ipsilateral hemisphere. This effect has been interpreted as a neural signature of actively suppressing irrelevant and possibly distracting input (Rihs et al., 2007; Capilla et al., 2012).

In a second approach, we focussed instead on the part of the EEG signal that was strictly phase-locked to the visual stimulation, the SSRs. As in classical frequency-tagging studies, we found spectrally distinct SSRs at the stimulation frequencies (here 10 and 12 Hz). These two concurrent rhythmic brain responses thus precisely reflected the temporal dynamics of the visual stimulation. Notably, SSR evoked power was between one to two orders of magnitude (10 – 100 times) lower than ongoing-alpha power, suggesting that the largest share of alpha activity did not synchronise with the stimulation. Smaller evoked power also explained why SSRs remained invisible in spectra of ongoing activity. They were likely masked by the broad alpha peak (*Figure 2*). Crucially, we examined SSRs for effects of focused spatial attention. Visual cortical regions contralateral to the respective driving stimuli showed maximum SSR evoked power. We expected to observe a decrease in SSR evoked power when the driving stimulus was attended (Kizuk and Mathewson, 2017) under the assumption that SSRs are frequency-specific neural signatures of a local entrainment of intrinsic alpha generators (Spaak et al., 2014; Notbohm et al., 2016) and exhibit similar functional characteristics. Instead, we found that SSR evoked power increased in line with earlier reports (Kim et al., 2007; Kashiwase et al., 2012; Keitel et al., 2013).

A caveat of the present data is that the attentional gain effects on SSR evoked power did not coincide topographically with scalp locations of the maximum SSR, or with scalp locations of the attention-related power decrease in ongoing alpha oscillations. Instead, they were most pronounced over hemispheres ipsilateral to the position of the respective driving stimuli. This could have been interpreted as a consequence of the attention-related ipsilateral alpha power increase that accompanied the contralateral power decrease (*Figure 3*). However, a control analysis failed to provide evidence for a positive relationship between ongoing alpha- and SSR evoked power increases that would have been expected in that case (*Figure 5*). We have described the apparent counter-intuitive lateralisation of this effect before (Keitel et al., 2017a) and suggested that attending to a stimulus could also lead to an additional recruitment of upstream visual cortices in contrast to a response gain confined to early visual cortices. More specifically, when comparing scalp distributions by means of Attended-minus-Unattended contrasts, expecting attention effects to emerge at sites of maximum SSR power entails the implicit assumption that attention only acts as a local response gain mechanism. Alternatively, neural representations of attended stimuli could access higher order visual processing (Lithari et al., 2016) and a gain in spatial extent could then produce seemingly ipsilateral effects when evaluating topographical differences as observed here (*Figure 4*). Considering that SSR inter-trial phase coherence showed yet another topographical distribution for gain effects (*Figure 4*), and that EEG only has a limited spatial resolution, warrants a dedicated analysis of underlying cortical sources generating these attention effects in further neuroimaging studies.

Opposing but co-occurring attention effects suggest interplay of distinct attention-related processes

Our analysis compared attention effects between "ongoing" spectral power within the alpha frequency band and a quantity that we have termed SSR "evoked power" and that is commonly used in frequency tagging research (Colon et al., 2012; Porcu et al., 2013; Stormer et al., 2014; Walter et al., 2016; Martinovic and Andersen, 2018). This term is somewhat misleading because it conflates a power estimate with the consistency of the phase of the SSR across trials of the experiment. Inter-trial phase consistency (ITC) is closely related to evoked power but involves an extra normalisation term that abolishes (or at least greatly attenuates) the power contribution[1] (Cohen, 2014; Gross, 2014) and has been used to quantify SSRs before (Ruhnau et al., 2016). The effects of attention on SSR evoked power and ITC are typically interchangeable (Covic et al., 2017; Keitel et al., 2017a). In fact, increased ITC, or phase synchronisation, has been considered the primary effect of attention on stimulus-driven periodic brain responses (Kim et al., 2007; Kranczioch, 2017). Looking at spectral power and ITC separately, as two distinct aspects of rhythmic brain activity, therefore resolves the attentional modulation conundrum: Seemingly opposing attention-related effects likely index different but parallel influences on cortical processing of rhythmic visual input. To avoid confusion, we therefore suggest opting for ITC (or other phase- synchronisation measures, e.g. the cosine similarity index (Chou and Hsu, 2018)) instead of "evoked power" to evaluate SSR attentional modulation.

Incorporating our findings into an account that regards SSRs primarily as stimulus-driven entrainment of intrinsic alpha rhythms would require demonstrating how exactly a decrease in alpha-band power (i.e. the contralateral alpha suppression) can co-occur with increased SSR phase synchronisation. Alternatively, SSRs could predominantly reflect an early cortical mechanism for the tracking of fluctuations in stimulus-specific visual input per se (Keitel et al., 2017a) without the need to assume entrainment (Capilla et al., 2011; Keitel et al., 2014). The underlying neural mechanism might similarly work for a range of rhythmic and quasi-rhythmic stimuli owing to the fact that visual cortex comprises a manifold of different feature detectors that closely mirror changes along the dimensions of colour, luminance, contrast, spatial frequency and more (Buracas et al., 1998; Blaser et al., 2000; Martinovic and Andersen, 2018). Most importantly, for (quasi-)rhythmic sensory input, attention to the driving stimulus may increase neural phase-locking to the stimulus to allow for enhanced tracking

---

[1] In a noisy, finite signal such as the typical second(s)-long EEG epoch, there will always be a positive relationship between the power and inter-trial phase consistency at any frequency as is shown by the greater than zero noise floor in our ITC spectra (*Figure 4*). Also note that ITC only measures SSRs meaningfully if the neurophysiological signal contains a periodic component at the stimulation frequency.

of its dynamics, i.e. increased fidelity. This effect has indeed been observed for quasi-rhythmic low-frequency visual speech signals (Crosse et al., 2015; Park et al., 2016; Hauswald et al., 2018) and task-irrelevant visual stimuli at attended vs ignored spatial locations (Keitel et al., 2017a).

Concurrent retinotopic biasing of visual processing through alpha suppression and stronger neural phase-locking to attended stimuli could therefore be regarded as complimentary mechanisms. Both could act to facilitate the processing of behaviourally relevant visual input in parallel. In this context, SSRs would constitute a special case and easy-to-quantify periodic signature of early visual cortices tracking stimulus dynamics over time. Intrinsic alpha suppression instead may gate the access of sensory information to superordinate visual processing stages (Jensen and Mazaheri, 2010; Zumer et al., 2014) and enhanced ipsilateral alpha power may additionally attenuate irrelevant and possibly distracting stimuli at ignored locations (Capilla et al., 2012). On a neuronal level, this could be implemented as follows: During rest or inattention, occipital neuronal populations synchronise with a strong internal, thalamocortical pacemaker such as alpha. During attentive processing of sensory input, retinotopic alpha suppression releases specific neuronal sub-populations from an internal reign and allows them to track the stimulus dynamics at attended locations. A related mechanism has been observed in the striatum, where local field potentials are dominated by synchronous oscillatory activity across large areas (Courtemanche et al., 2003). However, during task performance focal neuronal populations were found to disengage from these globally synchronous oscillations in a consistent and task-specific manner. At the level of EEG/MEG recordings, such a mechanism could lead to task-related decrease of oscillatory power but increase of coherence or ITC as observed in the current study and previously in the sensorimotor system (Gross et al., 2005; Schoffelen et al., 2005; Schoffelen et al., 2011).

Whereas such an account challenges the occurrence of strictly stimulus-driven alpha entrainment, it may still allow alpha to exert temporally precise top-down influences during predictable and behaviourally relevant rhythmic stimulation – a process that itself could be subject to entrainment (Thut et al., 2011; Nobre et al., 2012; Zoefel et al., 2018).

Conclusion

Our findings reconcile seemingly contradictory findings regarding spatial attention effects on alpha-rhythmic activity, assumed to be entrained by periodic visual stimulation, and SSRs. Focusing on spectral power or phase consistency of the EEG during visual stimulation yielded reversed attention effects in the same dataset. Our findings encourage a careful and consistent choice of measures of ongoing brain dynamics (here power) or measures of stimulus-related activity (here ITC), that should be critically informed by the experimental question, when studying the effects of visuo-spatial

selective attention on the cortical processing of dynamic (quasi-) rhythmic visual stimulation. Again, we emphasise that both data analysis approaches taken here can be equally valid and legitimate, yet they likely index distinct neural phenomena. These can occur simultaneously, as in our case, and may index distinct cortical processes that work in concert to facilitate the processing of visual stimulation at attended locations.

Competing interests

The authors declare no competing interests.

Author contributions

CK designed research, performed research, analysed data and wrote the article. JG designed research analysed data and wrote the article. AK, CSYB, CD and GT designed research and wrote the article.

Data accessibility

Pre-processed EEG data in Fieldtrip format that underlie all analyses reported here are available under osf.io/apsyf on the Open Science Framework (Keitel et al., 2017b).

References

Adrian ED, Matthews BH (1934) The interpretation of potential waves in the cortex. J Physiol 81:440-471.

Andersen SK, Muller MM (2015) Driving steady-state visual evoked potentials at arbitrary frequencies using temporal interpolation of stimulus presentation. BMC Neurosci 16:95.

Benwell CSY, Keitel C, Harvey M, Gross J, Thut G (2017a) Trial-by-trial co-variation of pre-stimulus EEG alpha power and visuospatial bias reflects a mixture of stochastic and deterministic effects. Eur J Neurosci.

Benwell CSY, Tagliabue CF, Veniero D, Cecere R, Savazzi S, Thut G (2017b) Prestimulus EEG Power Predicts Conscious Awareness But Not Objective Visual Performance. eNeuro 4.

Blaser E, Pylyshyn ZW, Holcombe AO (2000) Tracking an object through feature space. Nature 408:196-199.

Bonnefond M, Jensen O (2012) Alpha oscillations serve to protect working memory maintenance against anticipated distracters. Curr Biol 22:1969-1974.

Buracas GT, Zador AM, DeWeese MR, Albright TD (1998) Efficient discrimination of temporal patterns by motion-sensitive neurons in primate visual cortex. Neuron 20:959-969.

Capilla A, Pazo-Alvarez P, Darriba A, Campo P, Gross J (2011) Steady-state visual evoked potentials can be explained by temporal superposition of transient event-related responses. PLoS One 6:e14543.

Capilla A, Schoffelen J-M, Paterson G, Thut G, Gross J (2012) Dissociated α-Band Modulations in the Dorsal and Ventral Visual Pathways in Visuospatial Attention and Perception. Cerebral Cortex.

Chou EP, Hsu SM (2018) Cosine similarity as a sample size-free measure to quantify phase clustering within a single neurophysiological signal. J Neurosci Methods 295:111-120.

Cohen MX (2014) Analyzing neural time series data: theory and practice. Cambridge, Massachusetts: MIT Press.

Colon E, Nozaradan S, Legrain V, Mouraux A (2012) Steady-state evoked potentials to tag specific components of nociceptive cortical processing. Neuroimage 60:571-581.

Courtemanche R, Fujii N, Graybiel AM (2003) Synchronous, focally modulated beta-band oscillations characterize local field potential activity in the striatum of awake behaving monkeys. Journal of Neuroscience 23:11741-11752.

Covic A, Keitel C, Porcu E, Schroger E, Muller MM (2017) Audio-visual synchrony and spatial attention enhance processing of dynamic visual stimulation independently and in parallel: A frequency-tagging study. Neuroimage 161:32-42.

Crosse MJ, Butler JS, Lalor EC (2015) Congruent Visual Speech Enhances Cortical Entrainment to Continuous Auditory Speech in Noise-Free Conditions. J Neurosci 35:14195-14204.

Delorme A, Makeig S (2004) EEGLAB: an open source toolbox for analysis of single-trial EEG dynamics including independent component analysis. J Neurosci Methods 134:9-21.

Ferree TC (2006) Spherical splines and average referencing in scalp electroencephalography. Brain Topogr 19:43-52.

Groppe DM, Bickel S, Keller CJ, Jain SK, Hwang ST, Harden C, Mehta AD (2013) Dominant frequencies of resting human brain activity as measured by the electrocorticogram. Neuroimage 79:223-233.

Gross J (2014) Analytical methods and experimental approaches for electrophysiological studies of brain oscillations. J Neurosci Methods 228:57-66.

Gross J, Pollok B, Dirks M, Timmermann L, Butz M, Schnitzler A (2005) Task-dependent oscillations during unimanual and bimanual movements in the human primary motor cortex and SMA studied with magnetoencephalography. Neuroimage 26:91-98.

Gulbinaite R, van Viegen T, Wieling M, Cohen MX, VanRullen R (2017) Individual Alpha Peak
	Frequency Predicts 10 Hz Flicker Effects on Selective Attention. J Neurosci 37:10173-10184.

Hauswald A, Lithari C, Collignon O, Leonardelli E, Weisz N (2018) A Visual Cortical Network for
	Deriving Phonological Information from Intelligible Lip Movements. Curr Biol 28:1453-1459
	e1453.

Herrmann CS (2001) Human EEG responses to 1-100 Hz flicker: resonance phenomena in visual
	cortex and their potential correlation to cognitive phenomena. Exp Brain Res 137:346-353.

Iemi L, Chaumon M, Crouzet SM, Busch NA (2017) Spontaneous Neural Oscillations Bias Perception
	by Modulating Baseline Excitability. J Neurosci 37:807-819.

JASP-Team (2018) JASP. In, 0.8.2 Edition.

Jensen O, Mazaheri A (2010) Shaping functional architecture by oscillatory alpha activity: gating by
	inhibition. Front Hum Neurosci 4:186.

Kashiwase Y, Matsumiya K, Kuriki I, Shioiri S (2012) Time courses of attentional modulation in neural
	amplification and synchronization measured with steady-state visual-evoked potentials. J
	Cogn Neurosci 24:1779-1793.

Kayser J, Tenke CE (2015) On the benefits of using surface Laplacian (current source density)
	methodology in electrophysiology. Int J Psychophysiol 97:171-173.

Keitel A, Gross J (2016) Individual Human Brain Areas Can Be Identified from Their Characteristic
	Spectral Activation Fingerprints. PLoS Biology 14:e1002498.

Keitel C, Quigley C, Ruhnau P (2014) Stimulus-Driven Brain Oscillations in the Alpha Range:
	Entrainment of Intrinsic Rhythms or Frequency-Following Response? Journal of
	Neuroscience 34:10137-10140.

Keitel C, Thut G, Gross J (2017a) Visual cortex responses reflect temporal structure of continuous
	quasi-rhythmic sensory stimulation. Neuroimage 146:58-70.

Keitel C, Andersen SK, Quigley C, Muller MM (2013) Independent Effects of Attentional Gain Control
	and Competitive Interactions on Visual Stimulus Processing. Cerebral Cortex 23:940-946.

Keitel C, Benwell CSY, Thut G, Gross J (2017b) Alpha during quasi-periodic visual stimulation. In. Open
	Science Framework.

Keitel C, Benwell CSY, Thut G, Gross J (2018) No changes in parieto-occipital alpha during neural
	phase locking to visual quasi-periodic theta-, alpha-, and beta-band stimulation. Eur J
	Neurosci.

Kelly SP, Lalor EC, Reilly RB, Foxe JJ (2006) Increases in alpha oscillatory power reflect an active
	retinotopic mechanism for distracter suppression during sustained visuospatial attention. J
	Neurophysiol 95:3844-3851.

Kim YJ, Grabowecky M, Paller KA, Muthu K, Suzuki S (2007) Attention induces synchronization-based response gain in steady-state visual evoked potentials. Nat Neurosci 10:117-125.

Kizuk SA, Mathewson KE (2017) Power and Phase of Alpha Oscillations Reveal an Interaction between Spatial and Temporal Visual Attention. J Cogn Neurosci 29:480-494.

Kranczioch C (2017) Individual differences in dual-target RSVP task performance relate to entrainment but not to individual alpha frequency. PLoS One 12:e0178934.

Lithari C, Sanchez-Garcia C, Ruhnau P, Weisz N (2016) Large-scale network-level processes during entrainment. Brain Res 1635:143-152.

Maris E, Oostenveld R (2007) Nonparametric statistical testing of EEG- and MEG-data. J Neurosci Methods 164:177-190.

Martinovic J, Andersen SK (2018) Cortical summation and attentional modulation of combined chromatic and luminance signals. Neuroimage 176:390-403.

Mathewson KE, Prudhomme C, Fabiani M, Beck DM, Lleras A, Gratton G (2012) Making waves in the stream of consciousness: entraining oscillations in EEG alpha and fluctuations in visual awareness with rhythmic visual stimulation. J Cogn Neurosci 24:2321-2333.

Morgan ST, Hansen JC, Hillyard SA (1996) Selective attention to stimulus location modulates the steady-state visual evoked potential. Proceedings of the National Academy of Sciences of the United States of America 93:4770-4774.

Müller MM, Teder-Sälejärvi W, Hillyard SA (1998) The time course of cortical facilitation during cued shifts of spatial attention. Nat Neurosci 1:631-634.

Nobre AC, Rohenkohl G, Stokes M (2012) Nervous anticipation: top-down biasing across space and time. In: Cognitive Neuroscience of Attention, 2 Edition (Posner MI, ed), pp 159-186. New York: Guilford.

Nolan H, Whelan R, Reilly RB (2010) FASTER: Fully Automated Statistical Thresholding for EEG artifact Rejection. J Neurosci Methods 192:152-162.

Norcia AM, Appelbaum LG, Ales JM, Cottereau BR, Rossion B (2015) The steady-state visual evoked potential in vision research: A review. J Vis 15:4.

Notbohm A, Kurths J, Herrmann CS (2016) Modification of Brain Oscillations via Rhythmic Light Stimulation Provides Evidence for Entrainment but Not for Superposition of Event-Related Responses. Front Hum Neurosci 10:10.

Oostenveld R, Praamstra P (2001) The five percent electrode system for high-resolution EEG and ERP measurements. Clin Neurophysiol 112:713-719.

Oostenveld R, Fries P, Maris E, Schoffelen JM (2011) FieldTrip: Open source software for advanced analysis of MEG, EEG, and invasive electrophysiological data. Comput Intell Neurosci 2011:156869.

Park H, Kayser C, Thut G, Gross J (2016) Lip movements entrain the observers' low-frequency brain oscillations to facilitate speech intelligibility. Elife 5.

Perrin F, Pernier J, Bertrand O, Giard MH, Echallier JF (1987) Mapping of scalp potentials by surface spline interpolation. Electroencephalogr Clin Neurophysiol 66:75-81.

Porcu E, Keitel C, Muller MM (2013) Concurrent visual and tactile steady-state evoked potentials index allocation of inter-modal attention: a frequency-tagging study. Neurosci Lett 556:113-117.

Regan D (1966) Some characteristics of average steady-state and transient responses evoked by modulated light. Electroencephalogr Clin Neurophysiol 20:238-248.

Rihs TA, Michel CM, Thut G (2007) Mechanisms of selective inhibition in visual spatial attention are indexed by alpha-band EEG synchronization. Eur J Neurosci 25:603-610.

Rouder JN, Morey RD, Speckman PL, Province JM (2012) Default Bayes factors for ANOVA designs. Journal of Mathematical Psychology 56:356-374.

Rouder JN, Speckman PL, Sun DC, Morey RD, Iverson G (2009) Bayesian t tests for accepting and rejecting the null hypothesis. Psychonomic Bulletin & Review 16:225-237.

Ruhnau P, Keitel C, Lithari C, Weisz N, Neuling T (2016) Flicker-Driven Responses in Visual Cortex Change during Matched-Frequency Transcranial Alternating Current Stimulation. Front Hum Neurosci 10:184.

Samaha J, Iemi L, Postle BR (2017) Prestimulus alpha-band power biases visual discrimination confidence, but not accuracy. Conscious Cogn 54:47-55.

Samaha J, Bauer P, Cimaroli S, Postle BR (2015) Top-down control of the phase of alpha-band oscillations as a mechanism for temporal prediction. Proc Natl Acad Sci U S A 112:8439-8444.

Schoffelen JM, Oostenveld R, Fries P (2005) Neuronal coherence as a mechanism of effective corticospinal interaction. Science 308:111-113.

Schoffelen JM, Poort J, Oostenveld R, Fries P (2011) Selective movement preparation is subserved by selective increases in corticomuscular gamma-band coherence. J Neurosci 31:6750-6758.

Schonbrodt FD, Wagenmakers EJ (2017) Bayes factor design analysis: Planning for compelling evidence. Psychon Bull Rev.

Spaak E, de Lange FP, Jensen O (2014) Local entrainment of alpha oscillations by visual stimuli causes cyclic modulation of perception. J Neurosci 34:3536-3544.

Stormer VS, Alvarez GA, Cavanagh P (2014) Within-hemifield competition in early visual areas limits the ability to track multiple objects with attention. J Neurosci 34:11526-11533.

Thut G, Schyns PG, Gross J (2011) Entrainment of perceptually relevant brain oscillations by non-invasive rhythmic stimulation of the human brain. Front Psychol 2:170.

Thut G, Nietzel A, Brandt SA, Pascual-Leone A (2006) Alpha-band electroencephalographic activity over occipital cortex indexes visuospatial attention bias and predicts visual target detection. J Neurosci 26:9494-9502.

van Diepen RM, Mazaheri A (2018) The Caveats of observing Inter-Trial Phase-Coherence in Cognitive Neuroscience. Sci Rep 8:2990.

Vialatte FB, Maurice M, Dauwels J, Cichocki A (2010) Steady-state visually evoked potentials: focus on essential paradigms and future perspectives. Prog Neurobiol 90:418-438.

Wagenmakers EJ, Wetzels R, Borsboom D, van der Maas HL (2011) Why psychologists must change the way they analyze their data: the case of psi: comment on Bem (2011). J Pers Soc Psychol 100:426-432.

Walter S, Keitel C, Muller MM (2016) Sustained Splits of Attention within versus across Visual Hemifields Produce Distinct Spatial Gain Profiles. J Cogn Neurosci 28:111-124.

Walter WG, Dovey VJ, Shipton H (1946) Analysis of the Electrical Response of the Human Cortex to Photic Stimulation. Nature 158:540-541.

Worden MS, Foxe JJ, Wang N, Simpson GV (2000) Anticipatory biasing of visuospatial attention indexed by retinotopically specific alpha-band electroencephalography increases over occipital cortex. J Neurosci 20:RC63.

Zauner A, Fellinger R, Gross J, Hanslmayr S, Shapiro K, Gruber W, Muller S, Klimesch W (2012) Alpha entrainment is responsible for the attentional blink phenomenon. Neuroimage 63:674-686.

Zoefel B, Ten Oever S, Sack AT (2018) The Involvement of Endogenous Neural Oscillations in the Processing of Rhythmic Input: More Than a Regular Repetition of Evoked Neural Responses. Front Neurosci 12:95.

Zumer JM, Scheeringa R, Schoffelen JM, Norris DG, Jensen O (2014) Occipital alpha activity during stimulus processing gates the information flow to object-selective cortex. PLoS Biology 12:e1001965.